TFAM-dependent mitochondrial integrity in regulatory T cells preserves immune landscape and restrains inflammaging

Kai Guo[1, 2, $], Zhihan Wang[3, 4, $], Jappreet Singh Gill[3, $], Trishna Debnath[3], Benu Bansal[2, 3, 9], Rayansh Poojary[3], Jacquiline Kim Correa[3], Hakan Celik[3], Het Mehta[3], Eden Abrham[3], Zachery Even[3], Mansib Rahman[3], Abby Lund Da Costa[3], Shilpi Jain[3], Xusheng Wang[5], Gopal Murugaiyan[6], Adrian T. Ting[7], Junguk Hur[2], Nadeem Khan[8], Holly Brown-Borg[2], Donald A Jurivich[3], Ramkumar Mathur[3, *]

1. Department of Neurology, University of Michigan, Ann Arbor, MI 48109, USA
2. Department of Biomedical Sciences, School of Medicine and Health Sciences, University of North Dakota, Grand Forks, ND 58202, USA
3. Department of Geriatrics, School of Medicine and Health Sciences, University of North Dakota, Grand Forks, ND 58202, USA
4. West China School of Basic Medical Sciences & Forensic Medicine, Sichuan University, Chengdu, Sichuan, China
5. Department of Neurology, University of Tennessee Health Science Center, Memphis, TN 38163, USA
6. Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Boston and Harvard Medical School, Boston, MA 02115, USA
7. Department of Immunology, Mayo Clinic, Rochester, MN 55905, USA
8. Dept of Oral Biology, College of Dentistry, University of Florida, Gainesville, FL 32603, USA
9. Department of Biomedical Data Science, School of Applied Computational Sciences, Meharry Medical College, Nashville, TN 37208, USA

$Authors (Kai Guo, Zhihan Wang and Jappreet Singh Gill) contributed equally

*Correspondence
Ramkumar Mathur, Department of Geriatrics, School of Medicine and Health Sciences, University of North Dakota, Grand Forks, ND 58202, USA
Email: ramkumar.mathur@und.edu

Abstract

Foxp3⁺ regulatory T cells (Tregs) maintain immune homeostasis, yet the process that preserves their stability during aging remain unclear. Mechanistic progress has been hindered by models that ablate Tregs or delete *Foxp3*, which induce acute autoimmunity and prevent longitudinal study of physiological regulatory drift. Here, we establish a dose-dependent mitochondrial framework that preserves Treg lineage survival while permitting gradual metabolic attenuation. Using Treg-restricted TFAM modulation, a complementary haploinsufficient model, and whole-spleen single-cell profiling. We identify lineage-selective immune remodeling characterized by contraction of naïve CD8⁺ and follicular B-cell pools, alteration of CD4⁺ states, expansion of activated Tregs, and emergence of neuroimmune stress linked transcriptional modules that parallel physiological aging. Mechanistically, mitochondrial insufficiency is associated with functional loss of *FOXP3*-centered chromatin coordination and enrichment of NF-κB/NFAT/AP-1 inflammatory and senescence programs while lineage identity remains detectable. Partial mitochondrial attenuation within Tregs alone is sufficient to drive chronic low-grade systemic inflammation, neuromuscular decline, gut microbial restructuring, and elevated microglial responsiveness without Treg depletion. Pharmacologic and microbiota-directed interventions partially reduce inflammatory tone and improve functional metrics. Together, our findings identify TFAM as a key regulator of immune aging and reveal that healthy mitochondrial function in Tregs is essential for protecting against inflammaging and age-associated functional decline.

Introduction

Immune aging is increasingly defined by dysregulated inflammatory tone instead of a uniform reduction in leukocyte counts[1, 2, 3, 4]. Converging evidence from human cohorts and murine models indicates that subtle cytokine elevation, redistribution of lymphoid subsets, and gradual functional drift occur prior to the manifestation of overt immunodeficiency, suggesting that the aging immune system is actively recalibrated rather than passively depleted. The gradual increase of long-lasting, low-grade inflammation is often called inflammaging which reflects disturbed immune balance rather than loss of immune cells, suggesting that regulatory pathways remain present but slowly change in function over time.

Foxp3⁺ regulatory T cells are a metabolically and transcriptionally distinct lineage that rises predominantly within the CD4⁺ T cell compartment and enforce peripheral tolerance and suppress immunopathology[5, 6, 7, 8, 9]. In contrast, conventional effector T cells promote inflammation and primarily utilize glycolysis to support clonal expansion[10, 11]. The metabolic state of Tregs influences

their epigenetic landscape, particularly at the conserved non-coding sequence 2 (CNS2)
enhancer of the Foxp3 gene. Oxidative phosphorylation (OXPHOS) and fatty acid oxidation (FAO)
support the maintenance of an open chromatin configuration at CNS2, facilitating sustained Foxp3
expression[12, 13, 14, 15]. This epigenetic regulation is essential for preserving Treg identity and
suppressive function. Disruption of mitochondrial metabolism in Tregs can lead to decreased
Foxp3 expression and increased susceptibility to acquisition of effector-like programs.
Specifically, impaired OXPHOS and FAO may result in the upregulation of transcription factors
such as T-bet (TBX21), RORγt (RORC), and IRF4, which are associated with effector T cell
lineages. This shift undermines Treg stability and promotes inflammatory responses. Emerging
evidence, including from our own investigations, supports the concept that mitochondrial
dysfunction is not merely correlative, but acts as a proximal driver of Treg instability, fueling
systemic inflammation and age-associated immune remodeling[12, 16, 17, 18]. Progress in dissecting
these mechanisms has been constrained by experimental systems that eliminate Tregs entirely,
including scurfy mutations and diphtheria-toxin based *Foxp3* depletion models, which provoke
rapid systemic autoimmunity and early lethality [19, 20, 21]. Although these approaches firmly
establish the necessity of *Foxp3* immune tolerance, they do not reproduce the gradual metabolic
and transcriptional drift observed in physiological aging, where peripheral mature Tregs persist
but slowly alter function. As a result, distinguishing intrinsic regulatory adaptation from secondary
inflammatory collapse has remained challenging.
88
In recent years, mitochondrial genome integrity has emerged as an intrinsic checkpoint linking
cellular metabolism to immune longevity[9, 22, 23]. Mitochondrial transcription factor A (TFAM), a
nuclear-encoded high-mobility group (HMG) protein, is crucial for maintaining mitochondrial DNA
(mtDNA) integrity and regulating redox homeostasis[24, 25]. Beyond its structural role in mtDNA
packaging, TFAM modulates electron transport chain (ETC) function and limits mitochondrial
reactive oxygen species (ROS) production. Loss of TFAM expression leads to mtDNA depletion,
destabilization of oxidative phosphorylation (OXPHOS) complexes (CI-CIV), and activation of
mitonuclear stress responses These include the mitochondrial unfolded protein response
(UPR^{mt}), integrated stress response (ISR) via ATF4/DDIT3, and downstream inflammatory
signaling involving NF-κB and STAT1[25, 26, 27, 28]. The mitochondrial stress responses influence
chromatin and transcriptional regulators, including histone deacetylases (HDACs), SIRT1, ten-
eleven translocation (TET) enzymes, and polycomb repressive complex 2 (PRC2) components
like EZH2. These regulators, which modulate histone acetylation/methylation and DNA
demethylation states, are critical for *Foxp3* expression and the fidelity of the Treg lineage. Thus,

redox-sensitive epigenetic rewiring under mitochondrial stress may mechanistically compromise
Treg stability and function. However, whether TFAM-dependent mitochondrial integrity operates
as an upstream checkpoint safeguarding immune homeostasis remains an unresolved question.
106
Interestingly, *Tfam* disruption in CD4[+] T-cell compartment has been shown to accelerate systemic
inflammation and tissue decline[17]. Prior studies employing Foxp3-directed TFAM deletion further
demonstrate that mitochondrial impairment in regulatory T cells can precipitate acute
inflammatory pathology or alter tumor progression[12, 16]. However, whether declining mitochondrial
integrity confined specifically to *Foxp3[+]* Tregs is sufficient to reorganize whole-immune
architecture toward an aging-like trajectory rather than merely reproducing inflammatory
syndromes associated with overt regulatory dysfunction has remained unclear. Our single-cell
transcriptomic analyses revealed that *FOXP3[+]* T cells exhibit mitochondrial stress–associated
signatures that track with inflammatory skewing in human aging and frailty, implicating
mitochondrial dysfunction in progressive immune remodeling[18]. Consistent patterns reported in
independent cohorts further strengthen the association between mitochondrial perturbation and
age-related shifts in Treg state and function[30, 31, 32, 29]. Collectively, these single-cell studies
suggest that early mitochondrial dysfunction could involve in the gradual re-shaping of immune
trajectories.
121
Here, to dissect this relationship, we employed two Treg-specific TFAM mouse models: a
conditional knockout (Foxp3[+cre]Tfam[fl/fl]) and a heterozygous model (Foxp3[+cre]Tfam[f/+]), which
exhibits a moderate physiological reduction in TFAM expression. Together, these models
recapitulate aspects of the progressive mitochondrial decline observed during both normal and
pathological aging. Mitochondrial dysfunction in the Treg lineage led to a cascade of
immunological issues, including hyper-inflammatory CD4[+] T cell fate changes and altered gene
expression. Such shifts are consistent with hallmarks of inflammaging, which include chronic low-
grade inflammation, gut microbial dysbiosis, and neuroimmune activation. Together, our study
establishes a scalable experimental framework to directly examine how gradual mitochondrial
decline in persistent Tregs alters systemic inflammatory set-points. Our findings establish
mitochondrial integrity not only as a metabolic feature but also as a modifiable regulator of Treg
quality and immune homeostasis, highlighting mitochondrial maintenance as a potential avenue
to mitigate chronic age-associated inflammation.
135
136   **Results**

Treg-intrinsic TFAM loss in Foxp3+ Tregs reprograms CD4+ T cell lineage trajectories and enforces effector bias

To investigate how mitochondrial TFAM deficiency in *Foxp3+* Tregs perturb systemic CD4+ T cell homeostasis, we conducted single-cell RNA sequencing (scRNA-seq) on whole spleens from Foxp3^CreTfam^f/f (TFAM cKO) and Tfam^f/f (wild-type) (Fig. 1A). This model selectively impairs mitochondrial DNA maintenance in mature Tregs while largely preserving Foxp3 expression and lineage presence, enabling assessment of regulatory *functional* dysfunction rather than the severe autoimmune phenotypes produced by Foxp3-null/scurfy or diphtheria-toxin depletion systems [19, 20, 21]. Following stringent quality control, we retained transcriptomic profiles from 14,198 cells in TFAM cKO and 34,958 cells in wild-type mice, excluding erythroid lineage cells from downstream analyses (Fig. 1B). Based on canonical immune markers, we resolved 21 major immune subsets across 26 transcriptionally distinct clusters conserved across both genotypes (Fig. 1B; Fig. S1A-C). Notably, naïve-associated CD8+ T cells marked by Tcf7 and Lef1 were reduced, as were follicular B-cell populations expressing Cd79b and Fcer2a, whereas total CD4+ abundance remained relatively stable but underwent pronounced transcriptional redistribution. Importantly, the contraction of naïve CD8+ cells was not interpreted as direct mitochondrial impairment within CD8+ cells, but rather as a secondary consequence of altered regulatory set-points consistent with physiological immune aging in which naïve CD8+ attrition is a prominent and early feature of compartmental drift [1, 2].

TFAM deletion in Tregs elicited a striking inflammatory reprogramming of the CD4+ T cell compartment, marked by enrichment of TNF signaling, inflammation-associated aging pathways, and gene signatures linked to tumorigenic transformation (Fig. S1D; Table S1). Upregulation of pro-inflammatory mediators including *S100a6* and *Zfp3612*, coupled with diminished expression of homeostatic regulators such as *Cd79b* and *Fcer2a*, was consistent with premature immune aging (Fig. S1E). To examine how TFAM-deficient Tregs influence CD4+ T cell diversity at a steady state, we profiled subset heterogeneity based on canonical markers. Based on canonical lineage-defining markers, six transcriptionally defined CD4+ T cell subsets were identified: naïve (*Sell, Lef1, Tcf7*), effector memory (TEM; *S100a4, S100a6*), exhausted (*Tnfsf8, Tbc1d4*), conventional Tregs (nTregs; *Ikzf2, Il2ra, Cd74*), activated Tregs (aTregs; *Icos, Rgs1*), and cytotoxic CD4+ T cells (*Ccl5, Gzma, Gzmb*) (Fig. 1C, D). Comparative analysis revealed substantial shifts in subset proportions in TFAM cKO mice (Fig. 1E, Fig. S2A), including marked expansion of aTregs (+105.6%), cytotoxic CD4+ T cells (+250.3%), and TEM cells (+208.4%), alongside reduced frequencies of exhausted (-25.4%) and naïve (-29.8%) subsets, and a modest

reduction in nTregs (-6.4%). These shifts reflect an altered balance favoring pro-inflammatory and
terminally differentiated effector states. Thus, TFAM attenuation preserved lineage identity but
reshaped qualitative fate composition, aligning with aging datasets in which Tregs persist yet
adopt activated, cytokine-responsive phenotypes rather than disappearing [30, 31].

To gain mechanistic insights into these phenotypic alterations, differential gene expression and
pathway enrichment analyses in CD4+ T cells from TFAM cKO versus WT mice uncovered strong
upregulation of TNFα signaling, calcium flux, and cytokine/chemokine networks, with concurrent
downregulation of hematopoietic lineage maintenance programs (Fig. 1F, Fig. S2B, Table S2).
IL-17 signaling was particularly enriched across TEM, exhausted, aTregs, and nTregs, suggesting
a bias toward pro-inflammatory Th17-like conversion a hallmark of increased Treg plasticity.
Recognizing that CD4+ T cell diversity is governed by dynamic lineage trajectories, we next
applied trajectory inference to map differentiation states and fate decisions within CD4+ T cell
compartments. Trajectory inference using Monocle revealed a branched differentiation
architecture with three bifurcation points and two dominant paths (P1 and P2), covering seven
transcriptional states (Fig. 1G-I, Fig. S2C-D). CD4+ T cells from WT mice primarily followed the
P1 trajectory, occupying States 4-6, whereas TFAM cKO cells were enriched in States 1, 2, and
6, indicating a skewing toward the P2 trajectory and altered fate commitment. Naïve T cells, while
concentrated in States 4-6 in both groups, were aberrantly distributed across States 1, 2, and 7
in cKO mice. Notably, aTregs were enriched in State 1, while cytotoxic CD4+ T cells were depleted
in terminal States 1 and 7 (Fig. 1I). The shifts point to preferential lineage commitment along the
alternative P2 trajectory in TFAM-deficient cells, contrasting with the homeostatic P1 path favored
by wild-type counterparts. Next, we applied RNA velocity analysis to confirm the inferred
directionality of fate transitions, supporting Monocle-inferred pseudotime dynamics (Fig. 1J).
Gene expression kinetics along pseudotime identified four temporally distinct clusters. Early
expressed Cluster 1 genes (*Klf3, Sell, Bcl2, Satb1*) were enriched in hematopoietic progenitors,
while Cluster 2 genes (*Anxa2, S100a4, Ccl5, S100a6, Lgals1*) were associated with terminal
effector differentiation and inflammation (Fig. 1K). Strikingly, Cluster 2 was enriched for gene
signatures linked to neurodegenerative diseases, Alzheimer's, Parkinson's, and ALS, as well as
oxidative phosphorylation (Fig. 1K, Table S3). KEGG analysis across Monocle states further
demonstrated that State 1, enriched in TFAM-deficient cells, was associated with
neuroinflammatory and neurodegeneration-related programs, whereas hematopoietic renewal
pathways dominated the WT-favored P1 trajectory (Fig. 1K, Fig. S2D, Table S4). These findings
suggest that Treg mitochondrial dysfunction promotes effector CD4+ T cell programs enriched for

transcriptional features observed in neuroinflammatory and neurodegenerative conditions. Flow
cytometry confirmed the expansion of CD4$^+$CD44^{hi}CD62L^{lo} effector T cells and depletion of
CD4$^+$CD44^{lo}CD62L^{hi} naïve populations in secondary lymphoid organs of TFAM cKO mice (Fig.
1L-M). Functionally, TFAM-deficient Tregs exhibited reduced suppressive capacity, allowing for
enhanced proliferation of wild-type responder CD4$^+$ T cells in co-culture assays (Fig. 1N).
Collectively, these findings indicate that mitochondrial transcriptional attenuation restricted to
Foxp3$^+$ Tregs associates with coordinated splenic immune remodeling characterized by naïve
CD8$^+$ contraction, follicular B-cell reduction, diversification of CD4$^+$ states, selective expansion of
ICOS$^+$ activated Tregs, and enrichment of stress- and IL-17–linked transcriptional programs. In
aggregate, the resulting immune landscape aligns with aging-like immune drift redistribution and
functional reprogramming across compartments rather than a depletion-equivalent phenotype or
synchronized effector expansion typical of acute Treg ablation.

TFAM loss in Tregs unleashes a multi-lineage stress response coupled with inflammatory
and senescence signatures
To investigate how mitochondrial TFAM deficiency disrupts transcriptional networks in Foxp3$^+$
Tregs, particularly those governing inflammation-associated regulons and downstream TNF-
driven signaling implicated in Treg destabilization and effector skewing of CD4$^+$ T cells, we
performed gene regulatory network (GRN) inference using scRNA-seq data from TFAM cKO and
wild-type spleens. We identified 177 high-confidence regulons distributed across CD4$^+$ T cell
subsets (Fig. 2A). Canonical immunoregulatory transcription factors, including *Irf2, Irf8, Irf9,* and
Bcl11a, were preferentially enriched in wild-type Tregs, reflecting preservation of suppressive
lineage identity. In contrast, TFAM-deficient Tregs showed marked activation of inflammatory
regulons such as *Nfkb2, Nfatc1, Rora,* and *Runx2*, consistent with a breach in the mitochondrial
checkpoint that permits effector conversion (Fig. 2A, Fig. S3). Stress-adaptive regulons, including
Egr1, Junb, and *Xbp1*, were enriched in exhausted T cells, aTregs, and even nTregs in TFAM
cKO mice, pointing to a convergent transcriptional stress response. Enrichment of *Runx3* in
cytotoxic CD4$^+$ subsets further highlighted mitochondrial TFAM as a key repressor of cytolytic
plasticity. Notably, upregulation of *Ddit3* across all CD4$^+$ subsets indicated a shared mitochondrial
stress program (Fig. 2A). Targeted expression profiling of inflammatory regulators, including
Prdm1, Nfkb1, Nfkb2, and *Ahr*, showed strong induction in TFAM-deficient Tregs, reinforcing the
destabilization of the regulatory axis (Fig. 2B). The ensuing Systemic inflammation further
activates myeloid-derived cells, such as monocytes and macrophages, enhancing their oxidative
phosphorylation activity. This heightened metabolic state contributes to increased

neuroinflammation, and the activation of neurodegenerative pathways associated with
Parkinson's and Alzheimer's diseases (Fig. 2C). To assess downstream consequences, bulk
RNA-seq of TFAM cKO splenocytes revealed strong induction of cell death pathways, including
apoptosis and necroptosis (Fig. 2D). Immunoblotting confirmed caspase-3 and -8 cleavage, IκBα
degradation, and phosphorylation of RIPK1, RIPK3, and MLKL signatures of RIPK-mediated
immunogenic cell death (Fig. 2E). This was substantiated in situ by Annexin V/PI staining, which
revealed marked increases in apoptotic and necroptotic cell populations in both splenic and
cutaneous tissues (Fig. S4E), indicating systemic cytopathology secondary to mitochondrial
collapse in Tregs. Collectively, these findings lend support to regulatory destabilization caused by
chronic inflammatory injury rather than wholesale Treg lineage elimination.

To determine if Treg-specific mitochondrial dysfunction initiates systemic decline, we monitored
TFAM cKO mice longitudinally. While born at Mendelian ratios, these mice developed progressive
organismal deterioration from 4 weeks, including alopecia, kyphosis, lymphoid organomegaly,
impaired mobility, and early mortality (p<0.0001; Fig. S4A-B). By 8-10 weeks, they developed
severe skeletal muscle atrophy, adipose tissue loss, cortical bone thinning, and markedly reduced
survival relative to wild-type littermates (Fig. S4C). Histological analysis revealed inflammatory
infiltration in the lung, liver, colon, and kidney (Fig. 2F, S5A), and cortical bone loss (Fig. 2F, 2G).
Spleens from TFAM cKO mice exhibited splenomegaly, marginal zone disruption, and follicular
disorganization (Fig. 2H). These phenotypes were consistent with disrupted immune regulation
downstream of Treg-intrinsic mitochondrial dysfunction. Mitospy analysis showed mitochondrial
depolarization and increased oxidative stress in cKO Tregs (Fig. 2I, Fig. S4D, S4E). Multiplex
cytokine profiling revealed a pronounced pro-inflammatory signature characterized by elevated
levels of IL-17, IFN-γ, IL-12, IFN-β, GM-CSF, and IL-27, in parallel with significantly increased
circulating TNF-α and IL-6, reflecting systemic immune activation and inflammatory amplification
(Fig. 2J; Fig. S5B). At the molecular level, cKO mice exhibited upregulation of *Cdkn2a (p16)* and
Cdkn1a (p21) (Fig. 2K), and immunoblots confirmed elevated p53, CDK4, p16, and p21 (Fig. 2L),
indicative of a senescence-like program. Whole-spleen RNA-seq revealed enrichment of
inflammatory and aging-related networks (Fig. 2M). Gene Set Enrichment Analysis (GSEA) from
spleen, skin and brain revealed enrichment of neuroactive signaling, circadian disruption, and
protein trafficking defects pathways associated with immune aging and systemic decline (Fig. 2N,
Table S5). These findings establish TFAM as a critical mitochondrial checkpoint that safeguards
Treg identity, restrains inflammation, and maintains immune equilibrium.

Partial TFAM reduction in Tregs as a critical driver of aging pathologies, neuroinflammation, and cognitive decline

To model the progressive mitochondrial decline characteristic of physiological aging, we transitioned to a Treg-specific TFAM heterozygous mouse (TFAM Het; $Foxp3^{Cre}TFAM^{f/+}$). While complete TFAM ablation in Tregs provides mechanistic insight into the necessity of mitochondrial integrity for immune homeostasis, the rapid-onset pathology of this model precludes longitudinal interrogation of chronic immune aging and neurodegenerative processes. To better understand, the TFAM Het model, where one functional allele of the *Tfam* gene is deleted specifically in CD4$^+$Foxp3$^+$ Tregs, resulting in a sustained ~50% reduction in TFAM expression in CD4$^+$Foxp3$^+$ Tregs. This genetically defined model recapitulates subclinical mitochondrial insufficiency and mirrors the progressive mitochondrial decline characteristic of the human aging trajectory. Supporting its translational relevance, analysis of human PBMC transcriptomes revealed a significant age-dependent decline in TFAM expression, reinforcing its role as a mitochondrial-immune rheostat (Fig. 3A). By 8-10 months, TFAM Het mice exhibited classic hallmarks of aging, including kyphosis, weight loss, and reproductive decline (Fig. 3B-C, Fig. S6A-B).

Histopathological evaluation demonstrated that 8-month-old TFAM cKO mice exhibited pronounced structural degeneration in the spleen and lung, markedly exceeding that observed in both 8-week-old controls and TFAM Het counterparts. Notably, the extent of tissue deterioration in TFAM-deficient mice phenocopied the age-associated pathology typically seen in 2-year-old wild-type mice, underscoring the accelerated aging phenotype driven by Treg-intrinsic mitochondrial dysfunction (Fig. S6C). Senescence markers *Cdkn2a (p16^{Ink4a})* and *Cdkn1a (p21^{Cip1})* were significantly upregulated in splenic CD4$^+$ T cells (Fig. 3D), implicating mitochondrial stress induced senescence. Circulating TNF-α and IL-6 levels were markedly elevated (Fig. 3E), indicative of persistent low-grade inflammation. Neuromuscular evaluation revealed diminished grip strength (Fig. 3F), and behavioral assays demonstrated reduced exploratory activity and locomotor hypoactivity (Fig. 3G), consistent with CNS dysfunction. To interrogate the neuroanatomical consequences of peripheral Treg mitochondrial instability, cresyl violet staining of brain sections revealed substantial neuronal loss in the hippocampus and cortex of 8-month-old TFAM Het mice relative to 8-week counterparts (Fig. 3H-I). PCA plots from longitudinal hippocampal RNA-seq profiling revealed a clear difference between Het8M group (T) and Het 8W group (Y) (Fig. 3J), Notably, compared to both 8-week-old wild-type and age-matched 8-month-old controls we observed marked upregulation of neural stress–associated genes including *Gabra5*, *Slc38a5*, *Calcr*, and *Ganbare* (Fig. 3K). Further compared to wild 8W and comparable

8M TFAM Het mice shown KEGG pathway enrichment shown a marked elevated, inflammatory

signaling, Garabergic synapses neuro-active-ligand-receptor interaction and neurodegeneration-

linked pathways (Fig. 3L). To validate the functional relevance of these molecular changes, we

next assessed neuroimmune phenotypes at the cellular level. Iba1 immunohistochemistry

revealed widespread activation of amoeboid microglia in TFAM Het brains (Fig. 4A-B), with NeuN

co-staining confirming adjacent neuronal loss. Flow cytometric analysis revealed elevated MHC-

I expression on brain-resident CX3CR1$^+$ microglia in late-phase TFAM Het mice (Fig. 4C),

indicating enhanced antigen presentation. Upon TNF-α stimulation, purified CX3CR1$^+$ microglia

from TFAM Het mice exhibited increased expression of IL-1β, TNFR1, and IL-6 transcripts,

highlighting a heightened sensitivity to TNF-driven inflammatory activation compared to WT

controls (Fig. 4C-D). These findings collectively support a peripheral-to-central immunometabolic

cascade, wherein Treg-intrinsic mitochondrial insufficiency initiates systemic inflammation,

senescence, and CNS immune activation. The TFAM Het model reveals that gradual

mitochondrial decline in CD4$^+$ Tregs is sufficient to disrupt systemic immune homeostasis,

propagate neuroinflammation, and accelerate degenerative aging processes.

TFAM insufficiency in Tregs drives microbiota-lipid dysregulation and neuroinflammatory

reprogramming via the gut-brain-immune axis

Mitochondrial metabolism in immune cells is closely coupled to host lipid flux and microbiota-

derived metabolites[32]. We showed that advancing age is accompanied by progressive

dysregulation of host–microbial interactions, forming a bidirectional cycle linking metabolism,

microbial composition, and immune regulation[33]. Recently CD4 T cell therapy shown counteracts

inflammaging and senescence by preserving gut barrier integrity[34]. We therefore asked whether

gradual TFAM reduction within regulatory T cells also alters the intestinal metabolic environment

and downstream immune tone. Stable Treg identity depends on intact oxidative metabolism.

Short-chain fatty acids (SCFAs), especially butyrate and propionate, reinforce FOXP3 expression

through mitochondrial–epigenetic coupling[35, 36, 37]. It has remained unclear whether long-lived

Tregs undergoing slow mitochondrial decline can influence microbial ecology. Using a graded

TFAM-insufficiency strategy that preserves lineage survival while attenuating mitochondrial

transcription, and applying longitudinal 16S sequencing using our previously published protocol

in aging mouse model [33], we observed progressive, age-associated restructuring of gut microbial

communities in TFAM Het mice (Fig. 5A–B). Beta-diversity analyses separated groups by both

genotype and age (Fig. 5B). This pattern indicates gradual community redirection rather than

abrupt dysbiosis. At the phylum level, aged TFAM Het animals showed relative expansion of

Campylobacterota and contraction of Bacteroidota (Fig. 5C). Shared-taxa analysis revealed a
preserved core microbiota across age-matched groups (Fig. 5D). This overlap suggests that these
specific bacterial changes are associated with aging processes common to both genotypes,
highlighting potential age-related microbial shifts that occur irrespective of TFAM status.

Notably, differential abundance analysis identified 18 aging-associated microbial taxa altered
across genotypes, including the depletion of canonical SCFA-producing lineages, such as
Lactobacillus murinus, *Odoribacter*, and *Prevotellaceae_UCG-001* (Fig. 5E). Given that SCFAs
like butyrate are critical for Treg stability, intestinal barrier integrity, and peripheral immune
tolerance, these compositional shifts suggest a collapse in microbial-derived immunoregulatory
tone. To further contextualize these compositional alterations, we performed KEGG-based
predictive metagenomic analyses to infer functional capacities of the microbial communities. In 8-
month-old mice, predicted microbial gene modules were enriched for KEGG pathway annotations
broadly associated with aging- and immune-related processes, including longevity regulation,
RIG-I–like receptor signaling, FoxO signaling, MAPK signaling, and endocannabinoid-associated
pathways, relative to 8-week-old controls (Fig. 5F). Importantly, these annotations reflect
predicted microbial functional potential rather than direct activation of corresponding host
endogenous pathways. Within this framework, lipid metabolism emerged as a particularly
disrupted node to regulate Treg stability, with significant alterations in pathways governing
unsaturated fatty acid biosynthesis [36, 38, 39, 40, 41, 42, 43]. In TFAM Het mice, lipid metabolism was
profoundly disrupted, with significant alterations observed in pathways governing unsaturated
fatty acid biosynthesis, PI3K-Akt signaling, spinocerebellar ataxia pathways, sphingolipid
metabolism, primary bile acid biosynthesis, and fatty acid degradation (Fig. 5G). Concurrently,
neurotransmission-related pathways, including GABAergic and glutamatergic synapses, were
markedly downregulated in aged TFAM Het mice (Fig. 5G). This downregulation aligns with
observed behavioral dysfunction and hippocampal atrophy in 8-month-old TFAM Het mice (Fig
3H, 3I, 4A). These findings support a model wherein mitochondrial impairment in regulatory T
cells perturbs microbial generation of neuromodulator metabolites, disrupting gut-brain signaling
and predisposing the central nervous system to inflammatory remodeling[44, 45, 46, 47].

To mechanistically link microbial changes with host gene regulation, we performed an integrated
correlation analysis between differentially abundant bacterial taxa and host transcriptomes
obtained from brain. This analysis uncovered 323 significant microbiota-host gene interactions
(FDR-adjusted *p* <0.05), mapping a complex regulatory network that bridges microbial

composition and neuroimmune signaling (Fig. 5H). *Helicobacter* species exhibited a strong
positive correlation with *Ntsr2*, a neurotensin receptor involved in both neurotransmission and
microglial activation, suggesting microbial influence over neural inflammatory circuits.
Lactobacillus emerged as a central node, negatively correlating with neuroendocrine transcripts
(*Pomc*, *Sstr5*, *Ntsr2*) and neuronal developmental regulators (*Elk3*, *Sox4*), while positively
correlating with inflammatory mediators (*Sele*, *Mpo*, *Ifitm6*), linking its depletion to neuroimmune
destabilization. *Lachnospiraceae_NK4A136_group* was associated with complement-mediated
inflammation, showing a positive correlation with *C5ar2* and negative regulation of *Sstr5* and *Elk3*,
reinforcing its role in shaping neuroinflammatory tone. To assess downstream inflammatory
consequences associated with microbiota-linked alterations in TFAM Het mice, we quantified
inflammatory cytokine transcripts in peripheral tissues. IL-1β and TNFR1 expression were
significantly elevated in brain tissue (Fig. 5I), indicating the presence of sustained, low-grade
neuroinflammatory signaling. These inflammatory changes occurred in concert with marked shifts
in microbial community composition and predicted microbial metabolic capacity, including
pathways related to lipid and SCFA-associated processes. Together with concurrent shifts in
microbial composition and predicted metabolic capacity, these findings support a model in which
partial TFAM insufficiency in regulatory T cells destabilizes immune homeostasis and increases
susceptibility to age-associated systemic and neuroinflammatory remodeling.
392
Modulation of mitochondrial stress pathways attenuates systemic inflammation and
aging-associated pathology
To determine whether immune pathology associated with Treg-specific TFAM deficiency is
amenable to intervention, we evaluated resveratrol (RSV) treatment in TFAM cKO mice. RSV is
a pleiotropic SIRT1-associated compound with established effects on cellular stress responses
and inflammatory signaling. Prior studies have demonstrated that the protective effects of RSV
across multiple disease contexts are mediated, at least in part, through activation of the PGC-1α–
NRF-1–TFAM axis, thereby promoting mitochondrial biogenesis and transcriptional capacity in a
SIRT1-dependent manner[48, 49]. Accordingly, RSV was administered intraperitoneally at 7 mg/kg
every other day for four weeks (Fig. 6A). Consistent with these reports, RSV treatment was
associated with increased transcript levels of *PGC-1α*, *PGC-1β*, and the mitochondrial
transcription factors *TFAM* and *TFB2M*, indicating engagement of mitochondrial biogenic
programs. At the tissue level, RSV treatment was associated with altered mitochondrial
membrane potential, as reflected by MitoSpy staining in skin sections (Fig. 6B), and coincided
with a marked reduction in systemic inflammatory burden, evidenced by decreased circulating

TNF-α and IL-6 levels (Fig. 6C). In parallel, expression of senescence-associated genes *Cdkn2a*

(p16^{Ink4a}) and *Cdkn1a* (p21^{Cip1}) in splenocytes was reduced, consistent with attenuation of

inflammation-linked senescence programs (Fig. 6D-F). Histopathological analysis further

revealed diminished inflammatory infiltration and improved preservation of tissue architecture in

lungs, liver, and kidney (Fig. 6G).

Treg stability is also influenced by microbiota-derived metabolites, especially short-chain fatty

acids that support oxidative metabolism and chromatin accessibility at the *Foxp3* locus [16, 30, 36, 41,

43]. We therefore evaluated whether environmental metabolic inputs show similar corrective trends.

Fecal microbiota transplantation (FMT) was performed in TFAM heterozygous (TFAM Het) mice,

which exhibit a gradual, non-lethal phenotype more consistent with physiological immune aging

than the acute TFAM cKO model. Because regulatory T cell function and peripheral immune tone

are strongly influenced by microbiota-derived signals, we reasoned that partial TFAM insufficiency

may lower the threshold for microbiota-dependent modulation of host physiology. Antibiotic-

conditioned TFAM Het mice received fecal material from TFAM-sufficient donors over three days,

followed by a three-week recovery period (Fig. 6H). FMT was associated with improved

neuromuscular performance, including increased rotarod endurance and enhanced forelimb grip

strength (Fig. 6I). The observed functional improvements indicate that TFAM insufficiency renders

host physiology responsive to microbiota-dependent modulation. Together, pharmacologic and

microbiota-directed interventions collectively suggest that TFAM-linked mitochondrial

competence in Tregs serves as a modifiable nexus linking cellular bioenergetics, immune

modulation, and organismal aging. These results align with the perspective that mitochondrial and

metabolic support can mitigate the inflammatory and functional aspects of immune aging, rather

than effecting a conclusive reversal (Fig. 6J).

Discussion

The progressive decline in mitochondrial transcriptional integrity is a hallmark of aging, yet its

direct impact on immune regulation and cellular fate remains insufficiently characterized[50, 51, 52, 17,

32, 53]. Specifically, whether the decreasing mitochondrial transcriptional capacity within Tregs

directly initiates immune aging, or merely reflects broader senescent cues remain unresolved.

Here, we mechanistically establish TFAM as a central molecular rheostat of Treg identity and

systemic immune homeostasis. Using Treg-specific physiologically relevant graded TFAM

insufficiency, we demonstrate that progressive disruption of mitochondrial transcriptional output

rewires CD4[+] T cell fate, reshapes the gut microbial ecosystem, and primes the central nervous

system for neuroinflammatory remodeling.

A major limitation in immune aging research has been the absence of physiologically relevant

models that can selectively dissect the cell-intrinsic consequences of mitochondrial dysfunction

within Tregs across the aging trajectory. Existing models involving global TFAM deletion or tissue-

specific knockouts in high-turnover compartments often result in rapid lethality, systemic cytokine

storm, and developmental collapse, precluding longitudinal assessment of immune remodeling[12,]

[16, 24, 54, 55]. Similarly, Treg ablation models such as Foxp3[DTR] or DEREG mice induce severe

autoimmunity and fail to capture the progressive, transcriptionally encoded dysfunction of

suppressive Treg identity observed during physiological aging[19, 56]. Other aging models, including

Ercc1[-/Δ], BubR1[H/H], and Zmpste24[-/-], reflect accelerated DNA repair defects or nuclear envelope

instability, but do not reproduce the slow, cell-specific decline in mitochondrial transcriptional

output characteristic of aging Tregs[57, 58,59].

To overcome these constraints, we developed the Foxp3[Cre]Tfam[f/+] mouse, which induces a stable

~50% reduction in TFAM within Foxp3[+] Tregs, consistent with the age-related TFAM decline

reported in human PBMCs. Unlike Foxp3[Cre]Tfam[f/f] models that succumb to early mortality,

TFAMHet mice remain viable, thereby enabling longitudinal tracking of mitochondrial insufficiency

and its systemic consequences. With age, TFAMHet mice exhibit canonical features of

organismal aging kyphosis, sarcopenia, reproductive decline, and neuromuscular impairment

alongside immune aging hallmarks, including elevated p16[Ink4a], p21[Cip1], increased circulating IL-6

and TNF-α, and a shift of the CD4[+] T cell compartment toward CD44[hi]CD62L[lo] effector memory

phenotypes. In the CNS, TFAMHet mice show heightened microglial activation (CX3CR1[+]MHC-

I[hi]) and hippocampal neuronal loss, consistent with peripheral-to-central immunometabolic

disruption (Fig. 3-4). Collectively, this model provides a genetically precise, temporally scalable

system to dissect how mitochondrial transcriptional fidelity in Tregs orchestrates immunological

aging, gut–brain communication, and tissue integrity. The TFAMHet mouse thus offers a tractable

platform for probing mitochondria-driven immunoregulation with translational relevance to human

aging.

While mitochondrial metabolism is known to sustain FOXP3 expression and Treg function, the

direct role of mitochondrial genome integrity specifically TFAM-dependent transcriptional control,

in orchestrating systemic CD4[+] T cell fate has remained undefined. As aging coincides with both

mitochondrial erosion and effector CD4$^+$ expansion, the mechanistic bridge linking these
processes has not been elucidated. By leveraging single-cell RNA sequencing spleens from
TFAMcKO mice, we uncover a profound reshaping of the CD4$^+$ T cell landscape. TFAM loss in
Tregs drives expansion of cytotoxic Gzmb$^+$Ccl5$^+$ subsets, S100a6$^+$ effector memory (TEM) cells,
and Icos$^+$Rgs1$^+$ aged-like Tregs (aTregs), coupled with depletion of early-stage Tcf7$^+$Lef1$^+$ naïve
and Tnfsf8$^+$ activated/exhaustion-associated populations. These alterations are consistent with
transcriptionally encoded, lineage-intrinsic reprogramming rather than being explained solely by
a secondary inflammatory milieu, as revealed by pseudotime reconstruction and RNA velocity
mapping. Notably, TFAM deficiency enforces commitment along an alternative P2 trajectory
enriched for neuroinflammatory and Th17-like gene modules, including *Anxa2*, *S100a4*, and *Ccl5*,
while repressing transcriptional brakes such as *Satb1* and *Bcl2*, suggesting a premature exit from
naïve states and loss of regulatory restraint. Mechanistic interrogation through gene regulatory
network (GRN) analysis reveals a collapse of Foxp3-lineage stability in TFAM-deficient Tregs,
marked by activation of inflammatory regulons (*Nfkb2*, *Nfatc1*, *Runx2*), cytolytic transcriptional
drivers (*Runx3*, *Prdm1*), and stress-response regulators (*Ddit3*, *Egr1*, *Junb*). The induction of
Ddit3, a hallmark of mitochondrial unfolded protein response (UPR^{mt}), together with
RIPK1/RIPK3/MLKL phosphorylation and caspase activation, is consistent with activation of
mitochondrial stress linked cell-death programs (including necroptosis and apoptotic
components) that could propagate paracrine stress to surrounding CD4+ subsets. This
remodeling is accompanied by transcriptional enrichment for neurodegenerative signatures,
including PI3K-Akt, FoxO, and oxidative phosphorylation pathways supporting a molecular link
between mitochondrial erosion in Tregs and CNS-relevant inflammatory programming. Our
findings delineate TFAM-dependent mitochondrial transcription as a master regulator of CD4$^+$ T
cell fate diversification, acting upstream of immunosenescence and neuroinflammation. By
defining mitochondrial instability in Tregs as a transcriptional checkpoint that governs effector
reprogramming, our study uncovers a novel axis of immune aging in which fidelity of Treg
mitochondrial transcription actively restrains the emergence of pro-inflammatory and potentially
neurotoxic CD4$^+$ states.
503
A central unresolved question in immunometabolism and aging is whether mitochondrial
dysfunction in Tregs actively shapes gut microbiota composition and influences the gut–brain
axis, or whether microbiome alterations are merely downstream byproducts of systemic aging[35,
36, 37]. While prior studies have established that SCFAs, particularly butyrate and propionate,
produced by gut microbiota promote FOXP3 expression and enhance Treg function through

epigenetic remodeling, the reciprocal influence of Treg mitochondrial state on microbial ecology
has remained mechanistically undefined[36, 41, 42, 43, 46]. Our study addresses this critical gap by
demonstrating that partial loss of mitochondrial transcriptional control in Tregs via TFAM
haploinsufficiency is sufficient to disrupt microbial-metabolite circuits essential for immune
regulation and neuroimmune integrity. In TFAMHet mice, we observed a marked depletion of
SCFA-producing bacterial taxa, including *Lactobacillus murinus* and *Odoribacter*, which are
essential for maintaining Treg epigenetic programming and FOXP3 stability[36, 44,60]. This dysbiosis
coincided with broad remodeling of gut lipid metabolic pathways, particularly involving bile acid
biosynthesis, sphingolipid metabolism, and signaling cascades such as PI3K-Akt and FoxO, that
are frequently implicated in aging-related immune and neuronal dysfunction[61, 62].

Importantly, our integrative analysis combining single-cell RNA-seq, lipidomics, microbiome
profiling, and host–microbiota transcriptome correlation revealed that Treg-intrinsic TFAM
insufficiency is associated with a systems-level disruption of gut–brain communication.
Neurotransmission pathways, especially GABAergic and glutamatergic circuits critical for CNS
function, were significantly disrupted in TFAMHet mice, aligning with hippocampal atrophy and
behavioral decline. Furthermore, correlation mapping uncovered 323 robust associations
between microbial taxa and neuroimmune gene networks. Notably, *Helicobacter* spp. positively
correlated with *Ntsr2*, a neurotensin receptor linked to microglial activation, while *Lactobacillus*
abundance negatively correlated with *Pomc*, *Sstr5*, *Ntsr2*, and transcriptional regulators *Elk3* and
Sox4, underscoring a microbial influence on CNS signaling cascades[41, 63]. Elevated expression
of inflammatory mediators such as *Il1β* and *Tnfr1* in TFAMHet tissues further support the
emergence of a chronic, low-grade inflammatory state driven by Treg mitochondrial insufficiency.
Recent studies have shown gut–brain axis is a complex, bidirectional communication network that
links the gastrointestinal tract and the central nervous system through neural, hormonal, and
immune pathways. Together, these findings support a previously underappreciated framework
linking Treg mitochondrial integrity with lipid-associated metabolic programs and microbial
community structure. In this model, Treg-specific TFAM insufficiency is associated with
coordinated changes in microbial composition, reduced representation of SCFA-producing taxa,
and heightened inflammatory signaling, reflecting a destabilized immunometabolic state. Rather
than defining a closed causal feedback circuit or a resolved gut-brain immune equilibrium, our
data position mitochondrial fidelity in Tregs as a critical stabilizing influence on immune–microbial
interactions that shapes susceptibility to systemic and neuroinflammatory remodeling during
aging.

Finally, we provide a proof-of-concept that mitochondrial dysfunction in Tregs is therapeutically
actionable through redox restoration and microbiome modulation. In the TFAMcKO model,
resveratrol an established activator of SIRT1 and PGC-1α signaling enhanced mitochondrial
biogenesis, restored mitochondrial membrane potential, reduced senescence markers, and
suppressed systemic inflammation. These findings align with prior reports demonstrating
resveratrol's capacity to improve mitochondrial function and mitigate age-associated inflammation
across tissues[49, 64]. In the chronic TFAM Het model, fecal microbiota transplantation (FMT)
restored SCFA-producing taxa, improved neuromuscular performance, and normalized immune
signatures. While FMT is increasingly recognized for its ability to modulate systemic immunity and
neurocognitive health via the gut-brain axis[65, 66], its role in regulating Treg mitochondrial function
remains poorly defined. Our data suggests that microbiota-derived metabolites may enhance
peripheral tolerance or indirectly support Treg mitochondrial homeostasis. Together, these
findings position Treg-intrinsic mitochondrial decline as a reversible driver of inflammaging and
support the translational potential of combining metabolic and microbial interventions to combat
age-related immune dysfunction (Fig. 6).

In sum, our study redefines Treg mitochondrial integrity as a central axis of immunological aging.
By uncovering TFAM as a lineage-specific molecular rheostat that governs immune suppression,
microbial composition, and neuroimmune resilience, we provide a unifying framework for the
mitochondrial regulation of systemic aging. These findings open translational opportunities for
mitochondria-targeted immunotherapies in chronic inflammatory diseases, neurodegeneration,
and age-associated immune decline.

Methods

Animals

Foxp3[YFPCre] (Stock No: 016959) and TFAM[f/f] (Stock No: 026123) mice were obtained from
Jackson Laboratory (Bar Harbor, ME) and bred at the UND animal facility to create
Foxp3[Cre]TFAM[f/f]. All mice used in this study were from C57BL/6 backgrounds. Animals were kept
and bred in the UND Animal facility in a pathogen-free unit with a 12-hour light cycle, temperature,
and humidity control. Efforts are being made to reduce animal suffering and needless usage, in
accordance with the Institutional Animal Care and Use Committee's (IACUC) and National
Institutes of Health's Guide for the Care and Use of Laboratory Animals standards.
Histology, immunohistochemistry, and acquisition of images

Formalin-fixed, paraffin-embedded slices of various tissues were deparaffinized on slides using

xylene twice for 5 minutes each. Following that, they were switched to 100% ethanol twice for 3

minutes each and then to 95%, 75%, 65%, 50%, 35%, and 25% ethanol for 3 minutes each. Rinse

the slides for ten minutes under running tap water. The excess water was removed using a wiper

(KIMTECH SCIENCE) and 10% fetal bovine serum (FBS) in PBS was applied to the sections of

the slides and incubated for 30 minutes in the dark at room temperature (RT). FBS was gently

drained from the slides and cleaned with PBS (1X). MitoSpy (excitation (ex)/emission (em) at

577nm/598nm, MitoSpyTM Red CMXros (Cat#424801, BioLegend), After incubation, slides were

washed twice with 1X PBS and a single drop of mounting media (Fluoromount-G™ with DAPI,

Cat# 00-4958-02, Thermo Fisher Scientific) was applied to the tissue sections and the coverslip

was dipped into the mounting solution. For microglia (IBA1$^+$) and neuronal (NeuN$^+$)

immunostaining, mouse brains were fixed in 4% paraformaldehyde (PFA) in 1× PBS,

cryoprotected in 30% sucrose, and cryo-sectioned. Tissue sections were blocked in 0.2% Triton

X-100 and 5% normal goat serum in 1× PBS for 1 hour at room temperature, followed by

incubation with primary antibodies rabbit monoclonal anti-IBA1 (Cat# 17198, 1:1000) and mouse

monoclonal anti-Neuronal N (Cat# 94403, 1:1000) diluted in 0.2% Triton X-100 and 3% normal

goat serum in PBS for 24 hours at 4°C. After three 20-minute washes in PBS, sections were

incubated with appropriate secondary antibodies anti-Rabbit CF™594 antibody (Cat#

SAB4600107) and anti-mouse CF™488 antibody (Cat# SAB4600388) for 1 hour at room

temperature. Nuclei were counterstained using a single drop of Fluoromount-G™ with DAPI (Cat#

00-4958-02, Thermo Fisher Scientific) and sealed with nail polish. Fluorescence images were

acquired using a Leica Thunder Imaging System and an Olympus TIF fluorescence microscope.

Whole-slide histological scans were performed using a Hamamatsu NanoZoomer at the UND

Histology Core. Quantification of immunofluorescence signals was carried out using NIH ImageJ

software.

Western blot

Proteins were isolated from mouse tissue and/or single cells and homogenized in RIPA lysis

buffer containing a 1% protease inhibitor cocktail[67, 68, 69]. The protein concentration was

determined according to the manufacturer's instructions and our previously published protocol

using the Bio-Rad protein assay reagent. SDS-PAGE was used to extract a set quantity of cellular

protein and electroblotted onto an Immobilon-P transfer membrane (Cat# IPVH00010). The

immunological blot was blocked for 1 hour at room temperature with 5% nonfat dry milk in TBST

(25 mM Tris-HCL, pH 7.4, 1.5 M NaCl, 0.05% Tween-20), followed by overnight incubation with

primary antibodies at 4 °C. The next day, the membranes were washed five times at six-minute
intervals and incubated for one hour with HRP-conjugated secondary antibody (A16172,
Invitrogen) (1:1000) in 3% nonfat dry milk in TBST. The blots were then washed five further times
with TBST and developed with enhanced chemiluminescence. The following primary antibodies
were bought from Cell Signaling Technologies (Cell Signaling Technology, MA 01923, USA): RIP
(D94C12) Rabbit mAb #3493, MLKL (D2I6N) Rabbit monoclonal antibody #14993, RIPK3
(E1Z1D) Rabbit monoclonal antibody #13526, p-RIP (D1L3S) Rabbit monoclonal antibody
#65746, p-MLKL Rabbit monoclonal antibody (D6H3V) #91689, p-RIP3(D6W2T) Rabbit
monoclonal antibody #93654, p21 Waf1/Cip1 (12D1) rabbit monoclonal antibody #37543, p53
(1C12) Mouse monoclonal antibody #2524, CDK4 (D9G3E) Rabbit monoclonal antibody #12790,
PCNA (D3H8P) XP® Rabbit monoclonal antibody #13110, p16 INK4A (D3W8G) GAPDH
(D16H11) XP® Rabbit mAb #5174.

RNA isolation and qPCR
Total RNA was isolated from mouse tissues or single cells. The tissues were minced with a scissor
and added to 500 µL of TRIzol reagent (Ambion, cat# 15596018, Life Technologies, USA)[67, 68, 69].
A Fisher Scientific Sonic Dismembrator 100 was used to fully homogenize tissue solutions
(American Laboratory Trading, East Lyme, CT 06333. USA). The upper layer was removed, and
the RNA was precipitated with an equivalent volume of isopropanol and incubated at 55°C for 10
minutes before being resuspended in 50 µL of RNase-free water (Promega, Madison, WI, USA).
We determined the concentrations of RNA using a Smart-Spec plus spectrophotometer (Bio-
Rad). Then, using a High-Capacity cDNA Reverse Transcription Kit (Cat#: 4368814, Applied
Biosystem, Thermo Fisher Scientific), cDNA was generated from 0.5-1 µg of total RNA in a total
volume of 25 µL. The RT-PCR program was run for 5 minutes at 25 °C, 30 minutes at 42 °C, and
30 minutes at 95 °C. The cDNA templates were further diluted in water two-three times. For qPCR,
two microliters of diluted templates were utilized. A 96-well PCR plate (Cat#: MLL9601, Bio-Rad)
was used for the qPCR. All genes were amplified twice, and each sample was analyzed in
triplicate using 2X SYBR Green PCR mix (Cat#: B21202, Bimake.com) (Supplementary Table 6).
After normalization to *Gapdh*, the CT value was utilized to compute the fold change in RNA
abundance. qPCR reactions were performed using the Aria Mx Real-Time PCR equipment
(Agilent).

Flow cytometry and cytokine analysis

Single-cell suspensions were prepared from mouse spleen. To assess $CD4^+$ T cell activation, we
stained single-cell suspensions cell surface staining in 50µL FACS buffer (1% BSA, 2 mM EDTA,
and 0.1% sodium azide in PBS, pH 7.4) (5-10x10^5 cells/tube) and incubated with antibodies for
30 minutes at 37°C. (106 cells/50 mL) with anti-CD45 PercpCy5.5 (Cat# 103132, Clone#30F11,
BioLegend), anti-CD4 PE (Cat#100408, Clone#GK3.5, BioLegend), anti-CD44 Alexflour700
(Cat#103026,IM7, BioLegend), and anti-CD62L PeCy7 (Cat#104418, Clone# MEL/4, BioLegend)
on ice for 30 minutes (SupplementaryTable 6). Stained cells were then washed twice with 500µL
of ice-cold FACS buffer to eliminate binding antibodies before being analyzed immediately using
a BD FACSymphony A3Flow Cytometer. Annexin V APC staining cells were incubated with
Annexin V APC (Cat#640920, BioLegend) for 3 minutes in the ice-cold FACS buffer and then
washed. Followed by Propidium Iodide (Cat#6421301, BioLegend) added right before analyzing
the cells by FACS. MitoSpy (Cat# 424801, BioLegend) staining was carried out in accordance
with the manufacturer's recommendations. Cells were resuspended in MitoSpy in 300-400 µL of
warm 1X PBS and incubated at 37 °C for 30 minutes. Cells were then washed twice with 1X PBS
before being analyzed using a BD FACSymphony A3 Flow Cytometer on stained and unstained
samples (BD Biosciences). For inflammatory cytokines detection, sera were collected from mice
blood and measured by the LEGENDplex mouse inflammation panel according to the
manufacturer's instructions (Cat#740446, BioLegend). Samples were read on the BD
FACSymphony Flow Cytometer.
663

***In vitro* Treg suppression assay**
Tregs ($CD4^+CD25^+$) cells were isolated from the spleen of sex-matched TFAM cKO and TFAM$^{f/f}$
(control) mice and Tconv ($CD4^+CD25^-$) from TFAM control mice using BD FACS ARIA^{TM}Ilu at the
UND Flow Cytometry Core: anti-CD4 PE (Cat#100408, Clone#GK3.5, BioLegend), anti-CD25
(Cat#102030, clone PC61, BioLegend). Afterward, 5x10^4 Tconv ($CD4^+CD25^-$) cells from TFAM
control were labeled with CFSE Cell Division Tracker Kit (Cat# 423801, BioLegend) and cultured
with mitomycin-treated wild type splenocytes. To detect Treg suppressive effect on these labeled
Tconv ($CD4^+CD25^-$) cells culture, we added Tregs ($CD4^+CD25^+$) from the TFAM cKO and TFAM$^{f/f}$
(control) for 60 hours at 37°C with 2.5 mg/mL Dynabeads^{TM} Mouse T-Activator CD3/CD28 for T-
Cell Expansion and Activation bead (Cat# 1145). After 60 hours, the CD4 cell proliferation was
observed by FACS. Suppressive capacity was calculated based on the relative proportion of
proliferating Tconv cells compared with cultures lacking Tregs and is presented as percent
suppression, as our previously described study[69].
677

Cell isolation and sequencing library construction using the 10x Genomics Chromium platform

Cells were isolated from spleens from WT and cKO mice by grinding with 70 μm cell strainers (Cat#22363548, Fisher brand). ACK RBC lysis buffer (Cat#A10492-01, Gibco) was used to lyse RBC. Trypan blue (Cat#25900Cl, Corning) staining confirmed cell viability. The 10X Genomics Chromium system was used to create 3' single-cell gene expression libraries (v3.0) from WT and cKO spleen single-cell suspensions. On an Illumina HiSeq X Ten, pair-end 150 bp sequencing was used to generate high-quantity data (Illumina, San Diego, California, USA). Then, using Cell Ranger software v6.0.0, the output Illumina base call files from sequencing runs were combined and converted to FASTQ files using Illumina's bcl2fastq.

Single-cell RNA-seq data alignment and sample aggregating

RNA reads from the 10X Chromium platform with a sequence quality of less than 30 were filtered out and then mapped to the mm10 mouse reference genome using the CellRanger software (v6.0.0). Seurat (v4.0.2) was used to read the individual sample output files from CellRanger Count[70]. Cells were further filtered out of the downstream analysis using the following criteria: *nFeature RNA <200 & nFeature RNA >4000 & percent.mt >25*. Scrublet (v0.2.3)[71] was used to detect probable doublets and then *CellCycleScoring in Seurat was used* to define the cell cycle phase by using a published list of cell cycle genes[72]. *SCTransform* (v0.3.2) from Seurat was used to independently normalize the gene-cell count matrices with default parameters[73]. Dimension reductions, including principal component analysis (PCA), clustering, and t-distributed Stochastic Neighbor Embedding (t-SNE) projections (resolution =0.7) were then performed on the normalized data. By following a similar pipeline, subclusters for CD4$^+$ T cells were identified.

Cell type annotation and differential expression analysis

To determine the cell type of each cluster, the *FindAllMarkers* function in Seurat was used to find the cluster's gene markers. Cell types were allocated manually after consulting the CellMarker[74] and PanglaoDB[75] databases. MAST (v1.18.0)[76] was utilized to conduct differential analysis on each cell type to find the genes that were differentially expressed between WT and cKO cells with the adjusted p-value <0.05 and absolute log2FC >0.25. The Gene Set Enrichment Analysis (GSEA) with the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway annotation was done using the richR package (https://github.com/hurlab/richR), with a p-value of 0.01 used as the cutoff value to identify significant KEGG pathways.

Single-cell trajectory reconstruction for CD4⁺ T cell

The Monocle2 R package (v2.20.0, http://cole-trapnell-lab.github.io/monocle-release/) was used to reconstruct cell fate choices and pseudotime trajectories. The gene-cell matrix of UMI counts was sent to Monocle, which then created an object using its new Cell Data Set function (*lower Detection Limit = 0.5, expression Family = nonbinomial.size*). We evaluated the size factors and dispersion of gene expression. We next used the differentialGeneTest module to conduct differential expression analysis in order to determine the ordering of genes using the parameter "*fullModelFormulaStr = ~celltype*," and then maintained genes with q values less than 1E-3 for downstream analysis. Following the *setOrderingFilter* function, the *reduceDimension* module with the argument "*method = DDRTree*" was used to reduce the dimensionality of the data. When *orderCells* is executed, the root state variable may be adjusted to reflect the known biological background. Following that, the *plot_cell_trajectory* and *plot_pseudotime_heatmap* functions were used to visualize the data further.

RNA velocity analysis

Velocyto (v0.17.17) was used to analyze the velocity of RNA in cells[77]. This method utilizes the relative amount of unspliced and spliced mRNA as a predictor of future cell state. Annotation of spliced and unspliced reads was conducted using the Python script velocyto.py in the Cell Ranger output folder utilizing the Velocyto pipeline. Then, the scvelo (v0.2.3)[78] was used to determine the CD4⁺ T cell velocity. The previously normalized data served as the basis for calculating first and then order moments for each cell in relation to its closest neighbors (*scvelo.pp.moments(n pcs = 30, n neighbors = 30)*). Following that, the velocities were calculated, and the velocity graph was produced using the *scvelo.tl.velocity* and *scvelo.tl.velocity* graph functions with the mode set to 'stochastic'. Finally, using the *scvelo.tl.velocity* embedding function, the velocity field was projected onto the existing t-SNE coordinates.

Transcription factor regulons prediction

Gene regulatory network (GRN) was generated using pySCENIC package (v0.11.0)[79]. Briefly, the raw expression matrix for the cells of all samples was filtered by keeping genes with a total expression greater than 3*0.01*(number of cells). Then the *pyscenic.grn* command was used with *GRNboost2* method and default options and a fixed seed to derive co-expression modules between transcription factors and potential targets. The following database: cisTarget databases (*mm10__refseq-r80__10kb_up_and_down_tss.mc9nr.feather, mm10__refseq-r80__500bp_up_and_100bp_down_tss.mc9nr.feather*), the transcription factor motif annotation

database (*motifs-v9-nr.mgi-m0.001-o0.0.tbl*) and the list of mouse transcription factors
(*mm_mgi_tfs.txt*) were downloaded for the analysis. A regulon is a group of target genes regulated
by a common transcription factor. AUCell (Area Under the Curve) scores (regulon activities) in
each cell were computed with *pycenic.aucell* command (default parameters). These scores were
then binarized into either "on" or "off" states, by setting an optimized threshold on the distribution
of each regulon among all cells using the *skimage.filters.threshold_minimum* function.
752

Bulk RNA-seq data processing
Low quality (Q<30) reads, and sequencing adapters were removed with Trimmomatic software
v0.36[80] from raw reads. Clean reads were then mapped to the mouse reference genome (mm10)
by using HISAT2[81]. FeatureCounts[82] were used to summarize the unique mapped reads to mouse
genes. Differential gene expression analysis was performed using DESeq2[83] with adjusted p-
value <0.05 and fold change >2 as the cutoff for identifying differentially expressed genes (DEGs).
Gene Set Enrichment Analysis (GSEA) on KEGG pathway data was performed by using the richR
package and the adjusted p-value <0.05 was chosen as the cutoff value to select significant
KEGG pathways.
762

Microbiome analysis
Microbial DNA samples were collected as our vius published protocol[33]. Samples were sequenced
using Oxford Nanopore Technology. Raw sequencing data underwent quality control using
Porechop (v1.0.0) to remove adapter sequences, followed by length filtering with NanoFilt to
retain sequences between 1,200 and 1,800 base pairs. Quality metrics were assessed using
NanoStat. Chimeric reads were identified and removed through an all-versus-all alignment
approach using minimap2 (v2.17) with Oxford Nanopore-specific parameters (-x ava-ont -g 500),
followed by filtering with yacrd (coverage threshold 4, noise threshold 0.4). Cleaned reads were
mapped against a reference database using minimap2 (-x map-ont), and high-confidence
alignments were retained using custom filtering scripts from the Spaghetti pipeline. OTU tables
and taxonomy assignments were generated for downstream analysis.
774

775   Microbiome analysis was performed in R (version ≥4.0.0) using the phyloseq package. Alpha
776   diversity (Observed richness, Simpson, and Shannon indices) was calculated at the genus level.
777   Beta diversity was assessed using Bray-Curtis dissimilarity and visualized with Principal
778   Coordinates Analysis (PCoA). Differential abundance analysis between experimental groups
779   (P8M, P8W, PT, and PY) was conducted using ALDEx2 with 128 Monte Carlo instances. To

identify significantly differentially abundant bacterial taxa between groups, we used the ALDEx2 package[84] . This approach was chosen to properly handle the compositional nature of microbiome data. Four comparisons were performed: P8M vs P8W, PT vs PY, P8W vs PY, and PT vs P8M. For each comparison, Monte Carlo sampling with 128 Dirichlet instances was used to account for technical variation (mc.samples = 128), with all features included in the geometric mean calculation (denom = "all").Bacterial taxa were considered significantly differentially abundant if they met both thresholds: p-value < 0.05 (Welch's t-test) and effect size > 0.8. The effect size threshold represents cases where the between-group difference exceeds 80% of the maximum within-group dispersion[85] . Pathway functionality was predicted from the taxonomic data using PICRUSt2 (Phylogenetic Investigation of Communities by Reconstruction of Unobserved States), allowing for inference of KEGG pathways present in the microbial communities based on marker gene sequencing data. Significant pathways (p <0.05) were visualized using heatmaps.

Antibiotic Treatment and Fecal Microbiota Transplantation (FMT)

To deplete endogenous microbiota, mice were administered a broad-spectrum antibiotic cocktail in drinking water for two weeks: ampicillin (10 mg/mL, A1593, MilliporeSigma), metronidazole (10 mg/mL, PHR1052, MilliporeSigma), and vancomycin hydrochloride (5 mg/mL, SBR00001, MilliporeSigma). Fecal material for transplantation was collected from donor mice aged 8 weeks (young) or 2 years (aged), snap-frozen, and stored at -80°C. Following antibiotic treatment, recipient C57BL/6 mice were housed in sterile, autoclaved cages and orally gavaged with 300– 400 µL of freshly prepared fecal suspension from untreated donor mice. The suspension was generated by homogenizing fecal pellets in sterile water. Control groups received either sterile water or autologous fecal homogenates.

Grip strength analysis

Neuromuscular strength was assessed using the grip strength test. The forelimb and hindlimb grip strengths of mice were assessed by using a grip strength meter from San Diego Instruments. Each mouse was tested in three trials, and the force was recorded as grams-force (gf, 1 gf = 9.8 x 10^{-3} Newtons).

Open field test

Locomotor activity was measured in the open field test. A mouse was placed in the empty square cage from San Diego Instruments. Total distance (m) covered during a 10-min period was recorded and calculated by ANY-maze software (version 7.1).

814   **Data availability**
815   Datasets generated in this study by RNA-seq and scRNA-seq are accessible at GSE203143
816   (Reviewer access token: ozebasggjbyrpsf) and GSE197973 (Reviewer access token:
817   clajaakoztkhtyh), respectively. All data associated with this study are present in the paper or the
818   Supplementary Materials.
819
820   **Code availability**
821   All data analysis scripts are available on the GitHub (https://github.com/guokai8/TFAM).
822

Figure Legends
Figure 1. Treg-intrinsic TFAM loss reprograms CD4$^+$ T cell lineage fate toward pro-
inflammatory and terminal effector states.
(**A**) Schematic representation of the experimental workflow. Splenocytes were isolated from
Foxp3^{Cre}Tfam$^{f/f}$ (TFAM cKO) and Tfam$^{f/f}$ (WT) mice for single-cell RNA sequencing (scRNA-seq)
and downstream analysis.
(**B**) UMAP projection of 49,156 immune cells (WT: 34,958; cKO: 14,198), identifying 21 major
immune subsets across 26 transcriptional clusters. Erythroid lineage cells were excluded.
(**C**) UMAP projection of CD4$^+$ T cells (WT: 4,324; cKO: 3,497), revealing six transcriptionally
distinct subsets: naïve, effector memory (TEM), exhausted, natural Tregs (nTregs),
aged/dysfunctional Tregs (aTregs), and cytotoxic CD4$^+$ T cells.
(**D**) Track plot showing representative gene expression patterns across single CD4$^+$ T cells. Rows
represent genes; columns represent individual cells.
(**E**) Bar graph quantifying relative abundance of CD4$^+$ T cell subsets in WT and cKO mice. TFAM
deficiency promotes expansion of effector-like states (aTregs, cytotoxic, and TEM), with reduced
naïve and exhausted populations.
(**F**) Gene set enrichment analysis (GSEA) of KEGG pathways shows increased TNFα signaling,
cytokine/chemokine responses, and calcium flux in CD4$^+$ T cells from TFAM cKO mice.
(**G**) Pseudotime trajectory analysis (Monocle) of CD4$^+$ T cells reveal two dominant differentiation
paths (P1 and P2). TFAM-deficient cells preferentially adopt the alternative P2 trajectory, enriched
in transcriptional States 1 and 2.
(**H**) Monocle 2 trajectory cell states for all CD4$^+$ T cells, each branch represents one cell state
(left). Barplots representing cell proportions in the seven states of WT and cKO (right). Cells are
colored according to seven states, which partition the trajectory.

(**I**)**.** Relative proportion of CD4$^+$ T cells among different states of WT and cKO spleen, color stands

for cell type.

(**J**) RNA velocity analysis confirms directionality of lineage bifurcation, with streamlines overlaid

on UMAP embedding indicating transcriptional flow.

(**K**) Heatmap of four gene clusters ordered by pseudotime. Cluster 1 genes (e.g., Klf3, Bcl2) mark

early developmental states; Cluster 2 genes (e.g., Ccl5, S100a6) are linked to terminal effector

function and inflammation. KEGG pathway enrichment of cluster-specific genes is shown.

(**L**) Representative flow cytometry plots of CD4$^+$CD44^{hi}CD62L^{lo} effector and CD4$^+$CD44^{lo}CD62L^{hi}

naïve T cells in spleen and lymph nodes from WT and TFAM cKO mice.

(**M**) Quantification of mean fluorescence intensity (MFI) for effector and naïve CD4$^+$ T cells across

organs. TFAM cKO mice exhibit increased effector T cell signatures.

(**N**) Suppression assay showing reduced regulatory function of TFAM-deficient Tregs in vitro. Co-

culture with CFSE-labeled CD4$^+$ Tconv cells demonstrates impaired suppression of proliferation

upon CD3/CD28 stimulation.

Figure 2. Treg-intrinsic TFAM loss triggers a multi-lineage stress response, inflammatory

reprogramming, and systemic senescence.

(**A**) Gene regulatory network (GRN) inference from scRNA-seq data identified 177 regulons

across CD4$^+$ T cell subsets. TFAM cKO Tregs showed increased inflammatory activity (Nfkb2,

Nfatc1, Rora, Runx2), stress-responsive (Egr1, Junb, Xbp1), and cytotoxic (Runx3) regulons,

alongside pan-CD4$^+$ expression of the mitochondrial stress gene Ddit3.

(**B**) Dot plots showing upregulation of key inflammatory transcriptional regulators (Prdm1, Nfkb1,

Nfkb2, Ahr) in TFAM-deficient Tregs.

(**C**) Gene set enrichment analysis (GSEA) shows significant pathways in Monocytes (Mono),

Macrophages (Mac), dendritic cell (DC), Immature Dendritic Cells (iDC) and Plasmacytoid

Dendritic Cells (pDC) cells in TFAM cKO mice.

(**D**) Bulk RNA-seq of cKO spleens revealed enrichment of cell death pathways, including

apoptosis and necroptosis.

(**E**) Immunoblots confirmed caspase-3 and -8 cleavage, IκBα degradation, and phosphorylation

of RIPK1, RIPK3, and MLKL, indicative of RIPK-mediated immunogenic cell death.

(**F**) Histological analysis revealed multi-organ inflammatory infiltration in colon, liver, lung, and

kidney of TFAM cKO mice.

(**G**) H&E staining of femurs showed pronounced cortical bone thinning in TFAM cKO animals.

(**H**) Spleens from TFAM cKO mice exhibited splenomegaly, follicular disruption, and marginal
zone disorganization.
(**I**) Mitospy staining in sorted Tregs revealed mitochondrial depolarization and elevated oxidative
stress in TFAM-deficient cells.
(**J**) Elevated circulating levels of TNF-α, IL-6, and IFNγ in serum from TFAM cKO mice, confirming
systemic inflammation.
(**K**) RT-qPCR showed increased expression of Cdkn2a (p16) and Cdkn1a (p21);
(**L**) Immunoblotting of splenic lysates confirmed elevated levels of p53, CDK4, PCNA, p16, and
p21, consistent with senescence-associated gene expression.
(**M**) Whole-spleen RNA-seq revealed enrichment of aging- and inflammation-associated gene
networks.
(**N**) GSEA showed significant upregulation of pathways linked to neuroactive ligand–receptor
signaling, and downregulation of circadian rhythm disruption, and protein trafficking which are
hallmarks of systemic immune aging.
894
Figure 3. Partial TFAM reduction in Tregs promotes systemic aging features, peripheral
inflammation, and early neurodegenerative changes.
(**A**) Human PBMC transcriptome analysis reveals age-associated decline in TFAM expression,
supporting its role as a mitochondrial-immune rheostat.
(**B-C**) TFAM Het (Foxp3^{Cre}Tfam$^{f/+}$) mice develop aging phenotypes, including kyphosis and
reduced body weight, by 8-10 months of age.
(**D**) Circulating TNF-α and IL-6 levels are elevated in TFAM Het mice, indicating persistent low-
grade systemic inflammation.
(**E**) Splenic CD4$^+$ T cells isolated from TFAM Het mice display elevated expression of
senescence-associated markers, including Cdkn2a (*p16^Ink4a*), Cdkn1a (*p21^Cip1*), p53, and
CDK4, with GAPDH serving as the loading control.
(**F**) Neuromuscular assessment shows reduced forelimb grip strength in TFAM Het mice.
(**G**) Open field behavioral assay demonstrates decreased exploratory behavior and locomotor
hypoactivity, consistent with CNS dysfunction.
(**H–I**) Cresyl violet staining reveals neuronal loss in the hippocampus and cortex of 8-month-old
TFAM Het mice compared to 8-week controls.
(**J**) Longitudinal hippocampal RNA-seq profiling shows minimal transcriptional changes at 8
weeks but extensive neuroimmune remodeling by 8 months.

(**K**) Heatmap displaying upregulation of neuroinflammation-associated genes (Gabra5, Slc38a5,
Calcr, Ganbare) in 8-month TFAM Het hippocampus.

(**L**) KEGG pathway enrichment of 8-month hippocampal transcriptomes identify significant
enriched pathways of oxidative stress, inflammatory signaling, and neurodegeneration-linked
pathways.

Figure 4. TFAM insufficiency in Tregs triggers CNS immune activation and microglial
sensitization to TNF signaling.

(**A-B**) Iba1 and NeuN immunohistochemistry shows widespread activation of amoeboid microglia
in the cortex and hippocampus of 8-month-old TFAM Het mice, with concurrent neuronal loss.

(**C**) Flow cytometric analysis reveals increased MHC-I expression on brain-resident CX3CR1$^+$
microglia from TFAM Het mice, consistent with enhanced antigen presentation.

(**D**) Ex vivo stimulation with TNF-α induces elevated *Il1b*, *Tnfr1*, and *Il6* expression in CX3CR1$^+$
microglia from TFAM Het brains, indicating heightened inflammatory sensitivity.

Figure 5. TFAM insufficiency in Tregs drives microbiota–lipid dysregulation and
neuroinflammatory reprogramming via the gut–brain–immune axis.

(**A**) Longitudinal 16S rRNA sequencing reveals significantly elevated α-diversity (Shannon and
Simpson indices) in TFAM Het mice, peaking at 8 months of age.

(**B**) β-diversity analysis (Bray–Curtis dissimilarity) demonstrates genotype- and age-specific
microbial clustering, indicating progressive community remodeling.

(**C**) Phylum-level taxonomic profiling reveals expansion of *Campilobacterota* and depletion of
Bacteroidota in aged TFAM Het mice, signatures linked to inflammaging.

(**D**) VennDiagram depicting distribution and overlap of microbial species across genotypes and
age groups.

(**E**) Differential abundance analysis identifies 18 significantly altered taxa, including loss of key
SCFA-producing genera (*Lactobacillus murinus*, *Odoribacter*, *Prevotellaceae_UCG-001*) in
TFAM Het mice.

(**F**) KEGG-based predictive metagenomics shows TFAM Het microbiota enriched in age-related
pathways including FoxO, and spinocerebellar ataxia signaling.

(**G**) Lipid metabolism pathways, including unsaturated fatty acid biosynthesis, bile acid
biosynthesis, and sphingolipid metabolism, are significantly altered. Neurotransmission pathways
(GABAergic, glutamatergic synapses) are also enriched in aged TFAM Het mice.

(**H**) Microbiota–host gene co-correlation network analysis reveals 323 significant interactions. Notably, *Helicobacter* positively correlates with *Ntsr2*, while Lactobacillus inversely correlates with neuroendocrine (e.g., *Pomc*, *Sstr5*) and developmental genes (*Elk3*, *Sox4*) and positively correlates with inflammatory genes (*Mpo*, *Ifitm6*). *Lachnospiraceae_NK4A136_group* links to complement-mediated inflammation (*C5ar2*).

951 (**I**) RT-qPCR validation confirms elevated expression of *Il1b* and *Tnfr1* in peripheral tissues from TFAM Het mice, supporting a chronic inflammatory state.

Figure 6. Restoration of mitochondrial fitness mitigates inflammation and functional decline in models of immune aging.

(**A**) Schematic of therapeutic intervention in TFAM cKO mice. Mice received intraperitoneal resveratrol (RSV; 7 mg/kg) every other day for 4 weeks.

(**B**) Mitospy staining in skin tissue reveals enhanced mitochondrial membrane potential in RSV-treated TFAM cKO mice, indicating improved mitochondrial integrity.

(**C**) Serum levels of TNF-α and IL-6 were significantly reduced following RSV treatment, reflecting dampened systemic inflammation.

(**D-F**) Quantitative RT-PCR analysis of splenocytes shows reduced expression of senescence markers Cdkn2a (p16^{Ink4a}) and Cdkn1a (p21^{Cip1}) in RSV-treated mice, consistent with suppression of the senescence-associated secretory phenotype (SASP).

(**G**) Histological examination of lung, liver, and kidney shows decreased inflammatory infiltration and preserved tissue structure in RSV-treated animals.

(**H**) Experimental design for fecal microbiota transplantation (FMT) in TFAM Het mice. Mice were antibiotic-conditioned for 10 days and subsequently gavage feces from TFAM-sufficient donors over 3 days, followed by a 3-week recovery.

(**I**) FMT improved neuromuscular performance, demonstrated by increased rotarod endurance and forelimb grip strength. These improvements were associated with re-expression of Treg-associated genes and suppression of pro-inflammatory programs.

(**J**) Working model summarizing the role of TFAM as a central regulator of mitochondrial redox balance, microbiota composition, and immune aging. Restoration of mitochondrial integrity via RSV and microbial reprogramming via FMT ameliorates immunopathology and rejuvenates immune function.

Supplementary Fig. 1: Landscape of scRNA-seq analysis of WT and cKO spleen.

A. t-SNE projections of single-cell transcriptomes from WT and cKO mice before Erythroid cells were removed, annotated by cell type. Each dot denotes a distinct cell. **B.** Cell subset proportions (top) and numbers (bottom) for WT and cKO spleens. **C.** Dot plot of mean expression of canonical marker genes for 20 major lineages from tissues of each origin, as indicated. **D.** Gene Set Enrichment Analysis (GSEA) of KEGG pathways for all cell types between cKO and WT (Only age-related pathways are selected). Color stands for the up-regulated (red) or down-regulated (blue) in cKO spleen. Enriched terms were identified as significant at a p-value ≤ 0.01. **E.** Expression of aged-related genes (*Fcer2a*, *Cd79b*, *Zfp36l2*, and *S100a6*) on Figure 1B t-SNE embedding. Each dot corresponds to a single cell.

Supplementary Fig. 2: scRNA-seq results show the cellular heterogeneity in CD4$^+$ T cells.
A. Relative proportion of CD4$^+$ T cells among WT and cKO spleen, color stands for cell type. **B.** Violin plot showing the expression profile of age-related genes among all CD4$^+$ T cells. **C.** Monocle 2 prediction of CD4$^+$ T cell subsets developmental trajectory with pseudotime. Each dot represents an individual cell. **D.** Functional enrichment analysis of KEGG pathways with up-regulated differential expressed genes for the cell states. Only the top 20 of the enriched pathways were plotted from each state.

Fig. 3: t-SNE projection of binary regulon activity. t-SNE plots visualize the binarized regulon activity states of Irf8, Runx2, Bcl11a, and Nfkb1 across single cells, highlighting distinct transcriptional regulatory modules and cell-type-specific activation patterns.

Supplementary Fig. 4: TFAM deletion in Foxp3$^+$ Tregs induces systemic pathology and accelerates immune aging in Foxp3^{Cre}Tfam$^{f/f}$ mice. A. Representative photograph showing the deteriorated physical appearance of a cKO mouse (right) compared to an age- and sex-matched WT littermate (left) at 8 weeks of age. **B.** X-ray image depicting increased spinal curvature in the cKO mouse (right) relative to a WT littermate (left) of the same age and sex. **C.** Kaplan-Meier survival analysis comparing WT (n=10; M/F) and cKO (n=10; M/F) mice, revealing reduced lifespan in cKO animals. **E.** Flow cytometric analysis of Annexin V staining in spleen and skin cells from WT and cKO mice, indicating increased cell death in cKO tissues. **F**. Mitospy staining of splenocytes from WT and cKO mice, assessed by flow cytometry and presented as a fluorescence intensity histogram, showing mitochondrial alterations in cKO cells.

Supplementary Fig. 5: Increased cell infiltration and inflammation in different organs in Foxp3^CreTfam^f/f mice. A. Histology sections stained with H&E from lung, liver, colon, and kidney. **B.** Serum cytokines identified by Legend plex for IL-1β, IL-10, MCP1, IL-23, IL-1α, IL-17, IFNγ, IL-12, IFNβ, GM-CSF, and IL-27 with estimated values displayed. Data represented as mean ± SD. Statistical test used: t-test, WT (n=5-10; M/F) and cKO mice (n=5-10, M/F).

Supplementary Fig. 6: Restoration of mitochondrial function leads to mitigate pathologies. A-B. Body weight (**C**) and fold change (**D**) of young and old WT and cKO mice. Data represented as mean ± SD. Statistical test used: two-way ANOVA with post-hoc test (C). WT (n=5; M/F) and Het mice (n=5, M/F).

Supplementary Table 1: Gene Set Enrichment Analysis (GSEA) of KEGG pathway between cKO and WT for all cell types.

Supplementary Table 2: Gene Set Enrichment Analysis (GSEA) of KEGG pathway between cKO and WT for CD4 subsets.

Supplementary Table 3: Functional enrichment analysis of KEGG pathway for four clusters.

Supplementary Table 4: Functional enrichment analysis of KEGG pathway for Monocle states.

Supplementary Table 5: Gene Set Enrichment Analysis (GSEA) of KEGG pathway between cKO and WT from Brain, Spleen and Skin RNA-seq.

Supplementary Table 6: Primer sequences for RT-qPCR, antibodies for Western blots and flow cytometry, and other reagents.

Acknowledgments

We thank Kristian Herman, Reet Goyal, Risha Mathur, Leia Laurer, Sofie Robinson, and Harpreet Singh for assistance with cell culture, mouse experiments, and general laboratory support. Technical support was provided by Dr. Bony De Kumar and Damien Parrello (Genomics Core), Steve Anderson and Subha Nookala (Flow Cytometry Core), Neeta Adhikari and Donna Laturnus (Histology Core), Sarah Abrahamson (Imaging Core), Ellen Olson and Dr. Collin Combs (Behavior Core), and Dr. Brett McGregor (Computational Data Analysis Core) at the University of North Dakota.

Fundings

Research reported in this publication was supported by NIH Host–Pathogen Interactions COBRE Pilot Grant (NIGMS/NIH P20GM1034442 to X.W. and R.M.) and the DaCCOTA Pilot Grant (NIGMS/NIH U54GM128729 to R.M. and D.A.J.). TRANCENDS Pilot Grant NIGMS/NIH P20GM155890 to RM. E.A. and J.G. were supported by the Eva Gilbertson, M.D. Foundation Study Grant to J.D. Core facilities were supported by NIH/NIGMS awards P20GM113123 and U54GM128729. The graphical abstract was created using Servier Medical Art (https://smart.servier.com) under a Creative Commons Attribution 3.0 License.

Contributions

R.M. conceived and designed the study. K.G., R.M., J.G., and Z.W. analyzed and interpreted the scRNA-seq and RNA-seq data. T.D., Z.W., J.G., B.B., H.M., A.L., R.P., M.R., and S.J. performed the experiments. R.M., K.G., J.G., and Z.W. prepared the figures and wrote the manuscript. R.M., J. G., H.C., K.G., N.K., J.H., Z.W., D.J., H.B., M.G., and A.T. critically reviewed and revised the manuscript. All authors contributed to discussions and approved the final version.

Competing interests

The authors declare no competing interests.

References

1. Goronzy, J.J. & Weyand, C.M. Mechanisms underlying T cell ageing. *Nat Rev Immunol* **19**, 573-583 (2019).

2. Nikolich-Zugich, J. The twilight of immunity: emerging concepts in aging of the immune system. *Nat Immunol* **19**, 10-19 (2018).

3. Shaw, A.C., Goldstein, D.R. & Montgomery, R.R. Age-dependent dysregulation of innate immunity. *Nat Rev Immunol* **13**, 875-887 (2013).

4. Franceschi, C. *et al.* Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* **908**, 244-254 (2000).

5. Li, X., Liang, Y., LeBlanc, M., Benner, C. & Zheng, Y. Function of a Foxp3 cis-element in protecting regulatory T cell identity. *Cell* **158**, 734-748 (2014).

6. Olusanya, B.O. Surrogate measurements for identifying newborns with low birth weight in a community with predominant non-hospital births. *J Child Health Care* **14**, 386-395 (2010).

7. Josefowicz, S.Z., Lu, L.F. & Rudensky, A.Y. Regulatory T cells: mechanisms of differentiation and function. *Annu Rev Immunol* **30**, 531-564 (2012).

8. Sakaguchi, S. *et al.* Regulatory T Cells and Human Disease. *Annu Rev Immunol* **38**, 541-566 (2020).

9. Angelin, A. *et al.* Foxp3 Reprograms T Cell Metabolism to Function in Low-Glucose, High-Lactate Environments. *Cell Metab* **25**, 1282-1293 e1287 (2017).

10. Cascone, T. *et al.* Increased Tumor Glycolysis Characterizes Immune Resistance to Adoptive T Cell Therapy. *Cell Metab* **27**, 977-987 e974 (2018).

11. Klarquist, J. *et al.* Clonal expansion of vaccine-elicited T cells is independent of aerobic glycolysis. *Sci Immunol* **3** (2018).

12. Fu, Z. *et al.* Requirement of Mitochondrial Transcription Factor A in Tissue-Resident Regulatory T Cell Maintenance and Function. *Cell Rep* **28**, 159-171 e154 (2019).

13.   Zheng, Y. *et al.* Role of conserved non-coding DNA elements in the Foxp3 gene in regulatory T-cell fate. *Nature* **463**, 808-812 (2010).

14.   Grinberg-Bleyer, Y. *et al.* NF-kappaB c-Rel Is Crucial for the Regulatory T Cell Immune Checkpoint in Cancer. *Cell* **170**, 1096-1108 e1013 (2017).

15.   Oh, H. *et al.* An NF-kappaB Transcription-Factor-Dependent Lineage-Specific Transcriptional Program Promotes Regulatory T Cell Identity and Function. *Immunity* **47**, 450-465 e455 (2017).

16.   Chapman, N.M. *et al.* mTOR coordinates transcriptional programs and mitochondrial metabolism of activated T(reg) subsets to protect tissue homeostasis. *Nat Commun* **9**, 2095 (2018).

17.   Desdin-Mico, G. *et al.* T cells with dysfunctional mitochondria induce multimorbidity and premature senescence. *Science* **368**, 1371-1376 (2020).

18.   Gill, J.S. *et al.* Mitochondrial Oxidative Stress Regulates FOXP3+ T-Cell Activity and CD4-Mediated Inflammation in Older Adults with Frailty. *Int J Mol Sci* **25** (2024).

19.   Lahl, K. & Sparwasser, T. In vivo depletion of FoxP3+ Tregs using the DEREG mouse model. *Methods Mol Biol* **707**, 157-172 (2011).

20.   Hanakawa, S., Chow, Z. & Kabashima, K. Depletion of Treg by the Diphtheria Toxin System. *Methods Mol Biol* **2559**, 31-39 (2023).

21.   Brunkow, M.E. *et al.* Disruption of a new forkhead/winged-helix protein, scurfin, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* **27**, 68-73 (2001).

22.   Weinberg, S.E. *et al.* Mitochondrial complex III is essential for suppressive function of regulatory T cells. *Nature* **565**, 495-499 (2019).

23.   Soto-Heredero, G., Desdin-Mico, G. & Mittelbrunn, M. Mitochondrial dysfunction defines T cell exhaustion. *Cell Metab* **33**, 470-472 (2021).

24.   Larsson, N.G. *et al.* Mitochondrial transcription factor A is necessary for mtDNA maintenance and embryogenesis in mice. *Nat Genet* **18**, 231-236 (1998).

25.   West, A.P. & Shadel, G.S. Mitochondrial DNA in innate immune responses and inflammatory pathology. *Nat Rev Immunol* **17**, 363-375 (2017).

1143

1144   26.   Kuroda, R. *et al.* Loss of mitochondrial transcription factor A in neural stem cells leads to
1145   immature brain development and triggers the activation of the integral stress response in
1146   vivo. *PLoS One* **16**, e0255355 (2021).

1147

1148   27.   Lu, T. *et al.* TFAM deficiency in dendritic cells leads to mitochondrial dysfunction and
1149   enhanced antitumor immunity through cGAS-STING pathway. *J Immunother Cancer* **11**
1150   (2023).

1151

1152   28.   West, A.P. *et al.* Mitochondrial DNA stress primes the antiviral innate immune response.
1153   *Nature* **520**, 553-557 (2015).

1154

1155   29.   Elyahu, Y. *et al.* Aging promotes reorganization of the CD4 T cell landscape toward
1156   extreme regulatory and effector phenotypes. *Sci Adv* **5**, eaaw8330 (2019).

1157

1158   30.   Furman, D. *et al.* Chronic inflammation in the etiology of disease across the life span. *Nat
1159   Med* **25**, 1822-1832 (2019).

1160

1161   31.   Garg, S.K. *et al.* Aging is associated with increased regulatory T-cell function. *Aging Cell*
1162   **13**, 441-448 (2014).

1163

1164   32.   Amorim, J.A. *et al.* Mitochondrial and metabolic dysfunction in ageing and age-related
1165   diseases. *Nat Rev Endocrinol* **18**, 243-258 (2022).

1166

1167   33.   Wang, Z. *et al.* Age-related dysregulation of intestinal epithelium fucosylation is linked to
1168   an increased risk of colon cancer. *JCI Insight* **9** (2024).

1169

1170   34.   Gomez de Las Heras, M.M. *et al.* CD4 T cell therapy counteracts inflammaging and
1171   senescence by preserving gut barrier integrity. *Sci Immunol* **10**, eadv0985 (2025).

1172

1173   35.   Claesson, M.J. *et al.* Gut microbiota composition correlates with diet and health in the
1174   elderly. *Nature* **488**, 178-184 (2012).

1175

1176   36.   Smith, P.M. *et al.* The microbial metabolites, short-chain fatty acids, regulate colonic Treg
1177   cell homeostasis. *Science* **341**, 569-573 (2013).

1178

1179   37.   Thevaranjan, N. *et al.* Age-Associated Microbial Dysbiosis Promotes Intestinal
1180   Permeability, Systemic Inflammation, and Macrophage Dysfunction. *Cell Host Microbe*
1181   **21**, 455-466 e454 (2017).

1182

1183 38. Sanna, S. *et al.* Causal relationships among the gut microbiome, short-chain fatty acids and
1184   metabolic diseases. *Nat Genet* **51**, 600-605 (2019).

1185
1186 39. Zhou, D. *et al.* Sodium butyrate attenuates high-fat diet-induced steatohepatitis in mice by
1187   improving gut microbiota and gastrointestinal barrier. *World J Gastroenterol* **23**, 60-75
1188   (2017).

1189
1190 40. Stilling, R.M. *et al.* The neuropharmacology of butyrate: The bread and butter of the
1191   microbiota-gut-brain axis? *Neurochem Int* **99**, 110-132 (2016).

1192
1193 41. Furusawa, Y. *et al.* Commensal microbe-derived butyrate induces the differentiation of
1194   colonic regulatory T cells. *Nature* **504**, 446-450 (2013).

1195
1196 42. Louis, P. & Flint, H.J. Formation of propionate and butyrate by the human colonic
1197   microbiota. *Environ Microbiol* **19**, 29-41 (2017).

1198
1199 43. Arpaia, N. *et al.* Metabolites produced by commensal bacteria promote peripheral
1200   regulatory T-cell generation. *Nature* **504**, 451-455 (2013).

1201
1202 44. Fang, P., Kazmi, S.A., Jameson, K.G. & Hsiao, E.Y. The Microbiome as a Modifier of
1203   Neurodegenerative Disease Risk. *Cell Host Microbe* **28**, 201-222 (2020).

1204
1205 45. Wilmanski, T. *et al.* Gut microbiome pattern reflects healthy ageing and predicts survival
1206   in humans. *Nat Metab* **3**, 274-286 (2021).

1207
1208 46. Valdes, A.M., Walter, J., Segal, E. & Spector, T.D. Role of the gut microbiota in nutrition
1209   and health. *BMJ* **361**, k2179 (2018).

1210
1211 47. Johnson, E.L., Heaver, S.L., Walters, W.A. & Ley, R.E. Microbiome and metabolic
1212   disease: revisiting the bacterial phylum Bacteroidetes. *J Mol Med (Berl)* **95**, 1-8 (2017).

1213
1214 48. Wu, S.K., Wang, L., Wang, F. & Zhang, J. Resveratrol improved mitochondrial biogenesis
1215   by activating SIRT1/PGC-1alpha signal pathway in SAP. *Sci Rep* **14**, 26216 (2024).

1216
1217 49. Price, N.L. *et al.* SIRT1 is required for AMPK activation and the beneficial effects of
1218   resveratrol on mitochondrial function. *Cell Metab* **15**, 675-690 (2012).

1219
1220 50. Franceschi, C., Garagnani, P., Parini, P., Giuliani, C. & Santoro, A. Inflammaging: a new
1221   immune-metabolic viewpoint for age-related diseases. *Nat Rev Endocrinol* **14**, 576-590
1222   (2018).

1223

1224 51. Campisi, J. *et al.* From discoveries in ageing research to therapeutics for healthy ageing.
1225   *Nature* **571**, 183-192 (2019).

1226
1227 52. Lopez-Otin, C., Blasco, M.A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of
1228   aging. *Cell* **153**, 1194-1217 (2013).

1229
1230 53. Shin, B. *et al.* Mitochondrial Oxidative Phosphorylation Regulates the Fate Decision
1231   between Pathogenic Th17 and Regulatory T Cells. *Cell Rep* **30**, 1898-1909 e1894 (2020).

1232
1233 54. Liu, X. *et al.* Regulation of mitochondrial biogenesis in erythropoiesis by mTORC1-
1234   mediated protein translation. *Nat Cell Biol* **19**, 626-638 (2017).

1235
1236 55. Viader, A. *et al.* Aberrant Schwann cell lipid metabolism linked to mitochondrial deficits
1237   leads to axon degeneration and neuropathy. *Neuron* **77**, 886-898 (2013).

1238
1239 56. Kim, J. *et al.* Cutting edge: depletion of Foxp3+ cells leads to induction of autoimmunity
1240   by specific ablation of regulatory T cells in genetically targeted mice. *J Immunol* **183**,
1241   7631-7634 (2009).

1242
1243 57. Baker, D.J. *et al.* BubR1 insufficiency causes early onset of aging-associated phenotypes
1244   and infertility in mice. *Nat Genet* **36**, 744-749 (2004).

1245
1246 58. Niedernhofer, L.J. *et al.* A new progeroid syndrome reveals that genotoxic stress
1247   suppresses the somatotroph axis. *Nature* **444**, 1038-1043 (2006).

1248
1249 59. Osorio, F.G. *et al.* Nuclear envelope alterations generate an aging-like epigenetic pattern
1250   in mice deficient in Zmpste24 metalloprotease. *Aging Cell* **9**, 947-957 (2010).

1251
1252 60. Vacca, M. *et al.* The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms*
1253   **8** (2020).

1254
1255 61. Bettedi, L., Yan, A., Schuster, E., Alic, N. & Foukas, L.C. Increased mitochondrial and
1256   lipid metabolism is a conserved effect of Insulin/PI3K pathway downregulation in adipose
1257   tissue. *Sci Rep* **10**, 3418 (2020).

1258
1259 62. Martins, R., Lithgow, G.J. & Link, W. Long live FOXO: unraveling the role of FOXO
1260   proteins in aging and longevity. *Aging Cell* **15**, 196-207 (2016).

1261

1262   63.   Chai, J.N. *et al.* Helicobacter species are potent drivers of colonic T cell responses in
1263   homeostasis and inflammation. *Sci Immunol* **2** (2017).

1264
1265   64.   Lagouge, M. *et al.* Resveratrol improves mitochondrial function and protects against
1266   metabolic disease by activating SIRT1 and PGC-1alpha. *Cell* **127**, 1109-1122 (2006).

1267
1268   65.   Cryan, J.F. *et al.* The Microbiota-Gut-Brain Axis. *Physiol Rev* **99**, 1877-2013 (2019).

1269
1270   66.   O'Toole, P.W. & Jeffery, I.B. Gut microbiota and aging. *Science* **350**, 1214-1215 (2015).

1271
1272   67.   Mathur, R. *et al.* A mouse model of Salmonella typhi infection. *Cell* **151**, 590-602 (2012).

1273
1274   68.   Mathur, R., Alam, M.M., Zhao, X.-F., Huang, Y. & Zhu, X. Mechanistic insight into the
1275   development of TNBS-mediated intestinal fibrosis and evaluating the inhibitory effects of
1276   rapamycin. *JoVE (Journal of Visualized Experiments)*, e60067 (2019).

1277
1278   69.   Park, S.G. *et al.* T regulatory cells maintain intestinal homeostasis by suppressing
1279   gammadelta T cells. *Immunity* **33**, 791-803 (2010).

1280
1281   70.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821
1282   (2019).

1283
1284   71.   Wolock, S.L., Lopez, R. & Klein, A.M. Scrublet: Computational Identification of Cell
1285   Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289 (2019).

1286
1287   72.   Kowalczyk, M.S. *et al.* Single-cell RNA-seq reveals changes in cell cycle and
1288   differentiation programs upon aging of hematopoietic stem cells. *Genome Res* **25**, 1860-
1289   1872 (2015).

1290
1291   73.   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
1292   seq data using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

1293
1294   74.   Zhang, X. *et al.* CellMarker: a manually curated resource of cell markers in human and
1295   mouse. *Nucleic Acids Res* **47**, D721-d728 (2019).

1296
1297   75.   Franzén, O., Gan, L.M. & Björkegren, J.L.M. PanglaoDB: a web server for exploration of
1298   mouse and human single-cell RNA sequencing data. *Database (Oxford)* **2019** (2019).

1299

1300 76. Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional
1301   changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome*
1302   *Biol* **16**, 278 (2015).

1303
1304 77. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018).

1305
1306 78. Bergen, V., Lange, M., Peidli, S., Wolf, F.A. & Theis, F.J. Generalizing RNA velocity to
1307   transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).

1308
1309 79. Van de Sande, B. *et al.* A scalable SCENIC workflow for single-cell gene regulatory
1310   network analysis. *Nat Protoc* **15**, 2247-2276 (2020).

1311
1312 80. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
1313   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

1314
1315 81. Kim, D., Langmead, B. & Salzberg, S.L. HISAT: a fast spliced aligner with low memory
1316   requirements. *Nat Methods* **12**, 357-360 (2015).

1317
1318 82. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for
1319   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

1320
1321 83. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
1322   for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1323
1324 84. Fernandes, A.D. *et al.* Unifying the analysis of high-throughput sequencing datasets:
1325   characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by
1326   compositional data analysis. *Microbiome* **2**, 15 (2014).

1327
1328 85. Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V. & Egozcue, J.J. Microbiome Datasets
1329   Are Compositional: And This Is Not Optional. *Front Microbiol* **8**, 2224 (2017).

1330

1331

Figure 1. Treg-intrinsic TFAM loss reprograms CD4⁺ T cell lineage fate toward pro-inflammatory and terminal effector states. (**A**) Schematic representation of the experimental workflow. Splenocytes were isolated from Foxp3^Cre^Tfam^f/f^ (TFAM cKO) and Tfam^f/f^ (WT) mice for single-cell RNA sequencing (scRNA-seq) and downstream analysis. (**B**) UMAP projection of 49,156 immune cells (WT: 34,958; cKO: 14,198), identifying 21 major immune subsets across 26 transcriptional clusters. Erythroid lineage cells were excluded. (**C**) UMAP projection of CD4⁺ T cells (WT: 4,324; cKO: 3,497), revealing six transcriptionally distinct subsets: naïve, effector memory (TEM), exhausted, natural Tregs (nTregs), aged/dysfunctional Tregs (aTregs), and cytotoxic CD4⁺ T cells. (**D**) Track plot showing representative gene expression patterns across single CD4⁺ T cells. Rows represent genes; columns represent individual cells. (**E**) Bar graph quantifying relative abundance of CD4⁺ T cell subsets in WT and cKO mice. TFAM deficiency promotes expansion of effector-like states (aTregs, cytotoxic, and TEM), with reduced naïve and exhausted populations. (**F**) Gene set enrichment analysis (GSEA) of KEGG pathways shows increased TNFα signaling, cytokine/chemokine responses, and calcium flux in CD4⁺ T cells from TFAM cKO mice. (**G**) Pseudotime trajectory analysis (Monocle) of CD4⁺ T cells reveal two dominant differentiation paths (P1 and P2). TFAM-deficient cells preferentially adopt the alternative P2 trajectory, enriched in transcriptional States 1 and 2. (**H**) Monocle 2 trajectory cell states for all CD4+ T cells, each branch represents one cell state (left). Barplots representing cell proportions in the seven states of WT and cKO (right). Cells are colored according to seven states, which partition the trajectory. (**I**) Relative proportion of CD4+ T cells among different states of WT and cKO spleen, color stands for cell type. (**J**) RNA velocity analysis confirms directionality of lineage bifurcation, with streamlines overlaid on UMAP embedding indicating transcriptional flow. (**K**) Heatmap of four gene clusters ordered by pseudotime. Cluster 1 genes (e.g., Klf3, Bcl2) mark early developmental states; Cluster 2 genes (e.g., Ccl5, S100a6) are linked to terminal effector function and inflammation. KEGG pathway enrichment of cluster-specific genes is shown. (**L**) Representative flow cytometry plots of CD4⁺CD44^hi^CD62L^lo^ effector and CD4⁺CD44^lo^CD62L^hi^ naïve T cells in spleen and lymph nodes from WT and TFAM cKO mice. (**M**) Quantification of mean fluorescence intensity (MFI) for effector and naïve CD4⁺ T cells across organs. TFAM cKO mice exhibit increased effector T cell signatures. (**N**) Suppression assay showing reduced regulatory function of TFAM-deficient Tregs in vitro. Co-culture with CFSE-labeled CD4⁺ Tconv cells demonstrates impaired suppression of proliferation upon CD3/CD28 stimulation.

Figure 2. Treg-intrinsic TFAM loss triggers a multi-lineage stress response, inflammatory reprogramming, and systemic senescence. (**A**) Gene regulatory network (GRN) inference from scRNA-seq data identified 177 regulons across CD4+ T cell subsets. TFAM cKO Tregs showed increased activity of inflammatory (*Nfkb2*, *Nfatc1*, *Rora*, *Runx2*), stress-responsive (*Egr1*, *Junb*, *Xbp1*), and cytotoxic (*Runx3*) regulons, alongside pan-CD4+ expression of the mitochondrial stress gene *Ddit3*. (**B**) Dot plots showing upregulation of key inflammatory transcriptional regulators (*Prdm1*, *Nfkb1*,*Nfkb2*, *Ahr*) in TFAM-deficient Tregs. (**C**) Gene set enrichment analysis (GSEA) shows significant pathways in Monocytes (Mono), Macrophages (Mac), dendritic cell (DC), Immature Dendritic Cells (iDC) and Plasmacytoid Dendritic Cells (pDC) cells in TFAM cKO mice. (**D**) Bulk RNA-seq of cKO spleens revealed enrichment of cell death pathways, including apoptosis and necroptosis. (**E**) Immunoblots confirmed caspase-3 and -8 cleavage, IκBα degradation, and phosphorylation of RIPK1, RIPK3, and MLKL, indicative of RIPK-mediated immunogenic cell death. (**F**) Histological analysis revealed multi-organ inflammatory infiltration in colon, liver, lung, and kidney of TFAM cKO mice. (**G**) H&E staining of femurs showed pronounced cortical bone thinning in TFAM cKO animals. (**H**) Spleens from TFAM cKO mice exhibited splenomegaly, follicular disruption, and marginal zone disorganization. (**I**) Mitospy staining in sorted Tregs revealed mitochondrial depolarization and elevated oxidative stress in TFAM-deficient cells. (**J**) Elevated circulating levels of TNF-α, IL-6, and IFNγ in serum from TFAM cKO mice, confirming systemic inflammation. (**K**) RT-qPCR showed increased expression of Cdkn2a (p16) and Cdkn1a (p21). (**L**) Immunoblotting of splenic lysates confirmed elevated levels of p53, CDK4, PCNA, p16, andp21, consistent with senescence-associated gene expression. (**M**) Whole-spleen RNA-seq revealed enrichment of aging- and inflammation-associated gene networks. (**N**) GSEA showed significant upregulation of pathways linked to neuroactive ligand–receptor signaling, and downregulation of circadian rhythm disruption, and protein trafficking which are hallmarks of systemic immune aging.

Figure 3. Partial TFAM reduction in Tregs promotes systemic aging features, peripheral inflammation, and early neurodegenerative changes. (**A**) Human PBMC transcriptome analysis reveals age-associated decline in TFAM expression, supporting its role as a mitochondrial-immune rheostat. (**B-C**) TFAM Het (Foxp3CreTfamf/+) mice develop aging phenotypes, including kyphosis and reduced body weight, by 8-10 months of age. (**D**) Circulating TNF-α and IL-6 levels are elevated in TFAM Het mice, indicating persistent low-grade systemic inflammation. (**E**) Splenic CD4⁺ T cells isolated from TFAM Het mice display elevated expression of senescence-associated markers, including Cdkn2a (*p16^lnk4a*), Cdkn1a (*p21^Cip1*), p53, and CDK4, with GAPDH serving as the loading control. (**F**) Neuromuscular assessment shows reduced forelimb grip strength in TFAM Het mice. (**G**) Open field behavioral assay demonstrates decreased exploratory behavior and locomotor hypoactivity, consistent with CNS dysfunction. (**H–I**) Cresyl violet staining reveals neuronal loss in the hippocampus and cortex of 8-month-old TFAM Het mice compared to 8-week controls. (**J**) Longitudinal hippocampal RNA-seq profiling shows minimal transcriptional changes at 8 weeks but extensive neuroimmune remodeling by 8 months. (**K**) Heatmap displaying upregulation of neuroinflammation-associated genes (*Gabra5*, *Slc38a5*, *Calcr*, *Ganbare*) in 8-month TFAM Het hippocampus. (**L**) KEGG pathway enrichment of 8-month hippocampal transcriptomes identify significant enriched pathways of oxidative stress, inflammatory signaling, and neurodegeneration-linked pathways.

Figure 4. TFAM insufficiency in Tregs triggers CNS immune activation and microglial sensitization to TNF signaling. (**A-B**) Iba1 and NeuN immunohistochemistry shows widespread activation of amoeboid microglia in the cortex and hippocampus of 8-month-old TFAM Het mice, with concurrent neuronal loss.(**C**) Flow cytometric analysis reveals increased MHC-I expression on brain-resident CX3CR1+ microglia from TFAM Het mice, consistent with enhanced antigen presentation. (**D**) Ex vivo stimulation with TNF-α induces elevated *Il1b*, *Tnfr1*, and *Il6* expression in CX3CR1+ microglia from TFAM Het brains, indicating heightened inflammatory sensitivity.

Figure 5. TFAM insufficiency in Tregs drives microbiota–lipid dysregulation and neuroinflammatory reprogramming via the gut–brain–immune axis. (**A**) Longitudinal 16S rRNA sequencing reveals significantly elevated α-diversity (Shannon and Simpson indices) in TFAM Het mice, peaking at 8 months of age. (**B**) β-diversity analysis (Bray–Curtis dissimilarity) demonstrates genotype- and age-specific microbial clustering, indicating progressive community remodeling. (**C**) Phylum-level taxonomic profiling reveals expansion of *Campilobacterota* and depletion of *Bacteroidota* in aged TFAM Het mice, signatures linked to inflammaging. (**D**) VennDiagram depicting distribution and overlap of microbial species across genotypes and age groups. (**E**) Differential abundance analysis identifies 18 significantly altered taxa, including loss of key SCFA-producing genera (*Lactobacillus murinus*, *Odoribacter*, *Prevotellaceae_UCG-001*) in TFAM Het mice. (**F**) KEGG-based predictive metagenomics shows TFAM Het microbiota enriched in age-related pathways including FoxO, and spinocerebellar ataxia signaling. (**G**) Lipid metabolism pathways, including unsaturated fatty acid biosynthesis, bile acid biosynthesis, and sphingolipid metabolism, are significantly altered. Neurotransmission pathways (GABAergic, glutamatergic synapses) are also enriched in aged TFAM Het mice. (**H**) Microbiota–host gene co-correlation network analysis reveals 323 significant interactions. Notably, *Helicobacter* positively correlates with *Ntsr2*, while Lactobacillus inversely correlates with neuroendocrine (e.g., *Pomc*, *Sstr5*) and developmental genes (*Elk3*, *Sox4*) and positively correlates with inflammatory genes (*Mpo*, *Ifitm6*). *Lachnospiraceae_NK4A136_group* links to complement-mediated inflammation (*C5ar2*). (**I**) RT-qPCR validation confirms elevated expression of *Il1b* and *Tnfr1* in peripheral tissues from TFAM Het mice, supporting a chronic inflammatory state.

Figure 6. Restoration of mitochondrial fitness mitigates inflammation and functional decline in models of immune aging. (**A**) Schematic of therapeutic intervention in TFAM cKO mice. Mice received intraperitoneal resveratrol (RSV; 7 mg/kg) every other day for 4 weeks. (**B**) Mitospy staining in skin tissue reveals enhanced mitochondrial membrane potential in RSV-treated TFAM cKO mice, indicating improved mitochondrial integrity. (**C**) Serum levels of TNF-α and IL-6 were significantly reduced following RSV treatment, reflecting dampened systemic inflammation. (**D-F**) Quantitative RT-PCR analysis of splenocytes shows reduced expression of senescence markers Cdkn2a (p16Ink4a) and Cdkn1a (p21Cip1) in RSV-treated mice, consistent with suppression of the senescence-associated secretory phenotype (SASP). (**G**) Histological examination of lung, liver, and kidney shows decreased inflammatory infiltration and preserved tissue structure in RSV-treated animals. (**H**) Experimental design for fecal microbiota transplantation (FMT) in TFAM Het mice. Mice were antibiotic-conditioned for 10 days and subsequently gavage with feces from TFAM-sufficient donors over 3 days, followed by a 3-week recovery. (**I**) FMT improved neuromuscular performance, demonstrated by increased rotarod endurance and forelimb grip strength. These improvements were associated with re-expression of Treg-associated genes and suppression of pro-inflammatory programs. (**J**) Working model summarizing the role of TFAM as a central regulator of mitochondrial redox balance, microbiota composition, and immune aging. Restoration of mitochondrial integrity via RSV and microbial reprogramming via FMT ameliorates immunopathology and rejuvenates immune function.

Supplementary Fig. 1: Landscape of scRNA-seq analysis of WT and cKO spleen. A. t-SNE projections of single-cell transcriptomes from WT and cKO mice before Erythroid cells were removed, annotated by cell type. Each dot denotes a distinct cell. **B.** Cell subset proportions (top) and numbers (bottom) for WT and cKO spleens. **C.** Dot plot of mean expression of canonical marker genes for 20 major lineages from tissues of each origin, as indicated. **D.** Gene Set Enrichment Analysis (GSEA) of KEGG pathways for all cell types between cKO and WT (Only age-related pathways are selected). Color stands for the up-regulated (red) or down-regulated (blue) in cKO spleen. Enriched terms were identified as significant at a p-value ≤ 0.01. **E.** Expression of aged-related genes (*Fcer2a*, *Cd79b*, *Zfp36l2*, and *S100a6*) on Figure 1B t-SNE embedding. Each dot corresponds to a single cell.

Supplementary Fig. 2: scRNA-seq results show the cellular heterogeneity in CD4+ T cells.
A. Relative proportion of CD4+ T cells among WT and cKO spleen, color stands for cell type. **B.** Violin plot showing the expression profile of age-related genes among all CD4+ T cells. **C.** Monocle 2 prediction of CD4+ T cell subsets developmental trajectory with pseudotime. Each dot represents an individual cell. **D.** Functional enrichment analysis of KEGG pathways with up-regulated differential expressed genes for the cell states. Only the top 20 of the enriched pathways were plotted from each state.

Supplementary Fig. 3: t-SNE projection of binary regulon activity. t-SNE plots visualize the binarized regulon activity states of Irf8, Runx2, Bcl11a, and Nfkb1 across single cells, highlighting distinct transcriptional regulatory modules and cell-type-specific activation patterns.

Supplementary Fig. 4: TFAM deletion in Foxp3⁺ Tregs induces systemic pathology and accelerates immune aging in Foxp3^CreTfam^f/f mice. A. Representative photograph showing the deteriorated physical appearance of a cKO mouse (right) compared to an age- and sex-matched WT littermate (left) at 8 weeks of age. **B.** X-ray image depicting increased spinal curvature in the cKO mouse (right) relative to a WT littermate (left) of the same age and sex. **C.** Kaplan-Meier survival analysis comparing WT (n=10; M/F) and cKO (n=10; M/F) mice, revealing reduced lifespan in cKO animals. **D.** Flow cytometric analysis of Annexin V staining in spleen and skin cells from WT and cKO mice, indicating increased cell death in cKO tissues. **E**. Mitospy staining of splenocytes from WT and cKO mice, assessed by flow cytometry and presented as a fluorescence intensity histogram, showing mitochondrial alterations in cKO cells.

Supplementary Fig. 5: Increased cell infiltration and inflammation in different organs in Foxp3^{Cre}Tfam$^{f/f}$ mice. A. Histology sections stained with H&E from lung, liver, colon, and kidney. **B.** Serum cytokines identified by Legend plex for IL-1β, IL-10, MCP1, IL-23, IL-1α, IL-17, IFNγ, IL-12, IFNβ, GM-CSF, and IL-27 with estimated values displayed. Data represented as mean ± SD. Statistical test used: t-test, WT (n=5-10; M/F) and cKO mice (n=5-10, M/F).

Supplementary Fig. 6: Restoration of mitochondrial function leads to mitigate pathologies.
A-B. Body weight (**A**) and fold change (**B**) of young and old WT and cKO mice. Data represented as mean ± SD. Statistical test used: two-way ANOVA with post-hoc test (**B**). WT (n=5; M/F) and Het mice (n=5, M/F). (**C**) Representative images of H&E histology in spleen and lung from WT (8 weeks, 8 months, and 3 years) and Het (8 weeks and 8 months) mice.