Structure of the IL-27 quaternary receptor signaling complex

Nathanael A Caveney[1,2], Caleb R Glassman[1,2,3], Kevin M Jude[1,4], Naotaka Tsutsumi[1,4], K Christopher Garcia[1,3,4,*]

[1]Departments of Molecular and Cellular Physiology and Structural Biology, Stanford University School of Medicine, Stanford, CA 94305, USA
[2]These authors contributed equally
[3]Program in Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA
[4]Howard Hughes Medical Institute, Stanford University School of Medicine, Stanford, CA 94305, USA
*Correspondence: kcgarcia@stanford.edu

Abstract

Interleukin 27 (IL-27) is a heterodimeric cytokine that functions to constrain T cell-mediated inflammation and plays an important role in immune homeostasis. Binding of IL-27 to cell surface receptors IL-27Rα and gp130 results in activation of receptor-associated Janus Kinases and nuclear translocation of STAT1 and STAT3 transcription factors. Despite the emerging therapeutic importance of this cytokine axis in cancer and autoimmunity, a molecular blueprint of the IL-27 receptor signaling complex, and its relation to other gp130/IL-12 family cytokines, is currently unclear. We used cryogenic-electron microscopy (cryo-EM) to determine the quaternary structure of IL-27 (p28/Ebi3) bound to receptor subunits, IL-27Rα and gp130. The resulting 3.47 Å resolution structure revealed a three-site assembly mechanism nucleated by the central p28 subunit of the cytokine. The overall topology and molecular details of this binding are reminiscent of IL-6 but distinct from related heterodimeric cytokines IL-12 and IL-23. These results indicate distinct receptor assembly mechanisms used by heterodimeric cytokines with important consequences for targeted agonism and antagonism of IL-27 signaling.

Introduction

Cytokines are secreted factors that mediate cell-cell communication in the immune system (Spangler et al., 2015). Binding of cytokines to cell surface receptors leads to activation of receptor-associated Janus Kinase (JAK) proteins which phosphorylate each other as well as downstream Signal Transducer and Activator of Transcription (STAT) proteins, triggering nuclear translocation and regulation of gene expression. Cytokines can be classified by their use of shared receptors which transduce signals for multiple cytokines within a family. Interleukin-6 signal transducer (IL-6ST), also known as glycoprotein 130 (gp130), is a shared receptor that mediates signaling of multiple cytokines including IL-6, IL-11, and IL-27. Unlike other gp130 family cytokines, IL-27 is a heterodimeric cytokine consisting of a four-helix bundle, IL-27p28 (p28), with similarity to IL-6, complexed with a secreted binding protein, Epstein-Barr Virus-Induced 3 (Ebi3), with homology to type I cytokine receptors (Pflanz et al., 2002). IL-27 signals through a receptor complex consisting of IL-27Rα (TCCR/WSX-1) and gp130 expressed predominantly on T cells and NK cells (Pflanz et al., 2002, 2004) (Figure 1A). Binding of IL-27 to its receptor subunits triggers the activation of receptor-associated Janus Kinase 1 (JAK1) and JAK2 leading to phosphorylation of STAT1 and STAT3 (Lucas et al., 2003; Owaki et al., 2008; Wilmes et al., 2021). Functionally, IL-27 signaling serves to constrain inflammation by antagonizing differentiation of pro-inflammatory Th17 cells (Stumhofer et al., 2006), stimulating T-bet expression in regulatory T cells (Tregs) (Hall et al.,

49 2012), and inducing production of the anti-inflammatory cytokine IL-10 (Stumhofer et al.,
50 2007). The important role of IL-27 in constraining inflammation has prompted the development
51 of antagonist antibodies, one of which is currently being developed as a cancer
52 immunotherapy (Patnaik et al., 2021).
53
Although IL-27 is a member of the gp130 family, its heterodimeric composition is similar to IL-
12 and IL-23 which are cytokines that share a p40 subunit similar to Ebi3. In this system, the
p40 subunit of IL-12 and IL-23 directly engages the shared receptor IL-12Rβ1 in a manner
distinct from IL-6 and related cytokines (Glassman et al., 2021a). Given the diverse receptor
assembly mechanisms used by cytokine receptors, we sought to determine the structure of
IL-27 in complex with its signaling receptors, gp130 and IL-27Rα, by cryogenic-electron
microscopy (cryo-EM).
61
Here we report the 3.47 Å resolution structure of the complete IL-27 receptor complex. In this
structure, the central IL-27 cytokine engages IL-27Rα through a composite interface consisting
of p28 and the D2 domain of Ebi3. A conserved tryptophan residue at the tip of p28 then
engages the D1 domain of gp130 to facilitate quaternary complex assembly. This receptor
assembly mechanism bears striking resemblance to that of IL-6 but is distinct from IL-12 and
IL-23, indicating divergent receptor binding modes for heterodimeric cytokines.
68
69 **Results**
70
71 **CryoEM structure of the IL-27 quaternary complex**
72
IL-27 (p28/Ebi3) signals through a receptor complex composed of IL-27Rα and the shared
receptor gp130 (Figure 1A). The extracellular domain (ECD) of IL-27Rα consists of five
fibronectin type III (FNIII) domains (D1-5) of which the N-terminal D1-D2 constitute a cytokine-
binding homology region (CHR). gp130 has a similar domain architecture but with the addition
of an N-terminal immunoglobin (Ig) domain (D1). Initial attempts at reconstituting a soluble
receptor complex through mixing of IL-27, IL-27Rα D1-D2, and gp130 D1-D3 resulted in
dissociation on size-exclusion chromatography due to the low affinity of gp130. To stabilize
the complex, we expressed p28 fused to gp130 through a long (20 a.a.) flexible linker which
enabled us to purify the complete IL-27 receptor complex by size exclusion chromatography
(SEC). Importantly, the long linker did not constrain the binding mode but rather raised the
effective concentration of gp130. The complex was vitrified and subject to single-particle
cryoEM analysis. The quaternary complex was determined to a resolution of 3.47 Å (Table
S1, Supplementary Figure 1). All domains of IL-27 and IL-27Rα, as well as D1 of gp130 had
well resolved cryoEM density (Figure 1B-C, Supplemental Figure 1E), with localised resolution
estimates near and exceeding 3 Å in this region (Supplemental Figure 1B). The localised
resolution for the D2 density of gp130 is lower, yet able to be built with confidence, while D3
electron density was interpretable for domain placement (Figure 1C, Supplementary Figure
1B).
91
The IL-27 quaternary complex exhibits a molecular architecture containing "sites 1-3", as seen
in other gp130 family cytokines (Boulanger et al., 2003) (Figure 1D). In this structure, IL-27
bridges IL-27Rα and gp130 to initiate downstream signaling through JAK1 and JAK2 (Ferrao
et al., 2016; Pradhan et al., 2010). Within IL-27, the four-helical bundle of p28 packs against
the hinge between Ebi3 D1 and D2 in a site 1 interaction. Opposite Ebi3, IL-27Rα D1-D2

engages the helical face of p28 in an interaction stabilized by stem contact between the D2
domains of IL-27Rα and Ebi3 to form a site 2 interaction. On the posterior face of IL-27, the
D1 domain of gp130 binds to a conserved tryptophan at the tip of p28 to make a classical site
3 interaction (Boulanger et al., 2003).
101

102
103   **Figure 1. Composition and cryoEM structure of human IL-27 quaternary complex.** (**A**)
104   Cartoon representation of the components of the IL-27 quaternary signaling complex. gp130 (red),

p28 (blue), Ebi3 (purple), IL-27Rα (green), with domains excluded from the imaged constructs in grey. The D1 Ig domain of gp130 is represented by red, with an additional dark red outline to distinguish it from FNIII domains in gp130, Ebi3, and IL-27Rα. (**B**) Reference-free 2D averages from cryoEM of the IL-27 quaternary complex. (**C**) Refined and sharpened cryoEM density maps of IL-27 quaternary complex, colored as in **A**. (**D**) Ribbon representation of the atomistic modelling of IL-27 quaternary complex, colored as in **A**.

Site 1-3 interface architecture of IL-27 quaternary complex

A site 1 interface between p28 and Ebi3 is used to form the heterodimeric cytokine IL-27. In this interaction, the alpha helices A and D, the AB loop, and the C-terminus of p28 pack tightly against the hinge region between D1 and D2 of Ebi3 (Figure 2A-B). This interaction is mediated by a variety of hydrogen bonds between p28 and the hinge region of Ebi3, particularly along alpha helices A and D of p28. At the center of this interface is p28 R219 which is coordinated by D207 and T209 in Ebi3. Extending from this region, a hydrophobic patch of residues on the C-terminus and AB loop of p28 (F94, W97, L223, W232) packs against residues in the Ebi3 hinge (F97, F157, I160). The extensive and hydrophobic nature of this interface is consistent with the observation that co-expression of p28 and Ebi3 is required for efficient cytokine secretion (Pflanz et al., 2002).

Opposing the Ebi3 binding site, p28 engages the hinge region of IL-27Rα (Figure 2C, site 2a). This interaction is stabilized by stem-stem contacts between the D2 domains of IL-27Rα and Ebi3 (Figure 2C, site 2b). Alpha helices A and C of p28 form the bulk of the site 2a interaction. In contrast to the site 1 interaction which forms the holo-cytokine, the site 2a interaction is more limited both in terms of hydrophobicity and buried surface area (Site 1: 1339.4 $Å^2$, Site 2a: 810.4 $Å^2$) (Krissinel and Henrick, 2007). One notable feature of this interface is the contribution of a 13 amino acid polyglutamic acid region in p28 which forms an alpha helix (polyE helix) that contacts R74 and K77 in the IL-27Rα D1 domain. Cytokine binding is stabilized by "stem-stem" interactions between D2 FNIII domains of IL-27Rα and Ebi3. This site 2b interaction is characterized by a high degree of shape and charge complementarity with an arginine-rich patch of Ebi3 (R143, R171, R194) packing against the positively charged base of IL-27Rα D2 domain (D138, D142, E146). The composite nature of IL-27Rα binding explains in part the inability of p28 to mediate IL-27 signaling in the absence of Ebi3 (Stumhofer et al., 2010).

On the posterior face of the cytokine, a classical site 3 interface is formed through the interaction of the D1 Ig domain of gp130 associating with both p28 (site 3a) and Ebi3 (site 3b) (Figure 2D,E). Similar to IL-12 family members, IL-23 and IL-12 (Glassman et al., 2021a), and the gp130 family members IL-6 (Boulanger et al., 2003) and viral IL-6 (Chow et al., 2001), this interaction is anchored by the conserved W197 on p28 which packs tightly against the base of the D1 Ig domain of gp130 (Figure 2F). The site 3 interaction is extended by the AB loop of p28 which makes contact with both gp130 and Ebi3. The tip of gp130 D1 packs against the top of Ebi3 D1 in a limited interface centered around Ebi3 F118.

Figure 2. Binding interfaces of the human IL-27 quaternary complex. (**A**) Ribbon representation of the IL-27 quaternary signaling complex, containing gp130 (red), p28 (blue), Ebi3 (purple), and IL-27Rα (green). Regions containing sites 1, 2, and 3 are boxed in purple, green, and red respectively. (**B**) Two views of the site 1 interface, colored as in **A**. (**C**) The site 2 interface, comprised of a site 2a, p28 to IL-27Rα, interaction, and a site 2b, Ebi3 to IL-27Rα, interaction. All proteins are colored as in **A**. (**D,E**) The site 3 interface, comprised of a site 3a, p28 to gp130, interaction, and a site 3b, Ebi3 to IL-27Rα, interaction. All proteins are colored as in **A**. (**F**) Structural overlay of site 3 interacting domains from IL-27 complex, IL-6 complex (PDB 1P9M), and viral IL-6 (vIL-6) complex (PDB 1L1R). IL-27 quaternary complex colored as in **A**, IL-6 ternary complex in orange, and vIL-6 binary complex in teal.

159
Figure 2 - supplement 1. CryoEM density of binding interfaces of the human IL-27
quaternary complex. (**A**) Ribbon representation of the IL-27 quaternary signaling complex,
containing gp130 (red), p28 (blue), Ebi3 (purple), and IL-27Rα (green). Regions containing sites
1, 2, and 3 are boxed in purple, green, and red respectively. (**B**) CryoEM density of the site 1
interaction. Colored as in **A**. (**C**) CryoEM density of the site 2 interaction. Colored as in **A**. (**D**)
CryoEM density of the site 3 interaction. Colored as in **A**.

166
The heterodimeric nature of IL-27 has led some to classify it as an IL-12 family cytokine
(Trinchieri et al., 2003). However, comparison of the IL-27 receptor complex with that of IL-23
and IL-12 reveals striking differences (Figure 3A-C). In IL-27, the central p28 subunit engages
all receptor components. In contrast, each subunit of IL-23 and IL-12 engages a different
receptor in a modular interaction mechanism (Glassman et al., 2021a). The assembly of IL-
27 more closely resembles that of IL-6, in which the central four-helix bundle encodes binding
sites for all receptor components (Figure 3D). However, in the case of IL-6, this motif is
duplicated through a C2 symmetry axis due to a dual role of gp130 at site 2 and site 3. The
similarity between IL-27 and IL-6 is observed not only in overall architecture but also in
molecular detail, where IL-6, vIL-6, and IL-27 all engage gp130 using a highly convergent
interface in which a tryptophan from helix D of the cytokine (W197 in p28, W185 in IL-6, W166
in vIL-6) forms an aromatic anchor that is capped by Y116 of gp130 (Figure 2F, 3D-E).
Alignment of gp130 D1–(v)IL-6 pairs to the IL-27 structure reveals a tight correlation in gp130
binding pose with Cα RMSDs of 1.039 Å (IL-6) and 1.015 Å (vIL-6) across 149 and 124
trimmed residue pairs. This striking convergence of gp130 binding modes helps to explain the
ability of p28 to antagonize IL-6 signaling (Stumhofer et al., 2010) and contextualises its place
in respect to both gp130 and IL-12 family cytokines.

Figure 3. Comparison of IL-27 to IL-12 and IL-6 family complexes. (**A**) Ribbon representation of the IL-27 signaling complex, containing gp130 (red), p28 (blue), Ebi3 (purple), and IL-27Rα (green). Sites 1, 2, and 3 are noted with a numbered circle. (**B**) Ribbon representation of the IL-23 signaling complex (IL-12 family), containing IL-23R (red), IL-23p19 (blue), p40 (purple), and IL-12Rβ1 (green) (PDB 6WDQ). Sites 1 and 3 are noted as in **A**, with the unoccupied site 2 marked with a grey 'X' and the distinct IL-12/IL-23 site 2 marked with a circled number 2. (**C**) Ribbon representation of a model of the IL-12 signaling complex (IL-12 family), containing IL-12Rβ2 (red), IL-12p35 (blue), p40 (purple), and IL-12Rβ1 (green) (Baek et al., 2021; Glassman et al., 2021a). Sites 1 and 3 are noted as in **A**, with the unoccupied site 2 marked with a grey 'X' and the distinct IL-12/IL-23 site 2 marked with a circled number 2. (**D**) Ribbon representation of the IL-6 signaling complex (IL-6 family), containing gp130 (red and green), IL-6 (blue), and IL-6Rα (purple) (PDB 1P9M). Secondary copies of each protein in the complex are colored in white. Sites 1, 2, and 3 are noted as in **A**. (**E**) Ribbon representation of the viral IL-6 signaling complex (IL-6 family), containing gp130 (red and green) and (blue) (PDB 1I1R). Secondary copies of each protein in the complex are colored in white. Sites 2 and 3 are noted as in **A**, with the unoccupied site 1 marked with a grey 'X'.

Discussion

A key paradigm in cytokine structural biology is the use of shared receptor components that
mediate diverse signaling outputs. IL-27 extends this, given that it shares the gp130 subunit
with IL-6 but mediates distinct and often countervailing functional effects. Here we find that IL-
27 assembles a receptor complex reminiscent of IL-6 but distinct from other heterodimeric IL-
12/23 class cytokines Interestingly, the soluble form of IL-6Rα (sIL-6Rα) generated through
alternative splicing or proteolytic cleavage can complex with IL-6 to potentiate signaling in
gp130-expressing cells (Harbour et al., 2020). This IL-6/sIL-6Rα complex thus may be
functionally analogous to heterodimeric cytokine IL-27.
213
As in other types of cytokine signaling, there is context dependent therapeutic potential in both
the inhibition and potentiation of IL-27 signaling. The targeting of the heterodimeric cytokines
of the IL-12 family has been well explored, with various clinically approved inhibitors, targeting
either IL-23 signaling via p19 (Risankizumab, Guselkumab, Tildrakizumab) (Tait Wojno et al.,
2019) or IL-12 and IL-23 signaling via the shared p40 (Ustekinumab) (Luo et al., 2010). In
these cases, the inhibition of IL-12 family members antagonizes the pro-inflammatory effects
of these cytokines, while IL-27 plays an inverse role in the regulation of inflammation. Due to
the role of IL-27 in the regulation T cell mediated inflammation (Tait Wojno et al., 2019),
inhibition of IL-27 is currently being explored for its use in the inhibition of aberrant IL-27
signaling in cancer (Patnaik et al., 2021). The structure of the IL-27 quaternary signaling
complex provides ample opportunities for the design of IL-27 inhibitors targeting different steps
in receptor assembly to better regulate IL-27 signaling.
226
On the other end of the spectrum, IL-27 agonism has been explored for its therapeutic use in
inflammatory autoimmune dysregulation such as experimental autoimmune encephalomyelitis
(EAE) (Fitzgerald et al., 2013) and colitis (Hanson et al., 2014). Despite the promise of IL-27
as a therapeutic, the bipartite nature of IL-27 limits its usefulness in the clinic. Using the
interfacial information provided in the structure of the IL-27 quaternary complex, structure-
guided protein engineering techniques can now be used to improve the therapeutic potential
of IL-27. One such application may be to generate single agent IL-27 agonists by affinity
maturing the interactions between p28 and its receptor subunit, as has been done recently to
generate IL-6 which do not require IL-6Rα binding for efficient signaling (Martinez-Fabregas
et al., 2019). Conversely, attenuating the affinity of p28 for IL-27Rα might be used to generate
cell-type biased partial agonists with selective activity based on differences in IL-27Rα
expression across cell-type and activation state (Villarino et al., 2005) as has been done for
IL-2 (Glassman et al., 2021b), IL-10 (Saxton et al., 2021), and IL-12 (Glassman et al., 2021a).
240
241   Resolution of the complete IL-27 receptor complex reveals how this therapeutically important
242   cytokine engages its receptor subunits. The receptor assembly mechanism bears striking
243   resemblance to that of IL-6 but is distinct from IL-12 and IL-23, indicating divergent receptor
244   binding modes for heterodimeric cytokines. This structural insight further paves the way for
245   the continued development of therapeutics that modulate IL-27 signaling.
246

Methods

Cloning and protein expression
Human IL-27Rα (D1-2, residues 36-231), Ebi3 (residues 21-228), and p28 (residues 29-243)-
(GGGGS)$_4$-gp130 (D1-D3, residues 23-321) were cloned into the expression vector pD649.
IL-27Rα and Ebi3 were expressed with signal sequence of influenza hemagglutinin and N-

terminal 6-His tags. Proteins were co-expressed in Expi293F cells (GIBCO) maintained in
Expi293 Expression Media (GIBCO) at 37°C with 5% CO_2 and gentle agitation.
255
256 **Protein purification**
257 IL-27 quaternary complex was purified by Ni-NTA chromatography followed by size exclusion
258 chromatography with a Superdex 200 column (GE Lifesciences). Fractions containing pure IL-
259 27 complex were pooled and stored at 4°C until vitrification.
260
Cryo-electron microscopy
Aliquots of 3 μL of IL-27 quaternary complex were applied to glow-discharged Quantifoil®
(1.2/1.3) grids. The grids were blotted for 3 seconds at 100% humidity with an offset of 3 and
plunge frozen into liquid ethane using a Vitrobot Mark IV (Thermo Fisher). Grid screening and
preliminary dataset collection occurred at Stanford cEMc. Final grids were imaged on a 300
kV FEI Titan Krios microscope (Thermo Fisher) located at the HHMI Janelia Research
Campus and equipped with a K3 camera and energy filter (Gatan). Movies were collected at
a calibrated magnification of ×105,000, corresponding to a 0.839 Å per physical pixel. The
dose was set to a total of 50 electrons per $Å^2$ over an exposure of 50 frames. Automated data
collection was carried out using SerialEM with a nominal defocus range set from -0.8 to -2.0
μM. 18,168 movies were collected.
272
Image processing
All processing was performed in cryoSPARC (Punjani et al., 2017) unless otherwise noted
(Supplementary Figure 1). The 18,168 movies were motion corrected using patch motion
correction and micrographs were binned to 0.839 Å per pixel. The contrast transfer functions
(CTFs) of the flattened micrographs were determined using patch CTF and 6,387,370 particles
were picked using blob picking and subsequently template picking. A subset of 2,008,987
particles were used in reference-free 2D classification. A particle stack containing 109,536 2D
cleaned particles was used to generate three *ab initio* models, one of which resembled the IL-
27 quaternary complex. This good *ab initio* model was then used against two junk classes in
6 rounds of iterative heterogenous refinement to reduce the full particle stack to 548,147
particles. These particles were refined using cryoSPARC non-uniform refinement (Punjani et
al., 2020) followed by local refinement to achieve a global resolution of 3.47 Å. Resolution was
determined at a criterion of 0.143 Fourier shell correlation gold-standard refinement
procedure. The final map was sharpened using deepEMhancer (Sanchez-Garcia et al., 2020).
287
Model building and refinement
AlphaFold models (Jumper et al., 2021) of IL-27Rα, Ebi3, p28, and gp130 were docked into
the map using UCSF Chimera X (Pettersen et al., 2021). The resultant model was then refined
using Phenix real space refine (Adams et al., 2010) and manual building in Coot (Emsley and
Cowtan, 2004). The final model fit the map well (EMRinger (Barad et al., 2015) score 2.33)
and produced a favorable MolProbity score of 1.60 (Chen et al., 2010) with side-chain
rotamers occupying 97.37% Ramachandran favoured and 0.00% outliers (Supplementary
Table 1).
296
297 **Data Availability**
298 CryoEM maps and atomic coordinates for human IL-27 quaternary complex have been
299 deposited in the EMDB (EMD- 26382) and PDB (7U7N) respectively.

Acknowledgements

We thank Rui Yan at the HHMI Janelia CryoEM Facility for help in microscope operation and final data collection. We thank Liz Montabana and Stanford cEMc for microscope access for preliminary data collection. NAC is a CIHR postdoctoral fellow. KCG is an investigator with the Howard Hughes Medical Institute. KCG is supported by National Institutes of Health grant R01-AI51321, the Mathers Foundation, and the Ludwig Foundation.

Author Contributions

NAC contributed to conceptualization, methodology, investigation, analysis, writing – original draft, review, and editing. CRG contributed to conceptualization, methodology, investigation, analysis, writing – original draft, review, and editing. KMJ contributed to analysis, writing – review, and editing. NT contributed to methodology, investigation, and writing – review. KGC contributed to conceptualization, supervision, writing – review and editing, and funding acquisition.

Competing Interests

KCG is the founder of Synthekine.

References

Adams, P.D., Afonine, P. V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.-W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr. D. Biol. Crystallogr. *66*, 213–221.

Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G.R., Wang, J., Cong, Q., Kinch, L.N., Schaeffer, R.D., et al. (2021). Accurate prediction of protein structures and interactions using a three-track neural network. Science *373*, 871–876.

Barad, B.A., Echols, N., Wang, R.Y.R., Cheng, Y., Dimaio, F., Adams, P.D., and Fraser, J.S. (2015). EMRinger: Side chain-directed model and map validation for 3D cryo-electron microscopy. Nat. Methods *12*, 943–946.

Boulanger, M.J., Chow, D. chone, Brevnova, E.E., and Garcia, K.C. (2003). Hexameric structure and assembly of the interieukin-6/IL-6 α-receptor/gp130 complex. Science (80-.). *300*, 2101–2104.

Chen, V.B., Arendall, W.B., Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: All-atom structure validation for macromolecular crystallography. Acta Crystallogr. Sect. D Biol. Crystallogr. *66*, 12–21.

Chow, D.C., He, X.L., Snow, A.L., Rose-John, S., and Christopher Garcia, K. (2001). Structure of an extracellular gp130 cytokine receptor signaling complex. Science (80-.). *291*, 2150–2155.

Emsley, P., and Cowtan, K. (2004). Coot: Model-building tools for molecular graphics. Acta Crystallogr. Sect. D Biol. Crystallogr. *60*, 2126–2132.

Ferrao, R., Wallweber, H.J.A., Ho, H., Tam, C., Franke, Y., Quinn, J., and Lupardus, P.J. (2016). The Structural Basis for Class II Cytokine Receptor Recognition by JAK1. Structure *24*, 897–905.

Fitzgerald, D.C., Fonseca-Kelly, Z., Cullimore, M.L., Safabakhsh, P., Saris, C.J.M., Zhang, G.-X., and Rostami, A. (2013). Independent and Interdependent Immunoregulatory Effects of IL-27, IFN-β, and IL-10 in the Suppression of Human Th17 Cells and Murine Experimental Autoimmune Encephalomyelitis. J. Immunol. *190*, 3225–3234.

Glassman, C.R., Mathiharan, Y.K., Jude, K.M., Su, L., Panova, O., Lupardus, P.J., Spangler,

J.B., Ely, L.K., Thomas, C., Skiniotis, G., et al. (2021a). Structural basis for IL-12 and IL-23 receptor sharing reveals a gateway for shaping actions on T versus NK cells. Cell *184*, 983-999.e24.

Glassman, C.R., Su, L., Majri-Morrison, S.S., Winkelmann, H., Mo, F., Li, P., Pérez-Cruz, M., Ho, P.P., Koliesnik, I., Nagy, N., et al. (2021b). Calibration of cell-intrinsic interleukin-2 response thresholds guides design of a regulatory T cell biased agonist. Elife *10*.

Hall, A.O., Beiting, D.P., Tato, C., John, B., Oldenhove, G., Lombana, C.G., Pritchard, G.H., Silver, J.S., Bouladoux, N., Stumhofer, J.S., et al. (2012). The cytokines interleukin 27 and interferon-γ promote distinct Treg cell populations required to limit infection-induced pathology. Immunity *37*, 511–523.

Hanson, M.L., Hixon, J.A., Li, W., Felber, B.K., Anver, M.R., Stewart, C.A., Janelsins, B.M., Datta, S.K., Shen, W., McLean, M.H., et al. (2014). Oral delivery of il-27 recombinant bacteria attenuates immune colitis in mice. Gastroenterology *146*, 210-221.e13.

Harbour, S.N., DiToro, D.F., Witte, S.J., Zindl, C.L., Gao, M., Schoeb, T.R., Jones, G.W., Jones, S.A., Hatton, R.D., and Weaver, C.T. (2020). TH17 cells require ongoing classic IL-6 receptor signaling to retain transcriptional and functional identity. Sci. Immunol. *5*.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.

Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state. J. Mol. Biol. *372*, 774–797.

Lucas, S., Ghilardi, N., Li, J., and de Sauvage, F.J. (2003). IL-27 regulates IL-12 responsiveness of naive CD4+ T cells through Stat1-dependent and -independent mechanisms. Proc. Natl. Acad. Sci. U. S. A. *100*, 15047–15052.

Luo, J., Wu, S.-J., Lacy, E.R., Orlovsky, Y., Baker, A., Teplyakov, A., Obmolova, G., Heavner, G.A., Richter, H.-T., and Benson, J. (2010). Structural basis for the dual recognition of IL-12 and IL-23 by ustekinumab. J. Mol. Biol. *402*, 797–812.

Martinez-Fabregas, J., Wilmes, S., Wang, L., Hafer, M., Pohler, E., Lokau, J., Garbers, C., Cozzani, A., Fyfe, P.K., Piehler, J., et al. (2019). Kinetics of cytokine receptor trafficking determine signaling and functional selectivity. Elife *8*.

Owaki, T., Asakawa, M., Morishima, N., Mizoguchi, I., Fukai, F., Takeda, K., Mizuguchi, J., and Yoshimoto, T. (2008). STAT3 is indispensable to IL-27-mediated cell proliferation but not to IL-27-induced Th1 differentiation and suppression of proinflammatory cytokine production. J. Immunol. *180*, 2903–2911.

Patnaik, A., Morgensztern, D., Mantia, C., Tannir, N.M., Harshman, L.C., Hill, J., White, K., Chung, J.-K., Bowers, B., Sciaranghella, G., et al. (2021). Results of a phase 1 study of SRF388, a first-in-human, first-in-class, high-affinity anti-IL-27 antibody in advanced solid tumors. J. Clin. Oncol. *39*, 2551–2551.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci. *30*, 70–82.

Pflanz, S., Timans, J.C., Cheung, J., Rosales, R., Kanzler, H., Gilbert, J., Hibbert, L., Churakova, T., Travis, M., Vaisberg, E., et al. (2002). IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4+ T cells. Immunity *16*, 779–790.

Pflanz, S., Hibbert, L., Mattson, J., Rosales, R., Vaisberg, E., Bazan, J.F., Phillips, J.H., McClanahan, T.K., de Waal Malefyt, R., and Kastelein, R.A. (2004). WSX-1 and glycoprotein 130 constitute a signal-transducing receptor for IL-27. J. Immunol. *172*, 2225–2231.

Pradhan, A., Lambert, Q.T., Griner, L.N., and Reuther, G.W. (2010). Activation of JAK2-V617F by components of heterodimeric cytokine receptors. J. Biol. Chem. *285*, 16651–16663.

Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). CryoSPARC: Algorithms for rapid unsupervised cryo-EM structure determination. Nat. Methods *14*, 290–296.

Punjani, A., Zhang, H., and Fleet, D.J. (2020). Non-uniform refinement: adaptive

regularization improves single-particle cryo-EM reconstruction. Nat. Methods *17*, 1214–
1221.
Sanchez-Garcia, R., Gomez-Blanco, J., Cuervo, A., Carazo, J.M., Sorzano, C.O.S., and
Vargas, J. (2020). DeepEMhancer: A deep learning solution for cryo-EM volume post-
processing. BioRxiv.
Saxton, R.A., Tsutsumi, N., Su, L.L., Abhiraman, G.C., Mohan, K., Henneberg, L.T., Aduri,
N.G., Gati, C., and Garcia, K.C. (2021). Structure-based decoupling of the pro- and anti-
inflammatory functions of interleukin-10. Science *371*.
Spangler, J.B., Moraga, I., Mendoza, J.L., and Garcia, K.C. (2015). Insights into Cytokine–
Receptor Interactions from Cytokine Engineering. Annu. Rev. Immunol. *33*, 139–167.
Stumhofer, J.S., Laurence, A., Wilson, E.H., Huang, E., Tato, C.M., Johnson, L.M., Villarino,
A. V, Huang, Q., Yoshimura, A., Sehy, D., et al. (2006). Interleukin 27 negatively regulates
the development of interleukin 17-producing T helper cells during chronic inflammation of the
central nervous system. Nat. Immunol. *7*, 937–945.
Stumhofer, J.S., Silver, J.S., Laurence, A., Porrett, P.M., Harris, T.H., Turka, L.A., Ernst, M.,
Saris, C.J.M., O'Shea, J.J., and Hunter, C.A. (2007). Interleukins 27 and 6 induce STAT3-
mediated T cell production of interleukin 10. Nat. Immunol. *8*, 1363–1371.
Stumhofer, J.S., Tait, E.D., Quinn, W.J., Hosken, N., Spudy, B., Goenka, R., Fielding, C.A.,
O'Hara, A.C., Chen, Y., Jones, M.L., et al. (2010). A role for IL-27p28 as an antagonist of
gp130-mediated signaling. Nat. Immunol. *11*, 1119–1126.
Tait Wojno, E.D., Hunter, C.A., and Stumhofer, J.S. (2019). The Immunobiology of the
Interleukin-12 Family: Room for Discovery. Immunity *50*, 851–870.
Trinchieri, G., Pflanz, S., and Kastelein, R.A. (2003). The IL-12 family of heterodimeric
cytokines: new players in the regulation of T cell responses. Immunity *19*, 641–644.
Villarino, A. V, Larkin, J., Saris, C.J.M., Caton, A.J., Lucas, S., Wong, T., de Sauvage, F.J.,
and Hunter, C.A. (2005). Positive and negative regulation of the IL-27 receptor during
lymphoid cell activation. J. Immunol. *174*, 7684–7691.
Wilmes, S., Jeffrey, P.-A., Martinez-Fabregas, J., Hafer, M., Fyfe, P.K., Pohler, E., Gaggero,
S., López-García, M., Lythe, G., Taylor, C., et al. (2021). Competitive binding of STATs to
receptor phospho-Tyr motifs accounts for altered cytokine responses. Elife *10*.
436