Odors drive feeding through gustatory receptor neurons in *Drosophila*

Hong-ping Wei[1], Thomas Ka Chung Lam[2], and Hokto Kazama[1,3,4,*]

[1]RIKEN Center for Brain Science, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan.

[2]Neuroengineering Laboratory, Brain Mind Institute & Interfaculty Institute of Bioengineering, EPFL, Lausanne, Switzerland

[3]RIKEN CBS-KAO Collaboration Center, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan.

[4]Graduate School of Arts and Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8902, Japan.

*Correspondence: hokto.kazama@riken.jp

Summary

Odors are intimately tied to the taste system to aid food selection and determine the sensory experience of food. However, how smell and taste are integrated in the nervous system to drive feeding remains elusive. We show in *Drosophila* that odors alone activate gustatory receptor neurons (GRNs), trigger proboscis extension reflex (PER), a canonical taste-evoked feeding behavior, and enhance food intake. Odor-evoked PER requires the function of sugar-sensing GRNs but not olfactory organs. Calcium imaging and electrophysiological recording show that GRNs directly respond to odors. Odor-evoked PER is mediated by the Gr5a receptor, and is bidirectionally modulated by olfactory binding proteins. Finally, odors and sucrose co-applied to GRNs synergistically enhance PER and food consumption. These results reveal a cell-intrinsic mechanism for odor-taste multimodal integration that takes place as early as in GRNs, indicating that unified chemosensory experience is a product of layered integration in peripheral neurons and in the brain.

Keywords

Odor-taste multisensory integration, gustatory receptor neuron, feeding behavior, calcium imaging, tip recording, gustatory receptor, olfactory binding protein, *Drosophila*

Introduction

Feeding behavior is an outcome of intricate interactions among multiple external and internal senses (Boesveldt and de Graaf, 2017; McCrickerd and Forde, 2016; Spence, 2015). Of particular relevance is an interplay between the senses of smell and taste. Olfactory and gustatory systems prime feeding by signaling food presence, guiding food choice, and inducing anticipatory responses in digestive organs (Boesveldt and de Graaf, 2017; McCrickerd and Forde, 2016). During consumption, they synergistically define the sensory experience of food together with other cues (Spence, 2015). Although imaging studies have identified neural correlates of odor-taste interaction in the brain (Small, 2012), mechanistic understanding of chemosensory integration is still lacking.

Drosophila offers an attractive model to address this issue, where peripheral chemosensory systems are well-characterized (Scott, 2018; Wilson, 2013), and feeding behavior can be quantified by measuring the strength of proboscis extension reflex (PER) (Dethier, 1976) and the amount of subsequent food consumption. PER is an initial step of feeding evoked by the presentation of sugar to the proboscis or legs housing gustatory receptor neurons (GRNs). Previous studies reported that odors increase the rate of PER (Oh et al., 2021; Reisenman and Scott, 2019; Shiraiwa, 2008) as well as food intake (Reisenman and Scott, 2019), exemplifying olfactory enhancement of feeding in *Drosophila*. However, aside from the involvement of olfactory receptor neurons (Oh et al., 2021; Shiraiwa, 2008), biological mechanisms underlying this multisensory enhancement remain elusive.

Here we investigated the impact of odors on feeding behavior and found direct contribution of GRNs to odor-evoked PER, odor-taste integration, and odor-driven enhancement of food consumption, indicating that multisensory integration commences at the very periphery.

Results

Odors alone evoke PER

To examine how odors affect PER, we built a behavioral recording setup in which a tastant and multiple odors with diverse innate values (Badel et al., 2016) can be applied to individual, tethered flies (Figure 1A). To quantify the movement of proboscis, we used a deep learning-based, markerless pose estimation algorithm (Mathis et al., 2018) and tracked the trajectory of three segments constituting the proboscis. The angles made between the segments (rostrum, haustellum, and labellum angles) were subsequently calculated over time to detect PER (Figures

1A-1C; see Methods for the definition of PER). The results of this quantification and manual scoring of PER were comparable (Figure1-figure supplement 1C). Unexpectedly, we found that the odors alone evoked repetitive PER without an application of a tastant (Figures 1D-1G, and Figure 1-movie supplement1). Different odors evoked PER with different probability (Figure 1E), latency (Figure1-figure supplement 1A), and duration (Figures 1F, 1G, and Figure1-figure supplement 2). Stimulus-triggered PER was not observed in response to solvents or air (Figure 1D-1G). Odors evoked PER even at the concentration of 10^{-2}, which induce stimulus-specific activity in the olfactory pathway (Badel et al., 2016; Endo et al., 2020; Kato et al., 2023). The level of PER did not correlate with the innate value of odors, previously determined by quantifying the odor preference of flies in a behavioral arena (Badel et al., 2016; Pearson's R = 0.33, p = 0.52). Odor-evoked PER was observed in other genetic backgrounds and another *Drosophila* species as well albeit at a lower level with different tuning (Figure1-figure supplement 3).

Odor-evoked PER resembled taste-evoked PER in multiple aspects. First, its trajectory was similar to that of taste-evoked PER (Figure1-figure supplement 1B, Figure 1-movie supplement1) and was consistent with the reported movement of proboscis during sucrose-evoked PER (Schwarz et al., 2017). The trajectories were indistinguishable between the tested odors (Figure1-figure supplement1B). Second, it showed concentration dependency (Dethier, 1976; Wang et al., 2004). The probability and duration of PER increased as a function of stimulus concentration (Figures 1E and 1G). Third, it depended on the metabolic state just as taste-evoked PER (Inagaki et al., 2014). The duration of PER was higher in starved as compared to fed flies (Figure2-figure supplement 14).

In sum, odors initiate feeding behavior in *Drosophila*.

Odors evoke PER through GRNs

To confirm that this behavior is mediated by the olfactory system, we repeated the experiment after removing the sensory organs housing the olfactory receptor neurons, namely the antennae and the maxillary palps. However, this manipulation only attenuated and did not eliminate odor-evoked PER (Figure 2A), suggesting that the olfactory system modulates but is not required for odor-evoked PER.

We thus turned to another chemosensory system, the gustatory system. To examine its involvement in odor-evoked PER, we expressed Kir2.1(Paradis *et al.*, 2001) in different types of

GRNs using type-specific Gal4 drivers, each of which mediates a specific taste modality (Scott,

2018). Gr5a GRNs detect sugar and mediate food acceptance behaviors including PER

(Dahanukar et al., 2001; Wang et al., 2004). Suppression of Gr5a GRNs significantly reduced

odor-evoked PER, indicating that sweet-sensing neurons detect odors and drive PER (Figures 2B

and 2C). Bitter-sensing Gr66a GRNs, on the other hand, have been shown to inhibit sweet-

evoked PER (Wang et al., 2004). However, expression of Kir2.1 in these neurons had little effect

(Figures 2B and 2D). We hypothesized that this is due to our use of starved flies because Gr66a

pathway is reported to be downregulated in a starved condition (Devineni et al., 2019; Inagaki et

al., 2014). Indeed, flies of the same genotype showed increased PER to odors in a fed state

(Figure 2G), indicating that Gr66a GRNs counteract PER. Two other tastes, water and low salt

can also induce PER (Cameron et al., 2010; Wang et al., 2004), which are detected by Ppk28 and

Ir94e GRNs, respectively (Cameron et al., 2010; Jaeger et al., 2018). However, suppression of

these cell types did not affect odor-evoked PER (Figures 2B, 2E and 2F). Together, Gr5a GRNs

106   enhance whereas Gr66a GRNs inhibit odor-evoked PER.

As GRNs are housed in multiple external organs including the labella at the tip of

proboscis, the legs, the wing margins, and the ovipositor (Scott, 2018), we next sought to narrow

down the GRNs responsible for odor-evoked PER. We found that removal of the legs and wings

did not abolish the behavior (Figure2-figure supplement 2). Moreover, activation of the

ovipositor induces egg laying rather than PER (Dethier, 1976). These results suggest that GRNs

act predominantly in the labella to induce odor-evoked PER.

Gustatory receptors and olfactory binding proteins mediate odor-evoked PER

We have shown that Gr5a GRNs tuned to tastes trigger PER in response to a wide range of

odors. Because GRNs do not express olfactory receptors (Davie et al., 2018; Li et al., 2021; Scott

et al., 2001), either gustatory receptors or some other molecules are likely to interact with

odorants (Jones et al., 2007; Kwon et al., 2007). Gr5a receptor is a primary candidate as Gr5a

GRNs drive odor-evoked PER (Figures 2B and 2C). Consistent with this hypothesis, odor-

evoked PER was nearly undetectable in the Gr5a mutant but spared in the control (Figure 2H).

This demonstrates that Gr5a receptor is necessary for odor-evoked PER.

Odorant binding proteins (OBPs) are another set of molecules that could mediate odor

detection by GRNs. They are thought to bind and help transport hydrophobic odorants to

chemosensory receptors through aqueous lymph (Larter et al., 2016). OBPs are expressed in taste

as well as olfactory sensilla (Galindo and Smith, 2001) and modulate sugar-evoked PER (Swarup et al., 2014), raising the possibility that those in the taste sensilla also mediate odorant detection. To test this, we conducted a genetic screen by expressing individual Obp RNAi constructs with *tubulin-Gal4*. We targeted 9 Obp genes that are expressed highly in the labella (Cameron et al., 2010) but in trace amounts in the antenna (Larter et al., 2016) to examine the function of OBPs in the taste sensilla. Knockdown of *Obp18a*, *Obp57d/e*, and *Obp57e* genes decreased whereas that of *Obp49a* gene increased odor-evoked PER (Figures 2I and 2J, and Figure2-figure supplement 3). Knockdown of *Obp19b, Obp56g*, *Obp56h*, *Obp57a/c*, and *Obp83c* genes did not have a significant effect on odor-evoked PER (Figure 2K, and Figure2-figure supplement 3). The increased PER following *Obp49a* knockdown likely reflects disinhibition of sweet-sensing GRNs as Obp49a mediates the inhibitory impact of bitter chemicals on the activity of sweet-sensing GRNs (Jeong et al., 2013). These results suggest that OBPs modulate odor detection by GRNs.

GRNs directly respond to odors

To gain direct evidence that GRNs respond to odors, we performed two-photon calcium imaging and single-sensillum electrophysiological recording that have complementary properties (Figures 3 and 4). Calcium imaging of GRN axons using a cell-specific driver reports the population activity of neurons but can unambiguously ascribe the signals to genetically identified GRNs. On the other hand, while single-sensillum electrophysiology cannot uniquely ascribe the signals to genetically defined GRNs, it is sensitive and fast enough to detect individual spikes.

For imaging, we expressed a calcium indicator GCaMP6s (Chen et al., 2013) in Gr5a, Gr66a, Ppk28, or Ir94e GRNs and scanned the axons in the subesophageal zone of the brain (Figure 3A). Consistent with the contribution of Gr5a and Gr66a GRNs to odor-evoked PER, robust odor responses were observed in these neurons. They responded to all the tested odors but not to control stimuli, with distinct tuning (Figures 3B-3F, Figure3-figure supplement 1A to F and Figure3-figure supplement 2), in a concentration-dependent manner (Figure3-figure supplement 1G to H and Figure3-figure supplement 3B). A linear combination of Gr5a and Gr66a GRN responses well predicted the magnitude of odor-evoked PER with the former and the latter contributing to enhancement and suppression of PER, respectively (Figure 3G). Odor and sucrose responses were positively correlated in Gr5a GRNs (Figure3-figure supplement 1I). By

contrast, Ppk28 and Ir94e GRNs did not respond to any odors (Figure3-figure supplement 1J and K), matching their little contribution to odor-evoked PER (Figures 2B, 2F and 2G).

These responses could reflect the activity originating in GRNs or that from a central olfactory circuit impinging on GRN's presynaptic terminals. To distinguish between the two scenarios, we physically covered the maxillary palps with UV glue to block the input from the olfactory organs and applied odors (antennae were already removed to gain optical access to the subesophageal zone, see Methods). This manipulation had little effect on the odor responses (Figure 3D and 3F). On the other hand, the responses were eliminated after covering the labella (Figures 3D and 3F), demonstrating that GRNs themselves sense odors.

As we observed that Gr5a receptor was necessary for odor-evoked PER, we further asked whether it is required for Gr5a GRNs to respond to odors. We found that odor responses were absent in the Gr5a mutant but spared in control flies (Figure 3H). This confirms that odor responses in Gr5a GRNs depend on Gr5a receptor.

For electrophysiology, we measured odor-evoked responses from individual taste sensilla on the proboscis using tip recording, a conventional technique where a glass electrode placed on the tip of a sensillum delivers a stimulus as well as records neural responses (Figure 4; Hodgson et al., Science, 1955). We targeted large and intermediate type (L- and I-type) taste sensilla, and recorded responses to odors and sucrose. Odors were dissolved in tricholine citrate (TCC), an electrolyte that suppresses the responses of water-sensing neuron (Wieczorek and Wolff, 1989). L-type sensilla housing a sweet-sensing GRN robustly responded to sucrose, but also to two tested odors, 4-methylcyclohexanol and banana odor, at a rate higher than the solvent (Figures 4A and 4B). I-type sensilla housing sweet- and bitter-sensing GRNs also responded to odors (Figures 4A and 4C). These sensilla exhibited spikes with different amplitudes that may originate in different GRNs. This is consistent with our calcium imaging results showing that both Gr5a and Gr66a GRNs respond to odors (Figure 3). Here we counted all the spikes and did not sort them based on their amplitude and shape, because these characteristics change depending on the spiking frequency (Fujishiro et al., 1984).

In sum, recording from specific types of GRNs using calcium imaging and specific types of taste sensilla using tip recording together demonstrate that GRNs in the labella sense odors in line with a recent study using a different electrophysiological technique (Dweck and Carlson, 2023).

Odor-taste integration in single taste sensillum

We investigated whether GRNs integrate multisensory stimuli when they are presented together. We presented sucrose, banana odor or a mixture of the two to individual taste sensilla during tip recording. We found that the responses to a mixture were stronger than those to component stimuli (Figure 5). While the strength of responses varied across sensilla, responses to banana odor and sucrose recorded in the same sensillum were positively correlated (Figure5-figure supplement 1). These results suggest that GRNs function as an odor-taste multisensory integrator.

GRNs integrate multisensory input to enhance PER and food consumption

We finally asked if GRNs can enhance PER and subsequent food consumption through multisensory integration. We decided to apply a mix of banana odor and sucrose as a mimic of multisensory food stimuli in the natural environment. We confirmed that the banana odor can evoke PER and GRN responses in a concentration-dependent manner (Figure3-figure supplement 3). We then presented sucrose solution mixed with or without a low concentration (10^{-4}) of banana odor locally to the labella (Figure 6A). Although banana odor could not evoke PER on its own at this concentration (Figure3-figure supplement 3A), we found that the addition of banana odor increased PER to sucrose especially at low concentrations where sucrose alone can only induce unreliable PER (Figure 6B). Importantly, this enhancement was observed even after removing the olfactory organs (Figure 6B), indicating that the superadditive integration takes place in GRNs. Similar results were obtained when sucrose was mixed with different monomolecular odorants (Figures 6C and 6D).

We further examined if the amount of food consumption is also enhanced by odor-taste integration in GRNs. Using a previously reported multisensory feeding assay (Reisenman and Scott, 2019), we quantified the amount of sucrose consumed by freely behaving flies with or without odors (Figure 6E). We found that the presence of banana odor increased the amount of sucrose consumption in intact flies (Figure 6F). Critically, this effect was observed in olfactory organs-removed flies as well (Figure 6F), indicating the contribution of the gustatory system. A similar result was observed with ethyl butyrate, and a slight, though not significant, increase was also observed with 4-methylcyclohexanol (Figure 6G). These results suggest that GRNs serve as an initial node for odor-taste multisensory integration that shapes feeding behavior (Figure 6H).

Discussion

In terrestrial animals, it has been considered that there is a clear division of labor between the two types of chemosensory organs in sensing stimuli with distinct physical properties; the olfactory and gustatory organs sense volatile and non-volatile chemicals, respectively. Therefore, although odorants and tastants are defined by the identity of cognate organs, they are used interchangeably with volatile and non-volatile chemicals. However, we found using calcium imaging and electrophysiological recording that GRNs directly detect odorants, and this drives a classical taste-induced feeding behavior, PER, as well as increased food consumption. A previous study has also hinted that odors alone evoke PER (Oh et al., 2021). Thus, volatile chemicals are within the molecular receptive range of GRNs and should also be regarded as tastants based on the conventional definition. Critically, this indicates that GRNs are engaged in non-contact chemo-sensing as well, a behavioral strategy distinct from contact-based chemo-sensing.

A study applying base recording, an electrophysiological technique to record from individual taste sensilla, had reported that I-type sensilla do not respond to odorant molecules dissolved in solution (Hiroi et al., 2008). However, a recent study using the same method found that S-, I-, and L-type sensilla respond to a variety of odors applied as vapor (Dweck and Carlson, 2023). Here we found that GRNs respond to odors applied in both ways in agreement with Dweck and Carlson (2023). Although the reason behind the discrepancy is not entirely clear, one possibility is the difference between the preparation used in different studies; whereas Hiroi et al. used a severed head preparation, we as well as Dweck and Carlson used immobilized, intact flies. Because we noticed in preliminary experiments that stimulus-evoked GRN spikes become weaker over time in a head only preparation where hemolymph circulation is lost, the physiological condition of sensilla may be one of the factors underlying the discrepancy. Another possibility is a difference in genotype or genetic background. We found that the strength of odor-evoked PER is different between genetic backgrounds with w[1118] exhibiting weaker PER than Dickinson wild type (Figure1-figure supplement 3), implying that odor responses of GRNs vary across genetic backgrounds. Furthermore, odor tuning of sugar-sensing GRNs was different between our study and Dweck and Carlson (2023), which used different genetic lines. Although we have observed odor-evoked PER in *Drosophila simulans* as well, further study is required to examine the generality as well as the variability of GRN odor responses across genotypes and species.

Our results showed that odor responses of Gr5a GRNs are mediated by the Gr5a receptor.

This may not be totally enigmatic as gustatory and olfactory receptor genes belong to the same

ancient superfamily sharing a motif in the transmembrane region (Robertson et al., 2003; Scott et

al., 2001), and some gustatory receptors expressed outside of GRNs function in odor (Jones et

al., 2007; Kwon et al., 2007), temperature (Ni et al., 2013), and light detection (Montell, 2021;

Xiang et al., 2010). However, Gr5a is unique in that the same receptor can detect stimuli of two

different sensory modalities. Furthermore, we found that OBPs enriched in the labella modulate

odor-evoked PER, suggesting that OBPs in this taste organ not only link tastants with gustatory

receptors (Jeong et al., 2013; Swarup et al., 2014), but also mediate detection of odors.

PER was evoked by all the tested odors albeit to different extents, and this likely reflects

the broad odor tuning of Gr5a GRNs that drive PER (Fig. 3). The question, then, is why might it

be useful to exhibit PER to such a wide variety of odors including those that are normally

aversive? A likely benefit can be discussed by considering foraging behavior under natural

settings. During foraging in the air, olfactory receptor neurons that are more sensitive to odors

locate a potential food source together with other sensory systems. After landing on the potential

food, GRNs on the legs join to assess its edibility. In parallel, GRNs on the labella sense odors

and enhance the probability of PER. Given that the olfactory system most likely has already

guided the animal to palatable food, keeping the GRNs broadly tuned to odors is an efficient

mechanism to enhance PER. Once PER brings the labella into contact with the food, GRNs act

as a multisensory integrator to make a decision on food intake (Figure 6). Enhancing PER to

aversive odors might also be adaptive as animals often need to carry out the final check by

tasting a trace amount of potentially dangerous substances to confirm that those should not be

consumed. Because inputs from olfactory and gustatory organs interact centrally as well (Oh et

al., 2021; Shiraiwa, 2008) (Figure 2A), this indicates that unified chemosensory experience is

built through layered integration in different parts of the body.

Although odor detection by GRNs through gustatory receptors has not previously been

reported outside of *Drosophila melanogaster*, olfactory receptors are expressed in the proboscis

in multiple insects (Haverkamp et al., 2016; Kwon et al., 2006; Xia and Zwiebel, 2006). A recent

study reported that olfactory receptors are also expressed and functional in cultured human

fungiform and mouse taste papilla cells (Malik et al., 2019), implying that peripheral odor-taste

integration may be a process incorporated in various species. These and our results reveal that

GRNs should be further studied not as gustatory neurons but more broadly as chemosensory
integrators.
285

Acknowledgments

We thank Anupama Dahanukar, Graeme Davis, Michael Dickinson, Craig Montell, Kristin Scott,
the Bloomington Stock Center, and the Vienna Drosophila Resource Center for flies; RIKEN
CBS-Evident Collaboration Center for imaging equipment and software; Bor-Wei Cherng for
help with coding; members of the Kazama laboratory for their support and comments on the
manuscript. HP.W. was supported by Grant-in-Aid for JSPS Research Fellow (JP15F15384) and
RIKEN Special Postdoctoral Fellowship. This work was supported by a grant from RIKEN, Kao
Corporation, Asahi Corporation, Toray (23-6402), Japan Society for the Promotion of Science
(KAKENHI Grant JP18H02532, JP18K19502, JP21H04789), and Japan Science and
Technology Agency (CREST grant number JP24028095) to H.K.

Author contributions

298   HP.W. and H.K. designed the study. HP.W. performed all the experiments and analyzed data
299   except for part of the behavioral data analysis, which was performed by T.K.C.L. HP.W. and
300   H.K. wrote the paper with input from T.K.C.L.
301

Declaration of interests

303   Authors declare that they have no competing interests.
304

Methods

Resource availability

Lead contact

309   Further information and requests for resources should be directed to and will be fulfilled by the
310   Lead Contact, Hokto Kazama (hokto.kazama@riken.jp).
311

Materials availability

313   This study did not generate new unique reagents.
314

Data and code availability

All data reported in this paper will be shared by the lead contact upon request.

All original code has been deposited at GitHub, (https://github.com/hpwei82/2022_hpwei) and will be publicly available as of the date of publication.

Any additional information required to analyze the data reported in this study is available from the Lead Contact upon request.

Experimental model and subject details

Drosophila **strains**

Flies were maintained on standard cornmeal agar under a 12 h light, 12 h dark cycle at 25 °C. All experiments were performed on adult females 2–6 days after eclosion. Flies were obtained as described in the key resources table.

Fly stocks used in this study are as follows: Dickinson wild type (Michael Dickinson), *w^{1118}; P{y[+t7.7] w[+mC]=GAL4.1Uw}attP2* (Bloomington #68384), *w^{1118}; P{y[+t7.7] w[+mC]=p65.AD.Uw}attP40; P{y[+t7.7] w[+mC]=GAL4.DBD.Uw}attP2* (Bloomington #79603), *D.simulans* (Kyoto DRRG #900001), *tubulin-GAL4* (Bloomington #5138 (Lee and Luo, 1999)), *Orco-GAL4* (Bloomington #26818 (Larsson et al., 2004)), *Gr66a-GAL4* (Bloomington #57670 (Kwon et al., 2011)), *Ir94e-GAL4* (Bloomington #60725 (Jaeger et al., 2018)), *Ppk28-GAL4* (Bloomington #93020 (Cameron *et al.*, 2010)), *UAS-GCaMP6s* (Bloomington #42746, 42749 (Chen et al., 2013)), *UAS-Obp49a RNAi* (VDRC #330599 (Dietzl et al., 2007)), *UAS-Obp19b RNAi* (VDRC #1823), *UAS-Obp56g RNAi* (VDRC #23206), *UAS-Obp56h RNAi* (VDRC #102562), *UAS-Obp18a RNAi* (VDRC #101628), *UAS-Obp83c RNAi* (VDRC #106866), *UAS-Obp57d/e RNAi* (VDRC #101783), *UAS-Obp57e RNAi* (VDRC #105001), *UAS-Obp57a/c RNAi* (VDRC #107489), three OBP RNAi control lines *w^{1118}* (VDRC #60000, control for *UAS-Obp19b RNAi* and *UAS-Obp56g RNAi*), *w^{1118};P{VDRCsh60200attP40* (VDRC #60200, control for *UAS-Obp49a RNAi*), *y,w^{1118};P{attP,y[+],w[3']]}* (VDRC #60100, control for other *UAS-Obp RNAi* lines), *Gr5a-GAL4* (Kristin Scott), Gr5a mutant (Anupama Dahanukar (Dahanukar et al., 2007), *ΔEP(x)-5*) and its control (Anupama Dahanukar (Dahanukar et al., 2007), *EP(x)496*), and *UAS-Kir2.1^{AAE}-GFP* (Graeme Davis (Paradis et al., 2001)). All the lines except for *w^{1118}*, *D.simulans*, *Gr5a* mutant, its control, and the UAS-RNAi lines were backcrossed for 6 generations to the Dickinson wild type (Dickinson, 1999).

Detailed genotypes of flies used in each experiment are listed in Table 1.

Figure	Genotype
Figure 1B-1G, 2A, 4, 5, 6, Figure 1-figure supplement 1, Figure 1-figure supplement 2, Figure 2-figure supplement 1, Figure 3-figure supplement 3A	Dickinson Wild type
Figure 1-figure supplement 3A	w^{1118}; P{y[+t7.7] w[+mC]=GAL4.1Uw}attP2
Figure 1-figure supplement 3B	w^{1118}; P{y[+t7.7] w[+mC]=p65.AD.Uw}attP40; P{y[+t7.7] w[+mC]=GAL4.DBD.Uw}attP2
Figure 1-figure supplement 3C	D.simulans
Figure 2B	UAS-Kir2.1/+
Figure 2C	Gr5a-Gal4/UAS-Kir2.1 Gr5a-Gal4/+
Figure 2D	Gr66a-Gal4/UAS-Kir2.1 Gr66a-Gal4/+
Figure 2E	Ppk28-Gal4/UAS-Kir2.1 Ppk28-Gal4/+
Figure 2F	Ir94e-Gal4/UAS-Kir2.1 Ir94e-Gal4/+
Figure 2G	Gr66a-Gal4/UAS-Kir2.1 UAS-Kir2.1/+
Figure 2-figure supplement 2	Orco-Gal4>UAS-Kir2.1
Figure 2H	ΔEP(x)-5 EP(x)496
Figure 2I	tubulin-Gal4/y,w^{1118};P{attP,y[+],w[3']} tubulin-Gal4/UAS-Obp57d/e RNAi
Figure2-figure supplement 3C	tubulin-Gal4/y,w^{1118};P{attP,y[+],w[3']}
Figure 2J	tubulin-Gal4/w^{1118};P{VDRCsh60200}attP40 tubulin-Gal4/UAS-Obp49a RNAi

Figure 2K	*tubulin-Gal4/w^{1118}*
	tubulin-Gal4/UAS-Obp19b RNAi
Figure 2-figure supplement 3A	*tubulin-Gal4/w^{1118}*
Figure 2-figure supplement 3B	*tubulin-Gal4/UAS-Obp56g RNAi*
Figure 2-figure supplement 3D	*tubulin-Gal4/UAS-Obp18a RNAi*
Figure 2-figure supplement 3E	*tubulin-Gal4/UAS-Obp56h RNAi*
Figure 2-figure supplement 3F	*tubulin-Gal4/UAS-Obp83c RNAi*
Figure 2-figure supplement 3G	*tubulin-Gal4/UAS-Obp57e RNAi*
Figure 2-figure supplement 3H	*tubulin-Gal4/UAS-Obp57a/c RNAi*
Figure 3-figure supplement 1J	*UAS-GCaMP6s;Ppk28-Gal4*
Figure 3-figure supplement 1K	*UAS-GCaMP6s;Ir94e-Gal4*
Figure 3A-3D, 3G, Figure 3-figure supplement 1A-C, G, I, Figure3-figure supplement 2A, Figure 3-figure supplement 3B	*Gr5a-Gal4;UAS-GCaMP6s*
Figure 3E, 3F, Figure 3-figure supplement 1D-F, Figure 3-figure supplement 2B	*UAS-GCaMP6s;Gr66a-Gal4*
Figure 3H	*ΔEP(x)-5;Gr5a-Gal4;UAS-GCaMP6s*
	EP(x)496;Gr5a-Gal4;UAS-GCaMP6s

Table 1. Genotypes used to generate data presented in each figure

Examination of odor-evoked PER

Virgin females that had been raised on food for 1-3 days were starved for 24-28 h in vials with water supplied through a wet piece of Kimwipe. Individual flies were briefly anesthetized on ice and their dorsal side of the thorax was attached to a cover glass with ultraviolet-curing adhesive (NOA 63, Norland), after which the flies were allowed to rest for an hour. Prior to recording odor-evoked PER, flies were water satiated until they did not consume any more. Subsequently, 100 mM sucrose solution, which acted as a positive control stimulus, was applied either on the fly's legs or proboscis (without letting the fly consume it) to examine if the fly could exhibit

tastant-evoked PER. Flies were discarded if they did not exhibit PER to this stimulus. Flies

showing excessive spontaneous PER before the assay were also discarded. The tethered fly was

positioned horizontally in air facing an odor delivery tube (Figure 1A) except for the experiment

in Figure S2 where the fly was positioned vertically to mimic the preparation used in a previous

study (Oh et al., 2019). The tip of the odor tube was placed 10 mm away from the fly. A

monochrome camera (Lu070M, Lumenera corporation) taking a lateral view of the fly at 20 Hz

was used to record the movement of proboscis in response to odors.

To examine the contribution of olfactory organs to odor-evoked PER (Figures 2 and 6),

the third antennal segments and maxillary palps were removed with forceps while the flies were

anesthetized on ice. After the surgery, the flies were given an hour to recover before the

experiment.

To examine the contribution of GRNs on wings and legs to odor-evoked PER (Figure

S5), wings and tarsal segments were removed with forceps while the flies were anesthetized on

ice. After the surgery, the flies were given an hour to recover before the experiment.

Individual flies went through an experiment consisting of three blocks. In each block, 6

odors (see below), two solvent controls (mineral oil and water), and another control stimulus

(air) were applied for 2 s per trial in a randomized order with a 15 s inter-trial-interval.

Olfactory stimulation

Odors were delivered with a custom-made, multi-channel olfactometer controlled by a PC as

previously described (Badel et al., 2016). Briefly, an air stream (300 ml/min) was passed through

4 ml of odor solution diluted with mineral oil (nacalai tesque, 23334-85) or water. The

concentration of odor was varied between 10^{-4}, 10^{-2}, 10^{-1}, and 0.5 (v/v), but the default was 0.5

unless otherwise noted. The odorized air was further diluted by mixing it with a main air stream

(500 ml/min), a small portion of which was delivered frontally to the fly through an outlet placed

10 mm away from the fly. The speed of odorized air flow at the position of the fly was 0.6 m/s.

Using the photoionization detector (200B miniPID, Aurora Scientific Inc.), the time odors reach

the position of the fly was estimated to be 1.1 s after the odor valve opening. The odors used in

the study and their abbreviations are as follows: 2-pentanone (2PT, Wako, 13303743), 3-octanol

(OCT, Tokyo chemical industry, O0121), 4-methylcyclohexanol (MCH, Sigma-Aldrich,

153095), banana essence (Narizuka corporation), benzaldehyde (BNZ, Sigma-Aldrich, 418099),

ethyl butyrate (EBR, Sigma-Aldrich, E15701), and isopentyl acetate (IPA, Wako, 016-03646), 1-

hexanol (Tokyo chemical industry, H0130), ethyl acetate (Tokyo chemical industry, A0030),

yeast (nacalai tesque, M1E8921), fenchone (Tokyo chemical industry, F0164), isoamyl alcohol

(Tokyo chemical industry, I0289), phenethyl alcohol (Tokyo chemical industry, P0084).

Examination of multisensory PER

To examine how flies respond to odor-taste multimodal stimuli presented to GRNs in the labella,

individual flies were gently attached to a cover glass vertically by wrapping their thorax with a

piece of parafilm, and multimodal stimuli were applied by touching the ventral part of the

labellum with a wick made of Kimwipe immersed in a sucrose solution with or without odors

(Figure 4A; (Shiraiwa, 2008)). A constant air stream (0.5~0.7 m/s) was applied with an air pump

(eAir6000, GEX) from behind the wick. The legs of the flies were tucked under the parafilm to

prevent them from touching the wick. The flies were allowed to rest for an hour after fixation.

The concentration of sucrose was varied between 0.25-2% (v/v, Figure 6B) or fixed at 0.25%

(Figures 6C and 6D) depending on the experiment. When the concentration of sucrose was

varied, the stimuli were applied in an ascending order to avoid adaptation. The concentration of

the added odor was 10^{-4} for all the tested stimuli. Each stimulus was applied 3 times with an

inter-trial-interval of 15 s. PER was manually scored when the proboscis was fully extended

within ~2 s from the stimulation.

Quantification of PER

A markerless pose estimation algorithm DeepLabCut v2.0 was used to quantify the movement of

the proboscis. The position and orientation of three segments of the proboscis, the rostrum, the

haustellum, and the labellum were characterized by tracking six points, namely the proximal end

of the antenna, the distal end of the antenna, the rostrum apex, the rostrum-haustellum joint, the

haustellum-labellum joint, and the distal end of the labellum (Figures 1B and 1C). These six

points were manually labeled on the lateral view of the fly to generate the training dataset,

which consisted of 1465 labeled frames in video data from 19 flies. The ResNet50 based pose

estimation neural network was trained for about 200,000 iterations, after which the six points

were automatically tracked in all the frames in video data.

Following the pose estimation, data were analyzed using custom code written in Python.

To detect proboscis extensions, the rostrum angle (the angle made by the line passing through the

proximal and distal ends of the antenna and the line along the rostrum), the haustellum angle (the

angle between the rostrum and the haustellum), and the labellum angle (the angle between the

haustellum and the labellum) were calculated over time (see Figure 1B). The baseline of each

proboscis angle was calculated as the 25^{th} percentile of a rolling 5 s time window. The proboscis

extension was defined as PER if the haustellum angle exceeded 100° because it corresponded to

full extension by visual inspection. PER duration was defined as the time during which the fly

exhibited PER in each second. Integrated PER duration was defined as the sum of PER duration

over 4 s starting 1 s after the odor onset considering the time that odor requires to reach the fly

(see earlier section on Olfactory stimulation) and the time period during which the majority of

PER occurred. These values were averaged across three trials for each odor in each fly. PER

probability was defined as the percentage of trials in which PER was observed for each stimulus

in each fly.

Because repetitive, spontaneous PER occurring at regular intervals represented an

abnormal condition, flies showing such repetitive, spontaneous PER in more than 9 trials were

excluded from further analysis.

To examine if the movement of the proboscis is similar between odor-evoked and tastant-

evoked PER as well as between PER evoked by different odors, temporal sequences of the

rostrum and the haustellum angles during PER and partial proboscis extensions were

clustered with K-medoids clustering (Park and Jun, 2009). PER and partial proboscis

extensions were detected from the baseline subtracted rostrum angle using hysteresis

thresholding with a lower threshold of 5° and an upper threshold of 15°. Because the length of a

temporal sequence is different between PER, the data were converted to a distance metric using

dynamic time warping (Mearns et al., 2020; Sakoe and Chiba, 1978) with a warping window of

0.25 s prior to clustering. The optimal number of clusters was determined using the elbow

method.

Immunohistochemistry

To examine the expression pattern of Gal4 driver lines, we performed immunohistochemistry as

described previously (Badel et al., 2016) using rat anti-GFP (1:1,000, nacalai tesque, 04404-84)

and mouse nc82 (1:20, Developmental Studies Hybridoma Bank at Univ. of Iowa) as primary

antibodies, and anti-rat CF488A (1:250, Biotium, 20023) and anti-mouse CF633 (1:250,

Biotium, 20120) as secondary antibodies. Brains were dissected out from the head capsule in

phosphate-buffered saline (PBS, nacalai tesque, 2757531), fixed with 4% paraformaldehyde in

PBS (nacalai tesque, 915414) for 90 min on ice, and incubated in blocking solution containing

5% normal goat serum (Invitrogen, 50197Z) in PBST (0.2% Triton X-100 (nacalai tesque,

3550102) in PBS) for 30 min. Primary antibodies were then added and incubated at 4 °C for ~48

h. After removing antibodies and washing over an hour, the brains were incubated in a solution

of secondary antibodies at 4 °C for 24 h. The brains were immersed in Vectashield (Vector

laboratories, H-1000), sealed with a cover glass, and imaged with a confocal microscope

(FV3000, Olympus). Images were analyzed with Fiji (Schindelin et al., 2012).

Fly preparation for calcium imaging

Individual flies starved for 24-28 h with water were anesthetized on ice, and their dorsal side of

the thorax was fixed to a piece of parafilm with ultraviolet-curing adhesive (NOA 63, Norland).

The legs of the fly were covered with another piece of parafilm. The fly was subsequently

attached to a custom-made recording plate in a vertical position such that the anterior part of the

head was accessible through a small hole on the recording plate (Figure 3A). The proboscis was

pulled out gently with forceps and immobilized in an extended position using a strip of parafilm

with the labella exposed to the air. After covering the head with saline containing 103 mM NaCl,

3 mM KCl, 5 mM N-tris (hydroxymethyl) methyl-2-aminoethane-sulfonic acid, 8 mM trehalose,

10 mM glucose, 26 mM $NaHCO_3$, 1 mM NaH_2PO_4, 1.5 mM $CaCl_2$, and 4 mM $MgCl_2$

(osmolarity adjusted to 270-275 mOsm), the antennae and the associated cuticle were removed to

expose the subesophageal zone. Saline was bubbled with 95% O_2/5% CO_2 and perfused at a rate

of 2 ml/min during the recording. To examine the contribution of ORNs in the maxillary palps

and GRNs in the labellum, each of these sensory organs were covered with UV curable glue

(Figures 3D and 3F).

Two-photon calcium imaging

Calcium imaging of GRN axons in the subesophageal zone was conducted using a two-photon

microscope (LSM 7 MP, Zeiss) equipped with a piezo motor (P-725.2CD PIFOC, PI) that drives

a water immersion objective lens (W Plan-Apochromat, 20x, numerical aperture 1.0) along the z-

axis. The fluorophore was excited with a titanium:sapphire pulsed laser (Chameleon Vision II,

Coherent) mode-locked at 930 nm. The laser power measured at the back aperture of the

objective lens was below 20 mW. Fluorescence was collected with a GaAsP detector through a

bandpass emission filter (BP470-550). Five optical slices separated by 5 μm were scanned every

500 ms. The odor delivery system was identical to that used in behavioral experiments. Sucrose

was presented by a syringe whose movement was controlled by an actuator.

Image processing

Calcium imaging data were analyzed using custom code written in MATLAB (MathWorks).

Images were registered within and across trials to correct for movement in the x-y plane as well

as in depth by determining the shift along three dimensions that maximizes the correlation

between the images. The region of interest (ROI) was set to cover the GRN axons in the

subesophageal zone. The size of ROI was 80×60 μm for Gr5a GRNs, 60×40 μm for Gr66a

GRNs, 80×60 μm for Ir94e GRNs, and 80×40 μm for Ppk28 GRNs, respectively. The average

pixel intensity within the ROI was calculated for each time frame. The average of 5 frames

preceding a stimulus was used as the baseline signal to calculate $\Delta F/F$ for each time frame. The

peak stimulus response was quantified by averaging $\Delta F/F$ across five frames at the peak,

followed by averaging across three trials for each stimulus. Odor stimulation began at frame 11,

and the frames used for peak quantification were 12 to 16.

Quantifying the relationship between GRN activity and PER

A linear model in Figure 3G was generated using Python scikit-learn to quantify the relationship

between GRN odor responses and PER. The weights for Gr5a and Gr66a GRNs were estimated

using the LinearRegression function. The performance of the model was quantified by

calculating the coefficient of determination.

Electrophysiological recording from taste sensilla on the proboscis

Odor and taste responses of individual taste sensilla on the proboscis were measured using tip

recording (Hodgson et al., 1955). Virgin females that had been raised on food for 1-2 days were

starved for 24-28 h in vials with water supplied through a wet piece of Kimwipe. Individual flies

were immobilized at the end of a pipette tip. The proboscis was fixed at an extended position

using ultraviolet-curing adhesive (NOA 63, Norland). A saline-filled reference electrode was

inserted into the eye. Individual taste sensilla were stimulated for ~2 s with a capillary electrode

containing the stimulus and an electrolyte, 30 mM TCC. The recording electrode was connected

to an amplifier (Multiclamp 700B, Molecular Devices). Signals were bandpass filtered at 100-

1000 Hz, and digitally sampled at 10 kHz using a data acquisition module (NI cDAQ9178, NI

9215, National Instruments). The intensity of the response was measured by counting the number

of spikes generated between 200 and 700 ms after the stimulus contact following the convention

to avoid the contamination of motion artifact (Dahanukar and Benton, 2023) (Delventhal et al.,

2014) (Hiroi et al., 2002). Responses to sucrose, banana odor, MCH, and TCC were recorded

from L- and I-type sensilla. Sucrose and odors were dissolved in TCC (30 mM) solution. Each

stimulus was applied to the same sensillum for 2 or 3 times, with 1 min inter-stimulus-interval.

To examine odor-taste integration in each sensillum, banana odor and sucrose solution were

presented individually or simultaneously in a mixture. Recordings were made from L2 and L6

sensilla. Each stimulus was presented twice, with 1 min inter-stimulus-interval, in a random

order.

Food consumption assay

Food consumption was quantified in principle as described in a previous study (Reisenman and

Scott, 2019). Briefly, virgin females that had been raised on food for 1-2 days were starved for

24-28 h in vials with water supplied through a wet piece of Kimwipe. Flies were then transferred

to a custom-made vial (n = 10 to 15 per vial) with a filter paper (3 cm in diameter) on the top and

a small mesh tube attached laterally near the top (Figure 6E). The filter paper on the top was

impregnated with sucrose solution (0.1%, 150 μl) containing blue dye blue, kyoritsu-foods), and

a small piece of filter paper (0.5 x 1 cm) placed in the mesh tube was impregnated with odor

(1%, 10 μl) or control solution. Flies were allowed to consume sucrose solution for 5 min after

which the vials were placed in the freezer (-20 $^{\circ}$C) for at least 1 h. The extent of sucrose

consumption by individual flies was quantified under the microscope based on the amount of

blue dye in the abdomen, which was scored using the five-point scale ranging from 0 to 2

(Reisenman and Scott, 2019). The scorer was blind to the experimental condition (whether the

filter paper in the mesh tube was impregnated with odor or control solvent). For the examination

of olfactory organs-removed flies, their third antennal segments and maxillary palps were

removed with forceps before starvation.

Statistical analysis

Statistical analyses were preformed using Python (3.8.8) or R (4.0.2). The statistical tests used,

and the sample sizes are listed in figure legends and Table 2. We predetermined sample sizes not

using any statistical methods but based on effect sizes and sample-by-sample variability

observed in pilot experiments.

Supplemental information

Ten supplementary figures, one table and one movie.

References

Badel, L., Ohta, K., Tsuchimoto, Y., and Kazama, H. (2016). Decoding of Context-Dependent

Olfactory Behavior in Drosophila. Neuron *91*, 155-167. 10.1016/j.neuron.2016.05.022.

Boesveldt, S., and de Graaf, K. (2017). The Differential Role of Smell and Taste For Eating

Behavior. Perception *46*, 307-319. 10.1177/0301006616685576.

Cameron, P., Hiroi, M., Ngai, J., and Scott, K. (2010). The molecular basis for water taste in

Drosophila. Nature *465*, 91-95. 10.1038/nature09011.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr,

R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for imaging

neuronal activity. Nature *499*, 295-300. 10.1038/nature12354.

Dahanukar, A., and Benton, R. (2023). Recording from Fly Taste Sensilla. Cold Spring Harb Protoc

2023, pdb prot108064. 10.1101/pdb.prot108064.

Dahanukar, A., Foster, K., van der Goes van Naters, W.M., and Carlson, J.R. (2001). A Gr receptor

is required for response to the sugar trehalose in taste neurons of Drosophila. Nat Neurosci *4*, 1182-

1186. 10.1038/nn765.

Dahanukar, A., Lei, Y.T., Kwon, J.Y., and Carlson, J.R. (2007). Two Gr genes underlie sugar

reception in Drosophila. Neuron *56*, 503-516. 10.1016/j.neuron.2007.10.024.

Davie, K., Janssens, J., Koldere, D., De Waegeneer, M., Pech, U., Kreft, L., Aibar, S., Makhzami,

S., Christiaens, V., Bravo Gonzalez-Blas, C., et al. (2018). A Single-Cell Transcriptome Atlas of

the Aging Drosophila Brain. Cell *174*, 982-998 e920. 10.1016/j.cell.2018.05.057.

Delventhal, R., Kiely, A., and Carlson, J.R. (2014). Electrophysiological Recording From

Labellar Taste Sensilla. Jove-J Vis Exp. ARTN e51355

10.3791/51355.

Dethier, V.G. (1976). The hungry fly: A physiological study of the behavior associated with feeding.

Devineni, A.V., Sun, B., Zhukovskaya, A., and Axel, R. (2019). Acetic acid activates distinct taste

pathways in Drosophila to elicit opposing, state-dependent feeding responses. Elife *8*.

10.7554/eLife.47677.

Dickinson, M.H. (1999). Haltere-mediated equilibrium reflexes of the fruit fly, Drosophila

melanogaster. Philos Trans R Soc Lond B Biol Sci *354*, 903-916. 10.1098/rstb.1999.0442.

Dietzl, G., Chen, D., Schnorrer, F., Su, K.C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K.,

Oppel, S., Scheiblauer, S., et al. (2007). A genome-wide transgenic RNAi library for conditional

gene inactivation in Drosophila. Nature *448*, 151-156. 10.1038/nature05954.

Dweck, H.K.M., and Carlson, J.R. (2023). Diverse mechanisms of taste coding in Drosophila. Sci

Adv *9*. ARTN eadj7032

10.1126/sciadv.adj7032.

Endo, K., Tsuchimoto, Y., and Kazama, H. (2020). Synthesis of Conserved Odor Object

Representations in a Random, Divergent-Convergent Network. Neuron *108*, 367-381 e365.

10.1016/j.neuron.2020.07.029.

Fujishiro, N., Kijima, H., and Morita, H. (1984). Impulse Frequency and Action-Potential

Amplitude in Labellar Chemosensory Neurons of Drosophila-Melanogaster. J Insect Physiol *30*,

317-325. Doi 10.1016/0022-1910(84)90133-1.

Galindo, K., and Smith, D.P. (2001). A large family of divergent Drosophila odorant-binding

proteins expressed in gustatory and olfactory sensilla. Genetics *159*, 1059-1072.

10.1093/genetics/159.3.1059.

Haverkamp, A., Yon, F., Keesey, I.W., Mißbach, C., Koenig, C., Hansson, B.S., Baldwin, I.T.,

Knaden, M., and Kessler, D. (2016). Hawkmoths evaluate scenting flowers with the tip of their

proboscis. Elife *5*, e15039.

Hiroi, M., Marion-Poll, F., and Tanimura, T. (2002). Differentiated response to sugars among

labellar chemosensilla in. Zool Sci *19*, 1009-1018. DOI 10.2108/zsj.19.1009.

Hodgson, E.S., Lettvin, J.Y., and Roeder, K.D. (1955). Physiology of a Primary Chemoreceptor

Unit. Science *122*, 417-418. DOI 10.1126/science.122.3166.417-a.

Inagaki, H.K., Panse, K.M., and Anderson, D.J. (2014). Independent, reciprocal neuromodulatory

control of sweet and bitter taste sensitivity during starvation in Drosophila. Neuron *84*, 806-820.

10.1016/j.neuron.2014.09.032.

Jaeger, A.H., Stanley, M., Weiss, Z.F., Musso, P.Y., Chan, R.C., Zhang, H., Feldman-Kiss, D., and

Gordon, M.D. (2018). A complex peripheral code for salt taste in Drosophila. Elife *7*.

10.7554/eLife.37167.

Jeong, Y.T., Shim, J., Oh, S.R., Yoon, H.I., Kim, C.H., Moon, S.J., and Montell, C. (2013). An odorant-binding protein required for suppression of sweet taste by bitter chemicals. Neuron *79*, 725-737. 10.1016/j.neuron.2013.06.025.

Jones, W.D., Cayirlioglu, P., Kadow, I.G., and Vosshall, L.B. (2007). Two chemosensory receptors together mediate carbon dioxide detection in Drosophila. Nature *445*, 86-90. 10.1038/nature05466.

Kato, A., Ohta, K., Okanoya, K., and Kazama, H. (2023). Dopaminergic neurons dynamically update sensory values during olfactory maneuver. Cell Rep *42*. ARTN 113122
10.1016/j.celrep.2023.113122.

Kwon, H.W., Lu, T., Rutzler, M., and Zwiebel, L.J. (2006). Olfactory responses in a gustatory organ of the malaria vector mosquito Anopheles gambiae. Proc Natl Acad Sci U S A *103*, 13526-13531. 10.1073/pnas.0601107103.

Kwon, J.Y., Dahanukar, A., Weiss, L.A., and Carlson, J.R. (2007). The molecular basis of CO2 reception in Drosophila. Proc Natl Acad Sci U S A *104*, 3574-3578. 10.1073/pnas.0700079104.

Kwon, J.Y., Dahanukar, A., Weiss, L.A., and Carlson, J.R. (2011). Molecular and cellular organization of the taste system in the Drosophila larva. J Neurosci *31*, 15300-15309. 10.1523/JNEUROSCI.3363-11.2011.

Larsson, M.C., Domingos, A.I., Jones, W.D., Chiappe, M.E., Amrein, H., and Vosshall, L.B. (2004). Or83b encodes a broadly expressed odorant receptor essential for Drosophila olfaction. Neuron *43*, 703-714. 10.1016/j.neuron.2004.08.019.

Larter, N.K., Sun, J.S., and Carlson, J.R. (2016). Organization and function of Drosophila odorant binding proteins. Elife *5*. 10.7554/eLife.20242.

Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. Neuron *22*, 451-461. 10.1016/s0896-6273(00)80701-1.

Li, H., Janssens, J., De Waegeneer, M., Kolluru, S.S., Davie, K., Gardeux, V., Saelens, W., David, F., Brbić, M., and Leskovec, J. (2021). Fly Cell Atlas: a single-cell transcriptomic atlas of the adult fruit fly. bioRxiv.

Malik, B., Elkaddi, N., Turkistani, J., Spielman, A.I., and Ozdener, M.H. (2019). Mammalian Taste Cells Express Functional Olfactory Receptors. Chem Senses *44*, 289-301. 10.1093/chemse/bjz019.

Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci *21*, 1281-1289. 10.1038/s41593-018-0209-y.

McCrickerd, K., and Forde, C.G. (2016). Sensory influences on food intake control: moving

beyond palatability. Obes Rev *17*, 18-29. 10.1111/obr.12340.

Mearns, D.S., Donovan, J.C., Fernandes, A.M., Semmelhack, J.L., and Baier, H. (2020). Deconstructing Hunting Behavior Reveals a Tightly Coupled Stimulus-Response Loop. Curr Biol *30*, 54-69 e59. 10.1016/j.cub.2019.11.022.

Montell, C. (2021). Drosophila sensory receptors-a set of molecular Swiss Army Knives. Genetics *217*, 1-34. 10.1093/genetics/iyaa011.

Ni, L., Bronk, P., Chang, E.C., Lowell, A.M., Flam, J.O., Panzano, V.C., Theobald, D.L., Griffith, L.C., and Garrity, P.A. (2013). A gustatory receptor paralogue controls rapid warmth avoidance in Drosophila. Nature *500*, 580-584. 10.1038/nature12390.

Oh, S.M., Jeong, K., Seo, J.T., and Moon, S.J. (2021). Multisensory interactions regulate feeding behavior in Drosophila. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2004523118.

Paradis, S., Sweeney, S.T., and Davis, G.W. (2001). Homeostatic control of presynaptic release is triggered by postsynaptic membrane depolarization. Neuron *30*, 737-749. 10.1016/s0896-6273(01)00326-9.

Park, H.-S., and Jun, C.-H. (2009). A simple and fast algorithm for K-medoids clustering. Expert systems with applications *36*, 3336-3341.

Reisenman, C.E., and Scott, K. (2019). Food-derived volatiles enhance consumption in Drosophila melanogaster. J Exp Biol *222*. 10.1242/jeb.202762.

Robertson, H.M., Warr, C.G., and Carlson, J.R. (2003). Molecular evolution of the insect chemoreceptor gene superfamily in Drosophila melanogaster. Proc Natl Acad Sci U S A *100 Suppl 2*, 14537-14542. 10.1073/pnas.2335847100.

Sakoe, H., and Chiba, S. (1978). Dynamic programming algorithm optimization for spoken word recognition. IEEE transactions on acoustics, speech, and signal processing *26*, 43-49.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

Schwarz, O., Bohra, A.A., Liu, X., Reichert, H., VijayRaghavan, K., and Pielage, J. (2017). Motor control of Drosophila feeding behavior. Elife *6*. 10.7554/eLife.19892.

Scott, K. (2018). Gustatory Processing in Drosophila melanogaster. Annu Rev Entomol *63*, 15-30. 10.1146/annurev-ento-020117-043331.

Scott, K., Brady, R., Jr., Cravchik, A., Morozov, P., Rzhetsky, A., Zuker, C., and Axel, R. (2001). A chemosensory gene family encoding candidate gustatory and olfactory receptors in Drosophila.

Cell *104*, 661-673. 10.1016/s0092-8674(01)00263-x.

Shiraiwa, T. (2008). Multimodal chemosensory integration through the maxillary palp in

Drosophila. PLoS One *3*, e2191. 10.1371/journal.pone.0002191.

Small, D.M. (2012). Flavor is in the brain. Physiol Behav *107*, 540-552.

10.1016/j.physbeh.2012.04.011.

Spence, C. (2015). Multisensory flavor perception. Cell *161*, 24-35. 10.1016/j.cell.2015.03.007.

Swarup, S., Morozova, T.V., Sridhar, S., Nokes, M., and Anholt, R.R. (2014). Modulation of

feeding behavior by odorant-binding proteins in Drosophila melanogaster. Chem Senses *39*, 125-

132. 10.1093/chemse/bjt061.

Wang, Z., Singhvi, A., Kong, P., and Scott, K. (2004). Taste representations in the Drosophila brain.

Cell *117*, 981-991. 10.1016/j.cell.2004.06.011.

Wieczorek, H., and Wolff, G. (1989). The Labellar Sugar Receptor of Drosophila. J Comp Physiol

A *164*, 825-834. Doi 10.1007/Bf00616754.

Wilson, R.I. (2013). Early olfactory processing in Drosophila: mechanisms and principles. Annu

Rev Neurosci *36*, 217-241. 10.1146/annurev-neuro-062111-150533.

Xia, Y., and Zwiebel, L.J. (2006). Identification and characterization of an odorant receptor from

the West Nile virus mosquito, Culex quinquefasciatus. Insect Biochem Mol Biol *36*, 169-176.

10.1016/j.ibmb.2005.12.003.

Xiang, Y., Yuan, Q., Vogt, N., Looger, L.L., Jan, L.Y., and Jan, Y.N. (2010). Light-avoidance-

mediating photoreceptors tile the Drosophila larval body wall. Nature *468*, 921-926.

10.1038/nature09576.

Figure 1. Odors alone evoke PER

(**A**) Schematic for recording odor-evoked PER. (**B**) Six points on the head tracked by

DeepLabCut. The movement of the proboscis was characterized by the positions of three

segments of the proboscis and the angles between the segments, namely the rostrum angle (R),

haustellum angle (H), and labellum angle (L). (**C**) Example odor-evoked PER. Top, images of a

proboscis in a retracted (0.2 s), fully extended (3.1 s) and partially extended (5.5 s) state. Bottom,

the three angles over time. Green bar indicates an odor application period. Odor is ethyl butyrate.

(**D**) PER to nine odors in an example fly. Each row corresponds to a trial and each tick mark

indicates the timing of PER. The odors used are as follows: ethyl butyrate (EBR), 2-pentanone

(2PT), isopentyl acetate (IPA), 4-methylcyclohexanol (MCH), benzaldehyde (BNZ), 3-octanol

(OCT) and mineral oil (MO). (**E**) PER probability in wild type flies in response to different

concentrations of odors. The same stimulus was used for air, mineral oil, and water controls. n =

21, 26, 27, and 28 flies for 10^{-4}, 10^{-2}, 10^{-1}, and 0.5 concentration groups, respectively. Error bar,

standard error of the mean. PER probability was different between odor concentrations and

identities (p = 3.0e-05 and 2.0e-7 for odor concentration and identity factors, Scheirer–Ray–Hare

test). (**F**) PER duration in wild type flies. The black lines indicate the mean and the shaded areas

indicate standard error of the mean. n = 28 flies. (**G**) Integrated PER duration in wild type flies in

response to different concentrations of odors. The data is from the same flies as in **E**. Integrated

PER duration was different between odor concentrations and identities (p = 3.6e-13 and 3.5e-11

for odor concentration and odor identity factors, Scheirer–Ray–Hare test).

Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range

(bar).

Figure 2. Odors evoke PER through GRNs

(**A**) Integrated PER duration in intact (n = 28) and olfactory organs removed (n = 32) wild type flies. Flies with olfactory organs removed show less PER compared to control flies (p = 0.0079 and 0.10 for group and odor factors, Scheirer–Ray–Hare test). (**B-F**) Integrated PER duration in UAS control (*UAS-Kir2.1/+*, n = 20, **B**), GRN silenced (Gr5a-Gal4>UAS-Kir2.1, n = 25 for **C**; Gr66a-Gal4>UAS-Kir2.1, n = 24 for **D**; Ppk28-Gal4>UAS-Kir2.1, n = 28 for **E**; Ir94e-Gal4>UAS-Kir2.1, n = 23 for **F**) and Gal4 control flies (Gr5a-Gal4/+, n = 31 for **C**; Gr66a-Gal4/+, n = 21 for **D**; Ppk28-Gal4/+, n = 28 for **E**; Ir94e-Gal4/+, n = 30 for **F**). Each silenced line was compared to the UAS control in **B** (p = 4.9e-5, 0.49, 0.17, 0.11 and 0.12, 0.0014, 0.55, 1.2e-5 for genotype and odor factors for **C-F**, Scheirer–Ray–Hare test). Data were also compared between each silenced line and its Gal4 control (p = 1.0e-15, 0.28, 0.00062, 0.14 and 0.71, 0.0012, 0.56, 5.6e-9 for genotype and odor factors for **C-F**, Scheirer–Ray–Hare test). (**G**) Integrated PER duration in control (*UAS-Kir2.1/+*, n = 21) and Gr66a GRN silenced (Gr66a-Gal4>UAS-Kir2.1, n = 21) flies in a fed state. Silencing of Gr66a GRNs enhances odor-evoked PER in a fed state (p = 1.8e-4 and 0.041 for genotype and odor factors, Scheirer–Ray–Hare test). Odor concentration was 10^{-1}. (**H**) Integrated PER duration in control (n = 29) and Gr5a mutant (n = 29) flies. PER is severely reduced in mutant flies (p = 1.1e-4 and 0.32 for genotype and odor factors, Scheirer–Ray–Hare test). Control: *EP(x)496* flies. Mutant: *ΔEP(x)-5* flies. (**I-K**) Integrated PER duration in control and Obp RNAi flies. Obp57d/e RNAi flies show reduced (**I**, n = 33 and 25 for control and RNAi, p = 0.039 and 0.48 for genotype and odor factors, Scheirer–Ray–Hare test), Obp49a RNAi flies show enhanced (**J**, n = 24 and 20 for control and RNAi, p = 0.0066 and 0.16 for genotype and odor factors, Scheirer–Ray–Hare test), and Obp19b RNAi flies show similar level of PER (**K**, n = 24 and 21 for control and RNAi, p = 0.24 and 0.26 for genotype and odor factors, Scheirer–Ray–Hare test) as compared to their respective controls, which are *tubulin-Gal4/y,w^{1118};P{attP,y[+],w[3']}* (for **I**), *tubulin-Gal4/w^{1118};P{VDRCsh60200}attP40* (for **J**), and *tubulin-Gal4/w^{1118}* (for **K**). Box plots indicate the median (gray and red lines), mean (black and red dots), quartiles (box), and 5-95% range (bar).

Figure 3. Sweet- and bitter-sensing GRNs directly respond to odors

(**A**) Schematic for two photon calcium imaging of GRN axon termini in the subesophageal zone.

The top image shows an anterior view of a brain of Gr5a-Gal4>UAS-GCaMP6s fly. White

rectangle indicates the target region of calcium imaging. Scale bar: 50 μm. (**B**) Example response

(ΔF/F) to ethyl butyrate. Green bar indicates an odor application period. Odor concentration was

10^{-1} for all the odors in all the experiments described in this figure. (**C**) Summary responses of

Gr5a GRNs to nine odors. Green bar indicates an odor application period. ΔF/F of GCaMP6s

fluorescence is color coded according to the scale bar (n = 5). (**D**) Trial averaged peak responses

of Gr5a GRNs to individual odors. Covering the labella with glue reduced odor responses as

compared to control (n = 5 and 4 for control and labella covered, p = 2.6e-15 and 5.1e-3 for

organ condition and odor factors, mixed two-way ANOVA) but covering the maxillary palps had

no effects (n = 5 and 4 for control and palps covered, p = 0.69 and 0.055 for organ condition and

odor factors, mixed two-way ANOVA). (**E, F**) Same as in **C, D**, but for Gr66a GRNs. As in

Gr5a GRNs, covering the labella with glue reduced odor responses as compared to control (n = 6

and 5 for control and labella covered, p = 3.0e-16 and 1.1e-7 for organ condition and odor

factors, mixed two-way ANOVA) but covering the maxillary palps had no effects (n = 6 and 5

for control and palps covered, p = 0.48 and 4.2e-11 for organ condition and odor factors, mixed

two-way ANOVA). (**G**) A linear combination of Gr5a and Gr66a GRN responses well predicted

the magnitude of odor-evoked PER. PER was measured in Gr5a-Gal4>UAS-GCaMP flies. Each

dot represents an odor. Coefficient of determination = 0.81. Weights for Gr5a and Gr66a GRNs

are 5.0e-4 and −2.3e-5, indicating that the former and the latter contributes to enhancement and

suppression of PER, respectively. (**H**) Odor responses are severely reduced in Gr5a GRNs in a

Gr5a receptor mutant background (n = 5 and 8 for control and mutant, p = 8.0e-06 and 0.037 for

genotype and odor factors, mixed two-way ANOVA).

Box plots indicate the median (gray and red lines), mean (black and red dots), quartiles (box),

and 5-95% range (bar).

Figure 4. Individual taste sensilla sense odors

(**A**) Example responses of individual taste sensilla recorded using tip recording. Left, example responses to TCC (30 mM), MCH (1%), banana odor (1%), and sucrose (100 mM) recorded in the same L-type sensillum; Right, example responses to the same set of stimuli in the same I-type sensillum. Traces show the activity between 200 and 700 ms after the stimulus onset. Each gray dot indicates a spike. Bar indicates 50 ms. (**B**) Average responses of L-type sensilla. (n = 20, 6, 17, 20 flies for TCC, MCH, banana odor, and sucrose, respectively). Responses to stimuli are significantly larger than those to TCC (p = 3.0e-18, one-way ANOVA followed by Tukey's HSD, p = 0.0051, 0.0021, and 2.7e-7 for MCH, banana, and sucrose compared to TCC). (**C**) Average responses of I-type sensilla. (n = 30, 9, 22, 30 flies for TCC, MCH, banana odor, and sucrose, respectively). Responses to stimuli are significantly larger than those to TCC (p = 1.4e-12, one-way ANOVA followed by Tukey's HSD, p = 3.9e-6, 2.0e-7, and 5.9e-12 for MCH, banana, and sucrose compared to TCC).

Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).

Figure 5. Odor-taste integration in single taste sensillum

(**A**) Example responses of an L-type taste sensillum to sucrose (0.25%, top), banana odor (0.1%, middle) and the mixture of the two (bottom). Traces show the activity between 200 and 700 ms after the stimulus onset. Each gray dot indicates a spike. Bar indicates 50 ms. (**B**) Average responses of L-type sensilla. Responses to the mixture are significantly larger than those to components (n = 14 flies, p = 0.001, repeated measures ANOVA followed by paired t-tests with Bonferroni correction, p = 0.0030 and 0.0032 for sucrose and banana compared to mixture). Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).

Figure 6. GRNs integrate multisensory input to enhance PER and food consumption

(**A**) Schematic for examining PER in response to odor-taste multimodal stimuli. Stimuli are presented to the labellum with a wick (arrowhead) immersed in a sucrose solution with or without odors. A constant air stream was applied from behind the wick. (**B**) PER probability in response to sucrose at different concentrations with (green) or without (black) banana odor in

wild type flies. Odor concentration was 10^{-4}. Banana odor enhances sucrose-evoked PER not

only in intact flies (n = 13 and 11 for control and banana odor, p = 4.6e-4 and 2.1e-7 for odor

identity and sucrose concentration factors, Scheirer–Ray–Hare test) but also in flies without

olfactory organs (n = 11 and 11 for control and banana odor, p = 0.035 and 0.0070 for odor

identity and sucrose concentration factors, Scheirer–Ray–Hare test). Error bar, standard error of

the mean. (**C**) PER probability in response to sucrose is increased with EBR in both intact and

olfactory organs-removed flies (n = 25 and 25 for intact and olfactory organs-removed flies, p =

0.0052 and 0.0068 for intact and olfactory organ-removed flies, Wilcoxon signed-rank test). (**D**)

Same as in C, but for MCH (n = 21 and 36, p = 0.0072 and 0.00074 for intact and olfactory

organs-removed flies, Wilcoxon signed-rank test). The concentration of odors and sucrose was

10^{-4} and 0.25% for **C** and **D**. Error bar, standard error of the mean. (**E**) Schematic for quantifying

sucrose consumption in the presence or absence of odors. A filter paper on the top was

impregnated with sucrose solution containing blue dye and a small piece of filter paper placed in

the mesh tube near the top was impregnated with odor or control solution (see Methods). (**F**) The

amount of sucrose consumed by flies with or without the presence of banana odor was quantified

with a feeding score (see Methods). The amount of feeding was significantly larger in the

presence of odor for both intact (n = 20 and 22 for control and banana odors, p = 0.005,

Wilcoxon signed-rank test) and olfactory organs-removed flies (n = 13 and 15 for control and

banana odors, p = 0.007, Wilcoxon signed-rank test). (**G**) The same as in **F** but with EBR and

MCH odors and only for olfactory organs-removed flies (EBR experiment, n = 16 and 17 for

control and EBR, p = 0.03, Wilcoxon signed-rank test; MCH experiment, n = 11 and 14 for

control and MCH, p = 0.24, Wilcoxon signed-rank test). (**H**) Summary of mechanisms

underlying PER and food intake in a multisensory environment.

Error bars are standard error of the mean in bar plots. Box plots indicate the median (gray line),

mean (black dot), quartiles (box), and 5-95% range (bar).

Figure 1-figure supplement 1. Characteristics of PER in wild type flies

(**A**) Latency to the first odor-evoked PER in wild type flies. Odor concentration was 0.5 (n = 28 flies). Green bar indicates the odor application period. p = 1.0e-4, one-way ANOVA. Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar). (**B**) K-medoid clustering of odor- and sucrose-evoked proboscis extensions based on the rostrum and haustellum angles over time in wild type flies. All the trajectories during PER and partial proboscis extensions were clustered into five clusters (color coded), and the representative trajectory closest to each cluster medoid was plotted for each stimulus. The green trajectories represent full extensions that qualify the definition of PER (see Methods). Note that these green trajectories are observed in response to sucrose and all the odors (except for the controls), suggesting that the movement of proboscis during odor-evoked PER is similar to that during sucrose-evoked PER. (**C**) Integrated PER duration in response to different odors in wild type flies analyzed by DeepLabCut or manual scoring. The values measured by the two methods were similar (p = 0.94 and 5.5e-3 for scoring method and odor identity factors, Scheirer-Ray-Hare test). Odor concentration is 0.1.

Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).

853
854
Figure 1-figure supplement 2. PER evoked by additional odors
Integrated PER duration in wild type flies in response to 1-hexanol, ethyl acetate, yeast odor,
fenchone, isoamyl alcohol, phenethyl alcohol, air, mineral oil and water. In this experiment, flies
were fixed in a vertical position following a protocol in a previous study (Oh et al., 2021). Box
plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).
860
861

862
863
Figure 1-figure supplement 3. **Odor-evoked PER in flies with different genetic backgrounds and in another species**

866 Integrated PER duration in different flies (n = 20, 28 and 25 in A, B and C). Box plots indicate
867 the median (gray and red lines), mean (black and red dots), quartiles (box), and 5-95% range
868 (bar).
869
870
871
872

Figure 2-figure supplement 1. Dependence of odor-evoked PER on a feeding state

(**A**) Integrated PER duration in starved (n = 28) and fed (n = 19) wild type flies (p = 0.88 and 5.6e-3 for state and odor factors, Scheirer–Ray–Hare test). Odor concentration is 0.5. (**B**) Same as in a but for odor concentration of 10^{-1}. PER is enhanced in a starved state (p = 1e-6 and 0.74 for state and odor factors, Scheirer–Ray–Hare test). (**C**) PER probability in wild type fed flies in response to different concentrations of odors. n = 19 and 28 for 10^{-1} and 0.5 concentration groups, respectively. Error bar, standard error of the mean.

Box plots indicate the median (gray and red lines), mean (black and red dots), quartiles (box), and 5-95% range (bar).

Figure 2-figure supplement 2. Wings and legs are not required for odor-evoked PER

(**A**) Integrated PER duration in wing-removed Orco-Gal4>UAS-Kir2.1 flies (n = 26). GRNs in wings were dispensable for PER to certain odors (p = 6.0e-9, Kruskal-Wallis test, p = 0.0020, 0.0011, 0.75, 0.089, 0.00071, 0.58, 0.84, and 0.94 for comparison between mineral oil (MO) control vs EBR, 2PT, IPA, MCH, BNZ, OCT, Air, and Water, Dunn's post-hoc tests with Benjamini–Hochberg false discovery rate correction). (**B**) Integrated PER duration in wing- and leg-removed, Orco-Gal4>UAS-Kir2.1 flies (n = 27). GRNs in wings were dispensable for PER to certain odors (p = 0.004, Kruskal-Wallis test, p = 0.39, 0.029, 0.18, 0.18, 0.029, 0.44, 0.88, and 0.71 for comparison between mineral oil (MO) control vs EBR, 2PT, IPA, MCH, BNZ, OCT, Air, and Water, Dunn's post-hoc tests with Benjamini–Hochberg false discovery rate correction).

Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).

Figure 2-figure supplement 3. Contributions of OBPs to odor-evoked PER

(A, B) Integrated PER duration in control (*tubulin-Gal4/w^{1118}*) and Obp56g RNAi flies. These flies show similar level of PER (n = 24 and 23 for control and RNAi, p = 0.58 and 0.0017 for group and odor factors, Scheirer–Ray–Hare test). **(C-H)** Integrated PER duration in control (*tubulin-Gal4/y,w^{1118};P{attP,y[+],w[3']}*) and OBP RNAi flies. Obp18a RNAi and Obp57e RNAi flies show reduced level of PER as compared to control (n = 33, 25, 22, 25, 27, and 28 flies for **C-H**, p = 0.050, 0.36, 0.18, 1.9e-4, 0.92 and 0.10, 0.48, 0.063, 0.045, 0.022 for group and odor factors for **D-H**, Scheirer–Ray–Hare test).

Box plots indicate the median (gray and red lines), mean (black and red dots), quartiles (box), and 5-95% range (bar).

912
913

Figure 3-figure supplement 1. Responses of GRNs to odors and sucrose

(**A**) Image of a brain of Gr5a-Gal4>UAS-GCaMP6s fly, stained with anti-GFP to reveal Gr5a
neurons (green) and nc82 to reveal synapses (magenta). Scale bar, 50 μm. (**B**) Baseline
florescence intensity of GCaMP6s in Gr5a GRNs. Fluorescence (arbitrary unit) is color coded
according to the scale bar. (**C**) Peak florescence intensity of GCaMP6s in the same fly as shown
in B after stimulation with 4-methylcyclohexanol. Fluorescence is color coded according to the
scale bar. The ROI used for ΔF/F calculation is indicated by a white rectangle. Scale bar, 20 μm.
(**D-F**) Same as in **A-C**, but for Gr66a-Gal4>UAS-GCaMP6s fly and benzaldehyde stimulation.
(**G**) Trial averaged peak responses of Gr5a GRNs to different concentrations of ethyl butyrate

(EBR) and benzaldehyde (BNZ). n = 5 flies. (**H)** Same as in **G,** but for Gr66a GRNs. **(I)**
Responses to 4-methylcyclohexanol are positively correlated with that to sucrose (0.25%) in
Gr5a GRNs (R^2 = 0.73, p = 0.031). Each dot corresponds to a fly. (**J**) Trial averaged peak
responses of Ppk28 GRNs to individual odors. These neurons do not respond to odors. n = 5
flies. (**K)** Same as in **J**, but for Ir94e GRNs. n = 9 flies.
Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range
(bar).
931

932
933
Figure 3-figure supplement 2. **Individual responses of GRNs to odors**

(A) Time course of calcium responses (ΔF/F) to odors in Gr5a GRNs. Responses to nine odors in each fly are shown in one panel (n = 5 flies). Odor application period is 5 to 7 s, and each odor was applied three times (different trials are shown in different colors).

(B) Same as in **A**, but for Gr66a GRNs.

Figure 3-figure supplement 3. PER and GRN responses to various concentrations of banana odor

(**A**) Integrated PER duration in response to a banana odor presented at various concentrations in wild type flies. n = 21 flies. Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar). (**B**) Trial averaged peak responses of Gr5a GRNs in response to various concentrations of banana odor in Gr5a-Gal4>UAS-GCaMP6s flies. n = 5 flies. Error bar, standard error of the mean. Box plots indicate the median (gray line), mean (black dot), quartiles (box), and 5-95% range (bar).

949
950

Figure 5-figure supplement 1. Correlation between odor and taste responses in individual sensilla

Spike responses to banana odor (0.01%) are positively correlated with those to sucrose (0.25%) in L-type sensilla. Each dot corresponds to a fly ($R^2 = 0.62$, p = 0.00044).

957 **Table 2. Summary of statistical tests**

Results of multi-factorial statistical tests

Figure	Statistical test	concentration	identity	interaction
Figure 1E	Scheirer–Ray–Hare	3.0E-05	2.0E-07	2.0E-10
Figure 1G	Scheirer–Ray–Hare	3.6E-13	3.5E-11	2.8E-21
		method	**odor**	**interaction**
Figure1-figure supplement1C	Scheirer–Ray–Hare	0.94	5.5E-03	2.8E-03
		state	**odor**	**interaction**
Figure 2-figure supplement1A	Scheirer–Ray–Hare	0.88	5.6E-03	1.8E-03
Figure 2-figure supplement1B	Scheirer–Ray–Hare	1.0E-06	0.74	2.7E-03
		genotype	**odor**	**interaction**
Figure 2A	Scheirer–Ray–Hare	0.0079	0.10	0.0020
Figure 2B and C (comparison to UAS control)	Scheirer–Ray–Hare	4.9E-05	0.12	3.5E-05
Figure 2B and D (comparison to UAS control)	Scheirer–Ray–Hare	0.49	0.0014	6.6E-04
Figure 2B and E (comparison to UAS control)	Scheirer–Ray–Hare	0.17	0.55	0.027
Figure 2B and F (comparison to UAS control)	Scheirer–Ray–Hare	0.11	1.2E-05	2.0E-07
Figure 2C	Scheirer–Ray–Hare	1.0E-15	0.71	1.0E-16
Figure 2D	Scheirer–Ray–Hare	0.28	0.0012	3.0E-05
Figure 2E	Scheirer–Ray–Hare	6.2E-04	0.56	0.0036
Figure 2F	Scheirer–Ray–Hare	0.14	5.6E-09	1.7E-08
Figure 2G	Scheirer–Ray–Hare	1.8E-04	0.041	2.6E-05
Figure 2H	Scheirer–Ray–Hare	1.1E-04	0.32	0.00015
Figure 2I	Scheirer–Ray–Hare	0.039	0.48	0.0083
Figure 2J	Scheirer–Ray–Hare	0.0066	0.16	0.0051
Figure 2K	Scheirer–Ray–Hare	0.24	0.26	0.0032
Figure 2-figure supplement3A and B	Scheirer–Ray–Hare	0.58	0.0017	0.0011
Figure 2-figure supplement3C and D	Scheirer–Ray–Hare	0.050	0.10	0.0060
Figure 2-figure supplement3C and E	Scheirer–Ray–Hare	0.36	0.48	0.041
Figure 2-figure supplement3C and F	Scheirer–Ray–Hare	0.18	0.063	0.011
Figure 2-figure supplement3C and G	Scheirer–Ray–Hare	1.9E-04	0.045	8.0E-06
Figure 2-figure supplement3C and H	Scheirer–Ray–Hare	0.92	0.022	0.0072
		organ condition	**odor**	**interaction**
Figure 3D (control vs labella covered)	Mixed two way ANOVA	2.6E-15	5.1E-03	9.8E-03
Figure 3D (control vs palps covered)	Mixed two way ANOVA	0.69	0.055	0.91
Figure 3F (control vs labella covered)	Mixed two way ANOVA	3.0E-16	1.1E-07	1.9E-07
Figure 3F (control vs palps covered)	Mixed two way ANOVA	0.48	4.2E-11	0.90
Figure 3H	Mixed two way ANOVA	8.0E-06	0.037	0.070
		odor identity	**sucrose concentration**	**interaction**
Figure 6B (intact)	Scheirer–Ray–Hare	4.6E-04	2.1E-07	0.057
Figure 6B (olfactory organs removed)	Scheirer–Ray–Hare	0.035	0.0070	0.93

Figure 1-movie supplement 1. Odor-evoked PER

A movie capturing odor-evoked PER with the plot of haustellum angle over time. Six points on the head were tracked by DeepLabCut.