Putting life history theory to the test - the estimation of reproductive values

from field data

Mirjam J. Borger[1], Jan Komdeur[1], David S. Richardson[2], Franz J. Weissing[1]

[1]Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC,

Groningen, The Netherlands.

[2]Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University of East Anglia,

Norwich Research Park, NR4 7TJ Norwich, UK

Lay Summary

To study evolution in empirical systems it is important to accurately and precisely measure fitness. Here,

using simulations, we compare two methods for estimating fitness and test their accuracy and precision.

One estimates life-history parameters and calculates a fitness proxy using complex algebra. The other

method estimates genetic contribution to future generations by tracking descendants down the

pedigree. We conclude that the pedigree method is unreliable, while the algebraic method is accurate

and precise.

Abstract

Quantifying fitness is important to understand adaptive evolution. Reproductive values are useful for

making fitness comparisons involving different categories of individuals, like males and females. By

definition, the reproductive value of a category is the expected *per capita* contribution of the members

of that category to the gene pool of future generations. Life history theory reveals how reproductive

values can be determined via the estimation of life-history parameters, but this requires intricate algebraic

calculations. Recently, a more intuitive, pedigree-based method has become popular, which estimates

the genetic contributions of individuals to future generations by tracking their descendants down the

pedigree. Here we compare both methods. We implement various life-history scenarios (for which the

"true" reproductive values can be calculated) in individual-based simulations, use the simulation data to

estimate reproductive values with both methods, and compare the results with the true target values. We

show that the pedigree-based estimation of reproductive values is either systematically biased (and hence

inaccurate), or very imprecise. This holds even for simple life histories and under idealized conditions. In

contrast, the traditional algebraic method estimates reproductive values with high accuracy and precision.

Keywords

Class-structured population, fitness in a life history context, life-history model, pedigree-based

reproductive value

Introduction

Quantifying fitness is important to understand how natural selection affects evolution. Estimating fitness in the wild is, however, a difficult task. Studies often measure survival, recruitment rate, or the number of offspring reaching independence as a proxy for fitness. In some well-monitored populations, lifetime reproductive success can be determined. Expected lifetime reproductive success is often a good proxy for fitness (Brommer et al., 2004), but even this measure is incomplete as it does not account for the rate of reproduction and the fact that different types of offspring (e.g. females and males; different size classes) cannot just be added up, as they have a different potential for spreading their genes to future generations.

Reproductive value (RV) (Fisher, 1930) is a comprehensive fitness measure that copes with these problems. The reproductive value of a certain class of individuals is defined as the expected contribution of an individual in that class to the future gene pool of the population (Grafen, 2006). Reproductive values are influenced by all kinds of life-history decisions, like age at first reproduction, trade-offs between survival and fecundity, risk-taking behaviour, or the kind of offspring produced (e.g. sons or daughters). RVs are a particularly useful tool in behavioural ecology, as they allow the evolutionary costs and benefits of fitness-relevant individual decisions to be quantified. For example, Tinbergen and Daan (1990) used an RV approach to predict the optimal clutch size in passerine birds on the basis of the trade-off between current and future reproduction. Reproductive values also play an important role in predicting the optimal sex ratio (Pen and Weissing, 2002) and in the evolutionary theories of senescence (Hamilton, 1966; Baudisch, 2005).

While the definition of reproductive value is straightforward, its measurement is not. As briefly reviewed below, RVs can be derived from a life-history model by solving an eigenvector equation that contains all relevant life-history parameters (Caswell, 1982). In line with this, most applications of RVs in empirical systems first estimate life-history parameters and subsequently obtain RVs by inserting these parameters in the eigenvector equation (Leverich and Levin, 1979; Schulman and Chapais, 1980; Tinbergen and Daan, 1990; Newton and Rothery, 1997; Pen et al., 1999; Bonduriansky and Brassil, 2002; van de Pol et al., 2007; Bouwhuis et al., 2012). Hereafter, we will call the RVs thus obtained "model-based reproductive values" (mRV). Obtaining a good estimate of reproductive values along these lines can, however, be difficult in natural systems, since it requires a good understanding of all relevant life-history transitions and because life-history parameters can often only be estimated with limited accuracy and precision.

Recently, reproductive values have been estimated with a more intuitive method, based on genetic
pedigree data (Barton and Etheridge, 2011; Chen et al., 2019; Hunter et al., 2019; Reid et al., 2019). In this
method, the average *per capita* number of descendants of the members of a certain class of individuals
down the pedigree is used as an estimate of the reproductive value of that class. Hereafter, we will call
the RVs thus obtained "pedigree-based reproductive values" (pRV). The pRV method closely reflects
Fisher's definition of reproductive value. This method requires the availability of sufficiently deep and
complete pedigrees, but if such pedigrees are available it has the big advantage that no detailed
knowledge of life-history transitions or estimates of life-history parameters are required. Therefore, the
pRV method could potentially be a useful substitute for the mRV method. However, whether the pRV
method matches the mRV method in accuracy and precision when applied to field data has yet to be
determined.

In this theoretical study, we will compare the performance of both methods when applied to organisms
with a relatively simple life history. To this end, we use individual-based simulations to produce several
generations of individuals on the basis of a life-history model. The simulation data can be used to construct
a pedigree (complete or incomplete) and to estimate life-history data such as survival probabilities and
fecundities. Subsequently, reproductive values can be estimated by both methods (mRV and pRV) and
compared with the "true" reproductive values (which are well-defined for the simulated populations).
This allows us to judge the accuracy and precision of both estimates and to address questions like: how
deep and complete must a pedigree be in order to get a reliable pRV estimate? How well must the life-
history model be known, and how well do we need to know the life-history parameters in order to get
reliable mRV estimates?

Before approaching the estimation problem, we first give some theoretical background by briefly
reviewing some basal insights of life history theory.

Theoretical considerations: why are reproductive values useful?

In a population with discrete, non-overlapping generations and only one type of individual (e.g. no sex
differences), expected lifetime reproductive success (ELRS) is an adequate fitness measure, as alleles that
enhance the lifetime reproductive output of their bearers have a selective advantage. The situation is
different in populations with different classes of individuals (e.g. females and males; breeders and helpers;
workers, soldiers and reproductives). In such cases, alleles inducing their bearers to produce a maximal
number of offspring throughout their lifetime are not necessarily favoured. The reason is that different

types of offspring may differ in the efficiency with which they spread genes to future generations. For
example, in a population with a 4:1 female-biased sex ratio (4 females per male), the reproductive success
per male is on average four times greater than the reproductive success per female (because each
offspring has one mother and one father). Therefore, an allele inducing the lifetime production of two
sons has a lower ELRS than an allele inducing the lifetime production of four daughters, but a higher
"fitness": it spreads more efficiently to future generations, as the expected reproductive success of two
sons is twice the expected reproductive success of four daughters (for the given sex ratio). Similarly,
adding up potential offspring does not result in a suitable fitness measure when generations are
overlapping, and different age classes coexist. In this situation, the timing of reproduction matters: for
example, it may be advantageous to produce offspring as early in life as possible (even if this leads to a
lower lifetime production of offspring), as early-born offspring can more quickly contribute to the spread
of their parental genes.

Quantifying fitness in a life-history context (i.e., in a population with different categories of individuals) is
therefore not an easy task. Fortunately, life-history theory is a well-developed branch of evolutionary
biology (Roff, 1992; Stearns, 1992; Caswell, 2001). A typical life-history model starts with a life-cycle graph
(see Figure 1 for examples) that includes all life history "stages" (e.g. all age classes) and the transitions
between these stages (e.g. survival to the next age class; production of offspring of age one). If the number
of stages is finite, this graph can be translated into a "stage transition matrix" which in the case of three
stages looks like this:

115
$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$
(1)

116   Matrix element a_{ij} is the *per capita* contribution of a member of stage *j* to the number of individuals in
117   stage *i* in the next time step. For example, the stage transition matrix of an age-structured population with
118   three age classes has the special form of a "Leslie matrix":

119
$$\mathbf{A} = \begin{pmatrix} F_1 & F_2 & F_3 \\ P_1 & 0 & 0 \\ 0 & P_2 & 0 \end{pmatrix}.$$
(2)

Here, P_1 is the probability to survive from age class 1 to age class 2, P_2 is the probability to survive from
age 2 to age 3, and $F_1, F_2,$ and F_3 is the expected number of surviving offspring (surviving till age class 1)
produced per individual of the respective age class.

Life-history matrices are a very useful tool, as they determine the dynamics of a stage-structured

population (Stearns, 1992). In most cases, such a population will converge to a "stable stage distribution"

(in the case of an age-structured population: a stable age distribution; in the case of a population with

two sexes: a stable sex ratio), that is, to a state where the ratio between the number of individuals in the

stages ($n_1 : n_2 : n_3$) does not change anymore. Once the stable stage distribution is reached, the population

as a whole grows with a characteristic "population growth factor λ". When λ=1, the population size will

stay constant over time, while the population size will increase exponentially when λ>1 and decrease

exponentially when λ<1.

The population growth factor λ is the most fundamental fitness measure in a stage-structured population

(Brommer, 2000): if we consider a population with transition matrix **A** and corresponding λ, a mutant

inducing a slightly different transition matrix $\mathbf{A_m}$ with growth factor λ_m can be expected to successfully

invade the population if $\lambda_m > \lambda$, while mutants with $\lambda_m < \lambda$ will go extinct. Mathematically, λ is the

"dominant eigenvalue" of the stage-transition matrix **A**, and there are recipes to calculate λ for a given

matrix **A**. Unfortunately, the eigenvalue equation determining λ is an "implicit" equation (in case of an

age-structured population the so-called "Euler-Lotka equation" (Otto and Day, 2007)), which even for

simple life histories can only be solved numerically and does not provide much insight into the factors

governing evolution. It is therefore useful to look out for an alternative fitness measure.

At this point reproductive values come into play. Conceptually, the "reproductive value" v_i of a stage i is

the expected genetic contribution of a member of stage i to the gene pool of future generations (Grafen,

2006). Mathematically, the vector $\mathbf{v}^T = (v_1, v_2, v_3)$ of reproductive values satisfies the equation

$\mathbf{v}^T \cdot \mathbf{A} = \lambda \cdot \mathbf{v}^T$, which means that it is a "left eigenvector" with respect to the dominant eigenvalue λ.

Reproductive values are useful for making evolutionary predictions in two different ways. First, in many

life-history contexts, including those of age-structured populations, natural selection tends to maximize

the reproductive value of each age class (Schaffer, 1974). This is the basis of dynamic programming

(Houston and McNamara, 1999), a powerful technique for solving difficult problems in behavioural

ecology, such as finding the optimal allocation of resources to various activities, like growth and

reproduction. Second, and more important for us, the reproductive values in a given "resident" population

can be used to calculate the "selection gradient" $\partial\lambda/\partial x$, which tells us whether the population growth

factor λ (the most fundamental measure of fitness) will increase or decrease with a change in a

(behavioural) strategy x. In other words, the selection gradient tells us the direction of selection: if $\partial\lambda/\partial x$

is evaluated at the strategy x^* of the resident population, a positive selection gradient indicates that larger

values $x > x^*$ are selectively favoured, while smaller values $x < x^*$ are favoured in the case of a negative

gradient. Although λ is very difficult to calculate as a function of x, the sign of the selection gradient can

relatively easily be determined by the following equation (Taylor, 1990; Taylor and Frank, 1996; Brommer,

2000; Pen and Weissing, 2000b, 2002; Otto and Day, 2007; Lion, 2018):

158
$$\text{sign}\left(\frac{\partial \lambda}{\partial x}\right) = \text{sign}\left(\sum_{i,j} v_i \cdot \frac{\partial a_{ij}}{\partial x} \cdot n_j\right).$$
(3)

Here, the v_i and the n_i are the reproductive values and the relative abundances of the various stages in

the resident population (at demographic equilibrium) and the derivatives are evaluated at $x = x^*$.

Why is an equation like this useful? Assume that we want to know whether, in an age-structured

population, selection will favour an increase in the reproductive effort at age i. To this end, let x denote a

behavioural strategy that enhances the reproductive output at age i with a rate $\partial F_i / \partial x = b$ but reduces

survival to the next age with a rate $\partial P_i / \partial x = -c$, where b and c indicate the "benefit" and the "cost" of

an increase in x. Now the sum on the right-hand side of (3) has only two terms, and the sign of the selection

gradient is readily obtained

167
$$\text{sign}\left(\frac{\partial \lambda}{\partial x}\right) = \text{sign}\left(v_1 \frac{\partial F_i}{\partial x} n_i + v_{i+1} \frac{\partial P_i}{\partial x} \cdot n_i\right) = \text{sign}\left(b v_1 - c v_{i+1}\right).$$
(4)

168   Hence, the selection gradient is positive if $b v_1 > c v_{i+1}$, or equivalently:

169
$$b/c > v_{i+1}/v_1.$$
(5)

This inequality is known as the "asset protection principle": the age classes with the highest future fitness

expectation v_{i+1} (i.e. those age classes that have "much to lose") should be least inclined to take the

survival risks associated with increasing the reproductive effort (see Wolf et al., 2007 for various

implications of this principle).

As illustrated by these calculations, reproductive values are useful tools in translating intricate fitness

considerations into transparent cost-benefit comparisons (see Pen & Weissing 2000a, 2000b, 2002 for

other examples). Cost-benefit considerations are further simplified by the fact that reproductive values

are *relative* attributes that are only determined up to a constant factor. This implies that they can be

normalized in the most convenient manner, for example by equating one of the reproductive values to

one (e.g., $v_1 = 1$, which would simplify eqn (5)) or by equating the sum of all reproductive values to one.

Finally, we would like to remark that classical life history theory is ecologically not consistent, in that its
models predict exponential population growth (when λ>1) or exponentially population decline (when
λ<1). To cope with this problem, density dependence needs to be incorporated into a life history model.
This has to be done in an explicit manner, as the form of density dependence can have major implications
for the course and outcome of evolution (Mylius and Diekman, 1995; Pen and Weissing, 2000b). Below
we illustrate how to introduce density dependence in a life-history model. Rather than complicating the
model, this actually simplifies calculations, as the inclusion of density dependence has two benefits: (a) it
allows the assumption that λ=1 at ecological equilibrium and (b) it reduces the dimensionality of the
parameter space (thus making it easier to classify the evolutionary outcomes).

Methods

Simulations for the pedigree-based estimation of reproductive values

All individual-based simulations were based on a specific life-history scenario. Each scenario is specified
by a set of parameters (such as the age-dependent survival probabilities P_i and fecundities F_i characterizing
an age-structured population). For a given state X, all individuals in that state have the same state-specific
parameters. Accordingly, all individuals in that state have the same (expected) state-specific reproductive
value. Time proceeds in discrete steps, where one time step corresponds to a reproductive season. During
a time step t, each individual is in a certain state (either an age class or a breeding state), which can change
from one time step to the next. Reproduction takes place at the start of each time step. If, according to
the life history scenario, F_X is the expected fecundity in state X, the individuals in that stage produce F_X
offspring on average, while the actual number of offspring produced per individual is drawn from a
Poisson distribution with mean F_X. Offspring enter the population at time t+1, and they belong to age class
1 in the age-based scenarios and to the juvenile state in the breeder-helper scenario. After reproduction,
all individuals present at time t change their state stochastically (including death and staying in the same
state), according to the probability assigned to their state by the life-history model. Each simulation was
repeated for a fixed number of time steps T (typically T=100, of which the first 20 time steps are shown in
the figures). For each life-history scenario, simulations were run for various parameter combinations, and
in each case the simulations were repeated at least 100 times. To ensure that our results were not biased
by start-up effects, we initialized all populations in demographic equilibrium. Our results are therefore
not examples of "transient dynamics" (Hastings, 2004; Hastings et al., 2018). This was confirmed by
additional simulations that were first run for thousands of time steps (thus ensuring equilibration) before
the measurements were started.

We used a "gene-dropping" approach (MacCluer et al., 1986) to estimate pRVs. At the start of the

simulation (t=0), every individual is endowed with a unique marker (corresponding to a unique allele at a

locus with infinite alleles). To ensure that gene dropping happened in demographic equilibrium, we first

let each simulation run for 100 time steps (from t=-100 until t=-1) before gene dropping was initiated (at

t=0). At each reproduction event, the offspring inherit their marker from their parent. This allowed us to

identify the descendants of each individual of the initial population in all future time steps. For each life

history state X and each time t, we determined the average *per capita* number of descendants at time t

for those individuals in the initial population that started in state X. We interpret this number, $pRV_x(t)$, as

the pedigree-based estimate of the RV of individuals in state X: according to the definition of RV ("*per*

capita contribution to the gene pool of future generations") $pRV_x(t)$ should, for large t, approximate the

"true" reproductive value of state X. We then normalize all reproductive values so that the RV of the "first"

state (age class 1 in the age-structured scenarios; juvenile state in the helper-breeder scenario; young

female in the sex-and-age-structured scenario) is scaled to 1. To achieve this, the values $pRV_x(t)$ were

divided by $pRV_1(t)$, the estimated reproductive value of the first state. For simplicity, we assumed that

marker inheritance is based on asexual reproduction in most simulations, and that individuals are haploid.

However, in the simulations where sex was included, sexual reproduction and diploidy was assumed. As

shown in Appendix A in the Supplement, the expected value of $pRV_x(t)$ can be calculated mathematically.

To prevent exponential growth or exponential decline of the simulated populations, we made the

reasonable assumption that population sizes are kept in check by density dependent processes.

Therefore, we added density regulation by assuming that fecundity is density dependent:

231
$$F_X(N) = \frac{F_{X,0}}{1 + \alpha N},$$
(6)

Where $F_{X,0}$ is the baseline fecundity in state X, N is population size and α indicates the intensity of density

dependence. In other simulations (not shown), we also implemented other forms of density regulation

(via density-dependent fecundity of only a single state or via density-dependent survival of one or all

states); in all cases, the results agreed with those shown below. Unless indicated otherwise, we chose the

parameter α in such a way that the population size of the resulting stationary population was N=1,000.

Note that this means that about 1,000 individuals are present at every time step. If T is the depth of the

pedigree and E is the life expectancy of individuals, this implies that the pedigree encompasses about

$(T/E) \cdot 1,000$ individuals. Thus, in all our simulations with a time horizon $T = 20$ at least 10,000 individuals

were included in the simulation-based pedigree.

For the model-based estimation of reproductive values (mRV), the life-history parameters had to be

estimated from the life-history events (survival, death, state transition, offspring production) observed in

the simulations. To estimate $mRV_X(t)$ for a certain time t, we used all events observed on individuals in

state X up to time t to estimate the life-history parameters relevant for that state by averaging. For

example, the fecundity F_x in state X was estimated by the average number of offspring produced in state

X (per time step) by all individuals that had ever been in state X until time t. Subsequently, the $mRV_X(t)$

were estimated by calculating the left eigenvector of the matrix **A**(t) that is characterized by all parameter

estimates up to time t.

Four scenarios

In this manuscript we focus on four life history scenarios, which will be described in detail below. The life-

cycle graphs of these scenarios can be found in figure 1 and the corresponding transition matrices and

calculations on reproductive value can be found in Appendix B.

Scenario 1 – A population with two age classes. This simple scenario, with only two life-history states and

three life-history parameters, is characterized by Figure 1(a). Both one- and two-year-old individuals can

reproduce (with fecundity F_1 and F_2). One-year-old individuals survive to their second year with probability

P_1; all individuals die after their second year. A simple calculation (see Appendix B) reveals that a stationary

population ($\lambda=1$) is only achieved if the life-history parameters satisfy the requirement:

$$F_1 + P_1 \cdot F_2 = 1 . \qquad (7a)$$

This equation makes intuitive sense: the left-hand side corresponds to the expected lifetime production

of surviving offspring, which needs to be 1 in a stationary population. Condition (7a) for "ecological

consistency" will not be satisfied for arbitrary values of the life-history parameters. However, it will

eventually be satisfied, as, according to eqn (6), the fecundities decline with population density.

As shown in Appendix B, the "true" reproductive values are given by:

$$v_1 = 1, \ v_2 = F_2 . \qquad (7b)$$

Scenario 2 – A population with three age classes. This scenario, with three life-history states and four life-

history parameters, is described in Figure 1(b). Two- and three-year-old individuals can reproduce (with

fecundity F_2 and F_3). One- and two-year-old individuals can survive to their next year (with probabilities

Figure 1. The four life-history scenarios considered in this study. (a) A two-stage life cycle with two age classes. One-year-old individuals survive to their second year with probability P_1, second-year-old individuals die after reproduction. F_1 and F_2 denote the expected number of offspring (surviving to their first year) produced by one- and two-year-old individuals, respectively. **(b)** A three-stage life cycle with three age classes. Only two- and three-year-old individuals can reproduce (with expected fecundity F_2 and F_3), and one- and two-year-old individuals survive to the next year with probabilities P_1 and P_2, respectively. **(c)** A three-stage life cycle with behavioural stages. A juvenile can either stay juvenile for another time unit (probability P_j), promote to helper status (probability T_{hj}), promote to breeder status (probability T_{bj}), or die. A helper can either stay a helper for another time unit (probability P_h), promote to breeder status (probability T_{bh}), or die. A breeder can stay a breeder (probability P_b) or die. Both helpers and breeders can reproduce with expected fecundity (number of offspring surviving to the juvenile stage) F_h and F_b, respectively. **(d)** A four-stage life cycle including differences between sexes. Young females and young males survive to become an adult with probabilities P_{fy} and P_{my} respectively. Adult females and adult males survive to the next year (and stay in the same state) with probabilities P_{fa} and P_{ma} respectively. Young females and adult females have expected fecundities of F_{fy} and F_{fa}. They randomly pick a mate from the population of males. A newly produced offspring becomes a male with probability s and a female with probability 1-s, where s is the primary sex ratio.

268   P_1 and P_2); all individuals die after their third year. Now the condition for a stationary population and the

269   reproductive values in that population are given by:

$$P_1 \cdot \left(F_2 + P_2 F_2 \right) = 1 . \tag{8a}$$

$$v_1 = 1, \; v_2 = F_2 + P_2 F_3, \; v_3 = F_3 . \tag{8b}$$

Ecological consistency (a stationary population) is ensured by eqn (6).

<u>Scenario 3 – A population with three behavioural classes.</u> This scenario, which is characterized by Figure
1(c), includes three life-history states (juveniles, helpers, breeders) and a full set of nine life-history
parameters. All newly produced individuals start as juveniles. A juvenile either stays in juvenile state for
another time period (with probability P_j), or it becomes a helper (with probability T_{hj}) or a breeder (with
probability T_{bj}), or it dies (with probability $1 - P_j - T_{hj} - T_{bj}$). Likewise, a helper either stays in the helper
state for another period (probability P_h), or it becomes a breeder (probability T_{bh}), or it dies. A breeder
either stays in the breeding state for another time step (probability P_b), or it loses its position and becomes
a helper (probability T_{hb}), or it dies. Helpers and breeders can both produce offspring (fecundities F_h and
F_b per time step).

<u>Scenario 4 – A population with both sex and age classes.</u> This scenario, which is characterized by the life-
cycle graph in Figure 1(d), contains four life-history states: young females, young males, adult females and
adult males. Newly produced individuals start as young females or young males. The sex of an offspring is
assigned at random, where the probability *s* of becoming a male corresponds to the primary sex ratio. The
parameter *s* is the same for all individuals and constant throughout time. Young females survive to
become adults with probability P_{fy}, and young males survive to become adults with probability P_{my}. Adult
females and males survive to the next time step with probabilities P_{fa} and P_{ma} respectively. Both young
and adult individuals can reproduce. We assume that young and adult females produce on average F_{fy} and
F_{fa} offspring per time step, respectively, where these fecundities are density dependent according to (6).
For simplicity, we assume that each male has the same probability of siring any given offspring: whenever
an offspring was produced, a father was drawn at random from the set of all (young and adult) males.

Technical note

Simulations were ran in Visual Studio Enterprise 2019 (version 16.8). Figures were made using R 3.4.1 (R
Core Team, 2021), with the packages ggplot2 (Wickham, 2016) and cowplot (Wilke, 2019).

Results

Stochasticity in the number of descendants. For simulations based on Scenario 2 (three age classes), Figure 2 illustrates different sources of stochasticity in the number of descendants per individual. Figure 2a shows that, within a single simulation, stochasticity in individual reproductive success is extensive. For

Figure 2. Stochasticity in the number of descendants. Example simulations based on Scenario 2 (three age classes; Fig. 1b) illustrate stochasticity within and across simulations. **(a)** Number of descendants at a given time step (0 to 20) from nine individuals (three per age class) that were present in the same simulation at time zero. **(b)** For four replicate simulations, the panels show the number of descendants at a given time step of the three cohorts at the start of the simulation (t=0): one-year olds (green), two-year olds (blue), and three-year olds (red). **(c)** Summary of 100 simulations. Dashed horizontal lines correspond to the number of descendants expected by the "true" RVs. The dots connected by solid lines indicate the median of the 100 simulations. The 50% (respectively 90%) central values of the simulations are represented by the darker (respectively lighter) shaded area around the medians. Parameter values: $P_1=0.25$, $P_2=0.5$, $F_2=1$ and $F_3=6$.

example, two of the three individuals that started in age class 1 have not left any descendants after 20

years, while the third individual (Individual 2) leaves about 20 descendants. Similar patterns are seen for

the other age classes. Estimating RV based on a single individuals is therefore highly inaccurate, since

these values mainly reflect stochasticity, rather than the capability of a class of individuals to spread their

genes to future generations. When using estimates based on the averages of entire cohorts (or

behavioural groups in later scenarios), this stochasticity partially gets averaged out. Yet, as shown in Figure

2b, each simulation still creates a unique pattern of numbers of descendants per age group. To get an

impression of this variation, we ran 100 replicate simulations, resulting in Figure 2c, which shows the

median values and the 50% and 90% central values of these 100 simulations. Figure 2c shows that the

theoretically expected values (dashed lines) are approached in a "zig-zag" manner. As shown in Appendix

A, this is not caused by "transient dynamics" (short-term deviation from demographic equilibrium), but

an intrinsic property of the gene-dropping dynamics. Although the expected values are approached on a

longer-term perspective, the simulations still differ considerably in the number of descendants per cohort.

RV estimates in Scenario 2. In Figure 3, we show the RV estimates of Scenario 2. From now on, we follow

the standard practice in life history theory and report normalized reproductive values, where the

reproductive value of stage 1 is set to one (i.e., $v_1 = 1$). In Scenario 2, we can therefore focus on age classes

2 and 3. In Figure 3a-c it can be seen than individual simulations (comparable to estimating RV in a field

population) show considerable variation in pRV estimates, with some estimates being highly inaccurate.

In contrast, the mRV estimates are both accurate (they closely approximate the true RVs) and precise

(they do not show much variation between replicate simulations). The summary of 100 replicate

simulations in Figure 3d-f shows that these conclusions are hold in general: for Scenario 2, mRVs provide

an accurate and precise estimate of true RVs, while pRVs are quite inaccurate (in particular in the first 6

time steps) and/or unprecise (in particular in the later time steps. Note that one replicate simulation

corresponds to one field population of size N = 1,000, of which a complete pedigree is available. Figure 3

therefore exemplifies that, at least in Scenario 2, pedigree-based estimation of RVs is problematic: On a

short time horizon, the pRV estimates agree with each other reasonably well, but they deviate

substantially and systematically from their target value, the "true" RVs. These deviations occur despite of

the fact that the populations are reasonably large, and all simulations take place at demographic and

ecological equilibrium. In Appendix A of the Supplement, we show mathematically that, for small values

of t, large deviations are always to be expected. On a longer time horizon (t>10) the median pRVs

approximate the true RVs, but now the pRVs differ considerably between simulations.

Figure 3. mRV and pRV estimates in Scenario 2 (3 age classes). (a)-(c) Three replicate simulations (comparable to field populations) illustrating the diversity of outcomes. The dark-coloured lines (dark blue for two-year-olds and dark red for three-year-olds) represent pRV estimates, while the light-coloured lines (light blue and bright red for two- and three-year-olds, respectively) represent mRV estimates. The dashed lines (blue for two-year-olds and red for three-year-olds) represent the "true" values of the RVs. The simulations exemplify that (a) pRVs can substantially underestimate true RVs; (b) pRVs can approach true RVs; and (c) pRVs can substantially overestimate true RVs. In contrast, mRVs approximate true RVs very well in all three cases. **(d,e)** Summary of mRV and pRV estimates from 100 replicate simulations. In both cases, the dashed horizontal lines correspond to the true RVs. The dots connected by solid lines indicate the median of the 100 simulations. The 50% (respectively 90%) central values of the simulations are represented by the darker (respectively lighter) shaded area around the medians. **(f)** The pRV estimates of 100 simulations are plotted against the accompanying mRV estimates, all from time step t=20. To show the relationship between the pRV and the mRV more clearly, we normalized the RVs in such a way that their sum equates to 1 (instead of normalizing v_1 to 1). Here, the green dots represent the RV estimates of one-year-olds, the blue dots represent RV estimates of two-year-olds and the red dots represent RV estimates of three-year-olds. Parameter values: $P_1=0.25$, $P_2=0.25$, $F_2=2$ and $F_3=8$.

332   <u>RV estimates in Scenario 1.</u> Figure 4 shows that even for the extremely simple life history of Scenario 1,

333   pRV is either a systematically biased (on a short time horizon) or imprecise (on a longer time horizon)

334   estimator of RV. In contrast, "true" RV is, from the start, estimated with accuracy and precision by mRV.

Figure 4. mRV and pRV estimates in Scenario 1 (2 age classes). **(a)-(c)** Three example simulations (comparable to field populations). The dark blue lines represent pRV estimates (for two-year-olds), while the light blue lines represent mRV estimates. The dashed lines represent the "true" values of the RVs. The simulations exemplify that (a) pRVs can substantially underestimate true RVs; (b) pRVs can approach true RVs; and (c) pRVs can substantially overestimate true RVs. In contrast, mRVs approximate true RVs very well in all three cases. **(d,e)** Summaries of mRV and pRV estimates from 100 replicate simulations. In both cases, the dashed horizontal lines correspond to the true RVs. The dots connected by solid lines indicate the median of the 100 simulations. The 50% (respectively 90%) central values of the simulations are represented by the darker (respectively lighter) shaded area around the medians. **(f)** The pRV estimates of 100 simulations are plotted against the accompanying mRV estimates, all from time step t=20. To show the relationship between the pRV and the mRV more clearly, we normalized the RVs in such a way that their sum equates to 1 (instead of normalizing v_1 to 1). Here, the green dots represent the RV estimates of one-year-olds and the blue dots represent RV estimates of two-year-olds. Parameter values: $F_1 = 0.5$, $P_1 = 0.25$, $F_2 = 2$.

<u>RV estimates in Scenarios 3 and 4</u>. Figure 5 shows that the problems with pRV estimates also arise in other

life-history scenarios. In Scenario 3 (Fig. 5a), the pRV estimates can be two or even three times as large as

the true RVs in a considerable percentage of the simulations. Although the initial "zig-zag pattern"

observed in Scenarios 2 and 3 does not appear in Scenario 4 (Fig. 5b), pRV remains an inaccurate estimator

of true RV in the initial period; in fact, the systematic bias in this estimate only disappears after a long

period (in this case 10 time steps). Hence, the simulations in Fig. 5 confirm our earlier conclusions: in an

Figure 5. Accuracy and precision of pedigree-based estimates of RV in two more complex scenarios.
Pedigree-based estimation of two (normalised) RVs in 100 replicated simulations based on **(a)** Scenario 3 (juvenile-helper-breeder; Fig. 1(c)), and **(b)** Scenario 4 (young and old males and females; Fig 1(d)). The inserts show the life-cycle graphs. The "true" RVs are represented by dotted lines. Dots connected by solid lines indicate the median pRV of the 100 simulations; the 50% (resp. 90%) central values of the 100 pRVs are represented by the darker (respectively lighter) shaded area around the medians. In (a) blue represent the RV values of helpers and red represent the RV values of breeders. In (b) yellow represent the RV values of young males, while red and blue represent the RV values of adult females and males, respectively. Please note the difference in scale on the y-axis. Parameter values (a): F_j=0, F_h=0.05, F_b=12.5, T_{hj}=0.15, P_h=0.2, T_{hb} = 0, T_{bj}=0.01, T_{bh}=0.2, P_b=0.4 and (b): P_{fy}=0.1, P_{fa}=0.85, P_{my}=0.5, P_{ma}=0.9, F_{fy}=0.095, F_{fa}=2, s=0.3.

initial time period, the pRV values (including their median) differ substantially and systematically from the true RVs. On a longer time horizon, the accuracy of pRV increases (the median pRVs approach the true RVs), but now the estimates are very imprecise, as the individual simulations differ considerable from each other. As one simulation corresponds to a single field study, this results in the unfortunate conclusion that pedigree-based estimation of RVs is either inaccurate (in case of a shallow pedigrees including relatively few time steps) or imprecise (in case of deep pedigrees), even if the study population is relatively large (*N* = 1,000 in the simulations shown thus far), well-mixed, and in demographic and ecological equilibrium.

Effects of population size and time scale. In Appendix C of the Supplement, we also show the effects of population size and a longer time horizon on pRV estimates. When population sizes are small, pRV estimates are strongly affected by demographic stochasticity. After a while, all individuals present at a later time step are descendants of just one individual of the initial population. Accordingly, in each

simulation, eventually pRV is equal to zero for all but one of the categories considered. Therefore, in small

populations, the median pRV of 100 simulations converges to zero relatively rapidly (Supplementary

Figure C1(a)). Furthermore, the variation of the pRV estimates across simulations increases with the time

horizon (Supplementary Figs C1 and C2). In other words, the precision of pRV decreases with the depth of

the pedigree.

<u>Evolutionary predictions based on RV</u>. Above, we have seen that reproductive values play an important

role in predicting the course and outcome of life history evolution. One might speculate that the large

variance of pRV values that we observed in replicate simulations is associated with a corresponding

variance in evolutionary outcomes. To check this, we added an evolving parameter x (corresponding to

reproductive effort) to our Scenario 1 model with two age classes. To this end, we added a gene locus to

the model, with a continuum of alleles, ranging from 0 to 1. Individuals with allele x at this locus have life-

history parameters $P_1(x) = P_{1,0} \cdot (1 - \frac{1}{2}\gamma x^2)$, $F_1(x) = F_{1,0} \cdot (1 + \beta x)$, and $F_2(x) = F_{2,0} = \text{constant}$, where β and γ

are positive. Hence, one-year olds with a larger reproductive effort x have a higher reproductive output

F_1, but a lower survival probability P_1. We assume that x is transmitted from parent to offspring, subject

to rare mutations with small effect size (mutation rate 0.01, mutational variance 0.0025; see Netz et al.

2022). In the course of the generations, x should evolve to a value that optimizes the balance between

current and future reproduction. With the help of reproductive values, the optimal reproductive effort

can easily be calculated. At evolutionary equilibrium, the selection gradient is zero, which according to

eqn (4) corresponds to the equation $b/c = v_2/v_1$, where $b = F_1'(x) = F_{1,0}\beta$ and $c = -P_1'(x) = P_{1,0}\gamma x$, while

$v_2/v_1 = F_{2,0}$, in view of eqn (7b). Inserting all these terms and solving for x, we obtain the optimal

reproductive effort

374
$$x_{opt} = \frac{F_{1,0}}{P_{1,0} \cdot F_{2,0}} \cdot \frac{\beta}{\gamma} \ . \tag{9}$$

375   Figure 6 demonstrates that, for a set of parameters yielding the optimal value $x_{opt} = 0.5$, individual-based

376   simulations do indeed converge to this value, and that they stay close to this value, irrespective of whether

377   the pedigree-based estimate pRV_2/pRV_1 of v_2/v_1 is much larger or much smaller than the "true" value

378    $F_{2,0}$. We conclude that (a) reproductive values are indeed useful for making evolutionary predictions; but

379   that (b) pedigree-based estimates are too imprecise to be reliable predictors of the evolutionary outcome.

Figure 6: Evolutionary predictions based on reproductive values. (a) Evolution of reproductive effort *x* in Scenario 1 (two age classes) in two sets of 100 individual-based simulations; one set starting at $x = 0$ and the other starting at $x = 1$. shows the trajectory of x over time for 200 simulations (100 that started with a monomorphic population where x=0, and 100 that started with a monomorphic population where x=1). The solid lines indicate the median *x*-values of these simulations and the 50% (resp. 90%) central values of x are represented by the darker (resp. lighter) shaded areas around the medians. The dashed line indicates the optimal reproductive effort, which, based on the "true" reproductive values, is equal to $x_{opt} = 0.5$, according to eqn (9) and the parameter setting $P_{1,0} = 0.25$, $F_{1,0} = 0.5$, $F_{2,0} = 2$, $\beta = 0.5$, $\gamma = 1$. **(b)** and **(c)** show two example simulations that both start at equilibrium ($x = 0.5$). The evolutionary trajectories of *x* ((b2) and (c2)) stay close to this equilibrium value, while the pRV ratio pRV_2/pRV_1 either overestimated (b1) or underestimated (c1) the "true" RV ratio v_2/v_1 (dashed line).

Discussion

Reproductive values are a very useful theoretical tool, as they can help to answer questions like: what is

the optimal clutch size (Tinbergen and Daan, 1990)? What is the optimal investment in male versus female

offspring (Fisher, 1930; Pen and Weissing, 2002)? Under which circumstances should offspring stay on

their natal territory and help their parents raise additional offspring (Pen and Weissing, 2000c)? However,

we need to be able estimate RVs with accuracy and precision if we are to use them to test predictions of

life-history theory.

Reproductive values are increasingly estimated by making use of pedigree information from long term

studies (e.g. Barton and Etheridge, 2011; Chen et al., 2019; Hunter et al., 2019; Reid et al. 2019). In

essence, the *per capita* number of descendants of the members of a certain life-history stage is used to

estimate the reproductive value of that stage. This method is intuitively appealing, as it closely reflects

Fisher's (1930) definition of "reproductive value". However, our study clearly reveals that the method has

important drawbacks in practice, even if the pedigree includes many individuals (hundreds or even

thousands) per time step. The analytical arguments (Appendix A) and simulations outlined in the results

show that pedigree-based estimates are strongly and systematically biased (and therefore inaccurate) if

the pedigree encompasses a short time horizon (say, $t < 10$). On a longer time horizon, the median pRVs

of 100 replicate simulations match the "true" RVs reasonably well, but the individual simulations tend to

diverge considerably from each other with increasing "depth" of the pedigree. As one simulation

corresponds to one field study, this implies that the pedigree-based estimate of RV is typically way "off

target" if it is only based on one, or a small number of, field studies.

Our findings concur with the patterns reported in natural populations. In both Soay sheep (*Ovis aries*,

Hunter et al., 2019) and Florida Scrub-Jays (*Aphelocoma coerulescens*, Chen et al., 2019) estimates of RV

based on individuals did not stabilise, just as in our simulations. Moreover, the pedigree-based estimates

showed a similar initial "zig-zag" pattern as in our study (e.g. Fig. 2b, c, 3e, 4e, 5a). Last, but not least,

when the estimation was repeated for the same population by applying the gene-dropping method to

different cohorts, the RV estimates obtained varied a lot (Hunter et al., 2019).

Our study confirms that the more traditional, model-based calculation of reproductive values on the basis

of estimates of the life-history parameters has more desirable statistical properties. In our simulations,

we found that the mRV estimates are slightly biased, but the bias is too small to be visible in the figures.

mRV has the additional advantage that a sensitivity analysis (Caswell, 2019) can reveal how parameter

uncertainty affects the precision of the estimate. In the field, such uncertainty can be considerably larger

than in our simulations. Most importantly, we assumed that all stage transitions, including death, can be

observed in the study population. In real populations, the estimation of, say, survival probabilities will

often be unprecise and/or biased, as mortality cannot always be distinguished from emigration.

We purposely kept our model assumptions as simple as possible. We focussed on very simple life-

histories, assuming that estimation issues that arise in simple scenarios will most likely be even worse in

more complex life-histories. We assumed that our populations are closed (no emigration or immigration),

that pedigree information is complete, and that the life-history parameters remained constant over time.

Again, it is likely that RV estimates get worse when the situation is less ideal (as is the case for most field

studies). Most of our simulations assume asexual reproduction, but as shown in Figure 5b they equally

apply to populations with sexual reproduction and different sexes. However, they are not necessarily

representative for more complicated situations, such as kin interactions, inbreeding or inbreeding

avoidance, and sexual selection. But again, we would argue that methods that do not work well in a simple

context will most likely also fail in more intricate situations.

Life history theory is one of the most advanced branches of evolutionary biology, with sophisticated tools

and methods and a well-established fitness concept (Brommer, 2000). Individual-based simulations are

rarely used in life-history studies, perhaps because analytical techniques are readily available. Yet, such

simulations can provide valuable additional information. First, simulation models are very flexible and can

easily be tailored to the intricacies encountered in real-world situations. For example, it is quite difficult

to include sex- and age-structure, non-random mating, and kin interactions in a life history model without

compromising analytical tractability, while this is straightforward in a simulation approach. Second,

running replicated simulations gives researchers a good idea on the kind and degree of variation to be

expected. Individual-based simulations are therefore a useful tool for judging the validity of empirical

methods, such as the pedigree-based estimation of reproductive values.

Acknowledgements

We thank the MARM research group, the Seychelles warbler research group and the reviewers of a

previous version of this manuscript for their insightful comments on our study and the manuscript.

Funding

MJB was funded by ALW NWO Grant No. ALWOP.531 awarded to JK and DR. The work of FJW is supported

by the European Research council (ERC Advanced Grant No. 789240). The authors had no conflict of

interest.

Data availability
All data is archived on the University of Groningen Dataverse, which will be publicly accessible once this
manuscript is published.

References

Barton NH, Etheridge AM, 2011. The relation between reproductive value and genetic contribution. Genetics 188:953-973. doi: 10.1534/genetics.111.127555.

Baudisch A, 2005. Hamilton's indicators of the force of selection. Proceedings of the National Academy of Sciences of the United States of America 102:8263–8268.

Bonduriansky R, Brassil CE, 2002. Rapid and costly ageing in wild male flies. Nature 420:377.

Bouwhuis S, Choquet R, Sheldon BC, Verhulst S, 2012. The forms and fitness cost of senescence: age-specific recapture, survival, reproduction, and reproductive value in a wild bird population. The American Naturalist 179:E15-27. doi: 10.1086/663194.

Brommer JE, 2000. The evolution of fitness in life-history theory. Biological Reviews 75:377-404.

Brommer JE, Gustafsson L, Pietiäinen H, Merilä J, 2004. Single-generation estimates of individual fitness as proxies for long-term genetic contribution. The American Naturalist 163:505-517.

Caswell H, 1982. Optimal life histories and the maximization of reproductive value: a general theorem for complex life cycles. Ecology 63:1218-1222.

Caswell H, 2001. Matrix Population Models, second ed. Sunderland, Massachusetts: Sinauer Associates, Inc. Publishers.

Caswell H, 2019. Sensitivity Analysis: Matrix Methods in Demography and Ecology: Springer.

Chen N, Juric I, Cosgrove EJ, Bowman R, Fitzpatrick JW, Schoech SJ, Clark AG, Coop G, 2019. Allele frequency dynamics in a pedigreed natural population. Proceedings of the National Academy of Sciences of the United States of America 116:2158-2164. doi: 10.1073/pnas.1813852116.

Fisher RA, 1930. The Genetical Theory of Natural Selection. Oxford: Oxford University Press.

Grafen A, 2006. A theory of Fisher's reproductive value. Journal of Mathematical Biology 53:15-60. doi: 10.1007/s00285-006-0376-4.

Hamilton WD, 1966. The moulding of senescense by natural selection. Journal of Theoretical Biology 12:12-45.

Hastings A, 2004. Transients: the key to long-term ecological understanding? Trends in Ecology and Evolution 19:39-45. doi: 10.1016/j.tree.2003.09.007.

Hastings A, Abbott KC, Cuddington K, Francis T, Gellner G, Lai YC, Morozov A, Petrovskii S, Scranton K, Zeeman ML, 2018. Transient phenomena in ecology. Science 361. doi: 10.1126/science.aat6412.

Houston AI, McNamara JM, 1999. Models of Adaptive Behaviour. Cambridge, UK: Cambridge University Press.

Hunter DC, Pemberton JM, Pilkington JG, Morrissey MB, 2019. Pedigree-Based Estimation of Reproductive Value. Journal of Heredity 110:433-444. doi: 10.1093/jhered/esz033.

Leverich WJ, Levin DA, 1979. Age-specific survivorship and reproduction in *Phlox drummondii*. The American Naturalist 113:881-903.

Lion S, 2018. Class Structure, Demography, and Selection: Reproductive-Value Weighting in Nonequilibrium, Polymorphic Populations. The American Naturalist 191:620-637. doi: 10.1086/696976.

MacCluer JW, VandeBerg JL, Read B, Ryder OA, 1986. Pedigree analysis by computer simulation. Zoo Biology 5:147-160.

Mylius SD, Diekman O, 1995. On evolutionarily stable life histories, optimization and the need to be specific about density dependence. Oikos 74:218-224.

Netz C, Hildenbrandt H, Weissing FJ, 2022. Complex eco-evolutionary dynamics induced by the coevolution of predator–prey movement strategies. Evolutionary Ecology 36:1-17. doi: 10.1007/s10682-021-10140-x.

Newton I, Rothery P, 1997. Senescence and reproductive value in sparrowhawks. Ecology 78:1000-1008.

Otto SP, Day T, 2007. A Biologist's Guide to Mathematical Modeling in Ecology and Evolution. Princeton,
New Jersey: Princeton University Press.
Pen I, Weissing FJ, 2000a. Sex-ratio optimization with helpers at the nest. Proceedings of the Royal
Society London Biology 267:539-543.
Pen I, Weissing FJ, 2000b. Sexual selection and the sex ratio: an ESS analysis. Selection 1:111-121.
Pen I, Weissing FJ, 2000c. Towards a unified theory of cooperative breeding: the role of ecology and life
history re-examined. Proc R Soc Lond B 267:2411-2418. doi: 10.1098/rspb.2000.1299.
Pen I, Weissing FJ, 2002. Optimal sex allocation: steps towards a mechanistic theory. In: Hardy ICW,
editor. Sex Ratios Concepts and Research Methods Cambridge: Cambridge University Press.
Pen I, Weissing FJ, Daan S, 1999. Seasonal sex ratio trend in the European kestrel: an evolutionarily
stable strategy analysis. The American Naturalist 153.
R Core Team. 2021. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing.
Reid JM, Nietlisbach P, Wolak ME, Keller LF, Arcese P, 2019. Individuals' expected genetic contributions
to future generations, reproductive value, and short-term metrics of fitness in free-living song
sparrows (*Melospiza melodia*). Evolution Letters 3:271-285. doi: 10.1002/evl3.118.
Roff DA, 1992. The Evolution of Life Histories. New York: Chapman & Hall.
Schaffer WM, 1974. Selection for optimal life histories: the effects of age structure. Ecology 55:291-303.
Schulman SR, Chapais B, 1980. Reproductive value and rank relations among macaque sisters. The
American Naturalist 115:580-593.
Stearns SC, 1992. The Evolution of Life Histories. Oxford: Oxford University Press.
Taylor PD, 1990. Allele-frequency change in a class-structured population. The American Naturalist
135:95-106.
Taylor PD, Frank SA, 1996. How to make a kin selection model. Journal of Theoretical Biology 180:27-37.
Tinbergen JM, Daan S, 1990. Family planning in the great tit (*Parus major*): optimal clutch size as
integration of parent and offspring fitness. Behaviour 114:161-190.
van de Pol M, Pen I, Heg D, Weissing FJ, 2007. Variation in habitat choice and delayed reproduction:
adaptive queuing strategies or individual quality differences? The American Naturalist 170:530-
541. doi: 10.1086/521237.
Wickham, H 2016. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag.
Wilke, CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R package version
1.0.0.
Wolf M, van Doorn GS, Leimar O, Weissing FJ, 2007. Life-history trade-offs favour the evolution of
animal personalities. Nature 447:581-584. doi: 10.1038/nature05835.