MOSAIC: A Unified Trait Database to Complement Structured Population Models

Connor Bernard[1,†], Gabriel Silva Santos[1,2,3], Jacques Deere[1,4], Roberto Rodriguez-Caro[1,5], Pol

Capdevila[1,6] Erik Kusch[7], Samuel J L Gascoigne[1], John Jackson[1], Roberto Salguero-Gómez[1,7,8]

[1] Department of Zoology, University of Oxford, 11a Mansfield Rd, OX13SZ, Oxford, United Kingdom

[2] Department of Ecology, Rio de Janeiro State University, Rio de Janeiro, Brazil

[3] Instituto Nacional da Mata Atlântica, INMA, Santa Teresa, Brazil.

[4] Department of Evolutionary and Population Biology, Institute for Biodiversity and Ecosystem Dynamics,

University of Amsterdam, 1012 WX Amsterdam, Netherlands

[5] Departamento de Biología Aplicada. Universidad Miguel Hernández. Av. Universidad, s/n, 03202 Elche

(Alicante) Spain

[6] School of Biological Sciences, University of Bristol, 24 Tyndall Ave, Bristol BS8 1TQ, United Kingdom

[7] Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Section for Ecoinformatics &

Biodiversity, Department of Biology, Aarhus University

[7] Centre for Biodiversity and Conservation Science, University of Queensland, St. Lucia, QLD, Australia

[8] Evolutionary Demography Laboratory, Max Plank Institute for Demographic Research, Rostock, Germany

[†] Corresponding author: connor.bernard@zoo.ox.ac.uk

Page Count: 18

Word Count: 5,534

Figures: 4

Tables: 1

CONFLICT OF INTEREST STATEMENT

The authors report no conflicts of interest.

CONTRIBUTIONS

CB and RSG conceived the ideas; CB, RSG, GSS, JD, and PC designed the methodology; CB,

GSS, JD, PC, and SJLG integrated databases; CB, GSS, and JD collected primary records data;

EK led the climate data acquisition and analysis of geospatial data; and CB led the writing of the

manuscript with assistance from JJ. All authors contributed critically to the drafts and gave final

approval for publication.

DATA AVAILABILITY STATEMENT

The data in this paper and complete database are published open access in multiple formats (*e.g.*,

.csv, .pdf, .R, .tree) on the MOSAIC website https://mosaicdatabase.web.ox.ac.uk and associated

GitHub repository https://github.com/mosaicdatabase/mosaicdatabase.

Abstract (300/300)

1. The ecological sciences have joined the big data revolution. However, despite exponential growth in data availability, broader interoperability amongst datasets is still needed to unlock the potential of open access. The interface of demography and functional traits is well-positioned to benefit from said interoperability. Trait-based ecological approaches have been criticised because of their inability to predict fitness components, the core of demography; likewise, demographic approaches are data-hungry, and so using traits as ecological *shortcuts* to understanding and forecasting population viability could offer great value.

2. Here, we introduce MOSAIC, an open-access trait database that unlocks the demographic potential stored in the COMADRE, COMPADRE, and PADRINO open-access databases. MOSAIC data have been digitised and curated through a combination of existing datasets and additional taxonomic and/or trait records sourced from primary literature. In its first release, MOSAIC (v. 1.0.0) includes 14 trait fields for 300 animal and plant species: biomass, height, growth determination, regeneration, sexual dimorphism, mating system, hermaphrodism, sequential hermaphrodism, dispersal capacity, type of dispersal, mode of dispersal, dispersal classes, volancy, and aquatic habitat dependency. MOSAIC also includes species-level phylogenies for 1,359 species and population-specific climate data where locations are recorded.

3. Using MOSAIC, we highlight a taxonomic mismatch of widely used trait databases with existing structured population models. Despite millions of trait records available in open-access databases, taxonomic overlap between open-access demographic and trait databases is <5%. We identify where traits of interest to ecologists can benefit from

database integration and start to quantify traits that are poorly quantified (*e.g.,* growth

determination, modularity).

4. The MOSAIC database evidences the importance of improving interoperability in open-

access efforts in ecology as well as the need for complementary digitisation to fill

targeted taxonomic gaps. In addition, MOSAIC highlights emerging challenges

associated with the disparity between locations where different trait records are sourced.

KEYWORDS

COMADRE, COMPADRE, functional traits, gap of knowledge, integral projection model

(IPM), interoperability, matrix population model (MPM), open access, open traits, PADRINO.

"I'm not interested in your data; I'm interested in merging your data with other data.

73   Your data will never be as exciting as what I can merge it with."

74   —Sir Tim Berners-Lee (co-founder of the World Wide Web)

Introduction

The ecological sciences have recently joined the big data revolution (Farley, Dawson, Goring, &

Williams, 2018; Marx, 2013; Reichman, Jones, & Schildhauer, 2011). Total species distribution records

measure in the hundreds of millions (Maldonado et al., 2015; Troia & McManamay, 2016). Functional trait

data exist for tens of thousands of species across the globe (Díaz et al., 2016; Enquist et al., 2019; Maitner

et al., 2018). Similar gains have taken place in related fields, from big omics (Field et al., 2009) to global

state-of-the-art climate models (*e.g.*, ERA-5 [https://www.ecmwf.int/en/forecasts/dataset/ecmwf-

reanalysis-v5]) becoming available at fine spatial and temporal resolutions suitable for biological analyses

(Davy & Kusch, 2021; Hersbach et al., 2020), and from behavioural trait data (Rowcliffe, Jansen, Kays,

Kranstauber, & Carbone, 2016) to population data (Global Population Dynamics Database

[https://knb.ecoinformatics.org/view/doi:10.5063/AA/nceas.167.15]; Human Mortality Database (J.R.

Wilmoth, K. Andreev, D. Jdanov, 2007; mortality.org); Human Fertility Database (Jasilioniene et al., 2015;

humanfertility.org); AnAge Database (De Magalhães & Costa, 2009); DATLife [Max Plank Institute of

Demographic Research 2022; https://datlife.org/]). We have also seen growth datasets characterising

phenomena at levels of biological complexity above the organismal level, including community ecology

databases (BioTime (Dornelas et al., 2018); Web of Life (Fortuna, Ortega, & Bascompote, 2014);

metaCommunity Ecology: Species, Traits, Environment and Space (CESTES) (Jeliazkov et al., 2020);

Environmental Data Initiative [https://environmentaldatainitiative.org/]). The growth of records in ecology

datasets and complementary environmental datasets is enabling us now to test ecological theory at larger

and more complex scales. In line with open-access growth, however, there is a recognised need to improve

and coordinate data standards to guide the systematic collection and management of data across different

trait collection programmes (Gallagher et al., 2020; Kissling et al., 2018; Wilkinson et al., 2016).

Despite the increasing availability of biological data, combining different datasets into common

analyses is often hampered by the lack of complementarity between databases. Interoperable systems reflect

the transformation of constituent data types into a standardised format of spatial, temporal, and

measurement scales, accounting for differences in methods (Nadrowski et al., 2013). The need to improve

interoperability across datasets is demonstrated by the widescale emergence of data harmonisation

initiatives across fields of ecology (*e.g.,* Culina et al., 2018; Kattge et al., 2020; Schneider et al., 2019). In

the past decade, dozens of initiatives have taken shape to both centralise data from existing datasets and to

improve data interoperability: standardising units, scales, and terminology for comparative purposes

(Edwards, Lane, & Nielsen, 2000; Maurer, Firestone, & Scriver, 2000). Unifying data formats and

streamlining their integration unlocks the potential for existing datasets to answer questions that cut across

levels of biological complexity. Linking together levels of biological complexity is critical for identifying

how phenomena emerge and transmit across different levels of biological organisation, upscaling and

downscaling through biological systems. For example, the critical linkages between genetics and

biochemistry (The ENCODE Project Consortium, 2012; http://bigg.ucsd.edu/), biochemistry and

physiology (Vitousek, Johnson, & Husak, 2018), and physiology and demography (Laughlin, Gremer,

Adler, Mitchell, & Moore, 2020; Swenson et al., 2020) have benefitted from dataset integration.

The limitations of global-scale datasets are shifting away from data availability and toward data

interoperability. The global-scale integration of functional traits is headed in this direction. Momentum

toward data integration is evidenced by recent database initiatives (*e.g.,* Guerrero-Ramírez et al., 2021) and

global networks that, like the Open Trait Network (https://opentraits.org/), aim to standardise and integrate

trait data across taxa (Gallagher et al., 2020). Despite major improvements in the consolidation and

accessibility of trait data, there is not yet a single-source centralised database spanning behaviour,

physiology, habitat, and other trait data for a wide range of species. Existing databases are often linked by

taxonomy (*e.g.,* FishBase (Froese & Pauly, 2010), CoralTraits (Madin et al., 2016), MammalBase

(Lintulaakso, 2013), AmphiBio (Oliveira, São-Pedro, Santos-Barrera, Penone, & Costa, 2017)); trait type

(*e.g.,* TreeOfSex (The Tree of Sex Consortium 2014), TreeBase (Boettiger & Temple Lang, 2012), Xylum

Functional Traits (Borghetti, Gentilesca, Colangelo, Ripullone, & Rita, 2020)); data type (*e.g.,* GBIF

(https://www.gbif.org/), MOL (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012), TetraDensity (Santini,

Isaac, & Ficetola, 2018)); or a combination of the taxonomies and traits (WooDiv (Monnet et al., 2021),

CarniDiet (Middleton, Svensson, Scharlemann, Faurby, & Sandom, 2021)). A number of other databases

take a more general approach in their thematic scope, but are still constrained to a limited set of traits and

taxonomy (*e.g.,* Amniote (Myhrvold et al., 2015), Pantheria (Jones et al., 2009), BIEN (Enquist, Condit,

Peet, Schildhauer, & Thiers, 2016; Maitner et al., 2018), TRY (Kattge et al., 2020)). MOSAIC offers a

platform that integrates databases in the remit of species with structured population models across their

ecological traits.

Here, we introduce MOSAIC, a centralized database of trait data across the Tree of Life. MOSAIC

is an open-access database that complements the existing demographic data available in the COMPADRE

Plant Matrix Database (Salguero-Gómez et al., 2015), COMADRE Animal Matrix Database (Salguero-

Gómez et al., 2016), and the new PADRINO IPM Database (Levin et al., BioRxiv; doi:

10.1101/2022.03.02.482673; https://levisc8.github.io/RPadrino/). MOSAIC v. 1.0.0 includes 14 frequently

used traits that encompass morphological, reproductive, dispersal, and habitat type attributes for some 300

species of animals and plants. Additional traits will be added in the future (see *Future Direction, below*).

MOSAIC allows users to integrate structured population data to probe larger questions through the

collection, curation, and complementarity of relevant contextual data.

Methods

Scope and Coverage of MOSAIC

The MOSAIC database (Version 1.0.0) includes 14 key trait records across 300 species (Fig. 1).

MOSAIC is designed to provide complementary data for analysis in connection with structured population

models: matrix population models (MPMs, Caswell, 2000), where state variables are discrete (*e.g.*, age

(Morris et al., 2008), ontogeny/development (Smallegange, Deere, & Coulson, 2014), discrete classes of

size (Crouse, Crowder, & Caswell, 1987)), and integral projection models (IPMs, Easterling, Ellner, &

Dixon, 2000), where the state variables are continuous (*e.g.,* size (Jongejans, Shea, Skarpaas, Kelly, &

Ellner, 2011), mass (Ozgul et al., 2010), parasite load (Metcalf, Graham, Martinez-Bakker, & Childs,

2016)). The traits included in MOSAIC 1.0.0 were identified by the MOSAIC team as a set of physical,

physiological, geographic, and behavioural attributes of most immediate relevance to demographic research

(see Table 1; more information at https://mosaicdatabase.web.ox.ac.uk). Traits were also selected in

consideration of the lack of standardised and centralised databases for certain traits (*e.g.,* volancy,

modularity, and growth indeterminacy; see Fig. 2 for trait variance and taxonomic structure of select traits

excluded from existing databases). Importantly, we note that MOSAIC is not a general dataset for analysis

of functional traits, as this is available through other extensive repositories (*e.g.,* TRY (Kattge et al., 2020),

BIEN (Enquist et al., 2016; Maitner et al., 2018)). Instead, the focus of MOSAIC is on providing taxonomic

coverage to species with open-access structured population models available in the COMADRE (Salguero-

Gómez et al., 2016), COMPADRE (Salguero-Gómez et al., 2015), and PADRINO databases (Levin et al.

forthcoming) (See Fig. 3 and Fig. 4 for spatial scope and taxonomic scope with respect to structured

population databases, respectively). MOSAIC provides a much-needed interoperability between existing

databases that are germane to demography. By doing so, MOSAIC helps to fill critical data needs of

population ecologists and functional trait ecologists (see Fig. 5 for relevant covariance structure). Although

demographically motivated, MOSAIC affords opportunities for data to address novel questions more

broadly across ecology, evolution, and conservation biology.

Accessing MOSAIC

A guide for downloading the MOSAIC database can be accessed on the MOSAIC portal

(https://mosaicdatabase.web.ox.ac.uk). The MOSAIC database is open access and can be downloaded into

R using a few lines of code (see: https://mosaicdatabase.web.ox.ac.uk/download-database), which

download the MOSAIC data object as an S4 object (Rda file). The database can also be downloaded from

GitHub directly as an (.Rda) or comma separated value (.csv) file and loaded into R manually.

The MOSAIC portal hosts information pages and downloadable files about the database. Users can

download the *MOSAIC User Guide*, *vignettes*, and *data collection protocols* from the portal webpage

(https://mosaicdatabase.web.ox.ac.uk/vignettes and https://mosaicdatabase.web.ox.ac.uk/user-guide). The

User Guide specifies the classifications, precision, and data types for each trait field. The MOSAIC *User*

Guide details the metadata on the structure of data for each trait (*e.g.,* species, genus, mosaic index). An

updated list of the databases that are directly or indirectly addressed by the MOSAIC database is maintained

on the MOSAIC portal.

Data Sources

MOSAIC is both a *meta*-database (a database of databases) and a database in its own right,

containing new trait records from primary literature (Fig. 1). The MOSAIC database contains records

centralised from existing datasets where functional traits relevant to population ecologists can be openly

accessed and redistributed (*e.g.,* BIEN (Enquist et al., 2016; Maitner et al., 2018)). The records reflected in

the MOSAIC database do not encompass the entirety of the source databases but are instead partial

facsimiles of those databases that reflect records relevant to population databases (COMADRE,

COMPADRE, PADRINO). MOSAIC has three major components: (1) trait records sourced from existing

databases (22%); (2) trait records newly procured through searching the primary literature (71%); and (3)

trait record markers that direct users to additional records that are not included in the MOSAIC database

(7%). MOSAIC trait markers exist for one of two reasons: the database containing the records of interest

does not allow records to be accessed or limits redistribution rights behind individual registration and/or

specific use applications; or records in other databases contain multiple records for a species, which do not

currently fit within the data structure of version 1.0.0 of MOSAIC (see *Future Targets for MOSAIC*).

In addition to identifying whether a trait record is new, the "MOSAIC" attribute field also identifies

whether the attribute field (*i.e.,* trait name) is part of an existing database. For example, the MOSAIC field

might indicate that a record for specific leaf area is new for a specific species, and also that the attribute is

part of databases such as TRY (Kattge et al., 2020) or BIEN (Enquist et al., 2016; Maitner et al., 2018). By

contrast, a new species record for volancy would indicate that there are currently no databases that

systematically collect data on volancy attributes and therefore that all volancy records in the MOSAIC

database are new.

Organisation of Sources in MOSAIC

The MOSAIC attribute field is a factorial variable with three levels: MOSAIC-A, MOSAIC-B, and MOSAIC-C. The first of these levels, MOSAIC-A (Fig. 1), labels only records that reflect existing databases (*i.e.*, provenance of an existing data acquisition service); the second, MOSAIC-B, labels new records collected by the MOSAIC team that are in a trait field within the scope of an existing database initiative (*e.g.,* specific leaf area in BIEN); and the third, MOSAIC-C, labels new records collected by the MOSAIC team on traits for which there is not currently a database initiative centralising records. If a datum has been adopted from another dataset or database, then the relevant source is referenced in the *Database* attribute column. Note that this value will be "NA" for all MOSAIC-C records, logically. Over time, data sharing will move records in MOSAIC-B to MOSAIC-A as the MOSAIC-B traits are assimilated in the database networks that specialise on an existing trait (*i.e.,* data feedback; see Fig. 1).

Sources and Provenance of Records

Because of existing limitations on data access, some datasets cannot be transferred into MOSAIC. Where data exist outside of the MOSAIC platform, but have restricted access, the MOSAIC database directs users toward the appropriate database on a trait and taxa-specific level (see metaMOSAIC). The *MOSAIC User Guide* (Appendix S1) explains differences between data gaps that are yet to be reviewed and those that are true gaps (*e.g.,* volancy/flight capability of plants). Data were obtained through searching peer-reviewed records of ISI Web of Science, Scopus, and Google Scholar using key words pertinent to the species name and trait field in question (see Appendix S2 for a list of keywords queried). The MOSAIC team also searched the archives of data repositories, including the Dryad digital repository (http://datadryad.org; see Appendix S3 for a complete list of repositories reviewed – to be maintained hereafter on the MOSAIC portal), and reviewed in detail journal archives that have a high occurrence of data publishing, including *Nature Scientific Data*, *Methods in Ecology and Evolution*, and *Journal of Ecology*. MOSAIC also reflects a review of data from data aggregating servers, such as the open traits network (https://opentraits.org/), the ecological data wiki (https://ecologicaldata.org/), environmental data initiative (https://environmentaldatainitiative.org/), and databases that aggregate other databases (*e.g.,*

BIEN (Enquist et al., 2016; Maitner et al., 2018) and TRY (Kattge et al., 2020)). The complete set of key

words used in this review is detailed in a supplement to the *User Guide* posted on the MOSAIC portal. A

current list of databases reviewed in the development of MOSAIC is included in Appendix S3. Suggested

data sources and key terms can be submitted through the MOSAIC data portal.

Database Navigation

The *User Guide* contains detailed information on navigating the MOSAIC R data object. In general,

the data object can be queried with the syntax: `mosaic@trait.name@values[[mosaic.index]]`.

In the aforementioned syntax, the `trait.name` and `mosaic.index` should reflect one trait and one or

more species index values in the respective locations (*e.g.,* `mosaic@volancy@value[[93]]`). The

complete set of records for a trait can be viewed and assigned by calling the trait from the mosaic object

without an index specified (*e.g.,* `mosaic@volancy@value`). The mosaic index value reflects a unique

identifier that can be linked from COMADRE, COMPADRE, and PADRINO through the index field of

mosaic (*e.g.,* `comadre@data$MatrixID[comadre@data$SpeciesAccepted=="Alces`

`alces"]`). Associated metadata for each record can be queried by the following syntax:

`mosaic@trait.name@metadatafield[[mosaic.index]]`. Here, the `metadatafield`

should reflect one metadata element (*e.g.,* `mosaic@volancy@author[[93]]`). The complete

metadata for a species can be viewed using the `mos_meta()` function or searched manually by slots, or

by querying rows of the database in the comma separated value format

(https://mosaicdatabase.web.ox.ac.uk/navigating-mosaic). See the *User Guide* (Appendix S1) for more

information. The database is organised to be readily subsetted, including compatibility with Tidyverse

package (Wikham et al. 2019) syntax (see *Navigating MOSAIC*;

https://mosaicdatabase.web.ox.ac.uk/vignettes, also in Appendix S4). The MOSAIC object can be searched

in R by taxonomy, trait, locality, duration of study, sex, and other metadata fields applicable to COMADRE,

COMPADRE, and PADRINO (see *vignettes*, above, also Appendix SX). Trait values in the database are

measured in standardised units of measure for each trait (Table 1). Species names are specified in

conformance with the Catalogue of Life (www.catalogueoflife.org), also consistent with the COMPADRE,

COMADRE, and PADRINO databases. The *User Guide* provides specific guidelines for querying fields

within the data object in R and for negotiating the dataset in finer detail

(https://mosaicdatabase.web.ox.ac.uk/user-guide).

Error Reporting

The MOSAIC portal has an *Error Report* page for reporting potentially erroneous records, or to

query additional questions (https://mosaicdatabase.web.ox.ac.uk/suggested-additions-error-reporting; but

see also FAQ: https://mosaicdatabase.web.ox.ac.uk/frequently-asked-questions). Potential errors can be

reported anonymously or with contact information (*e.g.*, name, email). Decisions on reported errors will be

disclosed on the Error Report page (https://mosaicdatabase.web.ox.ac.uk/suggested-additions-error-

reporting) and to the reporting party if contact information is included in the request. If it is not possible to

submit errors via the portal, users can also to reach out by email to: mosaicdatabase@biology.ox.ac.uk.

Recommended Records

The MOSAIC portal has a *Recommended Records* page

(https://mosaicdatabase.web.ox.ac.uk/suggestion-additions) for reporting suggested records that are not

included in the MOSAIC database. Recommendations can be made with or without contact information.

Contact information will be used exclusively for clarifying questions and updating the commenter when

records are included. MOSAIC will report decisions on the *Recommendations Incorporated* page

(https://mosaicdatabase.web.ox.ac.uk/suggestion-additions). Users may also request new data fields to be

prioritised in future rollouts. Given the realities of limiting resources, The MOSAIC team will do their best

to include the requested records in future versions. Likewise, we also welcome submissions of data

collected by individuals via mosaicdatabase@biology.ox.ac.uk.

metaMOSAIC: licensed data, access limitations, and restricted redistribution of existing records and

databases

Not all datasets permit open use, dissemination, and redistribution of their trait data. Where

limitations on the data collation and redistribution apply, there may be application procedures, registration,

and other actions necessary for an individual to obtain access to specific trait records for analysis (*e.g.,*

TRY). MOSAIC centralises the metadata for datasets that do not allow data to be redistributed to help

navigate to relevant data resources outside the scope of open access. Records in limited access databases

can be searched in MOSAIC by taxonomic group and by trait. MOSAIC links researchers to application

materials for requesting access to those limited-access databases. The data access of licensed or non-open-

access databases is stored in a data object called `metaMOSAIC` that is an extension of the MOSAIC

database. The MOSAIC dataset thus provides data where it is accessible and `metaMOSAIC` guides

researchers to where data exist with registration. When searching fields in the MOSAIC database, the

`metaMOSAIC` adjunct dataset indicates if data are available in these ancillary sources (see *User Guide* in

Appendix S1 for more information) and provide links to pertinent sources and data through the provider.

`metaMOSAIC` is part of the main database object accessible through the MOSAIC portal.

Database Updates

The MOSAIC database will be updated as new data are added to the COMADRE, COMPADRE,

and PADRINO databases. Updates of the MOSAIC database will account for newly discovered data sources

and new literature that adds to or changes the species-level traits in the database, as well as correct errors

from earlier versions. New MOSAIC versions will be released periodically with a notice published on the

website, in the data object metadata, the mosaic GitHub page (https://github.com/mosaicdatabase), and

through updates in associated packages in conformance with standard semantic versioning (a three-part

version code reflecting major, minor, and patch updates, in respective positions). Updates will be published

to the mosaic portal (https://mosaicdatabase.web.ox.ac.uk), social media (Twitter: @mosaicdatabase), and

GitHub (https://github.com/mosaicdatabase). The MOSAIC portal includes a location for submission of

recommendations for additions or suggested changes to the database.

Future Targets of MOSAIC

Versioning

MOSAIC database updates and version history will be posted to the MOSAIC portal in the future.

Taxonomic Scope

Future versions of MOSAIC are scheduled for expansion to include 50 core traits across some

1,500 species (the current entirety of COMPADRE, COMADRE, and PADRINO; *MOSAIC target traits*;

see *User Guide* [Appendix S1]). The MOSAIC core traits list will guide the future roll-out of attributes

gathered and stewarded by the MOSAIC team (available on the MOSAIC portal website; see below), but

additional trait suggestions can be submitted through the online portal.

Interspecific Variation in Hierarchical Data Structures

Future versions of the MOSAIC database will adopt a hierarchical file structure that will

accommodate multiple records per species. Once records for COMADRE, COMPADRE, and PADRINO

are fully populated across the MOSAIC traits with mean, pooled, or other representative quantities (*e.g.,*

mean leaf size for all plants or adult bodymass of animals), secondary records will be added. Existing trait

databases may contain multiple records per species (see, for example, structure of COMADRE (Salguero-

Gómez et al., 2016), COMPADRE (Salguero-Gómez et al., 2015), BIEN (Enquist et al., 2016; Maitner et

al., 2018), TRY (Kattge et al., 2020)), although some databases host a single record per species' trait, such

as age and growth rate for animals in AnAge (De Magalhães & Costa, 2009). To facilitate research into

intra-specific trait variation (Albert, 2015; Messier, McGill, & Lechowicz, 2010; Siefert et al., 2015),

MOSAIC provides provenance of records, whether records were subject to selection or merger (means or

pooling), and fields that identify whether multiple records are known to exist. Where records for a given

species were isolated from existing databases, mean values are often retrieved, and the database sourcing

additional data is noted in the database under the attributed field ("Additional Trait Data Available" field).

citMOSAIC: Citizen Science

In addition to `metaMosaic`, which guides users to licenced data not reported in the MOSAIC

database, MOSAIC plans to roll out a database of identical structure to MOSAIC that gathers information

from citizen science datasets. Like MOSAIC, citMOSAIC will have three components (citMOSAIC-A,

citMOSAIC-B, citMOSAIC-C), reflecting the same relationship of databases and fields to MOSAIC (see

https://mosaicdatabase.web.ox.ac.uk/citizen-mosaic for more information). citMOSAIC will be kept

independent of the main MOSAIC database to avoid conflation of peer reviewed literature from citizen

science data. Where appropriate to use datasets together, the metadata, query functionality, and design of

citMOSAIC will mirror that of MOSAIC to promote interoperation of databases. citMOSAIC will be

downloadable from the MOSAIC portal website.

Cautionary Notes

Records in the MOSAIC database are gathered and standardised under the protocols detailed in the

User Guide (Appendix S1; also available through the MOSAIC portal). Users should be attentive to the

precision, levels, and context of data in MOSAIC when used for analysis. For those records in the dataset

that come from multiple individuals, we present them as statistical components (*e.g.,* minimum, maximum,

or mean trait values). Functional traits in MOSAIC may be estimated from populations studied in

COMADRE, COMPADRE, PADRINO, or other databases (Table S3). The potential temporal and spatial

mismatch between databases that are linked in an analysis merits close attention (Csergő et al., 2017). The

studies in the MOSAIC database also include research conducted by different investigators using

independent tools, technologies, sample designs, and study methods. The influence of research methods

and instruments on the error values in the dataset may require additional consideration for potential bias,

noise, and imprecision. Where more than one life history trait value exists for a given species, MOSAIC

users will need to determine whether averaging or selective filtering to one study is most appropriate in

view of the specifics of the given research question. In certain cases, trait values for a species might only

be available for a single st/age and therefore may not provide a complete picture of the trait variation

amongst st/ages. Users are encouraged to be cautious when contextualising the scope of representation of

the values in the database and their analyses.

Representation, Variance, and Bias in MOSAIC v1.0.0

MOSAIC can help to mechanistically examine the causes and correlates of demographic variability

across the thousands of animal and plant species housed in COMPADRE, COMADRE, and PADRINO. In

its version 1.0.0, MOSAIC contains a high degree of variance in trait values known to shape demographic

outcomes across major taxonomic groups. For example, determinant growth is present for 0% of

Amphibians and Birds and 100% of Bivalves, while volancy is present for 0% in Reptiles, Amphibians,

and Bony Fish and 77% in Birds (Fig. 2). Animal adult biomass and plant height follow lognormal trait

distributions (see Appendix S5). 11% of mammals are indeterminate growers *vs.* 54% of reptiles and 0%

birds, and 95% of mammals are monogamous compared to 100% of reptiles and 20% of birds in MOSAIC.

Recent studies have examined how different vital rates are explained by functional traits (e.g., Carmona,

Bello, Azcarate, Mason, & Peco 2019; Pistón et al. 2018; Adler et al. 2014; Poorter et al. 2008). However,

understanding how trait variation across taxa translates to demographically influential properties remains

underdeveloped.

MOSAIC's initial release (v1.0.0) includes records for all major regions of the globe for which we

have structured population models (Fig.3). Nevertheless, species trait values are not necessarily gathered

from the same localities as population models (see *future directions* for more information on systematising

spatial mismatch). This is an important consideration for users of MOSAIC (and more generally of trait-

based approaches) wishing to bring together functional traits and demographic rates, as traits and vital rates

are known to vary considerably within the same species across spatial scales (Violle et al. 2012; Albert et

al. 2010; Bolnick et al. 2011). Moreover, while there is at least one trait for each of these locations, the data

density remains variable. Thus, records are not necessarily representative of the global spread or the full

spatial scope of MOSAIC. For example, the highest complete coverage for MOSAIC traits is concentrated

toward localities with the with the longest-term demographic models (see COMPADRE locations

associated with MOSAIC records).

Phylogenetically, the initial release of MOSAIC is somewhat limited. Version 1.0.0 covers 300 of

the 1,400 species currently available in COMPADRE, COMADRE and PADRINO. However, MOSAIC

trait data are well distributed across clades (Fig. 4). While there is not a highly skewed phylogenetic

concentration with respect to the existing structured population models or clustering of records into small

groups across the Tree of Life, phylogenetic density of MOSAIC records remains low. Therefore,

information from the MOSAIC database may be limited for a given genus or order and, as such, should be

approached with caution. In future versions of MOSAIC, the phylogenetic bias is expected to diminish with

more samples and stronger phylogenetic representation.

Covariance across traits is also an important source of confound in existing analyses linking traits

and vital rates. Positive and negative correlations across traits that have independent influences on vital

rates can create apparent associations of demographic properties with traits, spuriously functionalising non-

functional traits (McElreath 2021). Disentangling the relevance of key axes of trait variation for their

demographic influences demands a clear quantification of the direction and strength networks of trait

associations, trade-offs, and demographic consequences. Population biologists seek to understand not only

how individual traits relate to different aspects of demographic performance (*e.g.,* population growth rate,

risk of quasi-extinction, *etc.*), but also understanding how trait syndromes shape those demographic

outcomes. MOSAIC presents a highly varied covariance structure in trait values for the examined 300

species. For example, without *a priori* expectation, indeterminate growth and regeneration traits show

strong correlation (r=0.51; Fig. 5), which could influence each other's effects on vital rates. The same could

be argued for the correlation between volancy and reproductive strategy (r=0.28 with monogamy; Fig. 5).

The MOSAIC database can be used as a platform to showcase the lack of overlap between trait and

vital rate data for the same species. This picture calls for a more systematic way address global biases in

ecological data quantification/collection. Even where we have complete information about species in the

COMADRE, COMPADRE, and PADRINO databases, we are subject to the constraints and biases of those

datasets, such as spatial bias toward high-GDP countries and the phylogenetic bias toward temperate

regional perennial plants (Kendall et al. 2019; Römer et al. 2021). The compounding of error and density

across datasets highlights the need to prioritise stronger representation of functional traits linked with

demography. The standardised framework of MOSAIC is an ideal platform to work from to achieve this

goal.

In view of potential biases introduced by low sampling density and the patchiness of cover in traits,

users of the database are advised to consult the literature about the representativeness and congruency

between MOSAIC data and related trait diversity within clades. Users need to be mindful of the scope of

the questions that they are setting out to answer and to be aware of the influence of sample sizes and

coherency or heterogeneity of traits across different taxonomic levels.

Discussion

From Databases to Data Networks

Broad aperture digitisation efforts (*e.g.,* BIEN (Enquist et al., 2016; Maitner et al., 2018) and TRY

(Kattge et al., 2020)) have helped resolve many answers to demographic questions. Examples include

whether there are trait spectra and key trade-off patterns amongst functional traits and whether these are

correlated with particular environments and life history strategies (Laughlin, 2018; Reich et al., 2003;

Wilkes et al., 2020), Trait-based ecology and Trait Driver Theory (Enquist et al., 2015) are indebted to such

opportunity-driven research programmes. More generally, however, the trait-based ecology paradigm has

failed to support clear answers to many research questions of central interest to demographers (Bellier,

Kéry, & Schaub, 2018; Laughlin et al., 2020). This limited reach of the functional trait programme coincides

with a dearth of species-specific overlap across the range of functional traits that are collected by the

functional trait databases.

The proliferation of databases and open data initiatives over the last two decades (Cheruvelil &

Soranno, 2018) evidences an interest in improving both data access and data usability (Gallagher et al.,

2020; Michener, 2006; Whitlock, 2011). While existing databases standardise trait fields, collate records,

and link associated metadata, existing databases often store data for simple, quantitative traits. Relatively

few ecological trait databases store diverse data types (such as rate arrays, population time series,

physiological rates at different structural levels, and habitat shape files) that may be associated with

multidimensional, ecological study systems (but see CESTES (Jeliazkov et al., 2020), GFBio (Diepenbroek

et al., 2014), DarwinCore (Wieczorek et al., 2012)).

The digitisation and standardisation of *existing* data and their integration with complementary, new

data presents a growing set of challenges and opportunities in ecology (Salguero-Gómez, Jackson,

Gascoigne 2021). Efforts to gap-fill records can leverage the value of existing datasets while expediting the

schedule for specific research outcomes. As trait datasets grow, the importance of targeted, gap-filling

initiatives to address bias and to capitalise on existing data will also increase (Conde et al., 2019). The value

of existing records is further enhanced through improvements in the interoperability of datasets. Much of

this work is done manually, at a high cost, and with little support from funding agencies (Salguero-Gómez,

Jackson, Gascoigne 2021), and yet it has been effective at facilitating research and creating new value for

old data. In recent years, initiatives have sought to improve the interoperability of datasets by guiding

prospective data structure or retroactively harmonising existing datasets. These include programmes that

develop universal standards to improve global interoperability (such as DarwinCore (Wieczorek et al.,

2012) and Frictionless [https://frictionlessdata.io/]data standards) or that contain guidelines for data

metastructure (such as the FAIR principles (findability, accessibility, interoperability, and reusability, *sensu*

Wilkinson et al., 2016) and the OpenTraits framework (Gallagher et al., 2020)). These initiatives address

emerging and scaling challenges of ecoinformatics, such as the protocols by which we share data, search

data, and preserve provenance within hierarchical data storage structures. These protocols will be essential

in centralising datasets as diverse as government monitoring datasets (*e.g.,* those stored in U.S. Data

clearing houses [https://www.data.gov; https://www.dataone.org/]; National Biodiversity Atlas

[https://nbnatlas.org/]); centralised monitoring and experimental networks (*e.g.,* LTER and NEON), raw or

reanalysed remote sensing datasets (*e.g.,* Landsat data, NASA EarthData datasets, ERA-5 data), and private

datasets (https://www.natureserve.org/) that will demand versatile and navigationally efficient data

structures.

Population ecology has benefitted from widespread open-access databases but requires further

dataset integration to answer its central questions. Understanding whether and how some morphological or

physiological traits predict demographic outcomes and why others fail to do so is of central interest to

questions in physiological, population, and community ecology (Adler et al., 2014; Chalmandrier et al.,

2021; Laughlin et al., 2020; Yang, Cao, & Swenson, 2018). Population ecologists routinely use data that

are distributed across a wide range of databases. Comparative and macroecological researchers use

phylogenies (*e.g.*, Daskalova, Myers-Smith, & Godlee, 2020; Levin, Crandall, Pokoski, Stein, & Knight,

2020; Roper, Capdevila, & Salguero-Gómez, 2021), adult bodymass (*e.g.*, Capdevila et al., 2020; Healy et

al., 2014; Terry, O'Sullivan, & Rossberg, 2022; Williams, McRae, Freeman, Capdevila, & Clements, 2021), and high-resolution, global climate information (*e.g.*, Daskalova, Bowler, Myers-Smith, & Dornelas, 2021; Paniw, Maag, Cozzi, Clutton-Brock, & Ozgul, 2019; Stenseth & Mysterud, 2002) to answer relevant biological, evolutionary and ecological questions and to contextualise their findings. Population ecologists frequently examine a subset of physiological, morphological, and behavioural attributes associated with demographic outcomes ("functional traits", *sensu* Violle et al., 2007). The trait-based research programme seeks to, among other aims (*e.g.*, Díaz et al., 2016), identify the intrinsic and extrinsic regulators of vital rates and the causes of variation and constraints on possible trait values (Buckley & Puy, 2022; Salguero-Gómez, Violle, Gimenez, & Childs, 2018). Not all traits predict demographic outcomes and functional traits may exercise influence on only a few demographic pathways (Pistón et al., 2019; Salguero-Gómez & Laughlin, 2021). The answers to these questions rely on the existence of vital rate and trait data, the overlap of which has been limited in the absence of targeted attention. For instance, of the hundreds of thousands of records available across thousands of plant species in TRY (Kattge et al., 2020) and over 345 plant species in COMPADRE (Salguero-Gómez et al. 2015, COMPADRE Ver 1.0.0), Adler et al. (2014) report functional trait-vital rate relationships for only 222 plant species due to their limited data overlap.

Ecological data are complex and their structures will need consistent rules to link datasets together. It will be important for future datasets to adopt hierarchical database designs that render large, thematically, and structurally diverse data to be readily locatable and usable without expert knowledge. Here, we show one such example in the scope of comparative research, using thematic groups and a strategy of achieving adequate record breadth before revisiting depth of records for specific species. The need for open access data, integrated workflows, and interoperable data systems is increasing with the scaling of data collection through use of robotics and technologies. The gaps in existing data systems, interoperability, and data acquisition can be filled strategically for specific applications, offering targeted and efficient dataset development. With data interoperability guiding the structure of new datasets, the modular development of area-specific datasets will enable more generalised use over time and help meet the aims of existing database initiatives.

ACKNOWLEDGEMENTS

This work was supported by a NERC IRF (NE/M018458/1) to RS-G. RCRC acknowledges postdoctoral

support from the Regional Valencian Government and the European Social Fund (APOSTD/2020/090).

ORCID

Connor Bernard: 0000-0003-2175-5712

Gabriel Santos: 0000-0001-7991-8807

Jacques Deere: 0000-0001-6736-2223

Roberto Rodriguez-Caro: 0000-0003-2321-9497

Pol Capdevila: 0000-0002-2842-4302

Erik Kusch: 0000-0002-4984-7646

Samuel J L Gascoigne: 0000-0002-2984-1810

John Jackson: 0000-0002-1401-3035

Roberto Salguero-Gómez: 0000-0002-6085-4433

References

Adler, P. B., Salguero-Gómez, R., Compagnoni, A., Hsu, J. S., Ray-Mukherjee, J., Mbeau-Ache, C., & Franco, M. (2014). Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(27), 10019. doi: 10.1073/pnas.1410430111

Albert, C. H. (2015). Intraspecific trait variability matters. *Journal of Vegetation Science*, *26*(1), 7–8. doi: 10.1111/jvs.12240

Albert, C.H., Thuiller, W., Yoccoz, N.G., Soudant, A., Boucher, F., Saccone, P. & Lavorel, S. (2010) Intraspecific functional variability: extent, structure and source of variation. *The Journal of Ecology*, 98, 604–613.

Bellier, E., Kéry, M., & Schaub, M. (2018). Relationships between vital rates and ecological traits in an avian community. *Journal of Animal Ecology*, *87*(4), 1172–1181. doi: 10.1111/1365-2656.12826

Boettiger, C., & Temple Lang, D. (2012). Treebase: An R package for discovery, access and manipulation of online phylogenies. *Methods in Ecology and Evolution*, *3*(6), 1060–1066. doi: 10.1111/j.2041-210X.2012.00247.x

Bolnick, D.I., Amarasekare, P., Araujo, M.S., Burger, R., Levine, J.M., Novak, M. et al. (2011). Why intraspecific trait variation matters in community ecology. *Trend Ecol. Evol.*, 24, 183–192.

Borghetti, M., Gentilesca, T., Colangelo, M., Ripullone, F., & Rita, A. (2020). Xylem Functional Traits as Indicators of Health in Mediterranean Forests. *Current Forestry Reports*, *6*(3), 220–236. doi: 10.1007/s40725-020-00124-5

Buckley, Y. M., & Puy, J. (2022). The macroecology of plant populations from local to global scales. *New Phytologist*, *233*(3), 1038–1050. doi: 10.1111/nph.17749

Capdevila, P., Beger, M., Blomberg, S. P., Hereu, B., Linares, C., & Salguero-Gómez, R. (2020). Longevity, body dimension and reproductive mode drive differences in aquatic versus terrestrial life-history strategies. *Functional Ecology*, *34*(8), 1613–1625. doi: 10.1111/1365-2435.13604

Carmona, C. P., de Bello, F., Azcarate, F. M., Mason, N. W. H., & Peco, B. (2019). Trait hierachies and intraspecific variability drive competitive interactions in Mediterranean annual plants. *Journal of Ecology*, 107, 2078–2089. https://doi.org/10.1111/1365-2745.13248

Caswell, H. (2001). *Matrix population models* (2nd editio). Sunderland, MA: Sinauer.

Chalmandrier, L., Hartig, F., Laughlin, D. C., Lischke, H., Pichler, M., Stouffer, D. B., & Pellissier, L. (2021). Linking functional traits and demography to model species-rich communities. *Nature Communications*, *12*(1). doi: 10.1038/s41467-021-22630-1

Cheruvelil, K. S., & Soranno, P. A. (2018). Data-intensive ecological research is catalyzed by open science and team science. *BioScience*, *68*(10), 813–822. doi: 10.1093/biosci/biy097

Conde, D. A., Staerk, J., Colchero, F., da Silva, R., Schöley, J., Maria Baden, H., … Vaupel, J. W. (2019). Data gaps and opportunities for comparative and conservation biology. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(19), 9658–9664. doi: 10.1073/pnas.1816367116

Consortium, E. P. (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature*, *489*(7414), 57–74. doi: 10.1038/nature11247

Crouse, D. T., Crowder, L. B., & Caswell, H. (1987). A stage-based population model for loggerhead sea turtles and implications for conservation. *Ecology*, *68*(5), 1412–1423.

Csergő, A. M., Salguero-Gómez, R., Broennimann, O., Coutts, S. R., Guisan, A., Angert, A. L., … Buckley, Y. M. (2017). Less favourable climates constrain demographic strategies in plants. *Ecology Letters*, *20*(8), 969–980. doi: 10.1111/ele.12794

Culina, A., Baglioni, M., Crowther, T. W., Visser, M. E., Woutersen-Windhouwer, S., & Manghi, P. (2018). Navigating the unfolding open data landscape in ecology and evolution. *Nature Ecology and Evolution*, *2*(3), 420–426. doi: 10.1038/s41559-017-0458-2

Daskalova, G. N., Bowler, D., Myers-Smith, I. H., & Dornelas, M. (2021). Representation of global change

drivers across biodiversity datasets. *EcoEvoRxiv*, 1–36. Retrieved from https://doi.org/10.32942/osf.io/db4s7

Daskalova, G. N., Myers-Smith, I. H., & Godlee, J. L. (2020). Rare and common vertebrates span a wide spectrum of population trends. *Nature Communications*, *11*(1), 1–13. doi: 10.1038/s41467-020-17779-0

Davy, R., & Kusch, E. (2021). Reconciling high resolution climate datasets using KrigR. *Environmental Research Letters*.

De Magalhães, J. P., & Costa, J. (2009). A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology*, *22*(8), 1770–1774. doi: 10.1111/j.1420-9101.2009.01783.x

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., … Gorné, L. D. (2016). The global spectrum of plant form and function. *Nature*, *529*(7585), 167–171. doi: 10.1038/nature16489

Diepenbroek, M., Glöckner, F. O., Grobe, P., Güntsch, A., Huber, R., König-Ries, B., … Triebel, D. (2014). Towards an Integrated Biodiversity and Ecological Research Data Management and Archiving Platform : The .... *Informatik 2014 – Big Data Komplexität Meistern. GI-Edition: Lecture Notes in Informatics (LNI) – Proceedings 232*, (November), 1711–1721. Retrieved from https://dl.gi.de/handle/20.500.12116/2782

Dornelas, M., Antão, L. H., Moyes, F., Bates, A. E., Magurran, A. E., Adam, D., … Zettler, M. L. (2018). BioTIME: A database of biodiversity time series for the Anthropocene. *Global Ecology and Biogeography*, *27*(7), 760–786. doi: 10.1111/geb.12729

Easterling, M. R., Ellner, S. P., & Dixon, P. M. (2000). Size-specific sensitivity: Applying a new structured population model. *Ecology*, *81*(3), 694–708.

Edwards, J. L., Lane, M. A., & Nielsen, E. S. (2000). Interoperability of biodiversity databases: Biodiversity information on every desktop. *Science*, *289*(5488), 2312–2314. doi: 10.1126/science.289.5488.2312

Enquist, B. J., Condit, R., Peet, R. K., Schildhauer, M., & Thiers, B. (2016). The Botanical Information and Ecology Network (BIEN): Cyberinfrastructure for an integrated botanical information network to investigate the ecological impacts of global climate change on plant biodiversity. *PeerJ Preprints*.

Enquist, B. J., Feng, X., Boyle, B., Maitner, B., Newman, E. A., Jørgensen, P. M., … McGill, B. J. (2019). The commonness of rarity: Global and future distribution of rarity across land plants. *Science Advances*, *5*(11), 1–14. doi: 10.1126/sciadv.aaz0414

Enquist, B. J., Norberg, J., Bonser, S. P., Violle, C., Webb, C. T., Henderson, A., … Savage, V. M. (2015). Scaling from traits to ecosystems: developing a general trait driver theory via integrating trait-based and metabolic scaling theories. In *Advances in ecological research* (pp. 249–318).

Farley, S. S., Dawson, A., Goring, S. J., & Williams, J. W. (2018). Situating ecology as a big-data science: Current advances, challenges, and solutions. *BioScience*, *68*(8), 563–576. doi: 10.1093/biosci/biy068

Field, D., Sansone, S., Collis, A., Booth, T., Dukes, P., Gregurick, S. K., … Remington, K. (2009). 'Omics Data Sharing. *Science*, *326*, 234–236.

Fortuna, M. A., Ortega, R., & Bascompote, J. (2014). The web of life. *ArXiv*. doi: 10.4324/9780203410134

Froese, R., & Pauly, D. (2010). *FishBase*. Retrieved from http://www.ices.dk/sites/pub/CM Doccuments/1992/L/1992_L10.pdf

Gallagher, R. V., Falster, D. S., Maitner, B. S., Salguero-Gómez, R., Vandvik, V., Pearse, W. D., … Enquist, B. J. (2020). Open Science principles for accelerating trait-based science across the Tree of Life. *Nature Ecology and Evolution*, *4*(3), 294–303. doi: 10.1038/s41559-020-1109-6

Guerrero-Ramírez, N. R., Mommer, L., Freschet, G. T., Iversen, C. M., McCormack, M. L., Kattge, J., … Weigelt, A. (2021). Global root traits (GRooT) database. *Global Ecology and Biogeography*, *30*(1), 25–37. doi: 10.1111/geb.13179

Healy, K., Guillerme, T., Finlay, S., Kane, A., Kelly, S. B. A., McClean, D., … Cooper, N. (2014). Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1784). doi: 10.1098/rspb.2014.0298

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., … Thépaut, J. N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*,

146(730), 1999–2049. doi: 10.1002/qj.3803

J.R. Wilmoth, K. Andreev, D. Jdanov, and D. A. G. (2007). Methods Protocol for the Human Mortality Database. *Database*, *2007*(June 2000), 1–80.

Jasilioniene, A., Jdanov, D. A., Sobotka, T., Andreev, E. M., Zeman, K., & Shkolnikov, V. M. (2015). Methods Protocol for the Human Fertility Database. In *Max Planck Institute for Demographic Research*. Retrieved from http://www.humanfertility.org/Docs/methods.pdf

Jeliazkov, A., Mijatovic, D., Chantepie, S., Andrew, N., Arlettaz, R., Barbaro, L., … Chase, J. M. (2020). A global database for metacommunity ecology, integrating species, traits, environment and space. *Scientific Data*, *7*(1), 1–15. doi: 10.1038/s41597-019-0344-7

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, *491*(7424), 444–448. doi: 10.1038/nature11631

Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D. L., … Purvis, A. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology*, *90*(9), 2648–2648. doi: 10.1890/08-1494.1

Jongejans, E., Shea, K., Skarpaas, O., Kelly, D., & Ellner, S. P. (2011). Importance of individual and environmental variation for invasive species spread: A spatial integral projection model. *Ecology*, *92*(1), 86–97. doi: 10.1890/09-2226.1

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., … Wirth, C. (2020). TRY plant trait database – enhanced coverage and open access. *Global Change Biology*, *26*(1), 119–188. doi: 10.1111/gcb.14904

Kissling, W. D., Walls, R., Bowser, A., Jones, M. O., Kattge, J., Agosti, D., … Guralnick, R. P. (2018). Towards global data products of Essential Biodiversity Variables on species traits. *Nature Ecology and Evolution*, *2*(10), 1531–1540. doi: 10.1038/s41559-018-0667-3

Laughlin, D. C. (2018). Rugged fitness landscapes and Darwinian demons in trait-based ecology. *New Phytologist*, *217*(2), 501–503. doi: 10.1111/nph.14908

Laughlin, D. C., Gremer, J. R., Adler, P. B., Mitchell, R. M., & Moore, M. M. (2020). The Net Effect of Functional Traits on Fitness. *Trends in Ecology and Evolution*, *35*(11), 1037–1047. doi: 10.1016/j.tree.2020.07.010

Levin, S. C., Crandall, R. M., Pokoski, T., Stein, C., & Knight, T. M. (2020). Phylogenetic and functional distinctiveness explain alien plant population responses to competition: Phylogeny and traits explain dominance. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1930). doi: 10.1098/rspb.2020.1070rspb20201070

Lintulaakso, K. (2013). *MammalBase—database of recent mammals*.

Madin, J. S., Anderson, K. D., Andreasen, M. H., Bridge, T. C. L., Cairns, S. D., Connolly, S. R., … Baird, A. H. (2016). The Coral Trait Database, a curated database of trait information for coral species from the global oceans. *Scientific Data*, *4*, 170174. doi: 10.1038/sdata.2017.174

Maitner, B. S., Boyle, B., Casler, N., Condit, R., Donoghue, J., Durán, S. M., … Enquist, B. J. (2018). The bien r package: A tool to access the Botanical Information and Ecology Network (BIEN) database. *Methods in Ecology and Evolution*, *9*(2), 373–379. doi: 10.1111/2041-210X.12861

Maldonado, C., Molina, C. I., Zizka, A., Persson, C., Taylor, C. M., Albán, J., … Antonelli, A. (2015). Estimating species diversity and distribution in the era of Big Data: To what extent can we trust public databases? *Global Ecology and Biogeography*, *24*(8), 973–984. doi: 10.1111/geb.12326

Marx, V. (2013). The big challenges of big data. *Nature*, *498*, 255–260.

Maurer, S. M., Firestone, R. B., & Scriver, C. R. (2000). Science's neglected legacy: Large, sophisticated databases cannot be left to chance and improvisation. *Nature*, *405*(6783), 117–120. doi: 10.1038/35012169

McElreath, R. (2021) *Statistical Rethinking.* London, UK: Chapman and Hall/CRC.

Messier, J., McGill, B. J., & Lechowicz, M. J. (2010). How do traits vary across ecological scales? A case for trait-based ecology. *Ecology Letters*, *13*(7), 838–848. doi: 10.1111/j.1461-0248.2010.01476.x

Metcalf, C. J. E., Graham, A. L., Martinez-Bakker, M., & Childs, D. Z. (2016). Opportunities and challenges of Integral Projection Models for modelling host-parasite dynamics. *Journal of Animal*

Ecology, *85*(2), 343–355. doi: 10.1111/1365-2656.12456

Michener, W. K. (2006). Meta-information concepts for ecological data management. *Ecological Informatics*, *1*(1), 3–7. doi: 10.1016/j.ecoinf.2005.08.004

Middleton, O., Svensson, H., Scharlemann, J. P. W., Faurby, S., & Sandom, C. (2021). CarniDIET 1.0: A database of terrestrial carnivorous mammal diets. *Global Ecology and Biogeography*, *30*(6), 1175–1182. doi: 10.1111/geb.13296

Monnet, A. C., Cilleros, K., Médail, F., Albassatneh, M. C., Arroyo, J., Bacchetta, G., … Leriche, A. (2021). WOODIV, a database of occurrences, functional traits, and phylogenetic data for all Euro-Mediterranean trees. *Scientific Data*, *8*(1), 1–11. doi: 10.1038/s41597-021-00873-3

Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S., … Menges, E. S. (2008). Longevity can buffer plant and animal populations against chnaging climatic variability. *Ecology*, *89*(1), 19–25. Retrieved from http://www.esajournals.org/doi/pdf/10.1890/07-0774.1

Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. K. M. (2015). An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles. *Ecology*, *96*(11). doi: 10.1890/15-0846r.1

Nadrowski, K., Ratcliffe, S., Bönisch, G., Bruelheide, H., Kattge, J., Liu, X., … Wirth, C. (2013). Harmonizing, annotating and sharing data in biodiversity-ecosystem functioning research. *Methods in Ecology and Evolution*, *4*(2), 201–205. doi: 10.1111/2041-210x.12009

Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017). AmphiBIO, a global database for amphibian ecological traits. *Scientific Data*, *4*, 1–7. doi: 10.1038/sdata.2017.123

Ozgul, A., Childs, D. Z., Oli, M. K., Armitage, K. B., Blumstein, D. T., Olson, L. E., … Coulson, T. (2010). Coupled dynamics of body mass and population growth in response to environmental change. *Nature*, *466*(7305), 482–485. doi: 10.1038/nature09210

Paniw, M., Maag, N., Cozzi, G., Clutton-Brock, T., & Ozgul, A. (2019). Life history responses of meerkats to seasonal changes in extreme environments. *Science*, *363*(6427), 631–635. doi: 10.1126/science.aau5905

Pistón, N., de Bello, F., Dias, A. T. C., Götzenberger, L., Rosado, B. H. P., de Mattos, E. A., … Carmona, C. P. (2019). Multidimensional ecological analyses demonstrate how interactions between functional traits shape fitness and life history strategies. *Journal of Ecology*, *107*(5), 2317–2328. doi: 10.1111/1365-2745.13190

Poorter, L., Wright, S.J., Paz, H., Ackerly, D.D., Condit, R., Ibarra-Manriques, G. et al. (2008) Are functional traits good predictors of demographic rates? Evidence from five Neotropical forests. *Ecology*, 89, 1908–1920.

Reich, P. B., Wright, I. J., Cavender-Bares, J., Craine, J. M., Oleksyn, J., Westoby, M., & Walters, M. B. (2003). The evolution of plant functional variation: Traits, spectra, and strategies. *International Journal of Plant Sciences*, *164*(SUPPL. 3). doi: 10.1086/374368

Reichman, O. J., Jones, M. B., & Schildhauer, M. P. (2011). Challenges and opportunities of open data in ecology. *Science*, *331*(6018), 703–705. doi: 10.1126/science.1197962

Roper, M., Capdevila, P., & Salguero-Gómez, R. (2021). Senescence: Why and where selection gradients might not decline with age. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1955). doi: 10.1098/rspb.2021.0851

Rowcliffe, J. M., Jansen, P. A., Kays, R., Kranstauber, B., & Carbone, C. (2016). Wildlife speed cameras: measuring animal travel speed and day range using camera traps. *Remote Sensing in Ecology and Conservation*, *2*(2), 84–94. doi: 10.1002/rse2.17

Salguero-Gómez, R., Jackson, J., & Gascoigne, S.J.L. (2021). Four key challenges in the open data revolution. *Journal of Animal Ecology*, *90(9), 2000-2004*. doi: 10.1111/1365-2656.13567

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Bein, C., de Buhr, H., Farack, C., … Vaupel, J. W. (2016). COMADRE: A global data base of animal demography. *Journal of Animal Ecology*. doi: 10.1111/1365-2656.12482

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Buckley, Y. M., Che-Castaldo, J., Caswell, H., … Vaupel, J. W. (2015). The compadre Plant Matrix Database: An open online repository for plant demography.

Journal of Ecology. doi: 10.1111/1365-2745.12334

Salguero-Gómez, R., & Laughlin, D. C. (2021). Not all traits are functional: the Panglossian paradigm. *Authorea*, (December 2021). doi: 10.22541/au.163940711.10447233/v1

Salguero-Gómez, R., Violle, C., Gimenez, O., & Childs, D. (2018). Delivering the promises of trait-based approaches to the needs of demographic approaches, and vice versa. *Functional Ecology*, *32*(6), 1424–1435. doi: 10.1111/1365-2435.13148

Santini, L., Isaac, N. J. B., & Ficetola, G. F. (2018). TetraDENSITY: A database of population density estimates in terrestrial vertebrates. *Global Ecology and Biogeography*, *27*(7), 787–791. doi: 10.1111/geb.12756

Schneider, F. D., Fichtmueller, D., Gossner, M. M., Güntsch, A., Jochum, M., König-Ries, B., … Simons, N. K. (2019). Towards an ecological trait-data standard. *Methods in Ecology and Evolution*, *10*(12), 2006–2019. doi: 10.1111/2041-210X.13288

Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., … Wardle, D. A. (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant communities. *Ecology Letters*, *18*(12), 1406–1419. doi: 10.1111/ele.12508

Smallegange, I. M., Deere, J. A., & Coulson, T. (2014). Correlative changes in life-history variables in response to environmental change in a model organism. *American Naturalist*, *183*(6), 784–797. doi: 10.1086/675817

Stenseth, N. C., & Mysterud, A. (2002). Climate, changing phenology, and other life history traits: Nonlinearity and match-mismatch to the environment. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(21), 13379–13381. doi: 10.1073/pnas.212519399

Swenson, N. G., Worthy, S. J., Eubanks, D., Iida, Y., Monks, L., Petprakob, K., … Zambrano, J. (2020). A reframing of trait–demographic rate analyses for ecology and evolutionary biology. *International Journal of Plant Sciences*, *181*(1), 33–43. doi: 10.1086/706189

Terry, J. C. D., O'Sullivan, J. D., & Rossberg, A. G. (2022). No pervasive relationship between species size and local abundance trends. *Nature Ecology and Evolution*, *6*(2), 140–144. doi: 10.1038/s41559-021-01624-8

Troia, M. J., & McManamay, R. A. (2016). Filling in the GAPS: evaluating completeness and coverage of open-access biodiversity databases in the United States. *Ecology and Evolution*, *6*(14), 4654–4669. doi: 10.1002/ece3.2225

Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., … Messier, J. (2012). The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology & Evolution*, *27*, 244–252.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! *Oikos*, *116*(5), 882–892. doi: 10.1111/j.2007.0030-1299.15559.x

Vitousek, M. N., Johnson, M. A., & Husak, J. F. (2018). Illuminating endocrine evolution: The power and potential of large-scale comparative analyses. *Integrative and Comparative Biology*, *58*(4), 712–719. doi: 10.1093/icb/icy098

Whitlock, M. C. (2011). Data archiving in ecology and evolution: Best practices. *Trends in Ecology and Evolution*, *26*(2), 61–65. doi: 10.1016/j.tree.2010.11.006

Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., … Vieglais, D. (2012). Darwin core: An evolving community-developed biodiversity data standard. *PLoS ONE*, *7*(1). doi: 10.1371/journal.pone.0029715

Wilkes, M. A., Edwards, F., Jones, J. I., Murphy, J. F., England, J., Friberg, N., … Brown, L. E. (2020). Trait-based ecology at large scales: Assessing functional trait correlations, phylogenetic constraints and spatial variability using open data. *Global Change Biology*, *26*(12), 7255–7267. doi: 10.1111/gcb.15344

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., … Mons, B. (2016). Comment: The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*, *3*, 1–9. doi: 10.1038/sdata.2016.18

Williams, N. F., McRae, L., Freeman, R., Capdevila, P., & Clements, C. F. (2021). Scaling the extinction

754   vortex: Body size as a predictor of population dynamics close to extinction events. *Ecology and*
755   *Evolution*, *11*(11), 7069–7079. doi: 10.1002/ece3.7555
756   Yang, J., Cao, M., & Swenson, N. G. (2018). Why Functional Traits Do Not Predict Tree Demographic
757   Rates. *Trends in Ecology and Evolution*, *33*(5), 326–336. doi: 10.1016/j.tree.2018.03.003
758
759

760   **Table 1**: Variable names and meaning contained in the MOSAIC Database, organised into seven

761   general trait domains. A more detailed description can be found in the MOSAIC user guide found

762   at: https://mosaicdatabase.web.ox.ac.uk/.

Aspect	Variable	Description	Units
Taxonomy (Parallels COM(P)ADRE)	1 *Species Author*	Taxonomic species name as used by the author(s) in the publication. When more than one study exists for the same species, these are given sequential numeric suffixes (e.g. Ursus_americanus, Ursus_americanus_2, etc.)	NA
	2 *Species Accepted*	Currently accepted taxonomic name according to the Catalogue of Life (www.catalogueoflife.org). See the Supplementary Online Material S3 for an R script to check accepted and synonym names from SpeciesAuthor above	NA
	3 *Common Name*	English common name of SpeciesAccepted	NA
	4 *Family*	Taxonomic family of study species	NA
	5 *Order*	Taxonomic order of study species	NA
	6 *Class*	Taxonomic class of study species	NA
	7 *Phylum*	Taxonomic phylum of study species	NA
	8 *Kingdom*	Taxonomic Kingdom of study species	NA
Species information General	9 *Organism type*	General plant/algae type, based mainly on architectural organisation. The species was assigned to one of these possible values using the description of plant growth type provided by the author and other external sources (e.g. other publications)	Categorical/ Factor
	10 *DicotMonocot*	Whether species is a dicot or monocot	Categorical/ Factor
	11 *AngioGymno*	Whether species is an angiosperm or a gymnosperm	Categorical/ Factor
Source information	12 *Authors*	Surname (family name) of all author	NA
	13 *Journal*	The document from which data were sourced.	NA
	14 *YearPublication*	Year of publication	NA
	15 *DOI*	Digital Object Identifier number	NA
Traits	01 *Biomass*	Mean mass of an individual/whole-organism. (Plants) Plant mass is measured as aboveground dry mass. See "Further information" for additional information on belowground biomass.	Grams
	02 Height	(Plants) Mean height of the whole organism/whole individual from surface (i.e. substrate) to tallest vertical extremity. (Animals) Mean ventro-dorsal length. (i.e. for terrestrial quadrupeds, vertical distance from the ground to the top of the shoulder. For marine bony fishes, from the top of the high point of the dorsal fin to the base of the ventral ridge or ventral fin, whichever is a longer distance perpendicular to the anterior-posterior axis.)	Centimetres
	03 Growth Determination	Whether a species exhibits growth determinacy or not.	Categorical/ Factorial
	04 Regeneration	Capacity for an individual to regenerate any substantial part of its body, including autotomy. Autotomy is defined as "The voluntary severance by an animal of a part of its body (commonly one of its own limbs), usually to escape capture	Categorical/ Factorial

		by a predator that has seized that part. The part then regrows."	
	05 Sexual Dimorphism	An indicator of whether sexual dimorphism is exhibited in the species. Sexual dimorphism is defined as "the occurrence of morphological differences (other than primary sexual characters) that distinguish males from females of a species of organism." (Oxford Dictionaries of Ecology and Zoology)	Categorical/ Factorial
	06 Mating System	System of mating; the organisation of sexual interactions of individuals within populations based on sex.	Categorical/ Factorial
	07 Hermaphrodism	Indicator of whether a species exhibits hermaphrodism or monoeicieosity. Hermaphrodism is defined as: "An individual that possesses both male and female sex organs; i.e. it is bisexual." (Oxford Dictionary of Zoology). Monoeciousness is defined as: "Applied to an organism in which separate male and female organs occur on the same individual (e.g. to a plant which bears male and female reproductive structures in the same flower or separate male and female flowers on the same plant, or to a hermaphrodite animal). Some authors restrict the term botanically to plants with separate male and female flowers; plants which bear male and female reproductive organs in the same flower are then called hermaphrodite."	Categorical/ Factorial
	08 Sequential Hermaphrodism	Indicator of whether a species is protogynous or protandrous.	Categorical/ Factorial
	09 Dispersal Capacity	An indicator for whether or not a species exhibits dispersal behaviour or at any stage in its life cycle. Where dispersing, a categorical description of whether dispersal is natal or breeding or otherwise. Dispersal is defined as "The tendency of an organism to move away, either from its birth site (natal dispersal) or breeding site (breeding dispersal): the opposite of philopatry." (Oxford Dictionary of Zoology).	Categorical/ Factorial
	10 Type of Dispersal	An indication of whether dispersal is a passive (requires assistance) or active (no assistance) event. See DispClasses for more information	Categorical/ Factorial
	11 Mode of Dispersal	An indicator of the mode of dispersal of the species (plant and animal specific terminology). (*e.g.,* phoretic dispersal in *Daphnia magna*).	Categorical/ Factorial
	12 Dispersal Classes	Age- or stage-classes of the species that are capable of dispersal.	Categorical/ Factorial
	13 Volancy	An indicator of whether a species is volant or non-volant (i.e., able to fly or not).	Categorical/ Factorial
	14 Aquatic Habitat Dependency	An indicator of whether a species is dependent on water or not. (e.g., limnic, lentic, lotic)	Categorical/ Factorial

Figures

Figure 1. Structure of the MOSAIC database (v1.0.0). The MOSAIC database comprises a combination of existing trait records centralised from data servers, databases, and datasets and new records collected from the primary literature by the MOSAIC team. Existing records are labelled, MOSAIC-A, in the mosaic metadata attribute field and include provenance in the metadata. New data records that adhere to existing data standards and that fit into the remit of an existing database (*i.e.*, a data gap filled by the MOSAIC team) are labelled MOSAIC-B. New records that do not have an existing data standard or for which there is not an existing database are labelled MOSAIC-C.

Figure 2. Trait fields in the MOSAIC database (v1.0.0). The MOSAIC database contains a combination of continuous and discrete trait fields. Six of the 14 trait fields are illustrated here for animals, and organised by trait level (all discrete fields) and taxonomic classification. Trait distributions vary strongly by trait and taxonomy. Observed variation in trait values by taxonomic group is a prerequisite condition to their potential value toward explaining reported variation in vital rates and resulting demographic properties across the Tree of Life. Species included in the MOSAIC database are limited to those for which stage-structured population data exist in the COMADRE, COMPADRE, and PADRINO databases.

Figure 3. The spatial distribution of MOSAIC (v1.0.0) records relative to the (a) COMADRE and (b) COMPADRE databases. The COMADRE and COMPADRE dataset include animal and plant demographic data, respectively, on all continents except Antarctica, with a substantial bias to North America and Central Europe. MOSAIC introduces trait records that include models in all major geographies of COMADRE and COMPADRE. The map shows the density of matrix population models globally (color graded by density per area in *ca.* 150 km^2 hexagonal cells). Cells bordered by white represent localities of population models for which there are trait records included in the first release of the MOSAIC trait database. Species prioritised in v1.0.0 of MOSAIC did not include the PADRINO dataset, so no distribution figure is shown for integral projection models.

Figure 4. Phylogenetic mapping of traits included in MOSAIC (v1.0.0). (a) Phylogenetic distribution of seven major traits for animals: (1) growth determination; (2) sexual dimorphism; (3) monogamy; (4) gonochory; (5) volancy; (6) dispersal; and (7) biomass. (b) Phylogenetic distribution of four traits in plants: (1) dispersal capability; (2) dispersal type; (3) dispersal mode; and (4) dispersal class. Figures developed in R using the ggtree package.

Figure 5. Trait covariance in the MOSAIC database (v1.0.0). Some traits exhibit high correlation in the MOSAIC database, such as between growth determination and between growth regeneration. Trait associations are expected to occur in the MOSAIC dataset and may reflect widespread constraints or statistical anomalies, particularly when dealing with small samples or specific taxonomic subgroups. Trait covariation can be symptomatic of biomechanical constraints (*e.g.*, flight and biomass), major growth characteristics (*e.g.*, modularity and growth determination), or other past or presently compelled associations (*e.g.*, height and vessel density).

805   **Figure 1**

807   **Figure 2**

809   **Figure 3**

(a) COMADRE

(b) COMPADRE

811 **Figure 4**

813 **Figure 5**

816   MOSAIC: A Unified Trait Database to Complement Structured Population Models

818   Connor Bernard, Gabriel Silva Santos, Jacques Deere, Roberto Rodriguez-Caro, Pol Capdevila,

819   Erik Kusch, Samuel J L Gascoigne, John Jackson, & Roberto Salguero-Gómez

821   **SUPPLEMENTARY ONLINE MATERIALS**

822   Table of Contents

830   S1: MOSAIC User Guide

<p style="text-align:center">USER GUIDE TO THE MOSAIC LIFE HISTORY DATABASE</p>

<p style="text-align:center">A Companion to COM(P)ADRE and PADRINO/A demographic databases</p>

<p style="text-align:center">*Working document – 06 February 2022*</p>

842   Table of Contents

843   User Guide Information

844   User Guide

845   Appendix

846   References

847   Table of Contents

897   User guide version information
898
899   Version 1.0.0
900
901   Release date: 6 February 2021
902
903   Contact: mosaicdatabase@biology.ox.ac.uk

General Instructions

Database Organization
The data associated with MOSAIC are provided in a single R data (extension.Rdata) and as a comma separated value (.csv) file format. The code for downloading the data can be located from the mosaic portal (https://mosaicdatabase.web.ox.ac.uk/download-database) In addition, these files are accompanied by R scripts and a nexus phylogeny available in the Supplementary Information of the manuscript introducing MOSAIC, and in our GitHub repository (https://github.com/mosaicdatabase/mosaicdatabase)

MOSAIC_v_1_0_0.RData: Contains basic information regarding the source of publication, as well as ecological, biogeographic, and taxonomic details of the demographic study for each study species, the demographic information (i.e., the matrix population model) and metadata.

Database Design
In developing the MOSAIC database, we balance level of detail with accessibility. A highly detailed, comprehensive profile of life history traits for species in COM(P)ADRE and PADRINO/A, if possible to collect, would be difficult to navigate. So rather than collating a wealth of information in many different formats, we designed this dataset to highlight a smaller collection of traits in a single format which is of most interest, and expressed, by COM(P)ADRE and PADRINO/A users. In MOSAIC's future updates, we plan to add additional detail and additional fields, but we initially took a limited approach and plan to keep the design minimal. If there are life history traits, alternative formats for existing variables, or other features you would like to see added to MOSAIC, please suggest them to us at:
mosaicdatabase@biology.ox.ac.uk

Understanding the diverse needs of users, we include in this guidance document additional direction on obtaining information on the variables we report in more detail. In this guide, we also highlight the scope of use, and caution against the most foreseeable abuses of data. We ask that all users approach this dataset with a caution and pay close attention to what variables and their statistical expression reflect.

The Meanings of NA and NDY in MOSAIC
NA in the MOSAIC data generally means that the data are not applicable. An example of where the data are not applicable is volancy within plants, as this trait does not occur in plants. NDY in the MOSAIC data means that the data have not yet been digitized. NF in the MOSAIC data means that the data are not available to date as no affirmative records were found upon review.

Disclaimer
The MOSAIC digitization team does its best to ensure data accuracy, and every piece of information goes through multiple error-checks prior to its release in www. mosaicdatabase.web.ox.ac.uk. However, we claim no responsibility for any damage that may arise from using MOSAIC. A list of error checks and potential issues in the use and interpretation of the database are described in the main manuscript. The end user is ultimately responsible for his/her interpretations of the data.

What is new in this version?

Version 1.0.0
- The first version of the database. No updates.

---

968 **Variables in MOSAIC**
969
970 The MOSAIC database is constructed of objects containing life history information across variables and
971 organized into themes to aid in navigation. The metadata object is the object containing information about
972 every study for which data is stored in the MOSAIC database. Every variable containing a value in the
973 MOSAIC database will have corresponding metadata.
974
975 Associated with every data record is a value and its corresponding metadata. The metadata details data
976 providence and relationships to existing databases. The fields within metadata are detailed below:
977
978
979 **Format Guide**
980
981
982
983 **[Index]**   *Variable Name*
984
985 **Definition**: [Definition of the variable]
986
987 **Possible values, [**cat. = categorical/discrete; cont. = continuous], [r variable class:
988   character, numeric, integer, complex, or logical.]
989 ● <XX>   [variable with two digits]
990   XXX – XXX   [discrete variable name] – [discrete variable definition]
991   <...>XXX<...>   [discrete variable with additional content on either side]
992 ● Units: [unit of measure (integer, percent, ratio, mm, km^2, g, etc.)]
993 ● Precision: [for continuous variables: scientific notation of the precision of the
994   measure - e.g., $1e^1$ for km = precise to the tens of km (3270 km), no
995   decimal; $1e^{-1}$ for temperature = precise to the tenth of a degree (10.4 °C)]
996 ● Error boundaries: [for continuous variables: boundaries beyond which values are errors]
997
998 **Usage Notes**: [Notes on boundaries of use – note that this is non-exhaustive and highlights major
999 potential errors]
1000
1001 **Additional Information:** [Guidance on databases containing more detailed information]
1002
1003 **Source Data: [**Datasets from which information was gathered; note that this will not include individual
1004 papers unless the papers are associated with a larger dataset/database. This category reflects databases.]
1005
1006 **Last updated**: [Date that information was retrieved from other databases or searches]
1007
1008

1009   **A1**   *Species Accepted*
1010
1011
1012   **Definition:** Currently accepted latin name.
1013
1014   **Possible values**, cat., factor
1015   ●   <Genus_species> - e.g., Taraxacum_officinale
1016
1017   **Usage Notes**: NA
1018
1019   **Additional Information:** NA
1020
1021   **Source Data:** This information is obtained from The Encyclopaedia of Life
1022
1023   **Last updated**: 24 January 2022
1024

1025
1026 **A2** *Kingdom*
1027
1028 **Definition:** Kingdom to which species belongs
1029
1030 **Possible values**, cat., factor
1031 ● <kingdom> - e.g., Plantae, Fungi, Rhodophyta, Chromista (yes, MOSAIC includes fungi and
1032 algae as well as plants)
1033
1034 **Usage Notes**: NA
1035
1036 **Additional Information:** NA
1037
1038 **Source Data:** The Encyclopaedia of Life
1039
1040 **Last updated**: 24 January 2022
1041

1042   **B1**   *Authors*
1043
1044   **Definition:** Surname (family name) of all authors
1045
1046   **Possible values**, NA, character
1047   ●   \<name(s)> - Separated with ";" e.g., "Smith; Jones"
1048
1049   **Usage Notes**: NA
1050
1051   **Additional Information:** NA
1052
1053   **Source Data:** NA
1054
1055   **Last updated**: 24 January 2022
1056

1057   **B2**   *Journal Name*

1058

1059   **Definition:** The document from which data were sourced.

1060

Possible values, cat., factor

● <abbreviated journal name> - Where the data come from a scientific journal article, the

abbreviated journal name is given. We use the standard abbreviation of the journal compliant

with the ISO-4 standard.

● Book - Records are from a book, or book chapter

● PhD thesis - Records are from a doctoral thesis

● MSc thesis - Records are from a masters thesis

● Report - Records are from a report

● Conference talk - Records are from a conference talk

● Conference poster - Records are from a conference poster

1071

1072   **Usage Notes**: NA

1073

1074   **Additional Information:** NA

1075

1076   **Source Data:** NA

1077

1078   **Last updated**: 24 January 2022

1079

1080   **B3**   *Year Publication*
1081
1082   **Definition:** Year of publication
1083
1084   **Possible values**, cont., numerical
1085   ● <yyyy> - e.g., 2012
1086
1087   **Usage Notes**: NA
1088
1089   **Additional Information:** NA
1090
1091   **Source Data:** NA
1092
1093   **Last updated**: 24 January 2022
1094

1095   **B4**   *DOI/ISBN Number*
1096
1097   **Definition:** Digital Object Identifier number
1098
1099   **Possible values**, NA, character
1100   ● <XXXXXXXXXXXXXX> - e.g., doi.org/10.1073/pnas.1506215112
1101
1102   **Usage Notes**: NA
1103
1104   **Additional Information:** NA
1105
1106   **Source Data:** NA
1107
1108   **Last updated**: 24 January 2022
1109

1110   **PRIMARY Variables**

1111

1112   **A1**   *Biomass*
1113
1114
1115   **Definition:** Maximum reported mass of adult individual/whole-organism. For plants, only aboveground
1116   dry mass is measured. See "Additional information" for additional information on belowground biomass.
1117
Possible values, cont., numerical
●   Units: Grams (g)
●   Precision: 0.000
●   Error Boundaries: 0-150,000,000g
●   NF – Body mass reviewed and inconclusive (no affirmative records found upon review)
●   NDY – Body mass not digitised yet (yet to be evaluated)
●   NA – not applicable
1125
1126   **Usage notes:** When both male and female data were reported, only the maximum value was considered
1127   independently of the gender of the individuals.
1128
1129   **Additional Information:** Belowground biomass is not reported because information availability is
1130   appreciably more limited than for aboveground biomass. The BIEN database includes information on
1131   belowground biomass that can be referenced and utilized where of interest.
1132
1133   **Source Data:** Amniote, TRY database.
1134
1135   **Last updated**: 24 January 2022
1136

1137 **A2** *Height*
1138

1139

1140 **Call:** -Morph$Height
1141

1142 **Definition:** (Plants) Maximum height of the whole organism/whole individual from surface (i.e.
1143 substrate) to tallest vertical extremity.
1144

1145 **Possible values**, cont., numerical
1146 ● Units: centimeters (cm)
1147 ● Precision: 0.000
1148 ● Error Boundaries: 0-1000cm
1149

1150 **Usage notes:**
1151 ● Depending on the species group, height can have profoundly different meanings. Height
1152 corresponds with embolic risk in some woody plants and corresponds with size-based fitness in
1153 others.
1154

1155 **Additional Information:** NA
1156

1157 **Source Data:** TRY database.
1158

1159 **Last updated**: 24 January 2022
1160

1161

1162 **A3**   *Growth Determination*
1163
1164 **Call:**   -Morph$Growth$Determination
1165
1166 **Definition:** Growth indeterminacy is defined by continuous growth of individuals throughout their
1167 lifetimes (measured by mass, length, bone ossification, or other indicators). This field reflects a binary
1168 classification of whether an individual is growth (in)determinate.
1169
Possible values, cat, factor
● Growth indeterminate – growth continues throughout an individual's lifespan
● Growth determinate – growth ceases or attenuates to negligibility before the end of an
individual's lifespan
● NF - Growth determination reviewed and inconclusive (no affirmative records found upon
review)
● NDY - Growth determination not digitised yet (yet to be evaluated)
● NA – Growth determination not applicable to the subject area
1178
1179 **Usage notes:** Additional classification systems for characterizing growth indeterminacy exist in the
1180 literature. Most well recognised is a six-type scheme describing growth and determination by Sebens
1181 (1987), which offers detailed characterization of growth patterns. Broad-scale information about growth
1182 and age for most species could not be located in the literature, and therefore a simplified schema is used.
1183 More resolved classifications might be incorporated into future versions of MOSAIC.
1184
1185 **Additional Information:** NA
1186
1187 **Source Data:** 24 January 2022
1188
1189 **Last updated**: NA
1190

1191 **A4**   *Regeneration*
1192
1193
1194 **Call:**   -Morphology$Growth$Regeneration
1195
1196 **Definition:** Capacity for an individual to regenerate any substantial part of its body, including autotomy.
1197 Autotomy is defined as "The voluntary severance by an animal of a part of its body (commonly one of its
1198 own limbs), usually to escape capture by a predator that has seized that part. The part then regrows."
1199
Possible values, cat., factor
● Regenerative – individuals are capable of regenerating
● Non-regenerative – individuals do not exhibit the capacity to regenerate tissues
● NF - regenerative abilities reviewed and inconclusive (no affirmative records found upon review)
● NDY - regeneration information not digitised yet (yet to be evaluated)
● NA – regenerative abilities not applicable to the subject area
1206
1207 **Usage notes:** There are a number of more resolved schemes for detailing whether regenerating different
1208 parts of the body. For particular questions pertaining to the nature of injury, the level of recovery, the role
1209 of depredation, and the consequences for reproduction, more detail may be appropriate. This dataset
1210 covers the most general applications and initially screens for regenerative capability (from poor recovery
1211 of appendages to complete regeneration of limbs).
1212
1213 **Additional Information:** NA
1214
1215 **Source Data:** NA
1216
1217 **Last updated**: 24 January 2022
1218

A5   *Sexual Dimorphism*

Call: -Morph$Dimorph

Definition: An indicator of whether sexual dimorphism is exhibited in the species. Sexual dimorphism is defined as "the occurrence of morphological differences (other than primary sexual characters) that distinguish males from females of a species of organism." (Oxford Dictionaries of Ecology and Zoology)

Possible values, cat., factor
- Sexually Dimorphic – species is sexually dimorphic
- Sexually Monomorphic – species is sexually monomorphic (i.e. non-dimorphic)
- NF - Sexual dimorphism reviewed and inconclusive (no affirmative records found upon review)
- NDY - Sexual dimorphism not digitised yet (yet to be evaluated)
- NA – Sexual dimorphism not applicable to the subject area

Usage Notes: NA

Additional Information:

Source Data: NA

Last updated: 24 January 2022

1244 **B1** *Mating system*
1245
1246
1247 **Call:** -Reproduction$MatingSystem
1248
1249 **Definition:** System of mating; the organization of sexual interactions of individuals within populations
1250 based on sex.
1251
Possible values, cat., factor
● Monogamy – exclusive mating between one male and one female
● Non-monogamy - Non-monogamy was assigned based on genetic or behavioural evidence
● NF - Mating system reviewed and inconclusive (no affirmative records found upon review)
● NDY - Mating system not digitised yet (yet to be evaluated)
● NA – Mating system not applicable to the subject area
1258
1259 **Usage notes:**
1260 ● Metric does not identify size of groups for plural mating systems
1261 ● Metric does not identify enforcement mechanisms for different mating systems
1262
1263 **Additional Information:** NA
1264
1265 **Source Data:** NA
1266
1267 **Last updated**: 24 January 2022
1268
1269
1270
1271

B2   *Sexual allocation*

Call: **-**Reproduction$Allocation

Definition: Indicator of whether a species exhibits hermaphroditism or monoeicieosity. Hermaphrodism is defined as: "An individual that possesses both male and female sex organs; i.e. it is bisexual." (Oxford Dictionary of Zoology). Monoeciousness is defined as: "Applied to an organism in which separate male and female organs occur on the same individual (e.g. to a plant which bears male and female reproductive structures in the same flower or separate male and female flowers on the same plant, or to a hermaphrodite animal). Some authors restrict the term botanically to plants with separate male and female flowers; plants which bear male and female reproductive organs in the same flower are then called hermaphrodite."

Possible values, cat., factor
- Hermaphroditic – species is hermaphroditic
- Monoecious – species is monoecious
- Dioecious or Gonochorous – species is dioecious (Gonochorous was adopted for animals)
- NF - Hermaphroditism reviewed and inconclusive (no affirmative records found upon review)
- NDY - Hermaphroditism not digitised yet (yet to be evaluated)
- NA – Hermaphroditism is not applicable to the subject area

Usage notes:
-

Additional Information: NA

Source Data: NA

Last updated: 24 January 2022

1306 **B3**   *Sequesntial hermaphroditism*
1307
1308
1309 **Call:**   Reproduction$SeqHermaph
1310
1311   **Definition:** Indicator of whether there is a sex switch during the organism's lifespan.
1312
Possible values, cat., factor
● Protogynous – species is protogynous: organisms that are female and at some point in their
lifespan change sex to male.
● Protandrous – species is protandrous: organisms that are male and at some point in their lifespan
change sex to male.
● NF - Protogyny/Protandry reviewed and inconclusive (no affirmative records found upon review)
● NDY - Protogyny/Protandry not digitised yet (yet to be evaluated)
● NA – Protogyny/Protandry is not applicable to the subject area.
1321
1322 **Usage notes:** NA
1323
1324 **Additional Information:** Note that is not rare flowers present protogyny/protandry but this was not
1325 considered in Mosaic so far
1326
1327 **Source Data:** NA
1328
1329 **Last updated**: 24 January 2022
1330

1331 **C1** *Dispersal Capability*
1332
1333
1334 **Call:** -Movement$Dispersal
1335
1336 **Definition:** An indicator for whether or not a species exhibits dispersal behaviour or at any stage in its life
1337 cycle. Where dispersing, a categorical description of whether dispersal is natal or breeding or otherwise.
1338 Dispersal is defined as "The tendency of an organism to move away, either from its birth site (natal
1339 dispersal) or breeding site (breeding dispersal): the opposite of philopatry." (Oxford Dictionary of
1340 Zoology).
1341
Possible values: cat., factor
● Dispersing – Exhibits at least one age-/stage-class which disperses; natal or breeding components
unknown.
● Natal Dispersal – Permanent dispersal of at least one age-/stage-class
● Breeding Dispersal – Dispersal of adults between breeding attempts in at least one age-/stage-
class
● Multi-Dispersal – Both natal and breeding dispersal reported in the species; see DispClasses for
more information
● Non-Dispersing – Species observed to have no dispersal traits/behaviour
● NF - dispersal capability reviewed and inconclusive (no affirmative records found upon review)
● NDY – dispersal capability unknown/not yet evaluated
● NA – not applicable
1354
1355 **Usage Notes:** NA
1356
1357 **Additional Information:** NA
1358
1359 **Source Data:** NA
1360
1361 **Last updated**: 24 January 2022
1362

1363 **C2**   *Type of Dispersal*
1364
1365
1366 **Call:**   -Movement$TypeDisp
1367
1368 **Definition:** An indication of whether dispersal is a passive (requires assistance) or active (no assistance)
1369 event. See DispClasses for more information.
1370
Possible values: cat., factor
● Active – organism utilises its own morphology for the dispersal event
● Passive – organism is unable to disperse through their own means and require an external factor
● Active and Passive – organism is able to disperse with assistance but can also use external
factors. Active and passive dispersal can occur within the same life stage or can occur in different
life stages.
● NF – type of dispersal reviewed and inconclusive (no affirmative records found upon review)
● NDY – type of dispersal unknown/not yet evaluated
● NA – not applicable
1380
1381 **Usage Notes:** NA
1382
1383 **Additional Information:** NA
1384
1385 **Source Data:** NA
1386
1387 **Last updated**: 24 January 2022
1388
1389

1390 **C3** *Mode of Dispersal*
1391
1392
1393 **Call:**   -Movement$ModeDisp
1394
1395 **Definition:** An indicator of the mode of dispersal of the species (plant and animal specific terminology).
1396
Possible values: cat., factor
● Motile – the dispersal of animal species without assistance
● Phoretic – the dispersal of animal species by attaching to another animal
● Water currents – the dispersal of animal species by water
● Motile and water currents – animal species that disperse without assistance and by water, both
forms of dispersal can occur within the same life stage or can occur in different life stages
● Anemochory – the dispersal of plant seeds by wind
● Anthropochory – the dispersal of plant seeds by humans
● Autochory – the dispersal of plant seeds without assistance from an external vector (e.g., by
gravity or ballistic dispersal)
● Hydrochory – the dispersal of plant seeds by water
● Zoochory – the dispersal of plant seeds by animals
● Anemochory and Anthropochory – plant seeds can be dispersed by wind and humans
● Anemochory and Autochory – plant seeds can be dispersed by wind and without the help of an
external vector
● Anemochory and Hydrochory – plant seeds can be dispersed by wind and water
● Anemochory and Zoochory – plant seeds can be dispersed by wind and animals
● Autochory and Hydrochory – plant seeds can be dispersed without the help of an external vector
and by water
● Autochory and Zoochory – plant seeds can be dispersed without the help of an external vector
and by animals
● Hydrochory and Zoochory – plant seeds can be dispersed by water and animals
● Autochory, Anthropochory and Zoochory – plant seeds can be dispersed without the help of an
external vector, by humans and by animals
● NF – mode of dispersal reviewed and inconclusive (no affirmative records found upon review)
● NDY – mode of dispersal unknown/not yet evaluated
● NA – not applicable
1424
1425 **Usage Notes:** Plant seed dispersal modes can be subdivided into further categories, however we collated
1426 lower order categories into the higher order categories identified here
1427
1428 **Additional Information:** NA
1429
1430 **Source Data:** NA
1431
1432 **Last updated**: 24 January 2022
1433
1434

1435 **C4**   *Dispersal Class*
1436
1437 **Call:**   -Movement$DispClass
1438
1439 **Definition:** Age- or stage-classes of the species that are capable of dispersal.
1440
Possible values: cat., factor
● Adult – dispersal stage is an individual that has reached maturity, we include sub-adults into this
category
● Egg – dispersal stage is a vessel within which an embryo develops and is expelled by an adult
allowing for dispersal
● Fertile material – dispersal stage is a part of an individual, or in some cases a complete individual,
that contains fertile material (e.g., the alga *Fucus vesiculosus*; detached floating
material/individual that contains gametes)
● Gamete – dispersal stage is the reproductive cell not within a vessel
● Juvenile – dispersal stage is an individual that has not reached maturity
● Larval – dispersal in a specific juvenile stage restricted to non-mammal species, species can have
multiple larval stages
● Seed – dispersal stage is fertilized, specific to plant species and, in our definition, references
seeds and/or fruits that are dispersed
● Sperm – dispersal stage is the male gamete
● Spore – dispersal stage is a single cell that only contains half of the chromosome of the adult, can
produce eggs or sperm
● Sporophyte – dispersal stage is a nonsexual phase of a species producing two diploid spores
● Zoospore – dispersal stage is a motile asexual spore
● Zygote – dispersal stage is a fused male and female gamete
● Adult and Juvenile – dispersal stage can be both the adult and juvenile stage
● Egg and Larval – dispersal stage can be both the egg and larval stage
● Gamete and Spore – dispersal stage can be both the gamete and spore
● Zoospore, Sperm and Sporophyte – dispersal stage can be a zoospore, sperm or sporophyte
● NF – dispersal class reviewed and inconclusive (no affirmative records found upon review)
● NDY – dispersal class unknown/not yet evaluated
● NA – not applicable
1468
1469 **Usage Notes:** Dispersal can occur in more than one age- or stage- class, where this occurs it is noted as
1470 such within the database
1471
1472 **Additional Information:** NA
1473
1474 **Source Data:** NA
1475
1476 **Last updated**: 24 January 2022
1477

1478   **C5**   *Volancy*
1479
1480
1481   **Call:**   -Movement$Volancy
1482
1483   **Definition:** An indicator of whether a species is volant or non-volant (i.e., able to fly or not).
1484
1485   **Possible values**, cat., factor
1486   ●   Volant – the species is volant
1487   (Most Birds (Class Aves), all Bats (Order Chiroptera), and some invertebrate species)
1488   ●   Non-volant – the species is non-volant
1489   ●   Semi-volant – the species has gliding abilities
1490   (e.g., Gliding lizards (*Draco* spp.); flying squirrels such as the Northern flying squirrel
1491   (*Glaucomys sabrinus*); flying fish (Exocoetidae); gliding frogs such as Wallace's flying frog
1492   (*Rhacophorus nigropalmatus*); and gliding ants such as *Cephalotes atratus*).
1493   ●   NF – volancy reviewed and inconclusive (no affirmative records found upon review)
1494   ●   NDY – volancy unknown/not yet evaluated
1495   ●   NA – not applicable
1496
1497   **Usage Notes:** NA
1498
1499   **Additional Information:** NA
1500
1501   **Source Data:** NA
1502
1503   **Last updated**: 24 January 2022

1504　S2: MOSAIC Trait Primary Search Key Words

1505　　　　　　　　MOSAIC: Keywords used in primary literature search

1506
1507　Taxonomic names (binomial nomenclature) were used in connection with the field-specific terms in the
1508　search of primary literature in identifying the records of interest. The below include names that were
1509　searched in review of the primary literature.
1510
1511
1512
1513　Biomass—
1514　　　　NA (database only)
1515
1516　Height—
1517　　　　NA (database only)
1518
1519　Growth determination—
1520　　　　Sources: *Web of Science; Google Scholar; Scopus*
1521　　　　Terms: Growth determinat*, growth determination, growth determinate, growth indetermination,
1522　　　　growth
1523
1524　Regeneration—
1525　　　　Sources: *Web of Science; Google Scholar; Scopus*
1526　　　　Terms: Regenerat*, regeneration, regenerate, rejuvenation
1527
1528　Sexual dimorphism—
1529　　　　Sources: *Web of Science; Google Scholar; Scopus*
1530　　　　Terms: Dimorphic
1531
1532　Mating system—
1533　　　　Sources: *Web of Science; Google Scholar; Scopus*
1534　　　　Terms: Monogamy, monogamous, polygyny, polygynous, polyandry, polyandrous, mating
1535　　　　system
1536
1537　Hermaphrodism—
1538　Mating system—
1539　　　　Sources: *Web of Science; Google Scholar; Scopus*
1540　　　　Terms: Hermaphrod*, hermaphroditic, hermaphrodism, hermaphrodite, gonochoric,
1541　　　　gynochorous, monoecious, dioecious, sexual differentiation,
1542
1543　Sequential hermaphrodism—
1544　　　　Sources: *Web of Science; Google Scholar; Scopus*
1545　　　　Terms: Sequential hermaphrodism, protandry, protogyny, protogynous hermaphroditism,
1546　　　　protandrous hermaphroditism
1547
1548　Dispersal capacity—
1549　　　　Sources: *Web of Science; Google Scholar; Scopus*
1550　　　　Terms: Dispers*, dispersal, dispersing, dispersal capacity, dispersal capability
1551
1552　Type of dispersal—

1553   Sources: *Web of Science; Google Scholar; Scopus*
1554   Terms: Dispers*, active, passive
1555
1556 Mode of dispersal—
1557   Sources: *Web of Science; Google Scholar; Scopus*
1558   Terms: Dispers*, motile, phoretic, currents, anemochory, anthropochory, autochory, hydrochory,
1559   zoochory, anthropochory
1560
1561 Dispersal classes—
1562   Sources: *Web of Science; Google Scholar; Scopus*
1563   Terms: Dispers*, adult, jouvenile, egg, fertile, seed, sperm, spore, sporophyte, zoospore, zygote,
1564   gamete
1565
1566 Volancy—
1567   Sources: *Web of Science; Google Scholar; Scopus*
1568   Terms: Volant, non-volant, flight, flying, flightless, non-flying, winged, glide, gliding, ground
1569
1570 Aquatic habitat dependency—
1571   Sources: *Web of Science; Google Scholar; Scopus*
1572   Terms: Anadromous, catandromous, estuarine, brackish, lotic, lentic, lemnitic, littoral, palagic,
1573   marine, freshwater, saltwater, sea, ocean.
1574
1575

S3. Databases Searched

MOSAIC: Databases Reviewed to Date for the MOSAIC Database

Type. Database Name. DOIs.

1. **Plants**. Botanical Information and Ecology Network (BIEN) (doi: 10.7287/peerj.preprints.2615v2)
2. **Plants**. TRY: Global Plant Trait Database (TRY) (doi: 10.1111/j.1365-2486.2011.02451.x)
3. **Phylogeny.** Open Tree of Life: A synthesis of phylogeny and taxonomy into a comprehensive tree of life (doi: 10.1073/pnas.1423041112)
4. **Vertebrates**. Amniote: An amniote life history database to perform comparative analysis with birds, mammals, and reptiles (doi: 10.1890/15-0846R.1)
5. **Amphibians**. AmphiBIO: a global database for amphibian ecological traits. (doi:L 10.1038/sdata.2017.123)
6. **Mammals**. PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals (doi: 10.1890/08-1494.1)
7. **Animals**. AnAge Database of Animal Ageing and Longevity. (doi: 10.1111/j.1474-9726.2008.00442.x)
8. **General (Sex)**. Tree of Sex: A database of sexual systems (10.1038/sdata.2014.15)
9. **General (Variable)**. Open Trait Network Databases (https://opentraits.org/)

1599 S4: Vignettes

1600

1601   Navigating MOSAIC

1602   Vignette #1 - Navigating MOSAIC

1603

1604

Welcome to the **MOSAIC database,** a database of functional traits for
comparative demography. The database, user, guide, and additional information
can be found and http://mosaicdatabase.web.ox.ac.uk.

MOSAIC is a database that aggregates existing databases and adds new records
for functional traits that currently do not have a database established. In
this vignette, we will show you how to download the dataset, search records,
and relate MOSAIC records with COMADRE, COMPADRE, and PADRINO databases.

Optional clearance of working space.

```
1613   rm(list = ls()) # Clear your environment

1614   if(!is.null(dev.list())) dev.off() # Clear plots/graphics

1615   cat("\014") # Clear console
```

Downloading MOSAIC

MOSAIC can be downloaded as an S4 object by running the below code in R. S4
data objects in R are an object oriented system in the R language that allow
control of constituent data fields Similar to S3 objects (which use the "$"
operator). S4 are comprised of objects that can be searched with the "@"
operator or slots, discussed in more detail below.

```
1622   library(devtools) # Compulsory package to pull down packages from GitHub. Ins
1623   tall if necessary.

1624   install_github("mosaicdatabase/Rmosaic")

1625   source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
1626   ain/mosaic_fetch.R") # Link to GitHub repo

1627   mosaic <- mos_fetch("v1.0.0") # Download version 1.0.0 (active version Feb 20
1628   22)

1629   library(Rmosaic)
```

Basics of manually navigating MOSAIC

Mosaic traits can be searched using the "@" operator. Attribute names
searched this way are analogous to the columns of a dataframe in a relational
database structure.

Once downloaded, you should be able to type statements
mosaicdatabase@[insertfield] (where [insertfield] is a particular trait). If
you are working in Rstudio, after the "@" a drop-down of the slots (traits)
should autopopulate.

```
1638  # Three examples of querying traits (easiest for navigation)

1639  mosaic@biomass

1640  mosaic@height

1641  mosaic@volancy
```

1642 The species corresponding with each index can be queried by prompting:

```
1643  # Three examples of querying traits (easiest for navigation)

1644  mosaic@species
```

1645 Data in mosaic can also be access using slots. Slots are the recommended mode
1646 of searching the database - though it has the disadvantage of not enabling
1647 the autopopulation of the attributes contained in the database (traits must
1648 be spelled out manually).

```
1649  # You can also search these by slot (recommended)

1650  slot(mosaic, "biomass")

1651  slot(mosaic, "height")

1652  slot(mosaic, "volancy")
```

Within each trait object, there are eight fields in mosaic. The first of
field is called "values" and contains the data. Values are unitless values
(either numeric or factorial) that are reported in units described in
the **Mosaic User Guide** http://mosaicdatabase.web.ox.ac.uk/user-guide. The
metadata is organised into additional attributes, reflecting the individual
elements of the metadata for a given record, including the authors, journal,
year of publication, databases from which data are sourced (if applicable)

mosaic@metaTaxa maps the complete taxanomic classification structure of a
species - from Kingdom to species for taxonomic clustering.

The other six attributes - "author", "year", "journal,"doi","database",
and"mosaic" - are metadata corresponding with each value record. For
instance:

```
1665  mosaic@species[[2]] # For this species, let us look at volancy (flight capaci
1666  ty) value

1667  mosaic@volancy@value[[2]] # or, equivalently:

1668  slot(slot(mosaic, "volancy"), "value")[[2]]
```

1669 Corresponds with the following metadata

```
1670  mosaic@volancy@author[[2]] # for "Acinonyx jubatus"
```

```
slot(slot(mosaic, "volancy"), "author")[[2]] # Author of the source publicati
on

slot(slot(mosaic, "volancy"), "year")[[2]] # And year of the source publicati
on

slot(slot(mosaic, "volancy"), "journal")[[2]] # The journal of the source pub
lication

#etc.
```

Using MOSAIC functions to quickly access files

A series of convenience functions can be sourced from the MOSAIC GitHub page
to facilitate navigating and working with the mosaic database that can be
accessed by running the follwing script.

```
source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
ain/navMosaic_46.R")
```

Below we highlight some of the basic queries for which the mosaic functions
can assist.

Is a species included in Mosaic?

```
spp_check("Fritillaria biflora")

## [1] TRUE

spp_check("Pagophilus groenlandicus")

## [1] FALSE
```

Can I see all records for a given trait?

```
traitAllSpp("biomass") # Only the first five records are shown for space

## [1] "NDY"    "50578" "ND"     "52500" "351000" "62000"
```

Can I see an overview of all records for a given species?

```
singSppTraitSummary("Aepyceros melampus")

##   biomass height   growthdet regen  dimorph        matsyst      hermaph se
qherm

## 1   52500    NDY Determinate   NDY Dimorphic Non-monogamous Gonochorous
NDY

##          dispcap disptype modedisp dispclass   volancy

## 1 Natal Dispersal   Active   Motile    Adult Non-volant
```

```
##                                              aquadep
## 1 Terrestrial, Water Habitat Independent
```

Can I see all records for more than one species?

```
sppAllTrait(c("Acinonyx jubatus", # you can also pass lists or dataframes to
this command
                "Acropora downingi",
                "Aepyceros melampus",
                "Alces alces",
                "Alligator mississippiensis"))
##                    sppnames biomass height   growthdet       regen   dim
orph
## 1          Acinonyx jubatus   50578    NDY Determinate         NDY Dimor
phic
## 2         Acropora downingi      ND    NDY         NDY Regenerative
NDY
## 3        Aepyceros melampus   52500    NDY Determinate         NDY Dimor
phic
## 4               Alces alces  351000    NDY Determinate         NDY Dimor
phic
## 5 Alligator mississippiensis   62000    NDY Determinate Regenerative
NDY
##         matsyst        hermaph seqherm         dispcap disptype       mod
edisp
## 1 Non-monogamous    Gonochorous     NDY Natal Dispersal   Active        M
otile
## 2            NDY Hermaphroditic     NDY Natal Dispersal  Passive Water cur
rents
## 3 Non-monogamous    Gonochorous     NDY Natal Dispersal   Active        M
otile
## 4 Non-monogamous    Gonochorous     NDY Natal Dispersal   Active        M
otile
## 5 Non-monogamous    Gonochorous     NDY Natal Dispersal   Active        M
otile
##          dispclass   volancy                                        aqua
dep
## 1        Juvenile Non-volant       Terrestrial, Water Habitat Independ
ent
## 2        Juvenile Non-volant                                        Mar
ine
```

```
## 3               Adult Non-volant       Terrestrial, Water Habitat Independ
ent
## 4          Juvenile Non-volant Terrestrial, Facultative Freshwater Depend
ent
## 5 Adult and Juvenile Non-volant  Terrestrial, Obligative Freshwater Depend
ent
```

Can I get a breakdown of counts/frequency of trait values?

```
traitFrequency("volancy")
##            counts  freq
## NDY          1329    NA
## Non-volant     82 0.882
## Semi-volant     1 0.011
## Volant         10 0.108
traitFrequency("growthdet")
##              counts  freq
## Determinate      31 0.596
## Indeterminate    21 0.404
## NDY            1370    NA
traitFrequency("hermaph")
##                          counts  freq
## Dioecious                    24 0.108
## Gonochorous                 106 0.475
## Hermaphroditic               72 0.323
## Hermaphroditic & Gonochorous  1 0.004
## Monoecious                   20 0.090
## NDY                        1199    NA
```

Can I get all metadata for one or more traits?

```
metadata("volancy", 14)
##           author year                 journal database mosaic
## 1 Campos, Z. et al. 2006 The Herpetological Journal     NDY    NDY
multiMetaRecords("volancy", c(14:20))
##                               author year
```

```
## 1                         Ekerna, L. S. & Cords, M. 2007
## 2                           Zimmerman, S. J. et al. 2019
## 3 Jack, K. M., Sheller, C. & Fedigan, L. M. 2012
## 4                                           DFO 2013
## 5                         Torres, R. T. et al. 2017
## 6                                          NDY  NDY
## 7                         Campos, Z. et al. 2006
##
journal
## 1                                                  A
nimal Behaviour
## 2
The Condor
## 3                                    American Journal
of Primatology
## 4 Canadian Science Advisory Secretariat Central and Arctic Region Science
Advisory Report
## 5
Oryx
## 6
NDY
## 7                                          The Herpeto
logical Journal
##    database mosaic
## 1        NDY     NDY
## 2        NDY     NDY
## 3        NDY     NDY
## 4        NDY     NDY
## 5        NDY     NDY
## 6        NDY     NDY
## 7        NDY     NDY
```

1807   Updated 22 February 2022

1808   Vignette #2 - Basic Regression Using MOSAIC & COMPADRE

1809
1810
MOSAIC was built with the intention of its use in comparative biodemography -
to strengthen the map of which functional traits predict vital rates and
which do not. Traits in the MOSAIC database (factorial and numeric variables)
are formatted to be easily accessed and used for regression-based analyses,
among other types of analysis. MOSAIC was also designed for quick integration
with the COMADRE, COMPADRE, and PADRINO structural population model
databases, as outlined in the model exercise below.

```
1818   library(devtools)

1819   library(tidyverse)

1820   library(Rcompadre) # package to access matrix population models

1821   library(Rage) # package to perform life-history calculations
```

1822   Accessing MOSAIC

1823   Download MOSAIC from the mosaic portal. For more information on the basics of
1824   downloading MOSAIC and navigating the data structure, see: Vignette #1:

```
1825   library(devtools) # Compulsory package to pull down packages from GitHub. Ins
1826   tall if necessary.

1827   install_github("mosaicdatabase/Rmosaic") # library of navigation-aiding funct
1828   ions

1829   source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
1830   ain/mosaic_fetch.R") # Link to GitHub repo

1831   mosaic <- mos_fetch("v1.0.0") # Download version 1.0.0 (active version Feb 20
1832   22)

1833   library(Rmosaic)
```

Regressing Biomass on Generation Time

In this exercise, the simple bivariate relationship of biomass on generation
time is explored through basic regression. Bare in mind, that the below offer
the building blocks for more complex multiple regression exercises and other
forms of analysis – such as geospatial interpolation and ordination-based
practices (see Vignette #4 for a brief vignette of PVA using mosaic).

#1: Extract biomass data for mammals

Extract the biomass values for mammals in the MOSAIC dataset.

```
1842  # Extracting all biomass data

1843

1844  mosaic_dataframe <- data.frame(mosaic@taxaMetadat,

1845                                 Biomass = as.numeric(mosaic@biomass@value),

1846                                 MOSAIC_index = 1:length(mosaic@index))

1847

1848  # Restrict mammal data with non NA biomass data

1849

1850  mosaic_mammal <- subset(mosaic_dataframe,

1851                          Class == "Mammalia" & is.na(Biomass) == F)

1852

1853  # Extracting the first COMPADRE matrix id for each entry

1854

1855  matrix_ids_full <- mosaic@index[mosaic_mammal$MOSAIC_index]

1856

1857  # Extract just the first matrix id using sapply

1858

1859  mammal_matrix_ids <- sapply(matrix_ids_full, `[[`, 1)

1860

1861  # Add matrix id to the mammal data

1862

1863  mosaic_mammal$MatrixID <- mammal_matrix_ids

1864

1865  # Add full species names

1866

1867  mosaic_mammal$Binomial <- paste0(mosaic_mammal$Genus, " ", mosaic_mammal$Spec
1868  ies)
```

1869 #2: Download COMADRE matrix population models

1870 A cdb_fetch() function is used to clone the Comadre database <www.compadre-
1871 db.org> - which is an s3 copy of a SQL-based relational database - into an S4
1872 data object locally manipulable in R.

Using ids associated with each matrix population model, we will subset the Comadre database of matrix population models for the animals. First we use the Rcompadre package to 'fetch' the Comadre database.

```
# Download the most recent database

comadre <- cdb_fetch("comadre")

# Rcompadre function flags potential issues with matrices, including NAs and
ergodicity

comadre_flag <- cdb_flag(comadre)

# Remove matrices with NAs and those that are non-ergodic (which throw errors
in generation time)

comadre_correct <- subset(comadre_flag,
                          check_NA_A == FALSE &
                          check_ergodic == TRUE)
```

The subsets of mosaic and Compadre can be overlapped in one line of code.

```
# Subset to mammal matrices

comadre_biomass <- subset(comadre_correct,
                          MatrixID %in% mammal_matrix_ids)
```

3. Calculating generation time

Rcompadre enables sub-setting of a matrix population model (life table) into constituent matrices representing the survival and fecundity/reproductive components (U and F matrices, respectively). Functions in the Rage package often require the specification of both the U and the F matrix, not the combined A matrix. The Rage package contains a suite of common demographic calculations - including GENERATION TIME - which are used to simplify this exercise. Note that other common demographic derivations, including transient indices can be extracted using the identical procedure.

```
# Calculate Generation Time

comadre_biomass$Generation_Time <- mapply(Rage::gen_time,
                                          matU(comadre_biomass),
```

```
1909                                              matF(comadre_biomass))
```

1910 4. Explore the relationship of generation time data and biomass data
1911 Matrix IDs can be used to bridge generation time from comaadre with mosaic
1912 records.

```
1913  # Extract matrix ID and generation time columns (excluding other matrix metad
1914  ata)
1915
1916  comadre_biomass <- as.data.frame(comadre_biomass)
1917
1918  comadre_biomass <- comadre_biomass[,c("MatrixID", "Generation_Time")]
1919
1920  # Join the datasets, using Matrix IDs to crosswalk the data
1921
1922  mosaic_biomass <- merge(mosaic_mammal, comadre_biomass,
1923                          by = "MatrixID", all.x = TRUE)
1924  # Remove infinity values or NA
1925
1926  mosaic_biomass <- mosaic_biomass[!(is.infinite(mosaic_biomass$Generation_Time
1927  ) |
1928                                     is.na(mosaic_biomass$Generation_Time)),]
1929
1930  # Log biomass and generation time
1931
1932  mosaic_biomass$log10_biomass <- log10(mosaic_biomass$Biomass)
1933  mosaic_biomass$log10_gent <- log10(mosaic_biomass$Generation_Time)
1934
1935
1936
1937  # The plot
1938
1939  plot(mosaic_biomass$log10_biomass, mosaic_biomass$log10_gent,
1940       pch=16, cex=1.5, col=alpha("black", 0.5),
1941       xlab=expression(paste(log[10], "Biomass")),
1942       ylab=expression(paste(log[10], "Generation time"))
```

```
)

# Simple linear regression (note that we are skipping model fitting in this b
rief vignette)

lmBioGenT = lm(log10_gent~log10_biomass, data = mosaic_biomass)

summary(lmBioGenT)
##
## Call:
## lm(formula = log10_gent ~ log10_biomass, data = mosaic_biomass)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.53325 -0.16107 -0.01597  0.19936  0.38382
##
## Coefficients:
##                Estimate Std. Error t value Pr(>|t|)
## (Intercept)    0.85966    0.25372   3.388  0.00253 **
## log10_biomass  0.02762    0.05728   0.482  0.63418
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2623 on 23 degrees of freedom
## Multiple R-squared:  0.01001,    Adjusted R-squared:  -0.03303
## F-statistic: 0.2326 on 1 and 23 DF,  p-value: 0.6342
abline(lmBioGenT, col="red", lty=2, lwd=2)
```

1971
1972

1973   S5: LogNormal Distribution of Mass and Height

1974

1975
1976   Raw heights from MOSAIC v1.0.0
1977
1978

1979
1980   Log Heights from MOSAIC v.1.0.0
1981

1982
1983   Raw mass from MOSAIV v1.0.0
1984

1985
Log mass from MOSAIC v1.0.0

Figures

Figure 1. Structure of the MOSAIC database (v1.0.0). The MOSAIC database comprises a combination of

existing trait records centralised from data servers, databases, and datasets and new records collected from

the primary literature by the MOSAIC team. Existing records are labelled, MOSAIC-A, in the mosaic

metadata attribute field and include provenance in the metadata. New data records that adhere to existing

data standards and that fit into the remit of an existing database (*i.e.*, a data gap filled by the MOSAIC team)

are labelled MOSAIC-B. New records that do not have an existing data standard or for which there is not

an existing database are labelled MOSAIC-C.

Figure 2. Trait fields in the MOSAIC database (v1.0.0). The MOSAIC database contains a combination of

continuous and discrete trait fields. Six of the 14 trait fields are illustrated here for animals, and organised

by trait level (all discrete fields) and taxonomic classification. Trait distributions vary strongly by trait and

taxonomy. Observed variation in trait values by taxonomic group is a prerequisite condition to their

potential value toward explaining reported variation in vital rates and resulting demographic properties

across the Tree of Life. Species included in the MOSAIC database are limited to those for which stage-

structured population data exist in the COMADRE, COMPADRE, and PADRINO databases.

Figure 3. The spatial distribution of MOSAIC (v1.0.0) records relative to the (a) COMADRE and (b)

COMPADRE databases. The COMADRE and COMPADRE dataset include animal and plant demographic

data, respectively, on all continents except Antarctica, with a substantial bias to North America and Central

Europe. MOSAIC introduces trait records that include models in all major geographies of COMADRE and

COMPADRE. The map shows the density of matrix population models globally (color graded by density

per area in *ca.* 150 km^2 hexagonal cells). Cells bordered by white represent localities of population models

for which there are trait records included in the first release of the MOSAIC trait database. Species

prioritised in v1.0.0 of MOSAIC did not include the PADRINO dataset, so no distribution figure is shown

for integral projection models.

Figure 4. Phylogenetic mapping of traits included in MOSAIC (v1.0.0). (a) Phylogenetic distribution of seven major traits for animals: (1) growth determination; (2) sexual dimorphism; (3) monogamy; (4) gonochory; (5) volancy; (6) dispersal; and (7) biomass. (b) Phylogenetic distribution of four traits in plants: (1) dispersal capability; (2) dispersal type; (3) dispersal mode; and (4) dispersal class. Figures developed in R using the ggtree package.

Figure 5. Trait covariance in the MOSAIC database (v1.0.0). Some traits exhibit high correlation in the MOSAIC database, such as between growth determination and between growth regeneration. Trait associations are expected to occur in the MOSAIC dataset and may reflect widespread constraints or statistical anomalies, particularly when dealing with small samples or specific taxonomic subgroups. Trait covariation can be symptomatic of biomechanical constraints (*e.g.*, flight and biomass), major growth characteristics (*e.g.*, modularity and growth determination), or other past or presently compelled associations (*e.g.*, height and vessel density).

41 **Figure 1**

43   **Figure 2**

45   **Figure 3**

(a) COMADRE

(b) COMPADRE

Figure 4

(a)

VOLANCY

VOLANT
SEMI-VOLANT
NON-VOLANT
NOT DIGITISED

SEQUENTIAL
HERMAPHRODISM

PROTOGYNY

PROTANDRY

HERMAPHRODISM

GONOCHOROUS
HERMAPHRODITIC
HERMAPH + GONO
NOT FOUND

GROWTH
DETERMINACY

DETERMINATE

INDETERMINATE

SEXUAL
DIMORPHISM

DIMORPHIC

MONOMORPHIC

SEXUAL
DIMORPHISM

MONOGAMOUS

NON-
MONOGAMOUS

DISPERSAL CLASS

ADULT   EGG + LARVAL   JUVENILE   NOT FOUND

(b)

DISPERSAL
CAPACITY

DISPERSING

NOT FOUND

DISPERSAL
TYPE

NOT FOUND

PASSIVE

DISPERSAL
CLASS

GAMETE + SPORE

SEED
SPORE
ZYGOTE

DISPERSAL MODE

ANEMOCHORY
(+SEMI-ANEMOCHORY)

AUTOCHORY
(+SEMI- AUTOCHORY)

HYDROCHORY
(+SEMI- HYDROCHORY)

49   **Figure 5**

50