MOSAIC: A Unified Trait Database to Complement Structured Population Models

Connor Bernard[1,†], Gabriel Silva Santos[1,2,3], Jacques A. Deere[1,4], Roberto Rodriguez-Caro[1,5], Pol

Capdevila[1,6] Erik Kusch[7], Samuel J L Gascoigne[1], John Jackson[1], Roberto Salguero-Gómez[1,7,8]

[1] Department of Zoology, University of Oxford, 11a Mansfield Rd, OX13SZ, Oxford, United Kingdom

[2] Department of Ecology, Rio de Janeiro State University, Rio de Janeiro, Brazil

[3] Instituto Nacional da Mata Atlântica, INMA, Santa Teresa, Brazil.

[4] Department of Evolutionary and Population Biology, Institute for Biodiversity and Ecosystem Dynamics,

University of Amsterdam, 1012 WX Amsterdam, Netherlands

[5] Departamento de Biología Aplicada. Universidad Miguel Hernández. Av. Universidad, s/n, 03202 Elche

(Alicante) Spain

[6] School of Biological Sciences, University of Bristol, 24 Tyndall Ave, Bristol BS8 1TQ, United Kingdom

[7] Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Section for Ecoinformatics &

Biodiversity, Department of Biology, Aarhus University

[7] Centre for Biodiversity and Conservation Science, University of Queensland, St. Lucia, QLD, Australia

[8] Evolutionary Demography Laboratory, Max Plank Institute for Demographic Research, Rostock, Germany

[†] Corresponding author: connor.bernard@zoo.ox.ac.uk

Page Count: 18

Word Count: 5,534

Figures: 4

Tables: 1

CONFLICT OF INTEREST STATEMENT

The authors report no conflicts of interest.

CONTRIBUTIONS

CB and RSG conceived the ideas; CB, RSG, GSS, JD, and PC designed the methodology; CB,

GSS, JD, PC, and SJLG integrated databases; CB, GSS, and JD collected primary records data;

EK led the climate data acquisition and analysis of geospatial data; and CB led the writing of the

manuscript with assistance from JJ. All authors contributed critically to the drafts and gave final

approval for publication.

DATA AVAILABILITY STATEMENT

The data in this paper and complete database are published open access in multiple formats (*e.g.*,

.csv, .pdf, .R, .tree) on the MOSAIC website https://mosaicdatabase.web.ox.ac.uk and associated

GitHub repository https://github.com/mosaicdatabase/mosaicdatabase.

Abstract (300/300)

1. The ecological sciences have joined the big data revolution. However, despite exponential growth in data availability, broader interoperability amongst datasets is still needed to unlock the potential of open access. The interface of demography and functional traits is well-positioned to benefit from said interoperability. Trait-based ecological approaches have been criticised because of their inability to predict fitness components, the core of demography; likewise, demographic approaches are data-hungry, and so using traits as ecological *shortcuts* to understanding and forecasting population viability could offer great value.

2. Here, we introduce MOSAIC, an open-access trait database that unlocks the demographic potential stored in the COMADRE, COMPADRE, and PADRINO open-access databases. MOSAIC data have been digitised and curated through a combination of existing datasets and additional taxonomic and/or trait records sourced from primary literature. In its first release, MOSAIC (v. 1.0.0) includes 14 trait fields for 300 animal and plant species: biomass, height, growth determination, regeneration, sexual dimorphism, mating system, hermaphrodism, sequential hermaphrodism, dispersal capacity, type of dispersal, mode of dispersal, dispersal classes, volancy, and aquatic habitat dependency. MOSAIC also includes species-level phylogenies for 1,359 species and population-specific climate data where locations are recorded.

3. Using MOSAIC, we highlight a taxonomic mismatch of widely used trait databases with existing structured population models. Despite millions of trait records available in open-access databases, taxonomic overlap between open-access demographic and trait databases is <5%. We identify where traits of interest to ecologists can benefit from

database integration and start to quantify traits that are poorly quantified (*e.g.,* growth

determination, modularity).

4. The MOSAIC database evidences the importance of improving interoperability in open-

access efforts in ecology as well as the need for complementary digitisation to fill

targeted taxonomic gaps. In addition, MOSAIC highlights emerging challenges

associated with the disparity between locations where different trait records are sourced.

KEYWORDS

COMADRE, COMPADRE, functional traits, gap of knowledge, integral projection model

(IPM), interoperability, matrix population model (MPM), open access, open traits, PADRINO.

"I'm not interested in your data; I'm interested in merging your data with other data.

Your data will never be as exciting as what I can merge it with."

74   —Sir Tim Berners-Lee (co-founder of the World Wide Web)

Introduction

The ecological sciences have recently joined the big data revolution (Farley, Dawson,

Goring, & Williams, 2018; Marx, 2013; Reichman, Jones, & Schildhauer, 2011). Total species

distribution records measure in the hundreds of millions (Maldonado et al., 2015; Troia &

McManamay, 2016). Functional trait data exist for tens of thousands of species across the globe

(Díaz et al., 2016; Enquist et al., 2019; Maitner et al., 2018). Similar gains have taken place in

related fields, from big omics (Field et al., 2009) to global state-of-the-art climate models (*e.g.*,

ERA-5 [https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5]) becoming available at

fine spatial and temporal resolutions suitable for biological analyses (Davy & Kusch, 2021;

Hersbach et al., 2020), and from behavioural trait data (Rowcliffe, Jansen, Kays, Kranstauber, &

Carbone, 2016) to population data (Global Population Dynamics Database

[https://knb.ecoinformatics.org/view/doi:10.5063/AA/nceas.167.15]; Human Mortality Database

(J.R. Wilmoth, K. Andreev, D. Jdanov, 2007; mortality.org); Human Fertility Database

(Jasilioniene et al., 2015; humanfertility.org); AnAge Database (De Magalhᾶes & Costa, 2009);

DATLife [Max Plank Institute of Demographic Research 2022; https://datlife.org/]). We have also

seen growth datasets characterising phenomena at levels of biological complexity above the

organismal level, including community ecology databases (BioTime (Dornelas et al., 2018); Web

of Life (Fortuna, Ortega, & Bascompote, 2014); metaCommunity Ecology: Species, Traits,

Environment and Space (CESTES) (Jeliazkov et al., 2020); Environmental Data Initiative

[https://environmentaldatainitiative.org/]). The growth of records in ecology datasets and

complementary environmental datasets is enabling us now to test ecological theory at larger and

more complex scales. In line with open-access growth, however, there is a recognised need to

improve and coordinate data standards to guide the systematic collection and management of data

across different trait collection programmes (Gallagher et al., 2020; Kissling et al., 2018;

Wilkinson et al., 2016).

Despite the increasing availability of biological data, combining different datasets into

common analyses is often hampered by the lack of complementarity between databases.

Interoperable systems reflect the transformation of constituent data types into a standardised

format of spatial, temporal, and measurement scales, accounting for differences in methods

(Nadrowski et al., 2013). The need to improve interoperability across datasets is demonstrated by

the widescale emergence of data harmonisation initiatives across fields of ecology (*e.g.,* Culina et

al., 2018; Kattge et al., 2020; Schneider et al., 2019). In the past decade, dozens of initiatives have

taken shape to both centralise data from existing datasets and to improve data interoperability:

standardising units, scales, and terminology for comparative purposes (Edwards, Lane, & Nielsen,

2000; Maurer, Firestone, & Scriver, 2000). Unifying data formats and streamlining their

integration unlocks the potential for existing datasets to answer questions that cut across levels of

biological complexity. Linking together levels of biological complexity is critical for identifying

how phenomena emerge and transmit across different levels of biological organisation, upscaling

and downscaling through biological systems. For example, the critical linkages between genetics

and biochemistry (The ENCODE Project Consortium, 2012; http://bigg.ucsd.edu/), biochemistry

and physiology (Vitousek, Johnson, & Husak, 2018), and physiology and demography (Laughlin,

Gremer, Adler, Mitchell, & Moore, 2020; Swenson et al., 2020) have benefitted from dataset

integration.

The limitations of global-scale datasets are shifting away from data availability and toward

data interoperability. The global-scale integration of functional traits is headed in this direction.

Momentum toward data integration is evidenced by recent database initiatives (*e.g.,* Guerrero-

Ramírez et al., 2021) and global networks that, like the Open Trait Network

(https://opentraits.org/), aim to standardise and integrate trait data across taxa (Gallagher et al.,

2020). Despite major improvements in the consolidation and accessibility of trait data, there is not

yet a single-source centralised database spanning behaviour, physiology, habitat, and other trait

data for a wide range of species. Existing databases are often linked by taxonomy (*e.g.,* FishBase

(Froese & Pauly, 2010), CoralTraits (Madin et al., 2016), MammalBase (Lintulaakso, 2013),

AmphiBio (Oliveira, São-Pedro, Santos-Barrera, Penone, & Costa, 2017)); trait type (*e.g.,*

TreeOfSex (The Tree of Sex Consortium 2014), TreeBase (Boettiger & Temple Lang, 2012),

Xylum Functional Traits (Borghetti, Gentilesca, Colangelo, Ripullone, & Rita, 2020)); data type

(*e.g.,* GBIF (https://www.gbif.org/), MOL (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012),

TetraDensity (Santini, Isaac, & Ficetola, 2018)); or a combination of the taxonomies and traits

(WooDiv (Monnet et al., 2021), CarniDiet (Middleton, Svensson, Scharlemann, Faurby, &

Sandom, 2021)). A number of other databases take a more general approach in their thematic

scope, but are still constrained to a limited set of traits and taxonomy (*e.g.,* Amniote (Myhrvold et

al., 2015), Pantheria (Jones et al., 2009), BIEN (Enquist, Condit, Peet, Schildhauer, & Thiers,

2016; Maitner et al., 2018), TRY (Kattge et al., 2020)). MOSAIC offers a platform that integrates

databases in the remit of species with structured population models across their ecological traits.

Here, we introduce MOSAIC, a centralized database of trait data across the Tree of Life.

MOSAIC is an open-access database that complements the existing demographic data available in

the COMPADRE Plant Matrix Database (Salguero-Gómez et al., 2015), COMADRE Animal

Matrix Database (Salguero-Gómez et al., 2016), and the new PADRINO IPM Database (Levin et

al., BioRxiv; doi: 10.1101/2022.03.02.482673; https://levisc8.github.io/RPadrino/). MOSAIC v.

1.0.0 includes 14 frequently used traits that encompass morphological, reproductive, dispersal, and

habitat type attributes for some 300 species of animals and plants. Additional traits will be added

in the future (see *Future Direction, below*). MOSAIC allows users to integrate structured

population data to probe larger questions through the collection, curation, and complementarity of

relevant contextual data.

Methods

Scope and Coverage of MOSAIC

The MOSAIC database (Version 1.0.0) includes 14 key trait records across 300 species

(Fig. 1). MOSAIC is designed to provide complementary data for analysis in connection with

structured population models: matrix population models (MPMs, Caswell, 2000), where state

variables are discrete (*e.g.*, age (Morris et al., 2008), ontogeny/development (Smallegange, Deere,

& Coulson, 2014), discrete classes of size (Crouse, Crowder, & Caswell, 1987)), and integral

projection models (IPMs, Easterling, Ellner, & Dixon, 2000), where the state variables are

continuous (*e.g.,* size (Jongejans, Shea, Skarpaas, Kelly, & Ellner, 2011), mass (Ozgul et al.,

2010), parasite load (Metcalf, Graham, Martinez-Bakker, & Childs, 2016)). The traits included in

MOSAIC 1.0.0 were identified by the MOSAIC team as a set of physical, physiological,

geographic, and behavioural attributes of most immediate relevance to demographic research (see

Table 1; more information at https://mosaicdatabase.web.ox.ac.uk). Traits were also selected in

consideration of the lack of standardised and centralised databases for certain traits (*e.g.,* volancy,

modularity, and growth indeterminacy; see Fig. 2 for trait variance and taxonomic structure of

select traits excluded from existing databases). Importantly, we note that MOSAIC is not a general

dataset for analysis of functional traits, as this is available through other extensive repositories

(*e.g.,* TRY (Kattge et al., 2020), BIEN (Enquist et al., 2016; Maitner et al., 2018)). Instead, the

focus of MOSAIC is on providing taxonomic coverage to species with open-access structured

population models available in the COMADRE (Salguero-Gómez et al., 2016), COMPADRE

(Salguero-Gómez et al., 2015), and PADRINO databases (Levin et al. forthcoming) (See Fig. 3

and Fig. 4 for spatial scope and taxonomic scope with respect to structured population databases,

respectively). MOSAIC provides a much-needed interoperability between existing databases that

are germane to demography. By doing so, MOSAIC helps to fill critical data needs of population

ecologists and functional trait ecologists (see Fig. 5 for relevant covariance structure). Although

demographically motivated, MOSAIC affords opportunities for data to address novel questions

more broadly across ecology, evolution, and conservation biology.

Accessing MOSAIC

A guide for downloading the MOSAIC database can be accessed on the MOSAIC portal

(https://mosaicdatabase.web.ox.ac.uk). The MOSAIC database is open access and can be

downloaded into R using a few lines of code (see: https://mosaicdatabase.web.ox.ac.uk/download-

database), which download the MOSAIC data object as an S4 object (Rda file). The database can

also be downloaded from GitHub directly as an (.Rda) or comma separated value (.csv) file and

loaded into R manually.

The MOSAIC portal hosts information pages and downloadable files about the database.

Users can download the *MOSAIC User Guide*, *vignettes*, and *data collection protocols* from the

portal webpage (https://mosaicdatabase.web.ox.ac.uk/vignettes and

https://mosaicdatabase.web.ox.ac.uk/user-guide). The *User Guide* specifies the classifications,

precision, and data types for each trait field. The MOSAIC *User Guide* details the metadata on the

structure of data for each trait (*e.g.,* species, genus, mosaic index). An updated list of the databases

that are directly or indirectly addressed by the MOSAIC database is maintained on the MOSAIC

portal.

Data Sources

MOSAIC is both a *meta*-database (a database of databases) and a database in its own right,

containing new trait records from primary literature (Fig. 1). The MOSAIC database contains

records centralised from existing datasets where functional traits relevant to population ecologists

can be openly accessed and redistributed (*e.g.,* BIEN (Enquist et al., 2016; Maitner et al., 2018)).

The records reflected in the MOSAIC database do not encompass the entirety of the source

databases but are instead partial facsimiles of those databases that reflect records relevant to

population databases (COMADRE, COMPADRE, PADRINO). MOSAIC has three major

components: (1) trait records sourced from existing databases (22%); (2) trait records newly

procured through searching the primary literature (71%); and (3) trait record markers that direct

users to additional records that are not included in the MOSAIC database (7%). MOSAIC trait

markers exist for one of two reasons: the database containing the records of interest does not allow

records to be accessed or limits redistribution rights behind individual registration and/or specific

use applications; or records in other databases contain multiple records for a species, which do not

currently fit within the data structure of version 1.0.0 of MOSAIC (see *Future Targets for*

MOSAIC).

In addition to identifying whether a trait record is new, the "MOSAIC" attribute field also

identifies whether the attribute field (*i.e.,* trait name) is part of an existing database. For example,

the MOSAIC field might indicate that a record for specific leaf area is new for a specific species,

and also that the attribute is part of databases such as TRY (Kattge et al., 2020) or BIEN (Enquist

et al., 2016; Maitner et al., 2018). By contrast, a new species record for volancy would indicate

that there are currently no databases that systematically collect data on volancy attributes and

therefore that all volancy records in the MOSAIC database are new.

Organisation of Sources in MOSAIC

The MOSAIC attribute field is a factorial variable with three levels: MOSAIC-A,

MOSAIC-B, and MOSAIC-C. The first of these levels, MOSAIC-A (Fig. 1), labels only records

that reflect existing databases (*i.e.*, provenance of an existing data acquisition service); the second,

MOSAIC-B, labels new records collected by the MOSAIC team that are in a trait field within the

scope of an existing database initiative (*e.g.,* specific leaf area in BIEN); and the third, MOSAIC-

C, labels new records collected by the MOSAIC team on traits for which there is not currently a

database initiative centralising records. If a datum has been adopted from another dataset or

database, then the relevant source is referenced in the *Database* attribute column. Note that this

value will be "NA" for all MOSAIC-C records, logically. Over time, data sharing will move

records in MOSAIC-B to MOSAIC-A as the MOSAIC-B traits are assimilated in the database

networks that specialise on an existing trait (*i.e.,* data feedback; see Fig. 1).

Sources and Provenance of Records

Because of existing limitations on data access, some datasets cannot be transferred into

MOSAIC. Where data exist outside of the MOSAIC platform, but have restricted access, the

MOSAIC database directs users toward the appropriate database on a trait and taxa-specific level

(see `metaMOSAIC`). The *MOSAIC User Guide* (Appendix S1) explains differences between data

gaps that are yet to be reviewed and those that are true gaps (*e.g.,* volancy/flight capability of

plants). Data were obtained through searching peer-reviewed records of ISI Web of Science,

Scopus, and Google Scholar using key words pertinent to the species name and trait field in

question (see Appendix S2 for a list of keywords queried). The MOSAIC team also searched the

archives of data repositories, including the Dryad digital repository (http://datadryad.org; see

Appendix S3 for a complete list of repositories reviewed – to be maintained hereafter on the

MOSAIC portal), and reviewed in detail journal archives that have a high occurrence of data

publishing, including *Nature Scientific Data*, *Methods in Ecology and Evolution*, and *Journal of*

Ecology. MOSAIC also reflects a review of data from data aggregating servers, such as the open

traits network (https://opentraits.org/), the ecological data wiki (https://ecologicaldata.org/),

environmental data initiative (https://environmentaldatainitiative.org/), and databases that

aggregate other databases (*e.g.,* BIEN (Enquist et al., 2016; Maitner et al., 2018) and TRY (Kattge

et al., 2020)). The complete set of key words used in this review is detailed in a supplement to the

User Guide posted on the MOSAIC portal. A current list of databases reviewed in the development

of MOSAIC is included in Appendix S3. Suggested data sources and key terms can be submitted

through the MOSAIC data portal.

Database Navigation

The *User Guide* contains detailed information on navigating the MOSAIC R data object.

In general, the data object can be queried with the syntax:

`mosaic@trait.name@values[[mosaic.index]]`. In the aforementioned syntax, the

`trait.name` and `mosaic.index` should reflect one trait and one or more species index values

in the respective locations (*e.g.,* `mosaic@volancy@value[[93]]`). The complete set of

records for a trait can be viewed and assigned by calling the trait from the mosaic object without

an index specified (*e.g.,* `mosaic@volancy@value`). The mosaic index value reflects a unique

identifier that can be linked from COMADRE, COMPADRE, and PADRINO through the index

field   of   mosaic   (*e.g.,*

`comadre@data$MatrixID[comadre@data$SpeciesAccepted=="Alces`

`alces"]))`. Associated metadata for each record can be queried by the following syntax:

`mosaic@trait.name@metadatafield[[mosaic.index]]`.   Here,   the

`metadatafield`   should   reflect   one   metadata   element   (*e.g.,*

`mosaic@volancy@author[[93]]`). The complete metadata for a species can be viewed

using the `mos_meta()` function or searched manually by slots, or by querying rows of the

database in the comma separated value format (https://mosaicdatabase.web.ox.ac.uk/navigating-

mosaic). See the *User Guide* (Appendix S1) for more information. The database is organised to be

readily subsetted, including compatibility with Tidyverse package (Wikham et al. 2019) syntax

(see *Navigating MOSAIC*; https://mosaicdatabase.web.ox.ac.uk/vignettes, also in Appendix S4).

The MOSAIC object can be searched in R by taxonomy, trait, locality, duration of study, sex, and

other metadata fields applicable to COMADRE, COMPADRE, and PADRINO (see *vignettes*,

above, also Appendix SX). Trait values in the database are measured in standardised units of

measure for each trait (Table 1). Species names are specified in conformance with the Catalogue

of Life (www.catalogueoflife.org), also consistent with the COMPADRE, COMADRE, and

PADRINO databases. The *User Guide* provides specific guidelines for querying fields within the

data   object   in   R   and   for   negotiating   the   dataset   in   finer   detail

(https://mosaicdatabase.web.ox.ac.uk/user-guide).

Error Reporting

The MOSAIC portal has an *Error Report* page for reporting potentially erroneous records,

or to query additional questions (https://mosaicdatabase.web.ox.ac.uk/suggested-additions-error-

reporting; but see also FAQ: https://mosaicdatabase.web.ox.ac.uk/frequently-asked-questions).

Potential errors can be reported anonymously or with contact information (*e.g.*, name, email).

Decisions on reported errors will be disclosed on the Error Report page

(https://mosaicdatabase.web.ox.ac.uk/suggested-additions-error-reporting) and to the reporting

party if contact information is included in the request. If it is not possible to submit errors via the

portal, users can also to reach out by email to: mosaicdatabase@biology.ox.ac.uk.

Recommended Records

The MOSAIC portal has a *Recommended Records* page

(https://mosaicdatabase.web.ox.ac.uk/suggestion-additions) for reporting suggested records that

are not included in the MOSAIC database. Recommendations can be made with or without contact

information. Contact information will be used exclusively for clarifying questions and updating

the commenter when records are included. MOSAIC will report decisions on the

Recommendations Incorporated page (https://mosaicdatabase.web.ox.ac.uk/suggestion-

additions). Users may also request new data fields to be prioritised in future rollouts. Given the

realities of limiting resources, The MOSAIC team will do their best to include the requested

records in future versions. Likewise, we also welcome submissions of data collected by individuals

via mosaicdatabase@biology.ox.ac.uk.

metaMOSAIC: **licensed data, access limitations, and restricted redistribution of existing**

records and databases

Not all datasets permit open use, dissemination, and redistribution of their trait data. Where

limitations on the data collation and redistribution apply, there may be application procedures,

registration, and other actions necessary for an individual to obtain access to specific trait records

for analysis (*e.g., TRY*). MOSAIC centralises the metadata for datasets that do not allow data to be

redistributed to help navigate to relevant data resources outside the scope of open access. Records

in limited access databases can be searched in MOSAIC by taxonomic group and by trait.

MOSAIC links researchers to application materials for requesting access to those limited-access

databases. The data access of licensed or non-open-access databases is stored in a data object called

`metaMOSAIC` that is an extension of the MOSAIC database. The MOSAIC dataset thus provides

data where it is accessible and `metaMOSAIC` guides researchers to where data exist with

registration. When searching fields in the MOSAIC database, the `metaMOSAIC` adjunct dataset

indicates if data are available in these ancillary sources (see *User Guide* in Appendix S1 for more

information) and provide links to pertinent sources and data through the provider. `metaMOSAIC`

is part of the main database object accessible through the MOSAIC portal.

Database Updates

The MOSAIC database will be updated as new data are added to the COMADRE,

COMPADRE, and PADRINO databases. Updates of the MOSAIC database will account for newly

discovered data sources and new literature that adds to or changes the species-level traits in the

database, as well as correct errors from earlier versions. New MOSAIC versions will be released

periodically with a notice published on the website, in the data object metadata, the mosaic GitHub

page (https://github.com/mosaicdatabase), and through updates in associated packages in

conformance with standard semantic versioning (a three-part version code reflecting major, minor,

and patch updates, in respective positions). Updates will be published to the mosaic portal

(https://mosaicdatabase.web.ox.ac.uk), social media (Twitter: @mosaicdatabase), and GitHub

(https://github.com/mosaicdatabase). The MOSAIC portal includes a location for submission of

recommendations for additions or suggested changes to the database.

Future Targets of MOSAIC

Versioning

MOSAIC database updates and version history will be posted to the MOSAIC portal in the

future.

Taxonomic Scope

Future versions of MOSAIC are scheduled for expansion to include 50 core traits across

some 1,500 species (the current entirety of COMPADRE, COMADRE, and PADRINO; *MOSAIC*

target traits; see *User Guide* [Appendix S1]). The MOSAIC core traits list will guide the future

roll-out of attributes gathered and stewarded by the MOSAIC team (available on the MOSAIC

portal website; see below), but additional trait suggestions can be submitted through the online

portal.

Interspecific Variation in Hierarchical Data Structures

Future versions of the MOSAIC database will adopt a hierarchical file structure that will

accommodate multiple records per species. Once records for COMADRE, COMPADRE, and

PADRINO are fully populated across the MOSAIC traits with mean, pooled, or other

representative quantities (*e.g.,* mean leaf size for all plants or adult bodymass of animals),

secondary records will be added. Existing trait databases may contain multiple records per species

(see, for example, structure of COMADRE (Salguero-Gómez et al., 2016), COMPADRE

(Salguero-Gómez et al., 2015), BIEN (Enquist et al., 2016; Maitner et al., 2018), TRY (Kattge et

al., 2020)), although some databases host a single record per species' trait, such as age and growth

rate for animals in AnAge (De Magalhães & Costa, 2009). To facilitate research into intra-specific

trait variation (Albert, 2015; Messier, McGill, & Lechowicz, 2010; Siefert et al., 2015), MOSAIC

provides provenance of records, whether records were subject to selection or merger (means or

pooling), and fields that identify whether multiple records are known to exist. Where records for a

given species were isolated from existing databases, mean values are often retrieved, and the

database sourcing additional data is noted in the database under the attributed field ("Additional

Trait Data Available" field).

citMOSAIC: Citizen Science

In addition to `metaMosaic`, which guides users to licenced data not reported in the

MOSAIC database, MOSAIC plans to roll out a database of identical structure to MOSAIC that

gathers information from citizen science datasets. Like MOSAIC, citMOSAIC will have three

components (citMOSAIC-A, citMOSAIC-B, citMOSAIC-C), reflecting the same relationship of

databases and fields to MOSAIC (see https://mosaicdatabase.web.ox.ac.uk/citizen-mosaic for

more information). citMOSAIC will be kept independent of the main MOSAIC database to avoid

conflation of peer reviewed literature from citizen science data. Where appropriate to use datasets

together, the metadata, query functionality, and design of citMOSAIC will mirror that of MOSAIC

to promote interoperation of databases. citMOSAIC will be downloadable from the MOSAIC

portal website.

Cautionary Notes

Records in the MOSAIC database are gathered and standardised under the protocols

detailed in the *User Guide* (Appendix S1; also available through the MOSAIC portal). Users

should be attentive to the precision, levels, and context of data in MOSAIC when used for analysis.

For those records in the dataset that come from multiple individuals, we present them as statistical

components (*e.g.,* minimum, maximum, or mean trait values). Functional traits in MOSAIC may

be estimated from populations studied in COMADRE, COMPADRE, PADRINO, or other

databases (Table S3). The potential temporal and spatial mismatch between databases that are

linked in an analysis merits close attention (Csergő et al., 2017). The studies in the MOSAIC

database also include research conducted by different investigators using independent tools,

technologies, sample designs, and study methods. The influence of research methods and

instruments on the error values in the dataset may require additional consideration for potential

bias, noise, and imprecision. Where more than one life history trait value exists for a given species,

MOSAIC users will need to determine whether averaging or selective filtering to one study is most

appropriate in view of the specifics of the given research question. In certain cases, trait values for

a species might only be available for a single st/age and therefore may not provide a complete

picture of the trait variation amongst st/ages. Users are encouraged to be cautious when

contextualising the scope of representation of the values in the database and their analyses.

Representation, Variance, and Bias in MOSAIC v1.0.0

MOSAIC can help to mechanistically examine the causes and correlates of demographic

variability across the thousands of animal and plant species housed in COMPADRE, COMADRE,

and PADRINO. In its version 1.0.0, MOSAIC contains a high degree of variance in trait values

known to shape demographic outcomes across major taxonomic groups. For example, determinant

growth is present for 0% of Amphibians and Birds and 100% of Bivalves, while volancy is present

for 0% in Reptiles, Amphibians, and Bony Fish and 77% in Birds (Fig. 2). Animal adult biomass

and plant height follow lognormal trait distributions (see Appendix S5). 11% of mammals are

indeterminate growers *vs.* 54% of reptiles and 0% birds, and 95% of mammals are monogamous

compared to 100% of reptiles and 20% of birds in MOSAIC. Recent studies have examined how

different vital rates are explained by functional traits (e.g., Carmona, Bello, Azcarate, Mason, &

Peco 2019; Pistón et al. 2018; Adler et al. 2014; Poorter et al. 2008). However, understanding how

trait variation across taxa translates to demographically influential properties remains

underdeveloped.

MOSAIC's initial release (v1.0.0) includes records for all major regions of the globe for

which we have structured population models (Fig.3). Nevertheless, species trait values are not

necessarily gathered from the same localities as population models (see *future directions* for more

information on systematising spatial mismatch). This is an important consideration for users of

MOSAIC (and more generally of trait-based approaches) wishing to bring together functional traits

and demographic rates, as traits and vital rates are known to vary considerably within the same

species across spatial scales (Violle et al. 2012; Albert et al. 2010; Bolnick et al. 2011). Moreover,

while there is at least one trait for each of these locations, the data density remains variable. Thus,

records are not necessarily representative of the global spread or the full spatial scope of MOSAIC.

For example, the highest complete coverage for MOSAIC traits is concentrated toward localities

with the with the longest-term demographic models (see COMPADRE locations associated with

MOSAIC records).

Phylogenetically, the initial release of MOSAIC is somewhat limited. Version 1.0.0 covers

300 of the 1,400 species currently available in COMPADRE, COMADRE and PADRINO.

However, MOSAIC trait data are well distributed across clades (Fig. 4). While there is not a highly

skewed phylogenetic concentration with respect to the existing structured population models or

clustering of records into small groups across the Tree of Life, phylogenetic density of MOSAIC

records remains low. Therefore, information from the MOSAIC database may be limited for a

given genus or order and, as such, should be approached with caution. In future versions of

MOSAIC, the phylogenetic bias is expected to diminish with more samples and stronger

phylogenetic representation.

Covariance across traits is also an important source of confound in existing analyses linking

traits and vital rates. Positive and negative correlations across traits that have independent

influences on vital rates can create apparent associations of demographic properties with traits,

spuriously functionalising non-functional traits (McElreath 2021). Disentangling the relevance of

key axes of trait variation for their demographic influences demands a clear quantification of the

direction and strength networks of trait associations, trade-offs, and demographic consequences.

Population biologists seek to understand not only how individual traits relate to different aspects

of demographic performance (*e.g.,* population growth rate, risk of quasi-extinction, *etc.*), but also

understanding how trait syndromes shape those demographic outcomes. MOSAIC presents a

highly varied covariance structure in trait values for the examined 300 species. For example,

without *a priori* expectation, indeterminate growth and regeneration traits show strong correlation

(r=0.51; Fig. 5), which could influence each other's effects on vital rates. The same could be argued

for the correlation between volancy and reproductive strategy (r=0.28 with monogamy; Fig. 5).

The MOSAIC database can be used as a platform to showcase the lack of overlap between

trait and vital rate data for the same species. This picture calls for a more systematic way address

global biases in ecological data quantification/collection. Even where we have complete

information about species in the COMADRE, COMPADRE, and PADRINO databases, we are

subject to the constraints and biases of those datasets, such as spatial bias toward high-GDP

countries and the phylogenetic bias toward temperate regional perennial plants (Kendall et al.

2019; Römer et al. 2021). The compounding of error and density across datasets highlights the

need to prioritise stronger representation of functional traits linked with demography. The

standardised framework of MOSAIC is an ideal platform to work from to achieve this goal.

In view of potential biases introduced by low sampling density and the patchiness of cover

in traits, users of the database are advised to consult the literature about the representativeness and

congruency between MOSAIC data and related trait diversity within clades. Users need to be

mindful of the scope of the questions that they are setting out to answer and to be aware of the

influence of sample sizes and coherency or heterogeneity of traits across different taxonomic

levels.

Discussion

From Databases to Data Networks

Broad aperture digitisation efforts (*e.g.,* BIEN (Enquist et al., 2016; Maitner et al., 2018)

and TRY (Kattge et al., 2020)) have helped resolve many answers to demographic questions.

Examples include whether there are trait spectra and key trade-off patterns amongst functional

traits and whether these are correlated with particular environments and life history strategies

(Laughlin, 2018; Reich et al., 2003; Wilkes et al., 2020), Trait-based ecology and Trait Driver

Theory (Enquist et al., 2015) are indebted to such opportunity-driven research programmes. More

generally, however, the trait-based ecology paradigm has failed to support clear answers to many

research questions of central interest to demographers (Bellier, Kéry, & Schaub, 2018; Laughlin

et al., 2020). This limited reach of the functional trait programme coincides with a dearth of

species-specific overlap across the range of functional traits that are collected by the functional

trait databases.

The proliferation of databases and open data initiatives over the last two decades

(Cheruvelil & Soranno, 2018) evidences an interest in improving both data access and data

usability (Gallagher et al., 2020; Michener, 2006; Whitlock, 2011). While existing databases

standardise trait fields, collate records, and link associated metadata, existing databases often store

data for simple, quantitative traits. Relatively few ecological trait databases store diverse data types

(such as rate arrays, population time series, physiological rates at different structural levels, and

habitat shape files) that may be associated with multidimensional, ecological study systems (but

see CESTES (Jeliazkov et al., 2020), GFBio (Diepenbroek et al., 2014), DarwinCore (Wieczorek

et al., 2012)).

The digitisation and standardisation of *existing* data and their integration with

complementary, new data presents a growing set of challenges and opportunities in ecology

(Salguero-Gómez, Jackson, Gascoigne 2021). Efforts to gap-fill records can leverage the value of

existing datasets while expediting the schedule for specific research outcomes. As trait datasets

grow, the importance of targeted, gap-filling initiatives to address bias and to capitalise on existing

data will also increase (Conde et al., 2019). The value of existing records is further enhanced

through improvements in the interoperability of datasets. Much of this work is done manually, at

a high cost, and with little support from funding agencies (Salguero-Gómez, Jackson, Gascoigne

2021), and yet it has been effective at facilitating research and creating new value for old data. In

recent years, initiatives have sought to improve the interoperability of datasets by guiding

prospective data structure or retroactively harmonising existing datasets. These include

programmes that develop universal standards to improve global interoperability (such as

DarwinCore (Wieczorek et al., 2012) and Frictionless [https://frictionlessdata.io/]data standards)

or that contain guidelines for data metastructure (such as the FAIR principles (findability,

accessibility, interoperability, and reusability, *sensu* Wilkinson et al., 2016) and the OpenTraits

framework (Gallagher et al., 2020)). These initiatives address emerging and scaling challenges of

ecoinformatics, such as the protocols by which we share data, search data, and preserve provenance

within hierarchical data storage structures. These protocols will be essential in centralising datasets

as diverse as government monitoring datasets (*e.g.,* those stored in U.S. Data clearing houses

[https://www.data.gov;   https://www.dataone.org/];   National   Biodiversity   Atlas

[https://nbnatlas.org/]); centralised monitoring and experimental networks (*e.g.,* LTER and

NEON), raw or reanalysed remote sensing datasets (*e.g.,* Landsat data, NASA EarthData datasets,

ERA-5 data), and private datasets (https://www.natureserve.org/) that will demand versatile and

navigationally efficient data structures.

Population ecology has benefitted from widespread open-access databases but requires

further dataset integration to answer its central questions. Understanding whether and how some

morphological or physiological traits predict demographic outcomes and why others fail to do so

is of central interest to questions in physiological, population, and community ecology (Adler et

al., 2014; Chalmandrier et al., 2021; Laughlin et al., 2020; Yang, Cao, & Swenson, 2018).

Population ecologists routinely use data that are distributed across a wide range of databases.

Comparative and macroecological researchers use phylogenies (*e.g.*, Daskalova, Myers-Smith, &

Godlee, 2020; Levin, Crandall, Pokoski, Stein, & Knight, 2020; Roper, Capdevila, & Salguero-

Gómez, 2021), adult bodymass (*e.g.*, Capdevila et al., 2020; Healy et al., 2014; Terry, O'Sullivan,

& Rossberg, 2022; Williams, McRae, Freeman, Capdevila, & Clements, 2021), and high-

resolution, global climate information (*e.g.*, Daskalova, Bowler, Myers-Smith, & Dornelas, 2021;

Paniw, Maag, Cozzi, Clutton-Brock, & Ozgul, 2019; Stenseth & Mysterud, 2002) to answer

relevant biological, evolutionary and ecological questions and to contextualise their findings.

Population ecologists frequently examine a subset of physiological, morphological, and

behavioural attributes associated with demographic outcomes ("functional traits", *sensu* Violle et

al., 2007). The trait-based research programme seeks to, among other aims (*e.g.*, Díaz et al., 2016),

identify the intrinsic and extrinsic regulators of vital rates and the causes of variation and

constraints on possible trait values (Buckley & Puy, 2022; Salguero-Gómez, Violle, Gimenez, &

Childs, 2018). Not all traits predict demographic outcomes and functional traits may exercise

influence on only a few demographic pathways (Pistón et al., 2019; Salguero-Gómez & Laughlin,

2021). The answers to these questions rely on the existence of vital rate and trait data, the overlap

of which has been limited in the absence of targeted attention. For instance, of the hundreds of

thousands of records available across thousands of plant species in TRY (Kattge et al., 2020) and

over 345 plant species in COMPADRE (Salguero-Gómez et al. 2015, COMPADRE Ver 1.0.0),

Adler et al. (2014) report functional trait-vital rate relationships for only 222 plant species due to

their limited data overlap.

Ecological data are complex and their structures will need consistent rules to link datasets

together. It will be important for future datasets to adopt hierarchical database designs that render

large, thematically, and structurally diverse data to be readily locatable and usable without expert

knowledge. Here, we show one such example in the scope of comparative research, using thematic

groups and a strategy of achieving adequate record breadth before revisiting depth of records for

specific species. The need for open access data, integrated workflows, and interoperable data

systems is increasing with the scaling of data collection through use of robotics and technologies.

The gaps in existing data systems, interoperability, and data acquisition can be filled strategically

for specific applications, offering targeted and efficient dataset development. With data

interoperability guiding the structure of new datasets, the modular development of area-specific

datasets will enable more generalised use over time and help meet the aims of existing database

initiatives.

ACKNOWLEDGEMENTS

We thank Mark Roper, Sara Middleton, Etienne Cousin, and Thomas Marrien for their

contributions to discussion at the beginning of the project, Jessica Hass and Lauren Hinchcliffe for

collecting data for the project, John Park for his contributions to the conception of citMOSAIC,

and Sam Levin for valuable comments on the Rmosaic package. This work was supported by a

NERC IRF (NE/M018458/1) to RS-G. RCRC acknowledges postdoctoral support from the

Regional Valencian Government and the European Social Fund (APOSTD/2020/090).

ORCID

Connor Bernard: 0000-0003-2175-5712

Gabriel Santos: 0000-0001-7991-8807

Jacques Deere: 0000-0001-6736-2223

Roberto Rodriguez-Caro: 0000-0003-2321-9497

Pol Capdevila: 0000-0002-2842-4302

Erik Kusch: 0000-0002-4984-7646

Samuel J L Gascoigne: 0000-0002-2984-1810

John Jackson: 0000-0002-1401-3035

Roberto Salguero-Gómez: 0000-0002-6085-4433

References

Adler, P. B., Salguero-Gómez, R., Compagnoni, A., Hsu, J. S., Ray-Mukherjee, J., Mbeau-Ache, C., & Franco, M. (2014). Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(27), 10019. doi: 10.1073/pnas.1410430111

Albert, C. H. (2015). Intraspecific trait variability matters. *Journal of Vegetation Science*, *26*(1), 7–8. doi: 10.1111/jvs.12240

Albert, C.H., Thuiller, W., Yoccoz, N.G., Soudant, A., Boucher, F., Saccone, P. & Lavorel, S. (2010) Intraspecific functional variability: extent, structure and source of variation. *The Journal of Ecology*, 98, 604–613.

Bellier, E., Kéry, M., & Schaub, M. (2018). Relationships between vital rates and ecological traits in an avian community. *Journal of Animal Ecology*, *87*(4), 1172–1181. doi: 10.1111/1365-2656.12826

Boettiger, C., & Temple Lang, D. (2012). Treebase: An R package for discovery, access and manipulation of online phylogenies. *Methods in Ecology and Evolution*, *3*(6), 1060–1066. doi: 10.1111/j.2041-210X.2012.00247.x

Bolnick, D.I., Amarasekare, P., Araujo, M.S., Burger, R., Levine, J.M., Novak, M. et al. (2011). Why intraspecific trait variation matters in community ecology. *Trend Ecol. Evol.*, 24, 183–192.

Borghetti, M., Gentilesca, T., Colangelo, M., Ripullone, F., & Rita, A. (2020). Xylem Functional Traits as Indicators of Health in Mediterranean Forests. *Current Forestry Reports*, *6*(3), 220–236. doi: 10.1007/s40725-020-00124-5

Buckley, Y. M., & Puy, J. (2022). The macroecology of plant populations from local to global scales. *New Phytologist*, *233*(3), 1038–1050. doi: 10.1111/nph.17749

Capdevila, P., Beger, M., Blomberg, S. P., Hereu, B., Linares, C., & Salguero-Gómez, R. (2020). Longevity, body dimension and reproductive mode drive differences in aquatic versus terrestrial life-history strategies. *Functional Ecology*, *34*(8), 1613–1625. doi: 10.1111/1365-2435.13604

Carmona, C. P., de Bello, F., Azcarate, F. M., Mason, N. W. H., & Peco, B. (2019). Trait hierachies and intraspecific variability drive competitive interactions in Mediterranean annual plants. *Journal of Ecology*, 107, 2078–2089. https://doi.org/10.1111/1365-2745.13248

Caswell, H. (2001). *Matrix population models* (2nd editio). Sunderland, MA: Sinauer.

Chalmandrier, L., Hartig, F., Laughlin, D. C., Lischke, H., Pichler, M., Stouffer, D. B., & Pellissier, L. (2021). Linking functional traits and demography to model species-rich communities. *Nature Communications*, *12*(1). doi: 10.1038/s41467-021-22630-1

Cheruvelil, K. S., & Soranno, P. A. (2018). Data-intensive ecological research is catalyzed by open science and team science. *BioScience*, *68*(10), 813–822. doi: 10.1093/biosci/biy097

Conde, D. A., Staerk, J., Colchero, F., da Silva, R., Schöley, J., Maria Baden, H., … Vaupel, J. W. (2019). Data gaps and opportunities for comparative and conservation biology. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(19), 9658–9664. doi: 10.1073/pnas.1816367116

Consortium, E. P. (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature*, *489*(7414), 57–74. doi: 10.1038/nature11247

Crouse, D. T., Crowder, L. B., & Caswell, H. (1987). A stage-based population model for

loggerhead sea turtles and implications for conservation. *Ecology*, *68*(5), 1412–1423.

Csergő, A. M., Salguero-Gómez, R., Broennimann, O., Coutts, S. R., Guisan, A., Angert, A. L.,
… Buckley, Y. M. (2017). Less favourable climates constrain demographic strategies in
plants. *Ecology Letters*, *20*(8), 969–980. doi: 10.1111/ele.12794

Culina, A., Baglioni, M., Crowther, T. W., Visser, M. E., Woutersen-Windhouwer, S., & Manghi,
P. (2018). Navigating the unfolding open data landscape in ecology and evolution. *Nature*
Ecology and Evolution, *2*(3), 420–426. doi: 10.1038/s41559-017-0458-2

Daskalova, G. N., Bowler, D., Myers-Smith, I. H., & Dornelas, M. (2021). Representation of
global change drivers across biodiversity datasets. *EcoEvoRxiv*, 1–36. Retrieved from
https://doi.org/10.32942/osf.io/db4s7

Daskalova, G. N., Myers-Smith, I. H., & Godlee, J. L. (2020). Rare and common vertebrates span
a wide spectrum of population trends. *Nature Communications*, *11*(1), 1–13. doi:
10.1038/s41467-020-17779-0

Davy, R., & Kusch, E. (2021). Reconciling high resolution climate datasets using KrigR.
Environmental Research Letters.

De Magalhães, J. P., & Costa, J. (2009). A database of vertebrate longevity records and their
relation to other life-history traits. *Journal of Evolutionary Biology*, *22*(8), 1770–1774. doi:
10.1111/j.1420-9101.2009.01783.x

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., … Gorné, L. D.
(2016). The global spectrum of plant form and function. *Nature*, *529*(7585), 167–171. doi:
10.1038/nature16489

Diepenbroek, M., Glöckner, F. O., Grobe, P., Güntsch, A., Huber, R., König-Ries, B., … Triebel,
D. (2014). Towards an Integrated Biodiversity and Ecological Research Data Management
and Archiving Platform : The .... *Informatik 2014 – Big Data Komplexität Meistern. GI-*
Edition: Lecture Notes in Informatics (LNI) – Proceedings 232, (November), 1711–1721.
Retrieved from https://dl.gi.de/handle/20.500.12116/2782

Dornelas, M., Antão, L. H., Moyes, F., Bates, A. E., Magurran, A. E., Adam, D., … Zettler, M. L.
(2018). BioTIME: A database of biodiversity time series for the Anthropocene. *Global*
Ecology and Biogeography, *27*(7), 760–786. doi: 10.1111/geb.12729

Easterling, M. R., Ellner, S. P., & Dixon, P. M. (2000). Size-specific sensitivity: Applying a new
structured population model. *Ecology*, *81*(3), 694–708.

Edwards, J. L., Lane, M. A., & Nielsen, E. S. (2000). Interoperability of biodiversity databases:
Biodiversity information on every desktop. *Science*, *289*(5488), 2312–2314. doi:
10.1126/science.289.5488.2312

Enquist, B. J., Condit, R., Peet, R. K., Schildhauer, M., & Thiers, B. (2016). The Botanical
Information and Ecology Network (BIEN): Cyberinfrastructure for an integrated botanical
information network to investigate the ecological impacts of global climate change on plant
biodiversity. *PeerJ Preprints*.

Enquist, B. J., Feng, X., Boyle, B., Maitner, B., Newman, E. A., Jørgensen, P. M., … McGill, B.
J. (2019). The commonness of rarity: Global and future distribution of rarity across land
plants. *Science Advances*, *5*(11), 1–14. doi: 10.1126/sciadv.aaz0414

Enquist, B. J., Norberg, J., Bonser, S. P., Violle, C., Webb, C. T., Henderson, A., … Savage, V.
M. (2015). Scaling from traits to ecosystems: developing a general trait driver theory via
integrating trait-based and metabolic scaling theories. In *Advances in ecological research* (pp.
249–318).

Farley, S. S., Dawson, A., Goring, S. J., & Williams, J. W. (2018). Situating ecology as a big-data

science: Current advances, challenges, and solutions. *BioScience*, *68*(8), 563–576. doi:
10.1093/biosci/biy068
Field, D., Sansone, S., Collis, A., Booth, T., Dukes, P., Gregurick, S. K., … Remington, K. (2009).
'Omics Data Sharing. *Science*, *326*, 234–236.
Fortuna, M. A., Ortega, R., & Bascompote, J. (2014). The web of life. *ArXiv*. doi:
10.4324/9780203410134
Froese, R., & Pauly, D. (2010). *FishBase*. Retrieved from http://www.ices.dk/sites/pub/CM
Doccuments/1992/L/1992_L10.pdf
Gallagher, R. V., Falster, D. S., Maitner, B. S., Salguero-Gómez, R., Vandvik, V., Pearse, W. D.,
… Enquist, B. J. (2020). Open Science principles for accelerating trait-based science across
the Tree of Life. *Nature Ecology and Evolution*, *4*(3), 294–303. doi: 10.1038/s41559-020-
1109-6
Guerrero-Ramírez, N. R., Mommer, L., Freschet, G. T., Iversen, C. M., McCormack, M. L.,
Kattge, J., … Weigelt, A. (2021). Global root traits (GRooT) database. *Global Ecology and*
Biogeography, *30*(1), 25–37. doi: 10.1111/geb.13179
Healy, K., Guillerme, T., Finlay, S., Kane, A., Kelly, S. B. A., McClean, D., … Cooper, N. (2014).
Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proceedings of*
the Royal Society B: Biological Sciences, *281*(1784). doi: 10.1098/rspb.2014.0298
Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., … Thépaut, J.
N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological*
Society, *146*(730), 1999–2049. doi: 10.1002/qj.3803
J.R. Wilmoth, K. Andreev, D. Jdanov, and D. A. G. (2007). Methods Protocol for the Human
Mortality Database. *Database*, *2007*(June 2000), 1–80.
Jasilioniene, A., Jdanov, D. A., Sobotka, T., Andreev, E. M., Zeman, K., & Shkolnikov, V. M.
(2015). Methods Protocol for the Human Fertility Database. In *Max Planck Institute for*
Demographic Research. Retrieved from http://www.humanfertility.org/Docs/methods.pdf
Jeliazkov, A., Mijatovic, D., Chantepie, S., Andrew, N., Arlettaz, R., Barbaro, L., … Chase, J. M.
(2020). A global database for metacommunity ecology, integrating species, traits,
environment and space. *Scientific Data*, *7*(1), 1–15. doi: 10.1038/s41597-019-0344-7
Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity
of birds in space and time. *Nature*, *491*(7424), 444–448. doi: 10.1038/nature11631
Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D. L., … Purvis, A. (2009).
PanTHERIA: a species-level database of life history, ecology, and geography of extant and
recently extinct mammals. *Ecology*, *90*(9), 2648–2648. doi: 10.1890/08-1494.1
Jongejans, E., Shea, K., Skarpaas, O., Kelly, D., & Ellner, S. P. (2011). Importance of individual
and environmental variation for invasive species spread: A spatial integral projection model.
Ecology, *92*(1), 86–97. doi: 10.1890/09-2226.1
Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., … Wirth, C. (2020). TRY
plant trait database – enhanced coverage and open access. *Global Change Biology*, *26*(1),
119–188. doi: 10.1111/gcb.14904
Kissling, W. D., Walls, R., Bowser, A., Jones, M. O., Kattge, J., Agosti, D., … Guralnick, R. P.
(2018). Towards global data products of Essential Biodiversity Variables on species traits.
Nature Ecology and Evolution, *2*(10), 1531–1540. doi: 10.1038/s41559-018-0667-3
Laughlin, D. C. (2018). Rugged fitness landscapes and Darwinian demons in trait-based ecology.
New Phytologist, *217*(2), 501–503. doi: 10.1111/nph.14908
Laughlin, D. C., Gremer, J. R., Adler, P. B., Mitchell, R. M., & Moore, M. M. (2020). The Net

Effect of Functional Traits on Fitness. *Trends in Ecology and Evolution*, *35*(11), 1037–1047.
doi: 10.1016/j.tree.2020.07.010
Levin, S. C., Crandall, R. M., Pokoski, T., Stein, C., & Knight, T. M. (2020). Phylogenetic and
functional distinctiveness explain alien plant population responses to competition: Phylogeny
and traits explain dominance. *Proceedings of the Royal Society B: Biological Sciences*,
287(1930). doi: 10.1098/rspb.2020.1070rspb20201070
Lintulaakso, K. (2013). *MammalBase—database of recent mammals*.
Madin, J. S., Anderson, K. D., Andreasen, M. H., Bridge, T. C. L., Cairns, S. D., Connolly, S. R.,
… Baird, A. H. (2016). The Coral Trait Database, a curated database of trait information for
coral species from the global oceans. *Scientific Data*, *4*, 170174. doi: 10.1038/sdata.2017.174
Maitner, B. S., Boyle, B., Casler, N., Condit, R., Donoghue, J., Durán, S. M., … Enquist, B. J.
(2018). The bien r package: A tool to access the Botanical Information and Ecology Network
(BIEN) database. *Methods in Ecology and Evolution*, *9*(2), 373–379. doi: 10.1111/2041-
210X.12861
Maldonado, C., Molina, C. I., Zizka, A., Persson, C., Taylor, C. M., Albán, J., … Antonelli, A.
(2015). Estimating species diversity and distribution in the era of Big Data: To what extent
can we trust public databases? *Global Ecology and Biogeography*, *24*(8), 973–984. doi:
10.1111/geb.12326
Marx, V. (2013). The big challenges of big data. *Nature*, *498*, 255–260.
Maurer, S. M., Firestone, R. B., & Scriver, C. R. (2000). Science's neglected legacy: Large,
sophisticated databases cannot be left to chance and improvisation. *Nature*, *405*(6783), 117–
120. doi: 10.1038/35012169
McElreath, R. (2021) *Statistical Rethinking.* London, UK: Chapman and Hall/CRC.
Messier, J., McGill, B. J., & Lechowicz, M. J. (2010). How do traits vary across ecological scales?
A case for trait-based ecology. *Ecology Letters*, *13*(7), 838–848. doi: 10.1111/j.1461-
0248.2010.01476.x
Metcalf, C. J. E., Graham, A. L., Martinez-Bakker, M., & Childs, D. Z. (2016). Opportunities and
challenges of Integral Projection Models for modelling host-parasite dynamics. *Journal of
Animal Ecology*, *85*(2), 343–355. doi: 10.1111/1365-2656.12456
Michener, W. K. (2006). Meta-information concepts for ecological data management. *Ecological
Informatics*, *1*(1), 3–7. doi: 10.1016/j.ecoinf.2005.08.004
Middleton, O., Svensson, H., Scharlemann, J. P. W., Faurby, S., & Sandom, C. (2021). CarniDIET
1.0: A database of terrestrial carnivorous mammal diets. *Global Ecology and Biogeography*,
30(6), 1175–1182. doi: 10.1111/geb.13296
Monnet, A. C., Cilleros, K., Médail, F., Albassatneh, M. C., Arroyo, J., Bacchetta, G., … Leriche,
A. (2021). WOODIV, a database of occurrences, functional traits, and phylogenetic data for
all Euro-Mediterranean trees. *Scientific Data*, *8*(1), 1–11. doi: 10.1038/s41597-021-00873-3
Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S., …
Menges, E. S. (2008). Longevity can buffer plant and animal populations against chnaging
climatic variability. *Ecology*, *89*(1), 19–25. Retrieved from
http://www.esajournals.org/doi/pdf/10.1890/07-0774.1
Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. K. M. (2015).
An amniote life-history database to perform comparative analyses with birds, mammals, and
reptiles. *Ecology*, *96*(11). doi: 10.1890/15-0846r.1
Nadrowski, K., Ratcliffe, S., Bönisch, G., Bruelheide, H., Kattge, J., Liu, X., … Wirth, C. (2013).
Harmonizing, annotating and sharing data in biodiversity-ecosystem functioning research.

Methods in Ecology and Evolution, *4*(2), 201–205. doi: 10.1111/2041-210x.12009

Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017). AmphiBIO, a global database for amphibian ecological traits. *Scientific Data*, *4*, 1–7. doi: 10.1038/sdata.2017.123

Ozgul, A., Childs, D. Z., Oli, M. K., Armitage, K. B., Blumstein, D. T., Olson, L. E., … Coulson, T. (2010). Coupled dynamics of body mass and population growth in response to environmental change. *Nature*, *466*(7305), 482–485. doi: 10.1038/nature09210

Paniw, M., Maag, N., Cozzi, G., Clutton-Brock, T., & Ozgul, A. (2019). Life history responses of meerkats to seasonal changes in extreme environments. *Science*, *363*(6427), 631–635. doi: 10.1126/science.aau5905

Pistón, N., de Bello, F., Dias, A. T. C., Götzenberger, L., Rosado, B. H. P., de Mattos, E. A., … Carmona, C. P. (2019). Multidimensional ecological analyses demonstrate how interactions between functional traits shape fitness and life history strategies. *Journal of Ecology*, *107*(5), 2317–2328. doi: 10.1111/1365-2745.13190

Poorter, L., Wright, S.J., Paz, H., Ackerly, D.D., Condit, R., Ibarra-Manriques, G. et al. (2008) Are functional traits good predictors of demographic rates? Evidence from five Neotropical forests. *Ecology*, 89, 1908–1920.

Reich, P. B., Wright, I. J., Cavender-Bares, J., Craine, J. M., Oleksyn, J., Westoby, M., & Walters, M. B. (2003). The evolution of plant functional variation: Traits, spectra, and strategies. *International Journal of Plant Sciences*, *164*(SUPPL. 3). doi: 10.1086/374368

Reichman, O. J., Jones, M. B., & Schildhauer, M. P. (2011). Challenges and opportunities of open data in ecology. *Science*, *331*(6018), 703–705. doi: 10.1126/science.1197962

Roper, M., Capdevila, P., & Salguero-Gómez, R. (2021). Senescence: Why and where selection gradients might not decline with age. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1955). doi: 10.1098/rspb.2021.0851

Rowcliffe, J. M., Jansen, P. A., Kays, R., Kranstauber, B., & Carbone, C. (2016). Wildlife speed cameras: measuring animal travel speed and day range using camera traps. *Remote Sensing in Ecology and Conservation*, *2*(2), 84–94. doi: 10.1002/rse2.17

Salguero-Gómez, R., Jackson, J., & Gascoigne, S.J.L. (2021). Four key challenges in the open data revolution. *Journal of Animal Ecology*, *90(9), 2000-2004*. doi: 10.1111/1365-2656.13567

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Bein, C., de Buhr, H., Farack, C., … Vaupel, J. W. (2016). COMADRE: A global data base of animal demography. *Journal of Animal Ecology*. doi: 10.1111/1365-2656.12482

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Buckley, Y. M., Che-Castaldo, J., Caswell, H., … Vaupel, J. W. (2015). The compadre Plant Matrix Database: An open online repository for plant demography. *Journal of Ecology*. doi: 10.1111/1365-2745.12334

Salguero-Gómez, R., & Laughlin, D. C. (2021). Not all traits are functional: the Panglossian paradigm. *Authorea*, (December 2021). doi: 10.22541/au.163940711.10447233/v1

Salguero-Gómez, R., Violle, C., Gimenez, O., & Childs, D. (2018). Delivering the promises of trait-based approaches to the needs of demographic approaches, and vice versa. *Functional Ecology*, *32*(6), 1424–1435. doi: 10.1111/1365-2435.13148

Santini, L., Isaac, N. J. B., & Ficetola, G. F. (2018). TetraDENSITY: A database of population density estimates in terrestrial vertebrates. *Global Ecology and Biogeography*, *27*(7), 787–791. doi: 10.1111/geb.12756

Schneider, F. D., Fichtmueller, D., Gossner, M. M., Güntsch, A., Jochum, M., König-Ries, B., … Simons, N. K. (2019). Towards an ecological trait-data standard. *Methods in Ecology and*

Evolution, *10*(12), 2006–2019. doi: 10.1111/2041-210X.13288

Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., … Wardle, D.
A. (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant
communities. *Ecology Letters*, *18*(12), 1406–1419. doi: 10.1111/ele.12508

Smallegange, I. M., Deere, J. A., & Coulson, T. (2014). Correlative changes in life-history
variables in response to environmental change in a model organism. *American Naturalist*,
183(6), 784–797. doi: 10.1086/675817

Stenseth, N. C., & Mysterud, A. (2002). Climate, changing phenology, and other life history traits:
Nonlinearity and match-mismatch to the environment. *Proceedings of the National Academy*
of Sciences of the United States of America, *99*(21), 13379–13381. doi:
10.1073/pnas.212519399

Swenson, N. G., Worthy, S. J., Eubanks, D., Iida, Y., Monks, L., Petprakob, K., … Zambrano, J.
(2020). A reframing of trait–demographic rate analyses for ecology and evolutionary biology.
International Journal of Plant Sciences, *181*(1), 33–43. doi: 10.1086/706189

Terry, J. C. D., O'Sullivan, J. D., & Rossberg, A. G. (2022). No pervasive relationship between
species size and local abundance trends. *Nature Ecology and Evolution*, *6*(2), 140–144. doi:
10.1038/s41559-021-01624-8

Troia, M. J., & McManamay, R. A. (2016). Filling in the GAPS: evaluating completeness and
coverage of open-access biodiversity databases in the United States. *Ecology and Evolution*,
6(14), 4654–4669. doi: 10.1002/ece3.2225

Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., … Messier,
J. (2012). The return of the variance: Intraspecific variability in community ecology. *Trends*
in Ecology & Evolution, 27, 244–252.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007).
Let the concept of trait be functional! *Oikos*, *116*(5), 882–892. doi: 10.1111/j.2007.0030-
1299.15559.x

Vitousek, M. N., Johnson, M. A., & Husak, J. F. (2018). Illuminating endocrine evolution: The
power and potential of large-scale comparative analyses. *Integrative and Comparative*
Biology, *58*(4), 712–719. doi: 10.1093/icb/icy098

Whitlock, M. C. (2011). Data archiving in ecology and evolution: Best practices. *Trends in*
Ecology and Evolution, *26*(2), 61–65. doi: 10.1016/j.tree.2010.11.006

Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., … Vieglais, D.
(2012). Darwin core: An evolving community-developed biodiversity data standard. *PLoS*
ONE, *7*(1). doi: 10.1371/journal.pone.0029715

Wilkes, M. A., Edwards, F., Jones, J. I., Murphy, J. F., England, J., Friberg, N., … Brown, L. E.
(2020). Trait-based ecology at large scales: Assessing functional trait correlations,
phylogenetic constraints and spatial variability using open data. *Global Change Biology*,
26(12), 7255–7267. doi: 10.1111/gcb.15344

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., … Mons,
B. (2016). Comment: The FAIR Guiding Principles for scientific data management and
stewardship. *Scientific Data*, *3*, 1–9. doi: 10.1038/sdata.2016.18

Williams, N. F., McRae, L., Freeman, R., Capdevila, P., & Clements, C. F. (2021). Scaling the
extinction vortex: Body size as a predictor of population dynamics close to extinction events.
Ecology and Evolution, *11*(11), 7069–7079. doi: 10.1002/ece3.7555

Yang, J., Cao, M., & Swenson, N. G. (2018). Why Functional Traits Do Not Predict Tree
Demographic Rates. *Trends in Ecology and Evolution*, *33*(5), 326–336. doi:

10.1016/j.tree.2018.03.003

Table 1: Variable names and meaning contained in the MOSAIC Database, organised into seven

general trait domains. A more detailed description can be found in the MOSAIC user guide found

at: https://mosaicdatabase.web.ox.ac.uk/.

Aspect	Variable	Description	Units
Taxonomy (Parallels COM(P)ADRE)	1 *Species Author*	Taxonomic species name as used by the author(s) in the publication. When more than one study exists for the same species, these are given sequential numeric suffixes (e.g. Ursus_americanus, Ursus_americanus_2, etc.)	NA
	2 *Species Accepted*	Currently accepted taxonomic name according to the Catalogue of Life (www.catalogueoflife.org). See the Supplementary Online Material S3 for an R script to check accepted and synonym names from SpeciesAuthor above	NA
	3 *Common Name*	English common name of SpeciesAccepted	NA
	4 *Family*	Taxonomic family of study species	NA
	5 *Order*	Taxonomic order of study species	NA
	6 *Class*	Taxonomic class of study species	NA
	7 *Phylum*	Taxonomic phylum of study species	NA
	8 *Kingdom*	Taxonomic Kingdom of study species	NA
Species information General	9 *Organism type*	General plant/algae type, based mainly on architectural organisation. The species was assigned to one of these possible values using the description of plant growth type provided by the author and other external sources (e.g. other publications)	Categorical/ Factor
	10 *DicotMonocot*	Whether species is a dicot or monocot	Categorical/ Factor
	11 *AngioGymno*	Whether species is an angiosperm or a gymnosperm	Categorical/ Factor
Source information	12 *Authors*	Surname (family name) of all author	NA
	13 *Journal*	The document from which data were sourced.	NA
	14 *YearPublication*	Year of publication	NA
	15 *DOI*	Digital Object Identifier number	NA
Traits	01 *Biomass*	Mean mass of an individual/whole-organism. (Plants) Plant mass is measured as aboveground dry mass. See "Further information" for additional information on belowground biomass.	Grams
	02 Height	(Plants) Mean height of the whole organism/whole individual from surface (i.e. substrate) to tallest vertical extremity. (Animals) Mean ventro-dorsal length. (i.e. for terrestrial quadrupeds, vertical distance from the ground to the top of the shoulder. For marine bony fishes, from the top of the high point of the dorsal fin to the base of the ventral ridge or ventral fin, whichever is a longer distance perpendicular to the anterior-posterior axis.)	Centimetres
	03 Growth Determination	Whether a species exhibits growth determinacy or not.	Categorical/ Factorial
	04 Regeneration	Capacity for an individual to regenerate any substantial part of its body, including autotomy. Autotomy is defined as "The voluntary severance by an animal of a part of its body (commonly one of its own limbs), usually to escape capture	Categorical/ Factorial

		by a predator that has seized that part. The part then regrows."	
	05 Sexual Dimorphism	An indicator of whether sexual dimorphism is exhibited in the species. Sexual dimorphism is defined as "the occurrence of morphological differences (other than primary sexual characters) that distinguish males from females of a species of organism." (Oxford Dictionaries of Ecology and Zoology)	Categorical/ Factorial
	06 Mating System	System of mating; the organisation of sexual interactions of individuals within populations based on sex.	Categorical/ Factorial
	07 Hermaphrodism	Indicator of whether a species exhibits hermaphrodism or monoeicieosity. Hermaphrodism is defined as: "An individual that possesses both male and female sex organs; i.e. it is bisexual." (Oxford Dictionary of Zoology). Monoeciousness is defined as: "Applied to an organism in which separate male and female organs occur on the same individual (e.g. to a plant which bears male and female reproductive structures in the same flower or separate male and female flowers on the same plant, or to a hermaphrodite animal). Some authors restrict the term botanically to plants with separate male and female flowers; plants which bear male and female reproductive organs in the same flower are then called hermaphrodite."	Categorical/ Factorial
	08 Sequential Hermaphrodism	Indicator of whether a species is protogynous or protandrous.	Categorical/ Factorial
	09 Dispersal Capacity	An indicator for whether or not a species exhibits dispersal behaviour or at any stage in its life cycle. Where dispersing, a categorical description of whether dispersal is natal or breeding or otherwise. Dispersal is defined as "The tendency of an organism to move away, either from its birth site (natal dispersal) or breeding site (breeding dispersal): the opposite of philopatry." (Oxford Dictionary of Zoology).	Categorical/ Factorial
	10 Type of Dispersal	An indication of whether dispersal is a passive (requires assistance) or active (no assistance) event. See DispClasses for more information	Categorical/ Factorial
	11 Mode of Dispersal	An indicator of the mode of dispersal of the species (plant and animal specific terminology). (*e.g.,* phoretic dispersal in *Daphnia magna*).	Categorical/ Factorial
	12 Dispersal Classes	Age- or stage-classes of the species that are capable of dispersal.	Categorical/ Factorial
	13 Volancy	An indicator of whether a species is volant or non-volant (i.e., able to fly or not).	Categorical/ Factorial
	14 Aquatic Habitat Dependency	An indicator of whether a species is dependent on water or not. (e.g., limnic, lentic, lotic)	Categorical/ Factorial

 Figures

Figure 1. Structure of the MOSAIC database (v1.0.0). The MOSAIC database comprises a

combination of existing trait records centralised from data servers, databases, and datasets and new

records collected from the primary literature by the MOSAIC team. Existing records are labelled,

MOSAIC-A, in the mosaic metadata attribute field and include provenance in the metadata. New

data records that adhere to existing data standards and that fit into the remit of an existing database

(*i.e.*, a data gap filled by the MOSAIC team) are labelled MOSAIC-B. New records that do not

have an existing data standard or for which there is not an existing database are labelled MOSAIC-

C.

Figure 2. Trait fields in the MOSAIC database (v1.0.0). The MOSAIC database contains a

combination of continuous and discrete trait fields. Six of the 14 trait fields are illustrated here for

animals, and organised by trait level (all discrete fields) and taxonomic classification. Trait

distributions vary strongly by trait and taxonomy. Observed variation in trait values by taxonomic

group is a prerequisite condition to their potential value toward explaining reported variation in

vital rates and resulting demographic properties across the Tree of Life. Species included in the

MOSAIC database are limited to those for which stage-structured population data exist in the

COMADRE, COMPADRE, and PADRINO databases.

Figure 3. The spatial distribution of MOSAIC (v1.0.0) records relative to the (a) COMADRE and

(b) COMPADRE databases. The COMADRE and COMPADRE dataset include animal and plant

demographic data, respectively, on all continents except Antarctica, with a substantial bias to

North America and Central Europe. MOSAIC introduces trait records that include models in all

major geographies of COMADRE and COMPADRE. The map shows the density of matrix

population models globally (color graded by density per area in *ca.* 150 km^2 hexagonal cells). Cells

bordered by white represent localities of population models for which there are trait records

included in the first release of the MOSAIC trait database. Species prioritised in v1.0.0 of

MOSAIC did not include the PADRINO dataset, so no distribution figure is shown for integral

projection models.

Figure 4. Phylogenetic mapping of traits included in MOSAIC (v1.0.0). (a) Phylogenetic

distribution of seven major traits for animals: (1) growth determination; (2) sexual dimorphism;

(3) monogamy; (4) gonochory; (5) volancy; (6) dispersal; and (7) biomass. (b) Phylogenetic

distribution of four traits in plants: (1) dispersal capability; (2) dispersal type; (3) dispersal mode;

and (4) dispersal class. Figures developed in R using the ggtree package.

Figure 5. Trait covariance in the MOSAIC database (v1.0.0). Some traits exhibit high correlation

in the MOSAIC database, such as between growth determination and between growth

regeneration. Trait associations are expected to occur in the MOSAIC dataset and may reflect

widespread constraints or statistical anomalies, particularly when dealing with small samples or

specific taxonomic subgroups. Trait covariation can be symptomatic of biomechanical constraints

(*e.g.*, flight and biomass), major growth characteristics (*e.g.*, modularity and growth

determination), or other past or presently compelled associations (*e.g.*, height and vessel density).

Figure 1

      Figure 3

(a) COMADRE

(b) COMPADRE

Figure 4

(a)

VOLANCY

VOLANT
SEMI-VOLANT
NON-VOLANT
NOT DIGITISED

SEQUENTIAL HERMAPHRODISM

PROTOGYNY

PROTANDRY

HERMAPHRODISM

GONOCHOROUS
HERMAPHRODITIC
HERMAPH + GONO
NOT FOUND

GROWTH DETERMINACY

DETERMINATE

INDETERMINATE

SEXUAL DIMORPHISM

DIMORPHIC

MONOMORPHIC

SEXUAL DIMORPHISM

MONOGAMOUS

NON-MONOGAMOUS

DISPERSAL CLASS

ADULT   EGG + LARVAL   JUVENILE   NOT FOUND

(b)

DISPERSAL CAPACITY

DISPERSING

NOT FOUND

DISPERSAL TYPE

NOT FOUND

PASSIVE

DISPERSAL CLASS

GAMETE + SPORE

SEED
SPORE
ZYGOTE

DISPERSAL MODE

ANEMOCHORY
(+SEMI-ANEMOCHORY)

AUTOCHORY
(+SEMI- AUTOCHORY)

HYDROCHORY
(+SEMI- HYDROCHORY)

880 **Figure 5**

883   MOSAIC: A Unified Trait Database to Complement Structured Population Models

885   Connor Bernard, Gabriel Silva Santos, Jacques Deere, Roberto Rodriguez-Caro, Pol Capdevila,

886   Erik Kusch, Samuel J L Gascoigne, John Jackson, & Roberto Salguero-Gómez

888   **SUPPLEMENTARY ONLINE MATERIALS**

889   Table of Contents

897   S1: MOSAIC User Guide

898   USER GUIDE TO THE MOSAIC LIFE HISTORY DATABASE
899
900   A Companion to COM(P)ADRE and PADRINO/A demographic databases

903   *Working document – 06 February 2022*

Table of Contents

User Guide Information

User Guide

Appendix

References

Table of Contents

Introduction

General Instructions

926 Format of the User Guide

930 **Metadata**

931 A. Species Name/Taxonomy

935 B. Study Information

942 **Primary User Guide/Variables**

943 * Applicable to plants

944 ** Applicable to animals

946 A. Morphometry & Growth

953 B. Reproductive traits

964   User guide version information
965
966   Version 1.0.0
967
968   Release date: 6 February 2021
969
970   Contact: mosaicdatabase@biology.ox.ac.uk

General Instructions

Database Organization
The data associated with MOSAIC are provided in a single R data (extension.Rdata) and as a comma separated value (.csv) file format. The code for downloading the data can be located from the mosaic portal (https://mosaicdatabase.web.ox.ac.uk/download-database) In addition, these files are accompanied by R scripts and a nexus phylogeny available in the Supplementary Information of the manuscript introducing MOSAIC, and in our GitHub repository (https://github.com/mosaicdatabase/mosaicdatabase)

MOSAIC_v_1_0_0.RData: Contains basic information regarding the source of publication, as well as ecological, biogeographic, and taxonomic details of the demographic study for each study species, the demographic information (i.e., the matrix population model) and metadata.

Database Design
In developing the MOSAIC database, we balance level of detail with accessibility. A highly detailed, comprehensive profile of life history traits for species in COM(P)ADRE and PADRINO/A, if possible to collect, would be difficult to navigate. So rather than collating a wealth of information in many different formats, we designed this dataset to highlight a smaller collection of traits in a single format which is of most interest, and expressed, by COM(P)ADRE and PADRINO/A users. In MOSAIC's future updates, we plan to add additional detail and additional fields, but we initially took a limited approach and plan to keep the design minimal. If there are life history traits, alternative formats for existing variables, or other features you would like to see added to MOSAIC, please suggest them to us at: mosaicdatabase@biology.ox.ac.uk

Understanding the diverse needs of users, we include in this guidance document additional direction on obtaining information on the variables we report in more detail. In this guide, we also highlight the scope of use, and caution against the most foreseeable abuses of data. We ask that all users approach this dataset with a caution and pay close attention to what variables and their statistical expression reflect.

The Meanings of NA and NDY in MOSAIC
NA in the MOSAIC data generally means that the data are not applicable. An example of where the data are not applicable is volancy within plants, as this trait does not occur in plants. NDY in the MOSAIC data means that the data have not yet been digitized. NF in the MOSAIC data means that the data are not available to date as no affirmative records were found upon review.

Disclaimer
The MOSAIC digitization team does its best to ensure data accuracy, and every piece of information goes through multiple error-checks prior to its release in www. mosaicdatabase.web.ox.ac.uk. However, we claim no responsibility for any damage that may arise from using MOSAIC. A list of error checks and potential issues in the use and

1016 interpretation of the database are described in the main manuscript. The end user is ultimately
1017 responsible for his/her interpretations of the data.

1024
1025
1026 **What is new in this version?**
1027
1028 Version 1.0.0
1029 ● The first version of the database. No updates.
1030
1031

1032
1033
1034

DIAGRAM OF THE MOSAIC DATABASE ARCHITECTURE

Variables in MOSAIC

The MOSAIC database is constructed of objects containing life history information across variables and organized into themes to aid in navigation. The metadata object is the object containing information about every study for which data is stored in the MOSAIC database. Every variable containing a value in the MOSAIC database will have corresponding metadata.

Associated with every data record is a value and its corresponding metadata. The metadata details data providence and relationships to existing databases. The fields within metadata are detailed below:

Format Guide

[Index]   *Variable Name*

Definition: [Definition of the variable]

Possible values, [cat. = categorical/discrete; cont. = continuous], [r variable class: character, numeric, integer, complex, or logical.]
- <XX>   [variable with two digits]
  XXX –XXX   [discrete variable name] – [discrete variable definition]
  <...>XXX<...>   [discrete variable with additional content on either side]
- Units: [unit of measure (integer, percent, ratio, mm, km^2, g, etc.)]
- Precision: [for continuous variables: scientific notation of the precision of the measure - e.g., $1e^1$ for km = precise to the tens of km (3270 km), no decimal; $1e^{-1}$ for temperature = precise to the tenth of a degree (10.4 °C)]
- Error boundaries: [for continuous variables: boundaries beyond which values are errors]

Usage Notes: [Notes on boundaries of use – note that this is non-exhaustive and highlights major potential errors]

Additional Information: [Guidance on databases containing more detailed information]

Source Data: [Datasets from which information was gathered; note that this will not include individual papers unless the papers are associated with a larger dataset/database. This category reflects databases.]

Last updated: [Date that information was retrieved from other databases or searches]

A1   *Species Accepted*

Definition: Currently accepted latin name.

Possible values, cat., factor
- <Genus_species> - e.g., Taraxacum_officinale

Usage Notes: NA

Additional Information: NA

Source Data: This information is obtained from The Encyclopaedia of Life

Last updated: 24 January 2022

1099
1100 **A2** *Kingdom*
1101
1102 **Definition:** Kingdom to which species belongs
1103
1104 **Possible values**, cat., factor
1105 ● <kingdom> - e.g., Plantae, Fungi, Rhodophyta, Chromista (yes, MOSAIC includes fungi
1106   and algae as well as plants)
1107
1108 **Usage Notes**: NA
1109
1110 **Additional Information:** NA
1111
1112 **Source Data:** The Encyclopaedia of Life
1113
1114 **Last updated**: 24 January 2022
1115

1116   **B1**   *Authors*
1117
1118   **Definition:** Surname (family name) of all authors
1119
1120   **Possible values**, NA, character
1121   ●   <name(s)> - Separated with ";" e.g., "Smith; Jones"
1122
1123   **Usage Notes**: NA
1124
1125   **Additional Information:** NA
1126
1127   **Source Data:** NA
1128
1129   **Last updated**: 24 January 2022
1130

1131   **B2**   *Journal Name*
1132
1133   **Definition:** The document from which data were sourced.
1134
Possible values, cat., factor
● <abbreviated journal name> - Where the data come from a scientific journal article, the
abbreviated journal name is given. We use the standard abbreviation of the journal
compliant with the ISO-4 standard.
● Book - Records are from a book, or book chapter
● PhD thesis - Records are from a doctoral thesis
● MSc thesis - Records are from a masters thesis
● Report - Records are from a report
● Conference talk - Records are from a conference talk
● Conference poster - Records are from a conference poster
1145
1146   **Usage Notes**: NA
1147
1148   **Additional Information:** NA
1149
1150   **Source Data:** NA
1151
1152   **Last updated**: 24 January 2022
1153

1154 **B3**   *Year Publication*

1155

1156 **Definition:** Year of publication

1157

1158 **Possible values**, cont., numerical

1159   ● <yyyy> - e.g., 2012

1160

1161 **Usage Notes**: NA

1162

1163 **Additional Information:** NA

1164

1165 **Source Data:** NA

1166

1167 **Last updated**: 24 January 2022

1168

1169   **B4**   *DOI/ISBN Number*
1170
1171   **Definition:** Digital Object Identifier number
1172
1173   **Possible values**, NA, character
1174   ● <XXXXXXXXXXXXXX> - e.g., doi.org/10.1073/pnas.1506215112
1175
1176   **Usage Notes**: NA
1177
1178   **Additional Information:** NA
1179
1180   **Source Data:** NA
1181
1182   **Last updated**: 24 January 2022
1183

1184   **PRIMARY Variables**
1185

1186 **A1** *Biomass*
1187
1188
1189 **Definition:** Maximum reported mass of adult individual/whole-organism. For plants, only
1190 aboveground dry mass is measured. See "Additional information" for additional information on
1191 belowground biomass.
1192
Possible values, cont., numerical
- Units: Grams (g)
- Precision: 0.000
- Error Boundaries: 0-150,000,000g
- NF – Body mass reviewed and inconclusive (no affirmative records found upon review)
- NDY – Body mass not digitised yet (yet to be evaluated)
- NA – not applicable
1200
1201 **Usage notes:** When both male and female data were reported, only the maximum value was
1202 considered independently of the gender of the individuals.
1203
1204 **Additional Information:** Belowground biomass is not reported because information availability
1205 is appreciably more limited than for aboveground biomass. The BIEN database includes
1206 information on belowground biomass that can be referenced and utilized where of interest.
1207
1208 **Source Data:** Amniote, TRY database.
1209
1210 **Last updated**: 24 January 2022
1211

A2   *Height*

Call: -Morph$Height

Definition: (Plants) Maximum height of the whole organism/whole individual from surface (i.e. substrate) to tallest vertical extremity.

Possible values, cont., numerical
- Units: centimeters (cm)
- Precision: 0.000
- Error Boundaries: 0-1000cm

Usage notes:
- Depending on the species group, height can have profoundly different meanings. Height corresponds with embolic risk in some woody plants and corresponds with size-based fitness in others.

Additional Information: NA

Source Data: TRY database.

Last updated: 24 January 2022

1237 **A3** *Growth Determination*
1238

1239 **Call:** -Morph$Growth$Determination
1240

1241 **Definition:** Growth indeterminacy is defined by continuous growth of individuals throughout
1242 their lifetimes (measured by mass, length, bone ossification, or other indicators). This field
1243 reflects a binary classification of whether an individual is growth (in)determinate.
1244

Possible values, cat, factor
● Growth indeterminate – growth continues throughout an individual's lifespan
● Growth determinate – growth ceases or attenuates to negligibility before the end of an
individual's lifespan
● NF - Growth determination reviewed and inconclusive (no affirmative records found
upon review)
● NDY - Growth determination not digitised yet (yet to be evaluated)
● NA – Growth determination not applicable to the subject area
1253

Usage notes: Additional classification systems for characterizing growth indeterminacy exist in
the literature. Most well recognised is a six-type scheme describing growth and determination by
Sebens (1987), which offers detailed characterization of growth patterns. Broad-scale
information about growth and age for most species could not be located in the literature, and
therefore a simplified schema is used. More resolved classifications might be incorporated into
future versions of MOSAIC.
1260

1261 **Additional Information:** NA
1262

1263 **Source Data:** 24 January 2022
1264

1265 **Last updated**: NA
1266

1267 **A4**   *Regeneration*
1268
1269
1270 **Call:** **-**Morphology$Growth$Regeneration
1271
1272 **Definition:** Capacity for an individual to regenerate any substantial part of its body, including
1273 autotomy. Autotomy is defined as "The voluntary severance by an animal of a part of its body
1274 (commonly one of its own limbs), usually to escape capture by a predator that has seized that
1275 part. The part then regrows."
1276
Possible values, cat., factor
● Regenerative – individuals are capable of regenerating
● Non-regenerative – individuals do not exhibit the capacity to regenerate tissues
● NF - regenerative abilities reviewed and inconclusive (no affirmative records found upon
review)
● NDY - regeneration information not digitised yet (yet to be evaluated)
● NA – regenerative abilities not applicable to the subject area
1284
1285 **Usage notes:** There are a number of more resolved schemes for detailing whether regenerating
1286 different parts of the body. For particular questions pertaining to the nature of injury, the level of
1287 recovery, the role of depredation, and the consequences for reproduction, more detail may be
1288 appropriate. This dataset covers the most general applications and initially screens for
1289 regenerative capability (from poor recovery of appendages to complete regeneration of limbs).
1290
1291 **Additional Information:** NA
1292
1293 **Source Data:** NA
1294
1295 **Last updated**: 24 January 2022
1296

1297 **A5** *Sexual Dimorphism*
1298
1299
1300 **Call:** -Morph$Dimorph
1301
1302 **Definition:** An indicator of whether sexual dimorphism is exhibited in the species. Sexual
1303 dimorphism is defined as "the occurrence of morphological differences (other than primary
1304 sexual characters) that distinguish males from females of a species of organism." (Oxford
1305 Dictionaries of Ecology and Zoology)
1306
Possible values, cat., factor
● Sexually Dimorphic – species is sexually dimorphic
● Sexually Monomorphic – species is sexually monomorphic (i.e. non-dimorphic)
● NF - Sexual dimorphism reviewed and inconclusive (no affirmative records found upon
review)
● NDY - Sexual dimorphism not digitised yet (yet to be evaluated)
● NA – Sexual dimorphism not applicable to the subject area
1314
1315 **Usage Notes:** NA
1316
1317 **Additional Information:**
1318
1319 **Source Data:** NA
1320
1321 **Last updated**: 24 January 2022
1322
1323

1324 **B1**   *Mating system*
1325
1326
1327 **Call:**   **-**Reproduction$MatingSystem
1328
1329 **Definition:** System of mating; the organization of sexual interactions of individuals within
1330 populations based on sex.
1331
Possible values, cat., factor
● Monogamy – exclusive mating between one male and one female
● Non-monogamy - Non-monogamy was assigned based on genetic or behavioural
evidence
● NF - Mating system reviewed and inconclusive (no affirmative records found upon
review)
● NDY - Mating system not digitised yet (yet to be evaluated)
● NA – Mating system not applicable to the subject area
1340
1341 **Usage notes:**
1342   ● Metric does not identify size of groups for plural mating systems
1343   ● Metric does not identify enforcement mechanisms for different mating systems
1344
1345 **Additional Information:** NA
1346
1347 **Source Data:** NA
1348
1349 **Last updated**: 24 January 2022
1350
1351
1352
1353

1354 **B2** *Sexual allocation*
1355
1356
1357 **Call:** -Reproduction$Allocation
1358
Definition: Indicator of whether a species exhibits hermaphroditism or monoeicieosity.
Hermaphrodism is defined as: "An individual that possesses both male and female sex organs;
i.e. it is bisexual." (Oxford Dictionary of Zoology). Monoeciousness is defined as: "Applied to
an organism in which separate male and female organs occur on the same individual (e.g. to a
plant which bears male and female reproductive structures in the same flower or separate male
and female flowers on the same plant, or to a hermaphrodite animal). Some authors restrict the
term botanically to plants with separate male and female flowers; plants which bear male and
female reproductive organs in the same flower are then called hermaphrodite."
1367
Possible values, cat., factor
● Hermaphroditic – species is hermaphroditic
● Monoecious – species is monoecious
● Dioecious or Gonochorous – species is dioecious (Gonochorous was adopted for animals)
● NF - Hermaphroditism reviewed and inconclusive (no affirmative records found upon
review)
● NDY - Hermaphroditism not digitised yet (yet to be evaluated)
● NA – Hermaphroditism is not applicable to the subject area
1376
1377 **Usage notes:**
1378   ●
1379
1380 **Additional Information:** NA
1381
1382 **Source Data:** NA
1383
1384 **Last updated**: 24 January 2022
1385
1386
1387
1388

1389 **B3** *Sequesntial hermaphroditism*
1390
1391
1392 **Call:** Reproduction$SeqHermaph
1393
1394 **Definition:** Indicator of whether there is a sex switch during the organism's lifespan.
1395
Possible values, cat., factor
● Protogynous – species is protogynous: organisms that are female and at some point in
their lifespan change sex to male.
● Protandrous – species is protandrous: organisms that are male and at some point in their
lifespan change sex to male.
● NF - Protogyny/Protandry reviewed and inconclusive (no affirmative records found upon
review)
● NDY - Protogyny/Protandry not digitised yet (yet to be evaluated)
● NA – Protogyny/Protandry is not applicable to the subject area.
1405
1406 **Usage notes:** NA
1407
1408 **Additional Information:** Note that is not rare flowers present protogyny/protandry but this was
1409 not considered in Mosaic so far
1410
1411 **Source Data:** NA
1412
1413 **Last updated**: 24 January 2022
1414

1415 **C1**   *Dispersal Capability*
1416
1417
1418 **Call:**   -Movement$Dispersal
1419
1420 **Definition:** An indicator for whether or not a species exhibits dispersal behaviour or at any stage
1421 in its life cycle. Where dispersing, a categorical description of whether dispersal is natal or
1422 breeding or otherwise. Dispersal is defined as "The tendency of an organism to move away,
1423 either from its birth site (natal dispersal) or breeding site (breeding dispersal): the opposite of
1424 philopatry." (Oxford Dictionary of Zoology).
1425
Possible values: cat., factor
● Dispersing – Exhibits at least one age-/stage-class which disperses; natal or breeding
components unknown.
● Natal Dispersal – Permanent dispersal of at least one age-/stage-class
● Breeding Dispersal – Dispersal of adults between breeding attempts in at least one age-
/stage-class
● Multi-Dispersal – Both natal and breeding dispersal reported in the species; see
DispClasses for more information
● Non-Dispersing – Species observed to have no dispersal traits/behaviour
● NF - dispersal capability reviewed and inconclusive (no affirmative records found upon
review)
● NDY – dispersal capability unknown/not yet evaluated
● NA – not applicable
1439
1440 **Usage Notes:** NA
1441
1442 **Additional Information:** NA
1443
1444 **Source Data:** NA
1445
1446 **Last updated**: 24 January 2022
1447

1448   **C2**   *Type of Dispersal*
1449
1450
1451   **Call:** -Movement$TypeDisp
1452
1453   **Definition:** An indication of whether dispersal is a passive (requires assistance) or active (no
1454   assistance) event. See DispClasses for more information.
1455
Possible values: cat., factor
● Active – organism utilises its own morphology for the dispersal event
● Passive – organism is unable to disperse through their own means and require an external
factor
● Active and Passive – organism is able to disperse with assistance but can also use
external factors. Active and passive dispersal can occur within the same life stage or can
occur in different life stages.
● NF – type of dispersal reviewed and inconclusive (no affirmative records found upon
review)
● NDY – type of dispersal unknown/not yet evaluated
● NA – not applicable
1467
1468   **Usage Notes:** NA
1469
1470   **Additional Information:** NA
1471
1472   **Source Data:** NA
1473
1474   **Last updated**: 24 January 2022
1475
1476

1477	**C3**	*Mode of Dispersal*
1478		
1479		
1480	**Call:**	-Movement$ModeDisp

1481

Definition: An indicator of the mode of dispersal of the species (plant and animal specific terminology).

1484

Possible values: cat., factor
- Motile – the dispersal of animal species without assistance
- Phoretic – the dispersal of animal species by attaching to another animal
- Water currents – the dispersal of animal species by water
- Motile and water currents – animal species that disperse without assistance and by water, both forms of dispersal can occur within the same life stage or can occur in different life stages
- Anemochory – the dispersal of plant seeds by wind
- Anthropochory – the dispersal of plant seeds by humans
- Autochory – the dispersal of plant seeds without assistance from an external vector (e.g., by gravity or ballistic dispersal)
- Hydrochory – the dispersal of plant seeds by water
- Zoochory – the dispersal of plant seeds by animals
- Anemochory and Anthropochory – plant seeds can be dispersed by wind and humans
- Anemochory and Autochory – plant seeds can be dispersed by wind and without the help of an external vector
- Anemochory and Hydrochory – plant seeds can be dispersed by wind and water
- Anemochory and Zoochory – plant seeds can be dispersed by wind and animals
- Autochory and Hydrochory – plant seeds can be dispersed without the help of an external vector and by water
- Autochory and Zoochory – plant seeds can be dispersed without the help of an external vector and by animals
- Hydrochory and Zoochory – plant seeds can be dispersed by water and animals
- Autochory, Anthropochory and Zoochory – plant seeds can be dispersed without the help of an external vector, by humans and by animals
- NF – mode of dispersal reviewed and inconclusive (no affirmative records found upon review)
- NDY – mode of dispersal unknown/not yet evaluated
- NA – not applicable

1514

Usage Notes: Plant seed dispersal modes can be subdivided into further categories, however we collated lower order categories into the higher order categories identified here

1517

Additional Information: NA

1519

Source Data: NA

1521

Last updated: 24 January 2022

1523
1524

C4   *Dispersal Class*

Call: -Movement$DispClass

Definition: Age- or stage-classes of the species that are capable of dispersal.

Possible values: cat., factor
- Adult – dispersal stage is an individual that has reached maturity, we include sub-adults into this category
- Egg – dispersal stage is a vessel within which an embryo develops and is expelled by an adult allowing for dispersal
- Fertile material – dispersal stage is a part of an individual, or in some cases a complete individual, that contains fertile material (e.g., the alga *Fucus vesiculosus*; detached floating material/individual that contains gametes)
- Gamete – dispersal stage is the reproductive cell not within a vessel
- Juvenile – dispersal stage is an individual that has not reached maturity
- Larval – dispersal in a specific juvenile stage restricted to non-mammal species, species can have multiple larval stages
- Seed – dispersal stage is fertilized, specific to plant species and, in our definition, references seeds and/or fruits that are dispersed
- Sperm – dispersal stage is the male gamete
- Spore – dispersal stage is a single cell that only contains half of the chromosome of the adult, can produce eggs or sperm
- Sporophyte – dispersal stage is a nonsexual phase of a species producing two diploid spores
- Zoospore – dispersal stage is a motile asexual spore
- Zygote – dispersal stage is a fused male and female gamete
- Adult and Juvenile – dispersal stage can be both the adult and juvenile stage
- Egg and Larval – dispersal stage can be both the egg and larval stage
- Gamete and Spore – dispersal stage can be both the gamete and spore
- Zoospore, Sperm and Sporophyte – dispersal stage can be a zoospore, sperm or sporophyte
- NF – dispersal class reviewed and inconclusive (no affirmative records found upon review)
- NDY – dispersal class unknown/not yet evaluated
- NA – not applicable

Usage Notes: Dispersal can occur in more than one age- or stage- class, where this occurs it is noted as such within the database

Additional Information: NA

Source Data: NA

Last updated: 24 January 2022

1569 **C5** *Volancy*
1570
1571
1572 **Call:** -Movement$Volancy
1573
1574 **Definition:** An indicator of whether a species is volant or non-volant (i.e., able to fly or not).
1575
Possible values, cat., factor
- Volant – the species is volant
(Most Birds (Class Aves), all Bats (Order Chiroptera), and some invertebrate species)
- Non-volant – the species is non-volant
- Semi-volant – the species has gliding abilities
(e.g., Gliding lizards (*Draco* spp.); flying squirrels such as the Northern flying squirrel
(*Glaucomys sabrinus*); flying fish (Exocoetidae); gliding frogs such as Wallace's flying
frog (*Rhacophorus nigropalmatus*); and gliding ants such as *Cephalotes atratus*).
- NF – volancy reviewed and inconclusive (no affirmative records found upon review)
- NDY – volancy unknown/not yet evaluated
- NA – not applicable
1587
1588 **Usage Notes:** NA
1589
1590 **Additional Information:** NA
1591
1592 **Source Data:** NA
1593
1594 **Last updated**: 24 January 2022

1595   S2: MOSAIC Trait Primary Search Key Words

1596   MOSAIC: Keywords used in primary literature search

1597
1598   Taxonomic names (binomial nomenclature) were used in connection with the field-specific terms
1599   in the search of primary literature in identifying the records of interest. The below include names
1600   that were searched in review of the primary literature.

1601
1602
1603
1604   Biomass—
1605   NA (database only)

1606
1607   Height—
1608   NA (database only)

1609
1610   Growth determination—
1611   Sources: *Web of Science; Google Scholar; Scopus*
1612   Terms: Growth determinat*, growth determination, growth determinate, growth
1613   indetermination, growth

1614
1615   Regeneration—
1616   Sources: *Web of Science; Google Scholar; Scopus*
1617   Terms: Regenerat*, regeneration, regenerate, rejuvenation

1618
1619   Sexual dimorphism—
1620   Sources: *Web of Science; Google Scholar; Scopus*
1621   Terms: Dimorphic

1622
1623   Mating system—
1624   Sources: *Web of Science; Google Scholar; Scopus*
1625   Terms: Monogamy, monogamous, polygyny, polygynous, polyandry, polyandrous,
1626   mating system

1627
1628   Hermaphrodism—
1629   Mating system—
1630   Sources: *Web of Science; Google Scholar; Scopus*
1631   Terms: Hermaphrod*, hermaphroditic, hermaphrodism, hermaphrodite, gonochoric,
1632   gynochorous, monoecious, dioecious, sexual differentiation,

1633
1634   Sequential hermaphrodism—
1635   Sources: *Web of Science; Google Scholar; Scopus*
1636   Terms: Sequential hermaphrodism, protandry, protogyny, protogynous hermaphroditism,
1637   protandrous hermaphroditism

1638
1639   Dispersal capacity—

1640   Sources: *Web of Science; Google Scholar; Scopus*
1641   Terms: Dispers*, dispersal, dispersing, dispersal capacity, dispersal capability
1642
1643 Type of dispersal—
1644   Sources: *Web of Science; Google Scholar; Scopus*
1645   Terms: Dispers*, active, passive
1646
1647 Mode of dispersal—
1648   Sources: *Web of Science; Google Scholar; Scopus*
1649   Terms: Dispers*, motile, phoretic, currents, anemochory, anthropochory, autochory,
1650   hydrochory, zoochory, anthropochory
1651
1652 Dispersal classes—
1653   Sources: *Web of Science; Google Scholar; Scopus*
1654   Terms: Dispers*, adult, jouvenile, egg, fertile, seed, sperm, spore, sporophyte, zoospore,
1655   zygote, gamete
1656
1657 Volancy—
1658   Sources: *Web of Science; Google Scholar; Scopus*
1659   Terms: Volant, non-volant, flight, flying, flightless, non-flying, winged, glide, gliding,
1660   ground
1661
1662 Aquatic habitat dependency—
1663   Sources: *Web of Science; Google Scholar; Scopus*
1664   Terms: Anadromous, catandromous, estuarine, brackish, lotic, lentic, lemnitic, littoral,
1665   palagic, marine, freshwater, saltwater, sea, ocean.
1666
1667

1668   S3. Databases Searched

1669
1670   MOSAIC: Databases Reviewed to Date for the MOSAIC Database

1671
1672   Type. Database Name. DOIs.

1673
1674   1. **Plants**. Botanical Information and Ecology Network (BIEN) (doi:
1675   10.7287/peerj.preprints.2615v2)
1676   2. **Plants**. TRY: Global Plant Trait Database (TRY) (doi: 10.1111/j.1365-
1677   2486.2011.02451.x)
1678   3. **Phylogeny.** Open Tree of Life: A synthesis of phylogeny and taxonomy into a
1679   comprehensive tree of life (doi: 10.1073/pnas.1423041112)
1680   4. **Vertebrates**. Amniote: An amniote life history database to perform comparative analysis
1681   with birds, mammals, and reptiles (doi: 10.1890/15-0846R.1)
1682   5. **Amphibians**. AmphiBIO: a global database for amphibian ecological traits. (doi:L
1683   10.1038/sdata.2017.123)
1684   6. **Mammals**. PanTHERIA: a species-level database of life history, ecology, and geography
1685   of extant and recently extinct mammals (doi: 10.1890/08-1494.1)
1686   7. **Animals**. AnAge Database of Animal Ageing and Longevity. (doi: 10.1111/j.1474-
1687   9726.2008.00442.x)
1688   8. **General (Sex)**. Tree of Sex: A database of sexual systems (10.1038/sdata.2014.15)
1689   9. **General (Variable)**. Open Trait Network Databases (https://opentraits.org/)
1690

1691 S4: Vignettes

1692

1693   Navigating MOSAIC

1694   Vignette #1 - Navigating MOSAIC

1695

1696

Welcome to the **MOSAIC database,** a database of functional traits for
comparative demography. The database, user, guide, and additional information
can be found and http://mosaicdatabase.web.ox.ac.uk.

MOSAIC is a database that aggregates existing databases and adds new records
for functional traits that currently do not have a database established. In
this vignette, we will show you how to download the dataset, search records,
and relate MOSAIC records with COMADRE, COMPADRE, and PADRINO databases.

Optional clearance of working space.

```
1705  rm(list = ls()) # Clear your environment
1706  if(!is.null(dev.list())) dev.off() # Clear plots/graphics
1707  cat("\014") # Clear console
```

Downloading MOSAIC

MOSAIC can be downloaded as an S4 object by running the below code in R. S4
data objects in R are an object oriented system in the R language that allow
control of constituent data fields Similar to S3 objects (which use the "$"
operator). S4 are comprised of objects that can be searched with the "@"
operator or slots, discussed in more detail below.

```
1714  library(devtools) # Compulsory package to pull down packages from GitHub. Ins
1715  tall if necessary.
1716  install_github("mosaicdatabase/Rmosaic")
1717  source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
1718  ain/mosaic_fetch.R") # Link to GitHub repo
1719  mosaic <- mos_fetch("v1.0.0") # Download version 1.0.0 (active version Feb 20
1720  22)
1721  library(Rmosaic)
```

Basics of manually navigating MOSAIC

Mosaic traits can be searched using the "@" operator. Attribute names
searched this way are analogous to the columns of a dataframe in a relational
database structure.

Once downloaded, you should be able to type statements
mosaicdatabase@[insertfield] (where [insertfield] is a particular trait). If
you are working in Rstudio, after the "@" a drop-down of the slots (traits)
should autopopulate.

```
1730  # Three examples of querying traits (easiest for navigation)
1731  mosaic@biomass
1732  mosaic@height
1733  mosaic@volancy
```

1734 The species corresponding with each index can be queried by prompting:

```
1735  # Three examples of querying traits (easiest for navigation)
1736  mosaic@species
```

1737 Data in mosaic can also be access using slots. Slots are the recommended mode
1738 of searching the database – though it has the disadvantage of not enabling
1739 the autopopulation of the attributes contained in the database (traits must
1740 be spelled out manually).

```
1741  # You can also search these by slot (recommended)
1742  slot(mosaic, "biomass")
1743  slot(mosaic, "height")
1744  slot(mosaic, "volancy")
```

Within each trait object, there are eight fields in mosaic. The first of
field is called "values" and contains the data. Values are unitless values
(either numeric or factorial) that are reported in units described in
the **Mosaic User Guide** http://mosaicdatabase.web.ox.ac.uk/user-guide. The
metadata is organised into additional attributes, reflecting the individual
elements of the metadata for a given record, including the authors, journal,
year of publication, databases from which data are sourced (if applicable)

mosaic@metaTaxa maps the complete taxanomic classification structure of a
species – from Kingdom to species for taxonomic clustering.

The other six attributes – "author", "year", "journal,"doi","database",
and"mosaic" – are metadata corresponding with each value record. For
instance:

```
1757  mosaic@species[[2]] # For this species, let us look at volancy (flight capaci
1758  ty) value
1759  mosaic@volancy@value[[2]] # or, equivalently:
1760  slot(slot(mosaic, "volancy"), "value")[[2]]
```

1761 Corresponds with the following metadata

```
1762  mosaic@volancy@author[[2]] # for "Acinonyx jubatus"
```

```
slot(slot(mosaic, "volancy"), "author")[[2]] # Author of the source publicati
on
```

```
slot(slot(mosaic, "volancy"), "year")[[2]] # And year of the source publicati
on
```

```
slot(slot(mosaic, "volancy"), "journal")[[2]] # The journal of the source pub
lication
```

```
#etc.
```

Using MOSAIC functions to quickly access files

A series of convenience functions can be sourced from the MOSAIC GitHub page
to facilitate navigating and working with the mosaic database that can be
accessed by running the follwing script.

```
source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
ain/navMosaic_46.R")
```

Below we highlight some of the basic queries for which the mosaic functions
can assist.

Is a species included in Mosaic?

```
spp_check("Fritillaria biflora")
## [1] TRUE
spp_check("Pagophilus groenlandicus")
## [1] FALSE
```

Can I see all records for a given trait?

```
traitAllSpp("biomass") # Only the first five records are shown for space
## [1] "NDY"    "50578" "ND"     "52500" "351000" "62000"
```

Can I see an overview of all records for a given species?

```
singSppTraitSummary("Aepyceros melampus")
##   biomass height   growthdet regen   dimorph       matsyst     hermaph se
qherm
## 1   52500   NDY Determinate   NDY Dimorphic Non-monogamous Gonochorous
NDY
##           dispcap disptype modedisp dispclass   volancy
## 1 Natal Dispersal   Active   Motile    Adult Non-volant
```

```
##                                    aquadep
## 1 Terrestrial, Water Habitat Independent
```

Can I see all records for more than one species?

```
sppAllTrait(c("Acinonyx jubatus", # you can also pass lists or dataframes to
this command
               "Acropora downingi",
               "Aepyceros melampus",
               "Alces alces",
               "Alligator mississippiensis"))
##                     sppnames biomass height    growthdet      regen   dim
orph
## 1          Acinonyx jubatus   50578    NDY Determinate        NDY Dimor
phic
## 2          Acropora downingi     ND    NDY         NDY Regenerative
NDY
## 3        Aepyceros melampus   52500    NDY Determinate        NDY Dimor
phic
## 4               Alces alces  351000    NDY Determinate        NDY Dimor
phic
## 5 Alligator mississippiensis   62000    NDY Determinate Regenerative
NDY
##           matsyst       hermaph seqherm         dispcap disptype      mod
edisp
## 1 Non-monogamous    Gonochorous    NDY Natal Dispersal   Active        M
otile
## 2           NDY Hermaphroditic    NDY Natal Dispersal  Passive Water cur
rents
## 3 Non-monogamous    Gonochorous    NDY Natal Dispersal   Active        M
otile
## 4 Non-monogamous    Gonochorous    NDY Natal Dispersal   Active        M
otile
## 5 Non-monogamous    Gonochorous    NDY Natal Dispersal   Active        M
otile
##           dispclass    volancy                                      aqua
dep
## 1      Juvenile Non-volant     Terrestrial, Water Habitat Independ
ent
## 2      Juvenile Non-volant                                         Mar
ine
```

```
## 3                Adult Non-volant        Terrestrial, Water Habitat Independ
ent
## 4          Juvenile Non-volant Terrestrial, Facultative Freshwater Depend
ent
## 5 Adult and Juvenile Non-volant  Terrestrial, Obligative Freshwater Depend
ent
```

Can I get a breakdown of counts/frequency of trait values?

```
traitFrequency("volancy")
##              counts  freq
## NDY            1329    NA
## Non-volant       82 0.882
## Semi-volant       1 0.011
## Volant           10 0.108
traitFrequency("growthdet")
##              counts  freq
## Determinate      31 0.596
## Indeterminate    21 0.404
## NDY            1370    NA
traitFrequency("hermaph")
##                            counts  freq
## Dioecious                      24 0.108
## Gonochorous                   106 0.475
## Hermaphroditic                 72 0.323
## Hermaphroditic & Gonochorous    1 0.004
## Monoecious                     20 0.090
## NDY                          1199    NA
```

Can I get all metadata for one or more traits?

```
metadata("volancy", 14)
##             author year                 journal database mosaic
## 1 Campos, Z. et al. 2006 The Herpetological Journal     NDY    NDY
multiMetaRecords("volancy", c(14:20))
##                              author year
```

```
## 1                    Ekerna, L. S. & Cords, M. 2007
## 2                       Zimmerman, S. J. et al. 2019
## 3 Jack, K. M., Sheller, C. & Fedigan, L. M. 2012
## 4                                         DFO 2013
## 5                    Torres, R. T. et al. 2017
## 6                                      NDY  NDY
## 7                    Campos, Z. et al. 2006
##
journal
## 1                                              A
nimal Behaviour
## 2
The Condor
## 3                              American Journal
of Primatology
## 4 Canadian Science Advisory Secretariat Central and Arctic Region Science
Advisory Report
## 5
Oryx
## 6
NDY
## 7                                      The Herpeto
logical Journal
##   database mosaic
## 1      NDY    NDY
## 2      NDY    NDY
## 3      NDY    NDY
## 4      NDY    NDY
## 5      NDY    NDY
## 6      NDY    NDY
## 7      NDY    NDY
```

1898

1899   http://mosaicdatabase.web.ox.ac.uk

1900   Updated 14 March 2022

1901   Vignette #2 – Tidyverse + MOSAIC

1902
1903   MOSAIC was built for use and integration with other datasets for
1904   comparative demography. Demographers are often interested in how traits
1905   scale across different taxonomic levels. The
1906   Tidyverse https://www.tidyverse.org/ collection of R packages (Dplyr in
1907   particular) are commonly used by ecologists to simplify workflows that
1908   subset and manipulate data. Here, we walk throught the basics of how Tidy
1909   tools can be use with the MOSAIC database to quickly isolate specific
1910   groups of organism or classes of records.

```
1911    library(tidyverse)
```

Accessing MOSAIC

1913   Download MOSAIC from the mosaic portal. For more information on the basics
1914   of downloading MOSAIC and navigating the data structure, see: Vignette #1:

```
1915    remotes::install_github("mosaicdatabase/Rmosaic")
1916    library(Rmosaic)
1917    mosaic <- mos_fetch("v1.0.0")
```

Using Tidyverse with MOSAIC

1919   Tidyverse allows for the easily filtering of outliers and NAs, and the
1920   isolation of records to particular groups of taxa in MOSAIC. For example,
1921   one can isolate the differences in biomass between birds and mammals and
1922   see how the decomposition of variance in biomass between the groups
1923   informs the representation of biomass values in MOSAIC

```
1924    # Bind the ID field, species, taxa, and biomass fields into a unified datafra
1925    me
1926
1927    biomass_table <- cbind(mosaic@mosaicID,
1928                           mosaic@species,
1929                           mosaic@taxaMetadat,
```

```
1930                                mosaic@biomass@value)

1931

1932   # Relabel the biomass field

1933

1934   names(biomass_table)[length(biomass_table)] <- "biomass"

1935

1936   # Coerce the field into a numeric values – transforms "NDY" into "NA"

1937

1938   biomass_table$biomass <- as.numeric(biomass_table$biomass)

1939

1940   # Removing an outlier and filtering out NAs

1941

1942   biomass_all <- biomass_table %>%
1943     drop_na(biomass) %>%
1944     filter(biomass < 12000000)

1945

1946   # Filter to mammals

1947

1948   biomass_mamm <- biomass_table %>%
1949     filter(Class == "Mammalia") %>%
1950     drop_na(biomass) # Remove results that are NA

1951

1952   # Filter to birds

1953

1954   biomass_aves <- biomass_table %>%
1955     filter(Class == "Aves") %>%
1956     drop_na(biomass)

1957

1958   # Plot all biomass

1959

1960   plot(biomass_all$`mosaic@mosaicID`, log(biomass_all$biomass), pch=16,
1961       xlab = "MOSAIC ID index", ylab = "Log Biomass", main = "Biomass")

1962

1963   # Highlight points for mammals
```

```
1964
1965   points(biomass_mamm$`mosaic@mosaicID`, log(biomass_mamm$biomass), pch=16, col
1966   ="red")
1967
1968   legend(2085, 16, legend=c("Mammals", "All Classes"),
1969          col=c("red", "black"), lty=1, cex=0.8)
```

```
1970
1971   # Generate histograms for all species on the same binning scale
1972
1973   Bin_all <- hist(log(biomass_all$biomass),
```

```
1974            breaks=seq(min(log(biomass_all$biomass)), max(log(biomass_all
1975   $biomass)),
1976                      length=15), main="Biomass (All Species)")
```

Biomass (All Species)

```
1977

1978   Bin_mamm <- hist(log(biomass_mamm$biomass),
1979            breaks=seq(min(log(biomass_all$biomass)), max(log(biomass_all
1980   $biomass)),
1981                      length=15), main = "Mammal Biomass")
```

1982

1983
1984
1985

1986

```
Bin_aves <- hist(log(biomass_aves$biomass),
                 breaks=seq(min(log(biomass_all$biomass)), max(log(biomass_al
l$biomass)),
                           length=15), main = "Aves Biomass")
```

Aves Biomass

Frequency

log(biomass_aves$biomass)

1987

```
# Ploting frequency-density curves for species

plot(Bin_all$mids, Bin_all$counts, pch=16, type="l", col=alpha("black", 0.5),
lwd=2, lty=2,
     ylim=c(0,12), ylab="record frequency", xlab="log biomass", main="Biomass
by Class")
points(Bin_mamm$mids, Bin_mamm$counts, pch=16, type="l", col=alpha("red", 0.5
), lwd=2)
points(Bin_aves$mids, Bin_aves$counts, pch=16, type="l", col=alpha("blue", 0.
5), lwd=3)
legend(12, 12.5, legend=c("Mammals", "Birds", "All Classes"),
```

```
1999              col=c("red", "blue", "gray"), lty=1:2, cex=0.8)
```

```
2000

2001    # Note how the peak in records of large mammals is driven by mammals, whereas
2002    the high record count in smaller values is sustained on the aves records. Ind
2003    ividually, the biomass distributions are semi-normal, whereas the joint distr
2004    ibution is more complex.
```

```
2005    Alternatively, the same data could be plotted by group with boxplots,
2006    illustrating the same sotry.
```

```
2007    bio_mamm.aves <- cbind(log(biomass_all$biomass), log(biomass_mamm$biomass), l
2008    og(biomass_aves$biomass))

2009    colnames(bio_mamm.aves) <- c("all", "mammals", "birds")
```

```
2010   boxplot(bio_mamm.aves, ylab="log biomass", xlab="class", main="Biomass by Cla
2011   ss")
```

Factor variables can similarly be parsed by class.

```
2014   # Bind the ID field, species, taxa, and volancy fields into a unified datafra
2015   me
2016
2017   bio_vol <- cbind(mosaic@mosaicID,
2018              mosaic@species,
2019              mosaic@taxaMetadat,
2020              mosaic@volancy@value)
```

```r
# Relabel the volancy field

names(bio_vol)[length(bio_vol)] <- "volancy"

# Assign into a factor class

bio_vol$volancy <- as.factor(bio_vol$volancy)

# Filter out NDYs

vol_all <- bio_vol %>%
  filter(!volancy == "NDY")

# Filter to Mammals

vol_mamm <- bio_vol %>%
  filter(Class == "Mammalia") %>%
  filter(!volancy == "NDY")

# Filter to birds

vol_aves <- bio_vol %>%
  filter(Class == "Aves") %>%
  filter(!volancy == "NDY")

# Look at the summary of factors to counts

summary(vol_mamm$volancy)
##         NDY  Non-volant Semi-volant      Volant
##           0          37           1           0
summary(vol_aves$volancy)
##         NDY  Non-volant Semi-volant      Volant
##           0           3           0          10
```

```r
# Store summaries

vol_all_summary <- summary(vol_all$volancy)
vol_mamm_summary <- summary(vol_mamm$volancy)
vol_aves_summary <- summary(vol_aves$volancy)

# Plot the points

plot(vol_all_summary[2:4],  type="b", pch=16, col=alpha("black", 0.5),
     ylab="record count", xlab="Volancy (1 = Nonvolant; 2 = Semivolant; 3 = V
olant)", main="Volancy by Class")
points(vol_mamm_summary[2:4], type="b", pch=16, col=alpha("red", 0.5))
points(vol_aves_summary[2:4], type="b", pch=16, col=alpha("blue", 0.5))
legend(2.5, 85, legend=c("Mammals", "Birds", "All Classes"),
       col=c("red", "blue", "gray"), lty=1:2, cex=0.8)
```

Volancy by Class

Volancy (1 = Nonvolant; 2 = Semivolant; 3 = Volant)

If one is more generally interested in the class breakdown of factors
across a trait field without a priori interest in a specific variable, one
can take advantage of the tools.

```
# Evaluating the variance breakdown within a given class

Volant_sum <- bio_vol %>%
  filter(volancy=="Volant")%>%
  group_by(Class, volancy)%>%
  count()

Semivolant_sum <- bio_vol %>%
```

```r
  filter(volancy=="Semi-volant")%>%
  group_by(Class, volancy)%>%
  count()

Nonvolant_sum <- bio_vol %>%
  filter(volancy=="Non-volant")%>%
  group_by(Class, volancy)%>%
  count()

volTable <- Nonvolant_sum # Generating DF

volTable <- volTable[,-2]
colnames(volTable)[2] <- "Nonvolant"
volTable[,3] <- rep(0, length(Nonvolant_sum$Class))
volTable[,4] <- rep(0, length(Nonvolant_sum$Class))
colnames(volTable)[3] <- "Volant"
colnames(volTable)[4] <- "Semivolant"
volTable[match(Volant_sum$Class, Nonvolant_sum$Class),3] <- Volant_sum$n
volTable[match(Semivolant_sum$Class, Nonvolant_sum$Class),4] <- Semivolant_sum$n

volTable
## # A tibble: 13 × 4
## # Groups:   Class [13]
##     Class           Nonvolant Volant Semivolant
##     <chr>               <int>  <dbl>      <dbl>
##  1 Actinopterygii          9      0          0
##  2 Amphibia                1      0          0
##  3 Anthozoa                5      0          0
##  4 Aves                    3     10          0
##  5 Bivalvia                3      0          0
##  6 Branchiopoda            2      0          0
##  7 Demospongiae            1      0          0
##  8 Elasmobranchii          1      0          0
##  9 Gastropoda              3      0          0
```

```
## 10 Insecta        2        0        0
## 11 Mammalia      37        0        1
## 12 Reptilia       8        0        0
## 13 <NA>           7        0        0
```

2122

2126   Vignette #3 - Basic Regression Using MOSAIC & COMPADRE

2127
2128
MOSAIC was built with the intention of its use in comparative biodemography -
to strengthen the map of which functional traits predict vital rates and
which do not. Traits in the MOSAIC database (factorial and numeric variables)
are formatted to be easily accessed and used for regression-based analyses,
among other types of analysis. MOSAIC was also designed for quick integration
with the COMADRE, COMPADRE, and PADRINO structural population model
databases, as outlined in the model exercise below.

```
2136  library(devtools)

2137  library(tidyverse)

2138  library(Rcompadre) # package to access matrix population models

2139  library(Rage) # package to perform life-history calculations
```

2140 Accessing MOSAIC

2141 Download MOSAIC from the mosaic portal. For more information on the basics of
2142 downloading MOSAIC and navigating the data structure, see: Vignette #1:

```
2143  library(devtools) # Compulsory package to pull down packages from GitHub. Ins
2144  tall if necessary.

2145  install_github("mosaicdatabase/Rmosaic") # library of navigation-aiding funct
2146  ions

2147  source_url("https://raw.githubusercontent.com/mosaicdatabase/mosaicdatabase/m
2148  ain/mosaic_fetch.R") # Link to GitHub repo

2149  mosaic <- mos_fetch("v1.0.0") # Download version 1.0.0 (active version Feb 20
2150  22)

2151  library(Rmosaic)
```

Regressing Biomass on Generation Time

In this exercise, the simple bivariate relationship of biomass on generation
time is explored through basic regression. Bare in mind, that the below offer
the building blocks for more complex multiple regression exercises and other
forms of analysis – such as geospatial interpolation and ordination-based
practices (see Vignette #4 for a brief vignette of PVA using mosaic).

#1: Extract biomass data for mammals

Extract the biomass values for mammals in the MOSAIC dataset.

```
# Extracting all biomass data

mosaic_dataframe <- data.frame(mosaic@taxaMetadat,
                                Biomass = as.numeric(mosaic@biomass@value),
                                MOSAIC_index = 1:length(mosaic@index))

# Restrict mammal data with non NA biomass data

mosaic_mammal <- subset(mosaic_dataframe,
                        Class == "Mammalia" & is.na(Biomass) == F)

# Extracting the first COMPADRE matrix id for each entry

matrix_ids_full <- mosaic@index[mosaic_mammal$MOSAIC_index]

# Extract just the first matrix id using sapply

mammal_matrix_ids <- sapply(matrix_ids_full, `[[`, 1)

# Add matrix id to the mammal data

mosaic_mammal$MatrixID <- mammal_matrix_ids

# Add full species names

mosaic_mammal$Binomial <- paste0(mosaic_mammal$Genus, " ", mosaic_mammal$Spec
ies)
```

#2: Download COMADRE matrix population models

A cdb_fetch() function is used to clone the Comadre database <www.compadre-
db.org> - which is an s3 copy of a SQL-based relational database - into an S4
data object locally manipulable in R.

Using ids associated with each matrix population model, we will subset the
Comadre database of matrix population models for the animals. First we use
the Rcompadre package to 'fetch' the Comadre database.

```
2194   # Download the most recent database
2195
2196   comadre <- cdb_fetch("comadre")
2197
2198   # Rcompadre function flags potential issues with matrices, including NAs and
2199   ergodicity
2200
2201   comadre_flag <- cdb_flag(comadre)
2202
2203   # Remove matrices with NAs and those that are non-ergodic (which throw errors
2204   in generation time)
2205
2206   comadre_correct <- subset(comadre_flag,
2207                             check_NA_A == FALSE &
2208                             check_ergodic == TRUE)
```

2209 The subsets of mosaic and Compadre can be overlapped in one line of code.

```
2210   # Subset to mammal matrices
2211
2212   comadre_biomass <- subset(comadre_correct,
2213                             MatrixID %in% mammal_matrix_ids)
```

3. Calculating generation time

Rcompadre enables sub-setting of a matrix population model (life table) into
constituent matrices representing the survival and fecundity/reproductive
components (U and F matrices, respectively). Functions in the Rage package
often require the specification of both the U and the F matrix, not the
combined A matrix. The Rage package contains a suite of common demographic
calculations - including GENERATION TIME - which are used to simplify this
exercise. Note that other common demographic derivations, including transient
indices can be extracted using the identical procedure.

```
2223   # Calculate Generation Time
2224
2225   comadre_biomass$Generation_Time <- mapply(Rage::gen_time,
2226                                             matU(comadre_biomass),
```

```
2227                                              matF(comadre_biomass))
```

2228   4. Explore the relationship of generation time data and biomass data

2229   Matrix IDs can be used to bridge generation time from comaadre with mosaic
2230   records.

```
2231   # Extract matrix ID and generation time columns (excluding other matrix metad
2232   ata)

2233

2234   comadre_biomass <- as.data.frame(comadre_biomass)

2235

2236   comadre_biomass <- comadre_biomass[,c("MatrixID", "Generation_Time")]

2237

2238   # Join the datasets, using Matrix IDs to crosswalk the data

2239

2240   mosaic_biomass <- merge(mosaic_mammal, comadre_biomass,

2241                           by = "MatrixID", all.x = TRUE)

2242   # Remove infinity values or NA

2243

2244   mosaic_biomass <- mosaic_biomass[!(is.infinite(mosaic_biomass$Generation_Time
2245   ) |

2246                                      is.na(mosaic_biomass$Generation_Time)),]

2247

2248   # Log biomass and generation time

2249

2250   mosaic_biomass$log10_biomass <- log10(mosaic_biomass$Biomass)

2251   mosaic_biomass$log10_gent <- log10(mosaic_biomass$Generation_Time)

2252

2253

2254

2255   # The plot

2256

2257   plot(mosaic_biomass$log10_biomass, mosaic_biomass$log10_gent,

2258       pch=16, cex=1.5, col=alpha("black", 0.5),

2259       xlab=expression(paste(log[10], "Biomass")),

2260       ylab=expression(paste(log[10], "Generation time"))
```

```
)

# Simple linear regression (note that we are skipping model fitting in this b
rief vignette)

lmBioGenT = lm(log10_gent~log10_biomass, data = mosaic_biomass)

summary(lmBioGenT)
##
## Call:
## lm(formula = log10_gent ~ log10_biomass, data = mosaic_biomass)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.53325 -0.16107 -0.01597  0.19936  0.38382
##
## Coefficients:
##               Estimate Std. Error t value Pr(>|t|)
## (Intercept)    0.85966    0.25372   3.388  0.00253 **
## log10_biomass  0.02762    0.05728   0.482  0.63418
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2623 on 23 degrees of freedom
## Multiple R-squared:  0.01001,    Adjusted R-squared:  -0.03303
## F-statistic: 0.2326 on 1 and 23 DF,  p-value: 0.6342
abline(lmBioGenT, col="red", lty=2, lwd=2)
```

2289
2290

2294   Vignette #4 – Dimension Reduction + MOSAIC

MOSAIC was built for integrated analyses across trait fields. Whether the
data are continuous (e.g., biomass) or categorical (e.g., dispersal
class), MOSAIC allows researchers to explore these dimensions to identify
the primary axes of variation across functional traits. Here, we will use
a Multiple Factor Analysis (MFA) to explore the primary axes of variation
between: biomass, dispersal capacity, growth determination, habitat type
and mating system.

Note: These traits are picked solely to illustrate the analysis method
that can be scaled for other traits in MOSAIC, demographic rates in
COMADRE, COMPADRE and PADRINO and climate data from ERA5-Land.

```r
library(devtools)

library(tidyverse)

library(FactoMineR) # Required for MFA.

library(factoextra) # Required for visualizing MFA.
```

Accessing MOSAIC

Download MOSAIC from the mosaic portal. For more information on the basics
of downloading MOSAIC and navigating the data structure, see: Vignette #1:

```r
remotes::install_github("mosaicdatabase/Rmosaic")

library(Rmosaic2.0)

mosaic <- mos_fetch("v1.0.0")

##

## Phylogenetic tree with 1359 tips and 1358 internal nodes.

##

## Tip labels:

##   Oxalis_acetosella, Rourea_induta, Euphorbia_telephioides, Euphorbia_font
queriana, Triadica_sebifera, Actinostemon_concolor, ...

## Node labels:

##   Node1, Node2, Node3, Node4, Node5, Node6, ...

##
```

```
## Rooted; includes branch lengths.
```

Generating Dataframe

```
# Create a dataframe storing the values of the traits of interest.

mosaic_dataframe <- data.frame(mosaic@taxaMetadat,
                               Biomass = as.numeric(mosaic@biomass@value),
                               Growth_Determination = as.factor(mosaic@growth
det@value),
                               Mating_System = as.factor(mosaic@matsyst@value
),
                               Dispersal_Capacity = as.factor(mosaic@dispcap@
value),
                               Habitat_Type = as.factor(mosaic@aquadep@value)
)

# Create a dataframe that only contains data for MFA.
# This step makes all dimension reduction functions easier.

mosaic_dataframe_subset <-  mosaic_dataframe[,c(8:12)] %>%
  drop_na() %>%
  filter(Growth_Determination != "NDY") %>%
  filter(Mating_System != "NDY") %>%
  filter(Dispersal_Capacity != "NDY") %>%
  filter(Habitat_Type != "NDY")
```

Perform MFA

```
# Run the multiple factor analysis using the MFA() function from FactoMineR.

mfa_mosaic <- MFA(mosaic_dataframe_subset, group = c(1, 1, 1, 1, 1),
    type = c("s", rep("n", 4)),
    name.group=c("Biomass","Growth_Determination","Mating_System","Dispersal_
Capacity", "Habitat_Type"),
    graph = FALSE)

```

```
# Note: Changing the graph argument to TRUE will return some diagnostic graph
s of the MFA.
```

Explore MFA

In this hypothetical example, we are interested in what traits covary with
the primary axes of variation in the MFA (i.e., Dim1 and Dim2). To
visualize these relationships, we can use the in-built fviz functions in
the factoextra package to illustrate (1) position of individuals, (2)
average effect of trait level (i.e. monogamous) on individual position
within the 2 dimensions and (3) average effect of trait type (i.e. mating
system).

```
fviz_mfa_ind(mfa_mosaic,
             repel = TRUE,
             col.ind = "dark blue")
```

2372

```
fviz_mfa_axes(mfa_mosaic,
              repel = TRUE)
```

```
fviz_hmfa_var(mfa_mosaic,
              repel = TRUE,
              choice = "group",
              col.var = "dark blue")
```

2380

2381

2386

MOSAIC was built with the intention of its use in comparative
biodemography - to strengthen the map of which functional traits predict
vital rates and which do not. Traits in the MOSAIC database (factorial and
numeric variables) are formatted to be easily accessed and used for
regression-based analyses, among other types of analysis. MOSAIC was also
designed with the analysis of demographic metrics and traits in their
environmental contexts. To this end, MOSAIC includes a separate object
(climate) which we will access and make use of in the following exercise.

```
2395   library(devtools)
2396   library(tidyverse)
2397   library(leaflet)
```

2398   Accessing MOSAIC

2399   Download MOSAIC from the mosaic portal. For more information on the basics
2400   of downloading MOSAIC and navigating the data structure, see: Vignette #1:

```
2401   remotes::install_github("mosaicdatabase/Rmosaic")
2402   library(Rmosaic2.0)
2403   mosaic <- mos_fetch("v1.0.0")
2404   ##
2405   ## Phylogenetic tree with 1359 tips and 1358 internal nodes.
2406   ##
2407   ## Tip labels:
2408   ##   Oxalis_acetosella, Rourea_induta, Euphorbia_telephioides, Euphorbia_font
2409   queriana, Triadica_sebifera, Actinostemon_concolor, ...
2410   ## Node labels:
2411   ##   Node1, Node2, Node3, Node4, Node5, Node6, ...
2412   ##
2413   ## Rooted; includes branch lengths.
```

2414   Selecting Target Matrices

To get this exercise started, we first need to select which part of MOSAIC we are interested in. For simplicity, we are here focusing on mammal biomass as was done in Vignette #2:

```r
# Extracting all biomass data
mosaic_dataframe <- data.frame(mosaic@taxaMetadat,
                               Biomass = as.numeric(mosaic@biomass@value),
                               MOSAIC_index = 1:length(mosaic@index))

# Restrict mammal data with non NA biomass data
mosaic_mammal <- subset(mosaic_dataframe,
                        Class == "Mammalia" & is.na(Biomass) == FALSE)

# Extracting the first COMPADRE matrix id for each entry
matrix_ids_full <- mosaic@index[mosaic_mammal$MOSAIC_index]

# Extract just the first matrix id using sapply
mammal_matrix_ids <- sapply(matrix_ids_full, `[[`, 1)

# Add matrix id to the mammal data
mosaic_mammal$MatrixID <- mammal_matrix_ids

# Add full species names
mosaic_mammal$Binomial <- paste0(mosaic_mammal$Genus, " ", mosaic_mammal$Species)
```

We now have a data frame containing MOSAIC IDs, taxonomic information, and biomass values for all mammals within MOSAIC. The corresponding matrix IDs are:

```r
mammal_matrix_ids
##  [1] 249092 249094 240296 240307 249126 240331 249159 240358 240362 249214
## [11] 249248 240402 249269 249273 249275 249286 240470 249399 249409 249455
## [21] 249504 240502 249524 249526 249597 249598 240504 249671 249674 249723
## [31] 249726 240674 249753 249759 249787 240511 249817 249821 249866 240565
## [41] 249878 240677 249909 240645
```

Accessing the Climate Data

With our matrix IDs at hand for analysis, we are now ready to tap into the
climate data pool contained within MOSAIC. The climate data hereafter, has
been obtained using the R-package KrigR and climate data is reported as
mean and standard deviation of monthly time-series belonging to the
location and study duration for each matrix contained within MOSAIC.

Climate Data in MOSAIC

The climate data within MOSAIC is stored in the climate object:

```
head(mosaic@climate)
```

```
##   X.1 X MatrixID      Lat       Lon StudyStart StudyEnd      GenTime      d
b
## 1   1 1   240296 61.79944 -150.3694       1976     1986 10.64398229 comadr
e
## 2   2 2   240297 61.79944 -150.3694       1976     1986        <NA> comadr
e
## 3   3 3   240298 61.79944 -150.3694       1976     1986        <NA> comadr
e
## 4   4 4   240299 61.79944 -150.3694       1976     1986        <NA> comadr
e
## 5   5 5   240300 61.79944 -150.3694       1976     1986  8.16781721 comadr
e
## 6   6 6   240301 61.79944 -150.3694       1976     1986  12.2931104 comadr
e
##   Download air_temperature air_temperature_SD total_precipitation
## 1        1          273.7485           9.862699          0.06464312
## 2        1          273.7485           9.862699          0.06464312
## 3        1          273.7485           9.862699          0.06464312
## 4        1          273.7485           9.862699          0.06464312
## 5        1          273.7485           9.862699          0.06464312
## 6        1          273.7485           9.862699          0.06464312
##   total_precipitation_SD volumetric_soil_water_layer_1
## 1              0.0420547                     0.3704328
## 2              0.0420547                     0.3704328
## 3              0.0420547                     0.3704328
## 4              0.0420547                     0.3704328
## 5              0.0420547                     0.3704328
## 6              0.0420547                     0.3704328
##   volumetric_soil_water_layer_1_SD volumetric_soil_water_layer_2
## 1                       0.02808088                     0.3684833
```

```
## 2                         0.02808088                          0.3684833
## 3                         0.02808088                          0.3684833
## 4                         0.02808088                          0.3684833
## 5                         0.02808088                          0.3684833
## 6                         0.02808088                          0.3684833
##   volumetric_soil_water_layer_2_SD volumetric_soil_water_layer_3
## 1                         0.02796726                            0.36928
## 2                         0.02796726                            0.36928
## 3                         0.02796726                            0.36928
## 4                         0.02796726                            0.36928
## 5                         0.02796726                            0.36928
## 6                         0.02796726                            0.36928
##   volumetric_soil_water_layer_3_SD volumetric_soil_water_layer_4
## 1                         0.02442513                          0.3888525
## 2                         0.02442513                          0.3888525
## 3                         0.02442513                          0.3888525
## 4                         0.02442513                          0.3888525
## 5                         0.02442513                          0.3888525
## 6                         0.02442513                          0.3888525
##   volumetric_soil_water_layer_4_SD     runoff  runoff_SD total_evaporation
## 1                         0.01339378 0.03377308 0.02755258       -0.02956482
## 2                         0.01339378 0.03377308 0.02755258       -0.02956482
## 3                         0.01339378 0.03377308 0.02755258       -0.02956482
## 4                         0.01339378 0.03377308 0.02755258       -0.02956482
## 5                         0.01339378 0.03377308 0.02755258       -0.02956482
## 6                         0.01339378 0.03377308 0.02755258       -0.02956482
##   total_evaporation_SD
## 1           0.03250795
## 2           0.03250795
## 3           0.03250795
## 4           0.03250795
## 5           0.03250795
## 6           0.03250795
```

As you can see, there are NA values for some matrix IDs in the above output. This results from either (1) missing geolocations of the underlying matrix models making extraction of environmental parameters impossible, (2) study durations of matrix models pre-dating climate data availability at monthly scale within ERA5-Land, or (3) geolocation of matrix models falling outside of land masses.

Extracting Climate Data

Since we are only interested in a subset of matrix models, we do not require the full climate data object and so we can refer to only the relevant subset of the climate object by matching MatrixID values:

```r
clim_df <- mosaic@climate[mosaic@climate$MatrixID %in% mammal_matrix_ids, ]
```

With the subsetted climate data, we can now create a data frame holding all necessary data for some exploratory analyses:

```r
analysis_df <- merge(mosaic_mammal, clim_df, by = "MatrixID")

analysis_df <- na.omit(analysis_df[analysis_df$Biomass < 1e6, ]) # we set a c
utoff on biomass here, to avoid influence of outliers. This is just a procedu
re for simplicity of this exercise
```

Let's plot the remaining data out in their geospatial context:

```r
cols <- rainbow(n = length(unique(analysis_df$Order)))

names(cols) <- unique(analysis_df$Order)

analysis_df$Col <- cols[match(analysis_df$Order, names(cols))]

Map <- leaflet(analysis_df, width = "100%")

Map <- addProviderTiles(Map, providers$Esri.WorldTopoMap)

Map <- addCircleMarkers(map = Map, lng = ~Lon, lat = ~Lat,

                        label = ~Order,

                        col = ~Col,

                        labelOptions = labelOptions(textsize = "12px")

)

Map
```

Regression & Analysis

Now we are ready to run analyses. The ones we are focussing on here are rudimentary at best and should simply serve as a demonstration of how easily one can make use of state-of-the-art ERA5-Land climate data with the MOSAIC data base.

All Mammals

First, let's assess the effect of temperature on biomass of all mammals:

```
summary(lm(data = analysis_df, Biomass ~ air_temperature))
##
## Call:
## lm(formula = Biomass ~ air_temperature, data = analysis_df)
##
## Residuals
##     Min      1Q  Median      3Q     Max
## -136457  -35516  -12225   17982  218728
##
## Coefficients:
##                 Estimate Std. Error t value Pr(>|t|)
## (Intercept)      1502659     532349   2.823  0.00991 **
## air_temperature    -5006       1850  -2.706  0.01290 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 79270 on 22 degrees of freedom
## Multiple R-squared:  0.2498, Adjusted R-squared:  0.2157
## F-statistic: 7.324 on 1 and 22 DF,  p-value: 0.0129
```

According to this, mammals in colder regions are heavier. This is in support of Bergmann's rule.

```
ggplot(data = analysis_df, aes(x = air_temperature, y = Biomass)) +
  geom_point() +
  stat_smooth(method = "lm") +
  labs(x = "Air Temperature [K]", y = "Biomass") +
  theme_bw()
```

2583

Mammals by Order

Now let's make use of the higher-resolution taxonomic information available in MOSAIC and assess the impact of air temperature on biomass by order of mammals:

```
summary(lm(data = analysis_df, Biomass ~ air_temperature * Order))
##
## Call:
## lm(formula = Biomass ~ air_temperature * Order, data = analysis_df)
##
## Residuals:
##     Min      1Q  Median      3Q     Max
## -122023  -10392       0   12873  143838
##
## Coefficients: (4 not defined because of singularities)
##                                        Estimate Std. Error t value Pr(>|t|)
## (Intercept)                             3047458    2548979   1.196    0.255
```

```
## air_temperature                            -10376      9154  -1.133     0.279
## OrderCarnivora                           -1985731   2765125  -0.718     0.486
## OrderCingulata                             -14654    148341  -0.099     0.923
## OrderDidelphimorphia                        -4262    158154  -0.027     0.979
## OrderDiprotodontia                        -3150686   3435359  -0.917     0.377
## OrderLagomorpha                             -99518     95702  -1.040     0.319
## OrderPrimates                             -2008837   3738418  -0.537     0.601
## OrderRodentia                              -219816    100383  -2.190     0.049
*
## air_temperature:OrderCarnivora                7092      9885   0.717     0.487
## air_temperature:OrderCingulata                  NA        NA      NA        NA
## air_temperature:OrderDidelphimorphia            NA        NA      NA        NA
## air_temperature:OrderDiprotodontia           10742     12153   0.884     0.394
## air_temperature:OrderLagomorpha                 NA        NA      NA        NA
## air_temperature:OrderPrimates                 6911     13007   0.531     0.605
## air_temperature:OrderRodentia                   NA        NA      NA        NA
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 75170 on 12 degrees of freedom
## Multiple R-squared:  0.6319, Adjusted R-squared:  0.2945
## F-statistic: 1.873 on 11 and 12 DF,  p-value: 0.148
```

Well, this is just a mess, but should serve to illustrate how many
different analyses can easily be performed with MOSAIC and the in-built
climate parameters.

```
ggplot(data = analysis_df, aes(x = air_temperature, y = Biomass, col = Order)
) +
  geom_point() +
  stat_smooth(method = "lm") +
  labs(x = "Air Temperature [K]", y = "Biomass") +
  theme_bw() + scale_color_manual(values = cols)
```

2631
2632

2633

2634   S5: LogNormal Distribution of Mass and Height

2635

2636
2637   Raw heights from MOSAIC v1.0.0
2638
2639

2640
2641   Log Heights from MOSAIC v.1.0.0
2642

Histogram of mass

2643
2644   Raw mass from MOSAIV v1.0.0
2645

Histogram of log(mass)

2646
2647   Log mass from MOSAIC v1.0.0