Tailoring CRISPR-Cas Immunity for the Degradation of Antibiotic Resistance Genes

Xin Li, Nan Bao, Zhen Yan, Xian-Zheng Yuan, Shu-Guang Wang *, Peng-Fei Xia *

Shandong Key Laboratory of Water Pollution Control and Resource Reuse, School of Environmental Science and Engineering, Shandong University, Qingdao 266237, China

* Correspondence:

Shu-Guang Wang

School of Environmental Science and Engineering, Shandong University, Qingdao 266237, China. Email: wsg@sdu.edu.cn

Peng-Fei Xia

School of Environmental Science and Engineering, Shandong University, Qingdao 266237, China. Email: pfxia@sdu.edu.cn

Abstract

The evolution and dissemination of antibiotic resistance genes (ARGs) are prompting severe health and environmental issues. While wastewater treatment processes are key barriers preventing the spread of ARGs, they are often sources of ARGs at the same time. An upgrading of wastewater treatment processes is imperative and urgent. ARGs confer antibiotic resistance based on the DNA sequences rather than the chemistry of DNA molecules. An ARG can be considered being degraded if its sequence was disrupted. Therefore, we present here that CRISPR-Cas immunity, an archaeal and bacterial immune system for eliminating invading foreign DNAs, can be repurposed and tailored for the degradation of ARGs. By engineering artificial IncP machinery, the designed system, namely VADER, can be successfully delivered via bacterial conjugation. Finally, we propose a new sector for ARG degradation to be implemented in wastewater treatment frameworks. In this endeavor, a prototype conjugation reactor was devised and demonstrated as the bridge connecting synthetic CRISPR-Cas immunity with wastewater treatment processes. By generating a win-win situation at the nexus of synthetic biology and environmental biotechnology, we believe that our work is not only an enterprise for tackling ARG problems but also a potential solution for managing undesired genetic materials in general in the future.

Keywords: Antibiotic resistance genes, CRISPR, Conjugation, Wastewater treatment, Synthetic biology

Introduction

Antibiotic resistance is a threatening issue for human health.[1, 2] Approximately, 4.95 million deaths were associated with antibiotic resistance in 2019, while 1.27 million deaths were direct consequences of antibiotic resistance.[2] Most antibiotic resistance results from antibiotic resistance genes (ARGs), which encode machinery that inactivate or pump out antibiotics. An ARG-harboring pathogen is more harmful to humans as the correlated antibiotic will not be effective anymore. In the environment, ARGs can disseminate via horizontal gene transfer or evolve under the selection of antibiotics, leading to increased health and environmental concerns.[1] Though many efforts have been made to understand and resolve ARG-related issues, fine control of ARGs has not yet been implemented.

Wastewater treatment is a key barrier that prevents ARGs from entering the environment.[3] It combats antibiotic resistance by degrading antibiotics, killing antibiotic resistance microbes, and directly removing ARGs. However, detection of various types of ARGs has been reported in and near wastewater treatment facilities,[4-7] indicating them potential sources for ARGs at the same time. To address this rising challenge, the fate of ARGs in engineered and natural systems has been interrogated,[8-13] and remediation methods have been developed.[14-16] Most of the current methods remove ARGs as biomolecules based on the physiochemistry of DNAs.[14] Yet, a fragment of DNA molecule is considered an ARG due to the sequence, which encodes proteins that confer resistance to the corresponding antibiotics. Thus, an ARG can be considered being degraded when the sequence was disrupted.

CRISPR-Cas system is an archaeal and bacterial immune system that has being repurposed as gene-editing tools.[17-20] As an immune system, the CRISPR-Cas system finds and cleaves a target DNA under the lead of a guide RNA (gRNA). Therefore, by applying CRISPR-Cas immunity, ARGs will be degraded. Pioneer studies have validated that CRISPR-Cas system can eliminate chromosomal ARGs for medical applications,[21-24] and the system can be delivered via conjugation.[24] These advances, however, have difficulties in finding a practical scenario. The bottlenecks mainly lie in delivery, as (1) conjugation needs physical contact between donor and recipient cells, and (2) a threshold

mount of bacterial cells and contacting time is necessary. The bottlenecks, though still challenging, may be addressed at the process-level.

We envision that a CRISPR-Cas system can be implemented in wastewater treatment processes for ARG removal. It is possible to instrument a controllable new sector (e.g., a conjugation reactor) to harness the CRISPR-Cas immunity for ARG degradation and to overcome the delivery issue. To do so, we devised and demonstrated VADER, an environmentally-aimed degradation system for eliminating ARGs, based on the *Streptococcus pyogenes* CRISPR-Cas system and the IncP conjugation machinery. Then, we constructed a prototype conjugation reactor and observed the successful delivery of VADER in the reactor. Our study shows a unique win-win situation. CRISPR-Cas system offers a novel path to tackle ARG problems in wastewater treatment, and in return, wastewater treatment provides process-level advantages in the integration of cutting-edge synthetic biology advances.

Methods

Stains and Media

Escherichia coli MG1655 (CGSC#6300) was used as a host of plasmid-borne ARG for evaluating VADER with functional methylation mechanisms. *E. coli* HST04 (Takara Bio.) was used as the methylation deficient strain to assess the effects of methylation on CRISPR-Cas mediated ARG degradation. *E. coli* S17-1 harboring the complete IncP conjugation machinery in the chromosome was used as the donor strain for conjugation.[23, 25] *E. coli* DH5α (Takara Bio.) was used for molecular cloning to generate and maintain plasmids. *E. coli* strains were cultivated in Luria-Bertani (LB) medium with ampicillin (100 μg/L), gentamycin (20 μg/L), tetracycline (25 μg/L), chloramphenicol (34 μg/L) when necessary.

Plasmid Construction

The plasmids used in this study are summarized in Table S1. pBBR1MCS5 was a lab stock plasmid,[26] and pCas9 (Addgene#42876) was a gift from Luciano Marraffini.[27] pMV-gRNA with gRNA cassette was synthesized by Beijing Liuhe BGI, and pBR322 was a

commercial plasmid from Takara Bio. The first gRNA plasmid pgRNA01 carrying gRNA01 cassette was generated via inverse PCR with a back-to-back primer pair. pCasEnv was constructed via DNA assembly with pBBR1MCS5 as the backbone and *cas9* from pCas9, and pBBR-LX01 was assembled with pBBR1MCS5 as the backbone and gRNA01 cassette. The first VADER plasmid harboring *cas9* and gRNA01 cassette was generated by fusing gRNA01 cassette from pgRNA01 to pCasEnv. Then, all VADER plasmids were directly constructed via inverse PCR to replace spacers. pELdu was constructed by fusing gRNA03 cassette to pEL11. pELnc and pEL03c were built by fusing *traJ* and *oriT* modules to pELnu and pEL03, respectively. All DNA assemblies used in plasmid generation were following protocols from the merchandiser (Takara Bio.), and inverse PCRs were following previously established protocols in the lab.[28] Plasmid extraction and purification were conducted using commercial kits from Tiangen Biotech. PCRs were performed with PrimeSTAR Max DNA Polymerase from Takara Bio. All primers used in this study were synthesized by Beijing Liuhe BGI and are listed in Table S2. The sequences of synthesized fragments are listed in Table S3. The gRNAs with spacers and PAMs are summarized in Table S4.

Bacterial transformation

Preparation of chemically competent cells and *E. coli* transformation were following the standard methods with minor modifications. For preparing competent cells, overnight culture of *E. coli* was inoculated to LB medium and grew until OD_{600} reaching 0.3. The cells were harvested at 4 °C and resuspended in 0.1 M of ice-cold $CaCl_2$, staying on ice for 30 min. After repeating the step, cells were resuspended and aliquoted in 0.1 M of $CaCl_2$ with 15% (v/v) of glycerol for storage at -80 °C. For transformation, the competent cells were first thawed on ice, and DNA was then added to the cells, following by incubation on ice for 30 min and a 60 s of heat shock at 42 °C. After recovery in LB medium for 1 h, all cells were centrifuged and resuspended in 100 μL of sterile water before plating on selection plates.

Bacterial conjugation

E. coli conjugation was performed in a 1.5-mL Eppendorf tube at a volume of 100 μL. First, donor and recipient cells were inoculated from overnight cultures and cultivated until OD_{600} reaching 0.8 – 1.0. Then, the donor and recipient cells were washed with LB medium, mixed together and kept at 37 °C for conjugation. After 1.5 h of conjugation, fresh LB medium was added and cultivated for another 1.5 h of recovery, then plating on selection plates with appropriate antibiotics. Colony forming units were measured to determine the numbers of transconjugants and recipients. Conjugation frequency was calculated according to following equation (*eq.1*).

$$\text{Conjugation frequency (per recipient)} = \frac{\text{transconjugants} \times \text{dilution rate}}{\text{recipient cell}} \times 100\% \quad (eq.1)$$

Analysis for degradation efficiency

Degradation efficiency was evaluated by checking transformants with VADER plasmids on selection plates with the corresponding antibiotic of the target ARG. In the present study, transformants were first selected on gentamycin plates to ensure the transformation of VADER plasmids. Then, at least 12 colonies from each of the plates were randomly picked and re-streaked on plates with tetracycline and ampicillin, respectively, to evaluate the degradation of *tetA* and the integrity of target plasmids. Biological duplication or triplication were performed. The degradation efficiency was calculated with *eq.2*.

$$\text{Degradation efficiency} = (1 - \frac{\text{number of colonies survived}}{\text{number of colonies picked}}) \times 100\% \quad (eq.2)$$

For conjugation experiments, the transconjugants were selected on double selective plates with gentamycin and tetracycline. As designed, transconjugants receiving a targeting VADER, with gRNA targeting *tetA*, will be dead and only escaper cells will survive. Thus, the degradation efficiency for conjugation experiments directly evaluated with tetracycline was calculated by comparing the conjugation frequencies between targeting and non-targeting VADER plasmids based on the hypothesis that they have similar conjugation frequencies.

$$\text{Degradation efficiency} = (1 - \frac{\text{conjugation frequency of targeting VADER}}{\text{conjugation frequency of non-targeting VADER}}) \times 100\% \ \ (eq.3)$$

With the assistance of pCmR giving an extra selection criterion, the transconjugants were able to be selected with gentamycin and chloramphenicol. The transconjugants were then tested with tetracycline to evaluate the VADER system. Similarly, a minimum of 12 colonies from the plates with gentamycin and chloramphenicol after conjugation were randomly picked and re-streaked on plates with tetracycline, and the number of survival colonies were recorded for calculation with *eq.2*.

Colony PCR

A colony was picked, resuspended in sterile water and then inoculated at 100 °C for 10 min. One microliter of the supernatant was used as the template for PCR amplification of designed fragments on pBR322. All primers used for colony PCR are listed in Table S2.

Prototype conjugation reactor

The conjugation reactor was demonstrated in a 30-mL glass bottle with inlet, outlet and a ventilation tube for air exchange. The reactor was operated as a sequencing batch reactor at 37 °C with a ratio of 8:1 between donor and recipient cells, including five steps: inlet, mixing, conjugation, separation and outlet. Samples were taken at time point 0, 1 h, 2 h and 3 h, and the samples were cultivated with fresh LB medium for one hour of recovery before plating to get survival colonies on selection plates. Then, the colonies were tested for tetracycline resistance to evaluated the degradation of *tetA*.

Results and Discussion

Modular design and demonstration of VADER

S. pyogenes CRISPR-Cas immunity works in a RNA-guided manner and cleaves the target DNA with the single effector Cas9 and a programmed gRNA (Figure 1A).[29] To employ this machinery, we designed VADER as a one-plasmid system with constitutive functions for simplicity and efficiency. First, we constructed a plasmid pCasEnv with (1)

cas9 from *S. pyogene*s as the ARG degradation module, (2) a selection marker (*gentR*), and (3) a broad-host *pBBR* origin of replication (ori) for expanding the application spectrum.[26] Meanwhile, gRNA01 was generated via inverse PCR with pMV-gRNA as a template (Table S1 and Table S3). The gRNA01 cassette consisting of the J23119 promoter, a spacer and the chimeric scaffold was fused to pCasEnv, generating pEL01 as the first VADER plasmid (Figure 1B). Then the following plasmids were generated in a streamlined method by inverse PCR with pEL01 as a template and programmable primer pairs to replace the spacer (Figure 1B).

As most of the ARGs are plasmid-borne,[1, 30] we focused on and evaluated VADER for eliminating ARGs on plasmids. We chose pBR322 as the target plasmid with a *tetA* gene conferring tetracycline resistance (Tet^R) and a *bla* gene conferring ampicillin resistance (Amp^R) (Figure 1C). *tetA* was regarded as the target ARG, while *bla* was used to check the integrity of the target plasmid. Three gRNAs, gRNA01, gRNA03 and gRNA06, were generated with spacers targeting different sites on the coding or non-coding strand of *tetA* (Table S4). We found that, by applying VADER, the *tetA* gene was efficiently inactivated. The resulting transformants receiving VADER plasmids pEL01 (gRNA01), pEL03 (gRNA03) and pEL06 (gRNA06) (Table S1 and Table S3) all lost Tet^R and Amp^R (Figure 1D and 1E), indicating the degradation of *tetA* and the elimination of pBR322. Under the current design, a VADER plasmid contains *pBBR* ori which shows 8 – 10 copies of plasmids in a cell, and the target plasmid pBR322 harbors *colE1* ori, which gives 15 – 20 copies of *tetA*.[31] To further explore the capability of VADER, we designed two plasmids with varied copy numbers by replacing *colE1* ori with *p15a* ori and *pUC* ori, generating pBR322-p15a and pBR322-pUC, respectively. pBR322-p15a presents 8 – 10 copies of *tetA* per cell, and pBR322-pUC gives 500 – 700 copies of *tetA* per cell.[32] We transformed pEL03 to *E. coli* with pBR322-p15a and pBR322-pUC and observed successful elimination of *tetA* on these plasmids (Figure 1E). Normally, mobilizable plasmids, such as RP4-2 and RK2, are low-copy plasmids with less than 10 copies in a cell.[30, 33] As VADER managed to degrade ARG carried by a plasmid with a copy number up to 700, it shows sufficient capability for degradation of plasmid-borne ARGs.

Figure 1. Design and demonstration of VADAR. (A) Working mechanism of CRISPR-Cas immunity on ARG degradation. (B) Design scheme of VADER plasmids and the streamlined generation of VADAR plasmids. (C) Maps of target plasmids harboring the target ARG *tetA* with different copy numbers. (D) Demonstration of VADER plasmids pEL01, pEL03 and pEL06 for *tetA* degradation and the validation of plasmid integrity with pCasEnv, pBBR-LX01 and pELnu as

controls. The transformants were first selected on plates with gentamycin to ensure the transformation of VADER plasmids. Then, colonies were randomly picked and streaked on plates with tetracycline and ampicillin. (E) Degradation efficiencies of VADER calculated via evaluation of tetracycline resistance and ampicillin resistance of randomly picked colonies from the transformants with VADER plasmids. (F) Electrophoresis documentation of PCR fragments with primer pairs designed for amplification of the intact *tetA*, small fragments nearby (192 bp of distance) and away from (1096 bp of distance) the cleavage site by VADER.

Furthermore, we tested whether *tetA* was actually degraded. Three pairs of primers were designed to detect (1) *tetA* as an intact gene with a 1093-bp fragment, (2) a small fragment (327 bp) near the cleavage site, and (3) another small fragment (304 bp) that has a distance of 1095 bp from the cleavage site (Figure 1F). We transformed non-targeting (pELnu) and targeting (pEL03) VADER plasmids to *E. coli* with either pBR322, pBR322-p15a or pBR322-pUC, and randomly picked the transformants for PCR analysis. We identified the PCR fragments with all three primer pairs for the transformants with the non-targeting VADER and for the positive controls (purified plasmids), but no PCR fragments for the colonies receiving the targeting VADER could be detected (Figure 1F). These results demonstrated that *tetA* was degraded via a single cleavage in the middle, and the resulting degradation covered the whole length of *tetA*.

Customizing and duplexing of gRNAs

Despite the demonstration of VADER, we encountered difficulties in generating VADER plasmids via inverse PCR. In some cases, the number of transformants was quite limited, and the plasmids from these transformants had mutations. These might result from the toxicity of foreign CRISPR-Cas systems in bacteria,[34, 35] and the toxicity as a selective pressure drove mutations in the CRISPR-Cas system. Previous reports highlighted the toxicity of CRISPR-Cas in bacteria,[35-37] and, specifically, Beisel and colleagues noticed a high mutation rate in generating plasmids for CRISPR-Cas-based base editing.[35] For better-streamlined plasmid generation, further optimization of VADER is necessary.

Figure 2. (A) Mutation patterns in J23119 promoter of mutated VADER plasmids. (B) Validation of re-generated pEL03 with J23119* and J23119** promoters. TATA box regions were highlighted in red. (C) Degradation efficiency of VADER plasmid pEL03 with J23119, J23119* and J23119** promoter driving gRNA03, respectively.

We examined the mutation patterns in the mutated plasmids and found most of the mutations lie in the TATA box of the J23119 promoter leading gRNAs in our design (Figure 2A). We tested a representative mutated plasmid when generating pEL14 (gRNA14, Table S3), with TAAAT instead of TATAAT as the TATA box and surprisingly observed high efficiency of ARG degradation as well (Figure S1). We hypothesized that the mutated J23119 promoters have lower strength and alleviate the toxicity of CRISPR-Cas systems, making them better candidate promoters. Two new promoters J23119* and J23119** were designed with one nucleotide deleted at the position -12 and -11, respectively. The pEL03 plasmid was re-constructed at ease with these new promoters (Figure 2B), and both constructs showed the same efficiency (100%) in *tetA* degradation compared to the original J23119 promoter (Figure 2C). We demonstrated the customized J23119** promoter with two more VADER plasmids, pEL12 and pEL13, both giving 100% efficiency of *tetA* degradation (Figure S2). The results suggest that promoter engineering for fine-tuning gRNA levels will be a solution for alleviating CRISPR-Cas toxicity in bacteria.

Figure 3. (A) Design scheme of tandem gRNAs for duplex VADER system with gRNA11 and gRNA03 as an example. Degradation efficiencies of VADER plasmids pEL11 (with gRNA11) and pELdu (with tandem gRNA11 and gRNA03) for *tetA* degradation (B) and plasmid elimination (C) in *E. coli* with functional methylation mechanisms and with methylation deficiency.

Moreover, we observed the low on-target efficiency of VADER for certain target sequences. A dual gRNA system is necessary for efficient degradation of ARGs in case one of the gRNAs is not functional. As an example, we generated a VADER plasmid pELdu with tandem gRNAs, gRNA11 and gRNA03 (Figure 3A). In a preliminary test, we found that pEL11 with gRNA11 cannot degrade *tetA* (Figure 3B). This may be a result of DNA methylation or protospacer specificity. When we employed pEL11 for *tetA* degradation in methylation deficient *E. coli* (*dam* and *dcm*), we found that *tetA* was efficiently eliminated in half of the transformants (Figure 3B), implying a combined mechanism underlying the low on-target efficiency for certain protospacers. Then, we applied pELdu in *E. coli* with functional methylation mechanism and methylation deficiency. One hundred percent efficiency of *tetA* degradation was observed in all tests, indicating that our duplex system worked well as double insurance for ARG degradation.

Integrating IncP conjugation module for delivery

Figure 4. Working mechanism of conjugative delivery of VADER (A) and rational design of artificial IncP modules (*traJ* and *oriT*) (B). (C) Conjugation frequency of VADER evaluated by a non-targeting VADER plasmid pELnc. (D) Degradation efficiency of VADER plasmid pEL03c at different ratio between donor and recipient cells. (E) Mutation rate of a non-targeting VADER plasmid pELnc determined by morphology of colonies and double checked by PCR and sequencing. (F) Experimental scheme for validation of VADER. For a direct selection by gentamycin (Gent^R) and tetracycline resistances (Tet^R), the transconjugants receiving a targeting VADER will, by design, be dead which making direct evaluation of VADER impossible. With the

assistance of an additional plasmid harboring a third selection criterion (Cm[R]), the transconjugants with a targeting VADER can be selected first by Gent[R] and Cm[R], then being evaluated on a second round by tetracycline resistance (Tet[R]). (G) Conjugation frequency of a targeting VADER evaluated directly by selection by Gent[R] and Cm[R]. (H) Degradation efficiency of VADER evaluated by a second round of selection on Tet[R]. Forty-eight colonies were randomly picked for evaluation (n = 48). (I) Demonstration of VADER on selection plates.

Transformation is, however, not suitable for the delivery of VADER in the real world due to the requirement of cell competence. One alternative is transduction depending on virus-mediated delivery via, for instance, bacteriophages.[23] Though transduction showed great potential in medical applications and eukaryotic gene-editing, it is often highly host-specific and has a limitation in cargo sizes. Conjugation is another possible way for delivery of genetic systems, as it has a broad spectrum of recipient cells, and the feasibility has been demonstrated in principle.[23-25] By applying conjugation machinery, when the donor cell with VADER contacts with the recipient cell harboring ARG, it delivers VADER and then VADER will degrade the target ARG in vivo (Figure 4A).

As such, we engineered VADER with artificial IncP machinery to allow conjugative delivery. We chose *E. coli* S17-1 strain as the host with the complete sequence of RP4-2 plasmid in the chromosome,[23] and incorporated the *traJ* and *oriT* modules into VADER plasmids to enable conjugative transfer (Table S3 and Figure 4B). To better fit further application scenarios, we performed conjugation in liquid medium by mixing donor and recipient cultures instead of filter-mating or plate-based methods. Efficient conjugation was observed (Figure 4C), and the conjugation frequency was increased along with the increased ratio between donor and recipient cells, reaching 1.27×10^{-3} transconjugants per recipient at the ratio of 8:1 between donor and recipient cells (Figure 4C). Then, the degradation efficiencies of VADER were calculated by dividing the conjugation frequency of the targeting VADER (pEL03c) by that of a non-targeting VADER (pELnc). We found similar and promising degradation efficiencies for all experiments (97.04% to 99.51%) (Figure 4D), while a complete degradation of ARGs was obstacled by escaper cells, shown as survival colonies on the selection plates, from the targeting VADER. These escaped colonies were analyzed, and IS-4 like transposon sequences were identified in

plasmids from all escaped colonies at the position of +22 in the *cas9* sequence, thus resulting in the lost function of VADER (Figure S2). Unexpectedly, we also found transposon insertions in non-targeting VADER plasmids (Figure S2), and the insertions occurred in a frequency of approximately 5% (2.60% to 5.44%) among all colonies (Figure 4E). A similar observation of unexpected insertions was also documented in a previous report about *E. coli* S17-1 mediated conjugation.[38] These insertions might result from the chromosomal RP4-2 sequence in *E. coli* S17-1, which contains *oriT* initiating the transfer of chromosome sequences.

The efficiency of VADER cannot be directly calculated, as all the cells receiving a targeting VADER, as designed, will be dead when being selected by the target ARG (Figure 4F). Therefore, the degradation efficiencies were calculated by comparing targeting and non-targeting VADER, based on the assumption that targeting and non-targeting VADER plasmids have similar conjugation frequencies. For a more concrete and direct validation, we transformed an assistant plasmid pCmR with chloramphenicol resistance (Cm^R) for the selection of living cells receiving a targeting VADER before testing for the target ARG. With this experimental design, we were able to select cells with Cm^R, harbored by the target cells besides the target ARG, and Gent^R, harbored by VADER plasmids, to get colonies receiving VADER plasmids. Then, the colonies were tested for Tet^R to evaluate the function of VADER for ARG degradation (Figure 4F). The results showed that the targeting and non-targeting VADER showed similar conjugation frequency (Figure 4G), which confirmed our assumption and data, and the targeting VADER exhibited high degradation efficiency (Figure 4H and 4I). Notably, we observed 100% efficiency in *tetA* degradation and did not find escaper cells from the targeting VADER, and no insertions in the non-targeting VADER were identified. Taking together, VADER can efficiently degrade plasmid-borne ARGs via conjugative delivery.

A prototype conjugation reactor for VADER

Wastewater treatment processes contain different sectors and remediates contaminates or recycles resources in designed units. For instance, the anaerobic-anoxic-aerobic (A2O) system shows process-level advantages to simultaneously remove organic contaminants,

nitrogen and phosphorus by setting up anaerobic, micro-aerobic, and aerobic regions.[39] At the process level, it is rational and possible to expand conventional processes by instrumenting a specialized sector for ARG degradation, and we propose a conjugation reactor as the additional unit. The conjugation reactor contains donor cells with targeting VADER, and when the recipient cells with target ARGs (e.g., microbes from activated sludge process) enter the reactor, VADER will be delivered to degrade the target ARGs. Finally, the ARG-free cells can be released for further treatment (e.g., sludge processing) or return to their assigned bioprocess.

Figure 5. (A) Configuration of the prototype conjugation reactor with a total volume of 30 mL and a working volume of 13.5 mL. (B) Operation scheme of the conjugation reactor following a sequencing batch mode in five stages: inlet, mixing, conjugation, separation, and outlet. (C) Conjugation frequency of VADER performed in the conjugation reactor. Samples were collected at different time points and cultivated in non-selective medium for one hour before plating. (D) Degradation efficiency of VADER in reactor. The efficiency was calculated by testing tetracycline resistance of randomly picked colonies (n = 48) of transconjugants with VADER.

As a proof of principle, we built a milliliter-level conjugation reactor (Figure 5A), and the conjugation reactor works in a sequencing batch mode. With pre-cultivated donor cells inside the reactor, the process can be divided into five stages: (1) the target recipient cells

as inlet enter the reactor, (2) stirring to mix the donor and recipient cells, (3) conjugation takes place in the reactor, (4) vigorous stirring to separate conjugated cells, and (5) discharge of ARG-free cells (Figure 5B). We tested our design using *E. coli* S17-1 with VADER plasmid pEL03c as the donor, and *E. coli* with pBR322 and pCmR as the target recipient. pCmR was included to enable the evaluation of ARG degradation with living cells (Figure 4F). By setting 8:1 as the ratio between the donor and recipient, we found efficient delivery of VADER after 2 h of conjugation (1.93×10^{-4} transconjugants per recipient), and the conjugation frequency reached 3.14×10^{-3} transconjugants per recipient at 3 h (Figure 5C). No longer than 3 h of evaluation was performed to minimize the exaggeration effect caused by cell growth. Then, we randomly picked colonies of recipients with the working VADER after 2 h and 3 h of conjugation, respectively, and tested for the Tet^R. As expected, all tested cells cannot grow on the selection plates, indicating successful degradation of the target ARG with 100% efficiencies (Figure 5D). Although our prototype reactor only has a working volume of 13.5 mL, which is way smaller than a real-world reactor, we scaled up the conjugation system from microliter-level to milliliter-level by over 100 times. While this is a small step towards implementing a practical conjugation reactor in wastewater treatment processes, it shows the first demonstration of milliliter-level conjugation and exhibited the great potential of a conjugation reactor for ARG removal.

Meanwhile, we do not want to overemphasize our work nor omit the drawbacks of VADER. We admit that (1) the conjugation frequency is still not high enough for practical applications, (2) mutations in VADER plasmids during conjugation can still occur, and (3) the conjugation reactor is currently a prototype under sub-optimal conditions. Further work will be taken place in developing more efficient and stable conjugation systems, and in optimizing and scaling up the conjugation reactor. Moreover, VADER is a genetic system with ARG as a selection marker and a foreign CRISPR-Cas system as the functional module. So, VADER requires a fine control to prevent leakage and distribution by combining physical containment of the reactor and pre-installed biocontainment modules.[40]

Beyond antibiotic resistance

Here, we demonstrated that the combination of tailored CRISPR-Cas immunity, IncP machinery and conjugation reactor can be employed for ARG degradation. For the first time, we put the CRISPR-Cas system itself rather than a CRISPR-engineered organism in a reactor for environmental application. As the nexus bridging synthetic biology and environmental bioprocess, the conjugation reactor provides a new sector for the wastewater treatment frameworks to allow integration of CRISPR-Cas immunity for resolving antibiotic resistance problems, and offers a scenario where conjugative delivery of genetic systems becomes practical. This unique win-win situation sheds light on a synthetic biology approach in the evolution of environmental biotechnology.

With the rising of synthetic biology, genetic materials, including but not limited to ARGs, have been used for generating genetically modified organisms and artificial genetic systems in all aspects of human lives,[40, 41] committing high probability of spread in the environment.[42] Genetic materials, especially DNAs, are biomolecules and more, where the DNA sequences determine their functions and impacts. They are stable, inheritable, transferable and sometimes self-recoverable, but the environmental and ecological impacts are still mysteries.[42] While pre-installed or external biocontainment approaches have been investigated and instigated to warrant biosafety, wastewater treatment and other environmental systems, as key barriers, will inevitably be challenged by undesired natural and synthetic genetic materials in the near future. Therefore, a new sector in wastewater treatment framework specialized for removal of genetic materials or genetic pollutants is crucial, and our work gives a paradigm in this endeavor. On the path towards a healthy and sustainable future, we believe the cross-disciplinary advantages of synthetic biology and environmental biotechnology are just awakening.

Conflict of Interest

The authors declare no conflict of interest

Associated Content

Supporting Information

Summary of plasmids and primers used in this study; summary of synthesized DNA fragments; summary of gRNAs with protospacer and PAMs; figures supporting the main results.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (U20A20146), Distinguished Young Scholar Program of Shandong Province (Overseas) and Qilu Young Scholar Program of Shandong University (to P.-F.X.).

References

(1) Larsson, D. G. J.; Flach, C. F., Antibiotic resistance in the environment. *Nat. Rev. Microbiol.* **2021**, 1-13.

(2) Collaborators, A. R., Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* **2022,** *399*, 629-655.

(3) Dai, D.; Brown, C.; Burgmann, H.; Larsson, D. G. J.; Nambi, I.; Zhang, T.; Flach, C. F.; Pruden, A.; Vikesland, P. J., Long-read metagenomic sequencing reveals shifts in associations of antibiotic resistance genes with mobile genetic elements from sewage to activated sludge. *Microbiome* **2022,** *10*, 20.

(4) Czekalski, N.; Diez, E. G.; Burgmann, H., Wastewater as a point source of antibiotic-resistance genes in the sediment of a freshwater lake. *ISME J.* **2014,** *8*, 1381-1390.

(5) Corno, G.; Yang, Y.; Eckert, E. M.; Fontaneto, D.; Fiorentino, A.; Galafassi, S.; Zhang, T.; Di Cesare, A., Effluents of wastewater treatment plants promote the rapid stabilization of the antibiotic resistome in receiving freshwater bodies. *Water Res.* **2019,** *158*, 72-81.

(6) Ju, F.; Beck, K.; Yin, X.; Maccagnan, A.; McArdell, C. S.; Singer, H. P.; Johnson, D. R.; Zhang, T.; Burgmann, H., Wastewater treatment plant resistomes are shaped by bacterial composition, genetic exchange, and upregulated expression in the effluent microbiomes. *ISME J.* **2019,** *13*, 346-360.

(7) Lee, J.; Ju, F.; Maile-Moskowitz, A.; Beck, K.; Maccagnan, A.; McArdell, C. S.; Dal Molin, M.; Fenicia, F.; Vikesland, P. J.; Pruden, A.; Stamm, C.; Burgmann, H., Unraveling the riverine antibiotic resistome: The downstream fate of anthropogenic

inputs. *Water Res.* **2021,** *197*, 117050.

(8) Choi, Y.; He, H.; Dodd, M. C.; Lee, Y., Degradation kinetics of antibiotic resistance gene mecA of methicillin-resistant *Staphylococcus aureus* (MRSA) during water disinfection with chlorine, ozone, and ultraviolet light. *Environ. Sci. Technol.* **2021,** *55*, 2541-2552.

(9) Gray, H. K.; Arora-Williams, K. K.; Young, C.; Bouwer, E.; Davis, M. F.; Preheim, S. P., Contribution of time, taxonomy, and selective antimicrobials to antibiotic and multidrug resistance in wastewater bacteria. *Environ. Sci. Technol.* **2020,** *54*, 15946-15957.

(10) Jong, M. C.; Harwood, C. R.; Blackburn, A.; Snape, J. R.; Graham, D. W., Impact of redox conditions on antibiotic resistance conjugative gene transfer frequency and plasmid fate in wastewater ecosystems. *Environ. Sci. Technol.* **2020,** *54*, 14984-14993.

(11) Zammit, I.; Marano, R. B. M.; Vaiano, V.; Cytryn, E.; Rizzo, L., Changes in antibiotic resistance gene levels in soil after irrigation with treated wastewater: a comparison between heterogeneous photocatalysis and chlorination. *Environ. Sci. Technol.* **2020,** *54*, 7677-7686.

(12) Zhu, L.; Zhou, Z.; Liu, Y.; Lin, Z.; Shuai, X.; Xu, L.; Chen, H., Comprehensive understanding of the plasmid-mediated colistin resistance gene *mcr-1* in aquatic environments. *Environ. Sci. Technol.* **2019,** *54*, 1603-1613.

(13) Pallares-Vega, R.; Hernandez Leal, L.; Fletcher, B. N.; Vias-Torres, E.; van Loosdrecht, M. C. M.; Weissbrodt, D. G.; Schmitt, H., Annual dynamics of antimicrobials and resistance determinants in flocculent and aerobic granular sludge treatment systems. *Water Res.* **2021,** *190*, 116752.

(14) Lu, X.; Hou, J.; Yang, K.; Zhu, L.; Xing, B.; Lin, D., Binding dorce and site-determined desorption and fragmentation of antibiotic resistance genes from metallic nanomaterials. *Environ. Sci. Technol.* **2021,** *55*, 9305-9316.

(15) Ren, S.; Boo, C.; Guo, N.; Wang, S.; Elimelech, M.; Wang, Y., Photocatalytic reactive ultrafiltration membrane for removal of antibiotic resistant bacteria and antibiotic resistance genes from wastewater effluent. *Environ. Sci. Technol.* **2018,** *52*, 8666-8673.

(16) Zhang, C.; Zhao, X.; Wang, C.; Hakizimana, I.; Crittenden, J. C.; Laghari, A. A., Electrochemical flow-through disinfection reduces antibiotic resistance genes and horizontal transfer risk across bacterial species. *Water Res.* **2022,** *212*, 118090.

(17) Liu, D.-F.; Li, W.-W., Genome editing techniques promise new breakthroughs in water environmental microbial biotechnologies. *ACS ES&T Water* **2020,** *1*, 745-747.

(18) Wu, J.; Cheng, Z. H.; Min, D.; Cheng, L.; He, R. L.; Liu, D. F.; Li, W. W., CRISPRi system as an efficient, simple platform for rapid identification of genes involved in pollutant transformation by *Aeromonas hydrophila*. *Environ. Sci. Technol.* **2020,** *54*, 3306-3315.

(19) Pickar-Oliver, A.; Gersbach, C. A., The next generation of CRISPR-Cas technologies and applications. *Nat Rev Mol Cell Biol* **2019,** *20*, 490-507.

(20) Wu, J.; Liu, D. F.; Li, H. H.; Min, D.; Liu, J. Q.; Xu, P.; Li, W. W.; Yu, H. Q.; Zhu, Y. G., Controlling pathogenic risks of water treatment biotechnologies at the source by genetic editing means. *Environ. Microbiol.* **2021**.

(21) Gomaa, A. A.; Klumpe, H. E.; Luo, M. L.; Selle, K.; Barrangou, R.; Beisel, C. L., Programmable removal of bacterial strains by use of genome-targeting CRISPR-Cas systems. *mBio* **2014,** *5*, e00928-13.

(22) Luo, M. L.; Jackson, R. N.; Denny, S. R.; Tokmina-Lukaszewska, M.; Maksimchuk, K. R.; Lin, W.; Bothner, B.; Wiedenheft, B.; Beisel, C. L., The CRISPR RNA-guided surveillance complex in *Escherichia coli* accommodates extended RNA spacers. *Nucleic Acids Res.* **2016,** *44*, 7385-94.

(23) Citorik, R. J.; Mimee, M.; Lu, T. K., Sequence-specific antimicrobials using efficiently delivered RNA-guided nucleases. *Nat. Biotechnol.* **2014,** *32*, 1141-5.

(24) Reuter, A.; Hilpert, C.; Dedieu-Berne, A.; Lematre, S.; Gueguen, E.; Launay, G.; Bigot, S.; Lesterlin, C., Targeted-antibacterial-plasmids (TAPs) combining conjugation and CRISPR/Cas systems achieve strain-specific antibacterial activity. *Nucleic Acids Res.* **2021,** *49*, 3584-3598.

(25) de la Cruz, F.; Frost, L. S.; Meyer, R. J.; Zechner, E. L., Conjugative DNA metabolism in Gram-negative bacteria. *FEMS Microbiol. Rev.* **2010,** *34*, 18-40.

(26) Kovach, M. E.; Elzer, P. H.; Hill, D. S.; Robertson, G. T.; Farris, M. A.; Roop, R. M., 2nd; Peterson, K. M., Four new derivatives of the broad-host-range cloning vector

pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **1995,** *166*, 175-6.

(27) Jiang, W.; Bikard, D.; Cox, D.; Zhang, F.; Marraffini, L. A., RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat. Biotechnol.* **2013,** *31*, 233-9.

(28) Xia, P. F.; Casini, I.; Schulz, S.; Klask, C. M.; Angenent, L. T.; Molitor, B., Reprogramming acetogenic bacteria with CRISPR-targeted base editing via deamination. *ACS Synth. Biol.* **2020,** *9*, 2162-2171.

(29) Jiang, F.; Doudna, J. A., CRISPR-Cas9 structures and mechanisms. *Annu. Rev. Biophys.* **2017,** *46*, 505-529.

(30) Dimitriu, T.; Matthews, A. C.; Buckling, A., Increased copy number couples the evolution of plasmid horizontal transmission and plasmid-encoded antibiotic resistance. *Proc. Natl. Acad. Sci. U. S. A.* **2021,** *118*, e2107818118.

(31) Lee, C.; Kim, J.; Shin, S. G.; Hwang, S., Absolute and relative QPCR quantification of plasmid copy number in *Escherichia coli*. *J. Biotechnol.* **2006,** *123*, 273-80.

(32) Jahn, M.; Vorpahl, C.; Hubschmann, T.; Harms, H.; Muller, S., Copy number variability of expression plasmids determined by cell sorting and droplet digital PCR. *Microb. Cell Fact.* **2016,** *15*, 211.

(33) Gerlitz, M.; Hrabak, O.; Schwab, H., Partitioning of broad-host-range plasmid RP4 is a complex system involving site-specific recombination. *J. Bacteriol.* **1990,** *172*, 6194-203.

(34) Selle, K.; Barrangou, R., Harnessing CRISPR-Cas systems for bacterial genome editing. *Trends Microbiol.* **2015,** *23*, 225-32.

(35) Vento, J. M.; Crook, N.; Beisel, C. L., Barriers to genome editing with CRISPR in bacteria. *J Ind Microbiol Biotechnol* **2019,** *46*, 1327-1341.

(36) Xia, P. F.; Ling, H.; Foo, J. L.; Chang, M. W., Synthetic biology toolkits for metabolic engineering of cyanobacteria. *Biotechnol. J.* **2019,** *14*, e1800496.

(37) Banno, S.; Nishida, K.; Arazoe, T.; Mitsunobu, H.; Kondo, A., Deaminase-mediated multiplex genome editing in *Escherichia coli*. *Nat. Microbiol.* **2018,** *3*, 423-429.

(38) Strand, T. A.; Lale, R.; Degnes, K. F.; Lando, M.; Valla, S., A new and improved host-independent plasmid system for RK2-based conjugal transfer. *PLoS One* **2014,** *9*, e90372.

(39) Wang, D.; Tooker, N. B.; Srinivasan, V.; Li, G.; Fernandez, L. A.; Schauer, P.; Menniti, A.; Maher, C.; Bott, C. B.; Dombrowski, P.; Barnard, J. L.; Onnis-Hayden, A.; Gu, A. Z., Side-stream enhanced biological phosphorus removal (S2EBPR) process improves system performance - A full-scale comparative study. *Water Res.* **2019,** *167*, 115109.

(40) Xia, P. F.; Ling, H.; Foo, J. L.; Chang, M. W., Synthetic genetic circuits for programmable biological functionalities. *Biotechnol. Adv.* **2019,** *37*, 107393.

(41) Auslander, S.; Auslander, D.; Fussenegger, M., Synthetic biology - the synthesis of biology. *Angew Chem Int Ed Engl* **2017,** *56*, 6396-6419.

(42) Power, M. E., Synthetic threads through the web of life. *Proc. Natl. Acad. Sci. U. S. A.* **2021,** *118*, e2004833118.