AlphaFold Models of Small Proteins Rival the Accuracy of Solution NMR Structures

Roberto Tejero[1,*], Yuanpeng J. Huang[2], Theresa A. Ramelot[2], and Gaetano T. Montelione[2.*]

[1] Departamento de Química Física, Universidad de Valencia, Dr. Moliner, 50 46100-Burjassot, Valencia, SPAIN

[2] Dept of Chemistry and Chemical Biology, Center for Biotechnology and Interdisciplinary Sciences, Rensselaer Polytechnic Institute, Troy, New York, 12180 USA

*Corresponding authors. Email: roberto.tejero@uv.es; monteg3@rpi.edu

Key words: Protein structure prediction, solution NMR, deep learning, AlphaFold

Roberto Tejero   orcid: 0000-0003-2504-5988   email: roberto.tejero@uv.es
Yuanpeng J. Huang   orcid: 0000-0002-3374-786X   email: huangy26@rpi.edu
Theresa A. Ramelot   orcid: 0000-0002-0335-1573   email: ramelt2@rpi.edu
Gaetano T. Montelione: orcid: 0000-0002-9440-3059   email: monteg3@rpi.edu

Abbreviations: AF – AlphaFold; AF2 – AlphaFold2; ANSURR – Accuracy of NMR Structures Using RCI and Rigidity; GDT – Global Distance Test; NOE – nuclear Overhauser effect; NOESY – NOE spectroscopy; PAG – PolyAcrylamide Gel (stretch); PEG – Polythylene Glycol; phage – PF1 phage; PSVS – Protein Structure Validation Software suite; RCI – Random Coil Index for assessing conformational flexibility from chemical shift data; RDC – Residual Dipolar Coupling; RMSD – Root Mean Squared Deviation; RPF-DP score – Recall, Precision, F-measure Discrimination Power score.

Abstract

Recent advances in molecular modeling using deep learning have the potential to revolutionize the field of structural biology. In particular, AlphaFold has been observed to provide models of protein structures with accuracy rivaling medium-resolution X-ray crystal structures, and with excellent atomic coordinate matches to experimental protein NMR and cryo-electron microscopy structures. Here we assess the hypothesis that AlphaFold models of small proteins have accuracies (based on comparison against experimental data) similar to experimental solution NMR structures. We selected six representative small proteins with structures determined by both NMR and X-ray crystallography, and modeled each of them using AlphaFold. Using several structure validation tools integrated under the Protein Structure Validation Software suite (PSVS), we then assessed how well these models fit to experimental NMR data, including NOESY peak lists (RPF-DP scores), comparisons between predicted rigidity and chemical shift data (ANSURR scores), and ^{15}N-^1H residual dipolar coupling data (RDC Q scores) analyzed by software tools integrated in the PSVS suite. Remarkably, the fits to NMR data for the protein structure models predicted with AlphaFold are generally similar, or better, than for the corresponding experimental NMR or X-ray crystal structures. These results document the value of PSVS for model *vs*. data assessment of protein NMR structures, and the potential for using AlphaFold models for guiding analysis of experimental NMR data and more generally in structural biology.

Introduction

Recent advances in protein structure prediction based on deep learning from experimental protein structure data have the potential to revolutionize structural biology. Building on advances in attention-based machine learning(Vaswani et al., 2017; Huang et al., 2019), contact prediction based on sequence co-variance (Marks et al., 2011; Morcos et al., 2011; Marks et al., 2012; Ovchinnikov et al., 2015; Ovchinnikov et al., 2016; Buchan and Jones, 2018), the massive and still growing databases of genomic sequence data, and the rapidly growing Protein Data Bank of experimental protein structures, these methods are being recognized as major advances enabling new structural biology research (Jones and Thornton, 2022).

In the 2020 Critical Assessment of Protein Structure Prediction (CASP14), the DeepMind AlphaFold2 (AF2) deep learning method (Jumper et al., 2021a; Jumper et al., 2021b) demonstrated outstanding performance in blind predictions of protein structure, delivering excellent structural matches to experimental models derived from X-ray crystallography, NMR and cryoEM data, over a wide range of target difficulty (Kryshtafovych et al., 2021). These AlphaFold2 model predictions had an unprecedented high accuracy, assessed by backbone atomic coordinate global distance test (GDT_TS) scores. On 96 CASP14 targets AF2 models had a mean GTD_TS of 0.88 ± 0.1, corresponding to a backbone atom root-mean-squared deviation (RMSD) between predicted and experimental protein structures of about 1.5 Å (Kryshtafovych et al., 2021). Buried sidechain conformations in these blind predictions of protein structure are also generally a remarkably good match between the predicted model and experimental structure (Pereira et al., 2021). Soon afterward, the related RosettaFold (Baek et al., 2021b) method also demonstrated excellent modeling accuracy on natural proteins, and was found to be particularly successful in modeling *de novo* designed proteins. These results have opened the door to innovative *de novo* protein design approaches using these platforms (Anishchenko et al., 2021). In addition, it was also quickly recognized that the structures of protein-protein complexes and multimeric assemblies can often be reliably modeled using modified approaches with these same AI platforms (Baek et al., 2021a; Baek et al., 2021b; Evans et al., 2021; Humphreys et al., 2021; Colman et al., 2022; Mondal et al., 2022). While

challenges remain, particularly for dynamic protein systems and complex multiprotein assemblies, these methods are already having a major impact on structural biology.

These advances are particularly relevant for structural studies of proteins using NMR data. In the 2017 CASP13 blind protein structure prediction experiment, we organized a "NMR-guided prediction" challenge for the CASP protein structure prediction community called CASP-NMR (Sala et al., 2019). In this project we provided 13 simulated and real NMR data sets for 10 small (80 to 326 residues) proteins, including interatomic contacts obtained (or, for simulated data, obtainable) from NOESY experiments and backbone dihedral restraints obtained (or obtainable) from backbone chemical shift data. These included NOESY data typical of that obtained for $^{15}N,^{13}C$-enriched, perdeuterated proteins up to about 40 kDa, which were simulated and used to generate tables of ambiguous contacts using simple NOESY peak assignments protocols. These Ambiguous Contact Lists were provided, together with backbone dihedral angle restraints obtainable from chemical shift data, to the CASP prediction community for data-assisted prediction. Real NMR data collected for a *de novo* designed protein were also used to generate ambiguous contact tables and chemical-shift based backbone dihedral angle restraints, that were then provided to CASP13 predictor groups, including one set of (ambiguous) experimental NMR-based contacts in which only backbone resonance (no sidechain) assignments were available. The CASP community was then challenged to use these data to "guide" blind protein structure predictions. These predictions were compared to NMR-based models generated from these data by experts using conventional methods, or against the NMR data. Remarkably, several CASP13 prediction groups provided models that matched the reference structures and/or fit these NMR data even better than the models generated by conventional expert NMR structure analysis. Notable among these top-performing NMR-guided prediction groups were methods using NMR-guided MELD (Robertson et al., 2019) and NMR-guided Rosetta (Kuenze and Meiler, 2019) methods. Amazingly at the time, some other CASP13 prediction groups provided pure prediction models, which did not use the NMR data at all, that also matched the reference structures better than the models generated with the data by expert data analysis (Sala et al., 2019). Among the top performing groups in this category were machine learning methods including AlphaFold2.

Even more exciting results came in late 2020 from the CASP14 blind protein structure prediction experiment. In CASP14, the next-generation AlphaFold2 methods demonstrated outstanding performance in protein structure prediction (Kryshtafovych et al., 2021; Pereira et al., 2021). Interesting results were observed for three CASP14 targets for which NMR data were available (Huang et al., 2021). For two of these, the pure prediction AlphaFold2 models were observed to fit real experimental NMR data as well or better than the reference structures provided by the experimental NMR groups. In a third case, target T1027, a protein exhibiting spectral properties indicating extensive conformational dynamics, the AlphaFold2 model did not fit the NMR data as well as the reference experimental NMR structure. The AlphaFold2 prediction model of a fourth NMR target, the 238-residue integral membrane protein MipA, also was an excellent fit to the experimental NMR data. This study also demonstrated how an AlphaFold2 model of CASP14 target T1029 could be used to guide reanalysis of the experimental NMR NOESY data to provide a revised experimental structure which better fit other NMR structure quality assessment metrics, including residual dipolar coupling (RDC) Q scores, and ANSURR scores. The conclusions of this CASP14 - NMR study, using blind predictions for targets not available to the prediction groups, are supported by two other recent studies of the modeling accuracy of AlphaFold, using as reference either previously deposited NMR structure coordinates (Zweckstetter, 2021), or X-ray crystal structures and experimental RDC data (Robertson et al., 2021).

The Protein Structure Validation Software suite (PSVS) (Bhattacharya et al., 2007) integrates multiple tools for protein structure validation, with a particular focus on protein NMR structure validation. PSVS provides several *knowledge-based protein structure validation* tools, including Molprobity clash and Ramachandran backbone analyses (Lovell et al., 2003; Chen et al., 2010), as well as Verify3D (Luthy et al., 1992) and ProsaII (Sippl, 1993) protein fold analysis tools. It also provides a *model vs data protein structure validation* analysis using the PDBStat (Tejero et al., 2013) software for distance and dihedral angle restraint violation analysis, supporting several common distance restraint formats and providing conversion of distance and dihedral angle restraints between these formats to allow interoperability between various NMR structure modeling software packages. Recently, PSVS has been updated (version 2.0) to include additional model vs data structure validation tools, including the RPF-DP score (Huang et al., 2005; Huang et al., 2012) comparing structure models against NOESY peak list data, RDC Q scores (Cornilescu et al., 1998; Clore and Garrett, 1999) comparing models against RDC data, and the ANSURR (Accuracy of NMR Structures Using RCI and Rigidity) score (Fowler et al., 2020) comparing measures of conformational rigidity for various regions of protein structure models with metrics of rigidity based on backbone chemical shift data.

Here, we test the hypothesis, based on our experiences in CASP13 and CASP14, that small protein structures modeled using the recently released AlphaFold platform fit experimental NMR NOE, RDC, and structural rigidity (chemical shift) data as well as models generated by experts using conventional NMR data analysis methods. Six proteins solved by expert NMR spectroscopists in the course of the PSI Structural Genomics Initiative were modeled by AlphaFold and then assessed against both experimental NMR data and knowledge-based statistical metrics using the PSVS software suite. AlphaFold modeling was done by excluding information from the deposited structure itself, or from any homologous templates, as input information. In all cases, the various PSVS structure quality scores for AlphaFold models document that these predicted structures fit the NMR data as well, or often better, than experimental structures deposited in the Protein Data Bank by expert spectroscopists. Overall, this study demonstrates the outstanding value of AlphaFold for modeling small, relatively rigid protein structures, and for providing atomic coordinates useful in guiding analysis of experimental data.

Methods
NMR and X-ray structure coordinate sets and NMR data. Experimentally-determined protein structure coordinates and NMR data were taken for proteins deposited in the Protein Data Bank (PDB) by the Northeast Structural Genomics Consortium (Montelione et al., 2013). For this study we used small proteins solved by both NMR and X-ray crystallography methods, and for which both nearly complete resonance assignment and NOESY peak list data are publicly available from the *NESG NMR / X-ray Pairs* web site (Everett et al., 2016) (https://montelionelab.chem.rpi.edu/databases/nmrdata/). For three of these proteins, ^{15}N-^{1}H residual dipolar coupling data are also available from this site. The proteins ranged in size from 58 to 158 residues (excluding short hexa-His purification tags). These same atomic coordinates and most of these NMR data are also available in the PDB and/or the BioMagResDataBase (BMRB). In addition, structures of these same six proteins that have been energy-refined using the NMR-restrained Rosetta refinement protocol (Mao et al., 2014) were also obtained from the *NESG / NMR X-ray Pair* web site. For the two structures for which RDCs were used in the original structures deposited in the PDB (SrG115C and RpR324), the Rosetta refinement was carried out with these RDC data. For a third target (SgR209C) RDC data is also available in the NESG database, but it was obtained only after the structure was deposited in the PDB, and was not used in the original structure determination nor in the Rosetta refinement.

Solution NMR structure determinations. For two cases required for this study, target protein structures were redetermined from the original NMR data following the standard methods of the NESG consortium (Liu et al., 2005; Montelione and Szyperski, 2010). The solution NMR structure ensemble for RpR324 was recalculated using previously described NMR data (Ramelot et al., 2012), including the resonance assignments, NOE, dihedral, and hydrogen bond restraints of PDB entry 2LPK, but excluding the RDC data, using CYANA (Güntert and Buchner, 2015) followed by refinement with CNS in explicit water. The structure of target SgR209C, PDB entry 2LO6 determined without RDC data, was redetermined using the NOE, dihedral, and hydrogen bond restraints from PDB entry but also including ^{15}N-^{1}H RDC RDC data measured on samples partially aligned in polyacrylamide stretched gel (PAG) and polyethylene glycol (PEG) alignment media that were not used in the original structure determination. These RDC data were also obtained from the *NESG NMR / Xray Pairs* web site. RDC data were manually assessed and excluded if they had peak overlap in the ^{1}H-^{15}N 2D plane or were in disordered regions of the protein. This resulted in 65/90 PAG and 69/103 PEG RDCs used in the refinement, carried out using the same standard methods outlined above for RpR324.

AlphaFold (AF) modeling. AF modeling was carried out using the AF-multimer software (Evans et al., 2021)installed on the NPL cluster in the Center for Computational Innovation at Rensselaer Polytechnic Institute. The system has 40 nodes each with 2x 20 core 2.5 GHz Intel Xeon Gold 6248 CPUs and 8x NVIDIA Tesla V100 GPUs, each with 32 GiB HBM, with 768 GiB RAM per node. This version of AF was trained using the PDB database of April, 2018 and did not use any NMR structures in the training data (Jumper et al., 2021a). PDB structures deposited after November 2005, including the subject structures themselves and the structures of any homologs detected with HMMPred (Soding et al., 2005) were excluded as modeling templates. X-ray crystal and NMR structures of the query target proteins were also excluded as templates. Hydrogen atoms were added by the AF prediction pipeline to each model using restrained refinement of the H atom positions with Amber (Case et al., 2021) with fixed heavy atom coordinates. The AlphaFold model was represented by five top-scored conformations along with estimates of prediction reliability, as described elsewhere (Jumper et al., 2021a).

Knowledge-based protein structure model validation. All structure quality statistical analyses were performed using the Protein Structure Validation Software (Bhattacharya et al., 2007) (PSVS ver 2.0-pre) (https://montelionelab.chem.rpi.edu/PSVS/PSVS2/). PSVS runs a suite of knowledge-based software tools including PDBStat (ver 5.21.6) (Tejero et al., 2013), ProCheck (ver 3.5.4) (Laskowski et al., 1993), MolProbity (mage ver 6.35.040409) (Chen et al., 2010), and an implementation of the algorithms of FindCore2 (Snyder et al., 2014) coded in PDBStat. The structure validation scores of these programs were used to calculate a normalized Z score relative to mean values and standard deviations obtained for a set of 252 reference X-ray crystal structures of < 500 residues, resolution < 1.80 Å, R-factor < 0.25 and R-free < 0.28; positive Z scores indicate 'better' scores.

RDC Q score analysis. ^{15}N-^{1}H residual dipolar couplings D_{calc} were calculated from model structures by single-value decomposition of the Saupe matrix (Losonczi et al., 1999) using PDBStat (Tejero et al., 2013), called from PSVS ver 2.0. Residual dipolar coupling Q scores were analyzed by PDBstat using both of the following methods. The most commonly used RDC-fit score Q1, described by Cornilescu et al (Cornilescu et al., 1998) is

$$Q1 = \sqrt{\frac{\sum_{i=1}^{N}(D_{i,exp} - D_{i.calc})^2}{\sum_{i=1}^{N}(D_{i,exp})^2}} \quad \text{Eqn 1}$$

where D_{exp} and D_{calc} are the measured and calculated values of the RDC, and N is the number of RDCs assessed. In addition, we also assessed models using RDC-fit score Q2, described by Clore et al (Clore and Garrett, 1999) and used by the DC: Servers for Dipolar Coupling Calculations (https://spin.niddk.nih.gov/bax/nmrserver/dc/svd.html).

$$Q2 = \sqrt{\frac{\sum_{i=1}^{N}(D_{i,exp} - D_{i.calc})^2}{N[D_a^2(4 + 3R_h^2)/5]}} \quad \text{Eqn 2}$$

where D_a is the axial component of the orientation tensor, and R_h is the rhombicity of the tensor. The Q2 score is preferable in case of a limited RDC sampling over all possible orientations (Clore and Garrett, 1999).

RPF-DP scores. RPF-DP scores are a set of fast and sensitive structure quality assessment measures which can be used to evaluate how well a 3D structure model fits with NOESY peak and resonance assignment lists, used to assess the correctness of the fold and accuracy of the structure (Huang et al., 2005; Huang et al., 2012). RPF-DP scores provide a type of NMR "R-factor", in which models are compared against unassigned NMR NOESY data. In this work, RPF-DP scores were computed with the program RPF called from PSVS. An RPF server is available online at https://montelionelab.chem.rpi.edu/rpf/. RPF-DP metrics have been described previously (Huang et al., 2005; Huang et al., 2012; Huang et al., 2021), but as they play a key role in this work, we provide a brief overview of these model vs data structure quality assessment metrics here; additional details are provided in the original paper (Huang et al., 2005).

The RPF-DP score algorithm is outlined schematically in Figure 1, adopted from Huang et al (Huang et al., 2005; Huang et al., 2012; Huang et al., 2021). Nodes represent all protons listed in the resonance assignment table. Edges connect the nodes and represent all potential associated NOEs from the NOESY peak lists, within a chemical shift match tolerance. In constructing G_{ANOE} (shown on right side of Figure 1) each NOESY cross peak (p) may be ambiguously assigned to one or more proton pairs, as determined by chemical shift degeneracies and match tolerances. The solution network, G_{NOE}, corresponding to the true 3D structure, is a subgraph of G_{ANOE}. Given complete NOESY peak lists and resonance assignments, for each NOESY cross peak p, at least one of its possible proton pair assignments has a corresponding edge in G_{NOE}. For each structure model (shown on left side of Figure 1), a distance network G is calculated from the summation distances (sum of inverse sixth powers of individual degenerate proton-proton distances), assuming uniform effects of nuclear relaxation processes. Nodes are connected by an edge in G if the corresponding interproton summation distance in the model structure is $\leq d_{NOE\_max}$, where d_{NOE_max} is the (estimated) maximum distance detected in the NOESY spectrum. Summation distances are used to address the lack of stereospecific assignments of prochiral methylene proton pairs, sets of protons that are degenerate (e.g. the three hydrogens of a methyl group, degenerate methylene protons, or degenerate resonances of Tyr or Phe), or combinations of these kinds of ambiguities (e.g. for prochiral isopropyl methyl groups of Leu or Val for which stereospecific assignments are not available). The default upper-bound observed distance, d_{NOE_max}, used in these metrics is 5 Å, but this can also be calibrated from the NOESY data. For models derived by X-ray

crystallography, protons are added with ideal covalent geometry using the program Reduce v2.14 (Chen et al., 2010).

As illustrated in Figure 1, proton pair short distances present in the atomic coordinates of a model structure, represented by the network G, may or may not be represented in the graphical representation of the NOESY peak list data G_{ANOE}. NOESY cross peaks represented in G_{ANOE} that are consistent with the short interproton distances in the network derived from the model, G, are defined as true positives (TPs), while NOESY peaks expected from the model (edges in G) that are not observed in the data, G_{ANOE}, are false positives (FPs). Since G_{ANOE} is an ambiguous network, a FN score is assigned to a NOESY peak only if none of the several possible short proton-proton distance consistent with the possible NOESY peak assignments are observed in G. In this context, recall (R) measures the fraction of NOE cross peaks that are consistent with the query model structures, while precision (P) measures the fraction of short proton pair distances in the query structure that are observed in the NOESY peak list (i.e., in G_{ANOE}), weighted by interproton distance to minimize the impact of weak NOEs arising from interproton distances near the edge of the defined distance cutoff (Huang et al., 2005). Hence "recall violations" are NOESY peaks (or "noise peaks") that cannot be explained by the model and resonance assignments, and "precision violations" are short distances in the model that are not supported by NOESY data, which can result from overpacking, incorrect sequence-specific assignments, or from exchange broadening. The F-measure is the harmonic mean of the recall and precision. Equations used to calculate recall (R), precision (P), and F-measure (F, also called the performance) are shown in Figure 1. The DP score is a scaled F-measure that accounts for lower-bound and upper-bound values of the F-measure. The lower-bound of F(G) is estimated by $F(G_{free})$, where G_{free} is a distance network graph computed from interproton distances in a freely rotating polypeptide chain model, as described by Flory and co-workers (Flory, 1969), and the upper-bound of F(G) is determined by assessing the completeness of the NOESY peak list data for the 3- and 4-bond connected H atoms which all have interproton distances < 5 Å.

RPF-DP scores can be calculated either for individual models from an ensemble of conformations, and averaged, or using average interproton distances across the ensemble. The ensemble DP score (<DP>) is usually 10 - 15% higher than average of individual DP scores (DP_{avg}). However, when the conformational ensemble is more diverse, larger differences between DP_{avg} and <DP> are observed. In various studies (Huang et al., 2005; Raman et al., 2010; Huang et al., 2012; Rosato et al., 2012; Rosato et al., 2013; Rosato et al., 2015; Sala et al., 2019; Huang et al., 2021), structures within 2.0 Å RMSD of the corresponding expert-derived "correct" structure have been observed to have <DP> scores > 0.70 for NMR ensembles, and DP_{avg} scores > 0.60, averaged over the individual conformers.

ANSURR scores. The Accuracy of NMR Structures Using RCI and Rigidity (ANSURR) method provides an independent assessment of model accuracy by comparing protein flexibility computed from backbone chemical shifts with protein flexibility predicted with a graph theory based measure of structural rigidity (Fowler et al., 2020). ANSURR provides two measures of similarity between these measures, a correlation score (corr) which assesses the correlation between peaks and troughs of observed and predicted structural flexibility along the sequence, and root-mean-squared deviation (RMSD) between these metrics. Both the corr and RMSD score are reported as a percentile score (ranging from 0 to 100). These scores were calculated using ANSURR program version 1.2.0.

Well-defined residue ranges and Global Distance Test scores. For NMR structure ensembles, the ranges of residues that are "well-defined" were determined by standard conventions instantiated in the programs

Cyrange (Kirchner and Güntert, 2011) and FindCore2 (Snyder et al., 2014). Following the recommendations of the wwPDB NMR Structure Validation Task Force (Montelione et al., 2013), residues were used in superimpositions and structure quality assessment if they are "well-defined" (i.e. well-converged) across the NMR structure ensemble and also "reliably predicted" by AlphaFold, as outline in our previous studies comparing NMR-derived and AlphaFold models (Huang et al., 2021). GDT_TS scores were computed using the representative "medoid" structure (Montelione et al., 2013; Tejero et al., 2013) of conformational ensembles of the NMR-derived or AlphaFold models within these "well defined" residue ranges by the method of Zemla (Zemla, 2003; Zhang and Skolnick, 2004). For brevity, GDT_TS scores are referred to throughout this paper as GDT scores.

Molecular Modeling. Molecular visualization and graphical representations for figures was done using *PyMol* (DeLano, 2002).

Results
RPF-DP and ANSURR scores for assessment of prediction and experimental NMR models in CASP14.
In previous studies, we have explored the value of RPF-DP scores in assessing models generated in the Critical Assessment of Protein Structure Prediction (CASP) rounds 13 and 14 (Sala et al., 2019; Huang et al., 2021). When an accurate model is used as the reference structure for computing GDT scores, there is a strong correlation between the DP score and GTD (Figure 2A and 2B). Prediction models from some 100+ prediction groups in each CASP edition, each contributing as many as 5 models, span a wide range of structural accuracy. Some poor prediction models can even provide negative DP scores (Figure 2) meaning that the agreement with the chemical shift and NOESY peak list data is poorer than what would be expected for a random coil polypeptide conformation (Huang et al., 2005). Interestingly, for CASP14 target T1055, the best prediction models (provided by AlphaFold2) have DP scores *higher than the experimental NMR structures* (Huang et al., 2021). For target T1027, the DP score is highly correlated with model prediction accuracy; however the best prediction models (again from AlphaFold2) have DP scores lower than the NMR-derived models. These results are analyzed in detail in Huang et al. (2021). Target T1029 presented an especially instructive case. The initial NMR structure provided for CASP model assessment, T1029_original, had a relatively poor DP score, and when used as a reference model for the GDT analysis resulted in a poor correlation between DP and GDT across 529 prediction models (Figure 2C). Recognizing a potential problem, the experimentalists reassessed their NOESY data for this target, and redetermined the NMR structure. The resulting models have much improved DP scores, marginally better than the AlphaFold2 prediction models, and when used as a reference structure for GDT analysis provide a strong correlation between DP and GDT as expected for an accurate structure (cf Figures 2C and 2D). These results demonstrate the value of DP scores in assessing model accuracy using experimental NOESY data.

The same CASP14 NMR structures were also assessed using ANSURR (Figures 3 and 4). A similar analysis of CASP14 prediction models has also recently been reported by Fowler and Williamson, the developers of ANSURR (Fowler and Williamson, 2022). Plots of ANSURR corr vs rmsd score (Figure 3) demonstrate the power of ANSURR in identifying accurate prediction models. For CASP14 target T1055, ANSURR scores for the AlphaFold2 models (blue dots with red outside circles) are somewhat better than for the experimental NMR structures (green dots with yellow outside circles), consistent with the conclusion of DP analysis. However, many other CASP14 prediction models (with lower GDT to the reference NMR structure) also have very good ANSURR scores (Figure 3A). A similar

conclusion can be made for target T1029 (Figure 3C), where the ANSURR scores clearly distinguish the original NMR structure (green dots with yellow outside circles) from the revised NMR structure (green dots with blue outside circles). In this case the revised NMR structures have somewhat better ANSURR scores than the AlphaFold2 models (blue dots with red outside circles), also consistent with DP analysis. For target T1027 the ANSURR scores are generally higher for the NMR structures than for the AlphaFold2 models (Figure 3B), which is also consistent with the DP analysis of Fig. 2B. However, this target has significant amounts of not-well-defined (Table 1) (apparently flexible) backbone structure (Wu et al., 2020) which can potentially dominate the ANSURR score. This sensitivity of the ANSURR score to removal of not-well-defined regions is evident by comparing Figures 3B (full-length T1027) and 3D (trimmed T1027).

An important feature of the DP analysis is the strong correlation between DP and GDT scores when an accurate model is used as the reference structure for calculating the GDT score. Using these CASP14 data, we also assessed if ANSURR scores provide a similar correlation (Figure 4). Generally speaking, the ANSURR scores (corr + rmsd) do not exhibit a strong correlation with GDT scores for this data set. Indeed, some incorrect prediction models with GDT scores as low as 50 have ANSURR scores similar to those of the best AlphaFold2 and NMR models. For the "trimmed" T1027 models, many inaccurate prediction models have better ANSURR scores than either the AlphaFold2 or NMR models (Figure 4D). Hence, while ANSURR is a powerful and convenient tool for model quality assessment, requiring only backbone chemical shift data, it is important to complement ANSURR scores with other metrics of structural accuracy.

NMR / X-ray pairs used for assessment of AF. Six targets from the *NESG NMR X-ray Pairs* data set (Everett et al., 2016) were selected for which NMR structures were determined using standard methods of the NESG Consortium (Liu et al., 2005; Montelione and Szyperski, 2010), but without any residual dipolar coupling data. These structures were NESG targets (with PDB / BMRB id's and PDB release dates): CtR107 (3E0H X-ray, released 2008-09-30; 2KCU / 16097 NMR, released 2009-01-20)*;* GmR137 (3CWI X-ray, released 2008-05-06; 2K5P / 15844 NMR released 2008-08-26); RpR324 (3LMO X-ray, released 2010-02-16; 2LPK / 18263 NMR, released 2012-02-29); SgR42 (3C4S X-ray, released 2008-02-12; 2JZ2 / 15604 NMR, released 2008-01-22); SgR209C (3OSJ X-ray, released 2010-10-06; 2L06 / 17031 NMR, released 2010-08-25); and SgR115C (3MA5 X-ray, released 2010-04-07; 2KCL / 16084 NMR released, 2009-01-06).

The solution NMR structure of target RpR324 is reported in multiple PDB entries, but all of the apo structures were refined with some RDC data. As part of the aim of this study was to assess the accuracy of AF models compared to NMR structures solved with and without RDC data, the deposited chemical shift and NOESY NMR data for RpR324 (BMRB id 18263) were used to re-determine its structure without any RDC data using our standard NMR structure analysis methods. This structure was deposited in December 2021 as PDB entry 7TZD.

Three of these NESG target proteins were also determined using standard methods that also included refinement with ^{15}N-^{1}H residual dipolar coupling data. For two protein targets, these data were obtained from the *NESG NMR / X-ray Pairs* web site (Everett et al., 2016): RpR324 (2LPK / 18263 NMR with ^{15}N-^{1}H RDCs, released 2012-02-29) and SrG115C (2KCV / 16821 NMR with ^{15}N-^{1}H RDCs, released 2009-01-20). In a third case, for target SgR209C, ^{15}N-^{1}H RDC data were available in our database but were not used in the original PDB deposition. For the purposes of this study, the structure

was re-determined using the original NOESY, dihedral, and hydrogen bond restraint data (BMRB id 17031) together with these RDC data. This structure and RDC data were deposited in December 2021 as PDB entry 7TZ8.

This process provided 9 solution NMR structures (NMR-based models consisting of ensembles of conformers) for six NESG targets, where all six were determined using chemical shift and NOESY data without ^{15}N-^{1}H RDC data, and 3 were determined using, in addition, ^{15}N-^{1}H RDC data. These 9 NMR structures are summarized in Table 1. In all three cases where RDC data is available, the RDCs were measured using two alignment media, either polyethylene glycol (PEG) and PF1 phage, or PEG and polyacrylamide stretched gels (PAG).

Well-defined and not-well-defined regions of models. For structure quality assessment and structural comparisons, it is important to identify "not well defined" regions of the protein NMR structure model; i.e. regions where the multiple models (typically 20) generated in the NMR modeling process do not converge, and hence the coordinates are not reliable. Such regions of the structure model will often also have poor knowledge-based structure quality scores and hence are generally excluded from Procheck, Molprobity packing, and Molprobity Ramachandran outlier analysis. Most importantly, not-well-defined regions of NMR structures should not be used for comparisons with prediction models. For identifying not-well-defined regions of these NMR model ensembles, we compared results of Cyrange (Kirchner and Güntert, 2011) and FindCore2 (Snyder and Montelione, 2005; Snyder et al., 2014). These results were generally consistent, and were used to identify consensus ranges of well-defined residue segments. In comparisons with AlphaFold models, we considered only polypeptide segments which are both well-defined in the solution NMR structures and also reliably predicted in the output of AlphaFold (Jumper et al., 2021b; Jumper and Hassabis, 2022). The unions of these two residue ranges, not-reliable based on AF and not-well-defined based on NMR structure convergence, were then excluded from the NMR, X-ray, and AF coordinates for structure quality assessment and structure comparisons (e.g. GDT scores). The resulting residue ranges of reliably comparable regions are summarized for each protein target in Table 1. For 5 of the targets, this process "trims" N-terminal and/or C-terminal polypeptide segments. However, for target SgR209C, this process also identified two internal polypeptide loops, of several residues each, for which atomic coordinates are not reliably modeled in the NMR and/or AF models, which were exclude from structural comparisons and knowledge-based structure quality assessment statistics.

Modeling with AlphaFold. The six protein targets were modeled with AlphaFold-multimer (Evans et al., 2021) implemented on CCI GPU clusters at RPI, and analyzed with PSVS ver 2.0 (and PDBStat). The resulting well-defined regions of the polypeptide backbone structures are compared for the NMR, AF, and X-ray crystal structures of the six protein targets in Figure 5. In these comparisons, the RDC-refined NMR structure models are shown where available. Generally, the AF, NMR, and X-ray models of these six protein targets are an excellent match. A backbone structural similarity matrix is provided below each superimposed set of backbone structure models (Figure 5). These pairwise comparison scores are between the medoid conformer of each structure ensemble (or the single conformer reported for the X-ray crystal structure). The upper diagonal elements in each matrix are Cα GDT scores, the lower diagonal Cα RMSD scores, and the diagonal values are mean pairwise RMSD between each member of the structure ensemble and the medoid conformer. In most cases, excluding target CtR107, the NMR, X-ray, and AlphaFold models have high pairwise similarity scores (GDT of 81.5 to 99.6) and low backbone RMSDs (0.48 to 1.68 Å). In particular, the AlphaFold models are in excellent agreement with RDC-refined NMR structures (GDT 91.3 – 99.4). For target RpR324, the AlphaFold models are a better match to the NMR

structure, while in the other cases they are a better match to the X-ray crystal structure. For RpR324, the AlphaFold models have high similarity with the NMR structures determined both without (GDT = 92.9, RMSD 1.13 Å) and with (GDT = 99.4, RMSD 0.59 Å) RDCs; the match to the X-ray crystal structure is significantly poorer (GDT = 86.7, RMSD 1.74 Å). The basis of this difference is discussed in more detail below. On the other hand, the AlphaFold model is a relatively poor match to the NMR model of target CtR107 (GDT = 64.7; RMSD 3.89 Å). The NMR model of CtR107 is particularly poorly converged, with backbone RMSD of well-defined regions within the NMR ensemble of 2.11 Å (compared to the five other targets with more typical ensemble backbone RMSDs of 0.4 to 1 Å). The AlphaFold model of CtR107 is, however, an excellent match to the corresponding X-ray crystal structure (GDT = 97.3; RMSD 1.09 Å).

Assessment of NMR and AF models for representative NESG structure using RPF and DP scores. The PSVS v2.0 / PDBStat server was next used to assess <DP> and DP_{avg} metrics (along with R_{avg}, P_{avg} and F_{avg} metrics) for the NMR, AlphaFold, and X-ray models (Table 2). Included in this analysis are the corresponding NMR structures refined using a NMR-data-restrained Rosetta modeling protocol (Mao et al., 2014). Generally, for most models and methods, the <DP> score is > 0.70 and DP_{avg} > 0.60, meeting criteria for good quality models (Huang et al., 2012; Huang et al., 2021). *Remarkably, in most cases the AlphaFold models have DP scores (i.e. how well the model fits the NMR data) similar to, or in some cases better than, the NMR structures generated from these same NOESY peak list data*. For each target, the scores for the best performing method is shown in bold red font. Considering DP_{avg} as the most discriminating score, compared to the NMR-based models the AF models had the best score (or tied for best score) for three of the six targets; for the other three targets the DP_{avg} score for the AF models was only slightly lower (0.1 units) than the best-scoring NMR-based models. For four of the six targets, the AF models fit the NMR data better than the corresponding X-ray crystal structures; for the remaining two targets the DP_{avg} for the AF models is only slightly lower (0.1 units) than for the corresponding X-ray crystal structure. *Hence, AlphaFold, using no sample-specific experimental data, provides models with an accuracy, based on the DP score, as good or better than the experimentally derived NMR or X-ray crystal structures*.

The one outlier in this analysis is, again, target CtR107. Comparing the <DP> and DP_{avg} provides information about how well individual conformers in the NMR ensemble fit the NMR data. The observation that DP_{avg} << <DP> indicates that while the average distances across the (relatively broad) conformer distribution are consistent with the NOESY data, no individual conformer is a good fit to these data. Rather, there appear to be multiple conformations in solution, providing inconsistent NOESY peak data for which no single model is well fit. None of the modeling methods (NMR, AF, or X-ray) provide models with DP_{avg} score > ~ 0.53. This suggests that more powerful ensemble-averaged modeling methods, fitting the data to multiple conformational states, are needed to optimally explain the observed NOESY data obtained for this protein in aqueous solution.

Assessment of NMR and AF models for representative NESG structure using knowledge-based metrics. A necessary, but not sufficient, condition for model accuracy is good knowledge-based structure quality scores. PSVS 2.0 integrates multiple software packages to assess backbone and sidechain dihedral angle distributions using Procheck (Laskowski et al., 1993), and overall packing scores and Ramachandran statistics using MolProbity (Lovell et al., 2003; Chen et al., 2010). ProCheck - backbone and ProCheck - all (backbone plus sidechain) dihedral angle distributions and MolProbity packing scores are normalized to Z scores, such that values of Z > 0 are better quality scores than the mean (Z = 0) obtained for 252 high-resolution X-ray crystal structures (Bhattacharya et al., 2007). Typically, good models have average

Z scores > -3 for these various structure quality scores. For all of the targets and all of the methods, almost all of the models have average Z scores > -3; for many of the targets and methods values of Z are > 0 (Table 2). Ramachandran analysis with Molprobity indicates nearly all backbone dihedral angles are in the allowed and generously allowed phi-psi regions. Generally speaking, the AlphaFold models (and Rosetta-refined NMR models) have excellent knowledge-based dihedral Z score, packing Z scores, and Ramachandran statistics for these metrics; in some cases the AlphaFold models have no Molprobity packing violations at all (in these cases MolProbity does not provide a proper packing score, and the Z score is defined as > +3.5 in Table 2).

Assessment of NMR and AF models for representative NESG structure using ANSURR. The models provided for these targets and methods were also assessed by ANSURR (Table 2). Using both full-length and trimmed (as defined in Table 1) models, the best (or second best) ANSURR scores (corr + rmsd) were returned for the AlphaFold models. The one exception was for target RpR324, where the RDC-refined experimental NMR models returned the highest ANSURR score. It is interesting to observe that ANSURR scores have a wide diversity across the models generated by the different modeling methods used for the same protein target, suggesting that it provides strong structural discrimination.

Assessment of NMR and AF models for representative NESG structure using RDC data. For three of these targets, ^{15}N-^{1}H RDC data was also available, allowing assessment of models against RDC data. These data are plotted in Figure 6, and the resulting analyses provided by PSVS /PDBStat are summarized in Table 3. For five of the six RDC data sets (three targets, each in two RDC alignment media) the NMR structures determined with the RDC data (with or without Rosetta refinement) were the best fit to the RDC data. For all six RDC data sets, the AlphaFold models fit the RDC data better than the NMR-based models generated from NOE data without RDC data. In five of six cases, the AlphaFold models do not fit the RDC data as well as models determined using these RDC data; however in the sixth case, target SgR209C in PEG alignment media, the AlphaFold models have an even better fit to RDC data (lower Q scores) than the corresponding NMR-derived models generated using these RDC data. The same conclusion is demonstrated by linear regression analysis (R^2) of calculated vs observed RDC values (Figure 6). *Overall, the AlphaFold models fit the experimental RDC data better than NMR structures generated without RDC data, and in some cases have RDC Q scores and linear regression (R^2) of calculated vs observed RDC values rivaling those obtained for NMR models refined with these RDC data.*

Discussion
For the nine data sets available for six protein targets, the AlphaFold models have remarkably good fit to the experimental NMR data. Across a wide range of structure validation methods, including both knowledge-based validations of dihedral angle distributions and packing scores, and model vs. data validation against experimental NOESY and RDC data, the AF models have similar, and in some cases, better structure quality scores compared with models generated using conventional structure generation methods in the hands of NMR experts using these same NMR data.

The DP score for assessment of NMR derived models has been used routinely in our laboratory, and by various scientists associated with the Northeast Structural Genomics Consortium, as a primary validation tool since its development as a "NMR R factor" in 2005 (Huang et al., 2005; Huang et al., 2006; Raman et al., 2010; Huang et al., 2012; Rosato et al., 2012; Rosato et al., 2013; Rosato et al., 2015; Sala et al., 2019; Huang et al., 2021) (https://montelionelab.chem.rpi.edu/rpf/). However, despite its sensitivity to model inaccuracies and its power for refining NOESY peak list data (Huang et al., 2012), it

has not been broadly adopted by the protein NMR community. Here we describe incorporation of RPF-DP analysis into the PSVS 2.0 software package and server. Hopefully this will provide broader access to these valuable tools.

In this study we outline the value of using a series of models, generated in this case by the CASP community, to evaluate the correlation between DP and GDT. In this analysis, carried out with good quality NMR data, an accurate reference structure provides a linear correlation (e.g. Fig. 2 A, B, and D), while an inaccurate reference structure provides a poor correlation (Fig 2C). Another valuable insight is provided by comparing the <DP> score, based on the average interproton distance within a model, and DP_{avg}, the average DP score computed for each model in the ensemble. In cases where the ensemble is tight and all the models fit the data, these two metrics are similar. However, as observed for the case of target CtR107, for dynamic systems with NOESY data arising from multiple conformations, no single conformer model explains all of the NOESY peak list data to provide a good DP score, and <DP> >> DP_{avg}. Such dynamic protein structures also challenge currently available deep-learning-based modeling methods like AlphaFold (Huang et al., 2021).

ANSURR scores are particularly powerful for protein structure quality assessment since they utilize backbone NMR resonance assignment data that is easier to obtain than NOESY data, particularly for larger proteins. Chemical shift data are also part of a PDB NMR structure deposition, ensuring that it is available for many NMR structures. The ANSURR scored can be affected significantly by large "not well defined" or disordered regions of the protein structure, as illustrated in Figures 3 and 4. Moreover, we observe that there is not a strong correlation between ANSURR and GDT score for the CASP14 models studied here, potentially complicating the interpretation of the ANSURR score. Many CASP14 prediction models have high ANSURR scores, but low GDT scores. However, the AlphaFold models returned the highest ANSURR scores compared to the experimentally-determined NMR or X-ray structures for most of the cases studied here.

Our analysis of NMR models generated with and without RDC data revealed that for six data sets for three targets, *AlphaFold models fit these RDC data better than the NMR structures determined without RDC data*. In some cases, the AlphaFold models have RDC Q scores rivaling those obtained for NMR structures determined with these same RDC data. These observations strongly support the concept of using AlphaFold models as a starting point for RDC-based structure refinement, without the need for generating NMR NOESY data (Cole et al., 2021).

Analysis of the AF predictions for target RpR324 is of particular interest because the NMR and X-ray crystal structure have notably different structures and also different oligomerization states (Ramelot et al., 2012). A detailed analysis of these structural details is presented in Figure 7. Movement of the α3 helix results in a more 'open' hydrophobic cavity in the X-ray structure than in the NMR 'closed' conformation (Fig 7A). The NMR sample used for this study was largely monomeric, and size-exclusion chromatography with multiple angle light scattering (SEC-MALS) measurements demonstrate less than 13% dimer in solution (Ramelot et al., 2012). Conversely, in the crystal lattice the protein forms a homodimer with an extensive buried interdomain interface. It remains unclear if there is biological significance for dimerization or for the open conformation, although it seems likely that this represents a functionally relevant conformation. We were interested in finding out whether the AlphaFold prediction would match the NMR or X-ray structure more closely and whether the modeling RpR324 as a

homodimer using AlphaFold-multimer would provide a model matching the X-ray crystal structure with a more open conformation.

We found that the monomeric AlphaFold models for RpR324 are an excellent match to NMR structures and NMR data, and are significantly better matches to the NMR structure than to the corresponding X-ray crystal structure (Figure 7A). The GDT_TS and RMSD analysis (Figure 5) shows that the AlphaFold model has best structural agreement with NMR structure 2LPK* refined with ^{15}N-^{1}H RDC data collected with two alignment media (Table 3 and Figure 7A). Based on the RPF-DP analysis, the AlphaFold prediction models agree with the NMR NOESY data just as well as the experimental NMR structure determined using both NOESY and RDC data (2LPK*), and has even better ProCheck and MolProbity clash scores as determined by PSVS analysis (Table 2). The agreement of AlphaFold models with the NMR ^{15}N-^{1}H RDC data as measured by Q scores is significantly better than that observed for the X-ray structure (Table 3). Taken together, it is clear that the AlphaFold models are more similar to the NMR structures than the X-ray structure, even though the AlphaFold AI was trained on a database of only X-ray structures.

The next noteworthy result is that the dimeric AlphaFold model, generated using AlphaFold multimer, correctly matches the dimer interface observed in the protein crystal, giving further support that this dimerization interface may have biological significance, rather than being a crystallization artifact. The resulting AlphaFold-predicted protomer from this dimer has an almost identical structural match with the monomeric AlphaFold model (Figure 7B), which matches closely the conformation in the monomeric NMR structure (7A). Differences between AlphaFold dimer models and the X-ray crystal structure are still apparent with the α3 helix being more open in the X-ray structure, resulting in small alterations at the dimer interface (Fig 7C, D). Hence, even when modeling the dimeric complex, AlphaFold does not predict the open structure observed in the X-ray crystal structure. The idea that the X-ray structure might represent a functional conformation of RpR324 is, however, supported by the expected biological function of this protein as a specialized acyl carrier protein (AcpXL) for the synthesis of very long-chain fatty acids (20-30 carbons). Although covalent modification of RpR324 on residue S37 by attachment of 4'-phosphopantetheine did not result in any significant changes to the NMR structure, it is believed that addition of a very long chain fatty acid to this carrier arm could favor dimerization or expansion of the hydrophobic cavity as observed in the X-ray structure (Ramelot et al., 2012). Modeling of multiple conformational states of proteins, and energy landscapes, remains an important challenge in the field of protein structure modeling.

This structure-function analysis of AcpXL RpR324 illustrates the power of AlphaFold models for developing specific testable hypotheses for driving structural biology research. Deep learning methods like AlphaFold and RosettaFold have the potential to generate novel insights into structure function relationships at an unprecedented rate, and on genomic and pangenomic scales. Considering this sea change in our ability to generate reliable protein structures, it is important to consider how to use these models to enable NMR data analysis, including the development of new methods to refine AlphaFold models against sparse NMR data, use of models in interpreting NMR data in terms of multiple conformational states of proteins, and for "inverse structure determination" (Huang et al., 2021) in which AlphaFold models are used to guide NMR data interpretation.

Conclusions

These studies provide an unambiguous demonstration that AlphaFold can predict structures of small single-domain proteins in solution, without structural templates, with an accuracy rivaling NMR studies. Since the AlphaFold AI has included X-ray crystal structures of these NESG proteins in its training data, it is possible that there is some kind of indirect memorization that enhances performance for these specific targets. However, for many of the protein targets predicted in CASP14, no homologs of known 3D structure were available in the training data used by AlphaFold. AlphaFold models predicted for this study using the platform available in the public domain fit our experimental NMR data (NOEs and RDCs) as well or better than NMR structures generated from these same data by experts using traditional methods. These results contradict the widely held misperception that AlphaFold cannot accurately model solution NMR structures. While AlphaFold and other deep learning methods do not yet have the ability to model multiple alternative conformations of proteins, protein dynamics, and various complex aspects of protein-biomolecular interactions, these models do provide a higher starting point with which to begin structure-dynamic-function studies of proteins.

Acknowledgements

RDC data for NESG proteins were recorded in the laboratory of Prof. James Prestegard. We thank all of the scientists of the NESG consortium who produced samples, determined X-ray crystal and NMR structures, and provided these to the community by deposition in the Protein Data Bank. We also thank Prof. M. Williamson and Dr. N. Fowler, and Drs. K. Banfa, B. Shurina, and G.V.T. Swapna, for helpful discussions and comments on the manuscript, and Sean Collen for his efforts in porting AlphaFold to the NPL CPU/GPU cluster of the RPI Center for Computing Innovation. This research was supported by grants R01-GM120574 and R35-GM141818 from the National Institutes of Health.

Author Contributions

RT, YJH, TAR and GTM analyzed data. RT and YJH developed computer codes. TAR determined experimental NMR structures. All authors contributed in writing and editing the manuscript.

Declaration of Interests

GTM is a founder of Nexomics Biosciences, Inc. This does not represent a conflict of interest for this study.

Table 1. Overview of data for nine solution NMR structures for six protein targets used in this study.

Sample	PDB ID for NMR Structure	Total number of residues[a]	Well-defined residue range[b]	RDC data used for structure determination[c]	Alignment media for RDC measurements
CtR107	2KCU	158	4-158	No	
GmR137	2K5P	70	1-63	No	
RpR324	7TZD	93	4-91	No	
RpR324	2LPK*,d	93	4-91	Yes	PEG / Phage
SgR42	2JZ2	58	1-56	No	
SgR209C	2L06	147	13-38, 47-134, 138-143	No	
SgR209C	7TZ8*	147	13-38, 47-134, 138-143	Yes	PEG / PAG
SrR115C	2KCL	92	2-92	No	
SrR115C	2KCV*	92	2-92	Yes	PEG / Phage

[a] Number of residues in deposited NMR structure, excluding disordered purification tags, if any.

[b] Consensus of well-defined residues in NMR structures determined by Cyrange (Kirchner and Güntert, 2011)) and FindCore2 (Snyder and Montelione, 2005; Snyder et al., 2014), and "reliably modeled" residues determined by AlphaFold.

[c] 'Yes' indicates that RDC data was used in the structure calculation and refinement. 'No' indicates RDC data were not used. RDC data files were taken from *NESG NMR / X-ray Pairs* web site (Everett et al., 2016).

[d] The asterisk (*) designates the structure was refined against ^{15}N-^{1}H RDC data using two alignment media.

Table 2. Structure quality statistics for experimental and predicted protein structures.

Sample	Method	$\langle DP \rangle^a$	DP_{avg}^a	R_{avg}^a	P_{avg}^a	F_{avg}^a	Pro-Check -bb[b]	Pro-Check -all[b]	Mol Probity[b]	Rama-chandran statistics[c]	ANSURR[d] full length	ANSURR[d] trimmed
CtR107	NMR - 2KCU	**0.72**[e]	0.39	0.96	0.83	0.89	− 1.30	− 3.02	− 1.21	96.8 / 3.1	91 ± 20[f]	82 ± 20[f]
	Rosetta	0.70	0.40	0.95	0.84	0.89	− 0.04	**+ 0.41**	+ 0.38	**99.3 / 0.6**	111± 25	104 ± 26
	AF	0.53	0.52	**0.96**	**0.87**	**0.91**	**+ 0.04**	+ 0.12	**+ 1.53**	98.4 / 1.6	**134 ± 6**	**129 ± 5**
	X-ray - 3E0H	-	**0.53**	0.92	0.91	0.91	− 0.24	−0.47	− 0.39	98.7 / 1.3	96	131
GmR137	NMR - 2K5P	**0.86**	0.73	0.98	0.88	0.93	− 1.97	- 2.84	− 0.14	95.0 / 4.6	58 ± 31	35 ± 24
	Rosetta	**0.86**	0.74	0.98	0.88	0.93	− 0.47	+ 0.47	+ 0.72	**98.5 / 1.3**	109 ± 32	75 ± 33
	AF	0.80	**0.78**	**0.98**	**0.90**	**0.94**	− 0.67	-0.89	**> 3.50**[g]	93.7 / 5.7	**127 ± 11**	**82 ± 13**
	X-ray - 3CWI	-	0.73	0.94	0.91	0.93	− 0.51	-0.77	+0.45	97.0 / 3.0	50	50
RpR324	NMR - 7TZD	**0.84**	0.79	0.95	0.89	0.92	+ 0.90	+ 0.24	**+ 1.39**	97.6 /1.3	139 ± 10	124 ± 11
	NMR - 2LPK*	0.82	0.79	**0.96**	0.88	0.92	+ 1.22	+ 0.65	− 0.91	98.4 / 1.4	**154 ± 13**	**142 ± 16**
	Rosetta*	0.82	**0.81**	**0.96**	0.89	0.92	**+ 1.97**	**+ 2.84**	+ 0.78	99.8 / 0.2	137 ± 15	120 ± 10
	AF	0.81	**0.81**	**0.96**	0.89	**0.93**	+ 1.69	+ 1.83	**+ 1.53**	98.8 / 1.2	125 ± 10	132 ± 8
	X-ray - 3LMO	-	0.80	0.95	**0.90**	0.92	+ 1.49	+ 1.60	+1.10	**100.0 / 0.0**	117	116
SgR42	NMR - 2JZ2	**0.75**	0.65	**0.97**	0.83	0.89	− 2.08	− 2.25	− 0.35	96.3 / 3.7	100 ± 33	84 ± 31
	Rosetta	0.73	0.68	**0.97**	0.84	0.90	− 1.14	**− 0.06**	+ 1.31	**99.2 / 0.8**	134 ± 20	116 ± 20
	AF	0.74	0.72	**0.97**	0.85	**0.91**	− 1.10	− 0.12	**> 3.50**[g]	96.9 / 3.1	**152 ± 7**	**139 ± 9**
	X-ray - 3C4S	-	**0.73**	0.96	**0.85**	0.90	**− 0.90**	− 0.47	+ 0.73	98.2 / 1.8	117	117
SgR209C	NMR - 2L06	**0.80**	0.67	**0.97**	0.89	0.92	+ 0.98	+ 0.53	− 1.95	98.1 / 1.7	104 ± 19	79 ± 20
	NMR - 7TZ8*	0.79	0.64	**0.97**	0.87	0.92	+ 0.87	+ 0.41	+ 1.37	96.7 / 2.8	122 ± 16	91 ± 18
	Rosetta	**0.80**	0.69	**0.97**	0.89	**0.93**	+ 1.65	**+ 2.37**	+ 0.04	**99.0 / 1.0**	114 ± 23	82 ± 11
	AF	0.74	**0.73**	**0.97**	0.90	**0.93**	**+ 1.69**	+ 2.13	**> 3.50**[g]	**99.0 / 1.0**	**140 ± 11**	**129 ± 6**
	X-ray - 3OSJ	-	0.70	0.94	**0.91**	**0.93**	+ 1.30	+ 0.83	− 0.39	98.5 / 1.5	133	99
SrR115C	NMR - 2KCL	**0.83**	**0.82**	**0.93**	0.97	**0.95**	+ 1.65	+ 1.30	+ 0.04	97.5 / 2.4	**152 ± 15**	144 ± 15
	NMR - 2KCV*	**0.83**	0.81	**0.93**	0.96	**0.95**	+ 1.85	+ 1.30	− 0.69	97.1 / 2.9	142 ± 17	132 ± 15
	Rosetta*	**0.83**	**0.82**	**0.93**	**0.97**	**0.95**	**+ 2.64**	**+ 3.02**	+ 1.35	98.8 / 1.2	130 ± 7	119 ± 6
	AF	0.82	0.81	0.92	0.97	**0.95**	+ 2.40	+ 2.60	**> 3.50**[g]	**98.9 / 1.1**	150 ± 6	**144 ± 6**
	X-ray - 3MA5	-	0.80	0.91	0.98	0.94	+ 0.67	− 1.18	**> 3.50**[g]	93.2 / 6.8	108	108

[a] The $\langle DP \rangle$ score is the DP score for the average interproton distances in the ensemble, while DP_{avg}, recall (R_{avg}), precision (P_{avg}), and F-measure (F_{avg}) scores are the average value of these metrics assessed individually for each member of the structure ensemble

[b] Z scores relative to the corresponding scores obtained from a data base of 252 high-resolution X-ray crystal structures solved at < 1.8 Å resolution (Bhattacharya et al., 2007).

[c] Percent of backbone residues in allowed / generously allowed regions of Ramachandran map (Lovell et al., 2003).

[d] ANSURR scores are the sum of ANSURR corr plus rmsd scores (Fowler et al., 2020), where the highest potential score is 200.00. ANSURR scores are reported for both the full-length protein (including disordered regions) and for the trimmed coordinates that exclude the not-well-defined regions of the protein structure as indicated in Table 1.

[e] For each metric and target, the best score (or tied best score) is indicated in red and bold font.

[f] As ANSUUR scores for each ensemble span a wide range, the ANSURR scores are reported ± standard deviation.

[g] MolProbity does not return a clash score for models with zero clashes; these are assigned a Z score of > 3.50.

Table 3. ^{15}N-^{1}H RDC Q Scores for experimental and AF models[a]

Sample	Method	Q1[b]	Q2[c]
RpR324 PEG	NMR - 7TZD	0.383 ± 0.022	0.474 ± 0.036
	NMR - 2LPK*,[d]	**0.122 ± 0.011[e]**	**0.144 ± 0.014[e]**
	Rosetta*	0.323 ± 0.016	0.364 ± 0.023
	AF	0.283 ± 0.021	0.331 ± 0.032
	X-ray - 3LMO	0.395	0.464
RpR324 Phage	NMR - 7TZD	0.454 ± 0.042	0.560 ± 0.074
	NMR - 2LPK*	0.225 ± 0.014	**0.239 ± 0.016**
	Rosetta*	**0.220 ± 0.015**	0.245 ± 0.018
	AF	0.227 ± 0.010	0.253 ± 0.012
	X-ray - 3LMO	0.459	0.564
SrR115C PEG	NMR - 2KCL	0.430 ± 0.024	0.368 ± 0.033
	NMR - 2KCV*	**0.135 ± 0.047**	**0.111 ± 0.043**
	Rosetta*	0.292 ± 0.019	0.257 ± 0.025
	AF	0.357 ± 0.007	0.327 ± 0.010
	X-ray - 3MA5	0.378	0.355
SrR115C Phage	NMR - 2KCL	0.250 ± 0.014	0.319 ± 0.022
	NMR - 2KCV*	**0.162 ± 0.016**	**0.205 ± 0.020**
	Rosetta*	0.170 ± 0.013	0.218 ± 0.022
	AF	0.235 ± 0.006	0.321 ± 0.010
	X-ray - 3MA5	0.275	0.348
SgR209C PEG	NMR - 2L06	0.471 ± 0.057	0.489 ± 0.070
	NMR - 7TZ8*	0.276 ± 0.036	0.277 ± 0.038
	Rosetta	0.499 ± 0.053	0.546 ± 0.078
	AF	**0.256 ± 0.009**	**0.266 ± 0.012**
	X-ray - 3OSJ	0.272	0.307
SgR209C PAG	NMR - 2L06	0.503 ± 0.065	0.433 ± 0.073
	NMR - 7TZ8*	**0.243 ± 0.030**	**0.191 ± 0.028**
	Rosetta	0.535 ± 0.061	0.501 ± 0.094
	AF	0.371 ± 0.010	0.303 ± 0.012
	X-ray - 3OSJ	0.323	0.273

[a] ^{15}N-^{1}H RDC Q scores were computed for "trimmed" residue ranges shown in Table 1.

[b] RDC Q1 scores computed by the method of Cornilescu et al (Cornilescu et al., 1998).

[c] RDC Q2 scores computed by the method of Clore et al. (Clore and Garrett, 1999).

[d] For each RDC Q score assessment in each alignment media, the NMR structures refined with the RDC data are indicated with an asterisk (*). NMR-restrained Rosetta refinement was carried out with ^{15}N-^{1}H RDC data for targets SrR115C and RpR324, but did not use RDC data for target SgR209C

[e] The best (lowest) score for each target and method is indicated in red and bold font.

References

Anishchenko, I., Pellock, S.J., Chidyausiku, T.M., Ramelot, T.A., Ovchinnikov, S., Hao, J., et al. (2021). De novo protein design by deep network hallucination. *Nature* 600(7889)**,** 547-552. doi:https://doi.org/10.1038/s41586-021-04184-w.

Baek, M., Anishchenko, I., Park, H., Humphreys, I.R., and Baker, D. (2021a). Protein oligomer modeling guided by predicted interchain contacts in CASP14. *Proteins* 89(12)**,** 1824-1833. doi:https://doi.org/10.1002/prot.26197.

Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G.R., et al. (2021b). Accurate prediction of protein structures and interactions using a three-track neural network. *Science*. doi:https://doi.org/10.1126/science.abj8754.

Bhattacharya, A., Tejero, R., and Montelione, G.T. (2007). Evaluating protein structures determined by structural genomics consortia. *Proteins* 66(4)**,** 778-795. doi:https://doi.org/10.1002/prot.21165.

Buchan, D.W.A., and Jones, D.T. (2018). Improved protein contact predictions with the MetaPSICOV2 server in CASP12. *Proteins* 86 Suppl 1**,** 78-83. doi:https://doi.org/10.1002/prot.25379.

Case, D.A., Kollman, P.A., and al., e. (2021). "Amber". University of California, San Francisco).

Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., et al. (2010). MolProbity: all-atom structure validation for macromolecular crystallography. *Acta crystallographica. Section D, Biological crystallography* 66(Pt 1)**,** 12-21. doi:https://doi.org/10.1107/S0907444909042073.

Clore, G.M., and Garrett, D.S. (1999). R-factor, free R, and complete cross-validation for dipolar coupling refinement of NMR structures. *J. Amer. Chem. Soc.* 121(39)**,** 9008-9012.

Cole, C.A., Daigham, N.S., Liu, G., Montelione, G.T., and Valafar, H. (2021). REDCRAFT: A computational platform using residual dipolar coupling NMR data for determining structures of perdeuterated proteins in solution. *PLoS Comput Biol* 17(2)**,** e1008060. doi:https://doi.org/10.1371/journal.pcbi.1008060.

Colman, D.R., Labesse, G., Swapna, G.V.T., Stefanakis, J., Montelione, G.T., Boyd, E.S., et al. (2022). Structural Evolution of the Ancient Enzyme, Dissimilatory Sulfite Reductase. *bioRxiv***,** 2021.2012.2028.474277. doi:https://doi.org/10.1101/2021.12.28.474277.

Cornilescu, G., Marquardt, J.L., Ottiger, M., and Bax, A. (1998). Validation of protein structure from anisotropic carbonyl chemical shifts in a dilute liquid crystalline phase. *J. Amer. Chem. Soc.* 120(27)**,** 6836-6837.

DeLano, W.L. (2002). The PyMOL Molecular Graphics System, Version 1.5.0.4 Schrödinger, LLC. *The PyMOL Molecular Graphics System, Version 1.5.0.4 Schrödinger, LLC.*

Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., et al. (2021). Protein complex prediction with AlphaFold-Multimer. *bioRxiv***,** 2021.2010.2004.463034. doi:https://doi.org/10.1101/2021.10.04.463034.

Everett, J.K., Tejero, R., Murthy, S.B., Acton, T.B., Aramini, J.M., Baran, M.C., et al. (2016). A community resource of experimental data for NMR / X-ray crystal structure pairs. *Protein Sci* 25(1)**,** 30-45. doi:https://doi.org/10.1002/pro.2774.

Flory, P.J. (1969). *Statistical Mechanics of Chain Molecules.* New York: Interscience Publishers.

Fowler, N.J., Sljoka, A., and Williamson, M.P. (2020). A method for validating the accuracy of NMR protein structures. *Nat Commun* 11(1)**,** 6321. doi:https://doi.org/10.1038/s41467-020-20177-1.

Fowler, N.J., and Williamson, M.P. (2022). The accuracy of protein structures in solution determined by AlphaFold and NMR. *bioRxiv***,** 2022.2001.2018.476751. doi:https://doi.org/10.1101/2022.01.18.476751.

Güntert, P., and Buchner, L. (2015). Combined automated NOE assignment and structure calculation with CYANA. *J Biomol NMR* 62(4)**,** 453-471. doi:https://doi.org/10.1007/s10858-015-9924-9.

Huang, Y.J., Powers, R., and Montelione, G.T. (2005). Protein NMR recall, precision, and F-measure scores (RPF scores): structure quality assessment measures based on information retrieval statistics. *Journal of the American Chemical Society* 127(6)**,** 1665-1674. doi:https://doi.org/10.1021/ja047109h.

Huang, Y.J., Rosato, A., Singh, G., and Montelione, G.T. (2012). RPF: a quality assessment tool for protein NMR structures. *Nucleic Acids Research* 40(Web Server issue)**,** W542-546. doi:https://doi.org/10.1093/nar/gks373.

Huang, Y.J., Tejero, R., Powers, R., and Montelione, G.T. (2006). A topology-constrained distance network algorithm for protein structure determination from NOESY data. *Proteins* 62(3)**,** 587-603. doi:https://doi.org/10.1002/prot.20820.

Huang, Y.J., Zhang, N., Bersch, B., Fidelis, K., Inouye, M., Ishida, Y., et al. (2021). Assessment of prediction methods for protein structures determined by NMR in CASP14: Impact of AlphaFold2. *Proteins* 89(12)**,** 1959-1976. doi:https://doi.org/10.1002/prot.26246.

Huang, Z., Wang, X., Huang, L., Huang, C., Wei, Y., and W., L. (Year). "CCNet: Criss-Cross Attention for Semantic Segmentation", in: *Proceedings of the IEEE/CVF International Conference on Computer Vision*).

Humphreys, I.R., Pei, J., Baek, M., Krishnakumar, A., Anishchenko, I., Ovchinnikov, S., et al. (2021). Computed structures of core eukaryotic protein complexes. *Science* 374(6573)**,** eabm4805. doi:https://doi.org/10.1126/science.abm4805.

Jones, D.T., and Thornton, J.M. (2022). The impact of AlphaFold2 one year on. *Nat Methods* 19(1)**,** 15-20. doi:https://doi.org/10.1038/s41592-021-01365-3.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., et al. (2021a). Highly accurate protein structure prediction with AlphaFold. *Nature*. doi:https://doi.org/10.1038/s41586-021-03819-2.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., et al. (2021b). Applying and improving AlphaFold at CASP14. *Proteins* 89(12)**,** 1711-1721. doi:https://doi.org/10.1002/prot.26257.

Jumper, J., and Hassabis, D. (2022). Protein structure predictions to atomic accuracy with AlphaFold. *Nat Methods* 19(1)**,** 11-12. doi:https://doi.org/10.1038/s41592-021-01362-6.

Kirchner, D.K., and Güntert, P. (2011). Objective identification of residue ranges for the superposition of protein structures. *BMC Bioinformatics* 12**,** 170. doi:https://doi.org/10.1186/1471-2105-12-170.

Kryshtafovych, A., Schwede, T., Topf, M., Fidelis, K., and Moult, J. (2021). Critical assessment of methods of protein structure prediction (CASP)-Round XIV. *Proteins* 89(12)**,** 1607-1617. doi:https://doi.org/10.1002/prot.26237.

Kuenze, G., and Meiler, J. (2019). Protein structure prediction using sparse NOE and RDC restraints with Rosetta in CASP13. *Proteins* 87(12)**,** 1341-1350. doi:https://doi.org/10.1002/prot.25769.

Laskowski, R.A., MacArthur, M.W., Moss, D.S., and Thornton, J.M. (1993). PROCHECK - a Program to Check the Stereochemical Quality of Protein Structures. *Journal of Applied Crystallography* 26**,** 283-291.

Liu, G., Shen, Y., Atreya, H.S., Parish, D., Shao, Y., Sukumaran, D.K., et al. (2005). NMR data collection and analysis protocol for high-throughput protein structure determination. *Proc Natl Acad Sci U S A* 102(30)**,** 10487-10492. doi:https://doi.org/10.1073/pnas.0504338102.

Losonczi, J.A., Andrec, M., Fischer, M.W., and Prestegard, J.H. (1999). Order matrix analysis of residual dipolar couplings using singular value decomposition. *J Magn Reson* 138(2)**,** 334-342. doi:https://doi.org/10.1006/jmre.1999.1754.

Lovell, S.C., Davis, I.W., Arendall, W.B., 3rd, de Bakker, P.I., Word, J.M., Prisant, M.G., et al. (2003). Structure validation by Calpha geometry: phi,psi and Cbeta deviation. *Proteins* 50(3)**,** 437-450. doi:https://doi.org/10.1002/prot.10286.

Luthy, R., Bowie, J.U., and Eisenberg, D. (1992). Assessment of protein models with three-dimensional profiles. *Nature* 356(6364)**,** 83-85. doi:https://doi.org/10.1038/356083a0.

Mao, B., Tejero, R., Baker, D., and Montelione, G.T. (2014). Protein NMR structures refined with Rosetta have higher accuracy relative to corresponding X-ray crystal structures. *J Am Chem Soc* 136(5)**,** 1893-1906. doi:https://doi.org/10.1021/ja409845w.

Marks, D.S., Colwell, L.J., Sheridan, R., Hopf, T.A., Pagnani, A., Zecchina, R., et al. (2011). Protein 3D structure computed from evolutionary sequence variation. *PloS one* 6(12)**,** e28766. doi:https://doi.org/10.1371/journal.pone.0028766.

Marks, D.S., Hopf, T.A., and Sander, C. (2012). Protein structure prediction from sequence variation. *Nature biotechnology* 30(11)**,** 1072-1080. doi:https://doi.org/10.1038/nbt.2419.

Mondal, A., Swapna, G.V.T., Hao, J., Ma, L., Roth, M.J., Montelione, G.T., et al. (2022). Structure determination of protein-peptide complexes from NMR chemical shift data using MELD. *bioRxiv*, 2021.2012.2031.474671. doi:https://doi.org/10.1101/2021.12.31.474671.

Montelione, G.T., Nilges, M., Bax, A., Guntert, P., Herrmann, T., Richardson, J.S., et al. (2013). Recommendations of the wwPDB NMR Validation Task Force. *Structure* 21(9)**,** 1563-1570. doi:https://doi.org/10.1016/j.str.2013.07.021.

Montelione, G.T., and Szyperski, T. (2010). Advances in protein NMR provided by the NIGMS Protein Structure Initiative: impact on drug discovery. *Curr Opin Drug Discov Devel* 13(3)**,** 335-349.

Morcos, F., Pagnani, A., Lunt, B., Bertolino, A., Marks, D.S., Sander, C., et al. (2011). Direct-coupling analysis of residue coevolution captures native contacts across many protein families. *Proceedings of the National Academy of Sciences of the United States of America* 108(49)**,** E1293-1301. doi:https://doi.org/10.1073/pnas.1111471108.

Ovchinnikov, S., Kinch, L., Park, H., Liao, Y., Pei, J., Kim, D.E., et al. (2015). Large-scale determination of previously unsolved protein structures using evolutionary information. *Elife* 4**,** e09248. doi:https://doi.org/10.7554/eLife.09248.

Ovchinnikov, S., Park, H., Kim, D.E., Liu, Y., Wang, R.Y., and Baker, D. (2016). Structure prediction using sparse simulated NOE restraints with Rosetta in CASP11. *Proteins* 84 Suppl 1**,** 181-188. doi:https://doi.org/10.1002/prot.25006.

Pereira, J., Simpkin, A.J., Hartmann, M.D., Rigden, D.J., Keegan, R.M., and Lupas, A.N. (2021). High-accuracy protein structure prediction in CASP14. *Proteins*. doi:https://doi.org/10.1002/prot.26171.

Raman, S., Huang, Y.J., Mao, B., Rossi, P., Aramini, J.M., Liu, G., et al. (2010). Accurate automated protein NMR structure determination using unassigned NOESY data. *J Am Chem Soc* 132(1)**,** 202-207. doi:https://doi.org/10.1021/ja905934c.

Ramelot, T.A., Rossi, P., Forouhar, F., Lee, H.W., Yang, Y., Ni, S., et al. (2012). Structure of a specialized acyl carrier protein essential for lipid A biosynthesis with very long-chain fatty acids in open and closed conformations. *Biochemistry* 51(37)**,** 7239-7249. doi:https://doi.org/10.1021/bi300546b.

Robertson, A.J., Courtney, J.M., Shen, Y., Ying, J., and Bax, A. (2021). Concordance of X-ray and AlphaFold2 Models of SARS-CoV-2 Main Protease with Residual Dipolar Couplings Measured in Solution. *J Am Chem Soc* 143(46)**,** 19306-19310. doi:https://doi.org/10.1021/jacs.1c10588.

Robertson, J.C., Nassar, R., Liu, C., Brini, E., Dill, K.A., and Perez, A. (2019). NMR-assisted protein structure prediction with MELDxMD. *Proteins* 87(12)**,** 1333-1340. doi:https://doi.org/10.1002/prot.25788.

Rosato, A., Aramini, J.M., Arrowsmith, C., Bagaria, A., Baker, D., Cavalli, A., et al. (2012). Blind testing of routine, fully automated determination of protein structures from NMR data. *Structure* 20(2)**,** 227-236. doi:https://doi.org/10.1016/j.str.2012.01.002.

Rosato, A., Tejero, R., and Montelione, G.T. (2013). Quality assessment of protein NMR structures. *Curr Opin Struct Biol* 23(5)**,** 715-724. doi:https://doi.org/10.1016/j.sbi.2013.08.005.

Rosato, A., Vranken, W., Fogh, R.H., Ragan, T.J., Tejero, R., Pederson, K., et al. (2015). The second round of Critical Assessment of Automated Structure Determination of Proteins by NMR: CASD-NMR-2013. *J Biomol NMR* 62(4)**,** 413-424. doi:https://doi.org/10.1007/s10858-015-9953-4.

Sala, D., Huang, Y.J., Cole, C.A., Snyder, D.A., Liu, G., Ishida, Y., et al. (2019). Protein structure prediction assisted with sparse NMR data in CASP13. *Proteins* 87(12)**,** 1315-1332. doi:https://doi.org/10.1002/prot.25837.

Sippl, M.J. (1993). Recognition of errors in three-dimensional structures of proteins. *Proteins* 17(4)**,** 355-362.

Snyder, D.A., Grullon, J., Huang, Y.J., Tejero, R., and Montelione, G.T. (2014). The expanded FindCore method for identification of a core atom set for assessment of protein structure prediction. *Proteins* 82 Suppl 2**,** 219-230. doi:https://doi.org/10.1002/prot.24490.

Snyder, D.A., and Montelione, G.T. (2005). Clustering algorithms for identifying core atom sets and for assessing the precision of protein structure ensembles. *Proteins* 59(4)**,** 673-686. doi:https://doi.org/10.1002/prot.20402.

Soding, J., Biegert, A., and Lupas, A.N. (2005). The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Res* 33(Web Server issue)**,** W244-248. doi:https://doi.org/10.1093/nar/gki408.

Tejero, R., Snyder, D., Mao, B., Aramini, J.M., and Montelione, G.T. (2013). PDBStat: a universal restraint converter and restraint analysis software package for protein NMR. *J Biomol NMR* 56(4)**,** 337-351. doi:https://doi.org/10.1007/s10858-013-9753-7.

Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A.N., et al. (2017). "Attention is all you need", in: *Proceedings of the 31st International Conference on Neural Information Processing Systems.* (Long Beach, California, USA: Curran Associates Inc.).

Wu, N., Kobayashi, N., Tsuda, K., Unzai, S., Saotome, T., Kuroda, Y., et al. (2020). Solution structure of Gaussia Luciferase with five disulfide bonds and identification of a putative coelenterazine binding cavity by heteronuclear NMR. *Sci Rep* 10(1)**,** 20069. doi:https://doi.org/10.1038/s41598-020-76486-4.

Zemla, A. (2003). LGA: A method for finding 3D similarities in protein structures. *Nucleic Acids Res* 31(13)**,** 3370-3374. doi:https://doi.org/10.1093/nar/gkg571.

Zhang, Y., and Skolnick, J. (2004). Scoring function for automated assessment of protein structure template quality. *Proteins: Structure, Function, and Bioinformatics* 57(4)**,** 702-710.

Zweckstetter, M. (2021). NMR hawk-eyed view of AlphaFold2 structures. *Protein Sci* 30(11)**,** 2333-2337. doi:https://doi.org/10.1002/pro.4175.

Fig. 1

Fig. 1. Schematic description of RPF-DP scores. In this analysis, the graph G with nodes corresponding to all ^{1}H's and edges representing all short (< 5 Å) ^{1}H-^{1}H distances in a structure model (left), is compared with a graph G_{ANOE} (right), in which nodes again correspond to all ^{1}H's and edges describe all possible assignments for each NOESY cross peak. True positives (TPs) are edges common to both G and G_{ANOE}, false positives (FPs) are edges present in G but not in G_{ANOE}, and false negatives (FNs) are the set of edges in G_{ANOE} representing the multiple possible assignments of a NOESY cross peak, none of which are present in G. These metrics are used to compute recall (R) , precision (P), and F-measure as shown in the figure and outlined in the Methods Section. The F-measure is the harmonic mean of the recall and precision. The Discriminating Power (DP) is a normalized F-measure corrected to account for the F-measure expected for a random-coil chain (DP = 0) and the best F-measure possible considering the completeness of the NMR data (DP = 1.0). Figure and legend are adopted from (Huang et al., 2021).

Fig. 2

Fig. 2. Plots of DP score vs. GDT for NMR and AlphaFold models. For each model, the DP score compares model vs NMR NOESY peak list data, and the GDT score is a measure of similarity to the NMR conformer with best DP score (Huang *et al.*, 2021). Plots are provided for (A) target T1055 (511 CASP14 models) (B) target T1027 (520 CASP14 models) (C) target T1029_original (529 CASP14 models), and (D) target T1029_revised (529 CASP14 models). Blue circles are values for CASP14 prediction models, green with blue border circles are the NMR structure models deposited in the PDB, and blue with red borders are AF prediction models. In panels C and D, the original NMR structures of target T1029, before revised analysis of NOESY data, are indicated by green circles with yellow borders. Negative DP scores are returned for models which fit the NMR data more poorly that expected for a random coil conformation (models with DP > -1 are shown). These data are replotted from reference (Huang et al., 2021).

Fig. 3. ANSURR Correlation vs RMSD scores for NMR and AlphaFold prediction models. (A) CASP14 target T1055, (B) T1027, (C) T1029, and (D) target T1027_trimmed (residues 36-75 and 96-145) in which coordinates are trimmed to remove the structurally not-well defined (i.e. unreliable) polypeptide segments. As in Fig, 2, in each panel, the blue circles are CASP 14 prediction models, green with blue border circles are the NMR structure models deposited in the PDB, blue with red borders are AF prediction models, and green circles with yellow borders (panels C and D) are for the original NMR structure of target T1029 before re-analysis.

Fig. 4. Plots of ANSURR composite score vs. GDT for NMR and AlphaFold models. The data of Fig. 3 were replot to compare the sum of correlation and RMSD ANSURR scores vs. GDT (Huang *et al.*, 2021). Plots are provided for CASP14 targets (A) T1055, (B) T1027, (C) T1029, and (D) T1027_trimmed (residues 36-75 and 96-145) in which coordinates are trimmed to remove the structurally not-well defined or unreliably predicted polypeptide segments. In each panel, the blue circles are the CASP 14 prediction models, green with blue border circles are the ensemble of NMR structure models deposited in the PDB, blue with red borders are AF prediction models, and green circles with yellow borders are for the original NMR structures of target T1029, before revised analysis of NOESY data.

CtR107

CtR107	2KCU	AF	X-ray
NMR 2KCU	2.11 Å	64.7	67.0
AF	3.89 Å	0.25 Å	97.3
X-ray 3E0H	3.62 Å	1.09 Å	----

GmR137

GmR137	2K5P	AF	X-ray
NMR 2K5P	0.78 Å	82.5	81.1
AF	1.52 Å	0.21 Å	98.8
X-ray 3CWI	1.59 Å	0.64 Å	----

RpR324

RpR324	7TZD	2LPK*	AF	X-ray
NMR 7TZD	0.54 Å	92.6	92.9	90.3
NMR 2LPK*	1.10 Å	0.45 Å	99.4	86.4
AF	1.13 Å	0.59 Å	0.27 Å	86.6
X-ray 3LMO	1.22 Å	1.68 Å	1.74 Å	----

SgR42

SgR42	2JZ2	AF	X-ray
NMR 2JZ2	0.66 Å	94.2	93.3
AF	0.86 Å	0.09 Å	99.6
X-ray 3C4S	0.97 Å	0.48 Å	----

SgR209C

SgR209C	2L06	7TZ8*	AF	X-ray
NMR 2L06	0.84 Å	82.7	82.7	82.2
NMR 7TZ8*	1.42 Å	0.94 Å	91.3	88.4
AF	1.51 Å	1.16 Å	0.11 Å	99.2
X-ray 3OSJ	1.53 Å	1.32 Å	0.52 Å	----

SrR115C

SrR115C	2KCL	2KCV*	AF	X-ray
NMR 2KCL	0.47 Å	92.9	93.4	87.4
NMR 2KCV*	1.03 Å	0.43 Å	95.6	95.8
AF	1.27 Å	1.28 Å	0.14 Å	94.7
X-ray 3MA5	1.35 Å	0.78 Å	1.13 Å	----

Fig. 5

Fig. 5. Comparison of AlphaFold, NMR and X-ray crystallography models. Superimposed backbone structures of solution NMR structures (NMR, blue), X-ray crystal structures (X-ray, grey), and AlphaFold prediction models (AF, red) for six proteins selected from the NESG NMR / X-ray pairs database. Below each superimposition is a matrix of backbone structurally-similarity statistics. The upper diagonal provides GDT-TS scores, and the lower diagonal $C\alpha$ backbone RMSDs. The diagonal (with values in red) are $C\alpha$ RMSD's within the corresponding superimposed conformer ensemble relative to the medoid conformer. These models are compared only for residues that are both "well-defined" in the NMR ensemble and "reliably predicted" in the AlphaFold models, as indicated in Table 1. For NMR and AF model ensembles, the coordinates of the medoid conformers are compared. For NMR structures refined with RDC data (i.e., targets RpR324, SgR209C, and SrR115C) the image provided is for the medoid conformer of the structure determined with these RDC data.

Fig. 6

Fig. 6. AF structures have excellent fit to RDC data. Comparison of experimentally measured ^{15}N-^{1}H RDC data (plotted on x-axis) and values computed from experimental or prediction models using PDBStat (Tejero et al., 2013). The data points are for (blue) NMR models determined without RDC data, (green) NMR models refined with ^{15}N-^{1}H RDC data, (red) AlphaFold prediction models, and (gold) X-ray crystal structures. For NMR and AlphaFold model ensembles, the medoid conformer of the well-defined regions (as indicated in Table 1) were compared. The linear correlation coefficient (R^2) for each data set is shown in the inset.

Fig. 7. Detailed comparison of solution NMR, X-ray crystal, and AF models of target RpR324. (A) Backbone ribbon representation of solution NMR structure of RpR324 (medoid conformer from PDB ID 2LPK, apo AcpXL) (blue), overlayed with AlphaFold structure calculated as monomer (red), and X-ray crystal structure (PDB ID 3LMO) (grey). (B) Overlay of AlphaFold structure calculated as monomer (red) and calculated as a dimer (orange) using AlphaFold-multimer software. (C) Comparison of (left) AlphaFold dimer (orange) with α3 helix highlighted in cyan (left), with (right) X-ray crystal structure (grey), with α3 helix highlighted in magenta. (D) Overlay of protomers from dimeric AlphaFold model (orange) with X-ray crystal structure (grey), illustrating the significant difference in orientation of α3 helix.