Research article

Genome-wide analysis of the non-coding RNA synthetic genetic network

reveals extensive plasticity and unique phenotypes in yeast

Running title: Mining yeast non-coding RNA synthetic interactions

Marcin G Fraczek[#,1,2], Soukaina Timouma[#,1,2], Laura Natalia Balarezo-Cisneros[#,1,2], Steven

Parker[1,2], Kobchai Duangrattanalert[1], Matthew Wenjie Feng[2], Sam Griffiths-Jones[2], Raymond

T O'Keefe[*,2] and Daniela Delneri[*,1,2]

[#]Joint authorship

[1]Manchester Institute of Biotechnology, Faculty of Biology, Medicine and Health, The

University of Manchester, Manchester, United Kingdom

[2]Division of Evolution, Infection and Genomics, Faculty of Biology, Medicine and Health, The

University of Manchester, Manchester, United Kingdom

*Corresponding authors: d.delneri@manchester.ac.uk; rokeefe@manchester.ac.uk

Running title: Analysis non-coding RNA synthetic interactions in yeast

ABSTRACT

Recent studies identified non-coding RNAs (ncRNAs) with unknown function that are

responsible for major fitness changes in yeast. To understand ncRNA interplay and aid their

functional assignment, the synthetic genetic array (SGA) methodology was employed to create

>15,000 double mutants and to score their epistasis in different environments. Unlike the

protein network, ncRNAs mostly displayed positive epistasis in rich medium. Interestingly, the

negative interactions significantly increased under stressors, showing environmental-

dependent functions for ncRNAs. No correlation was found between the network of ncRNAs

and that of their neighbouring genes, suggesting functional independence. The U3 paralogs,

SNR17A and SNR17B, share the majority of genetic interactions in rich medium as expected.

For example, SUT480 interacted with both paralogs and its function was linked to 18S rRNA

processing. However, under stressors, a large number of unique epistatic interactions were

observed, supporting the notion that SNR17A and SNR17B have diverged and sub-

functionalised after genome duplication.

INTRODUCTION

The majority of biological processes in the cell are performed by proteins, either independently or by interactions with other proteins. However, eukaryotic genomes pervasively transcribe sequences producing a wide range of non-coding RNAs (ncRNAs), which although are not translated into proteins, have important and often essential roles in the cell (*1-3*). Besides the well-known rRNAs and tRNAs, other classes of ncRNAs exist in eukaryotic cells. Approximately 98% of the human genome consists of non-protein-coding DNA sequences that were once regarded as non-functional evolutionary leftovers (*4*). In fact, most of the genome is transcribed and a number of ncRNAs are regulatory assisting in essential cellular processes, such as for example regulation of gene expression and transcription, apoptosis, telomere maintenance or RNA processing (*1-4*). Since ncRNAs are integral components of cellular regulatory networks, it is not surprising that many diseases in humans, ranging from cancer to neurological disorders, are affected by mutations or defects in many of these ncRNAs (*5*).

In *Saccharomyces cerevisiae*, approximately 85% of the genome is pervasively transcribed producing ~25% non-coding and ~75% protein-coding transcripts (*6*). *S. cerevisiae* constitutes an exception among Eukaryotes as it has lost small regulatory ncRNAs and conserved proteins required for RNA interference (RNAi) such as Argonaute and Dicer-like RNAses (*7, 8*). However, several classes of so called classical or housekeeping ncRNAs are present in the *S. cerevisiae* genome including tRNAs and rRNAs involved in protein synthesis as well as small nuclear RNAs (snRNAs) known to perform intron splicing and small nucleolar RNAs (snoRNAs) that process or direct chemical modifications of other RNAs. Novel functions for these classical ncRNAs are still emerging. Furthermore, a great number of ncRNAs with unknown or unclear functions have been discovered and classified into various groups depending on their half-life after transcription. The majority of ncRNAs are unstable (short-lived) as they are rapidly degraded by the RNA decay pathways. These include Cryptic Unstable Transcripts (CUTs) that were discovered in mutants of the nuclear exosome exonuclease component *RRP6* (Δ*rrp6*) (*9, 10*), Xrn1 exo-ribonuclease sensitive Unstable Transcripts (XUTs) (*2, 11*) and RNA-binding factor Nrd1 Unterminated Transcripts (NUTs) (*12*). There are also long-lived Stable Unannotated Transcripts (SUTs) that evade degradation in the nucleus and are processed in a similar manner to mRNA and exist in the wild type cell (*10*). The stability of SUTs implies that they are functional and involved in cellular processes. It is estimated that in the genome of *S. cerevisiae* there are >800 SUTs, >900 CUTs, >1800 XUTs and >1500 NUTs (*2, 10, 12*). These ncRNAs are transcribed from both sense and

antisense strands in relation to the coding genes from intra- and intergenic or open reading

frame overlapping regions. Many of these ncRNAs overlap with each other and some are

extended isoforms of other ncRNAs. Their molecular functions are still largely unknown.

Only a limited number of large-scale studies to identify the cellular functions ascribed to

ncRNAs have been performed (*13-16*). Libraries of 1502 barcoded ncRNA deletion mutants in

S. cerevisiae in both haploid (*MATa* and *MATα*) and diploid (homo- and heterozygote)

backgrounds were constructed as tools for ncRNA functional analysis (*15, 17*). These libraries

contain deletion mutants of tRNAs, snRNA, snoRNA, CUTs, SUTs and other annotated

ncRNAs with a total number of 443 unique ncRNAs, that are localised to intergenic regions

and do not overlap with protein coding genes, their promotors or terminators.

The yeast protein deletion collection has been successfully used to detect genetic interactions

(GIs) and scored epistasis between protein coding genes by synthetic genetic array (SGA)

screens (*18, 19*). Systematic mapping of genetic interaction was first automated in *S. cerevisiae*

by employing the single open reading frame deletion collection composed of ~4800 genes

crossed with 132 query strains using the SGA methodology (*20*). This approach led to

identification of ~4000 genetic interactions between ~1000 protein coding genes and the

construction of a large-scale genetic interaction map of the cell. Here, we employed the same

methodology to shed light on the function of ncRNAs in *S. cerevisiae* by generating double

ncRNA null mutants and analysing their genetic interactions by detecting double mutants with

altered phenotypes. A ncRNA mutant library encompassing 378 ncRNAs was created in the

SGA query strain Y7092 and an SGA study was conducted with a subset of 38 knock-out

deletion mutants crossed with 411 previously generated single ncRNA deletion mutants (*15,*

17). The phenotypic effect of the double ncRNA deletions, compared to the single mutants,

was then analysed in rich and non-fermentable media as well as in three stressor conditions,

including oxidative, high temperature and osmotic stresses. The interactive network fitness

data of the ncRNAs epistatic network has been deposited in the publicly accessible online

resource, Yeast ncRNA Analysis (YNCA; http://sgjlab.org/ynca/). We discovered a total of

1003 epistatic interactions in rich media, of which ca. 10% were synthetic sick or lethal.

Compared to the SGA protein-coding network, the ncRNAs mostly displayed positive

interactions, and did not correlate with the epistasis of their neighbouring protein coding gene.

We detected 395 "core" interactions shared in all condition tested and a significant number of

environmental dependent interactions, including a differential SGA network for the two U3

snoRNA genes, SNR17A and SNR17B, suggesting a functional divergence of these two

paralogs driven over time by different environmental pressures.

RESULTS AND DISCUSSION

Query ncRNA deletion mutant library construction and strain selection for SGA analysis

Single ncRNA deletion mutants in the Y7092 background (referred here as 'query') were

generated by replacing the ncRNA of interest with the *natMX4* marker that confers

nourseothricin resistance (clonNAT). A total number of 378 deletion mutant query strains were

generated (Supplementary Dataset S1) and SGA analysis was conducted with 38 of them.

Single mutant query strains with fitness defects are more likely to show a higher number of

epistatic interactions compared to query strains with low or no fitness deficiency (*21*). In fact,

there is a strong positive correlation between the number of genetic interaction and single-

mutant fitness ($r = 0.73$) (*21*). Based on previous fitness studies (*15, 16*) that scored both the

biomass (solid fitness of haploid ncRNA mutants) and the growth changes over time

(competitive fitness of diploid heterozygous ncRNA deletion mutants), we chose 34 single

ncRNA deletion strains which displayed either growth deficiency or haploinsufficiency under

obligatory respiratory condition, high temperature, osmotic stress conditions, alternative

carbon source such as sorbitol and melezitose or high concentration of ethanol (10%) and 4

strains with no significant fitness changes in any conditions (Supplementary Fig. 1).

SGA analysis of ncRNAs mutants reveals a prevalence of positive epistasis in the genetic

interaction network

We performed SGA screens by crossing 38 query strains of interest with the ncRNA array

library composed of 411 ncRNA deletion mutants (Supplementary Dataset S2), including 49

CUTs, 108 SUTs, 59 snoRNAs, 194 tRNAs and 1 unknown ncRNA RUF22. Approximately

6.4% (1003 interactions; Supplementary Dataset S3) of the 15,600 double mutant combinations

generated, displayed either positive or negative significant epistasis (Fig. 1A). Positive

epistasis was displayed by 905 (~90.2%) double mutants (Fig. 1B). In total, there were 98

(~9.8%) significant negative interactions of which 51 (~5.1%) and 47 (~4.7%) were synthetic

sick (Fig. 1C) and synthetic lethal (Fig. 1D), respectively. Overall, each query ncRNAs had an

average of 26 significant interactions, with numbers varying between 18 to 34 (Table 1,

Supplementary Dataset S3).

Table 1: Number of significant SGA interactions between query and array strains,
divided per ncRNA classes. The table shows the 1003 significant ncRNA-ncRNA interactions
for 38 query strains analysed (fold change $\geq$2 and q-value $\leq$0.001) broken down according to
type of ncRNAs. Positive and negative epistasis are represented with + and -, respectively.
Refer to Supplementary Dataset S3 for detailed interactions.

| | Number of interactions in the different ncRNA classes | | | | | | | | Total interactions | | |
| | CUTs | | SUTs | | SNRs | | tRNAs | | + | - | + & - |
Query	+	-	+	-	+	-	+	-			
CUT084-SUT068	4	0	9	1	0	1	10	2	23	4	27
CUT150	3	0	13	0	0	0	7	3	23	3	26
CUT244	1	0	8	1	0	1	6	7	15	9	24
SNR13	1	1	9	1	0	0	4	2	14	4	18
SNR17A	4	0	12	0	1	1	14	0	31	1	32
SNR17B	4	0	12	0	0	1	11	0	27	1	28
SNR34	3	0	10	0	0	0	12	2	25	2	27
SNR45	3	0	14	0	0	0	12	0	29	0	29
SNR71	3	0	12	0	1	0	11	0	27	0	27
SNR72	3	0	12	0	0	0	11	0	26	0	26
SNR72-78	3	0	12	0	0	0	11	2	26	2	28
SNR73	3	0	13	0	1	0	11	0	28	0	28
SNR74	3	0	12	0	1	0	11	0	27	0	27
SNR75	3	0	11	0	1	0	11	0	26	0	26
SNR76	2	0	11	0	0	0	10	0	23	0	23
SNR77	3	0	13	0	0	0	11	0	27	0	27
SNR78	3	0	13	0	1	0	11	0	28	0	28
SNR79	2	0	10	2	0	0	11	1	23	3	26
SNR8	3	0	13	0	0	0	11	1	27	1	28
SNR87	3	0	14	1	1	0	11	1	29	2	31
SUT083	3	0	13	0	0	0	12	0	28	0	28
SUT107	3	0	15	0	0	0	12	0	30	0	30
SUT193	3	2	7	2	0	1	9	2	19	7	26
SUT211	3	0	11	0	1	0	12	0	27	0	27
SUT289-SUT717	3	0	14	1	0	0	11	0	28	1	29
SUT361	0	1	9	0	0	0	8	2	17	3	20
SUT414	1	3	14	0	0	0	7	1	22	4	26
SUT420-CUT425	0	0	10	1	0	2	7	3	17	6	23
SUT457	3	1	13	0	1	0	11	0	28	1	29
SUT496	3	1	9	2	0	1	10	0	22	4	26
SUT532	1	3	6	2	0	3	9	3	16	11	27
SUT764	1	0	9	1	0	1	8	1	18	3	21
tD(GUC)M	2	0	13	1	0	0	9	1	24	2	26
tE(CUC)D	2	2	11	2	0	4	10	3	23	11	34
tE(UUC)M	1	1	9	2	0	1	9	0	19	4	23
tH(GUG)H	3	0	10	2	0	1	6	0	19	3	22
tI(AAU)N1	1	0	13	0	0	0	10	2	24	2	26
tV(AAC)G3	1	0	10	2	0	1	9	1	20	4	24
Total	91	15	429	24	9	19	376	40	905	98	1003

179
180

In contrast to protein-coding genes for which negative genetic interactions are approximately
two-fold more prevalent than positive interactions (*21*), here a surprisingly large number of
double mutants displayed positive epistasis, comprising ~90% of the significant interactions
(Fig. 1A, Table 1). The prevalence of positive epistasis in the SGA network was also seen when
interactions were broken down for each individual query strains. For example, SNR45, SNR71,
SNR72, SNR73, SNR74, SNR75, SNR76, SNR77, SNR78, SUT083, SUT107 and SUT211
only displayed positive interactions (Table 1). The snoRNAs SNR72, SNR73, SNR74, SNR75,
SNR76, SNR77 and SNR78 are in a tight cluster on the chromosome 13 and in the SGA are
detected as synthetic lethal interactions because it is unlikely that any recombination is
occurring between them. Such spurious inter-cluster interactions have been removed from our
dataset. We also carried out SGA using the entire SNR72-78 cluster null mutant. In this case
we detected two synthetic lethal interactions with two tRNAs on chromosome 2 and 12; both
coding for leucine (Supplementary Dataset S3). Moreover, SNR66 (C/D box snoRNA)
displayed 8 positive interactions, of which six were between snoRNAs (SNR71, SNR73,
SNR74, SNR75, SNR78 and SNR87) and two with SUTs (SUT211 and SUT457)
(Supplementary Dataset S3). Interestingly, SNR66 is involved in the same biochemical
function (methylation of rRNAs) as SNR71, SNR73, SNR74, SNR75, SNR78 and SNR87. It
has been reported previously that proteins that are part of the same complex and therefore
physically interact with each other are threefold more likely to show positive epistasis (*18*).
Hence, these snoRNA interactions may also reflect a physical relationship.
201
202 SUTs were responsible for the highest number of epistatic interactions, after normalisation for
203 the total cohort (p-value $=1.66e^{-53}$), whereas snoRNAs displayed the least number of
204 interactions (Table 1). These results are consistent with SUTs being transported to the
205 cytoplasm similarly to mRNAs and acting as functional transcripts affecting important cellular
206 functions (*16, 22, 23*).
207
208 The ncRNA genetic interaction networks, hosted in the Yeast NcRNA Analysis database
209 (YNCA), can be explored as an interactive map through the link http://sgjlab.org:3838/ynca/,
210 "interactive networks" tab.
211
212
213

Comparison between ncRNA and neighbouring protein interaction networks reveals poor overlap

To understand whether altered expression of genes flanking the ncRNAs indirectly influenced the ncRNA interaction network, we surveyed all genetic and physical interactions of the proteins encoded by the neighbouring genes, to determine the overlap between protein and ncRNA interaction networks. Intergenic ncRNAs could either work in *trans* and affect genes or transcription factors involved in cellular growth (*15, 16, 24*) or may provoke expressional changes in *cis* (*25-28*) and this expression change could cause the epistasis between the affected genes (*i.e.* synthetic interactions between the neighbouring genes due to expressional changes). Provided that the latter scenario is the most common, we should expect some overlap between the SGA network of ncRNAs and the SGA network of the neighbouring protein coding genes.

The BioGrid database was used to recover the interactions which are recorded as genetic or physical (*i.e.* protein-protein interactions scored either by affinity capture, protein-fragment complementation assay or two hybrid bait). We found that the overwhelming majority of the neighbouring genes (96.8%) flanking the ncRNAs involved in the ncRNA network had no recorded genetic or protein interactions (Fig. 2, and Supplementary Dataset S4). Recent data from transcriptome studies of ncRNA mutants revealed that the deletion of several SUTs has a global, rather than a local effect on gene transcription and can act on a large-scale by modulating expression of transcription factors (TFs) (*16*). For instance, in the absence of SUT532, SUT125 and SUT126 the expression of TFs that participate in the stress response, sporulation, cell cycle progression and cell integrity such as Xpb1, Rim1 Rgm1, Yox1, Rof1, Tos8, Msa1, and Tos4 was highly affected. Such modulation is scored as a growth advantage in the short term, although unlikely to render the cell fitter over time given that an accurate mitotic control is crucial to avoid multiple duplications and DNA damage.

Out of the 3.2% of ncRNAs which had neighbouring genes involved in interactions, ~64.5% were genetic and ~35.5% physical (Fig. 2). In this small subset most of the ncRNA genetic interactions were negative (21/24; 87.5%) leading to phenotypic suppression, synthetic growth defect and dosage lethality (Supplementary Dataset S4), mimicking the overall trend seen for the protein-coding gene network, where the majority of the interactions were negative(*29*). This neighbouring gene interaction suggests a potential *in cis* effect of the ncRNA on the flanking genes. Although when generating the ncRNA mutant collection (*15, 17*) every effort has been made to avoid deletion of either promotor or terminators of the adjacent protein coding genes,

it is possible that for some ncRNA double mutants the observed phenotype is due (or partially

due) to altered expression of flanking genes.

The ncRNAs genetic interaction data can also reveal neighbouring gene epistasis from their

overexpression (*i.e.* when altered expression of two genes together cause a phenotype but the

altered expression of the single mutants does not). For example, upon SUT259/691 deletion,

EMP46 and *GAL4* are both overexpressed causing a lethal phenotype (*15*). Such epistatic

interactions between proteins would not have been revealed by classical SGA analysis based

on gene deletions.

Network dynamics of ncRNAs in different stress conditions reveal more negative epistatic

interactions

Genetic epistasis is dependent on the genetic background and the environment(*30*) therefore,

we analysed all single and double ncRNA mutants under five different nutritional and stress

conditions (rich and non-fermentative media as well as oxidative, high temperature, and

osmotic stresses) to detect environmental-specific deviation of growth and infer changes in the

ncRNA network. We scored the fitness plasticity of the interaction networks with 31 query

strains encompassing ~12,700 mutant combinations (Fig. 3, Table 2 and Supplementary

Dataset S5). We observed that stress produces a significantly increased interactions, in

particular negative interactions. High temperature stress (37°C) displayed more negative than

positive epistatic interactions (Fig. 3B), an opposite trend than detected at 30°C (Fig. 3A). The

highest number of interactions, either positive or negative, was observed under respiratory

stress (YP+7% glycerol, 30°C; Fig. 3C) for which there were 40 times more negative

interactions than in rich medium. Overall, the different stress conditions displayed at least two

times more negative interactions than YPD at 30°C.

Interestingly, regardless of media and temperature, most interactions involving snoRNAs were

positive. In fact, there are conditions where snoRNAs display no negative interactions. This

result echoes the "monochromatic" nature observed in the protein-coding network, where

genetic interactions between defined pathways tend to be either exclusively positive or

exclusively negative (*21, 31*). Although the monochromatic effect is robust, the pattern of

interactions between the functional modules could be considerably affected by environmental

perturbations, as observed with a small number of snoRNA interactions when exposed to a

particular stress condition (Fig. 4 and Supplementary Dataset S5).

Table 2: Number of significant ncRNA interactions after fitness profiling in different environmental conditions.

Condition	Total number of significant interactions	Number of positive interactions	Number of negative interactions
YPD, 30°C	839	716	123
YPD, 37°C	831	398	433
YP+7% glycerol, 30°C	1525	844	681
YPD+1 M NaCl, 30°C	1027	653	374
YPD+4 mM H$_2$O$_2$, 30°C	1145	734	411

Next, we identified ncRNA interactions that were shared and unique to each of the environmental conditions tested (absolute fold change ≥2 and q-value ≤0.001) (Fig. 4A, Supplementary Dataset S6). There were 395 "core" ncRNA interactions shared between all five conditions tested. One hundred and fifty-nine ncRNA interactions were specific only to rich medium (YPD, 30°C), 392 to glycerol (respiratory stress), 24 to NaCl (osmotic stress), 19 to 37°C (temperature stress) and 12 to hydrogen peroxide (oxidative stress). Interestingly, in all the four stress conditions tested 21% of the core interactions changed directionality compared to rich medium (Supplementary Fig. 2). Among those that changed, 5 negative interactions became positive and 79 positive interactions became negative, with exception for 7 interactions in osmotic shock medium (Fig. 4B). A visual representation of the plasticity of the interactions in different media conditions of selected single and double mutants is shown in Fig. 4C. The observed sub-networks with shared common ncRNAs that change directionality in the stress conditions suggests that these ncRNAs may have a primary function in stress response. For instance, we identify a prevalence of negative interactions under obligatory respiratory condition (YP+7% glycerol, 30°C) among ncRNAs that have previously described to display a pivotal role in cell fitness (*16*). In fact, SUT125, SUT126 and SUT035 share similar phenotypic and transcriptome changes under different environmental conditions (*16*). These ncRNAs, along with SUT532, appear to modulate genes *in trans* that are involved in mitochondrial functions. Remarkably, we identified 22 epistatic interactions in common among SUT125, SUT126, SUT035 with the presence of predominantly negative interactions under respiratory stress, YP+7% glycerol at 30°C (Fig. 4C, Supplementary Fig. 2 and Supplementary Dataset S3). In the same stress condition, SUT532 displayed 14 unique interactions with other ncRNAs (Supplementary Dataset S6).

The potential correlation between phenotypic changes in a single ncRNA deletion mutant and

the number of SGA interactions was investigated using previously published fitness data (*15,*

16). We found that there is only a weak positive correlation between the fitness change of single

mutants and the number of interactions (Supplementary Fig. 3 and Supplementary Dataset S5).

We also compared the ncRNA interactions in YPD and the other conditions with the available

interaction network in minimal medium of the neighbouring protein coding genes. Again, only

a small subset of neighbouring genes/proteins displayed interactions either genetic or physical

(Supplementary Fig. 4 and Supplementary Dataset S7).

Analysis of the paralog snoRNAs, SNR17A and SNR17B

Next, we focused our attention on the yeast SNR17 (U3). In *S. cerevisiae* this snoRNA is

transcribed from two genetic loci *SNR17A* and *SNR17B* located on separate chromosomes, XV

and XVI, respectively. U3 is required for pre-rRNA processing, namely cleavage of 35S pre-

rRNA leading to mature 18S rRNA production (*32, 33*). An early study suggested that these

two paralogs may perform the same function in the cell since they share the same nuclear

localisation and bind to 35S pre-rRNA (*34*). Our results reveal that deletion of SNR17B does

not significantly change the phenotype compared to the WT, except in YP+7% glycerol, while

the SNR17A deletion displays decreased growth in YPD 30°C, YPD 37°C and YPD+4 mM

H_2O_2 (Supplementary Fig. 5). The double deletion of SNR17A and SNR17B is lethal (*34*) and

our SGA results confirm it their lethality together.

The SGA network of SNR17A and SNR17B can help to understand whether the function of

these two paralogs has diverged over evolutionary time. Thirty-one and 27 significant SGA

interactions were detected in our screens in YPD at 30°C with SNR17AΔ and SNR17BΔ,

respectively (Fig. 5A and Supplementary Dataset S8). As expected, the majority of interactions

were shared between these two snoRNAs reflecting that they are functional homologs (Fig.

5a). In fact, only six interactions were unique to SNR17A and two to SNR17B in YPD 30°C

and all displayed positive epistasis. However, surprisingly, a large increase in paralog specific

interactions was observed in other environmental conditions (Fig. 5, B to E) and such

specificity was not lost even using the most stringent q-value of 0.001 (Supplementary Fig. 6

and Supplementary Dataset S8). Interestingly, under obligatory respiration, oxidative,

temperature and osmotic stresses, the interaction network exclusive to SNR17BΔ was highly

expanded, revealing 5 to 10 times more specific interactions than SNR17AΔ. This difference
in the interaction network of SNR17A and SNR17B may indicate that since the duplication
event, the paralogs have sub-functionalised, and SNR17B might also be modulating different
cellular functions independently of its paralog SNR17A. Within the SNR17B specific network,
seven interactions were in common in all stress conditions, five of which involving tRNAs.
Interestingly, there is an overrepresentation of tRNAs in the specific network of SNR17B (p-
value=0.05 Pearson's Chi-squared test, Yates' continuity correction) which suggest a potential
role of SNR17B modulating/processing tRNAs. In other organisms, co-immunoprecipitation
and high-throughput sequencing of ligated RNAs have revealed interactions between
snoRNAs and other RNAs such as tRNA (*35*) and mRNAs (*36*). Moreover, snoRNAs have
been reported to participate in methylation of tRNAs (*37*), regulating 3′ mRNA processing (*38*)
and alternative splicing (*39*).
353
We tested the absolute expression of the two paralogs in three *S. cerevisiae* strains. Early
studies to detect snoRNA levels in the cell by labelled probe hybridisation to total RNA
revealed that SNR17A RNA was approximately 5 to 10-fold more abundant than SNR17B
implicating SNR17A as the major player in 18S rRNA processing (*34, 40*). Contrary to these
results, by using RT-qPCR, we observed that SNR17B RNA levels were three times higher
than SNR17A in two haploid WT strains BY4741 and Y7092 (Fig. 6A). This difference in
expression was also consistent when the snoRNA levels were measured in a natural diploid
isolate, *S. cerevisiae* 96.2 (courtesy of E. Barrio) (*41*), suggesting that it is not strain dependent.
For BY4741, we also tested the expression of the snoRNAs in presence of stressors (Fig. 6B).
Here, except for YP-7% glycerol at 30°C, we observed again the same trend with SNR17B
being the most highly expressed of the two paralogs. We also measured the expression of
SNR17A and SNR17B in the mutants SNR17BΔ and SNR17AΔ, respectively. In the standard
condition the levels of SNR17B were reduced almost to half when SNR17A was deleted, while
levels of SNR17A displayed no significantly change in the SNR17BΔ deletion mutant (Fig.
6C). These data confirm that the rRNA processing function of SNR17A, with respect to
SNR17B, is not due (or at least not only due) to its abundance, as previously suggested.
370

Deletion of SNR17A causes changes in 18S rRNA concentration
Since SNR17 is crucial for 35S pre-rRNA processing and maturation of 18S rRNA (*32, 33*),
we hypothesised that the deletion of the U3 snoRNA paralogues would reduce the

concentration of the 18S rRNA fraction in the cell. We tested our hypothesis by measuring the

concentration of 18S rRNA in the WT and deletion mutants grown in the yeast standard

condition and four stressors (high temperature, obligatory respiration, oxidative and osmotic

stresses) to determine if any 18S rRNA concentration changes are environmentally dependent.

Deletion of SNR17A resulted in a significant reduction of 18S rRNA fraction abundance when

compared to the WT strain in four out of five conditions tested (Fig. 7A). This reduction in 18S

rRNA concentration supports the established notion that SNR17A has a major role in pre-rRNA

processing. However, no significant changes in 18S rRNA abundance were observed in the

SNR17B deletion mutant, suggesting that SNR17B is not the key player in the maturation of

18S rRNA and may not mainly act as back up for SNR17A. Overall, these data suggest that

SNR17B may have sub-functionalised and be involved in other cellular functions.

Since our SGA screens revealed a positive epistatic interaction between SNR17A and SUT480,

most evident in the respiratory stress condition (Supplementary dataset S5), we investigated

how the concentration of 18S rRNA is affected by the epistatic interaction of SNR17A and

SUT480. We measured the concentration of 18S rRNA in both yeast standard condition and

YP+7% glycerol at 30°C for the WT, single SNR17A and SUT480 deletion mutants and the

double mutant SNR17AΔ/SUT480Δ (Fig. 7B). We found that the 18S rRNA concentration is

decreased in both the single mutants SNR17AΔ and SUT480Δ. The reduction of 18S rRNA

concentration in the single SUT480Δ mutant is more evident in the respiratory stress compared

to the standard growth media. No additive decrease of 18S rRNA concentration was observed

for the double mutant compared to single mutants. In fact, an increase in 18S concentration in

glycerol was observed compared to the SUT480Δ. Such data validate the positive epistatic

interaction detected in the SGA screen and suggest that these two ncRNAs synergistically

affect rRNA processing through the same pathway. qPCR data reveals that SUT480, SNR17A

and SNR17B do not affect each other's transcription levels in either of the lab strains tested

(Supplementary Fig. 7, A to C). Moreover, deletion of SUT480 does not influence the

expression of its neighbouring gene *RNH202*, which codes for the ribonuclease H2 subunit

required for RNA hydrolysis when annealed to a complementary DNA (*42*) (Supplementary

Fig. 7D), ruling out an indirect effect of the SUT480 phenotype acting by transcriptional

interference.

CONCLUSIONS

409
This study constitutes a pioneering effort to determine the genetic interactions between 5
classes of ncRNAs, namely CUTs, SUTs, snoRNAs, tRNAs and RUF, and provides insights
on their possible functions in yeast. We observed that the majority of genetic interactions
between ncRNAs in the rich condition are positive ($>\sim90\%$) as opposed to the predominantly
negative genetic interactions scored for the protein network (*18*). This positive epistasis
indicates that most of the ncRNA have presumably repressive regulatory functions in the cell.
The networks of ncRNAs and their neighbouring genes are primarily independent, showing
little overlap between each other. The small subsets of genetic networks that do overlap are
composed predominantly by negative interactions, suggesting a potential effect *in cis* of the
ncRNA on the flanking genes.

The environment has a significant impact on the ncRNA genetic interaction network. We
observed an increase in the number of negative interactions in all stress conditions tested, with
the highest number recorded in glycerol (up to 40 times more compared to rich media). This
result suggests that these ncRNAs have unique functions in the cell or are being differentially
transcribed depending on the environment.

The interaction data acquired in this study is also useful for teasing apart cellular roles of
ncRNA duplicates and to find whether their functions have diverged over evolutionary time,
as in the case of snoRNAs SNR17A and SNR17B. By measuring the expression of SNR17A
and SNR17B in the cell and the concentration of 18S rRNA in SNR17A and SNR17B mutants,
we confirmed that SNR17A is a major player in rRNA processing and this function is not
because of SNR17A abundance. Looking at SNR17A and SNR17B genetic networks in rich
medium, the two paralogues share the majority of genetic interactions, as expected. However,
under stress conditions, SNR17B displayed a large number of unique epistatic interactions
suggesting that they may have sub-functionalised as result of functional divergence that occurs
after genome duplication (*43, 44*), and has acquired other roles in the cell. We found that
SUT480 is genetically interacting with SNR17A, we linked its function to the processing of
18S rRNA, and validated the scored epistasis.

Overall, this study offers the first insight in the environmentally-dependent epistatic
interactions of ncRNAs in an eukaryotic organism.

MATERIALS AND METHODS

Strains and plasmids

We previously generated a library of ncRNA mutants in both haploid and diploid backgrounds (*15, 17*). The *MAT***a** library from this collection (Supplementary Dataset S2), referred here as 'array', was used in the crossing with the haploid *MATα* 'query' single deletion strains (generated in this study; Supplementary Dataset S1) to create double mutant according to the SGA protocol (*18*). The *MAT***a** array library was constructed in the BY4741 background (*his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) and all these mutants carry a *kanMX* resistance gene that replaced the ncRNAs of interest (*17*). The *MATα* query collection was generated in the Y7092 background (*can1Δ::STE2pr-Sp_his5 lyp1Δ ura3Δ0 leu2Δ0 his3Δ1 met15Δ0*; kindly provided by the Boone lab) (*18*). The query ncRNA deletion strains carry the *natMX4* resistance marker that replaced the ncRNAs of interest. This *natMX4* marker confers clonNAT (Nourseothricin; WERNER BioAgents GmbH) antibiotic resistance and was amplified from the pFA6-*natMX4* (*45*) plasmid in a single PCR step using primers partly complementary to the ncRNA flanking sequences. A natural diploid *S. cerevisiae* strain 96.2 (*41*) was used to test differential expression of SNR17A and SNR17B.

Query strain library construction

The query strains were constructed by substituting the ncRNA loci with the *natMX4* cassettes. Each cassette was amplified using a pair of barcoded primers containing genome complementary sequences (*46*) from pFA6-*natMX4* in the total of 25 µl PCR volume: 10 ng plasmid DNA, 5 µM of forward and 5 µM of reverse primers, 10 mM of each dNTP, 2.5 units of LongAmp polymerase (New England Biolabs) and 1x of reaction buffer. The cycling conditions were as follows: initial denaturation at 95°C for 2 min followed by 35 cycles of 94°C for 30 sec, 58°C for 30 sec and 65°C for 1.5 min and 1 step of final elongation of 65°C for 5 min. The amplified PCR products were ~1.5 kb and they were transformed into the Y7092 (*MATα*) background strain. Transformants were selected on YPD agar (1% yeast extract, 2% peptone, 2% dextrose, 2% agar) supplemented with 200 µg/ml of clonNAT (Werner BioAgents, Jena, Germany). After 4 days of growth at 30°C, 1 to 5 colonies were streaked and single colonies were confirmed by colony PCR for correct cassette integration. The list of all query strains constructed in this study can be found in Supplementary Dataset S1.

PCR confirmation of query strains

Correct ncRNA locus deletion was confirmed by colony PCR. Following transformation, streaked colonies were resuspended in 100 µl sterile H_2O. Three sets of primers were used: (i) confA-confNatR, (ii) confNatF-confD and (iii) confA-confB (Supplementary Fig. 8, Supplementary Table S1). Confirmation primers confA, confB and confD are listed in Parker *et al.*, 2017 (*17*). Primer pairs (i) and (ii) only generated PCR products if the ncRNA was replaced with the deletion cassette. Primer set (iii) generated PCR products if the ncRNA was intact and these colonies were discarded. The PCR was performed in the total volume of 25 µl and contained 5 µl of resuspended cells, 5 µM of forward and 5 µM of reverse primers, 10 mM of each dNTP, 2.5 units of LongAmp and 1x of reaction buffer. The same PCR conditions as above were used, generating PCR products between 0.6 kb and 1.2 kb, depending on the deleted ncRNA, which were analysed on a 1.5% agarose gel. Confirmed colonies were stored in YPD containing 15% glycerol at -80C until needed.

Single deletion ncRNA library formatting for SGA

The single deletion array library (*MATa*; ncRNA*::kanMX4*) is composed of 411 mutants encompassing 49 CUTs, 108 SUTs, 59 snoRNAs, 194 tRNAs and 1 RUF (Supplementary Dataset S2) (*17*). The library was first organised in 96 and 384 density formats and subsequently used as a source to generate the final working copy of two Plus Plates (Singer Instruments, UK) in a density of 1536 colonies per plate using the RoToR HDA system (Singer Instruments, UK). Each single deletion strain was plated in quadruplicates allowing replicates for consistent SGA analysis. The edges of each plate were excluded from the SGA analyses since colonies that are located on the edges and corners have more nutrient availability and decreased competition with the neighbouring colonies and hence often show increased growth (*47*). To minimise contamination, the single deletion ncRNA plates were grown on YPD agar supplemented with 200 µg/ml G418 (Sigma-Aldrich) or SD complete amino acid agar (0.67% yeast nitrogen base with amino acids (Merck), 2% glucose, 2% agar) supplemented with 200 µg/ml G418. The images of these plates were recorded using a Phenobooth (Singer Instruments) for subsequent SGA analyses.

Generation of the double ncRNA mutants using SGA

Thirty-eight query ncRNA deletion mutant strains (*MATα*; ncRNA*::natMX4*) (Supplementary Dataset S1) were selected to generate a library of double ncRNA deletion mutants. Prior to SGA, a small portion of -80°C cell cultures of these query strains was grown overnight in 5 ml

of liquid YPD at 30°C with shaking (200 rpm). Subsequently, 1 ml of the overnight cultures

was spread onto PlusPlates containing YPD supplemented with 200 µg/ml of clonNAT using

plastic cell spreaders and grown overnight at 30°C. Then, the query strains were crossed with

the array ncRNA deletion library in the 1536 format on YPD agar using a RoToR HDA robot

and disposable plastic replicators (pinning RePads, Singer Instruments, UK). SGA was carried

out according to Baryshnikova, *et al.,* 2010 (*18*) including the generation of the double ncRNAs

mutants library and subsequent analysis of epistasis.

Quality control of the double mutants created using SGA

Following SGA, 20 randomly selected haploid ncRNA double mutants were validated by PCR.

The following primer sets were used for ncRNA1::*kanMX4* validation: (i) confA+confKanR,

(ii) confD+confKanF and (iii) confA+confB. Primer sets (i) confA+confNatR, (ii)

confD+confNatF and (iii) confA+confB were used for ncRNA2::*natMX4* validation in the

same mutant (Supplementary Fig. 9, Supplementary Table S1).

SGA data analysis

Following SGA, images of the final selection plates containing generated double ncRNA

mutants were recorded in white light using a Phenobooth and cropped using Corel PaintShop

Pro X7. Subsequently, the images were processed using the high-throughput SGA data analysis

software *Balony* (*48*) that measures colony area in pixels. The initial normalisation was

performed in this software to correct for uneven plate growth according to the plate median

and the row/column correction was applied. Minimum and maximum spot sizes were set to

0.02 and 100, respectively. Following the initial image processing in the *Balony* software,

generated data was analysed using the R package version 3.4.4 (*49*). Recorded output size

values for single and double mutant colonies were used to generate SGA scores using a script

available on the SGATools website (http://sgatools.ccbr.utoronto.ca/) as well as in

Baryshnikova, *et al.* (*18*) and Wagih *et al.* (*50*)) colonies on the periphery of the plates (as

described above), (ii) single and double ncRNA mutants that showed inconsistent growth

patterns among the quadruplicates and (iii) linkage group of ncRNAs that were on the same

chromosome and were scored as lethal after the SGA as their phenotype was likely to be linked

to the lack of meiotic crossing-over between them. The data was processed using an in-house

R script. Assuming a normal distribution, p-values were calculated for each sample mean using

the Z-score. The Z-score was found by assuming that the null hypothesis was true, subtracting

the assumed mean, and dividing by the theoretical standard deviation. Q-values, which are

Bonferroni-adjusted p-values, were inferred using the "stats" R package (*49*). Absolute fold

change ≥2 and q-value ≤0.001 thresholds were applied to select most significant interactions.

The "UpSetR" R and "matplotlib_venn" python packages were used for data visualisation. The

R pipeline created for this analysis is available at https://github.com/Sookie-S/SGA-analysis.

SGA network generation

Genomic interaction networks were created from the data generated for all experimental

conditions tested. These included yeasts grown in rich media as well as on non-fermentable

media, oxidative, high temperature and osmotic stress environments. The networks were

generated using the "visNetwork" and "igraph" R libraries (*51, 52*). Nodes represent the

ncRNAs and the edges represent the genetic interaction between them. The size of the nodes

is proportional to the number of connections with the other nodes. The edges are coloured in

green and red to distinguish the positive and negative epistasis, respectively. The thickness of

the edges is proportional to the absolute value of fold change. The force-directed layout

(forceAtlas2Based solver) (*53*) has been used to organize the graph in a 2D space. The central

gravity of the networks is distance independent. The repulsion between the nodes is linear.

Phenotypic analysis of the mutants in stress conditions

The phenotypic analysis of both double and single deletion strains was performed in five

different environmental conditions on solid growth media. These included (i) yeast rich media,

YPD agar at 30°C, and four stress conditions: (ii) high temperature on YPD agar at 37°C, (iii)

obligatory respiratory on YP agar supplemented with 7% glycerol at 30°C, (iv) osmotic stress

on YPD agar supplemented with 1 M NaCl at 30°C and (v) oxidative stress on YPD agar

supplemented with 4 mM H_2O_2 at 30°C. The single and double deletion library was analysed

in the same 1536 format as described for SGA above. All cells were first grown on solid YPD

agar media for 48h prior to the experiment and subsequently transferred on the specific

experimental plates using the RoToR HDA system and grown for 48h at their relevant

temperatures before analyses. The images of these plates were recorded using a Phenobooth

and analysed using the *Balony* software in the similar manner as described for the SGA above.

We used the same cut-off analysis values (absolute fold change ≥2 and q-value ≤0.001) as

described above for the SGA analysis in standard yeast condition.

Genetic and physical interactions between neighbouring genes/proteins of interacting ncRNAs

Protein-protein physical interactions as well as genetic interactions between coding genes that were located in the neighbourhood of every interacting ncRNA pairs were analysed. Only significantly interacting ncRNA pairs were included (with absolute fold change ≥2 and q-value ≤0.001). A synthetic genome annotation file (BED file format) containing the names of protein coding genes and ncRNAs, their positions and strand information, was generated using an in-house Python 3.6 script. Gene annotations were extracted from the *Saccharomyces* genome database (SGD; www.yeastgenome.org/) and ncRNA annotations were added manually based on the information from Wery *et al*., 2016 (*2*). Subsequently, the genes and ncRNA were ordered by their start position and by the chromosome number using the BedTools sort tool (*54*). The flanking genes for each ncRNA were extracted using an in-house Python 3.6 script, generating 4 combinations of gene interactions for each pair of interacting double ncRNAs. A Python 3.6 script that uses the Representational State Transfer (REST) Application Programming Interface (API) of BioGRID database (https://thebiogrid.org/) was developed to query and access its data on the protein coding gene interaction data (physical and genetic) as well as phenotype observed and experiment type (https://github.com/Sookie-S/SGA-analysis).

Growth analysis of SNR17AΔ and SNR17BΔ mutants

A spot growth assay was performed on the single SNR17AΔ and SNR17BΔ mutants in the yeast standard media to reveal phenotypic effects of the snoRNA deletion. Three biological replicates were tested for each mutant. The growth was compared to the host strain Y7092. Cells for the spot assay were prepared in the following way: cells were grown overnight with shaking (200 rpm) and the cultures were centrifuged for 2 min at 2000 rpm. The pellet was resuspended in YPD and diluted to a starting OD_{600} of 1.0. Two and ten times dilutions were prepared and 1 and 5 µl of the cell suspensions was spotted on an agar plate which was incubated for 48h at 30°C or 37°C depending on the environmental condition. The growth was recorded every 12h of incubation.

SNR17AΔ and SNR17BΔ RNA extraction and RT-qPCR

Total RNA was extracted in the logarithmic growth phase using the RNeasy RNA isolation kit (Qiagen). Synthesis of complementary DNA was performed using QuantiTect Reverse Transcription Kit (Qiagen) following the manufacturer's instructions. Analysis of gene expression was performed using a The LightCycler® 480 real-time PCR System (Roche). The

gene *ACT1* was used as endogenous control. The Ct values obtained from three independent

biological replicates were used to calculate the relative gene expression of the target genes

according to the ΔΔCt method described by Livak and Schmittgen, 2001 (*55*). Statistical tests

were performed using Welch two sample t-test and multiple comparisons were analysed using

ANOVA followed by Dunnett's test or Tukey-Kramer test, according to the experiment. Error

bars denote standard deviations; *$p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$;* ns

= no significant change. The RT-qPCR primers are listed in Supplementary Table S1. The

SNR17A and SNR17B primers were designed to span the intron and their specificity were

verified by RT-qPCR.

Quantitative analysis of 18S rRNA processing in ncRNA deletion mutants

The concentration of processed 18S rRNA in the wild-type (WT) as well as in the single

ncRNA SNR17AΔ and SNR17BΔ mutants was performed in 5 different growth conditions.

Additionally, we analysed the 18S concentrations in the WT and SNR17AΔ, SUT480Δ,

SNR17A/SUT480Δ in the yeast standard growth condition and the respiratory stress. Total

RNA was extracted in the logarithmic growth phase as described above and the RNA was

processed using an RNA ScreenTape kit following the manufacturer's instructions. The RNA

samples were run on a TapeStation 2200 system (Agilent Technologies). The error bars were

calculated using the results from six technical replicates, and the p-values were determined

using Welch's *t*-test.

ACKNOWLEDGMENTS

This work was supported by the Wellcome Trust, under the grant number 094225 to RTO and

DD and 104981 to RTO, SGJ and DD. ST is supported by the European Research Council,

H2020-MSCA-ITN-2017, under the grant number 764364 awarded to DD

(https://cordis.europa.eu/project/id/764364). LNB was supported by The National Secretary of

Higher Education, Science, Technology and Innovation (SENESCYT,

http://siau.senescyt.gob.ec/), Ecuador and is currently funded by the Biotechnology and

Biological Sciences Research Council (BB/T002123/1) to DD. KD scholarship was funded by

the Royal Government of Thailand under the scheme for Development and Promotion of

Science and Technology Talent (DPST) projects. MWF was supported by The University of

Manchester - China Scholarship Council joint scholarship. The authors wish to thank Charlie

Boone, Michael Costanzo and Sondra Bahr for useful discussions, provision of the Y7092

strain and help with the SGA methodology. The authors also thank Diego Estrada-Rivadeneyra

for initial help with U3 snoRNA data acquisition.

DATA AVAILABILITY

648 The interactive network fitness data of the ncRNAs epistatic network has been added to the

649 publicly accessible online resource, the Yeast ncRNA Analysis (YNCA) available here:

650 http://sgjlab.org/ynca/

CODE AVAILABILITY

653 Codes used to analyse the data in this manuscript are available at https://github.com/Sookie-

654 S/SGA-analysis.

REFERENCES

1. T. H. Jensen, A. Jacquier, D. Libri, Dealing with pervasive transcription. *Mol Cell* **52**, 473-484
(2013).
2. M. Wery, M. Descrimes, N. Vogt, A. S. Dallongeville, D. Gautheret, A. Morillon, Nonsense-
Mediated Decay Restricts LncRNA Levels in Yeast Unless Blocked by Double-Stranded RNA
Structure. *Mol Cell* **61**, 379-392 (2016).
3. J. Wu, D. Delneri, R. T. O'Keefe, Non-coding RNAs in Saccharomyces cerevisiae: what is the
function? *Biochem Soc Trans* **40**, 907-911 (2012).
4. R. Boland, Non-coding RNA: It's Not Junk. *Dig Dis Sci* **62**, 1107-1109 (2017).
5. R. A. de Almeida, M. G. Fraczek, S. Parker, D. Delneri, R. T. O'Keefe, Non-coding RNAs and
disease: the classical ncRNAs make a comeback. *Biochem Soc Trans* **44**, 1073-1078 (2016).
6. U. Nagalakshmi, Z. Wang, K. Waern, C. Shou, D. Raha, M. Gerstein, M. Snyder, The
transcriptional landscape of the yeast genome defined by RNA sequencing. *Science* **320**, 1344-
1349 (2008).
7. I. A. Drinnenberg, D. E. Weinberg, K. T. Xie, J. P. Mower, K. H. Wolfe, G. R. Fink, D. P. Bartel,
RNAi in budding yeast. *Science* **326**, 544-550 (2009).
8. S. A. Shabalina, E. V. Koonin, Origins and evolution of eukaryotic RNA interference. *Trends Ecol*
Evol **23**, 578-587 (2008).
9. H. Neil, C. Malabat, Y. d'Aubenton-Carafa, Z. Xu, L. M. Steinmetz, A. Jacquier, Widespread
bidirectional promoters are the major source of cryptic transcripts in yeast. *Nature* **457**, 1038-
1042 (2009).
10. Z. Xu, W. Wei, J. Gagneur, F. Perocchi, S. Clauder-Munster, J. Camblong, E. Guffanti, F. Stutz,
W. Huber, L. M. Steinmetz, Bidirectional promoters generate pervasive transcription in yeast.
Nature **457**, 1033-1037 (2009).
11. E. L. van Dijk, C. L. Chen, Y. d'Aubenton-Carafa, S. Gourvennec, M. Kwapisz, V. Roche, C.
Bertrand, M. Silvain, P. Legoix-Ne, S. Loeillet, A. Nicolas, C. Thermes, A. Morillon, XUTs are a
class of Xrn1-sensitive antisense regulatory non-coding RNA in yeast. *Nature* **475**, 114-117
(2011).
12. D. Schulz, B. Schwalb, A. Kiesel, C. Baejen, P. Torkler, J. Gagneur, J. Soeding, P. Cramer,
Transcriptome surveillance by selective termination of noncoding RNA synthesis. *Cell* **155**,
1075-1087 (2013).

13.   Z. Bloom-Ackermann, S. Navon, H. Gingold, R. Towers, Y. Pilpel, O. Dahan, A comprehensive
tRNA deletion library unravels the genetic architecture of the tRNA pool. *PLoS Genet* **10**,
e1004084 (2014).
14.   F. Huber, D. Bunina, I. Gupta, A. Khmelinskii, M. Meurer, P. Theer, L. M. Steinmetz, M. Knop,
Protein Abundance Control by Non-coding Antisense Transcription. *Cell Rep* **15**, 2625-2636
(2016).
15.   S. Parker, M. G. Fraczek, J. Wu, S. Shamsah, A. Manousaki, K. Dungrattanalert, R. A. de
Almeida, E. Invernizzi, T. Burgis, W. Omara, S. Griffiths-Jones, D. Delneri, R. T. O'Keefe, Large-
scale profiling of noncoding RNA function in yeast. *PLoS Genet* **14**, e1007253 (2018).
16.   L. N. Balarezo-Cisneros, S. Parker, M. G. Fraczek, S. Timouma, P. Wang, R. T. O'Keefe, C. B.
Millar, D. Delneri, Functional and transcriptional profiling of non-coding RNAs in yeast reveal
context-dependent phenotypes and in trans effects on the protein regulatory network. *PLoS
Genet* **17**, e1008761 (2021).
17.   S. Parker, M. G. Fraczek, J. Wu, S. Shamsah, A. Manousaki, K. Dungrattanalert, R. A. de
Almeida, D. Estrada-Rivadeneyra, W. Omara, D. Delneri, R. T. O'Keefe, A resource for
functional profiling of noncoding RNA in the yeast Saccharomyces cerevisiae. *RNA* **23**, 1166-
1171 (2017).
18.   A. Baryshnikova, M. Costanzo, S. Dixon, F. J. Vizeacoumar, C. L. Myers, B. Andrews, C. Boone,
Synthetic genetic array (SGA) analysis in Saccharomyces cerevisiae and Schizosaccharomyces
pombe. *Methods Enzymol* **470**, 145-179 (2010a).
19.   A. H. Tong, C. Boone, Synthetic genetic array analysis in Saccharomyces cerevisiae. *Methods
Mol Biol* **313**, 171-192 (2006).
20.   A. H. Tong, G. Lesage, G. D. Bader, H. Ding, H. Xu, X. Xin, J. Young, G. F. Berriz, R. L. Brost, M.
Chang, Y. Chen, X. Cheng, G. Chua, H. Friesen, D. S. Goldberg, J. Haynes, C. Humphries, G. He,
S. Hussein, L. Ke, N. Krogan, Z. Li, J. N. Levinson, H. Lu, P. Menard, C. Munyana, A. B. Parsons,
O. Ryan, R. Tonikian, T. Roberts, A. M. Sdicu, J. Shapiro, B. Sheikh, B. Suter, S. L. Wong, L. V.
Zhang, H. Zhu, C. G. Burd, S. Munro, C. Sander, J. Rine, J. Greenblatt, M. Peter, A. Bretscher,
G. Bell, F. P. Roth, G. W. Brown, B. Andrews, H. Bussey, C. Boone, Global mapping of the yeast
genetic interaction network. *Science* **303**, 808-813 (2004).
21.   M. Costanzo, A. Baryshnikova, J. Bellay, Y. Kim, E. D. Spear, C. S. Sevier, H. Ding, J. L. Koh, K.
Toufighi, S. Mostafavi, J. Prinz, R. P. St Onge, B. VanderSluis, T. Makhnevych, F. J. Vizeacoumar,
S. Alizadeh, S. Bahr, R. L. Brost, Y. Chen, M. Cokol, R. Deshpande, Z. Li, Z. Y. Lin, W. Liang, M.
Marback, J. Paw, B. J. San Luis, E. Shuteriqi, A. H. Tong, N. van Dyk, I. M. Wallace, J. A. Whitney,
M. T. Weirauch, G. Zhong, H. Zhu, W. A. Houry, M. Brudno, S. Ragibizadeh, B. Papp, C. Pal, F.
P. Roth, G. Giaever, C. Nislow, O. G. Troyanskaya, H. Bussey, G. D. Bader, A. C. Gingras, Q. D.
Morris, P. M. Kim, C. A. Kaiser, C. L. Myers, B. J. Andrews, C. Boone, The genetic landscape of
a cell. *Science* **327**, 425-431 (2010).
22.   D. Kyriakou, E. Stavrou, P. Demosthenous, G. Angelidou, B. J. San Luis, C. Boone, V. J.
Promponas, A. Kirmizis, Functional characterisation of long intergenic non-coding RNAs
through genetic interaction profiling in Saccharomyces cerevisiae. *BMC Biol* **14**, 106 (2016).
23.   A. C. Tuck, D. Tollervey, A transcriptome-wide atlas of RNP composition reveals diverse classes
of mRNAs and lncRNAs. *Cell* **154**, 996-1009 (2013).
24.   J. Camblong, N. Beyrouthy, E. Guffanti, G. Schlaepfer, L. M. Steinmetz, F. Stutz, Trans-acting
antisense RNAs mediate transcriptional gene cosuppression in S. cerevisiae. *Genes Dev* **23**,
1534-1545 (2009).
25.   J. A. Martens, L. Laprade, F. Winston, Intergenic transcription is required to repress the
Saccharomyces cerevisiae SER3 gene. *Nature* **429**, 571-574 (2004).
26.   J. Camblong, N. Iglesias, C. Fickentscher, G. Dieppois, F. Stutz, Antisense RNA stabilization
induces transcriptional gene silencing via histone deacetylation in S. cerevisiae. *Cell* **131**, 706-
717 (2007).

27. P. Thebault, G. Boutin, W. Bhat, A. Rufiange, J. Martens, A. Nourani, Transcription regulation
by the noncoding RNA SRG1 requires Spt2-dependent chromatin deposition in the wake of
RNA polymerase II. *Mol Cell Biol* **31**, 1288-1300 (2011).

28. F. J. van Werven, G. Neuert, N. Hendrick, A. Lardenois, S. Buratowski, A. van Oudenaarden, M.
Primig, A. Amon, Transcription of two long noncoding RNAs mediates mating-type control of
gametogenesis in budding yeast. *Cell* **150**, 1170-1181 (2012).

29. M. Usaj, Y. Tan, W. Wang, B. VanderSluis, A. Zou, C. L. Myers, M. Costanzo, B. Andrews, C.
Boone, TheCellMap.org: A Web-Accessible Database for Visualizing and Mining the Global
Yeast Genetic Interaction Network. *G3 (Bethesda)* **7**, 1539-1549 (2017).

30. J. van Leeuwen, C. Pons, J. C. Mellor, T. N. Yamaguchi, H. Friesen, J. Koschwanez, M. M. Usaj,
M. Pechlaner, M. Takar, M. Usaj, B. VanderSluis, K. Andrusiak, P. Bansal, A. Baryshnikova, C. E.
Boone, J. Cao, A. Cote, M. Gebbia, G. Horecka, I. Horecka, E. Kuzmin, N. Legro, W. Liang, N.
van Lieshout, M. McNee, B. J. San Luis, F. Shaeri, E. Shuteriqi, S. Sun, L. Yang, J. Y. Youn, M.
Yuen, M. Costanzo, A. C. Gingras, P. Aloy, C. Oostenbrink, A. Murray, T. R. Graham, C. L. Myers,
B. J. Andrews, F. P. Roth, C. Boone, Exploring genetic suppression interactions on a global
scale. *Science* **354**, (2016).

31. D. Segre, A. Deluna, G. M. Church, R. Kishony, Modular epistasis in yeast metabolism. *Nat*
Genet **37**, 77-83 (2005).

32. E. S. Maxwell, M. J. Fournier, The small nucleolar RNAs. *Annu Rev Biochem* **64**, 897-934 (1995).

33. J. L. Woolford, Jr., S. J. Baserga, Ribosome biogenesis in the yeast Saccharomyces cerevisiae.
Genetics **195**, 643-681 (2013).

34. J. M. Hughes, D. A. Konings, G. Cesareni, The yeast homologue of U3 snRNA. *EMBO J* **6**, 2145-
2155 (1987).

35. A. M. Kiss, B. E. Jady, E. Bertrand, T. Kiss, Human box H/ACA pseudouridylation guide RNA
machinery. *Mol Cell Biol* **24**, 5797-5807 (2004).

36. E. Sharma, T. Sterne-Weiler, D. O'Hanlon, B. J. Blencowe, Global Mapping of Human RNA-RNA
Interactions. *Mol Cell* **62**, 618-626 (2016).

37. P. Vitali, T. Kiss, Cooperative 2'-O-methylation of the wobble cytidine of human elongator
tRNA(Met)(CAT) by a nucleolar and a Cajal body-specific box C/D RNP. *Genes Dev* **33**, 741-746
(2019).

38. C. Huang, J. Shi, Y. Guo, W. Huang, S. Huang, S. Ming, X. Wu, R. Zhang, J. Ding, W. Zhao, J. Jia,
X. Huang, A. P. Xiang, Y. Shi, C. Yao, A snoRNA modulates mRNA 3' end processing and
regulates the expression of a subset of mRNAs. *Nucleic Acids Res* **45**, 8647-8660 (2017).

39. S. Kishore, S. Stamm, The snoRNA HBII-52 regulates alternative splicing of the serotonin
receptor 2C. *Science* **311**, 230-232 (2006).

40. E. Myslinski, V. Segault, C. Branlant, An intron in the genes for U3 small nucleolar RNAs of the
yeast Saccharomyces cerevisiae. *Science* **247**, 1213-1216 (1990).

41. C. M. Paget, J. M. Schwartz, D. Delneri, Environmental systems biology of cold-tolerant
phenotype in Saccharomyces species adapted to grow at different temperatures. *Mol Ecol* **23**,
5241-5257 (2014).

42. H. S. Jeong, P. S. Backlund, H. C. Chen, A. A. Karavanov, R. J. Crouch, RNase H2 of
Saccharomyces cerevisiae is a complex of three proteins. *Nucleic Acids Res* **32**, 407-414 (2004).

43. A. Force, M. Lynch, F. B. Pickett, A. Amores, Y. L. Yan, J. Postlethwait, Preservation of duplicate
genes by complementary, degenerate mutations. *Genetics* **151**, 1531-1545 (1999).

44. A. Stoltzfus, On the possibility of constructive neutral evolution. *J Mol Evol* **49**, 169-181 (1999).

45. A. Wach, A. Brachat, R. Pohlmann, P. Philippsen, New heterologous modules for classical or
PCR-based gene disruptions in Saccharomyces cerevisiae. *Yeast* **10**, 1793-1808 (1994).

46. A. M. Chu, R. W. Davis, High-throughput creation of a whole-genome collection of yeast
knockout strains. *Methods Mol Biol* **416**, 205-220 (2008).

47. A. Baryshnikova, M. Costanzo, Y. Kim, H. Ding, J. Koh, K. Toufighi, J. Y. Youn, J. Ou, B. J. San
Luis, S. Bandyopadhyay, M. Hibbs, D. Hess, A. C. Gingras, G. D. Bader, O. G. Troyanskaya, G.

788   W. Brown, B. Andrews, C. Boone, C. L. Myers, Quantitative analysis of fitness and genetic
789   interactions in yeast on a genome scale. *Nat Methods* **7**, 1017-1024 (2010b).
790   48.   B. P. Young, C. J. Loewen, Balony: a software package for analysis of data generated by
791   synthetic genetic array experiments. *BMC Bioinformatics* **14**, 354 (2013).
792   49.   R. C. team. (2021).
793   50.   O. Wagih, M. Usaj, A. Baryshnikova, B. VanderSluis, E. Kuzmin, M. Costanzo, C. L. Myers, B. J.
794   Andrews, C. M. Boone, L. Parts, SGAtools: one-stop analysis and visualization of array-based
795   genetic interaction screens. *Nucleic Acids Res* **41**, W591-596 (2013).
796   51.   B. T. a. T. R. Almende B.V. and Contributors. (2021).
797   52.   N. T. Csardi G, The igraph software package for complex network research, InterJournal,
798   Complex Systems 1695. 2006. (2006).
799   53.   M. Jacomy, T. Venturini, S. Heymann, M. Bastian, ForceAtlas2, a continuous graph layout
800   algorithm for handy network visualization designed for the Gephi software. *PLoS One* **9**,
801   e98679 (2014).
802   54.   A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features.
803   *Bioinformatics* **26**, 841-842 (2010).
804   55.   K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
805   quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).
806
807

 FIGURE LEGENDS

FIG. 1. ncRNA SGA network for 38 query strains. The SGA network was generated based on
the absolute fold change ≥2 and q-value ≤0.001 cut off for 38 query ncRNA deletion strains.
The nodes are coloured by class of ncRNA. Orange: SUTs; red: CUTs; purple: snoRNAs;
green: tRNAs. (**A**) Total epistatic interaction network between ncRNAs. (**B**) Positive
interactions between ncRNA deletions. (**C**) Negative synthetic sick interactions. (**D**) Negative
synthetic lethal interactions. The node size is proportional to the number of interactions they
are involved in. Black and grey edges represent negative and positive epistasis, respectively.
The thickness of the edges is proportional to the strength of the interaction.

FIG. 2. Analysis of the interactions of genes neighbouring epistatic pairs of ncRNAs. (**A**)
Strategy to analyse genetic and physical interactions between the neighbouring protein coding
genes (green arrows) of ncRNA pairs showing significant epistasis (orange). (**B**) Pie charts
displaying the proportion of significantly interacting neighbouring genes (dark blue colour),
broken down in genetic (mid blue) and physical interactions (light blue).

FIG. 3. Visualization of the ncRNA network under different environmental condition. ncRNA
network generated for 31 query ncRNA deletion strains based on the absolute fold change ≥2
and q–value ≤0.001 cut off in (**A**) YPD, 30°C, (**B**) YPD, 37°C, (**C**) YP+7% Glycerol, 30°C,
(**D**) YPD+1 M NaCl, 30°C and (**E**) YPD+4 mM H_2O_2, 30°C. Network ncRNAs nodes (blue);
red edges – negative ncRNA genetic interactions, green edges – positive ncRNA genetic
interactions. The thickness of the edges is proportional to the absolute fold change.

FIG. 4. Environmental conditions elicit yeast ncRNA interaction network changes. (**A**) Upset
plot depicting environmental unique and shared ncRNA genetic interactions between the five
conditions tested (YPD, 37°C; YPD, 30°C; YPD+1 M NaCl, 30°C; YPD+4 mM H_2O_2, 30°C;
and YP+7% glycerol, 30°C). The bottom left hand side horizontal bar chart displays the total
number of significant interactions for each condition tested. The top vertical bar chart displays
the number of interactions unique or shared between conditions indicated by a black dot in the
bottom chart box. Unique interactions are represented by a single black dot between all 5
conditions tested. Several dots connected with a black line represent the number of interactions
shared in the selected conditions. (**B**) Heatmap depicting fold change of 86 ncRNA interactions
that display significant change in the five conditions tested compared to rich medium. Specific
interactions are arranged on the y-axis. Conditions are displayed on the x-axis. Negative

interactions are represented as shades of blue. Positive interactions are represented as shades of yellow. **c** Visual representation of fitness for selected single and double deletion mutants in the indicated conditions.

FIG. 5. Area proportional Venn diagrams displaying unique and shared interactions between SNR17A and SNR17B. Unique and shared interactions between SNR17A and SNR17B (q<0.1) are displayed in (**A**) YPD, 30°C; (**B**) YPD, 37°C; (**C**) YP+7% Glycerol, 30°C; (**D**) YPD+1 M NaCl, 30°C and (**E**) YPD+4 mM H_2O_2, 30°C. Light grey (centre) – shared interactions between SNR17A and SNR17B; mid grey (left) – unique interactions to SNR17A; dark grey (right) – unique interactions to SNR17B.

FIG. 6. Relative expression levels of SNR17A and SNR17B measured by RT-qPCR. (**A**) RNA levels measured in haploid WT strains BY4741 and Y7092, and in the natural diploid strain 96.2 in YPD, 30°C. (**B**) RNA levels measured in BY4741 under different conditions: YP+7% glycerol, 30°C; YPD, 37°C; YPD+1 M NaCl, 30°C and YPD+4 mM H_2O_2, 30°C. **c** relative expression of SNR17A and SNR17B in the SNR17AΔ and SNR17BΔ mutants in YPD, 30°C. The expression of snoRNAs in the deletion mutants was compared to the expression in the BY4741 background. The expression of SNR17B is significant higher (***p <0.001) in the SNR17AΔ strain compared to the WT strain. The relative quantities of SNR17A and SNR17B were calculated by the ΔΔCt method using *ACT1* as reference. Error bars represent standard deviation.

FIG. 7. Analysis of 18S rRNA concentrations in WT, SNR17AΔ, SNR17BΔ, SUT480Δ and SNR17AΔ/SUT480Δ. Concentrations of 18S rRNA in the WT and deletion mutants was analysed using the Tapestation 2200 system. (**A**) Comparison of 18S concentration for the WT, SNR17AΔ and SNR17BΔ in 5 different conditions. (**B**) Analysis of 18S concentrations in the SUT480Δ and SNR17AΔ/SUT480Δ mutants in two conditions.

SUPPORTING FIGURES

FIG. S1. Query strain selection for SGA. Heat-map of the solid fitness of ncRNA deletion strains used as query strains in the SGA. (**A**) Fitness of haploid deletion strains based on colony size normalized to WT. (**B**) Competitive fitness of diploid heterozygous ncRNAs grown under chemostat culture conditions. Rows represent the different growth conditions and columns

875 represent the ncRNAs deletions. The colour legends represent the colony size normalized to
876 the WT (**A**) and the difference of growth of the mutants between the end and the start of the
877 competition (**B**). Fitness reduction or increase is represented as shades of red or green,
878 respectively. The data were taken from (*15, 16*).
879

880 **FIG. S2.** Heatmap depicting 2-fold change of 395 ncRNA double ncRNA deletions that shared
881 genetic interactions between five environmental conditions tested. Deletion mutants are
882 arranged on the y-axis and conditions are displayed on the x-axis. Negative interactions are
883 represented as shades of blue and positive interactions are represented as shades of yellow.
884

885 **FIG. S3.** Scatter plot displaying correlation between fitness and the number of interactions.
886 Number of interactions were correlated with the fitness changes in (**A**) YPD, 30°C, (**B**) YPD,
887 37°C, (**C**) YP+7% glycerol, 30°C and (**D**) YPD+4 mM H_2O_2, 30°C.
888

FIG. S4. Genetic and physical interactions in environmental conditions. Pie chart
representation of genetic and physical interactions analysed between the neighbouring genes
of the significantly interacting ncRNAs**.** Dark blue colour in the larger pie charts represents the
percentage of ncRNAs protein-coding neighbouring genes with a significant type of
interactions. Smaller pie charts display the proportion of these interactions broken down into
genetic (mid blue) and physical interactions (light blue) in YPD, 30°C; YPD, 37°C; YP+7%
glycerol, 30°C; YPD+1 M NaCl, 30°C; and YPD+4 mM H_2O_2, 30°C.
896

897 **FIG. 5.** Spot assay to determine growth differences between SNR17A and SNR17B deletion
898 strains. Serial dilutions (2x for **A**, and 10x for **B** to **E**) of three biological replicates of haploid
899 SNR17AΔ and SNR17BΔ mutants along with a host strain Y7092 were spotted on YPD, 30°C
900 (**A**), YPD, 37°C (**B**), YP+7% glycerol, 30°C (**C**), YPD+4 mM H_2O_2, 30°C (**D**) and YPD+1 M
901 NaCl, 30°C (**E**). Growth was recorded after 24h.
902

FIG. S6. Area proportional Venn diagrams displaying unique and shared interactions between
SNR17A and SNR17B at different q-values. Unique and shared interactions between SNR17A
and SNR17B are displayed at q<0.01 and <0.001. (**A**) SGA data in YPD, 30°C; (**B**) YPD, 37°C;
(**C**) YP+7% Glycerol, 30°C; (**D**) YPD+1 M NaCl, 30°C and **e** YPD+4 mM H_2O_2, 30°C. Light

grey (centre) – shared interactions between SNR17A and SNR17B; mid grey (left) – unique

interactions to SNR17A; dark grey (right) – unique interactions to SNR17B.

FIG. S7. SUT480 does not altered SNR17A and SNR17B RNA levels. RNA levels of

SNR17A and SNR17B were analysed in the SUT480 deletion mutant in BY4741 (**A**) and

Y9072 background (**B**). Quantification of the RNA levels of SUT480 in *SNR17AΔ* and

SNR17BΔ strains was also carried out (**C**). mRNA levels of *RNH202* in a SUT480Δ deletion

in BY4741 and Y7092 strains (**D**). RNA was collected in YPD at 30$^{\text{o}}$C, RNA levels were

quantified by RT-qPCR and compared using t-test or ANOVA

FIG. S8. Construction and validation of query strains. (**A**) Strategy to construct and validate

ncRNA query deletion mutants. Briefly, ncRNA deletion cassette composed of the natMX4

marker and ncRNA complementary flanking regions were transformed into the MATα strain

(Y7092) to delete the ncRNAs of interest. Following transformation, single colonies were PCR

validated with primer sets: (i) confA + confNatR, (ii) confNatF + confD and (iii) confA +

confB. Primer sets (i) and (ii) only amplified PCR bands if the deletion cassettes were

integrated at the correct loci. Primer set (iii) amplified a PCR band if no integration event

occurred. (**B**) Agarose gel electrophoresis for the PCR validation of randomly selected ncRNA

query mutants. U: upstream flank PCR with primer set confA + confNatR; D: downstream

flank PCR with primer set confNatF + confD; N: control PCR for the native ncRNA, with

primer set confA + confB; M: 1 kb New England Biolabs ladder.

FIG. S9. Validation of a subset of SGA double deletion mutants. (**A**) Strategy to validate the

double ncRNA deletion mutants. Following SGA, randomly selected double deletion haploid

mutants were subjected to confirmation PCR. Primer sets (i) confA + confKanR and (ii)

confKanF + confD were used to test for the *kanMX4* deletion cassette integrations. Primer sets

(iii) confA + confNatR and (iv) confD + confNatF were used to test for the *natMX4* deletion

cassette integrations. Primers confA + confB provided additional ncRNA deletion

confirmation, only amplifying bands if the ncRNAs were intact. (**B**) Agarose gel

electrophoresis for PCR validation on a selection of mutants. U: upstream flank PCR with

primer set confA + kanB or confNatR; D: downstream flank PCR with primer set kanC or

confNatF + confD; N: control PCR for the native ncRNA with primer set confA + confB. M:

1 kb New England Biolabs ladder.

Fig. 1

Fig. 2

Fig. 3

a YPD, 30°C

b YPD, 37°C

c YP+7% Glycerol, 30°C

d YPD+1 M NaCl, 30°C

e YPD+4 mM H$_2$O$_2$, 30°C

○ ncRNA

━━ negative genetic interactions

━━ positive genetic interactions

Fig. 4

a

c

Positive interaction Negative interaction No change (not significant)

b

Fig. 5

Fig. 6

Fig. 7

Links to Supplementary Materials

(GitHub: https://github.com/Sookie-S/SGA-analysis)

Supplementary Dataset S1
Supplementary Dataset S2
Supplementary Dataset S3
Supplementary Dataset S4
Supplementary Dataset S5
Supplementary Dataset S6
Supplementary Dataset S7
Supplementary Dataset S8

Supplemetary Table S1

Supplemetary Fig. 1
Supplemetary Fig. 2
Supplemetary Fig. 3
Supplemetary Fig. 4
Supplemetary Fig. 5
Supplemetary Fig. 6
Supplemetary Fig. 7
Supplemetary Fig. 8
Supplementary Fig. 9