1 **Structural variation turnovers and defective genomes: key drivers for the *in***
2 ***vitro* evolution of the large double-stranded DNA koi herpesvirus (KHV)**
3
4

Nurul Novelia Fuandila[1], Anne-Sophie Gosselin-Grenet[2], Marie-Ka Tilak[1], Sven M Bergmann[3],
Jean-Michel Escoubas[4], Sandro Klafack[5], Angela Mariana Lusiastuti[6], Munti Yuhana[7], Anna-
Sophie Fiston-Lavier[1,8], Jean-Christophe Avarre[1*], Emira Cherif[1*]

[1]ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France
[2]DGIMI, Univ Montpellier, INRAE, Montpellier, France
[3]Institute of Infectology, Friedrich-Loeffer-Institut, Federal Research Institute for Animal Health,
Greifswald-Insel Riems, Germany
[4]IHPE, Univ Montpellier, CNRS, Ifremer, UPVD, Montpellier, France
[5]Department of Experimental Animal Facilities and Biorisk Management, Friedrich-Loeffer-
Institut, Federal Research Institute for Animal Health, Greifswald-Insel Riems, Germany
[6]Fish Health Laboratory, Research Institute for Freshwater Aquaculture and Fisheries
Extension, Bogor, Indonesia
[7] Faculty of Fisheries and Marine Sciences, Bogor Agricultural University, Indonesia
[8]Institut Universitaire de France (IUF)

* corresponding authors: jean-christophe.avarre@ird.fr; emira.cherif@ird.fr

Abstract

Structural variations (SVs) constitute a significant source of genetic variability in virus genomes.
Yet knowledge about SV variability and contribution to the evolutionary process in large double-
stranded (ds)DNA viruses is limited. Cyprinid herpesvirus 3 (CyHV-3), also commonly known as
koi herpesvirus (KHV), has the largest dsDNA genome within herpesviruses. This virus has
become one of the biggest threats to common carp and koi farming, resulting in high morbidity
and mortalities of fishes, serious environmental damage, and severe economic losses. A
previous study analyzing CyHV-3 virulence evolution during serial passages onto carp cell
cultures suggested that CyHV-3 evolves, at least *in vitro*, through an assembly of haplotypes
that alternatively become dominant or under-represented. The present study investigates the
SV diversity and dynamics in CyHV-3 genome during 99 serial passages in cell culture using,
for the first time, ultra-deep whole-genome and amplicon-based sequencing. The results
indicate that KHV polymorphism mostly involves SVs. These SVs display a wide distribution
along the genome and exhibit high turnover dynamics with a clear bias towards inversion and

deletion events. Analysis of the pathogenesis-associated ORF150 region in ten intermediate cell
passages highlighted mainly deletion, inversion and insertion variations that deeply altered the
structure of ORF150. Our findings indicate that SV turnovers and defective genomes represent
key drivers in the viral population dynamics and *in vitro* evolution of KHV. Thus, the present
study can contribute to the basic research needed to design safe live-attenuated vaccines,
classically obtained by viral attenuation after serial passages in cell culture.
41
42
43   **Keywords**
44   KHV, virus evolution, virulence, virus attenuation, structural variations, defective genome, carp
45
46
Introduction

Viruses have a remarkable ability to adapt to the complex and hostile host immune and
physiological constraints. Such capability is directly associated with the viral population's genetic
diversity, and deep characterization of this diversity is the cornerstone of our understanding of
virus evolutive response to a new cellular environment. The starkly evident examples are RNA
viruses. These viruses, such as HIV, hepatitis C, and influenza, display high mutation rates
which generate significant polymorphism levels, allowing the viral population to quickly adapt to
newly infected cellular environments and evolve strategies against vaccines and antiviral drugs
(Lauring and Andino 2010; Loiseau et al., 2020). However, the genetic diversity has been
thoroughly characterized in only a handful of viruses, mainly targeting SNPs (Single Nucleotide
Polymorphisms) in RNA viruses (Sanjuán and Domingo-Calap 2016). Although the mutation
rate of large double-stranded (ds) DNA viruses is up to four folds lower than that of RNA
viruses, due to the use of high-fidelity proofreading polymerases, and most SNPs in dsDNA
viruses are neutral and at low frequency, SNP-based approaches were chosen to analyze the
genetic diversity in Human cytomegalovirus (HCMV), Autographa californica multiple
nucleopolyhedrovirus (AcMNPV) and herpes simplex virus 2 (HSV-2), for example (Renzette et
al., 2015; Chateigner et al., 2015; Akhtar et al., 2019).
Structural variations (SVs) play a key role in viral evolutionary processes. Genome
rearrangements such as deletions, insertions, duplications and inversions can lead to defective
viral genomes (DVGs) (O'Hara et al., 1984; Molenkamp et al., 2000; Vignuzzi and López 2019).
Preben von Magnus first identified DVGs in the late 40's as incomplete influenza viruses that

can interfere with the wild-type virus replication (Vignuzzi and López 2019). Since then, the role

of DVGs in antiviral immunity, viral persistence and their negative impact on virus replication

and production has been established (Bull et al., 2003; Li et al., 2011; Vignuzzi and López 2019;

Loiseau et al., 2020). Nowadays, DVGs have been described in most RNA viruses and to a

lesser extent in dsDNA viruses (Vignuzzi and López 2019; Loiseau et al., 2020). Despite the

critical role of SVs in virus infection dynamics, the knowledge about structural variation diversity,

and their evolutionary impact in viral populations, especially those with large dsDNA, is limited.

The large dsDNA Cyprinid herpesvirus 3 (CyHV-3), more commonly known as koi herpesvirus

(KHV), is one of the most virulent viruses of fish. It is a lethally infectious agent that infects

common carp and koi (*Cyprinus carpio*) at all stages of their life (Hedrick et al., 2000; Haenen et

al., 2004). KHV infections are usually associated with high morbidities and mortalities (up to

95%), resulting in serious environmental damages and severe economic losses (Sunarto et al.,

2011; Rakus et al., 2013). This threatening virus had a rapid worldwide spread due to global fish

trade and international ornamental koi exhibitions (Gotesman et al., 2013). Classified within the

family *Alloherpesviridae*, genus *Cyprinivirus*, CyHV-3 is the subject of an increasing number of

studies and has become the archetype of alloherpesviruses (Boutier et al., 2015). Despite this

"status", only 19 isolates have been entirely sequenced so far (source: NCBI) since the release

of the first complete genome sequences in 2007 (Aoki et al., 2007). Such a low number of full

genomes impairs large-scale phylogenomic studies (Gao et al., 2018). On the other hand, KHV

infections have been shown to be the result of haplotype mixtures, both *in vivo* and *in vitro*

(Hammoumi et al, 2016; Klafack et al, 2019). If mixed-haplotype infections probably represent

an additional source of diversification for KHV (Renner and Szpara, 2018), they make genomic

comparisons more challenging.

KHV has the largest genome among all known herpesviruses, with a size of approximately 295

kb and 156 predicted open reading frames (ORFs) (Aoki et al. 2007). Several studies focusing

on the analysis of viral ORFs have shown the implication of some of them in KHV virulence

(Boutier et al., 2015; Fuchs et al. 2014). KHV isolates are known to carry mutations in ORFs

that are likely to alter gene functions, and these mutations may vary from virus to virus (Gao et

al., 2018) and even within viruses (Hammoumi et al., 2016). Aoki et al (2007) hypothesized that

virulent KHV would have arisen from a wild-type ancestor by loss of gene function. However,

nearly 15 years later, this hypothesis has still not been tested, probably because of the lack of

extensive genomic comparisons. SVs may play a key role in this gene function loss, as recently

shown by Klafack et al (2019). These authors conducted a comparative study of a cell culture-

propagated isolate that suggested that CyHV-3 evolves through an assemblage of haplotypes
whose composition changes within cell passages. This study revealed a deletion of 1,363 bp in
the ORF150 of the majority of haplotypes after 78 passages (P78), which was not detected after
99 passages. Furthermore, experimental infections showed that the virus passaged 78 times
was much less virulent compared to the original wild-type on the one hand and slightly less
virulent compared to the same virus passaged 99 times (P99), highlighting the potentially
important role of the ORF150 in the virulence of KHV. Besides, this study demonstrated that
haplotype assemblies evolve very rapidly along successive *in vitro* cell passages during
infectious cycles, and raised many questions regarding the mechanisms leading to such rapid
gene loss and gain *in vitro*.
The present study sought to characterize the SV diversity and dynamics in the KHV genome
using viruses propagated onto cell cultures. First, P78 and P99 whole virus genomes were
sequenced using ultra-deep long-read sequencing, a first with KHV. Then, the obviously
pathogenesis-associated ORF150 region (~5 kb) was sequenced in ten intermediate successive
cell passages through an Oxford nanopore® amplicon-based sequencing approach to gain
insights into the gene loss and gain mechanisms.
118

Material and methods

Extraction of high molecular weight DNA from P78 and P99 cell culture passages
The virus isolate used in this study was the same as that previously described in Klafack et al.
(2019), *i.e.* an isolate collected from an infected koi in Taiwan (KHV-T) and passed 99 times
onto common carp brain (CCB) cells. Considering previous results, a special focus was made
on passages 78 (P78) and 99 (P99). Genomic DNA was extracted from cell cultures stored at -
80°C, using the MagAttract HMW DNA Kit (Qiagen). Each frozen culture was thawed quickly in
a 37°C water bath, equilibrated to room temperature (25°C) and divided into 12 cell culture
aliquots of 250 µL. Tubes were centrifuged at 3,000× g for 1 minute and supernatants were
transferred into new 2-mL tubes containing 200 µL of proteinase K and RNase A solution. DNA
was subsequently extracted according to the manufacturer's recommendations and eluted in
200 µL distilled water provided in the kit. The 12 replicates of each sample were pooled together
and evaporated at room temperature using a vacuum concentrator, to reach a final volume of
around 60 µL. Concentrated DNA was quantified by fluorometry (Qubit, ThermoFisher Scientific)
and its quality was evaluated by spectrophotometry (Nanodrop 2100) and agarose gel
electrophoresis. The final concentration of P78 and P99 was 14.4 and 2.6 ng·µL^{-1}, respectively.

Quantitative PCR assays

Quantitative PCR (qPCR) was applied to evaluate cellular and viral DNA ratio. Two sets of primers were used: primers targeting the ORF150 of CyHV-3 (GenBank #AP008984.1, KHV-J, nt 259,965-260,110: 5'-GAGCGAGGAACTCTACACAAC-3' and 5'-GGTAAGGGTAAAGCAGACCATC-3') and primers targeting the glucokinase gene of *Cyprinus carpio* (GenBank #AF053332.2, nt 225-293: 5'-ACTGCGAGTGGAGACACAT-3' and 5'-TCAGGTGTGGAGGGGACAT-3'). Amplification reactions contained 1 µL of 2X SYBR Green I Master mix (Roche), 200 nM of each primer, and 1 µl of template DNA in a final volume of 10 µL. Amplifications were carried out in a LightCycler 480 (Roche) and cycling conditions consisted in an initial denaturation at 95°C for 5 min followed by 45 cycles of amplification at 95°C for 10 sec, annealing at 60°C for 20 sec and elongation at 72°C for 10 sec with a single fluorescence measurement. After amplification, a melting step was applied, which comprised a denaturation at 95°C for 5 sec, a renaturation at 65°C for 60 sec and a heating step from 65 to 97°C with a ramp of 0.1°C per second and a continuous fluorescence acquisition. Specificity of amplification was verified by visual inspection of the melting profiles, and the ratio between cellular and viral DNA was estimated as $2^{-\Delta Cq}$, assuming that each primer pair has an amplification efficiency close to 2 and that each amplicon is present as a single copy per genome.

Genomic library preparation and Oxford Nanopore whole genome sequencing

High-quality genomic DNA from the two samples (P78 and P99) was sequenced using Oxford Nanopore technology®. DNA libraries were prepared using the Ligation Sequencing Kit (SQK-LSK109) according to the manufacturer's instructions (Oxford Nanopore®). A total input amount of 48 µL (corresponding to 692 ng of P78 and 124 ng of P99 high molecular weight DNA) was used for sequencing library preparation. DNA was first end-repaired using NEBNext FFPE DNA Repair Mix and NEBNext Ultra II End repair, and then cleaned up with Agencourt AMPure XP beads (Beckman Coulter Inc) at a 1:1 bead to DNA ratio. Sixty-one µL of clean-up elution were transferred into a new 1.5-mL tube for subsequent adapter ligation. Adapter ligation was achieved using NEBNext Quick T4 DNA ligase adapter mix (AMX), ligation buffer (LNB), Long Fragment Buffer (LFB) and Elution Buffer (EB), following the provider's recommendations. The quantity of the retained DNA fragments was measured again by Qubit fluorometry. The final amount of P78 and P99 DNA was 346 ng and 94 ng, respectively. Each library was directly

168 sequenced on a separate R9.4.1 flow cell using a MinION sequencing device. Sequencing runs

169 were controlled with MinKNOW version 0.49.3.7 and operated for about 30 hours.

Amplicon-based Oxford nanopore sequencing

To specifically investigate ORF150 region, a fragment of ~4.3 kb encompassing the whole

ORF150 of CyHV-3 and part of the ORF149 and ORF 151 (nt 257,103-261,345 according to

GenBank #AP008984) was sequenced from various intermediate passages of KHV-T (P10,

P20, P30, P40, P50, P70, P78, P80, P90 and P99). A ligase-free protocol was used to limit the

risk of potential artifacts linked with sample preparation, e.g. the creation of chimeric sequences

during the end-repair or ligation steps (White et al., 2017). Genomic DNA was extracted from

cell cultures stored at -80°C, using the Nucleospin virus kit (Macherey-Nagel). Purified DNA was

subsequently   used   for   PCR   amplification   with   the   following   primers: 5'-

TGGGCGCAATCAAGATGT-3' (F) and 5'-TGAAGTGTCAGGGTCAGAGT-3' (R). PCR was

performed with GoTaq G2 DNA Polymerase (Promega) in a final volume of 40 µL containing 1

µL of total genomic DNA, 10 µL of Taq Buffer, 5 µL of dNTPs 2 mM, 2.5 µL of $MgCl_2$ and 1.25

µL of each primer (10 µM). Cycling conditions were as follows: initial denaturation at 95°C for 10

min, amplification with 40 cycles of 95°C for 10 sec, 60°C for 20 sec, 72°C for 3 min, and final

extension at 72°C for 5 min. PCR products were purified using 1X Agencourt AMPure XP

beads, tested for purity using the NanoDrop™ One spectrophotometer (ThermoFisher

Scientific), and quantified fluorometrically using the Qubit dsDNA High sensitivity kit. DNA

libraries   were   prepared   using   the   Rapid   Barcoding   kit   (SQK-RBK004),   following   the

manufacturer's instructions. For each sample, 400 ng of purified amplicon were adjusted with

nuclease-free water to a total volume of 7.5 µL and supplemented with 2.5 µL of Fragmentation

Mix RB01-4 (one for each sample). Four barcoded samples were combined with an equimolar

ratio by mixing 2.5 µL of each sample in a total volume of 10 µL. Furthermore, two additional

barcodes RB01-2 were used with the passages P78 and P99 as described above. Pooled

libraries were sequenced on 3 R9.4.1 flow cells (4 barcodes, 4 barcodes, 2 barcodes) for 24

hours and sequencing runs were controlled with MinKNOW version 0.49.3.7.

DNA sequence analysis

For each sample, bases from raw FAST5 files with a *pass* sequencing tag were recalled using

the high-accuracy model of ONT Guppy basecalling software version 4.0.15 which improves the

basecalling accuracy. The obtained fastq files were filtered to keep reads with a length ≥ 2 kb, in

order to span large structural variants. After the mapping step using minimap2 (Li, 2018), the
sequencing depth was calculated for each sample using the *plotCoverage* tool implemented in
deepTools2.0 tool suite (Ramírez et al*.,* 2016). Sequencing coverage was assessed with the
bamCoverage tool from the same tool suite and normalized using the RPGC (reads per genome
coverage) method (Figure 1).
205

Structural variant detection
To detect structural variants (SVs) in the P78 and P99 whole genomes, a mapping step followed
by BAM filtering was performed. Two aligners were used to map the raw long-reads against the
KHV-J AP008984.1 reference genome: minimap2 (Li, 2018) and NGMLR (Sedlazeck et al.,
2018). BAM files were then filtered using the option '-F' of Samtools view (Li et al., 2018) with
the flag "4" to keep only mapped reads and with the flag "0x800" to remove chimeric reads
(inconsistent/supplemental mapping). For P78 and P99, 99.33% and 97.77% of reads were
mapped, respectively. Chimeric reads represented 28.05% and 17.74% of the mapped reads in
P78 and P99, respectively. The resulting filtered BAM files from each mapper were used as
input data for SV caller, Sniffles (Sedlazeck et al., 2018). Only SVs ≥ 30 bp and supported by at
least 10 reads were kept in the final VCF files. A cross-validation step was performed using
SURVIVOR (Jeffares et al., 2017) by extracting common SVs from each mapper/caller
combination for each sample. Although the KHV-J reference genome used for the mapping is
phylogenetically close to the KHV-T isolate, some genetic diversity exists (Klafack et al. 2017).
Hence, to exclude inter-isolate SVs, a pairwise comparison between P78 and P99 was made.
The distribution of the different SVs along with P78 and P99 genomes was assessed by
estimating their occurrences using a 5 kb sliding window. SNPs and Indels variants were called
in P78 and P99 by *medaka_variant* implemented in medaka (1.4.4) using KHV-J AP008984.1
as a reference genome. To detect structural variants in the amplified region (257,103-261,345)
of P10 to P99 samples, a size filtering step using *guppyplex* was added to the steps described
above. Only reads from 1.5 kb to 8 kb were used for the analysis.
227

Results

Main features of sequencing data for P78 and P99
A total of 4,900,000 and 2,293,830 long-reads were obtained for P78 and P99, respectively.
After filtering, 462,982 long-reads with an average length of 4.96 kb were retained for P78, and
418,034 reads with an average length of 7.06 kb for P99 (Table 1, Table S1).100% of the

sampled bases from the P78 genome had at least 5,000 overlapping reads and 100% of the
sampled bases from the P99 genome had at least 7,500 overlapping reads (Figure S2). Both
P78 and P99 genomes were entirely covered by the sequencing data (Figure 1).
The ratio between cellular and viral DNA, calculated by qPCR, was 34800 and 33200 for P78
and P99, and the percent of mapped reads 99.33% and 97.77%, respectively.
238
239
240

241 **Figure 1.** Normalized sequencing coverage for P78 and P99 samples using the RPGC (reads
242 per genome coverage) method. Both P78 and P99 genomes were totally covered by the
243 sequencing. For P78, the coverage break (green triangle) corresponds to the 1.3 kb deletion.
250 **Table 1** Main features of reads obtained for each genome after the read length filtering step.
251 min_len= minimum length, avg_len= average length, max_len= maximum length.

Sample	reads	min_len	avg_len	max_len	N50	Q20(%)	Q30(%)
P78	462982	2000	4960.8	91494	6091	59.53	17.71
P99	418034	2000	7061.4	176512	8845	58.27	19.22

252 N50= the length for which all sequenced reads of that length or greater sum to 50% of the set's

253 total size Q20=1 in 100 probability of an incorrect base call, Q30= 1 in 1000 probability of an
254 incorrect base call. Here 59.53% of P78 bases=Q20 and 58.27% of P99 bases=Q20.
255
256
SV distribution in P78 and P99
For P78, the mapper/caller combination minimap2/Sniffles detected 731 structural variations
(SVs), and the combination NGMLR/Sniffles detected 460 SVs (Table S2). For P99, the
combination minimap2/Sniffles detected 210 SVs and NGMLR/Sniffles detected 397 SVs (Table
S2). Independently from the mapper/caller combination that was used, P78 showed more SVs
than P99 (Table S2). After the cross-validation step (extracting common SVs from each mapper/
caller combination), 236 and 87 SVs were kept for P78 and P99, respectively. For comparison,
the number of small variations (with a size < 30 nt, which were excluded from this study)
amounted to 57 and 77 for P78 and P99, respectively.
In both samples, inversions and deletions were the most prevalent SVs. In P78, inversions
represented 75% of the events and deletions 20% (Figure 2.A). In P99, 80% of the SVs were
inversions and 11% were deletions (Figure 2.A). Inversions were found along the entire
genome, with the highest number detected within the [70-75 kb] window in P78 and within the
[235-245 kb] and [270-275 kb] windows in P99 (Figure 2.B). In P78, deletions were mainly
detected within [10-40 kb], [70-100 kb] and [245-270 kb] windows (Figure 2.B). In P99, the two
major deletions were found within [10-20 kb] and [205-215 kb] regions (Figure 2.B). The
frequencies of these SVs were low and did not exceed 1% of the total reads (with a few
exceptions, Table S2). In spite of such low frequencies, it is interesting to note that the most
frequent SVs were located in ORFs potentially involved in DNA replication and encapsidation,
e.g. ORF33, 46, 47, 55 (Aoki et al, 2007; Table S2).
Altogether, these results highlight high SV turnover dynamics during the *in vitro* infection cycles
(from 78 passages to 99) with a clear trend, or bias, towards inversion and deletion events.

Dynamics and impacts of SVs in ORF150 region
Taking advantage of the high-resolution SV detection provided by the long-read sequencing, we
looked for the SV events around the potential virulence-linked ORF150 in P78 and P99 (nt
257,103-261,345 according to AP008984.1). Results confirmed that P99 had a reference-like
profile with an unmodified ORF150. In P78, the deletion (nt 258,154-259,517; D258153) was
found in 6,902 reads (100% of the reads), whereas the reference haplotype was also detected
in 30 reads, representing 0.44% of the total 6,734 supporting reads (Figure 3, Tables S3).
Surprisingly, 26 reads revealed a haplotype as yet unidentified (INV258153), consisting of an
inversion of the same length (1,363 bp) and at the same breakpoints as the deletion. The

inverted haplotype (INV258153) in P78 deeply altered the ORF150, by inverting the first 1200
bp of the ORF and 160 bp of the 5'UTR in the middle of the ORF (Figure 3).
In order to trace the unexpected dynamics of gain and loss of the full ORF150 along passages,
we searched for the SV turnovers during 10 intermediate passages (P10, P20, P30, P40, P50,
P70, P78, P80, P90, P99). This analysis revealed the presence of haplotype D258153 at low
frequency (from 0.05 to 0.15% of the reads) in passages P10 to P40 and a strong increase in its
frequency at P50 (88.7% of the reads) (Figure 4). The frequency of the haplotype D258153
reached a maximum at P78 (100% of the reads) then dropped quickly at P80 (30.7% of the
reads) to stabilize at low frequency (0.31% of readings) at P90., as during the first 40 passages
(Figure 4, Table S3). Interestingly, shorter deletions of 119 and 881 bp were observed near the
5' end of the ORF150 in P40 and P80, respectively, at low frequencies (0.42 % in P40 and
0.18% in P80) (Figure 4, Table S3). The haplotype D258153 completely disappeared at
passage 99 (Figure 4, Table S3).
This analysis also evidenced several other SVs that alter the structure of ORF150 and of its
upstream region, including the beginning of ORF149 (Figure 5). Besides the large deletion,
inversions and insertions were also observed in the ORF149-ORF150 region. Inversions were
at a low frequency (between 0.01% and 0.53% of the supporting reads) in all passages except
for P70, P90 and P99. P10 and P40 showed the lowest and the highest inversion frequencies,
respectively (Figure 5, Table S3). A large insertion of about 1 kb appeared in P50 and P70 at
moderate frequencies (14,34% and 16.01% of the supporting reads, respectively) to disappear
in P78 and re-appear at a lower frequency (6,79% of the supporting reads) in P80. The
consensus sequence of this insertion corresponds to the fragment 259,517-260,477 of the KHV
genome, with an identity of about 90%. In P90, an intriguing inverted-duplicated haplotype was
observed at a low proportion (0.054% of the supporting reads). Surprisingly, P99 exhibited a
unique reference-like, SV-free haplotype (Figure 5, Table S3). All the variations deeply impacted
the structure of ORF150 - and sometimes that of ORF149 as well - by shrinking or increasing its
size, causing the ORF149 and ORF150 fusion, inverting the ORF150 sequences and
duplicating the ORF150 with the deleted, inserted, inverted and inverted-duplicated haplotypes
(Figure 5).

324
325 **Figure 2.** Ratio and distribution of SVs detected in P78 and P99 genomes. **A)** Frequency of
326 each type of SV detected in P78 and P99. **B)** Occurrences and distribution of each SV type in
327 the P78 and P99 genomes using a 5kb window.

328
329
330

332 **Figure 3.** Impact of structural variations D258153 and INV258153 on the P78 genomic
333 structure. The inversion is highlighted by an inverted arrow compared to the reference
334 haplotype.

Percentage of deleted vs reference haplotypes

Reference haplotype ▢ (cyan)
Deleted haplotype D258153 ▢ (blue)
Shorter haplotype D258153 ▢ (magenta)

341 **Figure 4.** The prevalence of the deleted haplotype during the 10 intermediate passages (P10,
342 P20, P30, P40, P50, P70, P78, P80, P90, P99)
343

	Reference haplotype	CyHV3_ORF149 ◄ ◄ CyHV3_ORF150 ▷ ▷	
P10	Deleted haplotype (D258153)		0.15% *
	Inverted haplotype (INV257913)		0.023% *
P20	Deleted haplotype (D258153)		0.047% *
	Inverted haplotype (INV257941)		0.01% *
P30	Deleted haplotype (D258153)		0.31% *
	Inverted haplotype (INV257918)		0.089% *
	Inverted haplotype (INV258123-260721)		0.076% *
P40	Deleted haplotype (D258154-258272)		0.42% *
	Inverted haplotype (INV258155-260208)		0.41% *
	Inverted haplotype (INV257726-259517)		0.53% *
P50	Deleted haplotype (D258153)		88.97% *
	Inverted haplotype (INV257893-260177)		0.05% *
	Inserted haplotype (INS258155)		14.34% *
P70	Deleted haplotype (D258153)		82.25% *
	Inserted haplotype (INS258155)		16.01% *
P78	Deleted haplotype (D258153)		100% *
	Inverted haplotype (INV258153)		0.44% *
P80	Deleted haplotype (D258153)		30.68% *
	Deleted haplotype (D258155-259035)		0.18% *
	Inverted haplotype (INV257815-259517)		0.04% *
	Inverted haplotype (INV258076-261050)		0.03% *
	Inserted haplotype (INS258155)		6.79% *
P90	Deleted haplotype (D258153)		0.31% *
	InvDup haplotype (INVDUP258386-259486)		0.054% *
P99	Reference-like haplotype		100% *

Figure 5. Impact of SV dynamics on the ORF149-ORF150 structure in the successive passages
P10, P20, P30, P40, P50, P70, P78, P80, P90 and P99. The inversion is highlighted by an
inverted arrow compared to the reference haplotype. Red blocks correspond to an inserted
sequence.
InvDup = Inverted-duplicated haplotype.

*Percentage of reads covering the SVs at given breakpoints.
350
Discussion

SVs significantly impact the adaptation of viruses to their natural host and environment (Pérez-
Losada et al., 2015). Yet the role of SV diversity and dynamics in large DNA viruses is barely
known. Ultra-deep long-read sequencing opens unprecedented ways to gain insights into these
untapped viral genome polymorphisms. The present study started to tackle the impact of SVs in
the evolution of the large dsDNA KHV during cell culture serial passages using ultra-deep
whole-genome and amplicon-based sequencing. The sequence data showed a wide distribution
of various SVs along the genome associated with high SV turnover dynamics during the *in vitro*
infection cycles and a clear bias towards inversion and deletion events. Analysis of the
pathogenicity-associated ORF150 region in ten serial passages mainly highlighted deletions,
inversions and insertions that deeply altered the structure of ORF150.

Serial passages of viruses in cell culture may lead to the accumulation of mutations and gene
disruptions (Spatz 2010; Colgrove et al., 2014). These mutations can modify viral adaptation
and increase or decrease virulence (Boutier et al., 2017; López-Muñoz et al., 2021; Vancsok et
al., 2017). In the case of KHV, a previous work using short-read sequencing showed that 99
consecutive *in vitro* passages onto CCB cells resulted in the accumulation of less than 60 small
variations (<100 nt) (Klafack et al., 2019). It also showed that the haplotype composition can
quickly vary along with infection cycles of KHV *in vitro*. The present study unexpectedly
highlighted a high number of structural variations: 87 for P99 and 236 for P78. In contrast, the
accumulation of small variations was consistent with what had been observed with short-read
sequencing (Klafack et al, 2019). These findings illustrate that long-read sequencing is highly
suitable for genome-wide comparisons of viruses. Most importantly, they revealed a hidden
source of virus diversification, which had never been reported so far for KHV. They also
confirmed that P78 consists of a mixture of undeleted and deleted haplotypes revealing that the
undeleted haplotype did not correspond to the native one but to an inverted version of it. We
experimentally validated this inversion in P78 by PCR followed by sequencing. Moreover, the
inversion found in the middle of the reads excluded the formation of *in silico* chimeras, i.e.,
chimeras resulting from the basecaller when two molecules are sequenced in the same pore
that undergoes fast reloading (Martin and Legget, 2021). These structural variations form a
'mosaic' of viral subpopulations that seem to result from multiple rearrangement events, mainly
inversions and deletions and, to a lesser extent, insertions and duplications. Such sub-viral
variants often lead to Defective Viral Genomes (DVGs) (Vignuzzi and López 2019). Because of

their negative impact on viral replication, some forms of DVGs have been extensively studied,

and three pathogenesis-related functions have been well-described: interference with viral

replication, immunostimulation, and viral persistence (Marriott and Dimmock 2010; Vignuzzi and

López 2019). DVGs play a role in viral production interference by accumulating at higher rates

than the full-length viral genomes and consequently interfere with viral replication by taking up

the polymerase activity and competing for structural proteins (Calain and Roux 1995; Portner

and Kingsbury 1971). In addition, DVGs can act as primary stimuli and trigger antiviral immunity

by inducing the expression of some interleukins and pro-inflammatory cytokines (for review, see

Vignuzzi and López 2019; Xiao et al, 2021). However, the implication of DVGs for virus

persistence is complex and still unclear. Nevertheless, a combination of viral replication

interfering and cycle asynchrony between full-length viruses and DVGs seems to establish

chronical infection (for review, see Vignuzzi and López 2019).

The observed dynamics of structural variations in the KHV genomes during cell culture

passages where selection pressures are virtually non-existent may indicate an accumulation of

DVGs impacting the pathogenicity. Indeed, Klafack et al. (2019) showed that serial passages

significantly attenuated this infectious virus. This attenuation was associated with a truncated

KHV genome bearing a 1.3-kb deletion that removes the majority of the ORF150. A very recent

study showed that partial or complete depletion of ORF150 leads to a clear attenuation of the

virus. It also brought *in vivo* evidence that ORF150 may down-regulate the inflammatory

response of carp to enhance viral proliferation, thus confirming a key role of ORF150 product in

KHV virulence (Klafack et al., 2022 submitted). ORF150 encodes a protein with a Really

Interesting New Gene (RING) domain that is predicted to span 628 amino acids (Li et al, 2015;

Aoki et al, 2007). This RING domain contains the HC (C3HC4) type of RING structure, which is

involved in the ubiquitination pathway, by acting as a E3 ubiquitin-protein ligase (He and Kwang

2008). Viruses that encode RING finger like-ubiquitin ligases (E3s)may evade host immune

responses and also hijack the host's RING E3 to enhance their replication (Zhang et al.,2018).

In aquatic viruses, RING family genes have been reported to be involved in virus latency,

replication, and host protein degradation (Shekar and Venugopal 2019; Wang et al., 2021).

As previously observed, the most significant difference affecting all viral haplotypes between

P78 and P99 was around the ORF150. For this reason, we focused on this interesting region to

determine the passage number at which the presence of the deletion inversion first arose, by

sequencing selected representative passages (P10, P20, P30, P40, P50, P70, P80 and P90).

We found that deleted and inverted haplotypes appeared as soon as P10 and that their

proportion varied along the successive passages. The most striking feature was the rapid and

total disappearance of the deletion between P78 and P99, raising many questions regarding the mechanisms that led to the clearance of this major haplotype.

The rapid SV turnover of DNA viruses, including herpesviruses, likely involves recombination (Szpara and VanDoorslaer, 2021; Renner and Szpara, 2018; Cudini et al., 2019; Wilkinson and Weller, 2003; Kolb et al., 2017; Tomer et al., 2019), which is often linked to replication and DNA repair, as well as errors during viral genome replication (Kulkarni and Fortunato 2011; Xiaofei and Kowalik 2014). In the present case, the pretty good conservation of breakpoints around ORF150 may be a sign of homologous recombination. However, whether this recombination occurs within the same genomic entities or between different viruses remains open. It would be interesting to assess the involvement of each of these mechanisms in generating the observed structural diversity. The multiple mechanisms of DNA virus evolution beyond single nucleotide substitutions likely confer KHV a high level of evolutionary adaptability.

Classically, the generation of live-attenuated vaccines is achieved by passaging the virus in cell culture under different conditions (in different host species or at lower or higher temperatures), in order to induce mutation accumulation that supports viral adaptation to the specific conditions and provides viral attenuation (Minor, 2015, Hanley 2011). With the exception of the OPV polio vaccine viruses (Kew et al., 2005), the exact mechanisms by which these mutations lead to attenuated phenotypes are usually poorly characterized (Lauring et al., 2010). However, live-attenuated viruses can revert to virulent phenotypes either by reversions (as shown here between P78 and P99), introduction of compensatory mutations, or recombination with viruses belonging to the same genus (Cann et al., 1984; Bull et al., 2018; Muslin et al., 2019). Additionally, the combination of multiple live-attenuated viruses may result in competition or facilitation between individual vaccine viruses, resulting in undesirable increases in virulence or decreases in immunogenicity (Hanley 2011; Pereira-Gomez et al., 2021). Recently, genetic engineering has led to many novel approaches to generate live-attenuated virus vaccines that contain modifications to prevent reversion to virulence (Yeh et al., 2020) and improve interferences among multiple vaccine strains (Pereira-Gomez et al., 2021).

Conclusion

Our findings confirm that CyHV-3 can evolve rapidly during infectious cycles in cell culture, and SVs are a major component in the evolutionary process of this virus. SVs are extremely dynamic under *in vitro* controlled conditions, and it would now be interesting to evaluate their dynamics *in vivo*. The present study also contributes to the basic research on the mechanisms underlying attenuation, and may have important outcomes for the design of safe live-attenuated vaccine formulations.

Acknowledgements

Part of this work was supported by the "Laboratoire d'excellence" (LabEx) CeMEB, an ANR

"Investissements d'avenir" program (ANR-10-LABX-04-01), through the exploratory research

project HaploFit. This work benefited from the Montpellier Bioinformatics Biodiversity platform

supported by the LabEx CeMEB, an ANR "Investissements d'avenir" program (ANR-10-LABX-

04-01). Real-time PCR data were produced on the qPHD platform of Montpellier University, with

the support of LabEx CeMEB. N. N. F benefited from a CampusFrance fellowship for her PhD

thesis, and we are very grateful to the French Embassy of Indonesia. E. C received a post-

doctoral fellowship from IRD.

Funding

HaploFit project, an ANR "Investissements d'avenir" program (ANR-10-LABX-04-01).

CampusFrance fellowship.

IRD/Ecobio post-doctoral fellowship.

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Authors' contributions

Nurul Novelia Fuandila: Investigation; Methodology; Formal analysis; Writing – original draft;

Data Validation

Anne-Sophie Gosselin-Grenet: Investigation; Methodology; Data curation; Data Validation;
Writing – original draft
471
472 **Marie-Ka Tilak:** Investigation; Methodology ; Data Validation
473
474 **Sven M Bergmann**: Resources; Writing – review and editing
475
476 **Sandro Klafack:** Resources; Writing – review and editing
477
478 **Angela Lusiastuti:** Methodolgy; Writing – review and editing
479
Munti Yuhanna: Methodolgy; Writing – review and editing

Jean-Michel Escoubas: Methodology; Funding acquisition; Writing – review and editing

Anna-Sophie Fiston-Lavier: Writing – review and editing

Jean-Christophe Avarre: Conceptualization; Data curation; Methodology; Funding acquisition;
Data Validation; Writing – original draft

Emira Cherif: Conceptualization; Data curation; Formal analysis; Software; Writing – original
draft;

487
488 **Data availability**

References

Akhtar, L. N., Bowen, C. D., Renner, D. W., Pandey, U., Della Fera, A. N., Kimberlin, D. W., ... &
Szpara, M. L. (2019). Genotypic and phenotypic diversity of herpes simplex virus 2 within the
infected neonatal population. MSphere, 4(1), e00590-18.
https://doi.org/10.1128/mSphere.00590-18

Aoki, T., Hirono, I., Kurokawa, K., Fukuda, H., Nahary, R., Eldar, A., ... & Hedrick, R. P. (2007).
Genome sequences of three koi herpesvirus isolates representing the expanding distribution of
an emerging disease threatening koi and common carp worldwide. Journal of virology, 81(10),
5058-5065. DOI: 10.1128/JVI.00146-07

Boutier, M., Ronsmans, M., Rakus, K., Jazowiecka-Rakus, J., Vancsok, C., Morvan, L., ... &
Vanderplasschen, A. (2015). Cyprinid herpesvirus 3: an archetype of fish alloherpesviruses.
Advances in Virus Research, 93, 161-256. DOI: 10.1016/bs.aivir.2015.03.001

Boutier, Maxime; Gao, Yuan; Vancsok, Catherine; Suárez, Nicolás M.; Davison, Andrew J.;
Vanderplasschen, Alain (2017): Identification of an essential virulence gene of cyprinid
herpesvirus 3. In: *Antiviral research* 145, S. 60–69. DOI: 10.1016/j.antiviral.2017.07.002.

Bull, J. C., Godfray, H. C. J., & O'Reilly, D. R. (2003). A few-polyhedra mutant and wild-type
nucleopolyhedrovirus remain as a stable polymorphism during serial coinfection in Trichoplusia

ni. Applied and environmental microbiology, 69(4), 2052-2057.
https://doi.org/10.1128/AEM.69.4.2052-2057.2003

Bull, J. J., Smithson, M. W., & Nuismer, S. L. (2018). Transmissible viral vaccines. Trends in
microbiology, 26(1), 6-15. https://doi.org/10.1016/j.tim.2017.09.007

Calain, P., & Roux, L. Functional characterisation of the genomic and antigenomic promoters of
Sendai virus. Virology 212, 163–173 (1995). DOI: 10.1006/viro.1995.1464

Cann, A. J., Stanway, G., Hughes, P. J., Minor, P. D., Evans, D. M. A., Schild, G. T., & Almond,
J. (1984). Reversion to neurovirulence of the live-attenuated Sabin type 3 oral pollovirus
vaccine. Nucleic acids research, 12(20), 7787-7792. https://doi.org/10.1093/nar/12.20.7787

.

Chateigner, A., Bézier, A., Labrousse, C., Jiolle, D., Barbe, V., & Herniou, E. A. (2015). Ultra
deep sequencing of a baculovirus population reveals widespread genomic variations. Viruses,
7(7), 3625-3646. https://doi.org/10.3390/v7072788

Colgrove, R., Diaz, F., Newman, R., Saif, S., Shea, T., Young, S., ... & Knipe, D. M. (2014).
Genomic sequences of a low passage herpes simplex virus 2 clinical isolate and its plaque-
purified derivative strain. Virology, 450, 140-145. DOI: 10.1016/j.virol.2013.12.014.

Cudini, J., Roy, S., Houldcroft, C. J., Bryant, J. M., Depledge, D. P., Tutill, H., ... & Breuer, J.
(2019). Human cytomegalovirus haplotype reconstruction reveals high diversity due to
superinfection and evidence of within-host recombination. Proceedings of the National Academy
of Sciences, 116(12), 5693-5698. https://doi.org/10.1073/pnas.1818130116

Fuchs, Walter; Granzow, Harald; Dauber, Malte; Fichtner, Dieter; Mettenleiter, Thomas C.
(2014): Identification of structural proteins of koi herpesvirus. In: *Archives of virology* 159 (12),
S. 3257–3268. DOI: 10.1007/s00705-014-2190-4.

Gao, Y., Suárez, N. M., Wilkie, G. S., Dong, C., Bergmann, S., Lee, P. Y. A., ... & Boutier, M.
(2018). Genomic and biologic comparisons of cyprinid herpesvirus 3 strains. Veterinary
research, 49(1), 1-11. DOI: 10.1186/s13567-018-0532-z

Gotesman, M., Kattlun, J., Bergmann, S. M., & El-Matbouli, M. (2013). CyHV-3: the third
cyprinid herpesvirus. Diseases of aquatic organisms, 105(2), 163-174.
https://doi.org/10.3354/dao02614

Haenen, O. L. M., Way, K., Bergmann, S. M., & Ariel, E. (2004). The emergence of koi
herpesvirus and its significance to European aquaculture. Bulletin of the European Association
of Fish Pathologists, 24(6), 293-307.

Hammoumi, Saliha; Vallaeys, Tatiana; Santika, Ayi; Leleux, Philippe; Borzym, Ewa; Klopp,
Christophe; Avarre, Jean-Christophe (2016): Targeted genomic enrichment and sequencing of

CyHV-3 from carp tissues confirms low nucleotide diversity and mixed genotype infections. In:
PeerJ, e2516. DOI: 10.7717/peerj.2516.

Hanley, K. A. (2011). The double-edged sword: How evolution can make or break a live-
attenuated virus vaccine. Evolution: Education and Outreach, 4(4), 635-643. DOI:
10.1007/s12052-011-0365-y

He, F., & Kwang, J. (2008). Identification and characterization of a new E3 ubiquitin ligase in
white spot syndrome virus involved in virus latency. Virology journal 5, 151.
https://doi.org/10.1186/1743-422X-5-151

Hedrick, R. P., Gilad, O., Yun, S., Spangenberg, J. V., Marty, G. D., Nordhausen, R. W., ... &
Eldar, A. (2000). A herpesvirus associated with mass mortality of juvenile and adult koi, a strain
of common carp. Journal of Aquatic Animal Health, 12(1), 44-57. DOI: 10.1007/s12052-011-
0365-y

Jeffares, D. C., Jolly, C., Hoti, M., Speed, D., Shaw, L., Rallis, C., ... & Sedlazeck, F. J. (2017).
Transient structural variations have strong effects on quantitative traits and reproductive
isolation in fission yeast. Nature communications, 8(1), 1-11. DOI: 10.1038/NCOMMS14061

Kew, O. M., Sutter, R. W., de Gourville, E. M., Dowdle, W. R., & Pallansch, M. A. (2005). Vac-
cine-derived polioviruses and the endgame strategy for global polio eradication. Annu. Rev. Mi-
crobiol., 59, 587-635. DOI: 10.1146/annurev.micro.58.030603.123625

Klafack, S., Fiston-Lavier, A. S., Bergmann, S. M., Hammoumi, S., Schröder, L., Fuchs, W., ... &
Avarre, J. C. (2019). Cyprinid herpesvirus 3 evolves in vitro through an assemblage of
haplotypes that alternatively become dominant or under-represented. Viruses, 11(8), 754.
https://doi.org/10.3390/v11080754

Klafack, S., Schröder, L., Jin, Y., Lenk, M., Lee, P.Y., Fuchs, W, Avarre, J. C., Bergmann, S. M.
(2022). Development of an attenuated vaccine against Koi Herpesvirus Disease (KHVD)
suitable for oral administration and immersion. (submitted, NPJ Vaccines)

Klafack, S., Wang, Q., Zeng, W., Wang, Y., Li, Y., Zheng, S.C., Kempter, J., Lee, P.Y., &
Bergmann, S. M. (2017). Genetic Variability of Koi Herpesvirus In vitro—A Natural Event?.
Front. Microbiol., 8, 982. https://doi.org/10.3389/fmicb.2017.00982

Kolb, A. W., Lewin, A. C., Moeller Trane, R., McLellan, G. J., & Brandt, C. R. (2017).
Phylogenetic and recombination analysis of the herpesvirus genus varicellovirus. BMC
genomics, 18(1), 1-17. https://doi.org/10.1186/s12864-017-4283-4

Kulkarni, A. S., & Fortunato, E. A. (2011). Stimulation of homology-directed repair at I-SceI-
induced DNA breaks during the permissive life cycle of human cytomegalovirus. Journal of
virology, 85(12), 6049-6054. https://doi.org/10.1128/JVI.02514-10

Lauring, A. S., & Andino, R. (2010). Quasispecies theory and the behavior of RNA viruses.
PLoS pathogens, 6(7), e1001005. https://doi.org/10.1371/journal.ppat.1001005

Lauring, A. S., Jones, J. O., & Andino, R. (2010). Rationalizing the development of live
attenuated virus vaccines. Nature biotechnology, 28(6), 573-579.
\https://doi.org/10.1038/nbt.1635

Lecompte, L., Peterlongo, P., Lavenier, D., & Lemaitre, C. (2020). SVJedi: Genotyping structural
variations with long reads. Bioinformatics, 36(17), 4568-4575.
https://doi.org/10.1093/bioinformatics/btaa527

Li, D., Lott, W. B., Lowry, K., Jones, A., Thu, H. M., & Aaskov, J. (2011). Defective interfering
viral particles in acute dengue infections. PloS one, 6(4), e19447.
https://doi.org/10.1371/journal.pone.0019447

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics, 34(18),
3094-3100. DOI: 10.1093/bioinformatics/bty191

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & Durbin, R. (2009). The
sequence alignment/map format and SAMtools. Bioinformatics, 25(16), 2078-2079.
https://doi.org/10.1093/bioinformatics/btp352

Li, W., Lee, X., Weng, S., He, J., Dong, C. (2015). Whole-genome sequence of a novel Chinese
cyprinid herpesvirus 3 isolate reveals the existence of a distinct European genotype in East
Asia. Veterinary microbiology 175 (2-4), 185-194.
https://doi.org/10.1016/j.vetmic.2014.11.022

Loiseau, V., Herniou, E. A., Moreau, Y., Lévêque, N., Meignin, C., Daeffler, L., ... & Gilbert, C.
(2020). Wide spectrum and high frequency of genomic structural variation, including
transposable elements, in large double-stranded DNA viruses. Virus evolution, 6(1), vez060.
https://doi.org/10.1093/ve/vez060

López-Muñoz, A. D., Rastrojo, A., Martín, R., & Alcamí, A. (2021). Herpes simplex virus 2 (HSV-
2) evolves faster in cell culture than HSV-1 by generating greater genetic diversity. PLoS
pathogens, 17(8), e1009541. DOI: 10.1371/journal.ppat.1009541

Marriott, A. C., and Dimmock, N. J. (2010) 'Defective Interfering Viruses and Their Potential as
Antiviral Agents', Reviews in Medical Virology, 20: 51–62. DOI: 10.1002/rmv.641

Martin, S., & Leggett, R. M. (2021). Alvis: a tool for contig and read ALignment VISualisation
and chimera detection. BMC bioinformatics, 22(1), 1-10. https://doi.org/10.1186/s12859-021-
04056-0

Minor, P. D. (2015). Live attenuated vaccines: Historical successes and current challenges.
Virology, 479, 379-392. DOI: 10.1016/j.virol.2015.03.032

Molenkamp, R., Rozier, B. C., Greve, S., Spaan, W. J., & Snijder, E. J. (2000). Isolation and
characterization of an arterivirus defective interfering RNA genome. Journal of virology, 74(7),
3156-3165. https://doi.org/10.1128/JVI.74.7.3156-3165.2000

Muslin, C., Mac Kain, A., Bessaud, M., Blondel, B., & Delpeyroux, F. (2019). Recombination in enteroviruses, a multi-step modular evolutionary process. Viruses, 11(9), 859. DOI: 10.3390/v11090859

O'Hara, P. J., Nichol, S. T., Horodyski, F. M., & Holland, J. J. (1984). Vesicular stomatitis virus defective interfering particles can contain extensive genomic sequence rearrangements and base substitutions. Cell, 36(4), 915-924. https://doi.org/10.1016/0092-8674(84)90041-2

Pereira-Gómez, M., Carrau, L., Fajardo, Á., Moreno, P., & Moratorio, G. (2021). Altering Compositional Properties of Viral Genomes to Design Live-Attenuated Vaccines. Frontiers in Microbiology, 1706. https://doi.org/10.3389/fmicb.2021.676582

Pérez-Losada, M., Arenas, M., Galán, J. C., Palero, F., & González-Candelas, F. (2015). Recombination in viruses: mechanisms, methods of study, and evolutionary consequences. Infection, Genetics and Evolution, 30, 296-307. DOI:10.1016/j.meegid.2014.12.022

Portner, A., & Kingsbury, D. W. Homologous interference by incomplete Sendai virus particles: changes in virus-specifc ribonucleic acid synthesis. J. Virol. 8, 388–394 (1971). DOI: 10.1128/jvi.8.4.388-394.1971

Rakus, K., Ouyang, P., Boutier, M., Ronsmans, M., Reschner, A., Vancsok, C., ... & Vanderplasschen, A. (2013). Cyprinid herpesvirus 3: an interesting virus for applied and fundamental research. Veterinary Research, 44(1), 1-16. https://doi.org/10.1186/1297-9716-44-85

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., ... & Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic acids research, 44(W1), W160-W165. doi:10.1093/nar/gkw257.

Renner, D. W., & Szpara, M. L. (2018). Impacts of genome-wide analyses on our understanding of human herpesvirus diversity and evolution. Journal of virology, 92(1), e00908-17. DOI: 10.1128/JVI.00908-17

Renzette, N., Pokalyuk, C., Gibson, L., Bhattacharjee, B., Schleiss, M. R., Hamprecht, K., ... & Kowalik, T. F. (2015). Limits and patterns of cytomegalovirus genomic diversity in humans. Proceedings of the National Academy of Sciences, 112(30), E4120-E4128. https://doi.org/10.1073/pnas.1501880112

Sanjuán, R., & Domingo-Calap, P. (2016). Mechanisms of viral mutation. Cellular and molecular life sciences, 73(23), 4433-4448. https://doi.org/10.1007/s00018-016-2299-6

Sedlazeck, F. J., Rescheneder, P., Smolka, M., Fang, H., Nattestad, M., Von Haeseler, A., & Schatz, M. C. (2018). Accurate detection of complex structural variations using single-molecule sequencing. Nature methods, 15(6), 461-468. doi: 10.1038/s41592-018-0001-7

Shekar, M., & Venugopal, M. N. (2019). Identification and characterization of novel double zinc
fingers encoded by putative proteins in genome of white spot syndrome virus. Archives of
virology, 164(4), 961-969. DOI: 10.1007/s00705-019-04150-y

Spatz, S. J. (2010). Accumulation of attenuating mutations in varying proportions within a high
passage very virulent plus strain of Gallid herpesvirus type 2. Virus research, 149(2), 135-142.
DOI:10.1016/j.virusres.2020.01.007

Sunarto, A., McColl, K. A., Crane, M. S. J., Sumiati, T., Hyatt, A. D., Barnes, A. C., & Walker, P.
J. (2011). Isolation and characterization of koi herpesvirus (KHV) from Indonesia: identification
of a new genetic lineage. Journal of fish diseases, 34(2), 87-101. https://doi.org/10.1111/j.1365-
2761.2010.01216.x

Szpara, M.L., & Van Doorslaer, K. Mechanisms of dna virus evolution. In Dennis H. Bamford
and Mark Zuckerman, editors, Encyclopedia of Virology (Fourth Edition),pages 71–78.
Academic Press, Oxford, fourth edition edition, 2021. 3 https://doi.org/10.1016/B978-0-12-
809633-8.20993-X

Te Yeh, M., Bujaki, E., Dolan, P. T., Smith, M., Wahid, R., Konz, J., ... & Andino, R. (2020).
Engineering the live-attenuated polio vaccine to prevent reversion to virulence. Cell host &
microbe, 27(5), 736-751. https://doi.org/10.1016/j.chom.2020.04.003

Tomer, E., Cohen, E. M., Drayman, N., Afriat, A., Weitzman, M. D., Zaritsky, A., & Kobiler, O.
(2019). Coalescing replication compartments provide the opportunity for recombination between
coinfecting herpesviruses. The FASEB journal, 33(8), 9388-9403.
https://doi.org/10.1096/fj.201900032R

Vancsok, C., Peñaranda, M. M. D., Raj, V. S., Leroy, B., Jazowiecka-Rakus, J., Boutier, M., ... &
Vanderplasschen, A. F. (2017). Proteomic and functional analyses of the virion transmembrane
proteome of cyprinid herpesvirus 3. Journal of Virology, 91(21), e01209-17. DOI:
10.1128/JVI.01209-17

Vignuzzi, M., & López, C. B. (2019). Defective viral genomes are key drivers of the virus–host
interaction. Nature microbiology, 4(7), 1075-1087. https://doi.org/10.1038/s41564-019-0465-y

Wang, Z. H., Ke, F., Zhang, Q. Y., & Gui, J. F. (2021). Structural and functional diversity among
five ring finger proteins from Carassius auratus herpesvirus (CaHV). Viruses, 13(2), 254. DOI:
10.3390/v13020254

White, R., Pellefigues, C., Ronchese, F., Lamiable, O., & Eccles, D. (2017). Investigation of
chimeric reads using the MinION. F1000Research, 6.
https://doi.org/10.12688/f1000research.11547.2

Wilkinson, D., & Weller, S. (2003). The role of DNA recombination in herpes simplex virus DNA
replication. IUBMB life, 55(8), 451-458. https://doi.org/10.1080/15216540310001612237

Wilkinson, Dianna E., Weller, Sandra K. (2003): The role of DNA recombination in herpes
simplex virus DNA replication. In: *IUBMB life* 55 (8), S. 451–458. DOI:
10.1080/15216540310001612237

Xiao, Y., Lidsky, P. V., Shirogane, Y., Aviner, R., Wu, C. T., Li, W., ... & Andino, R. (2021). A
defective viral genome strategy elicits broad protective immunity against respiratory viruses.
Cell, 184(25), 6037-6051. DOI: 10.1016/j.cell.2021.11.02

Xiaofei, E., & Kowalik, T.F. The dna damage response induced by infection with human
cytomegalovirus and other viruses. Viruses, 6(5) :2155–2185, 2014.
https://doi.org/10.3390/v6052155

Zhang, Y., Li, L.F., Munir, M. Qiu, H. J. (2018). RING-Domain E3 Ligase-Mediated Host-Virus
Interactions: Orchestrating Immune Responses by the Host and Antagonizing Immune Defense
by Viruses. Frontiers in immunology 9, 1083. https://doi.org/10.3389/fimmu.2018.01083.

LTR

RTR

[0 - 1,78]

[0 - 1,71]