Reference genome choice and filtering thresholds jointly influence phylogenomic analyses

Jessica A. Rick[1,2*], Chad D. Brock[2,3], Alexander L. Lewanski[2], Jimena Golcher-Benavides[4], and Catherine E. Wagner,[2]

[1] *Program in Ecology, University of Wyoming, Laramie, WY, 82071, USA*
[2] *Department of Botany, University of Wyoming, Laramie, WY, 82071, USA*
[3] *Current: Department of Biological Sciences, Tarleton State University, Stephenville, TX 76401, USA*
[4] *Lees-McRae College, Banner Elk, NC 28604, USA*

Corresponding author: JA Rick, 1000 E University Dr, Laramie, WY 82070; jrick@uwyo.edu

Abstract

Molecular phylogenies are a cornerstone of modern comparative biology and are commonly
employed to investigate a range of biological phenomena, such as diversification rates,
patterns in trait evolution, biogeography, and community assembly. Recent work has
demonstrated that significant biases may be introduced into downstream phylogenetic
analyses from processing genomic data; however, it remains unclear whether there are
interactions among bioinformatic parameters or biases introduced through the choice of
reference genome for sequence alignment and variant-calling. We address these knowledge
gaps by employing a combination of simulated and empirical data sets to investigate to
what extent the choice of reference genome in upstream bioinformatic processing of
genomic data influences phylogenetic inference, as well as the way that reference genome
choice interacts with bioinformatic filtering choices. We demonstrate that more stringent
minor allele filters bias inferred trees away from the true species tree topology, and that
these biased trees tend to be more imbalanced and have a higher center of gravity than the
true trees. We find greatest topological accuracy when filtering sites for minor allele count
>3–4 in our 51-taxa data sets, while tree center of gravity was closest to the true value

₁₆ when filtering for sites with minor allele count >1–2. In contrast, filtering for missing data

₁₇ increased accuracy in the inferred topologies; however, this effect was small in comparison

₁₈ to the effect of minor allele filters and may be undesirable due to a subsequent mutation

₁₉ spectrum distortion. The bias introduced by these filters differs based on the reference

₂₀ genome used in short read alignment, indicating that choosing a reference genome for

₂₁ alignment is an important bioinformatic decision with implications for downstream

₂₂ analyses. These results demonstrate that care needs to be taken when choosing parameters

₂₃ for assembling and filtering large genomic data sets from short read data.

₂₄ *Key words*: phylogenomics, bioinformatics, diversification rate, minor allele frequency,

₂₅ imbalance, macroevolution

Phylogenetic history provides a critical framework for understanding the myriad components of a lineage's evolution, and this "tree thinking" approach is a cornerstone of modern comparative biology (O'Hara, 1997; Pagel, 1999). Large multilocus data sets have become increasingly important for inferring phylogenetic histories (e.g., Edwards, 2009; Lemmon and Lemmon, 2013; Chan et al., 2020; Cloutier et al., 2019; Huang et al., 2020), and our ability to resolve relationships among rapidly diversifying taxa has greatly improved as a result of advances in high-throughput sequencing methods (e.g., Wagner et al., 2013; Rokas et al., 2003; Philippe et al., 2004; Prasad et al., 2008). However, inferring phylogenetic relationships is not a trivial task, even with genomic data. These large data sets also present novel—and in some cases, unexplored—challenges to phylogenetic analyses (e.g., Kumar et al., 2012; Jeffroy et al., 2006).

Phylogenetic inferences may be sensitive to both sampling and analytical approaches, and previous work has demonstrated that factors such as taxon sampling (Rannala et al., 1998; Zwickl and Hillis, 2002; Cusimano and Renner, 2010; Brock et al., 2011; Heath et al., 2008a,b), missing data (Wiens, 1998, 2003; Pybus and Harvey, 2000; Wiens and Moen, 2008; Wiens and Tiu, 2012; Grievink et al., 2013; Lemmon et al., 2009), and method of phylogenetic reconstruction (Huelsenbeck and Kirkpatrick, 1996; Revell et al., 2005; Rüber and Zardoya, 2005; Grievink et al., 2013; Leaché et al., 2015) can affect phylogenetic accuracy and downstream inferences about the tempo and mode of evolution. While bioinformatic filters—such as those for read depth, missing data per site, missing data per individual, and minor allele frequency—are often employed to guard against sequencing and alignment errors, these choices can inadvertently bias the characteristics of sites that are retained for downstream analyses. These filters tend to reduce the number of loci in the data set and can bias the characteristics of those loci that are retained. In population genetics, it is known that rare variants (i.e., those at $<5\%$ frequency in a population) are common and often of interest (Biddanda et al., 2020), particularly in disease association studies (Nelson et al., 2012; Momozawa and Mizukami, 2021) and

analyses of fine-scale population structure (Slatkin, 1985; Novembre et al., 2008). The inclusion or removal of these low-frequency sites can affect estimates of gene flow (Slatkin, 1985) or differentiation among populations (Bhatia et al., 2013). Several recent studies have also demonstrated that filtering out sites with low minor allele frequencies will truncate the inferred allele frequency spectrum by removing the most recent mutations (Linck and Battey, 2019; Shafer et al., 2017), which can lead to biased estimates of historical demography and evolutionary processes, particularly in the recent past (Boitard et al., 2016) when using methods based on site frequency spectra.

Similarly, missing data in reduced-representation sequencing are expected to be non-randomly distributed across species and loci, with the amount of missing data proportional to the genetic distance between taxa (Eaton et al., 2017) and to locus-specific mutation rates. Filtering loci by amount of missing data thus disproportionately excludes loci with high mutation rates, which artificially truncates the mutational spectrum of the remaining loci (Huang and Knowles, 2016). This has important implications for analyses explicitly using the mutational spectrum for modeling evolutionary history (e.g., inferences of historical demography, Gutenkunst et al., 2009; Hotaling et al., 2018; Boitard et al., 2016; McTavish and Hillis, 2015; DeWitt et al., 2021), as well as for phylogenetic analyses where shared mutations provide information about evolutionary relationships and branch lengths. Loci with high mutation rates tend to be more phylogenetically informative, and therefore contribute more to accurate tree estimation (Lanier et al., 2014). Consequently, removing these loci may then lead to severe drops in tree estimation accuracy. Consistent with this, Molloy and Warnow (2018) demonstrated that filtering genes based on missing data generally reduces the accuracy of both species tree and concatenation-based phylogenetic analyses, with a greater effect in trees with more incomplete lineage sorting.

Recent advances in sequencing technologies along with falling sequencing costs have made it increasingly feasible to sequence and assemble reference genomes for non-model species (Feng et al., 2020; Kolora et al., 2021; Hotaling et al., 2021; Formenti et al., 2022;

Rhie et al., 2021). Non-model reference genomes can then be used to assemble short reads from reduced-representation or whole genome sequencing into shared loci across individuals, and an increasing number of studies use a reference genome of a focal study taxon for assembly. Assembly of reads to a reference genome is based on sequence similarity (Catchen et al., 2011), and reads with higher mutation rates and higher variability will tend to have lower alignment scores (Nielsen et al., 2011; Huang and Knowles, 2016). Therefore, the choice of an ingroup (i.e., closely-related) versus an outgroup (i.e., more distant) reference genome with which to align reads may have substantial consequences for downstream phylogenetic analyses. For example, there are likely to be more reads with lower alignment scores when aligning to a more distant genome than when using an ingroup reference, and conserved regions with low mutation rates are more likely to be kept post-filtering (McCormack et al., 2009; Huang and Knowles, 2016). A consequence of this is that DNA fragments carrying the reference allele will be more likely to map successfully and pass quality filters, and therefore be retained in analyses (Ros-Freixedes et al., 2018; Brandt et al., 2015). This bias is expected to increase with increasing genetic distance to the reference genome and to be more apparent in regions that are highly polymorphic. Following this prediction, several recent studies have demonstrated that reference genome choices can bias population genetic inferences, such as in estimates of historical demography, heterozygosity, and runs of homozygosity (Prasad et al., 2021; Günther and Nettelblad, 2019; Reid et al., 2021; Shafer et al., 2017). In addition, Bertels et al. (2014) have demonstrated that reference genome choice can bias both topological accuracy and branch length estimation in phylogenetic analyses in bacteria as a result of divergent regions failing to map when query sequences are "sufficiently divergent" from the reference sequence. With this knowledge, a more general treatment of biases introduced by reference genome choices in phylogenetics is warranted.

Such systematic biases in sites retained for analyses may hinder our ability to recover the true evolutionary history of a clade, and in doing so, may bias inferences made

about the tempo and mode of evolution. Small changes in the sites retained can have large influences on the inferred topology and branch lengths (Shen et al., 2017) and therefore the diversification patterns inferred from that topology. For example, more stringent minor allele frequency filters may remove a greater proportion of data from branches at the tips of a phylogeny, leading to the truncation of these branches and an erroneous signal of increasing diversification in the recent past. Employing a more distant reference genome may have a similar effect, as this could lead to lower alignment scores for the more rapidly evolving sites most likely to change in recently diverged groups. Alternatively, as missing data is likely to increase with increasing genetic distance between taxa (e.g., due to loss of restriction cut sites; Huang and Knowles, 2016; Eaton et al., 2017), missing data filters may disproportionately affect branches toward the base of the tree. Stringent filtering may reduce data set size, as well as the distribution of signal across a phylogeny. As phylogenetically informative sites are lost, increasing noise may lead to more imbalanced topologies, on average, because there are more ways for a tree to be unbalanced than balanced (Heard and Mooers, 1996; Huelsenbeck and Kirkpatrick, 1996).

Unfortunately, there is a lack of consensus on how best to process data for phylogenetic analysis. For example, some studies recommend filtering data stringently to ensure sequence accuracy (Davey et al., 2013). Others demonstrate that filtering can bias important features of the resulting data (Nielsen, 2000; Mastretta-Yanes et al., 2015; Huang and Knowles, 2016; Chan et al., 2020), and therefore recommend against filtering stringently. It may also be the case that there is not a universally "best" approach to these decisions (Nazareno and Knowles, 2021); variation in empirical data attributes such as divergence times, incomplete lineage sorting, and genetic diversity may mean that different filtering decisions are optimal in different scenarios. For example, Huang and Knowles (2016) show that high missing data filters are particularly harmful for data sets with short coalescent times and it is likely that the consequences of other filters are similarly context dependent. Consequently, there may not be a one-size-fits-all prescription that can be

₁₃₄ made for the treatment of data for phylogenetics and the proper filtering choices may be

₁₃₅ contingent on the focus of the analysis.

₁₃₆ Here, we assess the influence of bioinformatic choices on phylogenetic inference by

₁₃₇ combining simulations and empirical data sets to assess the interacting influences of

₁₃₈ reference genome choice, minor allele filters, and missing data filters on inferred

₁₃₉ phylogenies. We assess our ability to recover both the true tree and the true summary

₁₄₀ statistics describing tree shape and imbalance, and address three main questions: (1) To

₁₄₁ what extent do bioinformatic choices affect our ability to recover the 'true' phylogenetic

₁₄₂ relationships? (2) How do bioinformatic choices bias inferences of diversification patterns

₁₄₃ based on tree shape and node distribution? and (3) How do these biases depend on the

₁₄₄ diversification history within the clade? Our results demonstrate that decisions made by

₁₄₅ researchers during the assembly and processing of genomic data, as well as characteristics

₁₄₆ of the true divergence history among taxa, can bias the data set retained and the inferred

₁₄₇ phylogenetic tree in ways that are consequential for macroevolutionary analyses. Finally,

₁₄₈ while our inferences focus on single nucleotide polymorphism (SNP) data sets derived from

₁₄₉ reduced-representation data, we outline how these findings are also applicable to the whole

₁₅₀ genome data sets that are becoming more common in modern phylogenomics.

₁₅₁ <center>MATERIALS AND METHODS</center>

₁₅₂ <center>*Overview*</center>

₁₅₃ We inferred phylogenies for simulated data sets aligned to reference genomes at

₁₅₄ varying distances to the ingroup taxa. We quantified differences between the true species

₁₅₅ tree and trees inferred from 210 different combinations of alignment and filtering options

₁₅₆ for 10 simulated data sets, and for two different choices of reference genome in each

₁₅₇ simulation (Fig. 1). We quantified differences between phylogenies for all simulated and

₁₅₈ empirical trees using three different tree distance metrics. To summarize multi-dimensional

₁₅₉ tree shapes and branching patterns, we also calculate summary statistics for tree shape,

Fig. 1. We used simulated data and varying bioinformatic filters to compare inferred phylogenies to the true topology used to generate data. This schematic shows parameter choices that we iterated over (rounded shapes) and steps performed from species tree simulation to data analysis (square shapes). Each round of simulation resulted in 420 filtered phylogenetic trees for analysis, and the simulation was replicated for 10 different simulated species trees at each speciation rate. $QUAL = mapping \ quality; \ ILS = incomplete \ lineage \ sorting; \ VCF = variant \ call \ format$

imbalance, and overall bootstrap support. For describing tree shape and branching

patterns, we focus on three metrics: Colless' imbalance statistic (I_c; Colless, 1982), the

Sackin imbalance statistic (Sackin, 1972), and the γ statistic (Pybus and Harvey, 2000).

The Colless and Sackin statistics both increase with increasing tree imbalance (i.e., more

pectinate topologies). The γ statistic is a measure of the distribution of branching events

within a phylogeny; a constant-rate pure-birth tree has an expected $\gamma = 0$, and an increase

in γ corresponds to branching times in a tree being closer to the tips (i.e., a less

bottom-heavy topology with higher center of gravity), while a negative γ estimate

indicates a tree with more diversification events toward the root (i.e., a more bottom-heavy

topology).

Simulated Data

To replicate the various properties of reduced-representation

genotyping-by-sequencing (GBS) or restriction-associated-digest sequencing (RADseq), we

simulated 60 independent data sets using the following procedure (Fig. 1): (1) simulate a

species tree for 50 taxa plus one outgroup taxon at 3 different levels of incomplete lineage

sorting; (2) simulate 2,000 gene genealogies for 5,000bp loci under the multi-species

coalescent conditioned on the simulated species tree from (1); (3) simulate genome

sequences and 150bp fastq reads for each individual for each gene genealogy from (2); (4)

align simulated fastq reads to one ingroup and one outgroup reference genome; (5) call

variants separately on ingroup and outgroup reference data sets; (6) filter variants with

different parameter combinations; (7) infer the maximum likelihood phylogeny for each

dataset; and finally, (8) calculate summary statistics on the output trees and model the

contributions of each bioinformatic filter to differences in these statistics. Each of these

steps is explained in detail below.

Species and gene trees.—   We simulated 50-taxon species trees using `SimPhy`

(Mallo et al., 2016) and employing the Yule process with three different birth rates (-SB

0.00001, 0.000005, 0.000001) to represent three different levels of shared polymorphism due

to incomplete lineage sorting (ILS; high, medium, and low, respectively, sensu Mirarab and

Warnow 2015). All simulations had a tree-wide effective population size of 50,000 (`-SP`

`f:50000`) and a ratio of 10 between the ingroup height and the branch from the root to the

ingroup (`-SO f:10`). We then simulated genealogies for 2,000 genes of 5,000bp each,

conditioned on the species tree for the given simulation and using a tree-wide substitution

rate of 3×10^{-9} per generation (similar to that estimated in Lake Malawi cichlid fishes;

Malinsky et al., 2018) for converting branch lengths to the expected number of

substitutions per site. The ancestral gene sequences for these genes were pulled randomly

without replacement from the empirical *Lates calcarifer* reference genome (Vij et al., 2016)

(see Supplemental Methods). With these simulation parameters, ILS levels and gene tree

discordance varied from a mean normalized Robinson-Foulds (RF) distance among gene

trees of 0.061 to 0.845 (low ILS, mean = 0.165; medium ILS, mean = 0.568; high ILS,

mean = 0.761).

Generating fastqs.—   We simulated fastq reads at each gene for each individual

using the `TreeToReads` pipeline (McTavish et al., 2017). `TreeToReads` uses `Seq-Gen`

(v1.3.4; Rambaut and Grass, 1997) for simulating variable sites and `ART`

(vGreatSmokyMountains; Huang et al., 2012) to simulate Illumina reads. The pipeline

generates simulated Illumina short-read data for individuals along an input gene genealogy,

using a given number of variable sites, model of evolution, read length, and fragment

length (McTavish et al., 2017). We pulled the number of variable sites for each fragment

from a normal distribution with mean of 150 and standard deviation of 30 to achieve

reasonably-sized SNP data sets. The location of mutations in the genome are drawn from a

uniform distribution and generated using a generalized time reversible (GTR) model of

sequence evolution with gamma rate heterogeneity (Lanave et al., 1984; Yang, 1993). The

₂₁₁ read depth for each individual at each locus was drawn from a uniform distribution

₂₁₂ between 2 and 20, to generate heterogeneity in read depth and introduce missing data. We

₂₁₃ specified a sequencing error file parameterized using the empirical GBS data in Rick et al.

₂₁₄ (2022), to mimic empirical errors introduced in the sequencing process. We simulated

₂₁₅ 500bp (s.d. 50) fragment lengths, to reflect size selection for ∼500bp fragments in

₂₁₆ generating reduced-representation data. Reads were then simulated as 150bp single-end

₂₁₇ reads using the Illumina HiSeq2500 profile in `ART`.

₂₁₈ *Aligning fastqs and calling variants.—* For each simulated tree, we randomly chose

₂₁₉ an ingroup taxon to be the ingroup reference, and used the simulated outgroup as our

₂₂₀ outgroup reference. We aligned all of the simulated fastq reads to each of these two

₂₂₁ genomes using `bwa mem` (v0.7.17, Li and Durbin 2009) and `samtools mpileup` (v1.6, Li

₂₂₂ 2011) with default parameters. We then filtered to retain only variable sites and called

₂₂₃ variants using `bcftools` (v1.9, Li et al. 2009). We retained variants in the raw VCF file

₂₂₄ that were biallelic (`--min-alleles 2` and `--max-alleles 2`), had a genotype quality

₂₂₅ score (`--minGQ`) >10, and had a minimum mapping quality (`--minQ`) >40.

₂₂₆ We then further filtered variant sites using minor allele count (MAC) and missing

₂₂₇ data (miss) filters, based on parameter ranges observed in published studies. We chose

₂₂₈ seven different minimum minor allele count cutoffs (0, 1, 2, 3, 4, 5, 10) and five different

₂₂₉ missing data cutoffs (0, 0.25, 0.5, 0.75, 0.90, corresponding to the percent of genotyped

₂₃₀ taxa required for a site to be retained) in `vcftools` (v0.1.14, Danecek et al. 2011), and

₂₃₁ created SNP data sets using all combinations of these filtering parameters. Individual

₂₃₂ genotypes were only called with a read depth ⩾5 (`--minDP 5`). Following filtering, the

₂₃₃ resulting VCFs were converted to phylip format and invariant sites were removed using the

₂₃₄ `raxml_ascbias.py` script v1.0 from `https://github.com/btmartin721/raxml_ascbias`.

₂₃₅ As a more quantitative measure of divergence between the reference and ingroup lineages,

₂₃₆ we calculated for each unfiltered data set (MAC = 0, miss = 0) the mean pairwise

₂₃₇ genome-wide divergence, d_{xy} (or π_{xy}; Nei and Li, 1979) between the reference genome and

all ingroup individuals using the `parseVCF.py` and `distMat.py` scripts from Simon

Martin's `genomics_general` GitHub repository

(https://github.com/simonhmartin/genomics_general).

Tree estimation.—   We concatenated variant sites and used `RAxML` for inferring

phylogenies for each of the 4,200 simulation and filtering combinations. We used a

GTRCAT model of rate heterogeneity with Lewis-type ascertainment bias correction

(Leaché et al., 2015) in a full search for the best scoring ML tree with rapid bootstrap

analysis (100 bootstraps) in the hybrid MPI/Pthreads version of `RAxML` (v8.2.12,

Stamatakis, 2014).

To measure the effect that each of the bioinformatic choices has on downstream

phylogenetic analyses, we calculated the information-corrected normalized RF distance

(`InfoRobinsonFoulds` from the `TreeDist` package in `R`; Smith, 2020) between each

inferred tree and the corresponding species tree (the "true" tree). Due to the inability of

RF distance to capture all aspects of tree differences (Smith, 2020), we used two additional

metrics of distance to the true tree: quartet distance (Estabrook et al., 1985) and

clustering information distance (Smith, 2020). Quartet distance was calculated using

`QuartetDivergence` using the tqDist algorithm (Sand et al., 2014) from the `Quartet`

package in `R` (Smith, 2019), and is a count of the number of four-taxon statements

(quartets) that differ between two trees. Clustering information distance scores a pair of

trees based on the amount of information that one partition contains about the other and

is an information-based tree distance metric that recognizes similarity in tree structure

even when every possible pairing of splits conflicts (Smith, 2020), making it more sensitive

than RF distance when trees are very dissimilar from one another. We also calculated two

measures of imbalance, the Sackin statistic (Sackin, 1972) and Colless' imbalance statistic

(I_c, Colless, 1982), as well as the tree center of gravity via the γ statistic (`gammaStat` from

`ape`; Pybus and Harvey, 2000) and branch length distributions. These statistics are often

used in diversification rate analyses, and therefore any systematic bias in their direction or

magnitude may have important consequences for evolutionary inferences made in these types of analyses. The Sackin and Colless indices were standardized using the PDA model to allow comparison between trees of different sizes (Blum et al., 2006). We additionally calculated imbalance and γ statistics using only the ingroup taxa and standardized these statistics by subtracting the value of each statistic for the true tree from the value calculated for the inferred tree, thus making those values the deviation from the true statistic (sensu Stadler et al., 2016).

Statistical Analyses

All statistical analyses were conducted in R (v4.0.4, R Core Team, 2021) and plotted using ggplot2 (Wickham, 2016). Scripts associated with simulations and analyses, as well as simulation outputs, are available at https://github.com/jessicarick/refbias_scripts.

Biases in lost loci.— We first examined the effect of bioinformatic filters on the number of loci (size of the data matrix) and the corresponding mutational spectrum of retained SNPs. We quantified locus-specific mutation rate as the number of simulated variable sites for a given 5000bp locus. We then compared mutation rates for SNPs pre- and post-filtering to investigate whether there were biases in the mutation rates of SNPs lost via different filters. We used pairwise Kolmogorov-Smirnov tests (ks.test in R) to statistically compare distributions of mutation rates from lost SNPs at each MAC and missing data cutoff. We also calculated the distribution of mutation rates of loci lost with increasing MAC and missing data filters to determine whether loci with higher mutation rates are more likely to be lost first, as has recently been demonstrated in other studies (e.g., Linck and Battey 2019; Huang and Knowles 2016; Shafer et al. 2017).

To investigate whether there is a pattern to where lost SNPs are located in the phylogeny, we compared the mean number of descendants for nodes retained in each filtering step. We used the phytools package (Revell, 2012) to compare each output tree

to the original simulated tree (using the `matchNodes` function), which provides us with information about which nodes from the original tree are missing in the output tree. We then recorded the bootstrap support (from `RAxML` output) and number of node descendants (using `getDescendants` from `phytools`) for each node in the analysis. For each combination of ILS level, MAC, missing data, and reference genome choice, we then created a null distribution for the expected mean node descendants and expected mean nodal bootstrap support using 100 bootstrap replicates, by taking the mean of n values drawn from the possible values for each of these, where n is the number of nodes lost across all trees with the given parameters choices. From these values, we calculated a z-score for the observed mean to quantify deviation from the bootstrap mean value. We repeated the bootstrapping procedure for the nodes retained in each parameter combination, and calculated z-scores for these as well.

Tree space analyses.— To characterize dissimilarity among the inferred phylogenetic trees within each simulation, we calculated the information-corrected normalized (Smith, 2020) RF distances (Robinson and Foulds, 1981) between all of the output trees that had the same starting species tree (`InfoRobinsonFoulds` from `TreeDist`; Smith, 2020), and then performed a principal coordinates analysis (PCoA) on these distances (`pcoa` from `ape`, Paradis and Schliep 2019). Using a PCoA approach, we can visualize differences between trees in multidimensional "tree space" and determine how much separation among trees occurs with different parameter combinations. We then calculated the correlation between tree position on the first two PCoA axes and each of our filtering parameters and reference genome choice for each group of trees.

Linear modeling.— To characterize the relationship between parameter choices (predictors) and the resulting trees and tree statistics (responses), we used linear mixed models (implemented using `lme4`; Bates et al., 2015) in `R`. In our linear mixed models, for each tree height, our predictor variables included reference distance (mean d_{xy} between the

ingroup taxa and the reference taxon), minor allele count cutoff (MAC), and missing data

cutoff, as well as pairwise interaction terms between reference distance and each of MAC

and missing data. We additionally included a random intercept for simulation number, such

that the overall model was $X \sim d_{xy} + MAC + miss + d_{xy} : MAC + d_{xy} : miss + (1|sim)$,

where X is our tree statistic of interest, d_{xy} is mean distance to the reference genome,

MAC is minor allele count, $miss$ is missing data threshold, and sim is simulation replicate

number. We assessed model fit using pseudo-R^2 values and assessed the contribution of

each predictor variable to the response statistic using regression coefficients. We

additionally assessed bivariate correlations between all variable pairs using Pearson's

correlation coefficient, implemented using `cor.test` in `R`.

Accounting for the influence of data set size

Because each filtering step removes SNPs, and therefore decreases the amount of

input data for phylogenetic analyses, we performed further steps to parse out whether the

observed effects were simply due to the loss of power in smaller data sets, or if additional

effects could be attributed to the specific filters applied and a bias as to which sites are

retained. To investigate the relative influence of each filter on the three response variables

(RF distance to true tree, imbalance, and gamma), as well as the mutation spectrum, we

repeated filtering steps and then subsampled each data set to retain only 6,000 random

SNPs. We then ran phylogenetic analyses in `RAxML` as described above. With these

subsampled trees, we performed the same statistical and modeling steps described above

for the full data sets, and compared these results to those with the full data sets.

Empirical data

To investigate to what extent the same patterns observed in our simulated data

appear in empirical data sets, we used two sets of GBS data from clades with similar ages,

but different number of species and expected levels of gene tree discordance: the radiation

of *Lates* fishes endemic to Lake Tanganyika and the radiation of cichlid fishes in the tribe

Tropheini, also endemic to Lake Tanganyika. Both of these clades have an estimated origin

around ∼1–2 Ma (Koblmüller et al., 2021, 2010; Irisarri et al., 2018, although Ronco et al.

2021 estimates the origin of the tropheine radiation to be ∼3.5–5.2 Ma), but very different

species richness (n=4 *Lates spp.*; n≈40 described and undescribed tropheine spp., as

estimated in Ronco et al. 2020). Thus, the tropheines represent a clade with high

speciation rates, shorter internodes, and thus potentially higher amounts of ILS, while ILS

is expected to be lower in the more slowly diversifying *Lates* clade. Sequences for the *Lates*

spp. were generated as described in Rick et al. (2022) (NCBI Sequence Read Archive

PRJNA776855) and data for the tropheines were generated in a similar manner following

the GBS protocol in Parchman et al. (2012), as described in the Supplementary Methods.

    Using the same methods as for the simulated data, the *Lates* fastq reads were

aligned to the *L. calcarifer* (distant outgroup) and *L. mariae* (ingroup) reference genomes.

The *Tropheus* fastq reads were aligned to the *Oreochromis niloticus* (distant outgroup) and

Pundamilia nyerei (closely related outgroup) reference genomes (Brawand et al., 2015). We

randomly chose 49 individuals for each of 10 data sets for each clade out of the n=229

high-quality (i.e., >100,000 reads assembled to each reference genome) sequenced

individuals for the *Lates spp.* and n=463 high-quality sequenced individuals for the

tropheines. We then added the alignments for the two reference genome individuals to

these data sets, so that the total number of tip taxa was n=51, matching our simulations.

Following alignment of fastq reads to the reference genomes, we iterated through missing

data (miss = 0, 0.25, 0.50, 0.75, 0.90) and minor allele (MAC = 0, 1, 2, 3, 4, 5, 10) filters

and inferred the maximum likelihood tree for each filtered data set using `RAxML`, using the

same parameter choices as with the simulated data. We then calculated the RF distance

between all trees for each data set and visualized differences among the trees using PCoA

in `R`. We calculated tree height and measures of imbalance and γ, and again used linear

mixed models to assess the influence that each of our reference genome choices and

Fig. 2. Example simulated species trees from `SimPhy` for low ILS (speciation rate = 0.000001; yellow), medium ILS (speciation rate = 0.000005; green), and high ILS (speciation rate = 0.00001; red) trees (sensu Mirarab and Warnow, 2015), and summary statistics for all simulated species trees. All trees were simulated with 50 taxa and an outgroup (not shown in phylogenies), on which the ingroup clade was rooted. For each tree, the mean across 10 simulated species trees for the gene tree discordance (as measured by mean pairwise Robinson-Foulds distance between gene trees; RF_{gt}), mean ingroup gamma (γ), and mean Colless imbalance statistic (I_c), along with their ranges, are given. Note that scales (in coalescent units) are not consistent among the trees shown. In the righthand panel, the distribution of values across all 30 simulated "true trees" is shown for ingroup tree height, average branch lengths, mean gene tree discordance, ingroup Colless imbalance, and ingroup gamma.

bioinformatic filters had on these tree characteristics. With the empirical results, we treated the identity of the reference genome as a binary option (i.e., ingroup/close versus outgroup/distant reference), rather than as a continuous d_{xy} variable, such that our models were $X \sim ref + MAC + miss + ref : MAC + ref : miss + (1|sim)$, where ref indicates reference genome choice and the ingroup/close reference was coded as 1 and the outgroup/distant reference was coded as 0.

Results

Simulations

Our simulation scheme resulted in a total of 4200 phylogenetic trees across 10 replicate simulations, which we analyzed to determine the influence of reference genome

and filtering choices on topology and tree shape metrics. Following filtering and removal of

invariant sites, our simulated data sets retained an average of 93,138 SNPs (95% quantile

= 1,067–444,006). The mean outgroup reference genome distance (d_{xy}) was 0.39 (95%

quantile = 0.30–0.50), while the ingroup reference genomes were an average distance of d_{xy}

= 0.039 (95% quantile = 0.034–0.047) from the other ingroup taxa. As was expected based

on simulation parameters, the mean distance to the outgroup reference was greatest in the

low-ILS trees (mean=0.44, range 0.39–0.50), lower in the medium-ILS trees (mean=0.38,

range 0.33–0.46), and lowest in the high-ILS trees (mean=0.37, range 0.30–0.46), although

all of these distributions overlapped. The mean distance to the ingroup reference was

similar among all of the tree heights (low-ILS trees: mean=0.040, range 0.034–0.046;

medium-ILS trees: mean=0.039, range 0.035–0.047; high-ILS trees: mean=0.040,

range=0.035–0.044). Consistent with expectations, the average amount of gene tree

discordance (i.e., mean pairwise normalized Robinson-Foulds distance between all gene

trees) in the low-ILS trees was 0.177 (range=0.124–0.241), in medium-ILS trees was 0.606

(range=0.480–0.746), and in high-ILS trees was 0.771 (range=0.661–0.845). Our simulated

species trees had overlapping distributions of normalized ingroup imbalance and gamma

among tree heights (Fig. 2), with a mean ingroup I_c of 0.412 and mean ingroup γ of -0.159.

Example filtered phylogenies can be found in Fig. S1.

Correlations among all variables are shown in Figure S3. Among all of our filtered

data sets, minor allele count (MAC) threshold and number of SNPs retained were highly

negatively correlated (Pearson's $r(1888) = -0.779, p < 2.2e - 16$; Fig. S2,S3). Our two

measures of imbalance (i.e., Colless' imbalance and Sackin imbalance) were highly

correlated as well (Pearson's $r(1888) = 0.988, p < 2.2e - 16$), and these were positively

correlated with metrics of topological distance to the true tree (i.e., Robinson-Foulds

distance, quartet distance, and clustering information distance; mean Pearson's

$r(1888) = 0.244$, all $p < 0.001$), although quartet distance was less strongly correlated with

imbalance than the other topological distance metrics. Ingroup γ was inversely correlated

to tree height (Pearson's $r(1888) = -0.793, p < 2.2e - 16$), and slightly positively

correlated to imbalance and topological distance metrics (Colless imbalance: Pearson's

$r(1888) = 0.331, p < 2.2e - 16$; RF distance: Pearson's $r(1888) = 0.358, p < 2.2e - 16$).

Topological distance to the true tree was negatively correlated with average bootstrap

support (Pearson's $r(1888) = -0.756, p < 2.2e - 16$). Heatmaps visualizing the way in

which each of the tree characteristics varied across MAC and missing data thresholds can

be found in Fig. S4–S11.

Distance to true tree.—   In our analyses, the ability to recover the input species

tree topology in simulated data sets varied widely across filtering parameters and reference

genome distance. All four of our linear mixed models for Robinson-Fould's distance

explained a majority of the variation in the data (mean pseudo-$R^2 = 0.706$, range

0.647-0.737; Table S1), and minor allele count had the greatest effect on the distance of the

inferred phylogeny to the true tree. MAC was an important predictor variable in our linear

models for distance to the true tree across all three simulated tree heights, with increased

MAC corresponding to topologies more distant to the true tree (Fig. 3). Missing data had

less of an effect on topological accuracy, and was not an important predictor variable in

our models for all trees combined, or for each of the three tree heights individually,

although increasing the stringency of this filter tended to increase the distance to the true

tree in our model combining all tree heights together (Fig. 3a). While we only used RF

distance in our linear mixed models, we observed similar trends across MAC in quartet

and clustering information distances, such that all three topological distance statistics

increased with increasing minor allele count threshold (Fig. S14).

The influence of reference genome choice had the greatest effect in high-ILS trees

and the weakest effect in low-ILS trees (Fig. 3b). Across all three topological distance

metrics, trees constructed with the ingroup reference genome were generally closer to the

true tree than those using the outgroup reference genome, although this pattern interacted

with MAC thresholds and tree height (Fig. S14). High ILS trees aligned to the ingroup

Fig. 3. Minor allele count (MAC) had the greatest effect on the distance from an output tree to the true tree, as well as the imbalance and γ statistic of the output tree, with some interaction between minor allele frequency and distance to the reference genome and tree height (i.e., amount of incomplete lineage sorting in the true tree). Plots in (a) and (b) show coefficient estimates and 95% confidence intervals for each predictor variable from linear mixed models using the indicated parameter as the response variable. Response variables are Robinson-Foulds distance (RF Dist) to the true tree, the gamma statistic for tree shape (γ), and Colless' imbalance statistic (I_c). Filled in and darkened symbols indicate significant positive or negative relationships, while faded symbols are predictors with confidence intervals overlapping zero. Plots in (a) are the results for all tree types combined, while those in (b) were modeled individually for each of the three simulated ILS levels. Plots on the righthand side show patterns of variation in the three response variables (c, Robinson-Foulds distance to the true tree; d, standardized Colless imbalance of the ingroup; e, standardized γ statistic of the ingroup) across minor allele counts for the full data sets, with shapes indicating the reference used (EXT, external reference; INT, internal reference) and colors indicating the level of ILS. In (c)–(e), the statistics are standardized so that the true tree is 0, as indicated with the solid horizontal lines. *For model variables, mac = minor allele count, avg_dxy = mean distance from ingroup taxa to the reference genome, miss = maximum missing data allowed, and colons show interaction terms between the given variables.*

⁴³² reference and filtered using low MAC thresholds were more distant to the true tree than

⁴³³ outgroup reference trees, but at high MAC thresholds the ingroup reference trees are more

⁴³⁴ similar to the true tree (i.e., have higher topological accuracy). Robinson-Foulds and

⁴³⁵ clustering information distances identified the greatest difference between trees inferred

⁴³⁶ using ingroup versus outgroup reference genomes at high MAC thresholds. Quartet

⁴³⁷ distance metrics (Fig. S14c) indicate that at low MAC thresholds for medium and high ILS

⁴³⁸ trees, trees constructed from data aligned to an outgroup reference genome were generally

⁴³⁹ more similar to the true tree than those aligned to an ingroup reference genome. In

⁴⁴⁰ contrast, at high MAC thresholds, ingroup reference genomes produced topologies closer to

⁴⁴¹ the true species tree for medium and high ILS trees. Trees with low ILS had a smaller

⁴⁴² quartet distance difference between ingroup and outgroup reference genomes, and ingroup

⁴⁴³ reference genomes produced trees more similar to the true tree across all MAC thresholds

⁴⁴⁴ (Fig. S14c).

⁴⁴⁵   Using PCoA to visualize tree space (Fig. 4), we find that the trees closest to the

⁴⁴⁶ true tree were generally those that used an outgroup reference genome and a minor allele

⁴⁴⁷ count filter less than 5, as also supported by our modeling results. For trees with low ILS,

⁴⁴⁸ the first two PCoA axes for the majority of simulations (90%) are correlated with minor

⁴⁴⁹ allele count and reference genome choice (Fig. 4b), while only 10% of simulations had

⁴⁵⁰ missing data correlated with separation among trees on either of the first two PCoA axes.

⁴⁵¹ For medium ILS trees, results were similar to those in our low ILS trees, but minor allele

⁴⁵² count and reference genome were more strongly correlated with distance among trees on

⁴⁵³ the first two PCoA axes: minor allele count was strongly correlated with the first PCoA

⁴⁵⁴ axis for 100% of simulations and reference genome identity was correlated with one of the

⁴⁵⁵ first two PCoA axes for 90% of simulations. For high ILS trees, we found that the first two

⁴⁵⁶ axes were correlated with minor allele count and reference genome in 100% of simulations,

⁴⁵⁷ while missing data was correlated with one of the first two axes in 60% of simulations.

⁴⁵⁸   As with the full data sets, both MAC and the interaction between MAC and

⁴⁵⁹ reference genome distance were significant predictors of topological accuracy in our linear

⁴⁶⁰ models (Fig. 5a; Table S2). Within the subsampled data sets, there was an even clearer

⁴⁶¹ improvement in tree topology at moderate MAC thresholds over low or high MAC

⁴⁶² thresholds, with all three distance metrics indicating that the most accurate topologies for

⁴⁶³ all tree heights and both reference choices occurred at MAC = 3–4 (Fig. S14a,c,e). The

Fig. 4. Principal coordinate analyses (PCoAs) based on pairwise Robinson-Foulds distances between trees demonstrate the relationship between simulated and true trees in two dimensional tree space. (a) Example PCoA plot of trees from one iteration of simulated data, with the true tree indicated and correlations between PCoA axes and each of reference genome choice, minor allele count (MAC), and missing data threshold visualized via arrows. (b) Summary of correlations between each of the three bioinformatic choices and the first two PCoA axes for all simulated and empirical iterations, demonstrating the relative importance of minor allele count in the simulated data and missing data in the empirical data sets, while reference genome was important for all. Dotplots show the distribution of values, colored by correlation significance (p-value <0.01 in solid colors, p-value $\geqslant 0.01$ in empty circles), assessed using Pearson's correlation coefficients. Lines below dots show median, 95% quantiles (thick lines), and full data range (thin lines) for each group of analyses.

subsampled data sets also indicate that in data sets without a MAC filter (i.e., MAC=0),

trees constructed with the ingroup reference genome are closer to the true tree than those

using the outgroup reference genome for high and medium ILS trees (Fig. 5).

Phylogenetic tree imbalance.— Our inference of tree imbalance also varied widely

across filtering parameters and reference genome choices. For the full data sets, minor

allele count (MAC) was the strongest predictor variable in our models (Fig. 3a; Table S1)

and the linear mixed models explained slightly over half of the variation in the data (mean

pseudo-$R^2 = 0.566$, range 0.466-0.669). The average distance of the reference genome to

the ingroup taxa was only an important predictor in our models for imbalance when all

tree heights were combined, such that an increased distance to the reference genome

increased I_c. In high ILS trees the interaction between MAC and the distance to the

reference genome was also a significant predictor of tree imbalance (Fig. 3b). Filtering for

missing data tended to reduce imbalance in the inferred trees, but the effect of this filter

Fig. 5. Predictor variable coefficients from the subsampled data sets, which were consistent with the original data sets with variable numbers of SNPs. Plots show model coefficients for linear mixed models for (a) all tree heights, and (b) each group of tree heights, partitioned into high ILS (speciation rate = 0.00001), medium ILS (speciation rate = 0.000005), and low ILS (speciation rate = 0.000001) trees. In (a) and (b), color intensity indicates whether coefficients for the given variable in our linear mixed models were significant (filled in and stronger colors) or not significant (gray or faded). For Robinson-Foulds distance (RF Dist), negative coefficients are closer to the true tree (i.e., smaller distance to the true tree), while values for imbalance and gamma have been standardized so that the true trees have a $\gamma = 0$ and $I_c = 0$. For model variables, mac = minor allele count, avg_dxy = mean distance from ingroup taxa to the reference genome, miss = maximum missing data allowed, and colons show interaction terms between the given variables.

on ingroup imbalance was not significant. Both low and medium ILS trees were closest to

the true tree imbalance statistic at $0 \leqslant \text{MAC} \leqslant 2$, while the high ILS trees were closest to

the imbalance of the true tree at $3 \leqslant \text{MAC} \leqslant 4$ (Fig. 3d). These patterns were the same

for both Colless and Sackin measures of imbalance (Fig. S14).

In the subsampled data sets, MAC again was the largest driver of tree imbalance,

with higher MAC thresholds resulting in more imbalanced trees and trees farther from the

true tree statistic (Fig. 5a,b; Table S2). The interaction term between MAC and distance

to the reference genome was an important predictor again, but the relationship was the

inverse of the trend seen in the full data sets (i.e., the interaction term was negative in the

full data sets, but positive in the subsampled data sets). In addition, missing data had a

significant influence on imbalance in the subsampled data sets, such that increasing the

missing data threshold reduces tree imbalance (Fig. S17b,d), a trend that was weaker but

in the same direction in the full data sets (Fig. S15b,d). In the subsampled data sets, trees

of all three ILS levels and both reference genome types were closest to the true tree

imbalance at $3 \leqslant \text{MAC} \leqslant 4$, as well as when the missing data threshold was most

stringent. However, the range of I_c statistics across all missing data thresholds was only

$\sim$0.005 for high and medium ILS trees, and only $\sim$0.001 for low ILS trees, in contrast to

the greater variation in I_c for all tree heights across MAC thresholds ($\sim$0.05).

Influence of parameters on gamma.—   Similar to topological accuracy and

imbalance, the linear mixed models for γ of the ingroup explained a large proportion of the

variation in the data (mean pseudo-$R^2 = 0.782$, range 0.727-0.823) and had MAC as the

strongest predictor of deviation from the true tree's γ statistic (Fig. 3), and was the only

significant predictor variable in any of our γ models. Across all three tree heights, the

values for γ were closest to the true tree at $0 \leqslant \text{MAC} \leqslant 2$ and increased with increasing

MAC (Fig. 3e). The change in γ across MAC values was greatest in the high and medium

ILS trees, and lowest in the low ILS trees, although the low ILS trees were the most biased

away from the true γ statistic across all MAC thresholds. The reference genome had little

influence on γ on its own or in interactions with MAC or missing data. Increased missing

data thresholds tended to reduce γ estimates, although the relationship was weak and the

variation in γ statistics across missing data values was <1 (Fig. S15f).

We see similar patterns for γ statistics in the subsampled data sets, with the

addition of missing data as a strong predictor of deviation from the γ of the true tree (Fig.

5). As with the full data sets, increasing the stringency of the MAC filter increased ingroup

γ, while increasing the stringency of the missing data filter reduced γ. The bias introduced

by both filters was strongest in high and medium ILS trees, while low ILS trees had the

most consistent γ values across filter thresholds (Fig. S16f,S17f). Across all filtering

combinations, the inferred topologies tended to be biased to more positive γ values than

the true trees, which were simulated under a pure-birth process and thus had $\gamma \approx 0$, or

just smaller than 0 (Fig. 2). However, values for γ in medium and high ILS trees at $0 \leqslant$

MAC $\leqslant 2$ ranged from γ=-6.00 to γ=-1.00, with a mean γ=-4.4 in high ILS trees and

mean γ=-3.2 in medium ILS trees. At MAC=10, these means then became significantly

positive: in high ILS trees, mean γ=3.6 (range 1.2-6.8) and in medium ILS trees, mean

γ=5.25 (range 0.3-7.9). In contrast, low ILS trees only varied from a mean γ=-0.4 (range

-2.8-1.5) at $0 \leqslant$ MAC $\leqslant 2$ to a mean γ=1.9 (range -1.08-5.6) at MAC=10.

In the γ models including all tree heights, we find that the distance to the reference

genome and the interaction between MAC and distance to the reference genome were

strong predictors in our models (Fig. 5a). However, neither term is significant in any of our

three models for individual tree heights (Fig. 5b), suggesting that the strength in the

all-height model may be an artifact of the correlation between mean distance to the

reference genome and tree height (which is not included as a predictor in these models;

Fig. S3), rather than being due to the distance to the reference genome itself. This

hypothesis was supported by a lack of difference between γ for trees aligned to the ingroup

versus outgroup reference genomes within each tree height across MAC (Fig. S16f) or

missing data (Fig. S17f)

Influence of parameters on mutation spectra.— The patterns we observed in tree

statistics suggest that there may be a bias in the characteristics of SNPs lost across

filtering parameters, and therefore we examined the mutation rates of SNPs lost at each

MAC and missing data cutoff. We found that SNPs lost at low MAC and missing data

cutoffs were biased toward those located on high mutation rate loci (Fig. 6a), which

resulted in MAC and missing data filters truncating the mutation spectra of retained SNPs

toward those with lower mutation rates. The change in mean mutation rate of retained loci

Fig. 6. In our simulations, SNPs that were lost first were biased towards those on loci with higher mutation rates for both minor allele count (MAC; a) and missing data (b) filters. Mean values for distributions are indicated with dashed lines, with colors corresponding to minor allele or missing data threshold. Righthand panels show significance from Kolmogorov-Smirnov tests on the distribution of values for each MAC and missing data threshold, with asterisks denoting significant differences ($p < 0.01$). Note that some distributions are completely overlapping with one another, and thus not visible (i.e., MAC=1 and MAC=2; MAC=3 and MAC=4; miss=0 and miss=0.25). Mutation rates were calculated as the number of variable sites simulated per 5000bp locus.

was most dramatic from MAC=2 to MAC=3, and from miss=0.75 to miss=0.9 (Fig. 6).

We also found that the SNPs lost due to MAC filters were those that provided

support for nodes toward the tips of the tree (Fig. 7a). We found that the lost nodes had

fewer descendants than expected based on a null distribution, and the lost nodes were also

those that had lower nodal bootstrap support originally (Fig. 7b). As the MAC threshold

increased, more nodes toward the tips in the tree were lost, which then lowered the mean

number of node descendants for lost nodes (Fig. 7a). When examining the bootstrap

support for nodes that were lost, we find that nodes lost at low MAC cut-offs had lower

bootstrap support than those retained, and increasing the MAC cut-off resulted in losing

nodes with progressively higher bootstrap support. There were no clear patterns in the

number of node descendants for nodes lost across different missing data thresholds (Fig. 7c): the mean number of node descendants of lost nodes remained constantly lower than that of retained nodes independent of missing data threshold. In contrast, the mean bootstrap support of lost nodes decreased with increasing missing data thresholds (Fig. 7d), indicating that low-support nodes are being filtered out at high missing data thresholds, but not at lower thresholds.

Empirical analyses

General overview of empirical data sets.— The ten replicates of each of our two empirical clades resulted in 2,800 inferred trees, constructed from random 51-taxa data sets with an average of 82,978 SNPs (tropheines; range 311–550,214) and 730,239 SNPs (*Lates*; range 1,147–16,018,003). The average distance of ingroup taxa to the reference genome for the 40-species tropheine clade's distant reference (*Oreochromis niloticus*) was d_{xy}=0.517, and for the close reference genome (*Pundamilia nyerei*) was d_{xy}=0.161. For the 4-species *Lates* clade, the average reference genome distances were d_{xy}=0.602 (outgroup reference, *Lates calcarifer*) and d_{xy}=0.0669 (ingroup reference, *L. mariae*). As with our simulated data, the number of SNPs retained was highly correlated with MAC (Pearson's $r(1292) =$ -0.141, $p = 3.5e - 07$) and with missing data threshold (Pearson's $r(1292) = $ -0.237, $p < 2.2e - 16$). The number of SNPs retained was also correlated with overall tree height (Pearson's $r(1292) = 0.384$, $p < 2.2e - 16$). Ingroup tree height was correlated with ingroup Colless and Sackin imbalance, although the correlation was negative in the tropheines and positive in the *Lates* clade (tropheines: Pearson's $r(593) = $ -0.275, $p = 8.287e - 12$; *Lates*: Pearson's $r(698) = 0.114$, $p = 0.0024$). In contrast to the simulations, tree height was not correlated with ingroup γ in either empirical clade (tropheines: Pearson's $r(593) = 0.006$, $p = 0.883$; *Lates*: Pearson's $r(698) = $ -0.012, $p = 0.746$).

⁵⁷²   *Tropheine analyses.—* For the two empirical data sets, we modeled ingroup tree

⁵⁷³ height, ingroup Colless imbalance (I_c), and γ as a function of minor allele count, missing

⁵⁷⁴ data threshold (miss), and reference genome identity (ref), as well as the interaction

⁵⁷⁵ between reference genome and each of minor allele count and missing data thresholds. In

⁵⁷⁶ contrast to the simulation results, the reference genome was treated as a binary choice

⁵⁷⁷ (ingroup/close reference versus outgroup/distant reference), with the more distant genome

⁵⁷⁸ coded as 0 and the close genome coded as 1. In our phylogenetic inferences for the

⁵⁷⁹ tropheine data, our models explained the largest proportion of the variation in the data for

⁵⁸⁰ γ (pseudo-R^2=0.600), less variation for imbalance (pseudo-R^2=0.336), and the least

⁵⁸¹ proportion of variation for height (pseudo-R^2=0.191). We found that MAC threshold was

⁵⁸² an important predictor of ingroup tree height and ingroup γ, and missing data threshold

⁵⁸³ was an important predictor of ingroup imbalance and ingroup γ (Fig. 8a). As the MAC

⁵⁸⁴ cutoff became more stringent, the inferred ingroup tree height increased (Fig. 8b) and γ

⁵⁸⁵ increased to become more positive (Fig. 8f). Imbalance decreased as MAC increased from

⁵⁸⁶ MAC=0 to MAC=3, and then the imbalance increased again between MAC=5 and

⁵⁸⁷ MAC=10 (Fig. 8d). With increasingly stringent missing data thresholds, ingroup tree

⁵⁸⁸ height increased (Fig. 8c), ingroup imbalance decreased (Fig. 8e), and ingroup γ decreased

⁵⁸⁹ (Fig. 8g).

⁵⁹⁰   We additionally found an influence of reference genome on ingroup tree height, as

⁵⁹¹ well as interactions between the reference genome and our filtering parameters (Fig. 8a). In

⁵⁹² the tropheine topologies, using the ingroup reference genome tended to increase tree height

⁵⁹³ compared to the outgroup reference genome. Using the ingroup reference genome also

⁵⁹⁴ resulted in trees that were more imbalanced and had higher γ statistics than trees

⁵⁹⁵ constructed using the outgroup reference genome. These patterns additionally interacted

⁵⁹⁶ with MAC: at low MAC thresholds, trees using the ingroup reference were longer, but they

⁵⁹⁷ became shorter than trees using the outgroup reference as MAC threshold increased (Fig.

⁵⁹⁸ 8b). At low MAC thresholds, trees using the ingroup reference were also more tip-heavy

⁵⁹⁹ than those using the outgroup reference, but had lower γ values than trees using the

⁶⁰⁰ outgroup reference at higher MAC thresholds (Fig. 8f). Conversely, ingroup reference trees

⁶⁰¹ had lower γ statistics than outgroup reference trees when all missing data was allowed

⁶⁰² (miss = 0), but then had higher γ statistics than the outgroup trees at higher missing data

⁶⁰³ thresholds (Fig. 8g).

⁶⁰⁴   In analyses of tree space based on pairwise RF distances between tropheine trees,

⁶⁰⁵ trees generally clustered together primarily based on missing data, with some influence of

⁶⁰⁶ reference genome and MAC (Fig. 4). The first two PCoA axes were generally correlated

⁶⁰⁷ with missing data (axis 1) and reference genome choice (axis 2), and a few iterations also

⁶⁰⁸ had strong correlations between these first two axes and MAC (Fig. 4c).

⁶⁰⁹   *Lates analyses.—* In our analyses of the *Lates* topologies, we found that our

⁶¹⁰ models explained the largest amount of variation in the data for imbalance (pseudo-$R^2 =$

⁶¹¹ 0.857), less variation for height (pseudo-$R^2 = 0.552$), and the smallest amount of variation

⁶¹² for γ (pseudo-$R^2 = 0.203$). The MAC threshold was again an important predictor of

⁶¹³ ingroup tree height and ingroup γ (Fig. 8a). However, the relationship between MAC and

⁶¹⁴ γ was in the opposite direction to the tropheines and the simulated data: increased

⁶¹⁵ stringency in the MAC filter reduced ingroup tree height and reduced ingroup γ values,

⁶¹⁶ except for a brief increase in γ at MAC=3 (Fig. 8f). In addition, reference genome choice

⁶¹⁷ interacted with MAC to influence γ values, with trees using the outgroup reference

⁶¹⁸ becoming more negative at high MAC values than those using the ingroup reference.

⁶¹⁹ Ingroup imbalance decreased slightly from $0 \leqslant$ MAC $\leqslant 2$ to $3 \leqslant$ MAC $\leqslant 5$, and then

⁶²⁰ increased again at MAC=10 (Fig. 8d). In addition, reference genome choice had a strong

⁶²¹ influence on imbalance statistics: trees aligned to the outgroup reference genome tended to

⁶²² be more imbalanced than those aligned to the ingroup reference.

⁶²³   Missing data was an important predictor of ingroup height and ingroup imbalance

⁶²⁴ in the *Lates* data, such that increasingly stringent missing data filters resulted in shorter

⁶²⁵ and more imbalanced trees (Fig. 8c,e). The interaction between missing data and reference

Fig. 7. a) Nodes lost from trees tended to be those with fewer node descendants, which reduced the mean number of descendants of lost nodes as more nodes were lost at increasing minor allele thresholds. Mean number of node descendants for nodes lost (red) and retained (blue) at each minor allele count threshold, as well as the z-scores for the deviation of these means from a null distribution for nodes randomly drawn from the species tree. b) Nodes with lower bootstrap support were generally lost first at low MAC thresholds. Plots show the mean bootstrap support values for those nodes that are lost at each minor allele count threshold, as well as the distribution of z-scores for the deviation of these values from a null distribution of bootstrap support values. c) Mean number of node descendants for nodes lost at each missing data cutoff, demonstrating a generally lower number of node descendants for lost nodes but no clear trend as missing data cutoffs become more stringent. d) Mean bootstrap support values for nodes lost at each missing data cutoff, demonstrating a reduction in mean bootstrap support for lost nodes as the missing data threshold becomes more stringent. In the lefthand plots for each panel, means and standard errors are displayed, as calculated using the `mean_se` function in `ggplot2`. Horizontal dashed lines indicate the overall mean bootstrap support and node descendants across all nodes, respectively. In righthand panels, dashed lines indicate $|z| = 1.96$.

genome choice was not an important predictor in any of our models with the *Lates* data

(Fig. 8a).

Lates trees generally clustered together in PCoA tree space primarily based on

reference genome, with some influence of missing data and MAC. The first PCoA axis

generally correlated with reference genome choice, while the second axis was

predominantly correlated with missing data (Fig. 4c). The MAC filter only correlated with

one of the two main PCoA axes in one of the ten *Lates* iterations.

Fig. 8. In the empirical data sets, missing data, minor allele count, and the choice of reference genome affected the inferred tree height, imbalance, and γ statistics. Plots show model coefficients from linear mixed models for empirical data, with missing data (miss), minor allele count (mac), reference genome (ref), the interaction between missing data and reference genome choice (int:miss), and the interaction between minor allele count and reference genome choice (int:mac) as explanatory variables, and tree height, imbalance, and gamma (γ) in the inferred phylogenies as response variables. Results are shown for tropheine (orange) and *Lates* (teal) empirical data sets, with univariate responses across minor allele counts and missing data thresholds shown in B-I for our response variables of interest. For the *Lates*, the average D_{xy} between the ingroup taxa and the outgroup reference was 0.628, while the ingroup reference average D_{xy} was 0.0345. For the tropheines, these average values were $D_{xy} =$ 0.517 and $D_{xy} = 0.161$ for the more distant and more closely related genome, respectively.

Discussion

⁶³⁴   Single nucleotide polymorphism (SNP) data sets are used extensively in population

⁶³⁵ genomic and phylogenomic studies, but best practices in the bioinformatic choices made

⁶³⁶ while aligning and filtering these data remain unclear for both reduced-representation and

⁶³⁷ whole genome sequencing data sets. There is no generally accepted analysis pipeline for

⁶³⁸ these data, and it is unclear whether a "one size fits all" approach is warranted. Our

⁶³⁹ results from a combination of simulated and empirical analyses demonstrate how decisions

⁶⁴⁰ made by researchers during the assembly and processing of genomic data, as well as

⁶⁴¹ characteristics of the true divergence history among taxa, influence the final data set and

⁶⁴² the inferred phylogenetic and diversification history. We found that the value chosen for

⁶⁴³ minor allele filters interacts with the choice of reference genome and total tree height (and

thus ILS) to bias phylogenetic inferences away from the true divergence history. This suggests that removing rare variants reduces phylogenetic accuracy, particularly toward the tips of a tree. In contrast, missing data filters can reduce bias in the inferred phylogenetic tree. We also examined the effect these parameter choices have on tree balance and shape. In our simulations, we found that increasing the minor allele threshold generally has negative effects on the accuracy of tree shape and balance statistics, increasing the topological distance, increasing the imbalance, and increasing the γ statistic of inferred trees compared to the true tree. Our results complement and extend those from other recent studies—which focused mostly on single filtering and alignment parameters of interest (e.g., Linck and Battey, 2019; Huang and Knowles, 2016; Prasad et al., 2021; Günther and Nettelblad, 2019; Shafer et al., 2017; Molloy and Warnow, 2018; Lanier et al., 2014)—by demonstrating that the choice of reference genome can also bias phylogenetic analyses, as well as exacerbate or modify the effects of individual filtering parameters. Furthermore, a number of these effects are independent of the overall size of the final data sets, suggesting that it is the nature of the remaining data—not merely the volume—that biases phylogenetic inference.

Stringent minor allele filters bias topological inference and diversification statistics

Most variation in natural populations is rare, and rare variants are those that contain the most information about differentiation between individuals or populations (Slatkin, 1985; Nelson et al., 2012); thus it is perhaps unsurprising that we found a clear pattern that stringent minor allele filters bias topologies away from the true evolutionary history. This was true in both our full dataset, and in the datasets we subsampled to account for the influences of differences in dataset size on these inferences. Biased trees tended to be more imbalanced and more tip-heavy (i.e., had a less negative γ) than the true species trees. The topologies closest to that of the true tree were at minor allele count (MAC) 3–4, with a more dramatic improvement from MAC=1–2 to MAC=3–4 in the

subsampled trees than in the full data sets. The inferred topologies generally had imbalance statistics closest to that of the true tree at moderate MAC filters as well. This suggests that excluding sites with very low-frequency alleles (<3 copies across all samples in the data set, corresponding to a minor allele frequency of 1-3%) leads to more accurate results in phylogenetic analyses, likely by removing SNPs that are truly the result of sequencing or alignment errors. However, higher thresholds remove true phylogenetic signal from the data set, reducing data quality and biasing inferences toward trees that are more pectinate than the true phylogenetic history.

Inferred topologies had ingroup γ statistics closest to that of the true trees at our lowest MAC thresholds (MAC=0–2) and became much more biased at MAC=3–4. Our simulated trees were created using a pure birth process, which produced species tree γ statistics that follow a standard normal distribution (with a mean $\gamma \approx 0$ and sd=1). Thus γ values for trees which are <-1.96 or >1.96 fall outside the 95% CI and are considered significant evidence for rejecting a constant rate of diversification. The magnitude of change between phylogenies filtered with MAC=0 and MAC=10 was great enough to change γ from significantly negative ($\gamma = $ -5.73) to significantly positive ($\gamma = 7.84$; Fig. S4–S6), and the mean absolute change in γ was 3.28 for low ILS trees, 8.55 for medium ILS trees, and 8.02 for high ILS trees and thus ranges from nearly equal to the width of the 95% CI to more than double it. This suggests that the MAC filter alone could dramatically change inferences of diversification rates through time and, importantly, lead to incorrect macroevolutionary inferences at high MAC thresholds.

These findings follow from theoretical expectations that recent splits in the phylogeny have less shared variation supporting them (i.e., more low-frequency SNPs or rare variants), and thus are disproportionately lost as MAC increases (Linck and Battey, 2019). This is supported by our findings that the majority of nodes lost due to the minor allele filter were those with few node descendants (Fig. 7a) and therefore less hierarchical redundancy (sensu Eaton et al., 2017). The SNPs lost first at lower MAC filters also

tended to be those with higher mutation rates (Fig. 6). At $0 \leqslant \mathrm{MAC} \leqslant 2$, we thus gained

topological accuracy but reduced branch lengths near the tips of the trees, shifting the

tree's center of gravity tipward and increasing γ estimates. This process may have also

driven the increase in tree imbalance: the loss of more recently evolved SNPs also removes

lineage-specific variation from nested clades, which then may lead to more pectinate

topologies.

Some MAC-related bias was more apparent in our subsampled data sets than our

full simulations, suggesting an effect of the overall data set size on the general trends in

our parameters of interest described above. The most apparent of these was the shape of

the topological accuracy response across MAC thresholds: the full data sets showed a slight

increase in topological accuracy from MAC=1–2 to MAC=3–4 (Fig. 3c), while the

subsampled data sets showed a much more dramatic increase in accuracy at MAC=3–4

(Fig. 5c). This suggests that the deviation from the true tree when $\mathrm{MAC} \leqslant 2$ in the

subsampled data sets may have been due to having fewer SNPs (i.e., less data) than the

full data sets, and therefore the signal from any "erroneous" SNPs (e.g., those that result

from sequencing errors) was stronger and more likely to bias the inferred tree away from

the true topology. Thus, filtering out these true sequencing or alignment errors did improve

topological accuracy. In this way, we found that using a MAC filter is more important in

data sets with fewer overall SNPs, although this relationship will need to be explored

further in future studies.

These results together suggest that trade-offs exist when using a minor allele filter

between reducing SNP errors, improving topological accuracy, and improving accuracy of

diversification statistics. The general consistency observed after controlling for the size of

data sets in our subsampling inferences suggests that the numerical loss of SNPs when

using more stringent filters was not the only factor that drove these patterns, but rather

that there are biases in the characteristics of the SNPs being lost. This also suggests that

caution should be used in evaluating diversification statistics in trees inferred using heavily

filtered data. This echoes others' findings that filtering based on minor allele frequency can

bias population genetic inferences, especially for events in the recent past (Linck and

Battey, 2019; Shafer et al., 2017; Boitard et al., 2016), and provides further reason for

using caution when interpreting macroevolutionary patterns from phylogenies of extant

taxa (Louca and Pennell, 2020; Kubo and Iwasa, 1995; Louca and Pennell, 2021).

Some of the patterns in our empirical data mirror our observations from the

simulations: we found that MAC also had a strong influence on the inferred empirical

phylogenies. The change in γ with MAC was less evident in our empirical data sets;

however, ingroup γ decreased in the *Lates* phylogenies, transitioning from a non-significant

γ estimate to significant and negative γ at more stringent MAC thresholds (Fig. 8f,S11).

Thus, macroevolutionary inferences in this clade would differ widely at different MAC

thresholds. Estimates of γ increased slightly with increasing MAC filters in the tropheines,

although it remained significantly positive among all MAC thresholds and thus would not

change macroevolutionary inferences (Fig. 8f,S10). Whereas imbalance for the simulated

data sets increased across all MAC thresholds, first becoming more accurate before

exceeding the true value across most tree heights, imbalance actually decreased for both

empirical data sets from MAC=0 to MAC=3, and then increased again above MAC=5

(Fig. 8). Unlike the simulations, we do not know the true value for these data sets and

cannot know whether the values we estimated deviate away from the true statistic.

However, the consistency of the observed effect may indicate some common feature of the

empirical datasets affected by MAC thresholds.

In both the subsampled data sets and the empirical data sets, there were differences

between the responses at MAC<3 and MAC>4. In the subsampled data sets, this was a

general minimum in divergence from the true tree, while the truth is unknown for the

empirical data sets. That this threshold would be an inflection point is not surprising:

given that these are diploid datasets, SNPs filtered at MAC=3 but not MAC=2 must be

shared by at least two tip taxa. Consequently, this is the first MAC filter that must remove

shared variation, rather than alleles found in only a single taxon and thus it is expected to

have a broader influence on tree topology and tree metrics.

Missing data filters affect diversification statistics

In contrast to the patterns with minor allele filters, stringent missing data filters
(i.e., those that removed sites with lower amounts of missing data) were associated with an
improvement in the accuracy of imbalance and γ statistics, and had little effect on
topological accuracy. The subsampled data sets demonstrated a clear pattern that
increased missing data filters (i.e., allowing less missing data per site) resulted in less
deviance of both imbalance and γ from the true values (Fig. 5a). Our full data sets trended
the same direction, although missing data was not a statistically significant predictor in
our models (Fig. 3a).

Missing data had a stronger influence on imbalance and γ estimates in our
empirical data sets than in our simulated data sets. While we do not know truth for the
empirical data sets, we found a negative relationship between these two statistics and
missing data for both the *Lates* and the tropheines: the tropheine data set showed a strong
negative relationship between missing data cutoff and both imbalance and γ, and the *Lates*
phylogenies were also less imbalanced with more stringent missing data filters. This likely
reflects aspects of our empirical data that are not captured in our controlled simulations,
such as biases in which sites are most likely to have high levels of missing data (e.g., allelic
dropout due to restriction site mutations, which has been incorporated in other simulation
studies that find a strong bias due to missing data; Huang and Knowles, 2016; Eaton
et al., 2017). Our simulated data also had low overall rates of missing data: an average of
only 0.51% of sites in the unfiltered simulated data had >50% missing data (mean per-site
missing data = 2.21%). In contrast, our empirical data sets had an average of 97.0% of
sites with >50% missing data (mean per-site missing data = 94.3%). This discrepancy led
to fewer sites being lost due to missing data filters in the simulated data sets, which then

⁷⁷⁷ would lead to less variation between phylogenies inferred at different missing data

⁷⁷⁸ thresholds.

⁷⁷⁹   Despite the low amount of missing data present in our simulated data sets, we did

⁷⁸⁰ find that filtering for missing data improved the accuracy of our imbalance and γ estimates

⁷⁸¹ without much reduction in topological accuracy. Diversification statistic inferences were

⁷⁸² most accurate for simulated data at our most stringent threshold for missing data

⁷⁸³ (miss=0.9; Fig. S15,S17). This suggests that the reductions in imbalance and γ statistics

⁷⁸⁴ with increasingly stringent missing data filters in both of the empirical data sets (Fig. 8e,g)

⁷⁸⁵ may indicate more accurate inferences as well; however, without knowing the true value for

⁷⁸⁶ these statistics, we do not know whether these trends represent more or less accuracy in

⁷⁸⁷ macroevolutionary inferences. Even with the large number of sites being filtered due to

⁷⁸⁸ missing data in the empirical data sets, the difference in these statistics between miss=0.25

⁷⁸⁹ and miss=0.90 was not great enough to change macroevolutionary inferences based on γ or

⁷⁹⁰ imbalance statistics. In addition, we found that filtering sites for missing data biased the

⁷⁹¹ mutation spectrum of the retained loci by disproportionately removing SNPs on loci with

⁷⁹² the highest mutation rates (Fig. 6). Thus, the small possible improvement in imbalance

⁷⁹³ and γ accuracy may not outweigh the previously demonstrated drawbacks to filtering

⁷⁹⁴ stringently for missing data in real-world data sets (e.g., Huang and Knowles, 2016; Eaton

⁷⁹⁵ et al., 2017; Chan et al., 2020).

⁷⁹⁶   *Reference genome choice can bias topological inference and tree imbalance*

⁷⁹⁷   In addition to minor allele and missing data filters influencing the accuracy of

⁷⁹⁸ inferred trees, our results suggest that reference genome choice can strongly influence

⁷⁹⁹ phylogenetic insights gained from short read data. The difference between trees aligned to

⁸⁰⁰ an ingroup versus an outgroup reference genome was most evident in principal coordinate

⁸⁰¹ analyses of Robinson-Foulds distances among all trees that came from the same simulated

⁸⁰² species tree, where the choice of reference genome was correlated with the first or second

PCoA axis in the majority of analyses (Fig. 4b). In PCoA tree space, trees aligned to the outgroup reference generally clustered closer to the true tree than those aligned to the ingroup reference (e.g., Fig. 4a,S12), although this was not true in every case. However, our models revealed that the distance to the reference genome did not have a strong influence on topological accuracy on its own in our simulated data sets. This likely arose due to the statistics used in these inferences: absolute Robinson-Foulds distance to the true tree (used in our modeling) describes the one-dimensional magnitude of topological differences, but not *how* a tree is different from the true tree or whether it deviates in the same way that other trees deviate from the true tree. Thus, PCoAs pick up more detail about how similar the inferred trees are to one another, which allows inferences about which trees differ from the true tree in similar ways. In doing so, the PCoA results suggest that reference genome choice may be even more important to the shape of the inferred topology than our modeling results indicate.

Increasing the average distance of the reference genome to the ingroup taxa also increased imbalance in the inferred topologies, although the influence of reference genome on its own was not a significant predictor in either our full or subsampled simulations. These patterns may have arisen due to biases in the different characteristics of loci retained when using an ingroup versus outgroup reference genome: the former are more likely to retain low-frequency variation within the ingroup and the latter are more likely to retain conserved regions with low mutation rates (i.e., sites where the ingroup taxa are more similar to the reference genome; Ros-Freixedes et al., 2018; Brandt et al., 2015). The loss of low-frequency variation within the taxa of interest when using an outgroup reference genome may lead to a lack of support for nested clades within the phylogeny, which then leads to more imbalanced and pectinate topologies.

We find the same pattern in the *Lates* empirical data: trees aligned to the more distant reference genome tended to be more imbalanced than those aligned to the ingroup genome. Intriguingly, our tropheine empirical data trended in the opposite direction, with

the ingroup reference genome producing trees that were more imbalanced than those using

the outgroup reference genome. There are several possible explanations for this. First, we

do not know the true level of imbalance for the evolutionary history for these empirical

clades, and thus it is possible that higher imbalance in the tropheines and lower imbalance

in the *Lates* both produced less biased imbalance estimates, in which case these results

would agree with our simulated data. Second, the different responses between empirical

clades may have been due to the fact that the tropheine clade's closer reference

(*Pundamilia nyerei*) is not actually nested within the clade, but rather is a more closely

related outgroup than the "outgroup" reference (*Oreochromis niloticus*). Thus, the mean

distance to the close reference genome was greater in the tropheine clade (d_{xy}=0.161) than

for the *Lates* ingroup reference (d_{xy}=0.0669) or the simulation ingroup references

(d_{xy}=0.039). Third, the 40-species clade of tropheines is likely to have more low-frequency

variation (i.e., rare variants) captured than the 4-species clade of *Lates* and these

low-frequency alleles may be disproportionately lost when using a distant outgroup

reference genome. An extensive body of work already exists on biases introduced via taxon

sampling choices (e.g., Zwickl and Hillis, 2002; Heath et al., 2008b; Hohna et al., 2011;

Cusimano and Renner, 2010; Brock et al., 2011), and our results indicate that the choice of

reference genome may interact with taxon sampling schemes to bias phylogenetic inferences

and diversification statistics. We did find that there is a smaller difference in the mean

number of SNPs retained between data sets aligned to the close versus distant reference

genomes for the tropheine clade (difference in means = 9,552 SNPs, with more SNPs

retained with the close reference) than in the *Lates* clade (difference in means = 57,643

SNPs, with more SNPs retained with the outgroup reference). These differences in number

of SNPs may indicate differences in the characteristics of SNPs retained as well. Finally, it

is also possible that reference genome quality affects phylogenetic inferences, such that

more complete reference genomes retain more SNPs or that more fragmented genomes tend

to retain more conserved variation. The distant reference genomes used in our empirical

analyses here were both chromosome-level assemblies (*L. calcarifer*, 24 chromosomes; *O. niloticus*, 22 chromosomes), while the close reference assemblies are not yet assembled into chromosomes (*L. mariae*, 1,154 scaffolds; *P. nyererei*, 7,236 scaffolds). This pattern—having a distant, well-assembled genome and a closer, more fragmented genome available—is common when making the choice between a close versus distantly related reference genome. Thus some of the effect of reference genome distance that we observe in our empirical analyses (and perhaps some of the differences between our empirical and simulation results) may be linked to assembly completeness. While genome quality does not appear to bias population genomic demographic reconstruction methods (Patton et al., 2019), it remains unclear whether phylogenetic inferences are similarly unaffected.

If multiple reference genome choices are available, one way to potentially mitigate the bias introduced due to reference genome choice may be to map reads to multiple reference genomes, as recommended in Bertels et al. (2014), who found that the choice of reference taxon affected the percentage of incorrect topologies in their simulations, but demonstrated that this bias was reduced by replicating analyses over multiple reference genomes. Our results support this recommendation, as mapping to multiple reference genomes allows an exploration of how the topology changes when the data are aligned to one reference versus another. Another promising possibility to improve such bias is using multiple references to create a "pseudogenome" for the species of interest (Huang et al., 2014), as demonstrated in Sarver et al. (2017). Both of these options are increasingly possible due to the acceleration in genome sequencing for non-model taxa (Hotaling et al., 2021; Rhie et al., 2021; Formenti et al., 2022). However, not every clade of interest will have multiple reasonable reference genome options. In this case, our results can be used to inform filtering practices: stringent filtering thresholds (particularly for minor allele count) will bias the results more when using a distant reference genome than when using an ingroup (or closely related) reference genome.

Magnitude of bioinformatic biases depend on interactions and the true divergence history

No bioinformatic choice is made in isolation, and our results demonstrate that the magnitude of bias introduced by each decision depends both on the true evolutionary history of the clade of interest and on interactions among bioinformatic choices. Generally, our longest trees were the most resistant to filtering-related deviations in topology and imbalance (Fig. 3,5), suggesting that trees with low ILS are the least prone to bias arising from bioinformatic choices. In longer trees, larger internode distances mean that there are more redundant SNPs supporting each split in the tree and more time for lineage sorting to occur among ancestral polymorphisms, and this redundancy may allow for correct inferences despite biased patterns to the loss of SNPs. The low ILS trees were also generally further from the true γ statistic than medium or high ILS trees, despite also having more consistent γ statistics across MAC thresholds (Fig. 3e,5e). The deviation in γ at low MAC thresholds for low ILS trees may be due to proportionally more SNPs occurring—and also being lost—along the tip branches in the low ILS trees, but more work is needed to interrogate this effect.

Our two empirical data sets, which had different amounts of gene tree incongruence, different imbalances, and different diversification histories, demonstrate how these bioinformatic filters can interact with a clade's evolutionary history. Overall, we saw much more dramatic responses in tree characteristics in the *Lates* trees (particularly tree height and γ; Fig. 8c) than in the tropheine trees across minor allele thresholds, while missing data had a greater effect on the tropheine trees than the *Lates* (particularly imbalance; Fig. 8f,h,j). While we do not know the "true" statistics for the empirical analyses, we found that γ, imbalance, and overall tree height all showed an inflection point around MAC=3–4, as was the case with our simulated data. However, the shape of these statistics' responses to MAC across this inflection point differed between the tropheine, *Lates*, and simulated trees (Fig. 8, S14–S17). In the *Lates* analyses, the highest γ values were at moderate minor allele and moderate missing data thresholds. In the tropheine data set, on

the other hand, γ increased slightly with increasing minor allele thresholds and decreased with increasing missing data thresholds.

These interactions suggest that the best filtering scheme differs for data and clades with different characteristics (e.g., the rapidity of diversification within the clade, the amount of incomplete lineage sorting, the level of missing data or sequencing errors within the data set, and the overall size of the SNP data set). With smaller data sets and data sets with higher levels of ILS, moderate minor allele filters (i.e., MAC=3–4) likely improve topological inferences, but higher minor allele filters (i.e., MAC >5) will then reduce the accuracy of inference. However, in trees with low levels of ILS, little topological accuracy is gained with shifting from low ($0 \leqslant$ MAC $\leqslant 2$) to moderate (MAC=3–4) minor allele filters. When using an outgroup reference, high MAC filters will have a stronger effect in reducing the accuracy of inferences, while high MAC filters may result in more accurate trees when using an ingroup reference.

It is also worth noting that our simulations only included trees simulated under an equal-rates Yule model (with equal speciation rates and zero extinction across all lineages), resulting in balanced topologies with a true $\gamma \approx 0$. In clades with true divergence histories that are more pectinate or have a higher or lower center of gravity, it is possible that differences in the distribution of SNPs across the tree may cause response patterns to differ. Similarly, imbalanced taxon sampling across the topology (e.g., clustered sampling) may also affect the way that a data set responds to bioinformatic choices, and thus it will be important to investigate these patterns across a variety of different tree shapes to understand our ability to generalize these results.

Conclusions and Recommendations

Our results suggest that stringent bioinformatic filters tend to reduce the accuracy of phylogenetic analyses and should therefore be used with caution. However, best practices for filtering will depend on the specifics of the divergence history of a clade and

the reference genome(s) available for short read alignment. In particular, the rapidity of

diversification within a clade is predictive of the sensitivity of phylogenetic analyses to

biases introduced by minor allele and missing data filters, such that trees with high

amounts of gene tree incongruence require these filters to improve inferences, but also show

the most bias compared to the true topologies and true diversification statistics as filter

stringency increases. In addition, we find that data aligned to more distant reference

genomes are more sensitive to stringent filtering parameters. We further demonstrate that

minor allele filters greater than $\sim$4% reduce topological accuracy in our 51-taxon

phylogenies and increase both imbalance and γ statistics, suggesting that they are

removing important signal above this threshold rather than only removing sequencing or

alignment errors.

Intriguingly, we also demonstrate that the best filtering scheme for producing an

accurate topology and imbalance statistic may differ from the optimal scheme for accurate

γ inference. Thus the optimal filtering scheme likely differs depending on the intended

downstream inferences. While these general patterns are clear from our simulation results,

we recommend that researchers still replicate their own analyses over a series of

bioinformatic parameter combinations to ensure that results are in agreement across a

range of bioinformatic choices. While our results make clear that reference genome choice

and bioinformatic filters can bias phylogenetic inferences and downstream

macroevolutionary interpretations, the increasing availability of reference-quality genomes

makes it possible to implement multiple alignment and filtering schemes to assess the effect

that each choice has for the research question of interest. We encourage careful assessment

of these choices in empirical data sets.

Funding

This work was supported by a National Science Foundation grant to C.E.W. (grant

number DEB-1556963); by a Graduate Student Research Award from the Society of

Systematic Biologists to J.A.R.; by the Aven Nelson Fellowship in Systematic Botany from the University of Wyoming to J.A.R.; by a University of Wyoming NASA Space Grant Research Fellowship from the Wyoming NASA Space Grant Consortium to J.A.R. (NASA Grant number NNX15AI08H); by financial support provided by Robert B. Berry and the University of Wyoming to C.D.B.; and by the University of Wyoming INBRE Data Science Core, which is supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health (grant number 2P20GM103432).

Acknowledgements

The authors would like to thank E.J. McTavish for the use of and help with the TreeToReads pipeline. We also thank C.A. Buerkle for insightful discussions about the role of minor alleles and rare variants in evolutionary biology. We also thank members of the Wagner Lab at the University of Wyoming for many discussions related to this work that shaped the direction of this research. This work was made possible through the use of the University of Wyoming Advanced Research Computing Center's Mt. Moran (http://n2t.net/ark:/85786/m4159c) and Teton (https://doi.org/10.15786/M2FY47) computing environments. The findings of this paper are solely the responsibility of the authors and do not necessarily represent the official views of NIH, NSF, or other funding agencies.

Data Availability Statement

Simulated data, simulation scripts, and analysis scripts are available from the Zenodo Repository: `http://dx.doi.org/10.5281/zenodo.5940691`. Scripts used in simulations and analyses are also available on GitHub

`https://github.com/jessicarick/refbias_scripts`.

References

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67.

Bertels, F., O. K. Silander, M. Pachkov, P. B. Rainey, and E. Van Nimwegen. 2014. Automated reconstruction of whole-genome phylogenies from short-sequence reads. Molecular Biology and Evolution 31:1077–1088.

Bhatia, G., N. Patterson, S. Sankararaman, and A. L. Price. 2013. Estimating and interpreting FST: The impact of rare variants. Genome Research 23:1514–1521.

Biddanda, A., D. P. Rice, and J. Novembre. 2020. A variant-centric perspective on geographic patterns of human allele frequency variation. eLife 9:e60107.

Blum, M. G., O. François, and S. Janson. 2006. The mean, variance and limiting distribution of two statistics sensitive to phylogenetic tree balance. Annals of Applied Probability 16:2195–2214.

Boitard, S., W. Rodríguez, F. Jay, S. Mona, and F. Austerlitz. 2016. Inferring Population Size History from Large Samples of Genome-Wide Molecular Data - An Approximate Bayesian Computation Approach. PLOS Genetics 12.

Brandt, D. Y. C., V. R. C. Aguiar, B. D. Bitarello, K. Nunes, J. Goudet, and D. Meyer. 2015. Mapping Bias Overestimates Reference Allele Frequencies at the HLA Genes in the 1000 Genomes Project Phase I Data. G3 Genes—Genomes—Genetics 5.

Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, I. Keller, S. Fan, O. Simakov, A. Y. Ng, Z. W. Lim, E. Bezault, J. Turner-Maier, J. Johnson, R. Alcazar, H. J. Noh, P. Russell, B. Aken, J. Alföldi, C. Amemiya, N. Azzouzi, J. F. Baroiller, F. Barloy-Hubler, A. Berlin, R. Bloomquist, K. L. Carleton, M. A. Conte, H. D'Cotta, O. Eshel, L. Gaffney, F. Galibert, H. F. Gante, S. Gnerre, L. Greuter, R. Guyon, N. S. Haddad, W. Haerty, R. M. Harris, H. A. Hofmann, T. Hourlier, G. Hulata, D. B. Jaffe, M. Lara,

A. P. Lee, I. MacCallum, S. Mwaiko, M. Nikaido, H. Nishihara, C. Ozouf-Costaz, D. J. Penman, D. Przybylski, M. Rakotomanga, S. C. Renn, F. J. Ribeiro, M. Ron, W. Salzburger, L. Sanchez-Pulido, M. E. Santos, S. Searle, T. Sharpe, R. Swofford, F. J. Tan, L. Williams, S. Young, S. Yin, N. Okada, T. D. Kocher, E. A. Miska, E. S. Lander, B. Venkatesh, R. D. Fernald, A. Meyer, C. P. Ponting, J. T. Streelman, K. Lindblad-Toh, O. Seehausen, and F. Di Palma. 2015. The genomic substrate for adaptive radiation in African cichlid fish. Nature 513:375–381.

Brock, C. D., L. J. Harmon, and M. E. Alfaro. 2011. Testing for temporal variation in diversification rates when sampling is incomplete and nonrandom. Systematic Biology 60.

Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait. 2011. Stacks: Building and Genotyping Loci De Novo From Short-Read Sequences. G3: Genes, Genomes, Genetics 1:171–182.

Chan, K. O., C. R. Hutter, P. L. Wood, L. L. Grismer, and R. M. Brown. 2020. Larger, unfiltered datasets are more effective at resolving phylogenetic conflict: Introns, exons, and UCEs resolve ambiguities in Golden-backed frogs (Anura: Ranidae; genus Hylarana). Molecular Phylogenetics and Evolution 151.

Cloutier, A., T. B. Sackton, P. Grayson, M. Clamp, A. J. Baker, and S. V. Edwards. 2019. Whole-Genome Analyses Resolve the Phylogeny of Flightless Birds (Palaeognathae) in the Presence of an Empirical Anomaly Zone. Systematic Biology 68.

Colless, D. 1982. Review of Phylogenetics: the theory and practice of phylogenetic systematics. Systematic Zoology 31:100–104.

Cusimano, N. and S. S. Renner. 2010. Slowdowns in Diversification Rates from Real Phylogenies May Not be Real. Systematic Biology 59:458–464.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.

Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, and R. Durbin. 2011. The variant call format and VCFtools. Bioinformatics 27:2156–2158.

Davey, J. W., T. Cezard, P. Fuentes-Utrilla, C. Eland, K. Gharbi, and M. L. Blaxter. 2013. Special features of RAD Sequencing data: Implications for genotyping. Molecular Ecology 22:3151–3164.

DeWitt, W. S., K. D. Harris, A. P. Ragsdale, and K. Harris. 2021. Nonparametric coalescent inference of mutation spectrum history and demography. Proceedings of the National Academy of Sciences 118:e2013798118.

Eaton, D. A., E. L. Spriggs, B. Park, and M. J. Donoghue. 2017. Misconceptions on missing data in RAD-seq phylogenetics with a deep-scale example from flowering plants. Systematic Biology 66:399–412.

Edwards, S. V. 2009. Is a new and general theory of molecular systematics emerging? Evolution 63:1–19.

Estabrook, G. F., F. R. McMorris, and C. A. Meacham. 1985. Comparison of Undirected Phylogenetic Trees Based on Subtrees of Four Evolutionary Units. Systematic Biology 34.

Feng, S., J. Stiller, Y. Deng, J. Armstrong, Q. Fang, A. H. Reeve, D. Xie, G. Chen, C. Guo, B. C. Faircloth, B. Petersen, Z. Wang, Q. Zhou, M. Diekhans, W. Chen, S. Andreu-Sánchez, A. Margaryan, J. T. Howard, C. Parent, G. Pacheco, M.-H. S. Sinding, L. Puetz, E. Cavill, M. Ribeiro, L. Eckhart, J. Fjeldså, P. A. Hosner, R. T. Brumfield, L. Christidis, M. F. Bertelsen, T. Sicheritz-Ponten, D. T. Tietze, B. C. Robertson, G. Song, G. Borgia, S. Claramunt, I. J. Lovette, S. J. Cowen, P. Njoroge, J. P. Dumbacher, O. A. Ryder, J. Fuchs, M. Bunce, D. W. Burt, J. Cracraft, G. Meng, S. J. Hackett, P. G. Ryan, K. A. Jønsson, I. G. Jamieson, R. R. da Fonseca, E. L. Braun, P. Houde, S. Mirarab, A. Suh, B. Hansson, S. Ponnikas, H. Sigeman,

M. Stervander, P. B. Frandsen, H. van der Zwan, R. van der Sluis, C. Visser, C. N. Balakrishnan, A. G. Clark, J. W. Fitzpatrick, R. Bowman, N. Chen, A. Cloutier, T. B. Sackton, S. V. Edwards, D. J. Foote, S. B. Shakya, F. H. Sheldon, A. Vignal, A. E. R. Soares, B. Shapiro, J. González-Solís, J. Ferrer-Obiol, J. Rozas, M. Riutort, A. Tigano, V. Friesen, L. Dalén, A. O. Urrutia, T. Székely, Y. Liu, M. G. Campana, A. Corvelo, R. C. Fleischer, K. M. Rutherford, N. J. Gemmell, N. Dussex, H. Mouritsen, N. Thiele, K. Delmore, M. Liedvogel, A. Franke, M. P. Hoeppner, O. Krone, A. M. Fudickar, B. Milá, E. D. Ketterson, A. E. Fidler, G. Friis, M. Parody-Merino, P. F. Battley, M. P. Cox, N. C. B. Lima, F. Prosdocimi, T. L. Parchman, B. A. Schlinger, B. A. Loiselle, J. G. Blake, H. C. Lim, L. B. Day, M. J. Fuxjager, M. W. Baldwin, M. J. Braun, M. Wirthlin, R. B. Dikow, T. B. Ryder, G. Camenisch, L. F. Keller, J. M. DaCosta, M. E. Hauber, M. I. M. Louder, C. C. Witt, J. A. McGuire, J. Mudge, L. C. Megna, M. D. Carling, B. Wang, S. A. Taylor, G. Del-Rio, A. Aleixo, A. T. R. Vasconcelos, C. V. Mello, J. T. Weir, D. Haussler, Q. Li, H. Yang, J. Wang, F. Lei, C. Rahbek, M. T. P. Gilbert, G. R. Graves, E. D. Jarvis, B. Paten, and G. Zhang. 2020. Dense sampling of bird diversity increases power of comparative genomics. Nature 587:252–257.

Formenti, G., K. Theissinger, C. Fernandes, I. Bista, A. Bombarely, C. Bleidorn, C. Ciofi, A. Crottini, J. A. Godoy, J. Höglund, J. Malukiewicz, A. Mouton, R. A. Oomen, S. Paez, P. J. Palsbøll, C. Pampoulie, M. J. Ruiz-López, H. Svardal, C. Theofanopoulou, J. de Vries, A.-M. Waldvogel, G. Zhang, C. J. Mazzoni, E. D. Jarvis, M. Bálint, G. Formenti, K. Theissinger, C. Fernandes, I. Bista, A. Bombarely, C. Bleidorn, F. Čiampor, C. Ciofi, A. Crottini, J. A. Godoy, J. Hoglund, J. Malukiewicz, A. Mouton, R. A. Oomen, S. Paez, P. Palsbøll, C. Pampoulie, M. J. Ruiz-López, H. Svardal, C. Theofanopoulou, J. de Vries, A.-M. Waldvogel, G. Zhang, C. J. Mazzoni, E. Jarvis, M. Bálint, S. A. Aghayan, T. S. Alioto, I. Almudi, N. Alvarez, P. C. Alves, I. R. Amorim, A. Antunes, P. Arribas, P. Baldrian, P. R. Berg, G. Bertorelle, A. Böhne, A. Bonisoli-Alquati, L. L. Boštjančić, B. Boussau, C. M. Breton, E. Buzan, P. F.

Campos, C. Carreras, L. F. Castro, L. J. Chueca, E. Conti, R. Cook-Deegan, D. Croll, M. V. Cunha, F. Delsuc, A. B. Dennis, D. Dimitrov, R. Faria, A. Favre, O. D. Fedrigo, R. Fernández, G. F. Ficetola, J.-F. Flot, T. Gabaldón, D. R. Galea Agius, G. R. Gallo, A. M. Giani, M. T. P. Gilbert, T. Grebenc, K. Guschanski, R. Guyot, B. Hausdorf, O. Hawlitschek, P. D. Heintzman, B. Heinze, M. Hiller, M. Husemann, A. Iannucci, I. Irisarri, K. S. Jakobsen, S. Jentoft, P. Klinga, A. Kloch, C. F. Kratochwil, H. Kusche, K. K. Layton, J. A. Leonard, E. Lerat, G. Liti, T. Manousaki, T. Marques-Bonet, P. Matos-Maraví, M. Matschiner, F. Maumus, A. M. Mc Cartney, S. Meiri, J. Melo-Ferreira, X. Mengual, M. T. Monaghan, M. Montagna, R. W. Mysłajek, M. T. Neiber, V. Nicolas, M. Novo, P. Ozretić, F. Palero, L. Pârvulescu, M. Pascual, O. S. Paulo, M. Pavlek, C. Pegueroles, L. Pellissier, G. Pesole, C. R. Primmer, A. Riesgo, L. Rüber, D. Rubolini, D. Salvi, O. Seehausen, M. Seidel, S. Secomandi, B. Studer, S. Theodoridis, M. Thines, L. Urban, A. Vasemägi, A. Vella, N. Vella, S. C. Vernes, C. Vernesi, D. R. Vieites, R. M. Waterhouse, C. W. Wheat, G. Wörheide, Y. Wurm, and G. Zammit. 2022. The era of reference genomes in conservation genomics. Trends in Ecology & Evolution .

Grievink, L. S., D. Penny, and B. R. Holland. 2013. Missing data and influential sites: Choice of sites for phylogenetic analysis can be as important as taxon sampling and model choice. Genome Biology and Evolution 5.

Günther, T. and C. Nettelblad. 2019. The presence and impact of reference bias on population genomic studies of prehistoric human populations. PLOS Genetics 15.

Gutenkunst, R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante. 2009. Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. PLoS Genetics 5.

Heard, S. B. and A. Mooers. 1996. Imperfect information and the balance of cladograms and phenograms. Systematic Biology 45:115–118.

Heath, T. A., S. M. Hedtke, and D. M. Hillis. 2008a. Taxon sampling and the accuracy of phylogenetic analyses. Journal of Systematics and Evolution 46.

Heath, T. A., D. J. Zwickl, J. Kim, and D. M. Hillis. 2008b. Taxon Sampling Affects Inferences of Macroevolutionary Processes from Phylogenetic Trees. Systematic Biology 57:160–166.

Hohna, S., T. Stadler, F. Ronquist, and T. Britton. 2011. Inferring Speciation and Extinction Rates under Different Sampling Schemes. Molecular Biology and Evolution 28:2577–2589.

Hotaling, S., J. L. Kelley, and P. B. Frandsen. 2021. Toward a genome sequence for every animal: Where are we now? Proceedings of the National Academy of Sciences 118:e2109019118.

Hotaling, S., C. C. Muhlfeld, J. J. Giersch, O. A. Ali, S. Jordan, M. R. Miller, G. Luikart, and D. W. Weisrock. 2018. Demographic modelling reveals a history of divergence with gene flow for a glacially tied stonefly in a changing post-Pleistocene landscape. Journal of Biogeography 45:304–317.

Huang, H. and L. L. Knowles. 2016. Unforeseen consequences of excluding missing data from next-generation sequences: Simulation study of rad sequences. Systematic Biology 65:357–365.

Huang, J. P., J. V. G. Hill, J. Ortego, and L. L. Knowles. 2020. Paraphyletic species no more – genomic data resolve a Pleistocene radiation and validate morphological species of the Melanoplus scudderi complex (Insecta: Orthoptera). Systematic Entomology 45.

Huang, S., J. Holt, C.-Y. Kao, L. McMillan, and W. Wang. 2014. A novel multi-alignment pipeline for high-throughput sequencing data. Database 2014:bau057–bau057.

Huang, W., L. Li, J. R. Myers, and G. T. Marth. 2012. ART: A next-generation sequencing read simulator. Bioinformatics 28:593–594.

Huelsenbeck, J. P. and M. Kirkpatrick. 1996. Do phylogenetic methods produce trees with biased shapes? Evolution 50.

Irisarri, I., P. Singh, S. Koblmüller, J. Torres-Dowdall, F. Henning, P. Franchini, C. Fischer, A. R. Lemmon, E. M. Lemmon, G. G. Thallinger, C. Sturmbauer, and A. Meyer. 2018. Phylogenomics uncovers early hybridization and adaptive loci shaping the radiation of Lake Tanganyika cichlid fishes. Nature Communications 9.

Jeffroy, O., H. Brinkmann, F. Delsuc, and H. Philippe. 2006. Phylogenomics: the beginning of incongruence? Trends in Genetics 22:225–231.

Koblmüller, S., B. Egger, C. Sturmbauer, and K. M. Sefc. 2010. Rapid radiation, ancient incomplete lineage sorting and ancient hybridization in the endemic Lake Tanganyika cichlid tribe Tropheini. Molecular Phylogenetics and Evolution 55.

Koblmüller, S., C. A. Schöggl, C. J. Lorber, M. Van Steenberge, N. Kmentová, M. P. Vanhove, and L. Zangl. 2021. African lates perches (Teleostei, Latidae, Lates): Paraphyly of Nile perch and recent colonization of Lake Tanganyika. Molecular Phylogenetics and Evolution 160.

Kolora, S. R. R., G. L. Owens, J. M. Vazquez, A. Stubbs, K. Chatla, C. Jainese, K. Seeto, M. McCrea, M. W. Sandel, J. A. Vianna, K. Maslenikov, D. Bachtrog, J. W. Orr, M. Love, and P. H. Sudmant. 2021. Origins and evolution of extreme life span in Pacific Ocean rockfishes. Science 374:842–847.

Kubo, T. and Y. Iwasa. 1995. Inferring the rates of branching and extinction from molecular phylogenies. Evolution 49.

Kumar, S., A. J. Filipski, F. U. Battistuzzi, S. L. Kosakovsky Pond, and K. Tamura. 2012. Statistics and truth in phylogenomics. Molecular Biology and Evolution 29:457–472.

Lanave, C., G. Preparata, C. Sacone, and G. Serio. 1984. A new method for calculating evolutionary substitution rates. Journal of Molecular Evolution 20.

Lanier, H. C., H. Huang, and L. L. Knowles. 2014. How low can you go? The effects of mutation rate on the accuracy of species-tree estimation. Molecular Phylogenetics and Evolution 70:112–119.

Leaché, A. D., B. L. Banbury, J. Felsenstein, A. N. M. De Oca, and A. Stamatakis. 2015. Short tree, long tree, right tree, wrong tree: New acquisition bias corrections for inferring SNP phylogenies. Systematic Biology 64:1032–1047.

Lemmon, A. R., J. M. Brown, K. Stanger-Hall, and E. M. Lemmon. 2009. The effect of ambiguous data on phylogenetic estimates obtained by maximum likelihood and bayesian inference. Systematic Biology 58.

Lemmon, E. M. and A. R. Lemmon. 2013. High-Throughput Genomic Data in Systematics and Phylogenetics. Annual Review of Ecology, Evolution, and Systematics 44:99–121.

Li, H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics 27:2987–2993.

Li, H. and R. Durbin. 2009. Fast and accurate short-read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, and R. Durbin. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078–2079.

Linck, E. and C. J. Battey. 2019. Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. Molecular Ecology Resources 19:639–647.

Louca, S. and M. W. Pennell. 2020. Extant timetrees are consistent with a myriad of diversification histories. Nature 580:502–505.

Louca, S. and M. W. Pennell. 2021. Why extinction estimates from extant phylogenies are
so often zero. Current Biology 31:3168–3173.

Malinsky, M., H. Svardal, A. M. Tyers, E. A. Miska, M. J. Genner, G. F. Turner, and
R. Durbin. 2018. Whole-genome sequences of Malawi cichlids reveal multiple radiations
interconnected by gene flow. Nature Ecology and Evolution 2:1940–1955.

Mallo, D., L. De Oliveira Martins, and D. Posada. 2016. SimPhy: Phylogenomic
Simulation of Gene, Locus, and Species Trees. Systematic Biology 65:334–344.

Mastretta-Yanes, A., N. Arrigo, N. Alvarez, T. H. Jorgensen, D. Piñero, and B. C.
Emerson. 2015. Restriction site-associated DNA sequencing, genotyping error estimation
and de novo assembly optimization for population genetic inference. Molecular Ecology
Resources 15:28–41.

McCormack, J. E., H. Huang, and L. L. Knowles. 2009. Maximum likelihood estimates of
species trees: How accuracy of phylogenetic inference depends upon the divergence
history and sampling design. Systematic Biology 58:501–508.

McTavish, E. J. and D. M. Hillis. 2015. How do SNP ascertainment schemes and
population demographics affect inferences about population history? BMC Genomics
16:1–14.

McTavish, E. J., J. Pettengill, S. Davis, H. Rand, E. Strain, M. Allard, and R. E. Timme.
2017. TreeToReads - A pipeline for simulating raw reads from phylogenies. BMC
Bioinformatics 18:1–7.

Mirarab, S. and T. Warnow. 2015. ASTRAL-II: Coalescent-based species tree estimation
with many hundreds of taxa and thousands of genes. Bioinformatics 31:i44–i52.

Molloy, E. K. and T. Warnow. 2018. To Include or Not to Include: The Impact of Gene
Filtering on Species Tree Estimation Methods. Systematic Biology 67:285–303.

Momozawa, Y. and K. Mizukami. 2021. Unique roles of rare variants in the genetics of complex diseases in humans. Journal of Human Genetics 66:11–23.

Nazareno, A. G. and L. L. Knowles. 2021. There Is No 'Rule of Thumb': Genomic Filter Settings for a Small Plant Population to Obtain Unbiased Gene Flow Estimates. Frontiers in Plant Science 12.

Nei, M. and W. H. Li. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. Proceedings of the National Academy of Sciences of the United States of America 76:5269–5273.

Nelson, M. R., D. Wegmann, M. G. Ehm, D. Kessner, P. St. Jean, C. Verzilli, J. Shen, Z. Tang, S.-A. Bacanu, D. Fraser, L. Warren, J. Aponte, M. Zawistowski, X. Liu, H. Zhang, Y. Zhang, J. Li, Y. Li, L. Li, P. Woollard, S. Topp, M. D. Hall, K. Nangle, J. Wang, G. Abecasis, L. R. Cardon, S. Zöllner, J. C. Whittaker, S. L. Chissoe, J. Novembre, and V. Mooser. 2012. An Abundance of Rare Functional Variants in 202 Drug Target Genes Sequenced in 14,002 People. Science 337:100–104.

Nielsen, R. 2000. Estimation of population parameters and recombination rates from single nucleotide polymorphisms. Genetics 154:931–942.

Nielsen, R., J. S. Paul, A. Albrechtsen, and Y. S. Song. 2011. Genotype and SNP calling from next-generation sequencing data. Nature Reviews Genetics 12:443–451.

Novembre, J., T. Johnson, K. Bryc, Z. Kutalik, A. R. Boyko, A. Auton, A. Indap, K. S. King, S. Bergmann, M. R. Nelson, M. Stephens, and C. D. Bustamante. 2008. Genes mirror geography within Europe. Nature 456:98–101.

O'Hara, R. J. 1997. Population thinking and tree thinking in systematics. Zoologica Scripta 26.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Paradis, E. and K. Schliep. 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics 35:526–528.

Parchman, T. L., Z. Gompert, J. Mudge, F. D. Schilkey, C. W. Benkman, and C. A. Buerkle. 2012. Genome-wide association genetics of an adaptive trait in lodgepole pine. Molecular Ecology 21:2991–3005.

Patton, A. H., M. J. Margres, A. R. Stahlke, S. Hendricks, K. Lewallen, R. K. Hamede, M. Ruiz-Aravena, O. Ryder, H. I. McCallum, M. E. Jones, P. A. Hohenlohe, and A. Storfer. 2019. Contemporary Demographic Reconstruction Methods Are Robust to Genome Assembly Quality: A Case Study in Tasmanian Devils. Molecular Biology and Evolution 36:2906–2921.

Philippe, H., E. A. Snell, E. Bapteste, P. Lopez, P. W. Holland, and D. Casane. 2004. Phylogenomics of eukaryotes: Impact of missing data on large alignments. Molecular Biology and Evolution 21:1740–1752.

Prasad, A., E. D. Lorenzen, and M. V. Westbury. 2021. Evaluating the role of reference-genome phylogenetic distance on evolutionary inference. Molecular Ecology Resources 22:45–55.

Prasad, A. B., M. W. Allard, and E. D. Green. 2008. Confirming the phylogeny of mammals by use of large comparative sequence data sets. Molecular Biology and Evolution 25:1795–1808.

Pybus, O. G. and P. H. Harvey. 2000. Testing macro–evolutionary models using incomplete molecular phylogenies. Proceedings of the Royal Society of London. Series B: Biological Sciences 267.

R Core Team. 2021. R: A language and Environment for Statistical Computing.

Rambaut, A. and N. C. Grass. 1997. Seq-Gen: an application for the Monte Carlo simulation of DNA sequence evolution along phylogenetic trees. Bioinformatics 13.

Rannala, B., J. P. Huelsenbeck, Z. Yang, and R. Nielsen. 1998. Taxon sampling and the accuracy of large phylogenies. Systematic Biology 47:702–710.

Reid, B. N., R. L. Moran, C. J. Kopack, and S. W. Fitzpatrick. 2021. Rapture-ready darters: Choice of reference genome and genotyping method (whole-genome or sequence capture) influence population genomic inference in ¡i¿Etheostoma¡/i¿. Molecular Ecology Resources 21.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other things). Methods in Ecology and Evolution 3:217–223.

Revell, L. J., L. J. Harmon, and R. E. Glor. 2005. Underparameterized model of sequence evolution leads to bias in the estimation of diversification rates from molecular phylogenies. Systematic biology 54.

Rhie, A., S. A. McCarthy, O. Fedrigo, J. Damas, G. Formenti, S. Koren, M. Uliano-Silva, W. Chow, A. Fungtammasan, J. Kim, C. Lee, B. J. Ko, M. Chaisson, G. L. Gedman, L. J. Cantin, F. Thibaud-Nissen, L. Haggerty, I. Bista, M. Smith, B. Haase, J. Mountcastle, S. Winkler, S. Paez, J. Howard, S. C. Vernes, T. M. Lama, F. Grutzner, W. C. Warren, C. N. Balakrishnan, D. Burt, J. M. George, M. T. Biegler, D. Iorns, A. Digby, D. Eason, B. Robertson, T. Edwards, M. Wilkinson, G. Turner, A. Meyer, A. F. Kautt, P. Franchini, H. W. Detrich, H. Svardal, M. Wagner, G. J. P. Naylor, M. Pippel, M. Malinsky, M. Mooney, M. Simbirsky, B. T. Hannigan, T. Pesout, M. Houck, A. Misuraca, S. B. Kingan, R. Hall, Z. Kronenberg, I. Sović, C. Dunn, Z. Ning, A. Hastie, J. Lee, S. Selvaraj, R. E. Green, N. H. Putnam, I. Gut, J. Ghurye, E. Garrison, Y. Sims, J. Collins, S. Pelan, J. Torrance, A. Tracey, J. Wood, R. E. Dagnew, D. Guan, S. E. London, D. F. Clayton, C. V. Mello, S. R. Friedrich, P. V. Lovell, E. Osipova, F. O. Al-Ajli, S. Secomandi, H. Kim, C. Theofanopoulou, M. Hiller, Y. Zhou, R. S. Harris, K. D. Makova, P. Medvedev, J. Hoffman, P. Masterson, K. Clark, F. Martin, K. Howe, P. Flicek, B. P. Walenz, W. Kwak, H. Clawson, M. Diekhans,

L. Nassar, B. Paten, R. H. S. Kraus, A. J. Crawford, M. T. P. Gilbert, G. Zhang, B. Venkatesh, R. W. Murphy, K.-P. Koepfli, B. Shapiro, W. E. Johnson, F. Di Palma, T. Marques-Bonet, E. C. Teeling, T. Warnow, J. M. Graves, O. A. Ryder, D. Haussler, S. J. O'Brien, J. Korlach, H. A. Lewin, K. Howe, E. W. Myers, R. Durbin, A. M. Phillippy, and E. D. Jarvis. 2021. Towards complete and error-free genome assemblies of all vertebrate species. Nature 592:737–746.

Rick, J. A., J. Junker, I. A. Kimirei, E. A. Sweke, J. B. Mosille, C. Dinkel, S. Mwaiko, O. Seehausen, and C. E. Wagner. 2022. The genetic population structure of Lake Tanganyika's Lates species flock, an endemic radiation of pelagic top predators. Journal of Heredity .

Robinson, D. and L. Foulds. 1981. Comparison of phylogenetic trees. Mathematical Biosciences 53:131–147.

Rokas, A., B. I. Williams, N. King, and S. B. Carroll. 2003. Genome-scale approaches to resolving incongruence in molecular phylogenies. Nature 425:798–804.

Ronco, F., H. H. Büscher, A. Indermaur, and W. Salzburger. 2020. The taxonomic diversity of the cichlid fish fauna of ancient Lake Tanganyika, East Africa. Journal of Great Lakes Research 46:1067–1078.

Ronco, F., M. Matschiner, A. Böhne, A. Boila, H. H. Büscher, A. El Taher, A. Indermaur, M. Malinsky, V. Ricci, A. Kahmen, S. Jentoft, and W. Salzburger. 2021. Drivers and dynamics of a massive adaptive radiation in cichlid fishes. Nature 589.

Ros-Freixedes, R., M. Battagin, M. Johnsson, G. Gorjanc, A. J. Mileham, S. D. Rounsley, and J. M. Hickey. 2018. Impact of index hopping and bias towards the reference allele on accuracy of genotype calls from low-coverage sequencing. Genetics Selection Evolution 50.

58   REFERENCES

Rüber, L. and R. Zardoya. 2005. Rapid cladogenesis in marine fishes revisited. Evolution 59.

Sackin, M. 1972. "Good" and "Bad" Phenograms. Systematic Zoology 21:225–226.

Sand, A., M. K. Holt, J. Johansen, G. S. Brodal, T. Mailund, and C. N. Pedersen. 2014. TqDist: A library for computing the quartet and triplet distances between binary or general trees. Bioinformatics 30.

Sarver, B. A., S. Keeble, T. Cosart, P. K. Tucker, M. D. Dean, and J. M. Good. 2017. Phylogenomic Insights into Mouse Evolution Using a Pseudoreference Approach. Genome Biology and Evolution 9:726–739.

Shafer, A. B., C. R. Peart, S. Tusso, I. Maayan, A. Brelsford, C. W. Wheat, and J. B. Wolf. 2017. Bioinformatic processing of RAD-seq data dramatically impacts downstream population genetic inference. Methods in Ecology and Evolution 8:907–917.

Shen, X. X., C. T. Hittinger, and A. Rokas. 2017. Contentious relationships in phylogenomic studies can be driven by a handful of genes. Nature Ecology and Evolution 1:1–10.

Slatkin, M. 1985. Rare alleles as indicators of gene flow. Evolution 39:53–65.

Smith, M. R. 2019. Bayesian and parsimony approaches reconstruct informative trees from simulated morphological datasets. Biology Letters 15.

Smith, M. R. 2020. Information theoretic generalized Robinson–Foulds metrics for comparing phylogenetic trees. Bioinformatics 36.

Stadler, T., J. H. Degnan, and N. A. Rosenberg. 2016. Does Gene Tree Discordance Explain the Mismatch between Macroevolutionary Models and Empirical Patterns of Tree Shape and Branching Times? Systematic Biology 65.

Stamatakis, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312–1313.

Vij, S., H. Kuhl, I. S. Kuznetsova, A. Komissarov, A. A. Yurchenko, P. Van Heusden, S. Singh, N. M. Thevasagayam, S. R. S. Prakki, K. Purushothaman, J. M. Saju, J. Jiang, S. K. Mbandi, M. Jonas, A. Hin Yan Tong, S. Mwangi, D. Lau, S. Y. Ngoh, W. C. Liew, X. Shen, L. S. Hon, J. P. Drake, M. Boitano, R. Hall, C. S. Chin, R. Lachumanan, J. Korlach, V. Trifonov, M. Kabilov, A. Tupikin, D. Green, S. Moxon, T. Garvin, F. J. Sedlazeck, G. W. Vurture, G. Gopalapillai, V. Kumar Katneni, T. H. Noble, V. Scaria, S. Sivasubbu, D. R. Jerry, S. J. O'Brien, M. C. Schatz, T. Dalmay, S. W. Turner, S. Lok, A. Christoffels, and L. Orbán. 2016. Chromosomal-Level Assembly of the Asian Seabass Genome Using Long Sequence Reads and Multi-layered Scaffolding. PLoS Genetics 12:1–35.

Wagner, C. E., I. Keller, S. Wittwer, O. M. Selz, S. Mwaiko, L. Greuter, A. Sivasundar, and O. Seehausen. 2013. Genome-wide RAD sequence data provide unprecedented resolution of species boundaries and relationships in the Lake Victoria cichlid adaptive radiation. Molecular Ecology 22:787–798.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wiens, J. J. 1998. Does adding characters with missing data increase or decrease phylogenetic accuracy? Systematic Biology 47.

Wiens, J. J. 2003. Missing data, incomplete taxa, and phylogenetic accuracy. Systematic Biology 52:528–538.

Wiens, J. J. and D. S. Moen. 2008. Missing data and the accuracy of Bayesian phylogenetics. Journal of Systematics and Evolution 46:307–314.

Wiens, J. J. and J. Tiu. 2012. Highly incomplete taxa can rescue phylogenetic analyses from the negative impacts of limited taxon sampling. PLoS ONE 7:e42925.

1361 Yang, Z. 1993. Maximum-likelihood estimation of phylogeny from DNA sequences when

1362 substitution rates differ over sites. Molecular Biology and Evolution 10:1396–1401.

1363 Zwickl, D. J. and D. M. Hillis. 2002. Increased Taxon Sampling Greatly Reduces

1364 Phylogenetic Error. Systematic Biology 51:588–598.

APPENDIX

Supplementary Material

Supplemental methods

Reference genome for simulations.—   To create a reference genome to use as a base for our `TreeToReads` simulations, we started with the *Lates calcarifer* reference genome (Vij et al., 2016), as this was the most complete of the four reference genomes used in our empirical analyses. We randomly sampled 5,000bp windows without replacement using a custom bash script. For each random segment, we replaced any ambiguous bases (N) with a random base (A, C, G, or T), with equal weighting given to each option.

Genotyping-by-sequencing for Lake Tanganyika Tropheus cichlids.—   We sampled individuals in the cichlid tribe Tropheini in the Kigoma region of Tanzania in 2002, 2005, 2007, 2012, 2015, 2016, 2017, and 2018. Briefly, we collected individuals using gill nets while snorkeling in the rocky littoral zone at a 1–10 m depth. We took a fin clip from each individual for genetic analyses, which was stored in either DMSO-EDTA or 95% ethanol prior to DNA extraction. We retained all individuals as vouchers, which are deposited at the Cornell University Museum of Vertebrates (2002, 2005, and 2007 individuals) and the University of Wyoming Museum of Vertebrates (2015, 2016, 2017, and 2018 individuals).

We extracted DNA from the fin clips using DNeasy Blood & Tissue kits (Qiagen, Inc.) and prepared genomic libraries for high-throughput DNA sequencing following the GBS protocol described in Parchman et al. (2012). Briefly, we fragmented DNA using EcoRI and MseI restriction enzymes and then ligated a unique $8-10$ base pair (bp) barcode to each individual's DNA. We used polymerase chain reaction (PCR) to amplify the restriction/ligation products (two independent replicates per individual) and then combined the PCR products to create the final libraries. Prior to sequencing, the libraries were size-selected for $250-400$ bp fragments using BluePippin (Sage Science). Sequencing was completed on Illumina HiSeq 2500 and 4000 platforms (100 bp single-end) at the

₂₇ University of Texas Genome Sequencing and Analysis Facility (Austin, Texas, USA) and

₂₈ the University of Oregon Genomics and Cell Characterization Core Facility (Eugene,

₂₉ Oregon, USA). The individuals in this project were included in libraries containing

₃₀ samples of other cichlid species as part of a larger sequencing effort. Each library contained

₃₁ ∼100 individuals and was sequenced in its own lane.

₃₂ With the raw sequence data, we first assigned reads to individuals and subsequently

₃₃ removed the barcode sequences using a custom `perl` script. We then aligned the reads to

₃₄ the *Pundamilia nyererei* and *Oreochromis niloticus* reference genomes (Brawand et al.,

₃₅ 2015) using `bwa` v0.7.17 (v0.7.17 Li and Durbin, 2009) with default settings to produce `bam`

₃₆ alignment files, which were used for variant calling in downstream analyses.

Supplemental tables and figures

Table S1. Model coefficients from linear mixed models, including parameter estimates (Est), minimum 95% confidence interval (Min CI), upper 95% confidence interval (Max CI), and significance at $\alpha = 0.05$ (* indicates p <0.05). Parameters are listed for models for all trees, as well as the three ILS levels individually. Results are summarized graphically in Fig. 3). Predictor variables in each model are average distance from the ingroup taxa to the reference genome (avg_dxy), minor allele count threshold (maf), missing data threshold (missing), and the interaction among these terms, with simulation number as a random effect.

Variable	All Trees				Low ILS				Medium ILS				High ILS			
	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig
Standardized ingroup gamma																
avg_dxy	1.242	-0.103	2.590		0.032	-1.065	1.136		0.286	-1.087	1.664		0.094	-1.175	1.366	
maf	0.840	0.782	0.898	*	0.630	0.577	0.683	*	0.942	0.885	0.999	*	0.856	0.804	0.908	*
missing	-0.395	-0.941	0.152		-0.408	-0.905	0.088		-0.416	-0.954	0.121		-0.364	-0.856	0.127	
avg_dxy:maf	-0.061	-0.267	0.146		-0.018	-0.187	0.151		-0.022	-0.233	0.189		0.007	-0.188	0.202	
avg_dxy:missing	-0.096	-2.034	1.843		-0.076	-1.660	1.508		-0.097	-2.074	1.880		-0.106	-1.933	1.721	
Robinson-Foulds distance to true tree																
avg_dxy	-0.009	-0.029	0.011		0.006	-0.021	0.033		0.003	-0.024	0.029		-0.018	-0.051	0.014	
maf	0.013	0.012	0.013	*	0.015	0.014	0.017	*	0.012	0.011	0.013	*	0.011	0.010	0.013	*
missing	0.002	-0.007	0.010		0.001	-0.011	0.013		0.002	-0.009	0.012		0.001	-0.012	0.013	
avg_dxy:maf	0.003	0.000	0.006	*	0.000	-0.004	0.004		0.002	-0.002	0.006		0.006	0.001	0.011	*
avg_dxy:missing	-0.004	-0.033	0.025		-0.017	-0.056	0.022		-0.012	-0.050	0.027		0.020	-0.026	0.067	
Standardized ingroup Colless imbalance																
avg_dxy	0.017	0.001	0.033	*	0.006	-0.014	0.026		0.010	-0.019	0.038		0.025	0.000	0.049	
maf	0.009	0.009	0.010	*	0.007	0.006	0.008	*	0.009	0.008	0.011	*	0.011	0.010	0.012	*
missing	-0.004	-0.011	0.002		-0.001	-0.010	0.008		-0.005	-0.016	0.006		-0.006	-0.016	0.003	
avg_dxy:maf	-0.002	-0.004	0.001		0.001	-0.002	0.004		-0.001	-0.005	0.004		-0.004	-0.008	-0.001	*
avg_dxy:missing	-0.001	-0.024	0.023		-0.009	-0.038	0.020		0.002	-0.039	0.043		0.004	-0.031	0.040	

Table S2. Model coefficients from linear mixed models for subsampled data sets, including parameter estimates (Est), minimum 95% confidence interval (Min CI), upper 95% confidence interval (Max CI), and significance at α = 0.05 (* indicates $p < 0.05$). Parameters are listed for models for all trees, as well as the three ILS levels individually. Results are summarized graphically in Fig. 5). Predictor variables in each model are average distance from the ingroup taxa to the reference genome (avg_dxy), minor allele count threshold (maf), missing data threshold (missing), and the interaction among these terms, with simulation number as a random effect.

Variable	All Trees				Low ILS				Medium ILS				High ILS			
	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig
Standardized ingroup gamma																
avg_dxy	0.729	0.236	1.222	*	0.029	-0.435	0.493		0.313	-0.425	1.051		-0.074	-0.699	0.552	
maf	0.755	0.739	0.771	*	0.254	0.238	0.271	*	0.931	0.908	0.953	*	0.862	0.843	0.881	*
missing	-0.449	-0.645	-0.253	*	-0.034	-0.238	0.171		-0.590	-0.872	-0.309	*	-0.544	-0.780	-0.308	*
avg_dxy:maf	-0.127	-0.183	-0.071	*	0.021	-0.032	0.073		-0.075	-0.158	0.009		0.024	-0.047	0.095	
avg_dxy:missing	0.115	-0.579	0.809		-0.020	-0.673	0.632		0.036	-1.001	1.074		0.044	-0.836	0.924	
Robinson-Foulds distance to true tree																
avg_dxy	-0.014	-0.030	0.001		-0.012	-0.030	0.007		0.016	-0.006	0.039		-0.005	-0.031	0.020	
maf	0.006	0.006	0.007	*	0.015	0.014	0.016	*	0.005	0.005	0.006	*	0.002	0.002	0.003	*
missing	-0.005	-0.011	0.002		-0.003	-0.012	0.005		-0.001	-0.010	0.007		-0.009	-0.018	0.001	
avg_dxy:maf	0.003	0.001	0.005	*	0.001	-0.001	0.003		0.000	-0.002	0.003		0.002	-0.001	0.005	
avg_dxy:missing	0.003	-0.019	0.025		0.005	-0.022	0.031		-0.011	-0.043	0.020		0.016	-0.021	0.052	
Standardized ingroup Colless imbalance																
avg_dxy	-0.001	-0.011	0.009		0.007	-0.005	0.020		-0.010	-0.028	0.008		-0.001	-0.019	0.016	
maf	0.009	0.009	0.010	*	0.009	0.009	0.009	*	0.010	0.010	0.011	*	0.009	0.008	0.009	*
missing	-0.005	-0.009	-0.001	*	0.002	-0.004	0.007		-0.007	-0.014	0.000		-0.008	-0.015	-0.002	*
avg_dxy:maf	0.002	0.001	0.003	*	0.000	-0.001	0.002		0.004	0.002	0.006	*	0.001	-0.001	0.003	
avg_dxy:missing	-0.005	-0.019	0.010		-0.012	-0.030	0.006		-0.008	-0.034	0.017		0.002	-0.023	0.027	

Table S3. Model coefficients from linear mixed models for the empirical Tropheine and *Lates* data sets, including parameter estimates (Est), minimum 95% confidence interval (Min CI), upper 95% confidence interval (Max CI), and significance at $\alpha = 0.05$. Parameters are listed for models for all trees, as well as the three ILS levels individually. Results are summarized graphically in Fig. 8). Predictor variables in each model are reference genome choice (ref), minor allele count threshold (mac), missing data threshold (missing), and the interaction among these terms, with iteration number as a random effect. Reference genome choice (ref) was coded as 0=outgroup reference and 1=ingroup reference.

Variable	Tropheine Topologies				Lates Topologies			
	Est	Min CI	Max CI	Sig	Est	Min CI	Max CI	Sig
Ingroup gamma								
ref	0.111	-0.037	0.259		0.030	-0.390	0.450	
mac	0.262	0.246	0.278	*	-0.254	-0.299	-0.208	*
missing	-1.204	-1.357	-1.051	*	-0.345	-0.773	0.082	
ref:mac	-0.025	-0.048	-0.002	*	0.021	-0.043	0.086	
ref:missing	0.420	0.207	0.634	*	0.161	-0.444	0.766	
Ingroup tree height								
ref	0.013	-0.007	0.034		-0.006	-0.034	0.022	
mac	0.022	0.019	0.024	*	-0.032	-0.035	-0.029	*
missing	-0.018	-0.039	0.004		-0.062	-0.091	-0.033	*
ref:mac	-0.020	-0.024	-0.017	*	0.000	-0.005	0.004	
ref:missing	0.032	0.002	0.062	*	-0.008	-0.049	0.032	
Ingroup Colless imbalance								
ref	0.020	-0.025	0.064		-0.011	-0.031	0.009	
mac	-0.011	-0.016	-0.006	*	-0.001	-0.003	0.001	
missing	-0.224	-0.270	-0.178	*	-0.009	-0.029	0.012	
ref:mac	0.002	-0.005	0.009		0.002	-0.001	0.005	
ref:missing	0.006	-0.059	0.070		0.002	-0.027	0.031	

Fig. S1. Examples of inferred trees for a single simulation for high, medium, and low ILS species trees, where data were aligned to an ingroup (left) versus outgroup (right) reference genome. Trees are shown for no missing data cutoff and a minor allele count cutoff of 1, 3, 5, and 10, to demonstrate the changes in the trees that are summarized through the γ and imbalance statistics.

Fig. S2. Relationship within the full simulation data sets between (A) minor allele count threshold and the number of sites retained in a data set, and (B) missing data threshold and the number of sites retained in a data set, with colors indicating the tree height.

Fig. S3. Correlation matrix between output tree characteristics for the full simulation data sets. Color and size correspond to the magnitude of the correlation. Variables are overall tree height (tree.height), standardized Sackin imbalance of the ingroup (std.ingroup.sackin), standardized γ of the ingroup (std.ingroup.gamma), standardized Colless imbalance of the ingroup (std.ingroup.colless), Robinson Foulds distance to the true tree (RF.Dist.ML), quartet distance to the true tree (Q.Dist.ML), mean bootstrap support of all nodes in the tree (mean.support), height of the ingroup (ingroup.tree.height), and clustering information distance to the true tree (CI.Dist.ML).

Fig. S4. Heatmaps for the **low ILS trees** indicating how each of our tree descriptive parameters varies with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S5. Heatmaps for the **medium ILS trees** indicating how each of our tree descriptive parameters varies in two-dimensional space with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S6. Heatmaps for the **high ILS trees** indicating how each of our tree descriptive parameters varies in two-dimensional space with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S7. Heatmaps for the **subsampled low ILS trees** indicating how each of our tree descriptive parameters varies with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S8. Heatmaps for the **subsampled medium ILS trees** indicating how each of our tree descriptive parameters varies in two-dimensional space with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S9. Heatmaps for the **subsampled high ILS trees** indicating how each of our tree descriptive parameters varies in two-dimensional space with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. Three of the parameters, Robinson-Foulds (RF) distance, quartet distance, and clustering information (CI) distance, are measures of topological dissimilarity compared to the true simulated tree, and lower values indicate topologies closer to the true tree. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, with means for the true trees around I_c=0.4 and γ=0.

Fig. S10. Heatmaps for the **empirical Tropheine trees** indicating how each of our tree descriptive parameters varies in two-dimensional space with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, respectively.

Fig. S11. Heatmaps for the **empirical *Lates* trees** indicating how each of our tree descriptive parameters varies with minor allele count and missing data thresholds, with darker colors indicating lower values and lighter colors indicating higher values. The gamma and Colless imbalance statistics are measures of tree center of gravity and tree shape, respectively.

Fig. S12. Example plots of principal coordinates analysis (PCoA) visualizing distance between trees in tree space for (A) high ILS, (B), medium ILS, and (C) low ILS trees. For each plot, the true, simulated species tree is shown as a black star, while the other points are colored according to their minor allele count threshold and shape indicates the reference genome choice (circle = EXT/outgroup reference; triangle = INT/ingroup reference). Arrows on the plots show correlations of each of the three bioinformatic choices of interest with the first two PCoA axes, with the length of the arrow corresponding to the magnitude of correlation with the given bioinformatic choice.

Fig. S13. Example plots of principal coordinates analysis (PCoA) visualizing distance between trees in tree space for (a) Tropheine and (b) *Lates* trees. For each plot, the points are colored according to their minor allele count threshold and shape indicates the reference genome choice (circle = EXT/outgroup reference; diamond = INT/ingroup reference). Arrows on the plots show correlations of each of the three bioinformatic choices of interest with the first two PCoA axes, with the length of the arrow corresponding to the magnitude of correlation with the given bioinformatic choice.

Fig. S14. Patterns of variation for the full simulation data sets in the three metrics of distance to the true tree, two metrics of tree imbalance, and tree center of gravity (γ) across minor allele counts, with shapes indicating the reference used (EXT, outgroup reference; INT, ingroup reference) and colors indicating the level of ILS. For topological accuracy metrics (Robinson Foulds, quartet, and clustering information distances to the true tree), higher values indicate topologies more biased away from the true tree, and all have been normalized to be bounded by $0 \leqslant$ distance $\leqslant 1$. Measures of Colless imbalance, Sackin imbalance, and γ have been standardized so that a value of 0 is equal to the measure in the true tree.

Fig. S15. Patterns of variation for the full simulation data sets in the three metrics of distance to the true tree, two metrics of tree imbalance, and tree center of gravity (γ) across missing data thresholds, with shapes indicating the reference used (EXT, outgroup reference; INT, ingroup reference) and colors indicating the level of ILS. For topological accuracy metrics (Robinson Foulds, quartet, and clustering information distances to the true tree), higher values indicate topologies more biased away from the true tree, and all have been normalized to be bounded by $0 \leqslant$ distance $\leqslant 1$. Measures of Colless imbalance, Sackin imbalance, and γ have been standardized so that a value of 0 is equal to the measure in the true tree.

Fig. S16. Patterns of variation for the subsampled simulation data sets in the three metrics of distance to the true tree, two metrics of tree imbalance, and tree center of gravity (γ) across minor allele counts, with shapes indicating the reference used (EXT, outgroup reference; INT, ingroup reference) and colors indicating the level of ILS. For topological accuracy metrics (Robinson Foulds, quartet, and clustering information distances to the true tree), higher values indicate topologies more biased away from the true tree, and all have been normalized to be bounded by $0 \leqslant \text{distance} \leqslant 1$. Measures of Colless imbalance, Sackin imbalance, and γ have been standardized so that a value of 0 is equal to the measure in the true tree.

Fig. S17. Patterns of variation for the subsampled simulation data sets in the three metrics of distance to the true tree, two metrics of tree imbalance, and tree center of gravity (γ) across missing data thresholds, with shapes indicating the reference used (EXT, outgroup reference; INT, ingroup reference) and colors indicating the level of ILS. For topological accuracy metrics (Robinson Foulds, quartet, and clustering information distances to the true tree), higher values indicate topologies more biased away from the true tree, and all have been normalized to be bounded by $0 \leqslant$ distance $\leqslant 1$. Measures of Colless imbalance, Sackin imbalance, and γ have been standardized so that a value of 0 is equal to the measure in the true tree.

Fig. S18. Correlation matrix between output tree characteristics for the (A) Tropheine and (B) *Lates* empirical data sets. Color and size correspond to the magnitude of the correlation. Variables are ingroup tree height (ingroup.tree.height), Sackin imbalance of the ingroup (ingroup.sackin), γ of the ingroup (ingroup.gamma), Colless imbalance of the ingroup (ingroup.colless), and average branch lengths (Avg.BLs). White squares indicate pairs with non-significant Pearson's correlation coefficients at $\alpha = 0.05$.