Voltage-Sensing Phosphatase (VSP) Regulates Endocytosis-Dependent Nutrient

Absorption in Chordate Enterocytes

Adisorn Ratanayotha[1,5], Makoto Matsuda[1], Yukiko Kimura[2], Fumiko Takenaga[1], Tomoaki

Mizuno[3], Md. Israil Hossain[1], Shin-ichi Higashijima[2], Takafumi Kawai[1]*, Michio

Ogasawara[4], Yasushi Okamura[1, 6]*

[1] *Laboratory of Integrative Physiology, Department of Physiology, Graduate School of*

Medicine, Osaka University, Suita, Osaka, 565-0871, Japan

[2] *Exploratory Research Center on Life and Living Systems and National Institute for Basic*

Biology, National Institutes of Natural Sciences, Okazaki, Aichi 444-8787, Japan

[3] *Center for Medical Research and Education, Osaka University, Suita, Osaka, 565-0871,*

Japan

[4] *Department of Biology, Graduate School of Science, Chiba University, Inage-ku, Chiba,*

263-8522, Japan

[5] *Department of Anatomy, Faculty of Medicine Siriraj Hospital, Mahidol University,*

Bangkoknoi, Bangkok, 10700, Thailand

[6] *Graduate School of Frontier Biosciences, Osaka University, Suita, Osaka, 565-0871, Japan*

Manuscript contains 47 pages, 8 main figures, 9 supplemental figures, 2 supplemental videos,

2 tables.

* Correspondence:

Yasushi Okamura, MD, PhD

Laboratory of Integrative Physiology, Department of Physiology,

Graduate School of Medicine, Osaka University,

2-2 Yamada-Oka, Suita, Osaka, 565-0871, Japan

Tel: +81-6-6879-3310 Fax: +81-6-6879-3319

Email: vsop1@me.com

Takafumi Kawai, PhD

Laboratory of Integrative Physiology, Department of Physiology,

Graduate School of Medicine, Osaka University,

2-2 Yamada-Oka, Suita, Osaka, 565-0871, Japan

Tel: +81-6-6879-3311 Fax: +81-6-6879-3319

Email: kawai@phys2.med.osaka-u.ac.jp

Abstract

Voltage-sensing phosphatase (VSP) is a unique membrane protein that translates

membrane electrical activities into the changes of phosphoinositide profiles. VSP orthologs

from various species have been intensively investigated toward their biophysical properties,

primarily using a heterologous expression system. In contrast, the physiological role of VSP

in native tissues remains largely unknown. Here we report that zebrafish VSP (Dr-VSP) is

functionally expressed on the endomembranes of lysosome-rich enterocytes (LREs) that

mediate dietary protein absorption via endocytosis in the zebrafish mid-intestine. Dr-VSP-

deficient LREs were remarkably defective in forming endosomal vacuoles after initial uptake

of dextran and mCherry. Dr-VSP-deficient zebrafish exhibited growth restriction and higher

mortality during the critical period when zebrafish larvae rely primarily on exogenous

feeding via intestinal absorption. Furthermore, our comparative study on marine invertebrate

Ciona intestinalis VSP (Ci-VSP) revealed co-expression with endocytosis-associated genes

in absorptive epithelial cells of the *Ciona* digestive tract, corresponding to zebrafish LREs.

These findings signify a crucial role of VSP in regulating endocytosis-dependent nutrient

absorption in specialized enterocytes across animal species.

Keywords: Voltage-sensing phosphatase; VSP; Phosphoinositides; PIPs; endocytosis;

nutrient absorption; enterocyte; lysosome-rich enterocyte; zebrafish; ascidian

Introduction

Phosphoinositides (PIPs) are essential phospholipids that constitute eukaryotic

biological membranes. Differential phosphorylation of inositol head groups at the 3-, 4-, and

5-phosphate positions results in seven PIP species: three mono-phosphorylated PIPs, three bi-

phosphorylated PIPs, and a single tri-phosphorylated PIP. These interconvertible PIP species

are distinctly localized at the plasma membrane and subcellular compartments, regulating

specific membrane-associated activities within the cells, including membrane dynamics, and

cellular trafficking (1-5). Because of their essential roles, PIPs have attracted research

attention for their regulatory mechanism and fundamental interplay in cell physiology.

Voltage-sensing phosphatase (VSP) is among the key molecules that regulate PIPs'

homeostasis. VSP is a unique membrane with two functional domains: the voltage sensor

domain as typically found in voltage-gated ion channels; and the cytoplasmic catalytic region

sharing molecular similarity to the phosphatase and tensin homolog deleted on chromosome

10 (PTEN), a tumor-suppressing PIP phosphatase. VSP functions upon membrane

depolarization to exhibit voltage-dependent phosphatase activity towards PIPs; thus, directly

translating membrane electrical activities into intracellular PIP signals. Unlike PTEN which

exhibits rigid 3-phosphatase activity toward $PI(3,4,5)P_3$ and $PI(3,4)P_2$, VSP exhibits activities

of both 3- and 5-phosphatases, thereby mediating four subreactions: dephosphorylating

PI(3,4,5)P_3 to PI(4,5)P_2 or PI(3,4)P_2; and from PI(4,5)P_2 or PI(3,4)P_2 to PI(4)P (6-8). Among

four subreactions, the reaction with PI(4,5)P_2 to produce PI(4)P is the most robust (7, 9). VSP

orthologs are widely conserved across animal species, and their biophysical properties have

been intensively investigated using in vitro approach. VSP expression has been detected in

various animal tissues, including testis (10-12), epithelium of gastrointestinal tract and renal

tubules (13-16), neurons (6, 17) and blood cells (13). Our recent study has identified that

mouse VSP is required for normal sperm motility through its role in regulating the spatial

distribution of PI(4,5)P_2 in sperm flagellum (11). However, the physiological function of

VSP in most organs remains elusive.

Previous studies demonstrated that zebrafish *Danio rerio* VSP (Dr-VSP) shares

similar molecular architecture to other known VSP proteins and preserves their key

biophysical properties (18, 19). Zebrafish shows conserved physiology and anatomy with

mammals (20). Its optical transparency during early development enables non-invasive

monitoring and in vivo analysis of cellular behaviors (21, 22). By taking these advantages of

the zebrafish, we studied the physiological functions of VSP at the whole animal level. We

show that VSP is highly expressed in lysosome-rich enterocytes (LREs), which are

specialized enterocytes that contain large supranuclear vacuoles and absorb dietary protein

from the intestinal lumen (23-26). We present evidence from in vivo assays that VSP

contributes to the endocytosis of enterocytes. Comparative analysis on ascidian *Ciona*

intestinalis Type A also reveals co-expression of VSP (Ci-VSP) with the endocytic adaptor

Dab2 (Ci-Dab2) in absorptive epithelial cells of the digestive tract, suggesting that the

function of VSP in enterocytes is evolutionarily conserved. These results highlight an

unprecedented role of VSP in the endocytosis-dependent nutrient absorption of intestinal

epithelial cells.

Results

1. Dr-VSP is expressed in the zebrafish intestine.

RT-PCR analysis revealed significant expression of Dr-VSP mRNA in testis, spleen, intestine, kidney, and gills of adult zebrafish (Figure 1A). Dr-VSP mRNA was also detected in 7-dpf zebrafish larvae (Figure 1B). Whole-mounted in situ hybridization (WISH) showed specific expression of Dr-VSP mRNA in the mid-intestine (Figure 1C), with the signal detectable as early as 5 days after fertilization (dpf). The result was comparable to the EGFP signal observed in our *Tg(vsp:EGFP)* transgenic zebrafish (Figure 1, D and E), which were generated using CRISPR-Cas9-mediated knock-in transgenesis (27) to express EGFP recapitulating endogenous expression of *vsp*, the Dr-VSP-encoding gene (Figure 1-figure supplement 1). These findings suggest a potential role of Dr-VSP in the digestive tract from an early developmental stage.

2. Dr-VSP is spatially distributed inside zebrafish enterocytes.

Dr-VSP protein expression was examined using an anti-Dr-VSP antibody (NeuroMab clone N432/21). The immunostainings detected abundant expression of Dr-VSP protein in the epithelial cells of larval pronephros and posterior portion of the mid-intestine (Figure 2A), correlating with the results of WISH and *Tg(vsp:EGFP)* transgenesis.

Here we focused on Dr-VSP expression in the intestine. Principal enterocytes in this region have been characterized as lysosome-rich enterocytes (LREs) that mediate dietary protein absorption via endocytosis (23-26). Dr-VSP immunostaining in the intestine of *Tg(vsp:EGFP)* transgenic zebrafish revealed complete co-localization of Dr-VSP signal and EGFP-positive LREs (Figure 2B). Notably, Dr-VSP expression is absent in the intestinal epithelial cells (IECs), which morphologically differ from LREs by the absence of large supranuclear vacuoles (Figure 2, C and D) (28). Transcriptome database of zebrafish LREs

(Figure 2E) also supports that Dr-VSP is highly enriched in LREs, along with other genes

known to be associated with macromolecule absorption, such as *lamp2*, *cubn*, and *dab2* (23,

29).

The positive signal of the anti-Dr-VSP antibody was not entirely co-localized with

phalloidin staining, which represents actin filaments (F-actin) in microvilli (Figure 3, A and

B), indicating that Dr-VSP in LREs is preferentially distributed at intracellular region beneath

the apical surface rather than at the cell surface or microvilli. Quantification of fluorescence

(Figure 3C) demonstrates that Dr-VSP showed the most intense signal at a deeper region of

the cells than the microvillous F-actins. Immuno-electron microscopy analysis revealed a

particularly high density of immunoparticles at the subapical region of LREs, while it is

rarely detectable at microvilli (Figure 4A, Figure 4-figure supplement 1), corresponding to

the immunofluorescence results. To specify which intracellular structures contain Dr-VSP,

we used the MDCKII cells as a model for examining the intracellular compartments involved

in epithelial endocytosis (30), simulating zebrafish LRE function. In MDCKII cells, Dr-VSP

was co-expressed with (i) Rab5, an early endosome marker; or (ii) Rab11, a recycling

endosome marker. The Dr-VSP signal was broadly co-localized with Rab5-positive and

Rab11-positive intracellular compartments (Figure 4B, Figure 4-video supplement 1, Figure

4-video supplement 2), indicating that Dr-VSP is expressed on the membranes of early and

recycling endosomes.

3. Dr-VSP contributes to the survival and growth of zebrafish larvae.

To investigate Dr-VSP function under physiological conditions, we generated Dr-

VSP-deficient (Dr-VSP$^{-/-}$) zebrafish using CRISPR-Cas9-mediated mutagenesis, targeting the

early exon of the *vsp* gene. Mutant zebrafish lacking the transmembrane segments and

cytoplasmic catalytic region of Dr-VSP were obtained (Figure 5-figure supplement 1).

Dr-VSP[-/-] larvae showed normal gross morphology and behavior. However, we

noticed that the survival rate of Dr-VSP[-/-] mutants was significantly lower than that of wild-

type larvae at the age of 10 – 14 dpf (Figure 5A). During this period, the yolk is completely

reabsorbed, and zebrafish growth is generally dependent on exogenous feeding via the

digestive system (28, 31). Moreover, Dr-VSP[-/-] mutants that survived from the early period

exhibited long-term growth restriction, as evidenced by their smaller size compared to their

sibling wild-types (Figure 5, B and C). Despite the absence of obvious congenital deformities,

these results could be inferred that Dr-VSP function contributes to the survival and growth of

zebrafish larvae during early life. The remaining Dr-VSP[-/-] mutants thrived to adulthood and

reached productive maturity at around the same age as wild-type zebrafish without apparent

fertility defect.

4. Dr-VSP facilitates endocytic nutrient absorption in larval LREs.

Dr-VSP is abundant at the endomembranes inside LREs, which absorb nutrients via

endocytosis (23-26), suggesting that Dr-VSP could be involved in this specialized process.

To test this hypothesis, we compared the endocytosis efficiency between the wild type and

Dr-VSP[-/-] larvae by gavaging 6-dpf zebrafish larvae with either fluorescent dextran (fDex) or

mCherry solution directly into the digestive tract (Figure 6A) (32), and subsequently

observed endocytosis in LREs using confocal microscopy. fDex was used for assessing fluid-

phase endocytosis; and mCherry was used as a protein cargo for assessing receptor-mediated

endocytosis (23, 29).

At two hours after gavage, fDex and mCherry were absorbed into LREs in both wild-

type and Dr-VSP[-/-] zebrafish larvae (Figure 6, B and C). However, the uptake of fDex and

mCherry were reduced in Dr-VSP[-/-] LREs (Figure 6, D and E). In the experiment with fDex,

the fluorescent signals in Dr-VSP[-/-] LREs were mostly confined to apical vesicles, with less

signal found in larger endosomes and supranuclear vacuoles (Figure 6D). The average

percentage of LREs containing fDex-filled endosomes was nearly 80% in wild-types but less

than 30% in Dr-VSP$^{-/-}$ larvae (Figure 6F). We further analyzed the relative fDex

internalization profile of LREs by measuring fluorescent intensity within the cells (Figure

6G) (23, 29). As shown in Figure 6G, fDex uptake into Dr-VSP$^{-/-}$ LREs was clearly impaired

both in the apicobasal axis of individual cells and the anteroposterior axis across longitudinal

segments of LREs. Similar results were obtained in the mCherry experiment: the average

percentage of LREs containing mCherry-filled endosomes was greater than 80% in wild-

types versus 40% in Dr-VSP$^{-/-}$ larvae (Figure 6H). Spatial profile of the marker distribution in

the cells (Figure 6I) also showed reduced uptake of mCherry in Dr-VSP$^{-/-}$ LREs.

When we gavaged wild-type zebrafish with the mixture of fDex and mCherry, we

found that while most LREs contained both fluorescent cargoes in the same supranuclear

vacuoles, some vacuoles contained only fDex or mCherry (Figure 6-figure supplement 1A).

In Dr-VSP$^{-/-}$ mutants, on the other hand, mCherry was transported into some supranuclear

vacuoles, whereas fDex was mostly found in apical vesicles inside Dr-VSP$^{-/-}$ LREs (Figure 6-

figure supplement 1B). This suggests that the uptake defect in Dr-VSP$^{-/-}$ LREs was more

severe with fDex than with mCherry.

Taken together, these findings support our hypothesis that Dr-VSP is involved in the

endocytosis of LREs, specifically in the maturation of endosomal vacuoles following initial

endocytosis. The findings also suggest that Dr-VSP plays a greater role in fluid-phase

endocytosis, the primary mechanism of fDex uptake into cells.

5. Dr-VSP maintains intact LRE morphology.

We examined the LRE morphology of 14-dpf zebrafish larvae, the age around one

week after zebrafish enterocytes completed morphogenesis (Figure 7A) (28, 31). This age

also corresponds to the period when higher mortality was observed in Dr-VSP$^{-/-}$ larvae

(Figure 5A). We measured key parameters that represent cell morphology: central height,

apical width, basal width, and microvillus length (33). The measurement revealed that Dr-

VSP$^{-/-}$ LREs were significantly shorter in all other parameters except for the microvillus

length (Figure 7B). The fact that Dr-VSP deficiency primarily affected the cell body is

consistent with the idea that Dr-VSP plays a major role at the endomembranes within cells

than at the plasma membrane of microvilli. The defect in LRE morphology was not observed

at early stages, such as at 6 dpf (Figure 7C). We examined histological sections of 14-dpf

zebrafish LREs with toluidine blue staining and analyzed endosomal vacuoles, comparing

wild-type and Dr-VSP$^{-/-}$ mutants (Figure 7D). Although supranuclear vacuoles could be seen

in most cells, the measurement of vacuolar perimeter and area showed that their size was

apparently smaller in Dr-VSP$^{-/-}$ LREs (Figure 7, E and F). We also used transmission electron

microscopy (TEM) to define the ultrastructures of 14-dpf LREs, focusing on the subapical

region where Dr-VSP is typically expressed. TEM images of wild-type and Dr-VSP$^{-/-}$ LREs

(Figure 7G) revealed key features of absorptive enterocytes, such as membrane invaginations

at inter-microvillous spaces, cytoplasmic tubules, branching tubules, and endocytic vesicles

associated with invaginated membranes (tubule-vesicle complexes) (24, 34, 35). Interestingly,

these features appeared less prominent in Dr-VSP$^{-/-}$ LREs. Further observations suggested

that small endocytic vesicles in Dr-VSP$^{-/-}$ LREs are located at deeper regions than those in

wild-types, increasing the distance between the apical surface and vesicle-dense area (Figure

7-figure supplement 1). Moreover, branching tubules and tubule-vacuole complexes were

less frequently present in the subapical region of mutant LREs (Figure 7-figure supplement 2),

seemingly reflecting the disturbance in endosome maturation. Collectively, these results

suggest that LREs require Dr-VSP in regulating the formation and expansion of endosomal

vacuoles, which in turn maintain intact LRE morphology. Figure 7H summarizes how Dr-

VSP contributes to endosomal vacuoles and the morphology of LREs.

6. Endocytic role of VSP is potentially conserved in the ascidian digestive tract.

VSP gene is widely conserved among chordates (Figure 8-figure supplement 1, Figure

8-figure supplement 2). To determine if the role of VSP in the endocytosis of enterocytes is

conserved among animal species, we investigated the expression profile of sea squirt *C.*

intestinalis VSP (Ci-VSP) in the digestive tract. Sea squirt belongs to the basal chordates, and

the gene expression profiles of *C. intestinalis* Type A were studied in great detail with

reference to chordate evolution (36). WISH on transparent 2-week-old *Ciona* juveniles

revealed the expression of Ci-VSP mRNA in the stomach, mid-intestine, and posterior

intestine (Figure 8A), which were known as absorptive regions of the *Ciona* digestive tract

(37). Higher magnification showed a strip-like expression pattern both in the stomach and

bulged intestine. Similar patterns were found in the digestive tract of *Ciona* young adults,

with the expression signal being more robust in the stomach and mid-intestine than in the

posterior intestine (Figure 8B). Additional Ci-VSP expression was observed in the looped

region of the intestine (Figure 8B, white arrowhead), which corresponds to a region

expressing several absorptive genes; including peptide transporter (PEPT1) and

monosaccharide transporters (SGLT1, GLUT5) (Figure 8-figure supplement 3).

Transverse sections of the digestive tract obtained from *Ciona* juveniles (Figure 7C)

showed that the expression signals were present alternately in the digestive epithelial cells. In

the stomach, Ci-VSP signals were observed mainly in the inner folds, where the epithelial

cells predominantly express absorptive genes (38). In the outer-folds of the stomach, known

to express many pancreatic-related exocrine genes (39), Ci-VSP signals were occasionally

found in the regions which seemed to start invagination of newly generated inner-folds.

Similarly, in the bulged region of mid-intestine, circumferential epithelial cells expressed Ci-

VSP in a pattern corresponding to the alternating folding morphology as present in the

stomach.

253   We further examined whether Ci-VSP expression correlates with endocytosis-

associated genes. Several genes involved in enteric endocytosis in the zebrafish model have

previously been identified, including Cubilin (*cubn*), Amnionless (*amn*), and Dab2 (*dab2*)

(23). In *Ciona*, however, only *dab2* ortholog (Ci-Dab2) is expressed as a single homolog, so

Ci-Dab2 was selected to represent the endocytosis-associated gene in this study. In WISH on

2-week-old *Ciona* juveniles, the Ci-Dab2 expression pattern is highly comparable to that of

Ci-VSP (Figure 7, D – F), implying their concerted function in the *Ciona* digestive tract. Ci-

VSP is co-expressed with Ci-Dab2 in a strip-like pattern in the stomach and bulged intestine,

indicating that Ci-VSP is expressed in the epithelial cells with endocytosis-dependent

absorptive function.

Discussion

265   VSP expression was previously reported in the digestive organs of other animal

species, such as in the gut epithelial cells of invertebrate *C. intestinalis* Type A (13) and the

stomach of chicken embryos (17). However, the functional role of VSP in the enterocytes has

not been studied in detail. We report here the expression of VSP in specialized enterocytes

known as LREs in the mid-intestine of zebrafish. VSP in LREs is most likely expressed on

the endosomal membranes required for endocytosis. Loss of VSP activity impaired LRE

endocytosis and nutrient absorption, potentially accounting for growth restriction and

increased mortality in developing zebrafish. Dr-VSP was detectable in LREs as early as 5 dpf

(Figure 1 and 2), coinciding with the time when zebrafish intestine begins nutrient absorption

(40, 41). These findings highlight a novel physiological role of VSP in endocytosis-

dependent nutrient absorption besides its previously established role in sperm physiology

(11).

Localization of VSP on endomembranes

Previous research indicated that VSP is expressed on the plasma membrane, based on

studies of native expression in *Ciona* sperms (6), *Xenopus* renal epithelial cells (14, 16), and

mouse sperms (11), as well as in vitro experiments using heterologous expression system (6,

15, 42, 43). The same was reported for Dr-VSP when it was first characterized by whole-cell

patch-clamp recordings of mammalian cells heterologously expressing Dr-VSP protein (18).

However, our results of immunofluorescence staining and immuno-electron microscopy

revealed that native Dr-VSP in LREs was localized to the subapical region but not on

microvilli nor plasma membrane (Figure 3 and Figure 4A). Results from the MDCK model

indicate that Dr-VSP is enriched at the early and recycling endosomal membranes (Figure

4B), consistent with our recent in vitro study in which fluorescent signal from mCherry-fused

Dr-VSP expressed in HEK293T cells was also distributed throughout the intracellular

compartments (19). Although additional studies, including in vivo co-localization analysis

with endosome markers, are required, our current findings provide converging evidence that

Dr-VSP functions primarily on endomembranes at the subapical region of LREs.

PIP conversion has been well documented to regulate membrane trafficking in the

endosomal system, and this process is achieved by PIP-metabolizing enzymes, including PIP

kinases and phosphatases (3-5). Dr-VSP functions as 5-phosphatase and 3-phosphatase,

allowing it to hydrolyze $PI(3,4,5)P_3$, $PI(4,5)P_2$, and $PI(3,4)P_2$ (7, 9). Following gavage with

fDex and mCherry, we observed that endocytosis was remarkably decreased in Dr-VSP[-/-]

LREs, as evidenced by the reduction of subsequent endosomalz vacuoles and fluorescence

uptake into the cells (Figure 6). Histological and TEM findings show that loss of Dr-VSP

resulted in fewer and smaller vacuoles and, consequently, a reduction in the overall cell

volume of LREs. Besides, Dr-VSP$^{-/-}$ LREs show reduced numbers of endocytic vesicles,

cytoplasmic tubules, and tubule-vacuole complexes at the subapical region (Figure 7, Figure

7-figure supplement 1 and 2). These findings suggest that Dr-VSP is involved in endocytic

regulation, and Dr-VSP deficiency disrupts membrane trafficking in LREs.

PIPs enzyme activity of VSP in endomembranes of LREs

Dr-VSP deficiency may alter PIP homeostasis on various types of endomembranes,

which play a critical role in membrane trafficking. However, it remains a challenge to

determine which enzyme subreaction mediated by VSP contributes to the phenotypes in

LREs. Here we propose potential scenarios based on Dr-VSP's dual phosphatase activities.

One scenario considers a more robust Dr-VSP's 5-phosphatase activity toward PI(4,5)P$_2$ and

PI(3,4,5)P$_3$ (9). We speculate that intracellular Dr-VSP dephosphorylates such PIPs on the

membranes of newly enclosed endocytic vesicles during the transformation to early

endosomes (44-46). Defective endocytosis in Dr-VSP$^{-/-}$ LREs is likely caused by reduced

dephosphorylation of endosomal PI(4,5)P$_2$ and possibly PI(3,4,5)P$_3$, leading to ectopic

accumulation of these PIPs that prevent apical endocytic vesicles from transforming into, or

fusing with, subsequent intracellular compartments during endosomal maturation; thus

restricting most fDex and mCherry to the subapical region of Dr-VSP$^{-/-}$ LREs. It should be

noted that other 5-phosphatase, such as OCRL1 and INPP5B, can be detected in LREs,

although at lower levels according to RNA-seq data (23). These enzymes are involved in

early endocytic trafficking (46-49) and can function similarly as VSP during active

endocytosis. The presence of compensatory enzymes may account for the phenotypes of Dr-

VSP$^{-/-}$ LREs (Figure 6, F and H); some Dr-VSP$^{-/-}$ zebrafish larvae survived and thrived into

adulthood despite seemingly suffering from nutrient malabsorption.

Another scenario considers Dr-VSP's 3-phosphatase activity towards $PI(3,4)P_2$ and

$PI(3,4,5)P_3$ (9). Dr-VSP at recycling endosomes may be involved in the conversion of PI(3)P

to PI(4)P by dephosphorylating the intermediary $PI(3,4)P_2$ (45). Unless a compensatory

mechanism is available, the loss of Dr-VSP's 3-phosphatase activity can cause excessive

$PI(3,4)P_2$ retention and, as a result, disrupt the formation of recycling endosomes during

endocytic recycling in LREs. The defect may also reduce subsequent endocytosis when the

internalized membrane and specific receptors are unable to fully return to the plasma

membrane. Furthermore, based on a previous study showing that VSP exhibits small but

significant activity toward $PI(3,5)P_2$ by in vitro enzyme assay (50), Dr-VSP may contribute to

$PI(3,5)P_2$ homeostasis in the late endocytic pathway in LREs (5, 45). The precise mechanism

by which Dr-VSP is recruited to function at specific endomembranes has yet to be

determined, and this will be one of the important issues to be addressed in the future for

better understanding the molecular mechanisms underlying the physiological role of VSP in

LREs.

Two distinct endocytic pathways have been described in zebrafish LREs: (i) fluid-

phase endocytosis, for fDex uptake; and (ii) receptor-mediated endocytosis, for mCherry

uptake (23). In our study, Dr-VSP deficiency in zebrafish LREs affected fDex uptake more

remarkably than mCherry uptake. Furthermore, following the gavage with fDex-mCherry

mixture, the difference in the intracellular distribution of fDex and mCherry in LREs was

more pronounced in $Dr\text{-}VSP^{-/-}$ mutants (Figure 6-figure supplement 1). These results suggest

that the extent to which Dr-VSP contributes to LRE endocytosis differs between fluid-phase

and receptor-mediated endocytosis, possibly due to subtle differences in molecular

mechanisms, such as PIP conversion, between the two pathways.

Roles of VSP in other tissues in zebrafish

Our immunostaining results indicate that Dr-VSP was also expressed in other tissues than the intestine. Among them, the expression in testis and kidney is consistent with the results obtained from other species (11, 12, 14, 16), indicating that VSP is functionally conserved across animal species. We suspected that defective endocytosis in Dr-VSP$^{-/-}$ LREs is most responsible for growth restriction and higher mortality in zebrafish larvae (Figure 5), particularly during the critical period when zebrafish larvae rely primarily on exogenous feeding via intestinal absorption (28, 31). Although Dr-VSP is also expressed in larval pronephros (Figure 2A), Dr-VSP deficiency in this organ is less likely to be the primary cause of mortality. This is because larval pronephros begins to function at around 48 hpf (51), and defective pronephros should exhibit phenotypes earlier than what was observed in our study. We have previously reported that VSP regulates mouse sperm motility via its 5-phosphatase activity toward PI(4,5)P$_2$ on the plasma membrane during sperm capacitation (11). VSP deficiency in mice caused abnormal sperm movement and significantly reduced in vitro fertilization. Dr-VSP is expressed in zebrafish testis (Figure 1A) (7), but Dr-VSP$^{-/-}$ zebrafish have no apparent fertility defect. Further research may be necessary to determine whether Dr-VSP in zebrafish sperms serves a similar function as its ortholog in mouse sperms.

Stimulus to activate VSP

VSP is a membrane protein that functions upon membrane depolarization (6-8). Our study raises an open question of how Dr-VSP is activated in LREs. According to the in vitro experiment, the voltage required to activate Dr-VSP is beyond the physiological range of membrane potential in native cells (18). Unfortunately, there is no current in vivo method for measuring the membrane potential of intracellular vesicles in zebrafish LREs. However,

several possible mechanisms may induce Dr-VSP activation in endosomes. First, unlike in

vitro conditions, native Dr-VSP in LREs may function at physiological membrane potentials,

possibly through interactions with unknown endogenous auxiliary subunits, and a partial

activation could be sufficient to induce Dr-VSP phosphatase activity during active

endocytosis. This is similar to what may happen in mouse sperms, where physiological

membrane potential appears to activate mouse VSP (11). Second, membrane dissociation

during endosomal vesicle formation could alter membrane potential probably by changing

membrane capacitance, ultimately accelerating depolarization. Third, changes in micro-

environments such as membrane tension, luminal pH, and opening of endomembrane ion

channels (52-54) may facilitate Dr-VSP activation under physiological conditions. Further in

vivo analyses are required to explore the relationship among membrane potential, VSP

activation, and endosomal membrane trafficking. Potential future tests include rescue

experiments of Dr-VSP$^{-/-}$ mutants by expressing modified versions of VSP, such as a voltage-

insensitive or a voltage-hypersensitive version, to study how membrane potential affects

endocytosis in LREs. Additionally, optogenetics could be applied in zebrafish to manipulate

membrane potential by specifically targeting photoactivated ion transporting molecules to

specific membrane vesicles.

Role of VSP shared among chordates

Our comparative study reveals that VSP is conserved in the digestive tracts of

zebrafish and *C. intestinalis* Type A, as are other endocytosis-associated genes (Figure 2E

and Figure 8, D – F) (23, 55, 56), suggesting that Dr-VSP and Ci-VSP share a conserved

endocytic function in the absorptive epithelial cells. Endocytosis-dependent nutrient

absorption is conserved across species from unicellular organisms to vertebrates. Unicellular

organisms acquire nutrients from their surroundings by endocytosis, and many invertebrates

show LRE-like gastrodermal cells that line the digestive tract to absorb nutrients by

endocytosis (57). Endocytotic absorption through epithelial cells has also been reported in the

nutrient absorptive organs, such as the trophotaenia of viviparous fish embryos (58) and the

ileum of pre-weaning mammals (59, 60). Furthermore, LREs have also been proposed to be

involved in trans-cellular transport of antibodies and other antigens across enterocytes to

extra-intestinal tissues for immune modulation (23, 31, 60-62). Because both of LRE-like

cells and VSP gene are widely observed from Cnidaria to mammals (23, 57, 59, 60, 63), the

endocytic role of VSP may well be conserved among a wide range of animal species.

Materials and Methods

Key resources table

Experimental model and subject details

Zebrafish maintenance.

We used RIKEN wild-type zebrafish obtained from RIKEN Brain Science Institute

(Saitama, Japan), and *Tg(vsp:EGFP)* transgenic zebrafish that express EGFP under the

endogenous expression of *vsp*, the Dr-VSP-encoding gene. The fish were fed a non-calorie-

restricted standard diet twice daily and raised in an aquatic system maintained at 28°C with a

14 h/10 h light/dark cycle. All experiments were conducted under the regulation of the

Animal Care Facility, Center for Medical and Translational Research (CoMIT) at Osaka

University (Osaka, Japan).

Biological materials for Ciona intestinalis's experiments.

Matured and young adult *Ciona intestinalis* type A (*C. robusta*) were collected during

the spawning season at Port Chiba in Tokyo Bay (Chiba, Japan). *Ciona* Juveniles were

developed by artificial insemination using matured eggs and sperms obtained from dissected

gonoduct. The specimens of whole-mount juveniles and dissected post-pharyngeal region of

the digestive tracts of the young adult were prepared for in situ hybridization as previously

described (39).

Method details

RNA isolation and Reverse Transcription PCR (RT-PCR).

Total RNA was purified from zebrafish larvae and adult tissues using TRIzol^{TM} LS

reagent (Invitrogen, USA) with volume adjustment for small samples. Total RNA (200 ng)

extracted from each sample was reverse-transcribed using SuperScript^{TM} III First-Strand

Synthesis System (Invitrogen, USA). For semi-quantitative RT-PCR, fragments of *vsp* and *β-*

actin were amplified from cDNA using the primer sets listed in Supplemental Table S1.

Whole-mount in situ hybridization (WISH) for zebrafish larvae.

Zebrafish larvae were fixed in 4% PFA/PBS at 4ºC overnight. Next, the larvae were

washed twice with PBS for 5 min each time, followed by briefly washed with 100% MeOH

and kept in MeOH at -20ºC until the following steps. WISH was performed using a

digoxigenin (DIG)-labeled RNA probe according to the standard protocol. In this study, the

Dr-VSP cDNA was subcloned into pBluescript II KS (+) plasmid vector. The resultant

plasmid was linearized by XhoI enzyme and later used as a template for synthesizing DIG-

labeled RNA probe using T3 RNA polymerase. At the beginning of WISH process, fixed

zebrafish larvae were treated with proteinase K (10µg/mL) for 15 min at room temperature.

Prehybridization was done at 55ºC for 1 h. Larva samples were hybridized with 10 ng of

DIG-labeled Dr-VSP cRNA probe overnight. During posthybidization, larva samples were

washed with PBS containing 0.2% Tween-20 and then incubated with anti-DIG-alkaline

phosphatase (AP), Fab fragments (Roche, Mannheim, Germany) at 4ºC overnight. The

samples were subsequently washed with PBS containing 0.2% Tween-20 every 20 min for 3

h, to remove excessive antibody. Positive signals of the hybridized Dr-VSP mRNA transcript

were visualized by color development using BCIP/NBT solution (Wako, Japan) for 2 h at

room temperature. The stained larvae were observed under Zeiss Stemi SV11 Stereo

Microscope (Zeiss, Berlin, Germany).

In situ hybridization, detection, and observation of gene expression in Ciona intestinalis.

DIG-labeled antisense RNA probe for *Ciona* genes were synthesized from T7 RNA

polymerase promoter-attached amplified cDNA, and probes were purified by centrifugal

ultrafilter as previously described (13). WISH of the *Ciona* specimens was performed using

"InSitu Chip" (64). Gene expression was visualized using BCIP/NBT solution (Roche,

Mannheim, Germany). After color development, the digestive tracts of the *Ciona* juveniles

were sectioned transversely with a thickness of 10 μm (39). Whole-mount and sectioned

specimens of *Ciona* were observed under BX51 microscope (Olympus, Japan) or SZX12

Stereo Microscope (Olympus, Japan).

CRISPR-Cas9-mediated transgenesis and mutagenesis.

Tg(vsp:EGFP) zebrafish line was generated via CRISPR-Cas9-mediated knock-in

transgenesis (27) with adaptation to the *vsp* gene (Figure 1-figure supplement 1). The sgRNA

for genome digestion contained a target site (AAACCGCTACGTCTGCAGCAGG) derived

from the *vsp* exon sequence located between the transcriptional start site and translational

start site. Donor DNA for knock-in contained Mbait-hsp70 promoter-EGFP-polyA sequences.

Successful knock-in transgenesis was validated by sequence mapping for donor DNA

insertion and comparing EGFP signal with *vsp* expression pattern. Dr-VSP-deficient (Dr-

VSP$^{-/-}$) zebrafish were generated via CRISPR-Cas9-mediated mutagenesis. Potential CRISPR

targets for the *vsp* gene were identified via the online tool CHOPCHOP (65) based on

genomic DNA sequence obtained from the Ensembl zebrafish genome database version 9

(Zv9, https://asia.ensembl.org/Danio_rerio/). We selected three targets close to the translation

initiation site without potential off-target (Supplemental Table S2). In our setting, target T1 in

exon 6 (Figure 5-figure supplement 1) yielded the highest efficiency in producing adequate

frameshift mutation. Microinjection for Dr-VSP mutagenesis was performed base on the

reported protocol (66). All injected embryos were incubated at 28.5°C to support early

development. Subsequently, some injected embryos underwent genome extraction and

sequence analysis, while the remaining embryos were raised as the founder (F0) generation.

Genomic DNA extraction and sequence analysis.

Genomic DNA was isolated from whole larvae or fin-clips of adult zebrafish using the

HotSHOT DNA extraction procedure (67). Genomic DNA fragments covering CRISPR

targets were amplified by PCR using locus-specific primer sets listed in Supplemental Table

S2. We used BigDye^{TM} Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific,

Waltham, MA) for analyzing the sequences of PCR fragments.

Fish sectioning, immunohistochemistry (IHC), and toluidine staining.

Zebrafish larvae were fixed in 4%PFA/PBS, followed by 30% sucrose/PBS, and

embedded in 5% ultra-low gelling temperature agarose (Sigma-Aldrich, USA). Embedded

specimens were cryosectioned in 10 µm slices using Leica CM3050S cryostat (Leica,

Germany). Antibodies used for IHC are as follows: anti-Dr-VSP/TPTE (NeuroMab, USA),

Acti-stain™ 488 Phalloidin (Cytoskeleton, USA), Alexa Fluor^{TM} 594 Phalloidin (Invitrogen,

USA), and anti-Na^{+}-K^{+} ATPase (Ab76020; Abcam, USA). Images were taken under Zeiss

LSM880 confocal microscope with Airyscan (Zeiss, Berlin, Germany). For toluidine blue

staining, larvae were fixed in 4%PFA/PBS at 4°C overnight, followed by 2.5%GA at 4°C for

2 h and 1%GA overnight. Samples were subsequently dehydrated by ethanol substitution and

embedded in Epon for thin sectioning before the staining process.

Transmission electron microscopy (TEM).

Zebrafish were fixed in 4%PFA at 4°C overnight before en-bloc sectioning of the

mid-intestine region. Sectioned samples were subsequently fixed in 2.5%GA at 4°C for 2 h,

followed by 1%GA overnight. Specimen preparation and staining were done as standard

protocol, with technical support from Center for Medical Research and Education, Osaka

University. Briefly, the fixed samples were incubated with 1% osmium tetroxide, then

dehydrated in ethanol series and propylene oxide. The samples were then embedded in Epon

and ultra-thin sectioned, followed by contrast stained with uranyl acetate and lead citrate.

TEM images were acquired under the Hitachi H-7650 transmission electron microscope

(Tokyo, Japan) operating at 80 kV, and analyzed using ImageJ software (NIH, USA).

Pre-embedding immuno-electron microscopy (Immuno-EM)

Zebrafish larvae were fixed in 4%PFA/HEPES, or 4%PFA/0.1%GA/HEPES, before

being embedded in 5% ultra-low gelling temperature agarose (Sigma-Aldrich, USA) and

cryosectioned in 10 µm slices using Leica CM3050S cryostat (Leica, Germany). Sectioned

samples were incubated in 5% normal goat serum (NGS)/0.125% saponin/PBS for 30 min,

then incubated in anti-Dr-VSP/TPTE antibody (NeuroMab, USA) at 37°C for 2 h, followed

by Alexa Fluor® 488 FluoroNanogold antibody (Nanoprobes, USA) at room temperature for

2 h. Immunogold particles were intensified at room temperature with HQ SILVER^{TM}

enhancement kit (Nanoprobes, USA) and 5% selenium toner. Samples were further treated

with 1% osmium tetroxide, 0.5% uranyl acetate, and underwent the same process as

described in the TEM protocol.

Cell culture and transfection.

MDCK type 2 (MDCKII) cells were cultured in D-MEM high glucose (Wako, Japan)

containing 10% FBS at 37ºC under 5% CO_2. Transient expression was achieved by co-

transfecting mCherry-Dr-VSP plasmid with (i) EGFP-Rab5 plasmid (early endosome marker)

or (ii) EGFP-Rab11 plasmid (recycling endosome marker) into MDCKII cells using

Lipofectamine^{TM} 3000 transfection reagent (Invitrogen, USA).

Fish gavage and live imaging.

6-dpf zebrafish larvae were gavaged with Alexa Fluor 488-tagged dextran 10,000MW

(fDex) solution (Molecular Probes, USA) or purified mCherry solution (prepared in the

laboratory) following the established method (32) (see also Figure 6A). A microforged

capillary needle loaded with either solution was directly inserted through the larval mouth to

inject the solution into the anterior intestinal bulb. Gavaged larvae were incubated at 28.5°C

for 2 h to allow intestinal absorption before live image acquisition under Zeiss 710 confocal

microscope (Zeiss, Berlin, Germany) with temperature control at 28.5°C for entire sessions.

The following parameters were analyzed: (i) the percentage of vacuolated cells per section;

and (ii) relative fDex or mCherry internalization profile measured across the apicobasal axis

of individual cells and the longitudinal axis along the posterior portion of mid-intestine. The

data of (ii) were analyzed by measuring fluorescence intensity using Plot Profile function in

ImageJ software (NIH, USA) as previously described (23) and presented as means ± SEM

percentage of intracellular fluorescence intensity along the specified axes, with a 100%

indicates the maximal value of the average intensity on each axis.

Quantification and statistical analysis

Parameter analysis of larval enterocytes.

Quantification of cellular dimensions of LREs was done based on the previous

procedure (33). We performed immunostaining with Acti-stain^{TM} 488 Phalloidin

(Cytoskeleton, USA) and anti-Na$^+$-K$^+$ ATPase Antibody (Ab76020; Abcam, USA) to outline

individual cell (Figure 7A). In the mid-intestine region, each confocal image was selected

where enterocytes appeared the broadest, and the following parameters were analyzed:

central height, apical width, basal width, and microvillus length.

Body length and survival analysis.

Wild-type and Dr-VSP$^{-/-}$ zebrafish were raised in the standard system of the Animal

Care Facility, Center for Medical and Translational Research (CoMIT), Osaka University.

Body length and remaining number of larvae were measured periodically from the day of

fertilization to 28 dpf. All data were collected from 3 sets of independent experiments.

Analysis of gene expression level using RNA-seq data

Raw data of the RNA-seq experiments were retrieved from the GEO Datasets

database with accession number GSE124970 (23). Heatmap representing expression levels of

selected genes in IECs and LREs (Figure 2E) was analyzed using web-based application

iDEP v0.92 (http://bioinformatics.sdstate.edu/idep92) (68)

Image processing and statistical analysis.

Raw images were processed and analyzed using ImageJ software (NIH, USA).

Graphs and statistical analyses were performed using Prism 8 (Graphpad Software, San

Diego, CA). Statistical details for each experiment can be found in the figure legends.

Acknowledgments

We thank Dr. Shinji Kanda (University of Tokyo, Japan) for helpful advice on this

study. Our thanks are also due to Drs. Junji Takeda, Akihiro Harada, Shin-ichiro Yoshimura,

Atsushi Tamura, Hiroo Tanaka, Yoshifumi Okochi, Akira Kawanabe (Osaka University,

Japan), and all laboratory members for valuable suggestions and discussions. We thank Mr.

Eiji Oiki (Osaka University, Japan) for advice in electron microscopy; Mr. Masayuki

Yamagishi and Mr. Satoshi Nakayama (Chiba University, Japan) for advice and technical

assistance in the in-situ hybridization of the *Ciona* specimens. Technical support for this

study was provided by the Center for Medical Research and Education and the Center for

Medical and Translational Research (CoMIT) at Osaka University.

Additional information

Funding information

This work was supported by Grants-in-Aid from JSPS (12J01957, 15K18575) (to

T.K.), Grants-in-Aid from Ministry of Education, Culture, Sports, Science, and Technology

(MEXT) and JSPS (15H05901, 25253016 to Y.O.), JSPS (16H02617 to T.K. and Y.O.) and

Core Research for Evolutional Science and Technology, Japan Science and Technology

Agency (CREST, JST) (JPMJCR14M3 to Y.O.) M.M. was supported by Grans-in-Aid from

MEXT (A00H059010, A15H059010). A.R. was supported by Interdisciplinary Program for

Biomedical Sciences (IPBS), Osaka University

Author contributions (CRediT taxonamy)

Conceptualization and Methodology, A.R., T.K., and Y.O.; Investigation, A.R., M.M., S.H., Y.K., F.T., T.M., M.I.H., and M.O.; Formal Analysis, A.R., T.K., and Y.O.; Visualization and Writing - Original Draft, A.R.; Writing - Review & Editing, A.R., T.K., Y.O.; Funding Acquisition, T.K., Y.O., A.R.; Supervision, Y.O.

Ethics

Protocols used for the animal experiments in this study were approved by the Animal Research Committee of Osaka University, Japan (No. 27-079). All procedures were conducted in accordance with the regulation of the Animal Care Facility, Center for Medical and Translational Research (CoMIT) at Osaka University, Japan.

Competing interests

The authors declare that no competing interests exist.

Figure legends

Figure 1: Expression profile of the Dr-VSP-encoding gene (*vsp*) in zebrafish tissues.

(A) RT (+) *vsp* expression in adult zebrafish tissues: (1) testis, (2) spleen, (3) liver, (4) intestine, (5) kidney, (6) heart, (7) gill, (8) eyeballs, (9) brain, (10) ovaries. RT (-), negative control for all tissues in the same order. *β-actin*, positive control. M, marker.

(B) RT (+), *vsp* expression in 7-dpf zebrafish larva. RT (-), negative control. *β-actin*, positive control. M, marker.

(C) Whole-mounted in situ hybridization (WISH) of 6-dpf zebrafish larva showing *vsp*-positive signal (arrowhead) in enterocytes of mid-intestine.

(D) A 5-dpf *Tg(vsp:EGFP)* zebrafish larva expressing EGFP under the promotor of

the Dr-VSP-encoding gene in enterocytes of mid-intestine (arrowhead). The EGFP-positive

cells in this figure correspond to the *vsp*-positive cells in Figure 1C. Green, Dr-VSP

(*vsp:EGFP*). Scale bar = 200 µm.

(E) Confocal image of larval enterocytes at mid-intestine of a 5-dpf *Tg(vsp:EGFP)*

zebrafish larva (sagittal section). The enterocytes in this region are known as lysosome-rich

enterocytes (LREs) that contain large supranuclear vacuoles. Green, Dr-VSP (*vsp:EGFP*).

Blue, DAPI. Scale bar = 25 µm

Figure 2: Expression profile Dr-VSP protein in zebrafish tissues.

(A) Confocal images of Dr-VSP immunostaining in a 7-dpf wild-type zebrafish larva

(sagittal section). Dr-VSP is highly expressed in (I) larval pronephros and (II) enterocytes at

the posterior part of mid-intestine, which are defined as lysosome-rich enterocytes (LREs).

Green, Dr-VSP. Blue, DAPI. Scale bar = 500 µm (left) and 50 µm (middle and right)

(B) Dr-VSP immunostaining in LREs of a 7-dpf *Tg(vsp:EGFP)* transgenic zebrafish

larva (sagittal section). Dr-VSP expression signals were completely co-localized with EGFP

signals in LREs. Green, Dr-VSP *(vsp:EGFP)*. Red, Dr-VSP (antibody). Blue, DAPI. Scale

bar = 20 µm.

(C) Dr-VSP immunostaining in a 7-dpf *Tg(vsp:EGFP)* transgenic zebrafish larva

(sagittal section), showing a border between intestinal epithelial cells (IECs; VSP-negative)

and LREs (VSP-positive). LREs contain large supranuclear vacuoles in the cytoplasm,

contrasting to IECs. Green, Dr-VSP *(vsp:EGFP)*. Red, Dr-VSP (antibody). Blue, DAPI.

Scale bar = 20 µm. Background signal of EGFP channel was increased in the left figure to

enhance visibility of IECs.

(D) Dr-VSP immunostaining in IECs (left) and LREs (right) of a 7-dpf wild-type

zebrafish larva (sagittal section). Dr-VSP expression was restricted mainly to LREs, but not

detectable in other IECs. Green, Dr-VSP. Red, F-actin. Blue, DAPI. Scale bar = 10 µm.

Images were visualized under LSM880 confocal microscope with Airy Scan.

(E) Heatmap of RNA-seq data representing expression levels of selected genes in

IECs and LREs. Raw data of RNA-seq experiments were retrieved from the GEO Datasets

database with accession number GSE124970 (23), and analyzed using web-based application

iDEP v0.92 (68).

Figure 3: Expression profile Dr-VSP protein at subapical region of LREs.

(A) Confocal images of Dr-VSP immunostaining showing the subcellular distribution

of Dr-VSP in larval LREs. Dr-VSP is primarily localized in the cytoplasm underneath apical

surface, but barely detectable at the microvilli and plasma membrane of LREs. Green, Dr-

VSP. Red, F-actin, Blue, DAPI. *Vacuole. Scale bar = 10 µm.

(B) Confocal image of larval LREs in a higher magnification and their corresponding

schematic illustrations, focusing on the intracellular distribution of Dr-VSP. Green, Dr-VSP.

Red, F-actin. Blue, DAPI. V, vacuole. MV, microvilli. L, lumen. Scale bar = 5 µm.

(C) Fluorescence intensity of Dr-VSP (green) and F-actin (red) along the apico-basal

axis of individual LREs. Data are presented as mean ± SEM. Enterocytes, ≥ 16 cells from 4

larvae for each zebrafish line. Dotted line in the graph conceptually indicates the junction

between microvilli and cell body.

Figure 4: Expression profile Dr-VSP protein at subapical region of LREs.

(A) Pre-embedding silver-enhanced immunogold staining for Dr-VSP in LREs (top)

and IECs (bottom) of 14-dpf wild-type zebrafish larva. Right two images are the same as the

middle images but were highlighted with red and blue for microvilli and subapical region,

respectively. In LREs, immunoparticles for Dr-VSP are preferentially distributed at the

subapical region while rarely detectable at microvilli, consistent with results of Dr-VSP

immunofluorescence in Figure 3. In IECs, immunoparticles for Dr-VSP are rarely detectable

at both subapical region and microvilli, consistent with Dr-VSP immunofluorescence and

RNA-seq analysis in Figure 2. Scale bar = 2 µm.

(B) MDCKII cells co-expressing Dr-VSP-mCherry and (I) Rab5-EGFP for early

endosomes; or (II) Rab11-EGFP for recycling endosomes. Dr-VSP is localized at the

endosomal membranes of early and recycling endosomes. Red, Dr-VSP. Green, Rab5 or

Rab11. Scale bar = 20 µm.

Figure 5: Phenotypic investigation of Dr-VSP$^{-/-}$ zebrafish.

(A) Survival rates of wild-type and Dr-VSP$^{-/-}$ zebrafish larvae during the first month

of age. Zebrafish larvae, 300 at D0 for each zebrafish line. All statistical data were collected

from three independent experiments.

(B) Body length of wild-type and Dr-VSP$^{-/-}$ zebrafish larvae raised under non-calorie

restricted standard diet. Zebrafish larvae, ≥ 40 on the first day (D0) for each zebrafish line.

Error bars, means $\pm$ SD; *$P < 0.05$; **$P < 0.01$; unpaired Student's t-test.

(C) Brightfield images of the representative wild-type (top) and Dr-VSP$^{-/-}$ (bottom)

zebrafish larvae at 28 dpf, raised under the same conditions. Scale bar = 1 mm.

Figure 6: Functional analysis of Dr-VSP in larval enterocytes.

(A) Experimental approach: 6-dpf zebrafish larvae were gavaged with Alexa Fluor^{TM}

488-tagged dextran (fDex) or mCherry solutions into the anterior intestinal bulb. Live

imaging was performed at 2 hours after gavage.

(B - E) fDex and mCherry internationalized into zebrafish LREs after gavage. (B)

Live confocal images of wild-type (WT; left) and Dr-VSP$^{-/-}$ (right) zebrafish larvae showing

fDex internalization in enterocytes at the posterior portion of mid-intestine (arrowhead).

Scale bar = 200 µm

(C) Live confocal images of wild-type (WT; left) and Dr-VSP$^{-/-}$ (right) zebrafish

larvae showing mCherry internalization in enterocytes at the posterior portion of mid-

intestine (arrowhead). Scale bar = 200 µm

(D) Live confocal images of wild-type (left) and Dr-VSP$^{-/-}$ (right) enterocytes

showing fDex internalization. The images were presented with DIC (top) or without DIC

(bottom) to enhance visibility of the internalized fDex. Dotted lines conceptually indicate the

outlines of larval intestines. Scale bar = 20 µm

(E) Live confocal images of wild-type (left) and Dr-VSP$^{-/-}$ (right) enterocytes

showing mCherry internalization. The images were presented with DIC (top) or without DIC

(bottom) to enhance visibility of the internalized mCherry. Dotted lines conceptually indicate

the outlines of larval intestines. Scale bar = 20 µm

(F and G) Endocytosis efficiency analysis in LREs after fDex gavage. (F) Percentage

of the enterocytes containing fDex-filled lysosomal vacuoles at 2 hours following gavage.

Error bars, means ± SEM; ****$P < 0.0001$; unpaired Student's t-test. Number of enterocytes

= 344 cells from 14 wild-type and 359 cells from 13 Dr-VSP$^{-/-}$ zebrafish.

(G) Relative fDex internalization profile comparing between wild-type and Dr-VSP$^{-/-}$

zebrafish larvae at 2 hours following gavage. Data are means ± SEM percentage of

intracellular fDex intensity along the specified axes, with a 100% indicates the maximal

value of the average intensity on each axis. (Left) Apico-basal axis of individual cells.

(Right) Longitudinal axis along the posterior portion of mid-intestine. Number of fish = 14

719 for wild-type and 13 for Dr-VSP[-/-]. Data are presented as mean ± SEM, collected from 4

720 independent experiments for both wild-type and Dr-VSP[-/-].

721   (H and I) Endocytosis efficiency analysis in LREs after mCherry gavage. (H)

722 Percentage of the enterocytes containing mCherry-filled lysosomal vacuoles at 2 hours

723 following gavage. Error bars, means ± SEM; ****P < 0.0001; unpaired Student's t-test.

724 Number of enterocytes = 376 cells from 14 wild-type and 341 cells from 14 Dr-VSP[-/-]

725 zebrafish.

726   (I) Relative mCherry internalization profile comparing between wild-type and Dr-

727 VSP[-/-] zebrafish larvae at 2 hours following gavage. Data are means ± SEM percentage of

728 intracellular mCherry intensity along the specified axes, with a 100% indicates the maximal

729 value of the average intensity on each axis. (Left) Apico-basal axis of individual cells.

730 (Right) Longitudinal axis along the posterior portion of mid-intestine. Number of fish = 14

731 for each zebrafish line. Data are presented as mean ± SEM, collected from 4 independent

732 experiments for both wild-type and Dr-VSP[-/-].

734 **Figure 7: Morphological change in Dr-VSP[-/-] LREs.**

735   (A) Immunostaining showing cellular framework in sagittal sections of 14-dpf wild-

736 type (top) and Dr-VSP[-/-] (bottom) LREs (left); and their corresponding schematic illustration

737 (right). Green, F-actin. Red, Na^+-K^+ ATPase. Blue, DAPI. Scale bar = 10 µm.

738   (B) Cellular parameters of LREs at 14 dpf comparing between wild-type and Dr-VSP[-]

739 [/-] zebrafish larvae. (Left) Microvillus length. (Right) Cell height, apical width, and basal

740 width. Enterocytes, ≥ 120 cells from 5 larvae for each zebrafish line.

741   (C) Cellular parameters of LREs at 6 dpf comparing between wild-type and Dr-VSP[-/-]

742 zebrafish larvae. (Left) Microvillus length. (Right) Cell height, apical width, and basal width.

Enterocytes, ≥ 170 cells from 6 larvae for each zebrafish line. Error bars, means ± SD; ****P

< 0.0001; unpaired Student's t-test; ns, no statistically significant difference.

(D) Representative transverse sections at mid-intestine of 14-dpf wild-type (left) and

Dr-VSP$^{-/-}$ (right) zebrafish larvae, showing LREs with intracellular supranulcear vacuoles.

Sections were stained with toluidine blue. Scale bar = 20 µm.

(E – F) Vacuole parameters of 14-dpf LREs. (E) Vacuolar perimeter of wild-type

(WT) and Dr-VSP$^{-/-}$ LREs. **P = 0.0024; Mann-Whitney U test. Vacuoles, 108 for wild-type

and 95 for Dr-VSP$^{-/-}$. Data were collected from 3 larvae for each zebrafish line.

(F) Vacuolar area of wild-type (WT) and Dr-VSP$^{-/-}$ LREs from the same samples as in

Figure 6E. ***P = 0.0003; Mann-Whitney U test.

(G) Representative TEM images at mid-intestine of 14-dpf wild-type (left) and Dr-

VSP$^{-/-}$ (right) zebrafish LREs, showing the ultrastructures of microvilli and subapical region.

Key features of absorptive enterocytes are presented, including membrane invaginations at

inter-microvillous spaces (*), cytoplasmic tubules and tubule-vacuole complexes (arrowhead),

and numerous endocytic vesicles (Ve). In wild-type LRE, green represents the region upper

to supranuclear vacuole, which corresponds to the area with positive immunofluorescence

signal of Dr-VSP in Figure 3, A and B. Scale bar = 500 nm.

(H) Schematic diagram of wild-type (WT) and Dr-VSP$^{-/-}$ LREs. Dr-VSP (green) is

primarily localized in cytoplasm close to the apical surface of enterocytes, promoting

endosomal maturation during endocytosis-dependent nutrient absorption. N, nucleus. V,

Vacuoles.

Figure 8: Expression profile of Ci-VSP and Ci-Dab2 in ascidian *C. intestinalis*.

(A – C) Expression profile of Ci-VSP by WISH. (A) Whole-mount (left) 2-week-old

juvenile showing regional expressions in the stomach (st; arrow), bulged-region of the mid-

intestine (int; arrowhead), and posterior-intestine (gray arrowhead). Higher magnification

(right) reveals a stripe-like distribution of signals in the stomach and bulged intestine. (B)

Dissected post-pharyngeal region of the digestive tract of the 1.5-month-old young adult

showing an additional expression region (white arrowhead) in the looped-intestine. (C)

Transverse section specimens of the juvenile (white line in A) showing Ci-VSP expressions

in the epithelium of the digestive tract (left). Expression signals in the stomach epithelium are

restricted mainly in the inner-folds (white dotted lines), which contain several absorptive

enterocytes (38), as opposed to the outer-folds (dotted lines) with many pancreatic-related

exocrine genes expressions (39). Circumferences of the stomach (middle) and bulged

intestine (right) show alternating epithelial morphology and Ci-VSP expression.

(D–F) Expression profile Ci-Dab2. (D) Whole-mount (left) 2-week-old juvenile

showing regional expressions in the stomach (st; arrow), bulged-region of the mid-intestine

(int; arrowhead), and posterior-intestine (gray arrowhead). Higher magnification (right)

reveals a stripe-like distribution of signals in the stomach and bulged intestine. (E) Dissected

post-pharyngeal region of the digestive tract of the 1.5-month-old young adult showing an

additional expression region (white arrowhead) in the looped-intestine. (F) Transverse section

specimens of the juvenile (white line in D) showing Ci-Dab2 expressions in the epithelium of

the digestive tract (left). Expression signals in the stomach epithelium are restricted mainly in

the inner-folds (white dotted lines) as opposed to the outer-folds (dotted lines).

Circumferences of the stomach (middle) and bulged intestine (right) show alternating

epithelial morphology and Ci-Dab2 expression. Scale bars = 200 µm in (A, D), 1 mm in (B,

E), and 50 µm in (C, F). Additional abbreviations: os, oral siphon; as, atrial siphon; ph,

pharynx; en, endostyle; stg, stigma; es, esophagus; nc, neural complex; gd, gonad.

Figure legends for supplemental figures

Figure 1-figure supplement 1: Experimental schemes of CRISPR-Cas9-mediated

Tg(vsp:EGFP) **transgenesis**

Generation of *Tg(vsp:EGFP)* transgenic zebrafish. Donor DNA plasmid containing

the enhanced green fluorescence protein (EGFP) sequence was incorporated into *vsp* exon1,

which is located between transcription start site (exon1) and translation start site (exon3).

Successful transgenesis expresses EGFP recapitulating endogenous expression of *vsp* gene.

For detailed information, please refer to the Method Details, and Kimura et al (27). E1, exon

1. E2, exon2. hsp, hsp70 promoter. pA, polyA.

Figure 4-figure supplement 1: Spatial distribution of Dr-VSP within LREs.

Representative images of pre-embedding Dr-VSP immunogold staining in LREs from

14-dpf wild-type zebrafish larva, demonstrating the spatial distribution of Dr-VSP signals at

multiple areas within the cell. Dr-VSP is highly expressed at the subapical region (inset 1)

and gradually decreased in the middle (inset 2) and basal part (inset 3) of LREs,

corresponding to the immunostaining results in Figure 3. Scale bar = 2 µm. V, vacuole. N,

nucleus.

Figure 4-video supplement 1: Dr-VSP is localized at the endosomal membranes of early

endosomes.

MDCKII cells co-expressing Dr-VSP-mCherry and Rab5-EGFP for early endosomes.

Time-lapse images were acquired from the same samples as presented in Figure 4B. Red, Dr-

VSP. Green, Rab5. Scale bar = 20 µm.

816   **Figure 4-video supplement 2: Dr-VSP is localized at the endosomal membranes of**

817   **recycling endosomes.**

818   MDCKII cells co-expressing Dr-VSP-mCherry and Rab11-EGFP for recycling

819   endosomes. Dr-VSP is localized at the endosomal membranes of recycling endosomes. Time-

820   lapse images were acquired from the same samples as presented in Figure 4B. Red, Dr-VSP.

821   Green, Rab11. Scale bar = 20 μm.

823   **Figure 5-figure supplement 1: Experimental schemes of CRISPR-Cas9-mediated Dr-**

824   **VSP$^{-/-}$ mutagenesis.**

825   (A – E) Generation of heritable Dr-VSP$^{-/-}$ zebrafish. (A) Schematic diagram of the

826   zebrafish Dr-VSP-encoding gene (*vsp*), consisting of 21 exons. The CRISPR target with the

827   protospacer adjacent motif (PAM) sequence (TGG) is indicated in the box.

828   (B) Sequencing results illustrate microdeletion (arrowhead) in the genomic sequence

829   of Dr-VSP$^{-/-}$ zebrafish. WT, wild type.

830   (C) Alignment of genomic DNA sequences (top) and amino acid sequences (bottom)

831   comparing between wild-type and Dr-VSP$^{-/-}$ zebrafish. Microdeletion (red) within the

832   CRISPR target (gray shade) induced frameshift mutation in the subsequent amino acids of

833   Dr-VSP$^{-/-}$ zebrafish.

834   (D) Model of Dr-VSP illustrates the CRISPR-Cas9-mediated mutagenesis in

835   topological view. Mutation disrupted the functional domains of Dr-VSP located distal to the

836   CRISPR target.

837   (E) Immunostaining of wild-type and Dr-VSP$^{-/-}$ zebrafish. Anti-Dr-VSP antibody is

838   specific to the S4 transmembrane region and intracellular active center of the phosphatase

839   domain. Dr-VSP^{+} cell is undetectable in Dr-VSP$^{-/-}$ enterocytes. Green, Dr-VSP. Red, F-actin.

840   Blue, DAPI. Scale bar = 10 μm

Figure 6-figure supplement 1: Preliminary results of fDex-mCherry mixture

internalization into zebrafish LREs after gavage.

(A) Live confocal images of wild-type (WT) LREs showing fDex-mCherry mixture

internalization. Both fDex and mCherry signals are located in apical vesicles and

supranuclear vacuoles (arrowheads). Some mCherry signals did not overlap with fDex

signals (as indicated by asterisks).

(B) Live confocal images of Dr-VSP$^{-/-}$ LREs showing fDex-mCherry mixture

internalization. Notably, mCherry was transported deeply into some supranuclear vacuoles

(arrowheads), whereas fDex was mostly found in apical vesicles. Dotted lines indicate the

outlines of larval intestines. Many mCherry signals did not overlap with fDex signals (as

indicated by asterisks), indicating that mCherry and fDex were uptaken via different

mechanisms into distinct intracellular vesicles. The mismatch between mCherry and fDex

signals was more pronounced in Dr-VSP$^{-/-}$ than wild-type LREs. Scale bar = 20 µm.

Figure 7-figure supplement 1: Morphological structures of zebrafish LREs under

transmission electron microscopy.

(A) Representative TEM images (left) of LREs from 14-dpf wild-type (top) and Dr-

VSP$^{-/-}$ (bottom) zebrafish larvae; and their corresponding schematic illustration (right). Green

color represents the subapical region upper to supranuclear vacuoles, where the positive

immunofluorescence signal of Dr-VSP was observed in Figure 3, A and B. N, nucleus. V,

vacuole. Scale bar = 10 µm.

(B) Representative TEM images of LREs from 14-dpf wild-type (top) and Dr-VSP$^{-/-}$

(bottom) zebrafish larvae, showing the ultrastructures of microvilli and subapical region. Key

features of absorptive enterocytes are presented, including membrane invaginations at inter-

microvillous spaces, cytoplasmic tubules and tubule-vacuole complexes, and numerous

endocytic vesicles. In wild-type LRE, green color represents the subapical region upper to

supranuclear vacuole, where immunofluorescence signal of Dr-VSP was observed in Figure 3,

A and B. Scale bar = 2 μm (left) and 500 nm (right).

(C) Representative TEM images of LREs from 14-dpf wild-type (left) and Dr-VSP$^{-/-}$

(right) zebrafish larvae, showing the ultrastructures of microvilli and subapical region. Red

color represents the area beneath apical surface that contains cytoplasmic tubules but few

well-defined endocytic vesicles. The vesicles are more densely distributed in the areas deeper

than the red region and are frequently associated with larger endosomal vacuoles.

(D) Inter-edge distance between apical surface and vesicle-dense area in wild-type

and Dr-VSP$^{-/-}$ LREs, corresponding to the width of red shading areas shown in (C).

Enterocytes, > 50 cells from 25 TEM images (magnification = 15000x) for each zebrafish

line. Data were collected from 4 wild-types and 3 Dr-VSP$^{-/-}$ larvae. Average distances in each

TEM image were analyzed using ImageJ macro.

(E) Microvillous length of wild-type and Dr-VSP$^{-/}$ LREs measuring at TEM level

from the same samples as in (D). Error bars, means ± SD; ****P < 0.0001; unpaired

Student's t-test; ns, no statistically significant difference.

Figure 7-figure supplement 2: TEM image series of wild-type and Dr-VSP$^{-/-}$ LREs.

Representative TEM image series of LREs from 14-dpf wild-type (top) and Dr-VSP$^{-/-}$

(bottom) zebrafish larvae. Large vacuoles are distributed in the supranuclear region but are

fewer in Dr-VSP$^{-/-}$ LREs. Membrane invaginations at inter-microvillous spaces (*) can be

observed at higher magnification. Branching cytoplasmic tubules and tubule-vacuole

complexes (arrowheads) are more common in wild-type LREs. Scale bar at 6000x = 10 μm;

at 15000x = 2 μm; at 40000x = 500 nm.

Figure 8-figure supplement 1:

893   Multiple amino acid alignment of full-length VSP orthologs from human (*Homo*

sapiens, Hs-VSP), mouse (*Mus musculus*, Mm-VSP), zebrafish (*Danio rerio*, Dr-VSP),

common carp (*Cyprinus carpio*, Cc-VSP), Nile tilapia (*Oreochromis niloticus*, On-VSP),

Japanese medaka (*Oryzias latipes*, Ol-VSP), sea squirt (*Ciona intestinalis* Type A, Ci-VSP),

and Crown-of-thorns starfish (*Acanthaster planci*, Ap-VSP). Boxes indicate the putative

transmembrane helices (S1 - S4), VSD-PD linker, WPD loop, CX5R(T/S), gating loop, 515

loop/CBR3 loop (7). Amino acid sequences were retrieved from National Center for

Biotechnology Information (NCBI) public database. Analyses were performed using online

Constraint-based Multiple Alignment Tool (COBALT)

(https://www.ncbi.nlm.nih.gov/tools/cobalt/)

Figure 8-figure supplement 2:

905   Molecular phylogenetic tree showing the phylogenic relationships among VSP

orthologs from **mammals:** human (*Homo sapiens*, Hs-VSP1 and Hs-VSP2), mouse (*Mus*

musculus, Mm-VSP), rat (*Rattus norvegicus*, Rn-VSP), dog (*Canis lupus familiaris*, Clf-

VSP), cat (*Felis catus*, Fc-VSP); **amphibian:** American clawed frog (*Xenopus laevis*, Xl-

VSP1 and Xl-VSP2); **teleosts:** goldfish (*Carassius auratus*, Ca-VSP), common carp

(*Cyprinus carpio*, Cc-VSP), zebrafish (*Danio rerio*, Dr-VSP), channel catfish (*Ictalurus*

punctatus, Ip-VSP), Chinook salmon (*Oncorhynchus tshawytscha*, Ots-VSP), Atlantic

salmon (*Salmo salar*, Ss-VSP), Siamese fighting fish (*Betta splendens*, Bs-VSP), Japanese

puffer (*Takifugu rubripes*, Tr-VSP), turquoise killifish (*Nothobranchius furzeri*, Nf-VSP),

Japanese medaka (*Oryzias latipes*, Ol-VSP), Nile tilapia (*Oreochromis niloticus*, On-VSP);

invertebrates: sea squirt (*Ciona intestinalis* Type A, Ci-VSP), crown-of-thorns starfish

(*Acanthaster planci*, Ap-VSP), fresh-water polyp (*Hydra vulgaris*, Hv-VSP), sea anemone

(*Exaiptasia diaphana*, Ed-VSP), and smooth cauliflower coral (*Stylophora pistillata*, Spi-

VSP). Data were retrieved from the same source as Supplemental Figure S4, and analyzed

using Molecular Evolutionary Genetics Analysis (MEGA) X Software (Pennsylvania State

University, Pennsylvania, US).

Figure 8-figure supplement 3: Morphological structure of ascidian intestine and

expression profiles of intestine-related genes

(A – D) Expression profiles of intestine-related genes in *Ciona* juvenile (1st column)

and young adult (2nd column) revealed by WISH. In the juvenile, whole (left panel) and

magnificent of the post-pharyngeal region of the digestive tract including esophagus, stomach,

and intestine (right panel) were shown.

(A) Morphological structure of the digestive tract was noted on the expressions of

Mucin gene. Digestive tract of *C. intestinalis* Type A was divided into oral siphon (os),

pharynx (ph), esophagus (es), stomach (st), and intestine (int) (69). The intestine was

subdivided into four regions, R1 to R4 (70). Mucin gene was expressed broadly in the post-

pharyngeal region of the digestive tract of the juvenile, but not expressed in the stomach and

bulged-intestine (R3). In the young adult specimen, expression profile was similar to the

juvenile, but expression signal was disappeared between anterior intestine (R1) and looped-

intestine (R2) beside the developing gonad (gd).

(B) Absorption-related PEPT1 gene was invested as an example of the peptide

transporter genes. Expression signals of the *PEPT1* were observed in stomach (arrow) and

bulged-intestine (R3) of the mid-intestine (arrowhead) in both developmental stages of

juvenile and young adult. Additional expression (white arrowhead) was appeared between

anterior intestine (R1) and looped-intestine (R2) of the young adult intestine.

(C) Absorption-related SGLT1 gene was invested as a secondary active glucose

transporter. Transcripts of the *SGLT1* were detected in stomach (arrow), bulged-intestine

(R3) (arrowhead), and posterior-intestine (R4) (gray arrowhead) in juvenile and young adult

specimens.

(D) GLUT5 gene was also assess as a facilitated transporter of the monosaccharide.

Transcripts of the *GLUT5* were detected in stomach (arrow), bulged-intestine (R3)

(arrowhead), and posterior-intestine (R4) (gray arrowhead) in juvenile and young adult

specimens. Additional expression (white arrowhead) between anterior intestine (R1) and

looped-intestine (R2) was also found in the young adult. Scale bars = 200 µm in (left) and 1

mm in (right). Additional abbreviations: as, atrial siphon; en, endostyle. Latest gene model

IDs (HT version) of the *Ciona Mucin*, *PEPT1*, *SGLT1*, and *GLUT5* are KY.Chr2.2226,

KY.Chr13.370, KY.Chr10.1303, and KY.Chr7.810, respectively (71)

Table 1: RT-PCR Primers used in this study

Table 2: CRISPR target sequences of *vsp* and PCR primers used in this study

References (Vancouver)

1.   Falkenburger BH, Jensen JB, Dickson EJ, Suh BC, Hille B. Phosphoinositides: lipid

regulators of membrane proteins. J Physiol. 2010;588(Pt 17):3179-85.

2.   Jones DR, Ramirez IB, Lowe M, Divecha N. Measurement of phosphoinositides in

the zebrafish *Danio rerio*. Nat Protoc. 2013;8(6):1058-72.

3.   Di Paolo G, De Camilli P. Phosphoinositides in cell regulation and membrane

dynamics. Nature. 2006;443(7112):651-7.

4.   Li X, Garrity AG, Xu H. Regulation of membrane trafficking by signalling on

endosomal and lysosomal membranes. J Physiol. 2013;591(18):4389-401.

5.   Wallroth A, Haucke V. Phosphoinositide conversion in endocytosis and the

endolysosomal system. J Biol Chem. 2018;293(5):1526-35.

6.   Murata Y, Iwasaki H, Sasaki M, Inaba K, Okamura Y. Phosphoinositide phosphatase

activity coupled to an intrinsic voltage sensor. Nature. 2005;435(7046):1239-43.

7.   Okamura Y, Kawanabe A, Kawai T. Voltage-Sensing Phosphatases: Biophysics,

Physiology, and Molecular Engineering. Physiol Rev. 2018;98(4):2097-131.

8.   Okamura Y, Murata Y, Iwasaki H. Voltage-sensing phosphatase: actions and

potentials. J Physiol. 2009;587(3):513-20.

9.   Keum D, Kruse M, Kim DI, Hille B, Suh BC. Phosphoinositide 5- and 3-phosphatase

activities of a voltage-sensing phosphatase in living cells show identical voltage dependence.

Proc Natl Acad Sci U S A. 2016;113(26):E3686-95.

10.   Wu Y, Dowbenko D, Pisabarro MT, Dillard-Telm L, Koeppen H, Lasky LA. PTEN 2,

a Golgi-associated testis-specific homologue of the PTEN tumor suppressor lipid

phosphatase. J Biol Chem. 2001;276(24):21745-53.

11.   Kawai T, Miyata H, Nakanishi H, Sakata S, Morioka S, Sasaki J, et al. Polarized

PtdIns(4,5)P$_2$ distribution mediated by a voltage-sensing phosphatase (VSP) regulates sperm

motility. Proc Natl Acad Sci U S A. 2019;116(51):26020-8.

12.   Tapparel C, Reymond A, Girardet C, Guillou L, Lyle R, Lamon C, et al. The TPTE

gene family: cellular expression, subcellular localization and alternative splicing. Gene.

2003;323(1-2):189-99.

13.   Ogasawara M, Sasaki M, Nakazawa N, Nishino A, Okamura Y. Gene expression

profile of Ci-VSP in juveniles and adult blood cells of ascidian. Gene Expr Patterns.

2011;11(3-4):233-8.

14.   Neuhaus H, Hollemann T. Kidney specific expression of cTPTE during development

of the chick embryo. Gene Expr Patterns. 2009;9(8):568-71.

15.   Mutua J, Jinno Y, Sakata S, Okochi Y, Ueno S, Tsutsui H, et al. Functional diversity

of voltage-sensing phosphatases in two urodele amphibians. Physiol Rep. 2014;2(7):e12061-e.

16.   Ratzan W, Rayaprolu V, Killian SE, Bradley R, Kohout SC. The voltage sensing

phosphatase (VSP) localizes to the apical membrane of kidney tubule epithelial cells. PLoS

One. 2019;14(4):e0209056.

17.   Yamaguchi S, Aoki N, Kitajima T, Okamura Y, Homma KJ. Expression of the

voltage-sensing phosphatase gene in the chick embryonic tissues and in the adult cerebellum.

Commun Integr Biol. 2014;7(5).

18.   Hossain MI, Iwasaki H, Okochi Y, Chahine M, Higashijima S, Nagayama K, et al.

Enzyme domain affects the movement of the voltage sensor in ascidian and zebrafish

voltage-sensing phosphatases. J Biol Chem. 2008;283(26):18248-59.

19.   Kawanabe A, Mizutani N, Polat OK, Yonezawa T, Kawai T, Mori MX, et al.

Engineering an enhanced voltage-sensing phosphatase. J Gen Physiol. 2020;152(5):1-11.

20.   Gut P, Reischauer S, Stainier DYR, Arnaout R. Little fish, big data: zebrafish as a

model for cardiovascular and metabolic disease. Physiol Rev. 2017;97(3):889-938.

21.   Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of

embryonic development of the zebrafish. Dev Dyn. 1995;203(3):253-310.

22.   Vacaru AM, Unlu G, Spitzner M, Mione M, Knapik EW, Sadler KC. In vivo cell

biology in zebrafish - providing insights into vertebrate development and disease. J Cell Sci.

2014;127(Pt 3):485-95.

23.   Park J, Levic DS, Sumigray KD, Bagwell J, Eroglu O, Block CL, et al. Lysosome-

rich enterocytes mediate protein absorption in the vertebrate gut. Dev Cell. 2019;51(1):7-20

e6.

24.   Iwai T. Fine structure of gut epithelial cells of larval and juvenile carp during

absorption of fat and protein. Arch Histol Jpn. 1969;30(2):183-99.

25.   Stroband HWJ, van der Veen FH. Localization of protein absorption during transport

of food in the intestine of the grasscarp, *Ctenopharyngodon idella* (Val.). Journal of

Experimental Zoology. 1981;218(2):149-56.

26.   Rombout JH, Lamers CH, Helfrich MH, Dekker A, Taverne-Thiele JJ. Uptake and

transport of intact macromolecules in the intestinal epithelium of carp (*Cyprinus carpio L.*)

and the possible immunological implications. Cell Tissue Res. 1985;239(3):519-30.

27.   Kimura Y, Hisano Y, Kawahara A, Higashijima S. Efficient generation of knock-in

transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-mediated genome

engineering. Sci Rep. 2014;4:6545.

28.   Ng AN, de Jong-Curtain TA, Mawdsley DJ, White SJ, Shin J, Appel B, et al.

Formation of the digestive system in zebrafish: III. Intestinal epithelium morphogenesis. Dev

Biol. 2005;286(1):114-35.

29.   Rodriguez-Fraticelli AE, Bagwell J, Bosch-Fortea M, Boncompain G, Reglero-Real N,

Garcia-Leon MJ, et al. Developmental regulation of apical endocytosis controls epithelial

patterning in vertebrate tubular organs. Nat Cell Biol. 2015;17(3):241-50.

30.   Parton RG. Endocytosis in MDCK Cells. 1992:317-24.

31.   Wallace KN, Akhter S, Smith EM, Lorent K, Pack M. Intestinal growth and

differentiation in zebrafish. Mech Dev. 2005;122(2):157-73.

32.   Cocchiaro JL, Rawls JF. Microgavage of zebrafish larvae. J Vis Exp. 2013(72):e4434.

33.   Sidhaye J, Pinto CS, Dharap S, Jacob T, Bhargava S, Sonawane M. The zebrafish

goosepimples/myosin Vb mutant exhibits cellular attributes of human microvillus inclusion

disease. Mech Dev. 2016;142:62-74.

34.   Iida H, Shibata Y, Yamamoto T. The endosome-lysosome system in the absorptive

cells of goldfish hindgut. Cell and Tissue Research. 1986;243(2):449--52.

35.   Iida H, Yamamoto T. Intracellular transport of horseradish peroxidase in the

absorptive cells of goldfish hindgut in vitro, with special reference to the cytoplasmic tubules.

Cell and Tissue Research. 1985;240(3):553-60.

36.   Nakayama S, Sekiguchi T, Ogasawara M. Molecular and evolutionary aspects of the

protochordate digestive system. Cell Tissue Res. 2019;377(3):309-20.

37.   Yonge CM. Studies on the comparative physiology of digestion. Journal of

Experimental Biology. 1925;2(3):373-88.

38.   Hayashibe M, Nakayama S, Ogasawara M. Shared hemocyte- and intestine-dominant

expression profiles of intelectin genes in ascidian *Ciona intestinalis*: insight into the

evolution of the innate immune system in chordates. Cell Tissue Res. 2017;370(1):129-42.

39.   Nakayama S, Ogasawara M. Compartmentalized expression patterns of pancreatic-

and gastric-related genes in the alimentary canal of the ascidian *Ciona intestinalis*:

evolutionary insights into the functional regionality of the gastrointestinal tract in Olfactores.

Cell Tissue Res. 2017;370(1):113-28.

40.   Salinas I, Parra D. Fish mucosal immunity: Intestine: Elsevier Inc.; 2015. 135--70 p.

41.   Lu JW, Ho YJ, Ciou SC, Gong Z. Innovative disease model: zebrafish as an in vivo

platform for intestinal disorder and tumors. Biomedicines. 2017;5(4):1-17.

42.   Ratzan WJ, Evsikov AV, Okamura Y, Jaffe LA. Voltage sensitive phosphoinositide

phosphatases of *Xenopus*: their tissue distribution and voltage dependence. J Cell Physiol.

2011;226(11):2740-6.

43.   Yamaguchi S, Kurokawa T, Taira I, Aoki N, Sakata S, Okamura Y, et al. Potential

role of voltage-sensing phosphatases in regulation of cell structure through the production of

PI(3,4)P2. J Cell Physiol. 2014;229(4):422-33.

44.   Zoncu R, Perera RM, Balkin DM, Pirruccello M, Toomre D, De Camilli P. A

phosphoinositide switch controls the maturation and signaling properties of APPL endosomes.

Cell. 2009;136(6):1110-21.

45.   Wang H, Lo WT, Haucke V. Phosphoinositide switches in endocytosis and in the

endolysosomal system. Curr Opin Cell Biol. 2019;59:50-7.

46.   Vicinanza M, Di Campli A, Polishchuk E, Santoro M, Di Tullio G, Godi A, et al.

OCRL controls trafficking through early endosomes via PtdIns4,5P_2-dependent regulation of

endosomal actin. EMBO J. 2011;30(24):4970-85.

47.   Billcliff PG, Lowe M. Inositol lipid phosphatases in membrane trafficking and human

disease. Biochem J. 2014;461(2):159-75.

48.   Erdmann KS, Mao Y, McCrea HJ, Zoncu R, Lee S, Paradise S, et al. A role of the

Lowe syndrome protein OCRL in early steps of the endocytic pathway. Dev Cell.

2007;13(3):377-90.

49.   Cauvin C, Rosendale M, Gupta-Rossi N, Rocancourt M, Larraufie P, Salomon R, et al.

Rab35 GTPase triggers switch-like recruitment of the Lowe Syndrome lipid phosphatase

OCRL on newborn endosomes. Curr Biol. 2016;26(1):120-8.

50.   Matsuda M, Takeshita K, Kurokawa T, Sakata S, Suzuki M, Yamashita E, et al.

Crystal structure of the cytoplasmic phosphatase and tensin homolog (PTEN)-like region of

Ciona intestinalis voltage-sensing phosphatase provides insight into substrate specificity and

redox regulation of the phosphoinositide phosphatase activity. J Biol Chem.

2011;286(26):23368-77.

51.   Gerlach GF, Wingert RA. Kidney organogenesis in the zebrafish: insights into

vertebrate nephrogenesis and regeneration. Wiley Interdiscip Rev Dev Biol. 2013;2(5):559-

85.

52. Li P, Gu M, Xu H. Lysosomal ion channels as decoders of cellular signals. Trends

Biochem Sci. 2019;44(2):110-24.

53. Cang C, Bekele B, Ren D. The voltage-gated sodium channel TPC1 confers

endolysosomal excitability. Nat Chem Biol. 2014;10(6):463-9.

54. Wie J, Liu Z, Song H, Tropea TF, Yang L, Wang H, et al. A growth-factor-activated

lysosomal K^+ channel regulates Parkinson's pathology. Nature. 2021;591(7850):431-7.

55. Maurer ME, Cooper JA. Endocytosis of megalin by visceral endoderm cells requires

the Dab2 adaptor protein. J Cell Sci. 2005;118(Pt 22):5345-55.

56. Mishra SK, Keyel PA, Hawryluk MJ, Agostinelli NR, Watkins SC, Traub LM.

Disabled-2 exhibits the properties of a cargo-selective endocytic clathrin adaptor. EMBO J.

2002;21(18):4915-26.

57. Hartenstein V, Martinez P. Phagocytosis in cellular defense and nutrition: a food-

centered approach to the evolution of macrophages. Cell Tissue Res. 2019;377(3):527-47.

58. Iida A, Sano K, Inokuchi M, Nomura J, Suzuki T, Kuriki M, et al. Cubam receptor-

mediated endocytosis in hindgut-derived pseudoplacenta of a viviparous teleost (*Xenotoca*

eiseni). J Exp Biol. 2021;224(13).

59. Gonnella PA, Neutra MR. Membrane-bound and fluid-phase macromolecules enter

separate prelysosomal compartments in absorptive cells of suckling rat ileum. J Cell Biol.

1984;99(3):909-17.

60. Kraehenbuhl JP, Campiche MA. Early stages of intestinal absorption of specific

antibiodies in the newborn. An ultrastructural, cytochemical, and immunological study in the

pig, rat, and rabbit. J Cell Biol. 1969;42(2):345-65.

61. Brugman S. The zebrafish as a model to study intestinal inflammation. Dev Comp

Immunol. 2016;64:82-92.

62.   Kleinman RE, Walker WA. Antigen processing and uptake from the intestinal tract.

Clin Rev Allergy. 1984;2(1):25-37.

63.   Lickwar CR, Camp JG, Weiser M, Cocchiaro JL, Kingsley DM, Furey TS, et al.

Genomic dissection of conserved transcriptional regulation in intestinal epithelial cells. PLoS

Biol. 2017;15(8):e2002054.

64.   Ogasawara M, Satoh N, Shimada Y, Wang Z, Tanaka T, Noji S. Rapid and stable

buffer exchange system using InSitu Chip suitable for multicolor and large-scale whole-

mount analyses. Dev Genes Evol. 2006;216(2):100-4.

65.   Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a

CRISPR/Cas9 and TALEN web tool for genome editing. Nucleic Acids Res. 2014;42(Web

Server issue):W401-7.

66.   Kotani H, Taimatsu K, Ohga R, Ota S, Kawahara A. Efficient multiple genome

modifications induced by the crRNAs, tracrRNA and Cas9 protein complex in zebrafish.

PLoS One. 2015;10(5):e0128319.

67.   Meeker ND, Hutchinson SA, Ho L, Trede NS. Method for isolation of PCR-ready

genomic DNA from zebrafish tissues. Biotechniques. 2007;43(5):610, 2, 4.

68.   Ge SX, Son EW, Yao R. iDEP: an integrated web application for differential

expression and pathway analysis of RNA-Seq data. BMC Bioinformatics. 2018;19(1):534.

69.   Chiba S, Sasaki A, Nakayama A, Takamura K, Satoh N. Development of *Ciona*

intestinalis juveniles (through 2nd ascidian stage). Zoolog Sci. 2004;21(3):285-98.

70.   Yoshida R, Sasakura Y. Establishment of enhancer detection lines expressing GFP in

the gut of the ascidian *Ciona intestinalis*. Zoolog Sci. 2012;29(1):11-20.

71.   Satou Y, Nakamura R, Yu D, Yoshida R, Hamada M, Fujie M, et al. A nearly

complete genome of *Ciona intestinalis* Type A (*C. robusta*) reveals the contribution of

inversion to chromosomal evolution in the genus *Ciona*. Genome Biol Evol.

2019;11(11):3144-57.

1137

Figure 1

(A)

(B)

(C)

(D)

Tg(vsp:EGFP)

(E)

Tg(vsp:EGFP)

vsp:EGFP DAPI

Figure 2

(A)

(B)

(C)

(D)

(E)

Figure 3

(A)

(B)

(C)

Figure 4

(A)

(B)

I Rab5 (early endosomes)

II Rab11 (recycling endosomes)

Figure 5

(A)

(B)

(C) 28-dpf Zebrafish

Figure 6

Figure 7

(A)

F-actin Na⁺-K⁺-ATPase
WT

F-actin Na⁺-K⁺-ATPase
Dr-VSP⁻/⁻

(B) LRE parameters of 14-dpf larvae

(C) LRE parameters of 6-dpf larvae

(D) 14 dpf — WT / Dr-VSP⁻/⁻

(E) Vacuolar perimeter (μm)

(F) Vacuolar area (μm²)

(G) 14 dpf — WT / Dr-VSP⁻/⁻

(H) LREs — WT / Dr-VSP⁻/⁻

Figure 8