Research Article

Pulvinar Slits: Cellulose-deficient and De-esterified Homogalacturonan-rich Structures in a Legume Motor Cell

Masahiro Takahara[1*], Satoru Tsugawa[2,3], Shingo Sakamoto[4,5],

Taku Demura[2,6], Miyuki T Nakata[2,6*†]

*These authors contributed equally.

†For correspondence. E-mail: mtn.plant@bs.naist.jp

[1]Acacia Horticulture, Kizugawa, Kyoto, Japan

[2]Graduate School of Science and Technology, Nara Institute of Science and Technology (NAIST), Ikoma, Nara, Japan

[3]Department of Mechanical Engineering, Faculty of Systems Science and Technology, Akita Prefectural University, Akita, Japan

[4]Plant Gene Regulation Research Group, Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[5]Global Zero Emission Research Center, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[6]Center for Digital Green-innovation, Nara Institute of Science and Technology (NAIST), Ikoma, Nara, Japan

Short running title

Pulvinar Slits in a Legume Motor Cell

Affiliations

Acacia Horticulture, Kizugawa, Kyoto, Japan

Masahiro Takahara

Graduate School of Science and Technology, Nara Institute of Science and Technology (NAIST),

Ikoma, Nara, Japan

Department of Mechanical Engineering, Faculty of Systems Science and Technology, Akita

Prefectural University, Akita, 015-0055, Japan

Satoru Tsugawa

Plant Gene Regulation Research Group, Bioproduction Research Institute, National Institute of

Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

Global Zero Emission Research Center, National Institute of Advanced Industrial Science and

Technology (AIST), Tsukuba, Japan

Shingo Sakamoto

Graduate School of Science and Technology, Nara Institute of Science and Technology (NAIST),

Ikoma, Nara, Japan

Center for Digital Green-innovation, Nara Institute of Science and Technology (NAIST), Ikoma, Nara,

Japan

Taku Demura, Miyuki T Nakata

Corresponding author

Miyuki T Nakata

Nara Institute of Science and Technology

Graduate School of Science and Technology

Division of Biological Science

8916-5 Takayama, Ikoma, NARA 630-0192 JAPAN

E-mail: mtn.plant@bs.naist.jp

Abstract

The cortical motor cells (CMCs) in a legume pulvinus execute the reversible deformation in leaf movement that is driven by changes in turgor pressure. In contrast to the underlying osmotic regulation, the cell wall structure of CMCs that contributes to the movement has yet to be characterized in detail. Here, we report that the cell wall of CMCs has tangential slits with low levels of cellulose deposition. This structure is unique and distinct from any other primary cell wall reported so far; thus, we named them "pulvinar slits." Notably, we predominantly detected de-methyl esterified homogalacturonan inside pulvinar slits, with a low deposition of highly methyl-esterified homogalacturonan inside the slits, as with cellulose. In addition, Fourier-transform infrared spectroscopy analysis indicated that the cell wall composition of pulvini is different from that of other axial organs, i.e. petioles or stems. Moreover, monosaccharide analysis showed that pulvini are pectin-rich organs like developing stems. Computer modeling suggested that pulvinar slits facilitate anisotropic extension in the direction perpendicular to the slits in the presence of turgor pressure. Mechanical tests revealed that the proximo-distal extensibility of pulvini is superior to that of the other axial organs. In this study, we thus established the distinctive cell wall structure of CMCs that contribute to plant movement, adding new knowledge to the structural diversity and function of the plant cell wall.

Abbreviations

CMC, a cortical motor cell; FTIR, Fourier-transform infrared spectroscopy; PCW, primary cell wall; SCW, secondary cell wall; HG, homogalacturonan; PD, proximo-distal; CBM3a, a carbohydrate-binding module 3; FEM, the finite-element method; PME, pectin methylesterases.

Introduction

A legume pulvinus is a joint-like thickening at the base of a leaf, leaflet, or rachis that plays a central role in leaf movement. Pulvinus-actuated leaf movements include circadian rhythm-driven movements completed over the scale of hours (Darwin and Darwin, 1880), blue light-induced movements in minutes (Koller, 1990), and touch-stimulated movements of Mimosoideae in seconds (Weintraub, 1952). During movement, a pulvinus is repeatedly and reversibly deformed by changing cell volume due to variation in the osmotic gradient. These repetitive and reversible steps are a unique feature that distinguish pulvinus-based movements from other types of organ movement driven by irreversible growth or by mechanical instability (Dumais and Forterre, 2012; Geitmann, 2016).

In general, osmotic gradients between the inside and the outside of a cell trigger directional water transport resulting in a change in cell volume. Cells swell when exposed to a hypotonic solution, while they contract in a hypertonic solution. Importantly, excessive expansion or contraction is blocked by the physical scaffold provided by the cell wall. When the cell volume increases in a hypotonic solution, cell walls act as a load-bearing structure against tensile stress. Therefore, the time derivative of cell volume depends on turgor pressure or cell wall extensibility under variable osmotic conditions, unless plasmolysis occurs (Geitmann and Ortega, 2009). In growing cells, cell walls are weak enough to yield under turgor, yet strong enough to offer mechanical resistance against bursting (Smithers et al., 2019).

The importance of the cell wall has been thoroughly investigated during the irreversible extension of growing cells (Cosgrove, 2005). Growing cells are surrounded by a primary cell wall (PCW) consisting mainly of cellulose, xyloglucan, and pectin, and they lack a secondary cell wall (SCW). Among PCW components, multiple cellulose polymers are bundled into microscale fibers, namely, cellulose microfibrils, to provide tensile stiffness and strength to the PCW. In axially expanding cells, the orientation of cellulose microfibrils often exhibits a bias perpendicular to the direction of expansion. In addition, xyloglucan and pectin contribute to cell wall extensibility by interacting with cellulose microfibrils (Cosgrove, 2005; Smithers et al., 2019). Xyloglucan crosslinks the hydrophobic segments of cellulose microfibrils to form a complex network, while pectin is a component of the matrix that fills the space between cellulose microfibrils by binding to their hydrophilic fraction (Cosgrove, 2005; Geitmann and Ortega, 2009; Zheng et al., 2018; Cosgrove, 2018). Thus, how the orientation, deposition, and structure of the cellulose microfibril network control cell wall extensibility and irreversible growth is well established and supported.

Unlike organ movement due to irreversible growth, pulvinus-driven movements are independent from growth (Darwin and Darwin, 1880). The pulvinus is comprised of several mature tissues, including the epidermis, cortical motor cells (CMCs), and vascular tissues (Fleurat-Lessard and Bonnenmain, 1978; Fleurat-Lessard and Roblin, 1982; Moysset and Simon, 1991). CMCs have a central role in the reversible deformation of the pulvinus. The CMCs that shrink when the pulvinus flexes are called extensor motor cells, while CMCs on the opposite side are called flexor cells. In the

early 1900s, changes in turgor of CMCs were proposed to be critical for leaf movements (Weintraub,

1952). Later, studies using runner bean (*Phaseolus coccineus*) reported anisotropic extension to the

proximo-distal (PD) direction in an intact pulvinus during nyctinastic movement and in slices of CMCs

exposed to different concentrations of mannitol (Mayer et al., 1985). Based on an estimation of the

bulk moduli of extensibility of CMCs, this work concluded that the osmotic concentration in the

apoplast determines by how much the CMC volume will change and actuate leaf movement. Therefore,

the changes in osmotic concentration of CMCs have received much recent attention in various legume

species with pulvini (Moran, 2007).

In this study, we focused on the structure of the CMC cell wall, which is not well characterized in

contrast to the osmotic regulation of CMCs. Cellulose microfibrils are "tangentially" aligned in the

cell wall of *P. coccineus* CMCs (Mayer et al., 1985); that is, they are aligned perpendicular to the PD

axis. Indeed, the orientation of cellulose microfibrils appears to be consistent with the extension

anisotropy of slices of CMCs as described above. However, the shape of each CMC does not show

the same bias relative to the orientation perpendicular to that of cellulose microfibrils as seen in the

case of irreversible growth. In addition, the orientation of cellulose microfibrils is closely associated

with a slow irreversible extension referred to as "creep," but it is unclear whether the orientation alone

is sufficient to explain the relatively rapid and reversible deformation of CMCs during leaf movement.

In this study, we investigated the cell wall structure of CMCs in detail and discovered a hitherto

undescribed slit-like structure that we designated pulvinar slits. Computational simulations suggested

that the presence of pulvinar slits promotes extension in the PD direction, which we tested empirically

by performing a mechanical test to compare the extensibility in the PD direction in pulvini with

pulvinar slits and other axial organs lacking the slits. Our data suggest that the function of pulvinar

slits is associated with PD extensibility and mechanical reversibility of CMCs.

Results

CMCs have tangential slits with less cellulose

To identify the cell wall structure of CMCs, we stained pulvini with calcofluor white, which stains

cell wall components such as cellulose, in longitudinal and cross sections using pulvini freshly

harvested from panicled-leaf ticktrefoil (*Desmodium paniculatum*), belonging to the Papilionoideae

family (Figure 1A-F). Notably, in CMC longitudinal sections, we observed several slits with no

significant fluorescence when stained by calcofluor white that run "tangentially" (Figure 1C, G). In

vertical sections, we noticed small spots with no staining but no slits (Figure 1D, G). To determine

whether tangential slits are specific to CMCs, we repeated the staining protocol on cortical cells

collected from the boundary between a pulvinus and a petiole and from a longitudinal section of a

petiole (Figure 1E, F). In both cases, we visualized small spots lacking staining but did not see

tangential slits (Figure 1E, F). Since we only observed these tangential slits in pulvinus CMCs, we

designated the tangential slits as "pulvinar slits."

To investigate the extent of conservation of pulvinar slits among legumes, we stained longitudinal

sections of pulvini from various legume species with calcofluor white. The Fabaceae family comprises

two major subfamilies: the Papilionoideae and the Caesalpinioideae (Azani et al., 2017). Importantly,

we detected slit structures in pulvinar CMCs from all 12 species selected and tested from these two

subfamilies (Supplemental Table 1, Figure 1H). In addition, we observed pulvinar slits in both adaxial

and abaxial pulvinar CMCs (Supplemental Figure 1). To make sure that the slits were not artifacts

arising during sectioning, we performed whole-mount staining. Again, we clearly observed slits in

CMCs from whole-mount-stained samples, with the slits located at the cell periphery (Figure 1I). We

therefore concluded that pulvinar slits are an innate cell wall structure of CMCs.

To validate the observation that a pulvinar slit is a region with lower cellulose deposition than its

surrounding region, we used cellulose-specific dyes, namely, pontamine and immunolabeling with an

antibody against type A carbohydrate-binding module 3 (CBM3a). With pontamine staining, we

detected tangential slits similar to those seen with calcofluor white staining in pulvinar CMCs (Figure

2A). Immunolabeling with CBM3a predominantly stains crystalline cellulose as speckles, but we also

found slit-like structures with lower signals (Figure 2B). Double staining with CBM3a and calcofluor

white indicates that slits lacking the CBM3a signal were colocalized with pulvinar slits, the structures

unstained by calcofluor white (Figure 2B, arrowheads). These observations indicated that pulvinar

slits are regions characterized by an extremely low cellulose deposition.

Taken together, we concluded that pulvinar slits are a structure specifically formed on the PCW of

CMCs, they have less cellulose than the surrounding cell wall, and they are widely conserved among

legume species.

De-methyl esterification of homogalacturonan is associated with pulvinar slits

To further characterize pulvinar slits, we examined the spatial pattern of other major PCW

components by immuno-histological analysis. Homogalacturonan (HG) is the most abundant pectin

and xyloglucan is the main hemicellulose detected in the PCW. HG is methyl-esterified during its

biosynthesis and later de-methyl-esterified during maturation. We detected a strong signal with the

LM19 antibody, which labels de-methyl-esterified HG, inside pulvinar slits with a stronger signal

close to the boundary with the rest of the PCW (Figure 3A, B). By contrast, the LM20 antibody, which

labels highly methyl-esterified HG, showed no signal in pulvinar slits (Figure 3A, C). We also labeled

XXXG-type xyloglucan with the LM15 antibody, which revealed dispersed signals throughout CMCs

(Figure 3A). We detected a faint signal with the 2F4 antibody, which labels calcium-crosslinked de-

methyl-esterified HG, although the signal appeared to be excluded from pulvinar slits (Supplemental

Figure 2A). We observed no signal when sections were incubated only with the secondary antibody

(Supplemental Figure 2B). These results thus suggested that de-methyl esterification of

homogalacturonan, probably without calcium cross-linking, is associated with pulvinar slits.

To better assess the composition of the pulvinus cell wall, we used Fourier-transform infrared

spectroscopy (FTIR) and monosaccharide analysis to compare the cell wall from pulvini to that of

other axial organs. Using the absorbance profile from FTIR, we identified several peaks (e.g. 778,

1317 cm^{-1}) and a range of 1620–1550 cm^{-1} that are specific to pulvini (Figure 4A). A principal

component analysis revealed that the cell wall properties obtained from FTIR can be classified into

three groups (shown as the dashed circles in Figure 4B), with pulvinar organs forming a group distinct

from all of other axial organs (Figure 4B). The monosaccharide composition of pulvinar organs

(primary pulvinus and secondary pulvinus) was also analyzed in detail. The xylose content mainly

related to xylan in secondary cell wall of pulvini and developing stems was lower than that of petioles

and mature stems (Figure 4C, Table 1). In contrast, pectin-related monosaccharides, such as galactose,

arabinose, rhamnose and galacturonic acid were more abundant in pulvini (Figure 4C, Table 1).

Especially, the abundance of galacturonic acid of pulvini was superior to developing stems (Figure

4C). Therefore, these results suggest that, despite being mature organs, pulvini have pectin-rich cell

wall and resemble developing stems rather than mature organs (petioles and mature stems).

Pulvinar slits in the cell wall have a potential effect on the PD extensibility of pulvini

To explore the mechanical role of pulvinar slits in the turgor-triggered morphological changes of

CMCs, we implemented the finite-element method (FEM) to determine how cell shape might be

affected by the slits under turgor pressure. A previous study using a single cell layer sheet of onion

(*Allium cepa*) peels indicated that cellulose mainly contributes to in-plane rigidity and plasticity, while

pectin contributed very little (Zhang et al., 2019). Therefore, we hypothesized that pulvinar slits would

present a mechanical defect when CMCs deform under turgor. We modeled CMCs as a hexagonal cell

with several elliptical holes (Figure 5A). To set an accurate slit size for simulations, we measured cell

length and the length and width of the slits in 59 randomly selected slits from eight species, yielding

slit lengths ranging from 2.03 to 10.06 μm and a slit width from 0.32 to 2.26 μm. With a cell length

of 20.6-46.1 μm, slit length was between 7.2% and 25.4% of overall cell length, with a mean of about

15%, while slit width was between 0.9% and 9.9% of cell length, with a mean of 3% (Figure 5B). We

also calculated the slit index (length/width), which was 2.46–16.60 with a mean of about 6.14.

Based on these numbers as a standard slit property, we generated several cell models with different

slit properties (Figure 5C-I). Using FEM, we simulated cell deformation under an isotropic pressure

and analyzed the final cell shape. In this calculation, we evaluated the principal direction of growth of

each triangle demonstrated in Figure 5C, representing a vertical elongation by blue and a horizontal

elongation by red and in-between by white. With this setting, we can capture the local direction of

growth by color diagram (Figure 5D, F, H). We explored the cell growth anisotropy as functions of

slit shapes (Figure 5E), slit lengths (Figure 5G) and slit numbers (Figure 5I), respectively. These

simulations revealed that higher slit anisotropy, larger number of slits and larger slit area enhanced a

horizontal cell elongation. This result indicated that the slit structure promotes cell expansion in the

direction perpendicular to the slit. Pulvinar slits aligned in the tangential direction thus have a potential

effect on anisotropic cell extension to the PD direction upon turgor pressure.

The predictions by computer modeling were consistent with a previous finding that the extension of

P. coccineus pulvini shows a bias in the PD direction of the cell (Mayer et al., 1985). To confirm that

pulvini facilitate extensibility along the PD axis, we performed a compression test. In a uniaxial

compression test, a material generally expands in directions perpendicular to the direction of

compression, which is called the Poisson's effect. Using this principle, the continuous deformation in

the PD direction was recorded while compressing axial organs of *D. paniculatum* from a direction

perpendicular to the PD axis (Supplemental Figure S3A, B), and compared the PD extensibility of

pulvini with other axial organs. In addition, because pulvini are thought to be homologous to petiole-

like organs, petiolules, based on the *petiolule-like pulvinus* mutant phenotype in barrel clover

(*Medicago truncatula*) (Zhou et al., 2012), we also performed the compression test especially on

pulvini and petioles using samples from three other species. In *D. paniculatum*, we detected load

oscillation between approximately 0.4 and 0.6 of strain (equivalent to a reduction in height from the

original height to 40% and 60%, respectively) in almost all tested pulvini (Supplemental Figure S4A).

This load oscillation was similar to oscillation seen in stress-strain curve during brittle breaking under

compression. We observed a load oscillation in developing stems between approximately 0.3 and 0.8

of strain, but not in petioles and rarely in mature stems (Supplemental Figure S4A). The length of

pulvini along the PD axis slowly increased until the strain reached 0.3 but extended much more for

strain values above 0.4 (Figure 6A). The length of developing stems increased, but not as much as that

of pulvini (Figure 6A). We did not observe a statistically significant extension of petioles, while mature

stems only slightly extended under greater strain (Figure 6A). For a strain value of 0.5, the extension

rate of pulvini was higher than that of other axial organs (Figure 6B). Similarly, we determined that

pulvini display load oscillation and substantial PD extension in three other legume species, for a strain

value of 0.4 (Figure 6C, Supplemental Figure S4B, C). These results support the idea that pulvini have

excellent PD extensibility in comparison to other axial organs.

To test whether pulvinus extensibility is due to the PD extension of CMCs, we characterized the

deformation of pulvini. There were many tiny wrinkles aligned perpendicularly to the PD axis on the

pulvinar surface (Figure 6D, Supplemental Figure S3B, S5A), which gradually stretched along with

the PD extension during compression (Figure 6D, Supplemental Figure S3B). This phenomenon

resembled that previously reported during the leaf movements of *Mimosa pudica* (Song et al., 2014).

In cross-sectional views, CMCs became separated from the vascular bundles after compression

(Supplemental Figure S5B, C). These results indicate that the PD extension of a pulvinus by

compression results from the extension of CMCs and the spreading of tiny wrinkles in the epidermis,

together with the separation of CMCs and the vascular bundle (Supplemental Figure S5D).

Discussion

The pulvinus is a specialized organ that facilitates dynamic leaf movement. Here, we discovered that

CMCs have a unique cell wall structure, designated the pulvinar slit, with a very low content of

cellulose or highly methyl-esterified HG while being rich in de-methyl-esterified HG. In addition, we

employed computational simulations to predict that the slits bestow CMCs with extensibility in the

perpendicular direction. Finally, we demonstrated that the pulvinus has superior PD extensibility when

compared to other axial organs. We summarize below the properties of the cell wall and speculate on

the mechanical and physiological roles of pulvinar slits. We also discuss the similarities and

differences between unique cell wall structures seen in pulvini and in other tissues. Finally, we place

the discovery of pulvinar slits in context for a greater understanding of plant movement and cell wall

mechanics.

We identified pulvinar slits as tangential slits after calcofluor white staining and established that the

slits are widely conserved in legume species. Whole-mount staining revealed that the structures are

not technical artifacts during sectioning but *bona fide* structures. Staining with pontamine and CBM3a

lent support to our hypothesis that pulvinar slits reflect the pattern of cellulose deposition. De-methyl-

esterified HG, probably without calcium crosslinking, appeared to accumulate abundantly inside

pulvinar slits. By contrast, both cellulose and highly esterified HG were excluded from the pulvinar

slits. FTIR analysis and monosaccharide analysis emphasized the distinct cell wall properties of

pulvini relative to other axial organs and suggested the abundant deposition of pectin in pulvini. Mayer

et al. showed that cellulose microfibrils are oriented "tangentially" (Mayer et al., 1985). Observations

of pulvinar ultrastructure by transmission electron microscopy identified curving walls and pits

including plasmodesmata on CMCs in transverse sections (Fleurat-Lessard and Roblin, 1988;

Machado and Rodrigues, 2004; Rodrigues and Machado, 2006). We also observed similar pits here.

Longitudinal sections of pulvini from *G. max* and *M. truncatula* were previously observed by

microscopy after staining with toluidine blue O (Gao et al., 2017; Du et al., 2020), which visualizes

various components of the cell wall including pectin (O'Brien et al., 1964; Mitra and Loqué, 2014).

However, none of these earlier studies observed the slit-like structures, nor did they report on variation

in cell wall thickness. The de-methyl-esterification of HG by pectin methylesterases (PMEs) was

recently reported to enhance wall hydration and to increase cell wall thickness (Wang et al., 2020).

Taken together, we propose that de-methyl-esterified HG has a role in thickening the cell wall within

the slit, which accumulates little cellulose.

To explore the potential mechanical role played by pulvinar slits in turgor-triggered extension of

CMCs, we performed a FEM simulation based on the assumption that the slits have a mechanical

effect in a specific direction. The FEM results suggested that pulvinar slits have a potential effect on

anisotropic cell extension in the PD direction when the cell is under turgor pressure. This indicates

that the cell can control the mobility or reversible extensibility only by modifying the shape and size

of the slits structure. In this context, the shape anisotropy, number and size of the slits might be

meaningful because they determine the degree of how the cell deforms and deforms back. In a future

work, it should be a target to detect the correlations of the shape anisotropy, number and size of the

slits with a flexibility or bending property of the pulvini.

Previous studies using onion epidermal cells suggested that de-methyl-esterification of HG by PME

treatment causes increased in-plane plasticity of the cell wall but does not affect cell wall creep (Wang

et al., 2020). In addition, pectin enhances the strength of the cell wall perpendicular to the plane (Zhang

et al., 2019; Wang et al., 2020). However, the addition of calcium blocks the increase of plasticity

imparted by PME (Wang et al., 2020). Taken together, we propose that filling the slit with de-methyl-

esterified HG without calcium crosslinking achieves both in-plane plasticity and cell wall integrity. In

our experiments, we detected a faint signal with the 2F4 antibody (recognizing calcium cross-linked

HG) that did not appear to be associated with pulvinar slits. As calcium density in the cell wall can

change during leaf movement (Moysset et al., 2019), further analyses will be required to determine

whether the subcellular localization of calcium has an effect on the mechanical properties of the slits,

including a mechanical test at the cellular level.

In terms of a physiological role, pulvinar slits may also regulate turgor pressure. Pits observed in

previous studies in cross sections were proposed to contribute to volume changes of CMCs and to be

related to water transport, like xylem cell pits (Machado and Rodrigues, 2004; Rodrigues and Machado,

2006). Cell wall components, especially the pattern of cellulose deposition, are indeed responsible for

the localization and/or mobility of transmembrane proteins (Martinière et al., 2012; Li et al., 2021).

Like pits, pulvinar slits may also affect the localization of ion-channel proteins and/or aquaporins to

facilitate water exchange across the cell membrane and modulate cell volume. To clarify the detailed

physiological role of pulvinar slits, molecular biological analyses will be needed.

Considering all the data described here, pulvinar slits can be characterized as a structurally and

mechanically unique property of the cell wall. A unique cell wall structure is linked with the function

of several tissues. For example, cellulose and other components of the SCW are deposited in spiral,

reticulate, scalariform, or pitted patterns in xylem vessels (Oda and Fukuda, 2012a). The cellulose

synthase complex assembles a single cellulose polymer at the plasma membrane and associates with

microtubules (Paredez et al., 2006). SCW-type cellulose synthase complexes localize at the

presumptive area of the SCW in xylem vessels (Watanabe et al., 2015). The deposition pattern of the

SCW is controlled by MICROTUBULE DEPLETION DOMAIN 1 and a Rho-GTPase ROP in

metaxylem cells (Oda and Fukuda, 2012b). By analogy, the unique cellulose pattern seen in the CMCs

of pulvini might be controlled by the localization of cellulose synthase complexes as well. However,

unlike xylem vessels with a thicker SCW, variation in cell wall thickness has not been reported in the

CMCs of pulvini. These differences may be derived from the high accumulation of de-methyl-
esterified HG in the slits, which would suggest that the underlying mechanism establishing the unique
cell wall pattern of CMCs may be considerably different from that of xylem vessels.

The repeatability and reversibility of legume leaf movements are in sharp contrast to other types of
organ movement driven by growth or by mechanical instability. The cellular basis of the repeated and
reversible stomatal movements driven by turgor has been extensively studied. Stomata are composed
of a pair of guard cells on the leaf epidermis that control gas exchange and repeatedly adjust their
aperture by modulating the activity of transmembrane ion channels in response to environmental
signals (Kollist et al., 2014). Guard cells have unique morphological features, for example, a kidney-
like shape in many eudicots or a dumbbell-like shape in grasses. Major components of primary cell
walls, e.g., cellulose, xyloglucan, and pectin, contribute to stomatal movements (Rui and Anderson,
2016; Amsbury et al., 2016; Yi et al., 2018), and computational model-based analyses have
demonstrated that cellulose-based anisotropy and polar stiffening of stomata are important factors
(Carter et al., 2017; Woolfenden et al., 2017; Woolfenden et al., 2018). The cell wall of guard cells is
low in highly methyl-esterified HG and rich in de-esterified HG, a composition similar to that of the
inside of pulvinar slits. Notably, de-methyl-esterification of homogalacturonan is required for stomatal
function (Amsbury et al., 2016). As with guard cells, we propose that the unique deposition pattern of
cell wall components plays an important role in the repeated and reversible deformation of the CMC
cell wall that is driven by turgor pressure during leaf movement. Meanwhile, the spatial arrangement
of cell wall components significantly different between CMCs and guard cells supports the idea that
a unique slit-like pattern has a role in conversion of non-directional inputs (turgor pressure) into
directional outputs (biased cell deformations).

Finally, it is worth noting that we observed slits in all pulvini analyzed. This finding raises the
possibility that the presence of the slit itself might be one of the physical prerequisites for dynamic
leaf movement; slits may therefore cooperate with the regulatory mechanism of turgor pressure that
has been elucidated in recent years. Furthermore, from a broader perspective, it is intriguing that
CMCs with pulvinar slits are structurally analogous to kirigami (Japanese art of paper cutting), a
typical kind of which has many straight cuttings, i.e., slits. Slits of kirigami increase extensibility of
the sheet and prevent unpredictable local structural failures (Shyu et al., 2015; Isobe and Okumura,
2016). In conclusion, we propose that pulvinar slits are an example of specialization in motor organs.
Our findings provide a new insight into the structural diversity of the cell wall and its association with
the mechanical regulation of plant movement.

Methods

Plant materials

Plants used in this study were prepared by four different conditions; outdoor collection of wild plants,

outdoor cultivation, indoor cultivation and greenhouse cultivation (Supplemental Table 1). Wild plants

of *D. paniculatum*, Persian silk tree (*Albizia julibrissin*), and *white clover (Trifolium repens)* were

collected outdoors in Nara, Japan (34°43.5'N, 135°44.1'E), in the summer of 2019 or 2021. Adzuki

bean (*Vigna angularis*) Dainagon (TOHOKU SEED Co., Ltd, Japan), cowpea (*Vigna unguiculate*)

Sansyaku (TOHOKU SEED Co., Ltd, Japan), common bean (*Phaseolus vulgaris*) L. cv *Morocco*

(Atariya Noen Co. Ltd, Japan or TAKII & Co., Ltd, Japan) and sword bean (*Canavalia gladiate*)

(Atariya Noen Co. Ltd, Japan or TAKII & Co., Ltd, Japan) were grown outdoors and collected in the

summer of 2019. Sensitive plant (*Mimosa pudica*) Wase (SAKATA SEED CORPORATION, Japan),

common licorice (*Glycyrrhiza glabra*) (purchased at a garden store) and carob (*Ceratonia siliqua*)

(purchased at a garden store) were grown under SMD 5050 white LED light (Samsung) at 100-200

μmolm^{-2}s^{-1} under long-day condition (16h light/ 8h dark). Plants of soybean (*Glycine max*) and *Lotus*

japonicus were grown in the greenhouse of NAIST adjusted to long-day condition by supplementary

light. Except for species collected outdoors, potted plants of all tested species were grown on culture

soil (Kohnan Shoji Co.,Ltd. or Nihon Hiryo Co.,Ltd). Samples were harvested at Zeitgeber time (ZT)

2–7.

Staining and microscopy

Leaves with pulvini were fixed in FAA solution (5% [v/v] formaldehyde, 5% [v/v] acetic acid, 45%

[v/v] ethanol) or PFA solution (2.22% [w/v] paraformaldehyde, 0.5 M mannitol, 50 mM PIPES, pH

6.9) and stored at 4°C. For the analysis of *D. paniculatum* (Figure 1) and nine other species (excluding

L. japonicus), fixed samples were transferred into distilled water several hours before analysis and

sliced with a single-edged razor. For analyses of *D. paniculatum* (Figures 2, 3, Supplemental Figure

2) and *L. japonicus*, fixed samples were washed in distilled water several times, embedded in 5% (w/v)

agarose gel, and sliced into 100-μm sections with a MicroSlicer Zero 1 (DOSAKA EM, Kyoto, Japan).

For calcofluor white staining, the sections were placed on glass slides, treated with one drop of

calcofluor white stain solution (Sigma-Aldrich, USA) and one drop of 10% (w/v) potassium hydroxide

for 1 min, mounted in distilled water, and covered by a cover slip, before observations with a confocal

laser scanning microscope (LSM800, Zeiss, Germany) equipped with a Plan-Apochromat 20x/0.8

M27 objective lens (Zeiss, Germany; excitation 405 nm; emission 400–560 nm). For whole-mount

observations of *L. japonicus* pulvini, fixed samples were transferred into 1x phosphate buffered saline

(PBS), pH 7.5 (130 mM NaCl, 7 mM Na$_2$HPO$_4$, and 3 mM NaH$_2$PO$_4$), washed in 1x PBS twice, soaked

in ClearSee solution (10% [w/v] xylitol, 15% [w/v] sodium deoxycholate, and 25% [w/v] urea)

(Kurihara et al., 2015) for 1 week, stained with 2.5 μg/mL calcofluor white in ClearSee solution for 5

d, mounted in ClearSee solution, and observed under a LSM800 as described above.

For pontamine staining, the sections were incubated with 50 mg/L Direct Fast Scarlet 4BS (Wako,

Japan) for 20 min, rinsed with 100 mM NaPO$_4$ buffer several times, mounted in distilled water, and

covered by a cover slip, before observations with a LSM800 equipped with a Plan-Apochromat
20x/0.8 M27 objective lens (excitation 488 nm; emission 560–605 nm).

For immunostaining, sections were rinsed with Tris-buffered saline (TBS, 20 mM Tris-HCl, 150 mM
NaCl, pH 7.5) three times, incubated in blocking solution (1% [w/v] bovine serum albumin, 1% [w/v]
normal goat serum (abcam ab7481) in TBS with 0.2% [v/v] Tween-20 [TBST]), rinsed with TBST
three times, treated with TBS (for no antibody control) and with primary antibodies (JIM7, LM19,
2F4, PlantProbes, diluted 1:10) in TBS at 4°C for 1 h, rinsed with TBST three times, incubated with
secondary antibodies (anti-rat IgG-CFL488, anti-mouse IgG-Alexa488, anti-His tag IgG-CFL488, all
diluted 1:100) at 4°C for 1 h, and rinsed with TBST five times. The samples were mounted with 1 mL
TBST with 10 μL calcofluor white stain solution and 10 μL of 10% (w/v) potassium hydroxide,
covered by a cover slip, and observed with a LSM800 equipped with a Plan-Apochromat 20x/0.8 M27
or Plan-Apochromat 100x/1.46 Oil objective lens (excitation 488 nm; emission 400–575 nm;
excitation 405 nm; emission 400–510 nm).

Cell wall analysis

416   Primary pulvini, secondary pulvini, petioles, young stems, and mature stems of *D. paniculatum* were
417 harvested for analysis of cell wall composition. Preparation of cell wall residues and monosaccharide
418 composition analysis were performed as previously described (Sakamoto et al., 2015). Fourier-
419 transform infrared spectroscopy (FTIR) analysis was performed as described previously (Sakamoto et
420 al., 2018).

Computational modeling

The finite-element method (FEM) was implemented based on previously reported FEM models
(Bonazzi et al., 2014; Hong et al., 2016; Tsugawa, 2020). Using FEM, cell deformation under uniform
pressure was confirmed, which indicates a mechanically unstable orientation depending on the internal
structure. To adjust the models with our observations, the initial internal structure was set as one of
the following three types: 1) cells with four slits with different slit anisotropy; 2) cells with different
numbers of slits but with the same slit anisotropy; or 3) cells with four slits with different slit areas.
The cell contour was set to be a regular hexagon, and the side of the hexagon was set to 10 arbitrary
units (au), which was assumed to be micrometers in live cells. With these initial conditions and certain
mechanical properties (Young's modulus 1 MPa and Poisson's ratio 0.35), the cells were uniformly
pressurized with turgor pressure. Increases in the parameters $E = 1, 5, 10$ (MPa) and $\nu =$
$0.35, 0.4, 0.45$ resulted in small deformations, but the overall trend of all three cases was qualitatively
similar. We also assumed that the small deformation and the constitutive law follow the generalized
Hooke's law. When cells were pressurized with a turgor pressure $P = 0.1$ (MPa) to match the typical
scale of previously reported turgor pressure, the cells slightly expanded depending on the slit structure.

As with changes in the mechanical parameters, changes in turgor pressure ($P = 0.2, 0.3$) did not

qualitatively differ from those with $P = 0.1$.

Compression test of pulvini and other axial organs

Each organ was excised into segments of approximately 2 mm in length just before the test. Load

while giving a constant-speed deformation (0.05 mm/s) was measured with a RHEONERII creep

meter (Yamaden, Japan; RE2-3305C) equipped with a 20 N load cell. Excised specimens were placed

on a moving platform with the apex-base axis as the horizontal axis and compressed by sandwiching

each specimen between the moving platform and a cylindrical plunger with a diameter of 3 mm. Strain

was calculated by dividing the deformation by the height at time 0. Analyses were conducted in the

daytime on the same day as sample collection.

To measure the PD extension, videos were captured during the compression test with a 10–200x

magnification USB digital microscope (Giwox, China) and the PhotoBooth App (Apple, USA).

Captured videos were divided into frames with ffmpeg software (https://ffmpeg.org/). Images at the

time of each 10% strain were selected based on their frame rate and the height at time 0. The PD length

of each organ was manually measured with ImageJ (Fiji; https://fiji.sc/) and the extension ratio was

calculated by the following equation in Microsoft Excel.

$$Extension\ ratio = \frac{L_n - L_0}{L_0}$$

where L_0 represents the length when strain is 0 and L_n represents the length when strain is n (0.1-

0.8).

For the observation of the spreading of tiny wrinkles in the epidermis upon compression, videos were

captured with a HiVision Microscope FZ200BA equipped with HDMI recorder X Capture MINI

(Shodensha, Japan).

For observations of loaded pulvini, prior to observation, tissues were manually compressed with a

10 N load while measuring with an electronic balance. For observations by scanning electron

microscopy, tissues were frozen in liquid nitrogen for 15 s and were observed with a FEI Quanta 250

scanning electron microscope (ThermoFisher Scientific, USA). For observations by confocal laser

microscopy, fresh samples embedded in 6% (w/v) agarose gel were sliced into 50-μm sections with a

MicroSlicer Zero 1. For calcofluor white staining, the 50-μm sections were observed by the same

procedure as when observing CMCs. For visualization of lignin autofluorescence, unstained sections

were mounted in distilled water and a cover glass. Sections were observed with a LSM800 as described

above.

Statistical analyses

Microsoft Excel for Mac v16.16.11 (Microsoft, USA) or the Python language v3.7.2

(https://www.python.org/) and Python packages/libraries including Pandas v0.23.4

(https://pandas.pydata.org/), NumPy v1.16.4 (https://www.numpy.org/), Matplotlib v2.2.3

(https://matplotlib.org/), Seaborn v0.9.0 (https://seaborn.pydata.org/), and Statsmodels v0.9.0

(https://www.statsmodels.org/stable/) were used for statistical analyses.

References

Amsbury S, Hunt L, Elhaddad N, Baillie A, Lundgren M, Verhertbruggen Y, Scheller HV, Knox JP,

Fleming AJ, Gray JE (2016) Stomatal Function Requires Pectin De-methyl-esterification of the

Guard Cell Wall. Curr Biol **26:** 2899-2906

Azani N, Babineau M, Bailey D, C., Banks H, Barbosa R, Ariane, Pinto B, Rafael, Boatwright S,

James, Borges M, Leonardo, Brown K, Gillian, Bruneau A, Candido E, Cardoso D, Chung K-

F, Clark P, Ruth, Conceição S, de, Adilva, Crisp M, Cubas P, Delgado-Salinas A, Dexter G,

Kyle, Doyle J, Jeff, Duminil J, Egan N, Ashley, Estrella l, de, Manuel, Falcão J, Marcus,

Filatov A, Dmitry, Fortuna-Perez P, Ana, Fortunato H, Renée, Gagnon E, Gasson P, Rando G,

Juliana, Tozzi A, de, Goulart, Maria, Ana, Gunn B, Harris D, Haston E, Hawkins A, Julie,

Herendeen S, Patrick, Hughes E, Colin, Iganci RV, João, Javadi F, Kanu A, Sheku,

Kazempour-Osaloo S, Kite C, Geoffrey, Klitgaard B, Bente, Kochanovski J, Fábio, Koenen

JM, Erik, Kovar L, Lavin M, Roux l, Marianne, Lewis P, Gwilym, Lima d, C., Haroldo,

López-Roberts C, Maria, Mackinder B, Maia H, Vitor, Malécot V, Mansano F, Vidal, Marazzi

B, Mattapha S, Miller T, Joseph, Mitsuyuki C, Moura T, Murphy J, Daniel, Nageswara-Rao

M, Nevado B, Neves D, Ojeda I, Dario, Pennington T, R., Prado E, Darién, Prenner G,

Queiroz d, Paganucci, Luciano, Ramos G, Filardi R, L., Fabiana, Ribeiro G, Pétala, Rico-Arce

L, de, María, Sanderson J, Michael, Santos-Silva J, São-Mateus MB, Wallace, Silva JS,

Marcos, Simon F, Marcelo, Sinou C, Snak C, Souza d, R., Élvia, Sprent J, Steele P, Kelly,

Steier E, Julia, Steeves R, Stirton H, Charles, Tagane S, Torke M, Benjamin, Toyama H, Cruz

d, Trabuco, Daiane, Vatanparast M, Wieringa J, Jan, Wink M, Wojciechowski F, Martin,

Yahara T, Yi T, Zimmerman E (2017) A new subfamily classification of the Leguminosae based

on a taxonomically comprehensive phylogeny: The Legume Phylogeny Working Group (LPWG).

Taxon **66:** 44-77 , DOI = 10.12705/12661.12703

Bonazzi D, Julien JD, Romao M, Seddiki R, Piel M, Boudaoud A, Minc N (2014) Symmetry breaking in

spore germination relies on an interplay between polar cap stability and spore wall mechanics. Dev

Cell **28:** 534-546

Carter R, Woolfenden H, Baillie A, Amsbury S, Carroll S, Healicon E, Sovatzoglou S, Braybrook S,

Gray JE, Hobbs J, Morris RJ, Fleming AJ (2017) Stomatal Opening Involves Polar, Not Radial,

Stiffening Of Guard Cells. Curr Biol **27:** 2974-2983.e2972

Cosgrove DJ (2005) Growth of the plant cell wall. Nat Rev Mol Cell Biol **6:** 850-861

Cosgrove DJ (2018) Diffuse Growth of Plant Cell Walls. Plant Physiol **176:** 16-27

Darwin C, Darwin F (1880) The power of movement in plants, Chapter VII. John Murray, London

Du J, Lu S, Chai M, Zhou C, Sun L, Tang Y, Nakashima J, Kolape J, Wen Z, Behzadirad M, Zhong T,
Sun J, Zhang Y, Wang ZY (2021) Functional characterization of *PETIOLULE-LIKE PULVINUS*
(*PLP*) gene in abscission zone development in *Medicago truncatula* and its application to genetic
improvement of alfalfa. Plant Biotechnol J **19:** 351-364

Dumais J, Forterre Y (2012) "Vegetable dynamicks" : the role of water in plant movements. Annu Rev
Fluid Mech **44:** 453-478 , DOI = 410.1146/annurev-fluid-120710-101200

Fleurat-Lessard P, Bonnemain JL (1978) Structural and ultrastructural characteristics of the vascular
apparatus of the sensitive plant (*Mimosa pudica* L.). Protoplasma **94:** 127-143

Fleurat-Lessard P, Roblin G (1982) Comparative histocytology of the petiole and the main pulvinus in
Mimosa pudica L. Annals of Botany **50:** 83-92

Fleurat-Lessard P, Roblin G (1988) Structural and ultrastructural features of cortical cells in motor organs
of sensitive plants. Biol Rev **63:** 1-22

Gao J, Yang S, Cheng W, Fu Y, Leng J, Yuan X, Jiang N, Ma J, Feng X (2017) GmILPA1, Encoding an
APC8-like Protein, Controls Leaf Petiole Angle in Soybean. Plant Physiol **174:** 1167-1176

Geitmann A (2016) Actuators Acting without Actin. Cell **166:** 15-17

Geitmann A, Ortega JK (2009) Mechanics and modeling of plant cell growth. Trends Plant Sci **14:** 467-478

Hong L, Dumond M, Tsugawa S, Sapala A, Routier-Kierzkowska AL, Zhou Y, Chen C, Kiss A, Zhu
M, Hamant O, Smith RS, Komatsuzaki T, Li CB, Boudaoud A, Roeder AH (2016) Variable
Cell Growth Yields Reproducible Organ Development through Spatiotemporal Averaging. Dev Cell
38: 15-32

Isobe M, Okumura K (2016) Initial rigid response and softening transition of highly stretchable kirigami
sheet materials. Sci Rep **6:** 24758

Koller D (1990) Light-driven leaf movements. Plant Cell Environ **13:** 615-632 , DOI = 610.1111/j.1365-
3040.1990.tb01079.x

Kollist H, Nuhkat M, Roelfsema MR (2014) Closing gaps: linking elements that control stomatal
movement. New Phytol **203:** 44-62

Kurihara D, Mizuta Y, Sato Y, Higashiyama T (2015) ClearSee: a rapid optical clearing reagent for
whole-plant fluorescence imaging. Development **142:** 4168-4179

Li H, von Wangenheim D, Zhang X, Tan S, Darwish-Miranda N, Naramoto S, Wabnik K, De Rycke R,
Kaufmann WA, Gütl D, Tejos R, Grones P, Ke M, Chen X, Dettmer J, Friml J (2021) Cellular
requirements for PIN polar cargo clustering in *Arabidopsis thaliana*. New Phytol **229:** 351-369

Machado SR, Rodrigues TM (2004) Anatomia e ultra-estrutura do pulvino primário de Pterodon pubescens
Benth. (Fabaceae – Faboideae). Rev Bras Bot **27:** 135-147

Martinière A, Lavagi I, Nageswaran G, Rolfe DJ, Maneta-Peyret L, Luu DT, Botchway SW, Webb SE,

Mongrand S, Maurel C, Martin-Fernandez ML, Kleine-Vehn J, Friml J, Moreau P, Runions J
(2012) Cell wall constrains lateral diffusion of plant plasma-membrane proteins. Proc Natl Acad Sci
U S A **109:** 12805-12810

Mayer WE, Flach D, Raju MV, Starrach N, Wiech E (1985) Mechanics of circadian pulvini movements
in *Phaseolus coccineus* L. : Shape and arrangement of motor cells, micellation of motor cell walls,
and bulk moduli of extensibility ([Formula: see text]). Planta **163:** 381-390

Mitra PP, Loqué D (2014) Histochemical staining of *Arabidopsis thaliana* secondary cell wall elements. J
Vis Exp

Moran N (2007) Osmoregulation of leaf motor cells. FEBS Lett **581:** 2337-2347

Moysset L, Llambrich E, Simón E (2019) Calcium changes in *Robinia pseudoacacia* pulvinar motor cells
during nyctinastic closure mediated by phytochromes. Protoplasma **256:** 615-629

Moysset L, Simon E (1991) Secondary pulvinus of *Robinia pseudoacacia* (leguminosae): structural an
ultrastructural features. Am J Bot **78:** 1467-1486, DOI = 1410.2307/2444972

O'Brien TP, Feder N, McCully ME (1964) Polychromatic Staining of Plant Cell Walls by Toluidine Blue.
Protoplasma **59:** 368-373

Oda Y, Fukuda H (2012a) Secondary cell wall patterning during xylem differentiation. Curr Opin Plant
Biol **15:** 38-44

Oda Y, Fukuda H (2012b) Initiation of cell wall pattern by a Rho- and microtubule-driven symmetry
breaking. Science **337:** 1333-1336

Paredez AR, Somerville CR, Ehrhardt DW (2006) Visualization of cellulose synthase demonstrates
functional association with microtubules. Science **312:** 1491-1495

Rodrigues TM, Machado SR (2006) Anatomia comparada do pulvino prim · rio de leguminosas com
diferentes velocidades de movimento foliar. Rev Bras Bot **29:** 709-720

Rui Y, Anderson CT (2016) Functional Analysis of Cellulose and Xyloglucan in the Walls of Stomatal
Guard Cells of Arabidopsis. Plant Physiol **170:** 1398-1419

Sakamoto S, Somssich M, Nakata MT, Unda F, Atsuzawa K, Kaneko Y, Wang T, Bågman AM,
Gaudinier A, Yoshida K, Brady SM, Mansfield SD, Persson S, Mitsuda N (2018) Complete
substitution of a secondary cell wall with a primary cell wall in Arabidopsis. Nat Plants **4:** 777-783

Sakamoto S, Yoshida K, Sugihara S, Mitsuda N (2015) Development of a new high-throughput method to
determine the composition of ten monosaccharides including 4-O-methyl glucuronic acid from plant
cell walls using ultra-performance liquid chromatography. Plant Biotechnol **32:** 55-63

Shyu TC, Damasceno PF, Dodd PM, Lamoureux A, Xu L, Shlian M, Shtein M, Glotzer SC, Kotov NA
(2015) A kirigami approach to engineering elasticity in nanocomposites through patterned defects.
Nat Mater **14:** 785-789

Smithers ET, Luo J, Dyson RJ (2019) Mathematical principles and models of plant growth mechanics:
from cell wall dynamics to tissue morphogenesis. J Exp Bot **70:** 3587-3600

Song K, Yoem E, Lee SJ (2014) Real-time imaging of pulvinus bending in *Mimosa pudica*. Sci. Rep. **4:**
6466

Tsugawa S (2020) Suppression of soft spots and excited modes in the shape deformation model with spatio-
temporal growth noise. J Theor Biol **486:** 110092

Wang X, Wilson L, Cosgrove DJ (2020) Pectin methylesterase selectively softens the onion epidermal wall
yet reduces acid-induced creep. J Exp Bot **71:** 2629-2640

Watanabe Y, Meents MJ, McDonnell LM, Barkwill S, Sampathkumar A, Cartwright HN, Demura T,
Ehrhardt DW, Samuels AL, Mansfield SD (2015) Visualization of cellulose synthases in
Arabidopsis secondary cell walls. Science **350:** 198-203

Weintraub M (1952) Leaf movements in *Mimosa pudica* L. New Phytol **50:** 357-382

Woolfenden HC, Baillie AL, Gray JE, Hobbs JK, Morris RJ, Fleming AJ (2018) Models and
Mechanisms of Stomatal Mechanics. Trends Plant Sci **23:** 822-832

Woolfenden HC, Bourdais G, Kopischke M, Miedes E, Molina A, Robatzek S, Morris RJ (2017) A
computational approach for inferring the cell wall properties that govern guard cell dynamics. Plant J
92: 5-18

Yi H, Rui Y, Kandemir B, Wang JZ, Anderson CT, Puri VM (2018) Mechanical Effects of Cellulose,
Xyloglucan, and Pectins on Stomatal Guard Cells of. Front Plant Sci **9:** 1566

Zhang Y, Tang H, Vavylonis T, Cosgrove DJ (2019) Disentangling loosening from softening: insights into
primary cell wall structure. Plant J **100:** 1101-1117

Zheng Y, Wang X, Chen Y, Wagner E, Cosgrove DJ (2018) Xyloglucan in the primary cell wall:
assessment by FESEM, selective enzyme digestions and nanogold affinity tags. Plant J **93:** 211-226

Zhou C, Han L, Fu C, Chai M, Zhang W, Li G, Tang Y, Wang ZY (2012) Identification and
characterization of petiolule-like pulvinus mutants with abolished nyctinastic leaf movement in the
model legume *Medicago truncatula*. New Phytol. **196(1):** 92-100

Acknowledgements and funding
We thank Kiyotaka Okada (Ryukoku Univ., Japan) and Nobutaka Mitsuda (AIST, Japan) for helpful
comments, Atsunobu Suzuki (NAIST, Japan), Hitomi Ichikawa (NAIST, Japan) and Aeni Hosaka
(AIST, Japan) for technical support. We thank Tadashi Kunieda (NAIST, Japan), Ko Kato (NAIST,
Japan) and members of the Demura lab for general support during experiments. *G. max* seeds were
obtained from LegumeBase (http://www.legumebase.brc.miyazaki-u.ac.jp/), which is supported by
the National BioResource Project (NBRP) in Japan. We thank Plant Editors (https://planteditors.com/)
for English-language editing.
This work was supported by JSPS KAKENHI Grant Numbers JP20K15832 (to S.T.), JP19K16174
(to S.S.), JP19K23753 (to M.N.) and JP20K06707 (to M.N.), and by MEXT KAKENHI Grant-in-Aid
for Scientific Research on Innovative Areas "Plant-Structure Optimization Strategy" Grant Numbers

JP18H05484 and JP18H05489 (to T.D.).

Author Contributions

M.T and M.T.N. contributed conception and design of the study, performed

biological/histochemical/mechanical experiments and image/statistical analyses, and wrote the first

draft of the manuscript. S.T. performed FEM analysis and wrote the section of FEM simulation. S.S.

performed FTIR and monosaccharide analysis, contributed the data interpretation, and wrote the

section of cell-wall assay. T.D. contributed significantly the revision of the experimental design and

the modification of the draft. All authors discussed the results and contributed to the final manuscript.

Ethics declarations

Competing interests

The authors declare no competing interest.

Supplemental information

Supplemental figures and table

Figure legends

Figure 1. Calcofluor-white staining of pulvinar cortical motor cells (CMCs) in various legume species.

(A) Schematic diagram (left) and photography (right) of a *Desmodium paniculatum* leaf. Pu, primary

pulvinus; SPu, secondary pulvinus; Pe, petiole; Ds, developing stem; Ms, mature stem. (B) Schematic

diagrams of the collected pulvini and petiole samples for microscopy. Letters C to F indicate the

positions visualized in the corresponding panels. (C-F) Sections of calcofluor white-stained *D.*

paniculatum pulvini and petioles by confocal laser scanning microscopy. AdAb, adaxial-abaxial axis;

PD, proximo-distal axis; ML, medial-lateral axis; ep in F indicates epidermis. (G) A model of pulvinar

slits and small spots in the cell wall of a CMC. TPu, a tertiary pulvinus. (H) Vertical sections of

calcofluor white-stained pulvinar CMCs from 11 legume species. The vertical direction of all images

corresponds to the adaxial-abaxial axis. (I) Whole-mount projection of calcofluor white-stained *L.*

japonicus CMCs (top). Middle, magnified view of the orange square. Bottom, optical Z-section shown

as the blue line in the middle panel. Arrowheads indicate pulvinar slits. Bars, 2 cm (A), or 20 μm (C-

F, H, I).

Figure 2. Visualization of cellulose in *D. paniculatum* pulvinar CMCs.

(A) Pontamine-stained CMCs of a *D. paniculatum* pulvinus (vertical section). (B) Co-localization of

calcofluor white signal (CF) and staining with the CBM3a antibody (recognizing crystalline cellulose)

in CMCs in a vertical section of a *D. paniculatum* pulvinus. Arrowheads, pulvinar slits. Bars, 20 μm.

Figure 3. Immuno-histological analyses of *D. paniculatum* pulvinar CMCs.

(A) Immuno-labeled CMCs with antibodies against primary cell wall (PCW) components in vertical

sections. These samples were co-stained with calcofluor white (CF). Asterisk indicates a tannin

vacuole. (B-C) Magnified views of dashed boxes in (A) (left) and profiles of relative intensities for

calcofluor white (blue) and antibody signal (green) along the yellow lines in the magnified images

(right). Bars, 20 μm (A).

Figure 4. Cell wall assay of *D. paniculatum* axial organs.

(A) FTIR spectra of cell wall components obtained from five different axial organs: primary pulvini

(Pu), secondary pulvini (SPu), petioles (Pe), developing stems (Ds), and mature stems (Ms), as six

biological replicates. Data are shown as the mean and 95% confidence interval. (B) Principal

component analysis applied to the FTIR spectra shown in A. (C) The heatmap with dendrogram for

the relative amount of each monosaccharide in each organ. Color indicates the ratio to the mean in

each monosaccharide. Gal, galactose; GalA, galacturonic acid; Ara, arabinose; Fuc, fructose; GlcA,

glucuronic acid; Man, mannose; Rha, rhamnose; Glc, glucose; Xyl, xylose.

Figure 5. Computational modeling of the expansion of a hexagonal cell with slits.

(A) Schematic diagram of the hexagonal cells with definition of terms used. (B) Relative length and

width of 59 slits from eight species: *D. paniculatum*, *P. vulgaris*, *V. angularis*, *V. unguiculata*, *G. max*

(Jack), *A. julibrissin*, *M. pudica*, and *C. siliqua*. (C) Color key for principal direction of growth shown

in D, F, and H. 0, horizontal plane; $\pi/2$, vertical plane. (D-I) Results of FEM simulations to clarify the

effect of slit direction (D, E), slit number (F, G), or slit size (H, I). In D, F, and H, the upper panels

indicate finite element mesh, and the lower panels indicate the final shape and the principal direction

of growth. er_x, extension ratio along the x (horizontal) axis; er_y, extension ratio along the y (vertical)

axis; r_x, slit length along the x-axis; r_y, slit length along the y-axis.

Figure 6. Compression test of pulvini and other axial organs.

(A) Ratio of the PD extension of the axial organs for each strain applied. N = 18 (pulvini, Pu), 10

(petioles, Pe), 15 (developing stems, Ds), 10 (mature stems, Ms, other than strain = 0.7), and 8 (mature

stems, strain = 0.7). Different letters indicate statistical differences (Tukey's HSD test, family-wise

error rate = 0.05). (B) Ratio of the PD extension of the axial organs in *D. paniculatum* when the applied

strain is 0.5. (C) Ratio of the PD extension of axial organs in three other species when the applied

strain is 0.5. N = 15 (*P. vulgaris* petioles), 16 (*P. vulgaris* pulvini), 12 (*V. angularis*), and 11 (*V.*

unguiculata). $***p < 0.001$ by two-sided Welch's *t*-test. The calculated t-statistics for *P. vulgaris*, *V.*

angularis, and *V. unguiculata* are 12.20, 9.04, and 9.05, respectively. (D) Serial images of side views

of *D. paniculatum* pulvini during compression with a creep meter.

Figure 1. Calcofluor-white staining of pulvinar cortical motor cells (CMCs) in various legume species.

(A) Schematic diagram (left) and photography (right) of a *Desmodium paniculatum* leaf. Pu, primary pulvinus; SPu, secondary pulvinus; Pe, petiole; Ds, developing stem; Ms, mature stem. (B) Schematic diagrams of the collected pulvini and petiole samples for microscopy. Letters C to F indicate the positions visualized in the corresponding panels. (C-F) Sections of calcofluor white-stained *D. paniculatum* pulvini and petioles by confocal laser scanning microscopy. AdAb, adaxial-abaxial axis; PD, proximo-distal axis; ML, medial-lateral axis; ep in F indicates epidermis. (G) A model of pulvinar slits and small spots in the cell wall of a CMC. TPu, a tertiary pulvinus. (H) Vertical sections of calcofluor white-stained pulvinar CMCs from 11 legume species. The vertical direction of all images corresponds to the adaxial-abaxial axis. (I) Whole-mount projection of calcofluor white-stained *L. japonicus* CMCs (top). Middle, magnified view of the orange square. Bottom, optical Z-section shown as the blue line in the middle panel. Arrowheads indicate pulvinar slits. Bars, 2 cm (A), or 20 μm (C-F, H, I).

Figure 2. Visualization of cellulose in *D. paniculatum* pulvinar CMCs.

(A) Pontamine-stained CMCs of a *D. paniculatum* pulvinus (vertical section). (B) Co-localization of calcofluor white signal (CF) and staining with the CBM3a antibody (recognizing crystalline cellulose) in CMCs in a vertical section of a *D. paniculatum* pulvinus. Arrowheads, pulvinar slits. Bars, 20 μm.

Figure 3. Immuno-histological analyses of *D. paniculatum* pulvinar CMCs.

(A) Immuno-labeled CMCs with antibodies against primary cell wall (PCW) components in vertical sections. These samples were co-stained with calcofluor white (CF). Asterisk indicates a tannin vacuole. (B-C) Magnified views of dashed boxes in (A) (left) and profiles of relative intensities for calcofluor white (blue) and antibody signal (green) along the yellow lines in the magnified images (right). Bars, 20 μm (A).

Figure 4. Cell wall assay of *D. paniculatum* axial organs.

(A) FTIR spectra of cell wall components obtained from five different axial organs: primary pulvini (Pu), secondary pulvini (SPu), petioles (Pe), developing stems (Ds), and mature stems (Ms), as six biological replicates. Data are shown as the mean and 95% confidence interval. (B) Principal component analysis applied to the FTIR spectra shown in A. (C) The heatmap with dendrogram for the relative amount of each monosaccharide in each organ. Color indicates the ratio to the mean in each monosaccharide. Gal, galactose; GalA, galacturonic acid; Ara, arabinose; Fuc, fructose; GlcA, glucuronic acid; Man, mannose; Rha, rhamnose; Glc, glucose; Xyl, xylose.

Figure 5. Computational modeling of the expansion of a hexagonal cell with slits.

(A) Schematic diagram of the hexagonal cells with definition of terms used. (B) Relative length and width of 59 slits from eight species: *D. paniculatum*, *P. vulgaris*, *V. angularis*, *V. unguiculata*, *G. max* (Jack), *A. julibrissin*, *M. pudica*, and *C. siliqua*. (C) Color key for principal direction of growth shown in D, F, and H. 0, horizontal plane; π/2, vertical plane. (D-I) Results of FEM simulations to clarify the effect of slit direction (D, E), slit number (F, G), or slit size (H, I). In D, F, and H, the upper panels indicate finite element mesh, and the lower panels indicate the final shape and the principal direction of growth. er_x, extension ratio along the x (horizontal) axis; er_y, extension ratio along the y (vertical) axis; r_x, slit length along the x-axis; r_y, slit length along the y-axis.

Figure 6. Compression test of pulvini and other axial organs.

(A) Ratio of the PD extension of the axial organs for each strain applied. N = 18 (pulvini, Pu), 10 (petioles, Pe), 15 (developing stems, Ds), 10 (mature stems, Ms, other than strain = 0.7), and 8 (mature stems, strain = 0.7). Different letters indicate statistical differences (Tukey's HSD test, family-wise error rate = 0.05). (B) Ratio of the PD extension of the axial organs in *D. paniculatum* when the applied strain is 0.5. (C) Ratio of the PD extension of axial organs in three other species when the applied strain is 0.5. N = 15 (*P. vulgaris* petioles), 16 (*P. vulgaris* pulvini), 12 (*V. angularis*), and 11 (*V. unguiculata*). ***$p < 0.001$ by two-sided Welch's *t*-test. The calculated t-statistics for *P. vulgaris*, *V. angularis*, and *V. unguiculata* are 12.20, 9.04, and 9.05, respectively. (D) Serial images of side views of *D. paniculatum* pulvini during compression with a creep meter.

Table 1. The amount of cell-wall components (µg/gAIR) in aerial axial organs

	Primary pulvinus		Secondary pulvinus		Petiole		Developing stem		Mature stem	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Glc***	228.97	8.27 [ab]	180.03	5.46 [a]	345.59	20.16 [bc]	221.73	15.08 [ab]	401.62	19.84 [c]
GalA***	157.30	5.98 [a]	126.41	8.79 [ab]	109.45	6.01 [abc]	96.89	4.89 [bc]	66.94	10.33 [c]
Xyl***	51.73	9.12 [ab]	33.49	3.43 [a]	108.06	8.60 [b]	30.05	1.65 [a]	143.79	10.76 [b]
Gal***	46.61	1.73 [ab]	41.87	3.17 [abc]	30.51	3.02 [bc]	61.40	1.66 [a]	20.64	2.59 [c]
Man***	41.92	1.97 [ab]	39.57	1.49 [abc]	34.40	1.73 [bc]	43.92	1.06 [a]	30.57	2.76 [c]
Ara***	38.03	1.67 [a]	23.29	2.09 [ab]	21.58	1.82 [bc]	27.84	0.76 [ac]	16.25	1.77 [b]
Rha***	14.08	0.34 [a]	11.17	0.92 [ab]	10.25	0.67 [bc]	12.03	0.14 [ac]	8.13	0.99 [b]
GlcA***	2.86	0.18 [a]	2.17	0.18 [ab]	2.51	0.28 [ab]	2.67	0.12 [a]	2.04	0.17 [b]
Fuc***	2.88	0.29 [a]	2.41	0.12 [ab]	2.14	0.26 [bc]	2.52	0.23 [ac]	1.33	0.15 [b]
Total***	584.37	15.82 [ab]	460.40	17.74 [a]	664.48	30.48 [b]	499.05	23.41 [a]	691.31	36.70 [b]

n = 6; ***$p < 0.001$ by the Kruskal-Wallis H-test; a-c, different characters indicate significantly different ($p<0.05$) by pairwise test for multiple comparisons of mean rank sums (Dunn's test; Holm's method)