Prokaryotic genome design is ruled by the mechanical properties and complexity

of the DNA sequence

Tsuneyuki Tatsuke[1*]

[1] Faculty of Advanced Science and Technology, Kumamoto University, Kumamoto, Japan

* Corresponding author

E-mail address: tatsuke@kumamoto-u.ac.jp (TT)

Short title: The macro-level design rules of the prokaryotic genome

Abstract

Although we represent the genetic code using four letters, namely A, T, G, and C, it is a chemical entity with sequence-dependent physical properties and a spatial extent. In other words, it can be said that genetic information is compressed and recorded in the genomic DNA as a storage medium with a spatial spread. This study analyzed prokaryotic genomes in terms of their physical properties and complexity. The findings clearly indicated that a mathematic equation can be used to express the relationship between the length of a genome and the upper limit of its rigidity. Furthermore, many prokaryotes have been shown to maximize DNA complexity under various physical constraints. Here, this relationship between rigidity and complexity was found to be shaped like an entropy function. It was also clarified that codon usage tendencies are affected by the physical properties and complexity of the DNA sequence. Moreover, the analysis of codons encoding two consecutive synonymous amino acids suggests that many prokaryotic codon combinations tend to increase genomic DNA complexity. This study revealed the macro-level rules for packing genomic DNA into prokaryotic cells.

Introduction

DNA sequences are usually represented with four letters: A, T, G, and C. The actual genomic DNA is, however, a heterogeneous chain with a double-helical structure consisting of four types of chemical components. The genome is, therefore, an information storage medium with a

spatial extent and distinct physical properties [1]. The sequence-dependent properties of

double-helical DNA have been extensively studied. At the simplest level, the sequence–structure

relationships of dinucleotides as the fundamental building blocks of DNA polynucleotides have been

reported [2]. There have also been reports on the sequence-dependent properties of

tetranucleotide-based sequences, and the structure prediction models for long DNA sequences

containing either are in good agreement with the X-ray crystal structure data of DNA [3].

Furthermore, *in silico* genomic DNA structure prediction of whole-genome folding in yeast based on

these prediction models and the DNA-encoded nucleosome organization reported by Kaplan et al.

(2009) [4] reflects the actual structure of genomic DNA including the chromatin structure [1].

Therefore, genomic DNA records structural information in addition to genetic information, and it can

be said that genetic information and the mechanical properties of genomic DNA are inextricably

linked.

In the last decade, there have been reports of the activation of a prokaryote from chemically

synthesized genomic DNA [5], the creation of artificial bacteria with minimal genomic DNA [6], and

the generation of *Escherichia coli* bearing recoded genome using only 61 codons by synonymous

codon compression [7]. A fully engineered bacterial genome, however, has not yet been created

because the dynamic behavior of the entire biological system is still not fully understood. A

systems-level understanding of dynamic processes such as metabolism, protein interactions, RNA

transcriptional control, and the multi-layered control of all these processes is required for the

complete design of biological systems, and this is an essential area of research in systems biology

and synthetic biology. On the other hand, the understanding of genomic DNA based on its physical

properties is also required for the complete design of biological systems, and this can be analyzed by

mainly assuming a static state. In eukaryotes, the state of genomic DNA within the nucleosomes in

the nucleus has been revealed using the Hi-C method [8], and it is now known that there are

three-dimensional, self-interacting genomic regions, termed topologically associating domains [9]. In

prokaryotes, on the other hand, approximately 85% of the genomic DNA is composed of

uninterrupted protein-coding sequences, i.e., open reading frames (ORFs), without nucleosomes and

other DNA-binding proteins, and their mechanical properties are expected to be firmly ruled by

trinucleotides—the codons. Indeed, A-tract clusters, which are the sequence motifs introducing the

most pronounced DNA curvature, have been reported to facilitate DNA packaging in the bacterial

nucleoid [10]. Besides, it is necessary to pack genomic DNA together with RNA and protein in the

limited space available within prokaryotic cells. That is, genomic DNA is considered to hold not only

genetic information but also structural information. Therefore, understanding of the storage of

prokaryotic genome-wide genetic information requires an approach based on physical ideas.

In this study, the macro-level rules for packing genomic DNA into prokaryotic cells were

revealed by focusing on the mechanical properties and sequence diversity of DNA as a

heterogeneous polymer. This study was mainly based on an analysis of the mechanical properties of

prokaryotic genomic DNA in units of 4 base pairs (bp) [3, 11], and diversity analysis using the

Shannon entropy score [12]. From the analysis of the relationship between mechanical properties and genome length using the complete genomic DNA sequences of approximately 3,000 prokaryotic organisms registered in the National Center for Biotechnology Information (NCBI) GenBank database, the formula describing the upper limit of rigidity, a mechanical property of genomic DNA, was obtained. This formula also explains the upper limit of the size of prokaryotic genomic DNA. Of course, bottom-up approaches should consider the presence of nucleoid-constituting proteins such as HU. However, since this study aimed to discover the rules governing genomic DNA at the macro-level in a top-down manner, the effects of nucleoid-constituting proteins are also encompassed within the obtained formula. The entropy score of the genomic DNA sequence complexity showed that prokaryotes tend to maximize genomic DNA complexity as much as possible under the constraints of the mechanical properties of the genome. Moreover, the present study identified mechanical properties and sequence diversity as the driving forces that correlate codon usage with genomic DNA size and species. This study has revealed that genetic information, the mechanical properties of genomic DNA, and the complexity of DNA sequences are inseparable. These results are expected to contribute toward the creation of a fully engineered bacterial genome in the future.

Results

The mechanical properties of DNA limit the spatial extent of

prokaryotic genomes

In most prokaryotes, the protein-coding region accounts for approximately 80%–90% of the genomic DNA (Fig 1). In other words, the codon context firmly rules most sections of the prokaryotic genome. DNA sequences are known to have mechanical properties that depend on the arrangement of the nucleotides [3]. Therefore, it was expected that prokaryotic genomic DNA, which is mostly composed of ORFs in which codons consisting of three bases are arranged regularly without interruption, would have specific mechanical characteristics. Therefore, an analysis was conducted focusing on the physical properties of the DNA including the length of the genomic DNA, using the method described in Fukue et al. (2004) [11], and the parameters described in Packer et al. (2000) [3] (see Methods for more details). Non-linear regression analysis (Fig 2A) revealed that the upper limit of the rigidity of prokaryotic genomic DNA and its size can be described by the following equation:

$$L_{max} = \frac{c}{1 + a \cdot exp\big(b(\bar{R} - d)\big)} + e \# (1)$$

where L_{max} indicates the maximum size of the genomic DNA, $\bar{R}$ refers to the average rigidity of the genomic DNA, and a, b, c, d, and e are the values obtained by non-linear regression analysis: $a = 0.00854163$, $b = 2.178543$, $c = 10.5789$, $d = 12.14316$, and $e = 0.058$. At present, however, it is difficult to explain the physical meaning of these parameters. Equation (1) shows that the upper size limit of prokaryotic genomic DNA, L_{max}, is ruled by its average rigidity, $\bar{R}$. In equation (1), $\bar{R}$ represents an average value of the energy required to deform a section of the

genomic DNA per bp ($kJ \cdot mol^{-1} \cdot Å^{-2}/bp$) [3]. Since the relationship between persistent length of a

DNA fragment y (nm) and DNA rigidity x ($kJ \cdot mol^{-1} \cdot Å^{-2}$) can be expressed by the equation

$y = 2.1246x + 10.566$ [1], higher $\bar{R}$ values indicate a stiffer DNA sequence and higher space

costs, while lower $\bar{R}$ values indicate a more flexible DNA sequence and lower space costs. Contrary

to expectations, the randomly sorted genomic DNA sequences of 2,943 prokaryotes by single- or

triple-base units did not significantly change the distribution of species in the relationship between

their rigidity and length (Figs 2B and C), comparing to the randomly generated genomic DNA (Figs

2D and E). Therefore, it is suggested that the set of bases that compose the genome sequences of

2,943 prokaryotes themself have the property of generating upper and lower limits that are inherent

to organisms. Furthermore, such properties were not observed for genomic DNAs comprising

sequences that were randomly generated with a computer (Figs 2D and E), indicating that the

relationship described by equation (1) is inherent to organisms. These findings are probably

important in considering the relationship between genomic DNA length and genomic DNA folding in

prokaryotes by macro-level, including effect by nucleoid-related proteins.

Fig 1. Ratios of protein-coding regions in genomic DNAs of different lengths from 2,943

prokaryotic species. Each data point corresponds to one species.

Fig 2. Relationship between the mechanical properties of average genomic sequence rigidity

and genome length. (A) Relationship between the average rigidity and length of the genomic DNA

of 2,943 prokaryotes. Each data point corresponds to one species. The blue line indicates the upper

limit of the rigidity of the genomic DNA, while the orange line represents the lower limit. (B and C)

Genomic DNA sequences of 2,943 prokaryotes were randomly sorted by various base unit sizes

(single nucleotides, i.e., ATTGCCGAC to TGACAGCCT, or codon units, i.e., ATTGCCGAC to

GACATTGCC). Genomic DNA sequences were sorted by single-base units (B), or triple-base units

(C). (D and E) *In silico*, 2,943 genomic DNA sequences were randomly generated using single-base

units (D), or triple-base units (E). The relationship between their rigidity and length was plotted.

Each data point corresponds to one randomly generated or sorted genomic DNA sequence.

Solving equation (1) for the average rigidity of the genomic DNA yields:

$$\bar{R} = \frac{1}{b} log \left(\frac{1}{a} \left(\frac{c}{L_{max} - e} - 1 \right) \right) + d \# (2)$$

136   Equation (2) implies that when the size of the genome increases by a particular value, its rigidity

137   decreases by a specific value to mitigate the space cost. In other words, prokaryotic genomic DNAs

138   have evolved according to design rules that allow more compact storage as their size increases. The

139   maximum theoretical size of prokaryotic genomic DNA deduced from these equations is

140   approximately 10 million base pairs (Mbp).

The prokaryotic genome maximizes the complexity of its nucleotide combinations under the constraints of the mechanical properties of DNA

So how does genomic DNA complexity occur under these mechanical constraints? The answer to this question lies in the diversity of the DNA sequences carried by genomic DNA. The complexity of the prokaryotic genomic DNAs was assessed based on Shannon entropy and plotted in comparison with their average rigidity. The complexity of genomic DNA and the entropy score was found to be proportional. The genomic DNA sequences of 2,943 prokaryotes produced a curve similar to the entropy function in terms of average rigidity and entropy (Fig 3A). Further, this curve indicated that the majority of prokaryotic species lie within the curve maxima.

Focusing on the lower limit of the rigidity of prokaryotic genomic DNA, this value takes a fixed value of $12.14316\,\mathrm{kJ} \cdot \mathrm{mol}^{-1} \cdot \mathrm{Å}^{-2}$. *In silico*, 2,943 genomic DNA sequences were randomly sorted under various conditions to confirm this lower limit. When the genomic DNA sequences were sorted using units of a single base strung together randomly, the rigidity did not reach this lower limit (Fig 3B). However, when the genomic sequence was randomly sorted using units comprising triple bases (corresponding to codons), the rigidity almost reached this lower limit (Fig 3C). This value is considered to derive from the genomic regions consisting of three-base units, which account for approximately 85% of prokaryotic genomic DNA. In other words, the lower limit of the rigidity of prokaryotic genomic DNA is derived from the major structural unit of genomic DNA.

 Fig 3. Relationship between the complexity and rigidity of genomic DNA sequences. (A) The

rigidity and complexity of genome DNA sequences of 2,943 prokaryotes. Each data point

corresponds to one species. (B and C) Genomic DNA sequences of 2,943 prokaryotes were randomly

sorted by various base unit sizes (single nucleotides, i.e., ATTGCCGAC to TGACAGCCT, or codon units,

i.e., ATTGCCGAC to GACATTGCC). Genomic DNA sequences were sorted by single-base units (B), or

triple-base units (C). (D and E) *In silico*, 2,943 genomic DNA sequences were generated using

various base unit sizes (single nucleotides or codons). Genomic DNA sequences were generated

using single-base units (D), triple-base units (E). The relationships between their complexity and

rigidity were plotted. Each data point corresponds to one randomly sorted or generated genomic

DNA sequence. In all plots, the vertical axis represents Shannon entropy as a proxy for the

complexity of the genomic DNA sequence, and the horizontal axis represents the average rigidity of

the genomic DNA sequence. The orange dashed line indicates the lower limit of the rigidity as

determined in Fig 2.

Comparing the plot of the genomic DNA sequences of 2,943 prokaryotes (Fig 3A) with

plots of randomly sorted genomic DNA sequences (Figs 3B and C), the distribution shape of the plot

of DNA sequences shuffled by single-base units tends to be thin and more rigid (Fig 3B), while the

distribution shape of the plot of them shuffled by triple-base units is close to that of the plot of

organisms (Fig 3C). This result is possibly due to about 85% of prokaryotic genomic DNA consists

of protein-coding regions with triple-base units. The original genomic DNA shows a more diverse

distribution than randomly sorted genomes, presumably because of the rules inherent to organisms

for arranging DNA nucleotides. Also, comparing these plots with plots of randomly generated DNA

sequences (Figs 3D and E), it was apparent that the DNA sequences derived from organisms tend to

maximize diversity as much as possible and that this is probably a property inherent in the set of

bases that compose genomic DNA of organisms. In the case of genomic DNAs randomly generated

by a computer and comprising single-base units, the entropy values extended downward (Fig 3D).

This indicated that these DNA sequences tended to be simpler. In contrast, when the genomic DNAs

were randomly generated using triple-base units, the entropy values were biased upward (Fig 3E).

These results suggest that prokaryotes adapt codon usage and the codon context by

diversifying the genomic DNA sequence within the structures of many limitations, including

mechanical limitations, environmental factors, and the size of the genetic system encoding the

genomic DNA. This would help to avoid monotony in the genetic code and create genomic DNA

diversity in its stead.

The mechanical properties and complexity of DNA sequences rule the codon context

The prokaryotic preferences for codon usage were analyzed to determine codon usage biases

when encoding the same amino acid at two consecutive codons. If the genetic code functions only as

a "letter," there should be no differences in the order of codon combinations when encoding the same

amino acid at two consecutive positions. However, if there is an interplay between the mechanical

properties of genomes and their genetic code, it is expected that the order of such combinations

would be affected. To investigate this, the directionality index (DI) was used to analyze whether the

combination of codons coding the same amino acid was biased toward rigid or flexible [10]. The

average rigidity of triplet repeat sequences was calculated using the method described in Fukue et al.

(2004) [11] and the parameters described in Packer et al. (2000) [3] (see Methods for more details).

Here, the Cys-Cys amino acid sequence encoded by the combination of tgc and tgt was chosen as an

example of a combination that is rarely used by most species. The Ala-Ala amino acid sequence

encoded by the combination of gcc and gcg was chosen as an example of a combination with

different frequencies of use among species. Lastly, the Leu-Leu amino acid sequence encoded by the

combination of ctt and ttg, which exerts a largely different effect on rigidity depending on the order

of the codon combination, was also chosen.

The Cys-Cys amino acid sequence encoded by the combination of tgc and tgt was found to

be used infrequently in either of the two consecutive codon combinations and was considered to have

little structural effect on the genomic DNA. For this reason, DI bias was observed in many species

(Fig 4A). On the other hand, the Ala-Ala amino acid sequence encoded by the combination of gcc

and gcg is frequently used in many species and tended to show a less biased DI in the species in

which this combination is frequently used (Fig 4B). Also, in the species with a low frequency of use

of the combination of gcc and gcg, DI is similar to Cys-Cys encoded by the combination, tgc and tgt.

Because it is the nature of organisms to try to diversify their genomic DNA sequences as much as

possible, these relationships between frequency of use and bias appear reasonable. In addition, in the

Leu-Leu amino acid sequence with a large difference in rigidity depending on the order of the

combination of ctt and ttg, it was observed that many species tended to be biased toward the flexible

ttg-ctt side (Fig 4C). When such a large difference in rigidity is observed according to the order of

codon combination, it is considered that each codon combination tends to have its own unique DI

pattern.

Fig 4. Codon usage bias when the same amino acid is encoded by two different consecutive codons. (A) When Cys-Cys is encoded by the tgc and tgt codons, the rigidity of tgc-tgt is 12.73 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}, while the rigidity of tgt-tgc is 11.80 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}. (B) When Ala-Ala is encoded by the gcg and gcc codons, the rigidity of gcg-gcc is 15.00 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}, while the rigidity of gcc-gcg is 11.97 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}. (C) When Leu-Leu is encoded by the ctt and ttg codons, the rigidity of ctt-ttg is 24.20 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}, while the rigidity of ttg-ctt is 10.97 kJ $\cdot$ mol$^{-1} \cdot$ Å^{-2}. For all plots, the directional index (DI) represents bias toward the usage of a particular combination of codons. DI takes a value from -1 to 1, with highly rigid codon combinations having a positive DI and more flexible codon combinations having a negative DI. Each data point corresponds to one

species. The vertical axis represents the DI, and the horizontal axis represents the ratio of the usage

frequency of the codons encoding the two target consecutive amino acids to that of all two-codon

combinations. Δ_{rigidity} represents the difference in rigidity caused by the differing order of the codons

encoding the same two consecutive amino acids.

These results suggest that sequences with a high usage frequency of specific codon

combinations are less biased in their directionality of use, while those with a low usage frequency

tend to be directionally biased toward one of the two possible combinations. This strategy appears

reasonable because if the usage frequency of two codon combinations is biased, the entropy is low,

but if both are used in equal proportions, the entropy is maximized. Moreover, when there is a large

difference in rigidity, there is a corresponding difference observed in the distribution. The opposite

tendency may be observed when the difference in rigidity is small. In most cases, however, the

frequency of use of these two-codon combinations encoding the same amino acid was low relative to

the total frequency of all two-codon combinations, so it is possible that the structure of the genomic

DNA as a whole is not significantly affected by these combinations. It is expected that similar

relationships would be observed for codon combinations encoding other amino acids, as the genetic

code and its physical properties are both sides of the same coin.

Discussion

Prokaryotic genomic DNA is packed within the limited space of prokaryotic cells, and the

amount of space available for packing prokaryotic genomes is difficult to expand from the viewpoint

of the biochemical reactions occurring within, but it is often necessary to increase the genome size

from the viewpoint of evolution. Besides, since 80%–90% of prokaryotic genomic DNA comprises

protein-coding regions using three-base units, it is considered likely that the arrangement of genetic

codes—the codon context—and the physical properties of genomic DNA have coevolved. The

findings of this study indicate that even when different codons encode two synonymous amino acids,

a bias in usage frequency is observed due to the differences in rigidity arising from the different

codon combinations. These results suggest that the genetic code avoids simplifying codon

combinations and evolves in concert with mechanical properties such as DNA rigidity, all while

maintaining complexity.

From the above hypothesis, it would appear that prokaryotic genomic DNA maintains the

evolutionary robustness of an organism by diversifying the combination of genetic codes under the

limitations of restricted storage space in prokaryotic cells. It is possible that the evolution of tRNAs

is also significantly affected by this aspect. In cases where codon combinations are affected by the

storage space afforded to genomic DNA, the evolution of tRNAs as the decoder of codons may also

be affected by the physical properties of the genetic code. It is assumed that such characteristics are

driving forces in producing differences in codon usage among species.

Furthermore, the differences in genomic DNA size among prokaryotic species are due to

differences in the total number of genes encoded by the genomic DNA. In other words, differences in

the quantity of genetic information and the structure thereof create the observed species differences

in genomic DNA. These differences could be the result of the environmental adaptation and

diversification of prokaryotes. Some prokaryotes have a larger genome than *Saccharomyces*

cerevisiae (1.2×10^7 bp), and the average rigidity of their genomic DNAs is around the lower limit

of the rigidity of prokaryotic genomic DNA determined here (shown in Fig 2A). From this

perspective, it is possible that eukaryotes have acquired introns and exons to overcome the

limitations exerted on codon placement strategies by the mechanical constraints evident in

prokaryotic genomes. It is possible that eukaryotic genomic DNA was released from these

mechanical restraints by dividing the genome into ORFs, and even further diversification was made

possible by acquiring nucleosomes that store genomic DNA efficiently. Therefore, the rigidity of

genomic DNA may have been one of the driving forces in the evolution from prokaryote to

eukaryote.

In recent years, researchers in synthetic biology have synthesized prokaryotic total genomic

DNAs [5-7]. However, full, made-from-scratch prokaryotic genomes have not yet been designed.

The findings of the current study should facilitate the future design of synthetic whole prokaryotic

genomes.

Methods
293

Source of complete prokaryotic genome sequences

The NCBI GenBank (https://www.ncbi.nlm.nih.gov/genbank/) and RefSeq (https://www.ncbi.nlm.nih.gov/refseq/) assemblies, July 08, 2015, updated version, were used as sources of the prokaryotic genomic sequences, including those of eubacteria and archaea. After excluding duplicates and incomplete sequences, 2,943 sequences were obtained.

Data processing

All data processing was performed using Python (https://www.python.org/). The SciPy library (https://www.scipy.org/) was used for regression analysis and the Matplotlib library (https://matplotlib.org/) was used for plotting.

Calculations of DNA rigidity

The calculations were performed with all 136 tetranucleotide step flexibility parameters based on a potential energy surface model of a tetranucleotide [3]. The flexibility of each tetranucleotide step in a DNA sequence was calculated with a sliding step length of 1 bp [11]. After the tetranucleotide step calculations, the average value of the results was determined for a whole genomic DNA sequence in each species or for two repeated codons.

Entropy scoring of genomic DNA sequences

313   In this study, Shannon entropy was used as the entropy score for the complexity and

314   diversity of the genomic DNA of each species. The Shannon entropy, $H(P)$, of a whole genomic

315   DNA sequence from each species was determined from various tetranucleotides' Ω obtained from

316   the genomic DNA sequence with a sliding step length of 1 bp, based on the following formula [12]:

$$H(P) = - \sum_{A \in \Omega} P(A) log_2 P(A) \# (3)$$

317   where A refers to the count of individual tetranucleotides, $P(A)$ indicates the ratio of A in Ω that

318   consists of genomic DNA sequences sliced by tetranucleotides.

Directionality indexes of codon pairs encoding the same amino acid

321   To analyze bias in the selectivity of codon pairs encoding the same two consecutive amino

322   acids, the directionality index (DI) was used. The DI was calculated using the following formula [9]:

$$DI_{raw} = \left(\frac{B - A}{|B - A|} \right) \left(\frac{(A - E)^2}{E} + \frac{(B - E)^2}{E} \right) \# (4)$$

where A and B are the proportion of each codon combination in codon pairs encoding the same

two consecutive amino acids, and E is the expected ratio of A and B under the null hypothesis

and is equal to $(A + B)/2$. Furthermore, B represents the codon pair that is the reverse

combination of that comprising A, with B being the more rigid of the two codon combinations. The

normalized DI was defined by DI_{raw} in the range from −1.0 to 1.0. A positive DI indicated a bias

toward more rigid codon combinations, while a negative DI indicated a bias toward more flexible

codon combinations. Thus, the DI values indicated both the bias strength and direction.

Acknowledgments

I thank Dr. Hiroyuki Hamada (Kyushu University) for advice on mathematical model construction and other aspects of the study and his insights regarding the manuscript content. I thank Natasha Beeton-Kempen, Ph.D., from Edanz Group (https://en-author-services.edanzgroup.com/) for editing a draft of this manuscript.

References

1. Kimura H, Shimooka Y, Nishikawa J, Miura O, Sugiyama S, Yamada S, et al. The genome folding mechanism in yeast. J Biochem. 2013;154(2):137-47. doi: 10.1093/jb/mvt033.

2. Packer MJ, Dauncey MP, Hunter CA. Sequence-dependent DNA structure: Dinucleotide conformational maps. J Mol Biol. 2000;295(1):71-83. doi: 10.1006/jmbi.1999.3236.

3. Packer MJ, Dauncey MP, Hunter CA. Sequence-dependent DNA structure: Tetranucleotide conformational maps. J Mol Biol. 2000;295(1):85-103. doi: 10.1006/jmbi.1999.3237.

4. Kaplan N, Moore IK, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, Field Y, et al. The DNA-encoded nucleosome organization of a eukaryotic genome. Nature. 2009;458(7236):362-6. doi: 10.1038/nature07667.

5. Gibson DG, Glass IJ, Lartigue C, Noskov VN, Chuang RY, Algire MA, et al. Creation of a bacterial cell controlled by a chemically synthesized genome. Science. 2010;329(5987):52-6.

doi: 10.1126/science.1190719.

6.   Hutchison 3rd CA, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman HM, et al.

Design and synthesis of a minimal bacterial genome. Science. 2016;351(6280):aad6253. doi:

10.1126/science.aad6253.

7.   Fredens J, Wang K, Torre D de la, Funke LFH, Robertson WE, Christova Y, et al. Total synthesis

of *Escherichia coli* with a recoded genome. Nature. 2019;569(7757):514-518. doi:

10.1038/s41586-019-1192-5.

8.   Lieberman-Aiden E, Berkuman NL van, Williams L, Imakaev M, Ragoczy T, Telling A, et al.

Comprehensive mapping of long-range interactions reveals folding principles of the human

genome. Science**.** 2009;326(5950):289-93. doi: 10.1126/science.1181369.

9.   Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, et al. Topological domains in mammalian

genomes identified by analysis of chromatin interactions. Nature**.** 2012;485(7398):376-80. doi:

10.1038/nature11082.

10. Tolstorukov MY, Virnik KM, Adhya S, Zhurkin VB. A-tract clusters may facilitate DNA

packaging in bacterial nucleoid. Nucleic Acids Res. 2005;33(12):3907-18. doi:

10.1093/nar/gki699.

11. Fukue Y, Sumida N, Nishikawa J, Ohyama T. Core promoter elements of eukaryotic genes have

a highly distinctive mechanical property. Nucleic Acids Res. 2004;32(19):5834-40. doi:

10.1093/nar/gkh905.

12. Shannon CE. A mathematical theory of communication. Bell System Technical Journal. 1948;27:

379-423, 623-56.

Supporting information

S1 File. The analysis dataset of genomic DNA sequences of 2,943 prokaryotes. This dataset

includes accession number, description, length of genomic DNA, average rigidity, standard

deviations of rigidity, entropy score, and %GC of the genomic DNA sequences of 2,943 prokaryotes.

S2 File. The analysis dataset of the randomly sorted genomic DNA sequences of 2,943

prokaryotes by single-base unit. This dataset includes accession number, description, length of

genomic DNA, average rigidity, and entropy score of the randomly sorted genomic DNA sequences

of 2,943 prokaryotes by single-base unit.

S3 File. The analysis dataset of the randomly sorted genomic DNA sequences of 2,943

prokaryotes by triple-base unit. This dataset includes accession number, description, length of

genomic DNA, average rigidity, and entropy score of the randomly sorted genomic DNA sequences

of 2,943 prokaryotes by triple-base unit.

Figure 1

Figure 2

Figure 3

Figure 4