Programming bacteria to sense environmental DNA for multiplexed pathogen detection

Yu-Yu Cheng[1]*, Zhengyi Chen[1]*, Xinyun Cao[1], Tyler D. Ross[1],
Tanya G. Falbel[2], Briana M. Burton[2], and Ophelia S. Venturelli[1,2,3]†

[1]Department of Biochemistry, University of Wisconsin - Madison, WI, United States
[2]Department of Bacteriology, University of Wisconsin - Madison, WI, United States
[3]Department of Chemical & Biological Engineering, University of Wisconsin-Madison

*These authors contributed equally to this work

†To whom correspondence should be addressed: venturelli@wisc.edu

ABSTRACT

DNA is a universal and programmable signal of living organisms. Here we developed Cell-based DNA sensors (Cell-Sens) by engineering the naturally competent bacterium *Bacillus subtilis* (*B. subtilis*) to detect specific DNA sequences in the environment. The DNA sensor strains can identify diverse bacterial species including major human pathogens with high specificity and sensitivity. Multiplexed detection of combinations of different species genomic DNA was achieved by coupling the sensing mechanism to orthogonal fluorescent reporters. We demonstrate that the DNA sensors can detect presence of species in a microbial community without requiring DNA purification of the donor species. In sum, the modularity of the living cell-based DNA sensing mechanism and simple measurement procedure could enable programmable DNA sensing for broad applications.

INTRODUCTION

Chemical and electrical signaling in microbial communities play key roles in biofilm development, activation of virulence pathways and symbioses with multicellular organisms[1,2]. These signals can be exploited to control the collective growth or gene expression of the population or mediate interactions between constituent community members[3,4]. For example, circuits have been designed in engineered organisms to sense specific signals produced by pathogens for selective inhibition of growth[5,6]. However, there are limited well-characterized and orthogonal chemical signals systems for building communication networks between strains due to signal crosstalk[7,8]. In addition, there are challenges to engineering these chemical signals for inter-species communication[4,9]. Therefore, new versatile mechanisms are needed for sensing diverse species in microbial communities.

Towards this goal, sensing of bacterial pathogens is a critical and unsolved challenge as new pathogens can emerge[10]. Current methods for pathogen detection include quantitative polymerase chain reaction (qPCR), immunology-based testing, selective culturing, and Next-Generation Sequencing (NGS)[11,12]. Further, diagnostic tools based on CRISPR-Cas nucleases have also been developed[13,14]. While qPCR is sensitive to the concentration of the target sequence, this method requires specialized equipment and trained personnel which may limit its broad deployment. Immunological detection methods have lower sensitivity and specificity than PCR-based techniques but have a faster turnaround time. Due to these limitations, new cost-effective, sensitive, generalizable, and easy to implement pathogen detection methods are needed.

DNA is a universal and programmable biological signal that selectively distinguish among distinct organisms[15]. For example, mammalian immune system can detect the presence of bacterial DNA and execute protective molecular programs[16,17]. We develop a living cell-based sensor (Cell-Sens) by exploiting the natural competence ability of *B. subtilis* to detect unique DNA

sequences present in the environment. The uptake and integration of the programmable target

DNA sequence onto the *B. subtilis* genome enables growth of the DNA sensor by eliminating a

kill switch via homologous recombination (**Figure 1A**). We demonstrate that Cell-Sens can detect

diverse bacteria including major human pathogens (*Escherichia coli* or *E. coli*, *Salmonella*

typhimurium or *S. typhimurium*, *Staphylococcus aureus* or *S. aureus*, and *Clostridium difficile* or

C. difficile)[18–20]. Cell-Sens can detect the presence of donor species in a microbial community,

thus obviating the need for purification of DNA from the target species. In addition, we show that

Cell-Sens can perform multiplexed detection of purified genomic DNA (gDNA) from multiple

species. In sum, Cell-Sens is a low-cost and versatile method for the selective detection of

bacteria present in the environment.

RESULTS

Construction of a living DNA sensor strain

To build a living DNA sensor strain, we exploited the natural competence ability of the well

characterized soil bacterium *B. subtilis*[21]. The natural competence ability of *B. subtilis* enables the

uptake of environmental DNA and integration of specific sequences with sufficient homology into

the genome via homologous recombination[22]. The recombination efficiency of homologous

sequences depends stringently on the sequence length and percent identity, which can be

exploited to build a highly specific DNA sensor[23,24].

To control the ability to enter the natural competence state and enhance transformation

efficiency, we introduced a xylose-regulated (P_{xylA}) master regulator of competence *comK* onto

the *B. subtilis* genome[25]. To select for the transformed sub-population, we designed a circuit for

negative growth selection consisting of a kill switch that is eliminated by homologous

recombination with an target DNA sequence. To this end, we constructed a kill switch composed

of the Isopropyl β-D-1-thiogalactopyranoside (IPTG) regulated toxin-antitoxin system *txpA-ratA*[26]

flanked on the 3' and 5' ends by homology to the target DNA sequence (**Figure S1A**). In the

presence of IPTG, *txpA* inhibits cell growth by inducing cell lysis. TxpA is predicted to have activity

on the membrane and block cell wall synthesis but the exact mechanism of growth inhibition is

unknown. Homologous recombination with the target DNA sequence eliminates the kill switch and

the LacI transcriptional repressor. The LacI repressor regulates both the expression of *txpA-ratA*

and GFP integrated at a different location in the *B. subtilis* genome (**Figure S1B and S1C**). In the

presence of IPTG, the transformed sub-population that has eliminated *txpA-ratA* is able to grow

exponentially and amplifies the GFP fluorescence signal (**Figure 1A and S2A**).

Using this system, we constructed a DNA sensor strain to detect the presence of *E. coli*
(EC sensor) by introducing homology to the *E. coli* MG1655 *xdhABC* operon, encoding genes for
purine catabolism[27]. To characterize the DNA sensor, the EC sensor was exposed to purified *E.*
coli genomic DNA (gDNA) in media supplemented with 50 mM xylose to induce natural
competence (**Figure 1B**). The transformation frequency of the EC sensor was quantified by
antibiotic selection in liquid or solid media in the presence of IPTG. We investigated the
transformation frequency of the EC sensor as a function of homology length to the target DNA
sequence by constructing a set of EC sensor strains that varied in length of homology to the target
DNA sequence. The transformation frequencies of EC sensors with greater than 1 kbp homology
on the 3' and 5' ends of the kill switch were higher in the presence of *E. coli* gDNA than in the
absence of gDNA (**Figure 1C**). In addition, transformation frequency increased with the homology
length (**Figure 1C and Figure S2B**).

A moderate number of transformants was observed in the absence of gDNA and the
transformation frequency in this condition did not vary with the EC sensor homology length. These
transformants consisted of both GFP OFF and ON colonies due to mutations in either *txpA* and/or
lacI (**Figure S3**). The GFP ON colonies contained mutations in *txpA*, suggesting that these
mutations reduced the growth inhibitory activity of TxpA. The GFP OFF colonies contained
mutations in *lacI* near the ligand-binding site, suggesting that they may reduce binding affinity of
IPTG to LacI, which in turn could reduce the expression of TxpA and GFP[28,29]. Using the 2.5 kbp
EC sensor that maximized the DNA sensor transformation frequency, we found that the
transformation frequency increased and saturated by 10 hr (**Figure 1D and S2A**).

Building living DNA sensors to sense major pathogens
Since the target DNA sequences are programmable and modular, we constructed DNA sensors
to detect the opportunistic intestinal pathogens *Salmonella typhimurium*[30] (*S. typhimurium*) or
Clostridioides difficile[31] (*C. difficile*) or skin pathogen *Staphylococcus aureus*[32] (*S. aureus*). We
introduced 2.5 kbp sequences homologous to the pathogenicity island *sipBCDA* in *S. typhimurium*
(ST sensor), the heme biosynthesis pathway *hemEH* in *S. aureus* (SA sensor), or phenylalanyl-
tRNA synthetase *pheST* in *C. difficile* (CD sensor)[33–35]. The target sequences used to construct
each pathogen DNA sensor were either linked to virulence activities of the pathogen or critical for
fitness.

The SA sensor transformation frequency was approximately one order of magnitude
higher than the other sensor strains, suggesting that the identity of the target sequence in addition
to the homology length determines the transformation frequency for each DNA sensor (**Figure**

2A). To test the specificity of the sensors, we performed time-series measurements of absorbance

at 600 nm (OD600) and GFP for each sensor in the presence of individual gDNA (100 ng mL^{-1})

from each donor species (**Figure S4**). To quantify the specificity of each sensor strain for the

gDNA with homology, we analyzed GFP expression at 12 hr using a fluorescence plate reader

(**Figure 2B**). Our results showed significantly higher GFP expression at 12 hr in the presence of

the gDNA with homology to the given DNA sensor strain than in the absence of gDNA (p-

value<0.01). By contrast, the GFP signal in the presence of gDNA without homology to the given

DNA sensor strain was not significantly different from the negative control lacking gDNA except

for the SA sensor in the presence of *S. typhimurium* gDNA (p-value=0.0415). These data indicate

that each DNA sensor strain has high specificity to the correct target DNA sequence.

We evaluated the sensitivity of each sensor strain to individual gDNA with homology by

quantifying the temporal response of each DNA sensor strain to a range of gDNA concentrations

(**Figure S5**). The GFP response time decreased with gDNA concentration for all sensor strains

and the steady-state GFP expression increased with gDNA concentration for the EC, ST and SA

sensor strains (**Figure 2C-2F**). We determined the sensitivity of each DNA sensor strain to the

target sequence by evaluating the statistical significance of the difference in GFP expression at

12 hr with the negative control lacking gDNA (**Figure 2G-2J**). Due to the variation in the temporal

response of each sensor strain to its target sequence, we analyzed the GFP expression in the

saturated fluorescence regime at 12 hr for each sensor strain. Our results showed that the EC

and ST sensors displayed the highest sensitivity of 1 ng mL^{-1} gDNA, whereas the sensitivities of

the SA and CD sensors were lower (4 ng/mL and 62.5 ng mL^{-1}, respectively). The observed

sensitivities of the EC and ST sensors to the target sequence are sufficient for detection of DNA

release in cell culture or cell extracts (**Figure 2K and S6**)[36–38].

We compared sensitivities of Cell-Sens to qPCR using primers that anneal to each target

sequence. Our results showed that qPCR was approximately one order of magnitude more

sensitive than the EC and ST sensors with the highest sensitivity to the target sequence. In sum,

each sensor displayed high sensitivity to the target DNA sequence except for the CD sensor which

displayed moderate sensitivity. Growth of the non-transformed sub-population and GFP ON

escape mutants in liquid culture contributed to the GFP background signal (**Figure S4 and S5**),

which in turn may reduce the sensitivity and specificity of Cell-Sens.

Cell-Sens can perform multiplexed detection of bacterial genomic DNA

To determine whether the sensor strains could detect gDNA from multiple donor species

simultaneously, we used three orthogonal fluorescent reporters including the EC-GFP (EC-G),

ST-RFP (ST-R) and SA-BFP (SA-B) sensor strains. We introduced all combinations of gDNA (100

ng mL^{-1}) from the pathogen donor species into a mixed culture containing EC-G, ST-R and SA-B.

The mixed culture was transferred onto agar plates and the fluorescent imaging was used to

determine the number of RFP, BFP and GFP colonies in each condition (**Figure 3A, 3B, and S7**).

We found that the mixed DNA sensor culture could accurately detect the presence and absence

of each donor species gDNA across all conditions (**Figure 3C-E**). In sum, a mixture of DNA sensor

strains each individually labeled with a unique fluorescent reporter enabled accurate detection of

all possible combinations of donor species gDNA derived from diverse bacterial pathogens.

Cell-Sens can detect diverse donor species in co-culture

Specific bacterial species have been shown to release extracellular DNA (eDNA) in response to

certain environmental stimuli but the mechanisms leading to DNA release and prevalence of DNA

release in microbial communities are unknown[39]. To determine if the DNA sensors could detect

the presence of the donor cells without performing a DNA extraction, we performed high-

throughput co-culture experiments by combining each DNA sensor strain with its corresponding

donor species into a single culture. The cultures contained the antibiotic spectinomycin which

specifically inhibited the growth of the donor species and not the DNA sensor strain to reduce

ecological competition (**Figure 4A**). Following 10 hr, the transformed DNA sensor sub-populations

were selected on agar plates containing antibiotics that inhibit growth of the donor species and

IPTG. By quantifying the fluorescence intensities of the colonies, we demonstrated that all four

DNA sensors can detect the presence each donor species in co-culture without requiring gDNA

purification (**Figure 4B-4E and S8**).

The fluorescence intensity of each DNA sensor strain co-cultured with individual donor

species that did not have homology to the given DNA sensor strain was similar to the negative

control lacking gDNA (**Figure 4B-4E**). In addition, the fluorescence intensity of the EC and CD

sensors co-cultured with each corresponding donor species with homology displayed similar

fluorescence levels to the positive control (PC) supplemented with purified gDNA containing the

target sequence (100 ng mL^{-1}), suggesting that the transformation was efficient in co-culture

(**Figure 4B and 4E**). Time-series eDNA measurements using qPCR demonstrated that each

donor species released eDNA into the environment (**Figure S9A**). The eDNA concentration

released by each donor species in co-culture with the individual DNA sensor strains ranged

between 0.5 ng mL^{-1} for *S. aureus* to 42 ng mL^{-1} for *C. difficile* after 10 hr and was thus lower than

the concentration of purified gDNA used to characterize the DNA sensor strains (**Figure S9A**).

Therefore, the SA sensor was able to detect a lower concentration of gDNA in co-culture with *S.*

aureus than in monoculture supplemented with *S. aureus* gDNA, indicating that the presence of

certain donor species can enhance the sensitivity of Cell-Sens (**Figure 2I and 4D**)[40]. To determine

if the DNA sensing mechanism occurred via natural competence, we introduced DNase I into the

co-culture containing each donor species and the corresponding DNA sensor strain. The

presence of DNase I reduced the transformation frequency to a similar background fluorescence

level of each DNA sensor as the negative control lacking gDNA (**Figure S9B**). In sum, our results

demonstrate that Cell-Sens can accurately detect the presence of diverse species in a microbial

community.

DISCUSSION

By exploiting the natural competence ability of *B. subtilis*, we developed a cell-based DNA sensor

to selectively detect programmable DNA sequences in the environment. We demonstrated that

the DNA sensors achieved high sensitivity to the concentration of target DNA and specificity for

diverse bacteria including major human pathogens. Cell-Sens requires a simple procedure to

detect a given target sequence. First, purified DNA or donor cells are introduced into a culture

containing the DNA sensor strain(s), incubated for a period and then readouts include colony

forming units (CFU) counting or fluorescence measurements. This workflow could be parallelized

using high-throughput automation techniques to simultaneously process many samples[41]. In

addition, the design of the DNA sensor is modular since any target sequence of interest can be

used.

Since homologous recombination has stringent requirements for target sequence length

and percent identity, Cell-Sens could potentially achieve higher specificity than standard

techniques such as PCR due to off-target primer annealing[42]. In addition, new DNA sensor strains

can be constructed and tested within week timescales for emerging pathogens. Finally, the unique

properties of *B. subtilis* enables detection of donor cells without DNA purification. Since DNA

released in co-culture may be more easily taken up and/or recombined onto the recipient

genome[43], the sensitivity of Cell-Sens may be enhanced in a microbial community containing the

target bacterial species than purified DNA.

The performance of the DNA sensor mechanism could be further optimized for real-world

applications. To reduce the background signal, proteins promoting mutagenesis such as *mfd* can

be deleted to reduce the number of escaped mutants once *txpA* is overexpressed[44]. Combining

different selection mechanisms such as toxin overexpression and essential protein degradation

through proteolysis tagging could also improve the strength of negative selection of the non-

transformed sub-population[45]. In addition, GFP could be expressed from a unique promoter

controlled by an orthogonal transcriptional repressor to directly couple the expression of GFP to
homologous recombination with the target DNA sequence. To achieve autonomous sensing
without inducer induction, the kill switch may be coupled to a quorum-sensing systems such as
agrBDCA so that non-transformed cells will be eliminated after the population reaches a threshold
density allowing only the transformed sub-population to survive[46]. Due to the transformation
efficiency and doubling time of the cells, the total experimental time for Cell-Sens to perform
detection of a target DNA sequence was approximately one day. To speed up the DNA detection
workflow, flow cytometry or fluorescence microscopy could be used to quantify fluorescent cells
within hours following transformation before selection[47]. Future work will investigate the
generalizability of the DNA sensor to sensing other organisms beyond bacteria, including viruses,
fungi or cancer cells[48].
219

MATERIALS AND METHODS
Plasmid and strain construction
All sensor strains were derived from *B. subtilis* PY79. The pAX01-comK plasmid was used to
introduce P_{xyl}-*comK* at the *lacA* locus in *B. subtilis* PY79 to enhance the transformation efficiency
in Luria broth (LB)[25,43]. Reporter plasmids (pOSV00170, pOSV00455 and pOSV00456) were
constructed to introduce $P_{hyperspank}$-*gfp*, $P_{hyperspank}$-*rfp*, or $P_{hyperspank}$-*bfp* at the *ycgO* locus[21,49] (**Table
S1**). The kill switch plasmid pOSV00157 was composed of *lacI*, $P_{hyperspank}$-*txpA-ratA*, flanked by
two target sequences, and was introduced at the *amyE* locus. The toxin-antitoxin system *txpA-*
ratA was amplified from purified gDNA of *B. subtilis* 168 by PCR and cloned onto the *amyE*
integration plasmid creating pOSV00157. The target sequences *xdhABC* (Location: 3001505-
3004004 and 3004005-3006504) were PCR amplified from the purified gDNA of *E. coli* MG1655
(NCBI Reference Sequence: NC_000913.3), *sipBCDA* (Location: 3025979-3028478 and
3028479-3030978) from *Salmonella enterica* serovar Typhimurium LT2 ATCC 700720 (NCBI
Reference Sequence: NC_003197.2), *hemEH* (Location: 553-2770 and 2864-5638) from *S.*
aureus DSM 2569 (GenBank: LHUS02000002.1), and *pheST* (Location: 770923-773144 and
773157-775686) from *C. difficile* DSM 27147 (GenBank: FN545816.1). These sequences were
used to construct a set of plasmids (pOSV00169, pOSV00205, pOSV00206, pOSV00207,
pOS00208, pOSV00292, pOSV00459 and pOSV00475) using restriction enzymes BamHI-HF
(New England Biolabs) and EcoRI-HF (New England Biolabs) or Golden Gate Assembly Mix (New
England Biolabs). A nucleotide BLAST search (megablast with default setting) was used to
determine if the target sequences were unique to each of the species used in our study.

Plasmids were cloned in *E. coli* DH5α and transformed into *B. subtilis* using MC medium[50].
Plasmids were extracted using Plasmid Miniprep Kit (Qiagen). MC medium is composed of 10.7
g/L potassium phosphate dibasic (Chem-Impex International), 5.2 g/L potassium phosphate
monobasic (MilliporeSigma), 20 g/L glucose (MilliporeSigma), 0.88 g/L sodium citrate dihydrate
(MilliporeSigma), 0.022 g/L ferric ammonium citrate (MilliporeSigma), 1 g/L Oxoid casein
hydrolysate (Thermo Fisher Scientific), 2.2 g/L potassium L-glutamate (MilliporeSigma), and 20
mM magnesium sulfate (MilliporeSigma). *B. subtilis*, *E. coli*, *S. typhimurium*, and *S. aureus* were
cultured in Lennox LB medium (MilliporeSigma) for gDNA extraction. *C. difficile* was cultured in
YBHI medium in an anaerobic chamber (Coy Laboratory). YBHI medium is Brain-Heart Infusion
Medium (Acumedia Lab) supplemented with 0.5% Bacto Yeast Extract (Thermo Fisher Scientific),
1 mg/mL D-Cellobiose (MilliporeSigma), 1 mg/mL D-maltose (MilliporeSigma), and 0.5 mg/mL L-
cysteine (MilliporeSigma). The gDNA from each species was extracted using DNeasy Blood &
Tissue Kit (Qiagen). For *S. aureus* gDNA extraction used for PCR target sequence or
transformation experiments, 0.1 mg/mL Lysostaphin (MilliporeSigma) was added in the
pretreatment step. For building the DNA sensor strains, a double crossover and thus stable
integration of each plasmid introduced into *B. subtilis* PY79 was confirmed by the replacement of
a different antibiotic resistance gene at the integration locus. gDNA of modified *B. subtilis* was
then extracted and transformed into another modified *B. subtilis* to introduce multiple
modifications into the genome. Bacterial strains are listed in **Table S2**. DNA sequences of genetic
parts are listed in **Table S3**.

gDNA detection experiments
The DNA sensor strains were inoculated from a -80 °C glycerol stock into LB with 100 µg/mL
spectinomycin and incubated at 37 °C with shaking (250 rpm) for 14 hours. The OD600 was
measured by NanoDrop One (Thermo Fisher Scientific) and then the DNA sensor strains then
diluted to OD600 of 0.1 in 1 mL LB in 14 mL Falcon™ Round-Bottom Tube (Thermo Fisher
Scientific) supplemented with 50 mM xylose (Thermo Fisher Scientific) and 100 µg/mL
spectinomycin (Dot Scientific). Xylose was added to induce the competence and spectinomycin
was used to prevent the contamination of other bacteria. Purified gDNA from each donor species
was quantified by a Quant-iT dsDNA Assay Kit (Thermo Fisher Scientific) and added to the
cultures with known concentration. The positive control in all experiments was 100 ng/mL gDNA
from a given donor species. The DNA sensor strains cultures were incubated at 37°C with shaking
(250 rpm) for 10 hours and then transferred to liquid LB with a 1:20 dilution or 20 µL was plated
on an LB plate for CFU counting. The liquid or solid LB media was supplemented with 2 mM IPTG

(Bioline), 5 µg/mL chloramphenicol (MilliporeSigma), and MLS (1 µg/mL erythromycin from

Sigma-Aldrich and 25 µg/mL lincomycin from Thermo Fisher Scientific). In liquid culture, OD600

and fluorescence were measured using a SPARK Multimode Microplate Reader (TECAN). In

these experiments, 100 uL of the 1:20 diluted cultures of transformed DNA sensors were

transferred to 96-well black and clear-bottom CELLSTAR® format sterile cell culture microplates

(Greiner Bio-One). Plates were sealed with Breathe-Easy Adhesive Microplate Seals (Thermo

Fisher Scientific) and incubated in the microplate reader at 37 °C with shaking for time-series OD

and GFP measurements. On agar plate, CFU of transformed *B. subtilis* and total *B. subtilis* were

counted for both fluorescent and non-fluorescent colonies. To determine the CFU of transformed

B. subtilis, 20 µL of cell culture was plated on LB agar plate with 2 mM IPTG, 5 µg/mL

chloramphenicol and MLS. To determine the CFU of total *B. subtilis*, cell culture was diluted in

phosphate-buffered saline (PBS) (Dot Scientific) and plated on LB agar plate with 5 µg/mL

chloramphenicol and MLS. Colonies were imaged using an Azure Imaging System 300 (Azure

Biosystems) using Epi Blue LED Light Imaging with 50 millisecond exposure time to determine

the fluorescence expression. Transformation frequency was defined as the ratio of fluorescent or

non-fluorescent transformed *B. subtilis* to the total number of *B. subtilis* in **Figure 1C and 1D**.

Transformation frequency was defined as the ratio of fluorescent transformed *B. subtilis* to the

total number of *B. subtilis* in **Figure 2A and S9B**.

Multiplexed gDNA detection experiment

The DNA sensor strains EC-G, ST-R, and SA-B sensors were diluted to an OD600 of 0.1, 0.1,

and 0.01, respectively, in 1 mL LB supplemented with 50 mM xylose to induce expression of *comK*

(Thermo Fisher Scientific) and 100 µg/mL spectinomycin (Dot Scientific). Different combinations

of 100 ng/mL purified gDNA from each donor species were introduced into a mixed culture

containing different DNA sensor strains (EC-G, ST-R and SA-B). The DNA sensor strain mixtures

were incubated at 37 °C with shaking (250 rpm) for 10 hours and 5 µL of each culture was plated

on 6-well plates (Thermo Fisher Scientific). The 6-well plates were large enough to visualize

approximately a hundred colonies in each well and could be imaged by the Nikon Ti II Microscope.

In these plates, each well contained solidified LB agar supplemented with 2 mM IPTG, 5 µg/mL

chloramphenicol, and MLS. These plates were incubated at 37 °C overnight.

On the next day, each well was imaged using a Nikon Eclipse Ti2-E Microscope.

Brightfield images were collected at 4X magnification using the built-in transilluminator of the

microscope. The epifluorescence light source was the SOLA-III Light Engine (Lumencor) and

standard band filter cubes (Nikon) including DAPI (Excitation: 375/28, Emission: 460/50), FITC

(Excitation: 480/30, Emission: 535/40) and TRITC (Excitation: 540/25, Emission: 605/55) filter

cubes were used to image BFP, GFP, and RFP, respectively. The exposure times for BFP, GFP,

RFP were 500 ms, 9 ms, and 50 ms, respectively. Since the DAPI filter was not optimized for BFP,

a strong BFP signal was observed from the LB agar. To reduce the autofluorescence from LB

agar, images were processed as follows: complete images of each LB agar well were generated

from multipoint images using the stitching function in the Nikon NIS Elements software. Once the

full images were assembled, each of the four channels were mapped to unity by the minimum

and maximum pixel values of all images for that channel using ImageJ. Fluorescence channels

were further processed to remove background signal from agar fluorescence by masking out all

signals not associated with a colony. Masks for this step were generated from the DIC images

using the pixel classification tool in Ilastik[51]. Any remaining noise in the fluorescence channels

was reduced by clipping each image by the average noise value for that channel.

Co-culture experiments

E. coli, *S. typhimurium*, and *S. aureus* and the corresponding DNA sensor strains were each

inoculated from a -80 °C glycerol stock into LB media and incubated at 37 °C with shaking (250

rpm) overnight. *C. difficile* was cultured in YBHI medium in an anaerobic chamber (Coy

Laboratory). On the next day, the cultures were diluted to an OD600 of 0.1 each in 100 µL LB

supplemented with 50 mM xylose and 100 µg/mL spectinomycin into 96-well plates at 37 °C with

shaking (250 rpm). Overnight cultures of target bacteria were also diluted in PBS (Dot Scientific)

and plated onto an LB or YBHI agar plate to determine the initial CFU of target bacteria in the co-

culture, which was 2.54×10^6 cells per mL, 1.19×10^7 cells per mL, 1.22×10^7 cells per mL, and

1.05×10^6 cells per mL for *E. coli*, *S. typhimurium*, *S. aureus*, and *C. difficile*, respectively.

Following 10 hours of incubation, cell culture were diluted into PBS with 1:10 dilution and 10 uL

diluted cell culture were spotted on a large LB agar plate (140 mm diameter and 20 mm height

petri dish, Thermo Fisher Scientific). The LB agar was supplemented with 2 mM IPTG, 5 µg/mL

chloramphenicol, and MLS and incubated overnight at 37 °C. On the next day, each plate was

imaged with an Azure Imaging System 300 (Azure Biosystems) using the Epi Blue LED Light

Imaging with 50 millisecond exposure time. The pixel fluorescence intensity of the colony spots

was quantified by ImageJ[52].

To determine if the detection of target bacteria was via transformation, 1 unit/mL DNase I

(Thermo Fisher Scientific) was added to the 1 mL of each co-culture and CFU of both transformed

B. subtilis and total *B. subtilis* were counted to calculate the transformation frequency change. To

quantify the extracellular DNA (eDNA), a given DNA sensor strain and donor species were diluted

into OD0.1 each in 3 mL LB containing 50 mM xylose and 100 µg/mL spectinomycin and
incubated at 37 °C with shaking (250 rpm) for 10 h. Because the sensor also contained the same
DNA sequence targeted by the qPCR primers, to quantify the eDNA released from only the target
bacteria but not the sensor, a sensor strain (msOSV00409) without homologous sequence was
used in the co-culture instead. At every two hours, supernatants were collected and filtered with
0.2 µm Whatman Puradisc Polyethersulfone Syringe Filter (GE Healthcare) to remove the cells,
and stored in the -20 °C freezer for qPCR measurements.
350
qPCR measurements
To determine the detection limit of qPCR, purified gDNA from *E. coli*, *S. typhimurium*, *S. aureus*,
or *C. difficile* was extracted using DNeasy Blood & Tissue Kit (Qiagen) and quantified by Quant-
iT dsDNA Assay Kit (Thermo Fisher Scientific). The purified gDNA samples were serially diluted
using a 1:10 dilution from 10 µg/mL to 10^{-5} µg/mL and mixed with 2X SsoAdvanced Universal
SYBR® Green Supermix (Bio-Rad) and 500 nM of each primer (**Table S4**). Primers were designed
and synthetized by Integrated DNA Technologies to target the sequences used to construct each
DNA sensor strain. The qPCR reactions were performed on the CFX Connect Real-Time PCR
Detection System (Bio-Rad). The gDNA concentration produced at least 95% positive replicates
(no missing Ct values) is defined as the detection limit[53]. To quantify the rate of eDNA release of
E. coli, *S. typhimurium*, *S. aureus*, or *C. difficile* in each co-culture, gDNA standards previously
quantified by a Quant-iT dsDNA Assay Kit (Thermo Fisher Scientific) were included in the qPCR
run and the gDNA concentration of the supernatant collected at different time points was inferred
from the standard curve based on a titration of purified gDNA.
365

366 **DATA AVAILABILITY**
367 All data presented in this manuscript will be available from Zenodo. Plasmids and strains are
368 available upon request.
369

370 **CODE AVAILABILITY**
371 Code for data analysis will be available from Github. The microscopic images of multiplexed
372 detection were processed by ImageJ and ilastik to reduce the background fluorescence from the
373 agar using the basic features such as segmentation and pixel classification and no code was
374 generated in this process.
375

ACKNOWLEDGEMENTS

We would like to thank Dr. Scott Coyle and Zhejing Xu for providing access to key equipment
used in this study. This work was supported by the Defense Advanced Research Projects Agency
(DARPA) Grant under grant number HR0011-19-2-0002, Multi University Research Initiative
(MURI) under grant number W911NF-19-1-0269, National Institutes of Health under grant number
R35GM124774, National Institute of Biomedical Imaging and Bioengineering under grant number
R01EB030340.
383

AUTHOR CONTRIBUTIONS

Y.Y.C. and O.S.V. conceived the research. Y.Y.C., and Z.C. performed the experiments. Y.Y.C., Z.C., and X.C. constructed the sensors. T.D.R. developed custom code for data analysis. T.G.F. assisted with the strain construction. Y.Y.C. and O.S.V. wrote the manuscript. O.S.V., B.M.B., T.G.F., and Z.C. analyzed the data. All authors provided feedback on the manuscript. O.S.V. and B.M.B. secured the funding.

COMPETING INTERESTS

O.S.V., Y.Y.C. and Z.C. are inventors on a provisional patent application filed by the Wisconsin Alumni Research Foundation (WARF) with the US Patent and Trademark Office, which describes and claims concepts disclosed herein (Application No. 63/290,442 Filing Date: 12/16/2021). All other authors declare no conflict of interest.

REFERENCES

1. Prindle, A. *et al.* Ion channels enable electrical communication in bacterial communities. *Nature* **527**, (2015).
2. Miller, M. B. & Bassler, B. L. Quorum sensing in bacteria. *Annual Review of Microbiology* **55**, (2001).
3. Danino, T., Mondragón-Palomino, O., Tsimring, L. & Hasty, J. A synchronized quorum of genetic clocks. *Nature* **463**, (2010).
4. Du, P. *et al.* De novo design of an intercellular signaling toolbox for multi-channel cell–cell communication and biological computation. *Nat. Commun.* **11**, (2020).
5. Saeidi, N. *et al.* Engineering microbes to sense and eradicate Pseudomonas aeruginosa, a human pathogen. *Mol. Syst. Biol.* **7**, (2011).
6. Borrero, J., Chen, Y., Dunny, G. M. & Kaznessis, Y. N. Modified lactic acid bacteria detect and inhibit multiresistant enterococci. *ACS Synth. Biol.* **4**, (2015).
7. Scott, S. R. & Hasty, J. Quorum Sensing Communication Modules for Microbial Consortia. *ACS Synth. Biol.* **5**, (2016).
8. Sexton, J. T. & Tabor, J. J. Multiplexing cell-cell communication. *Mol. Syst. Biol.* **16**, (2020).
9. Marchand, N. & Collins, C. H. Peptide-based communication system enables Escherichia coli to Bacillus megaterium interspecies signaling. *Biotechnol. Bioeng.* **110**, (2013).

10. Morens, D. M. & Fauci, A. S. Emerging Pandemic Diseases: How We Got to COVID-19.
Cell **182**, (2020).
11. Iqbal, S. S. *et al.* A review of molecular recognition technologies for detection of biological
threat agents. *Biosens. Bioelectron.* **15**, (2000).
12. Chiu, C. Y. & Miller, S. A. Clinical metagenomics. *Nature Reviews Genetics* **20**, (2019).
13. East-Seletsky, A. *et al.* Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA
processing and RNA detection. *Nature* **538**, (2016).
14. Gootenberg, J. S. *et al.* Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science
(80-.).* **356**, (2017).
15. Ortiz, M. E. & Endy, D. Engineered cell-cell communication via DNA messaging. *J. Biol.
Eng.* **6**, (2012).
16. Hemmi, H. *et al.* A Toll-like receptor recognizes bacterial DNA. *Nature* **408**, (2000).
17. Krieg, A. M. CpG motifs in bacterial DNA and their immune effects. *Annual Review of
Immunology* **20**, (2002).
18. Fàbrega, A. & Vila, J. Salmonella enterica serovar Typhimurium skills to succeed in the
host: Virulence and regulation. *Clinical Microbiology Reviews* **26**, (2013).
19. Tong, S. Y. C., Davis, J. S., Eichenberger, E., Holland, T. L. & Fowler, V. G.
Staphylococcus aureus infections: Epidemiology, pathophysiology, clinical
manifestations, and management. *Clin. Microbiol. Rev.* **28**, (2015).
20. Rupnik, M., Wilcox, M. H. & Gerding, D. N. Clostridium difficile infection: New
developments in epidemiology and pathogenesis. *Nature Reviews Microbiology* **7**,
(2009).
21. Popp, P. F., Dotzler, M., Radeck, J., Bartels, J. & Mascher, T. The Bacillus BioBrick Box
2.0: Expanding the genetic toolbox for the standardized work with Bacillus subtilis. *Sci.
Rep.* **7**, (2017).
22. Dubnau, D. Genetic competence in Bacillus subtilis. *Microbiological Reviews* (1991).
23. Ji, M. *et al.* Engineering Bacillus subtilis ATCC 6051a for the production of recombinant
catalases. *J. Ind. Microbiol. Biotechnol.* **48**, (2021).
24. Carrasco, B., Serrano, E., Martín-González, A., Moreno-Herrero, F. & Alonso, J. C.
Bacillus subtilis MutS modulates RecA-mediated DNA strand exchange between
divergent DNA sequences. *Front. Microbiol.* (2019). doi:10.3389/fmicb.2019.00237
25. Zhang, X.-Z. & Zhang, Y.-H. P. Simple, fast and high-efficiency transformation system for
directed evolution of cellulase in Bacillus subtilis. *Microb. Biotechnol.* **4**, 98–105 (2011).
26. Silvaggi, J. M., Perkins, J. B. & Losick, R. Small untranslated RNA antitoxin in Bacillus
subtilis. *J. Bacteriol.* **187**, (2005).
27. Xi, H., Schneider, B. L. & Reitzer, L. Purine catabolism in Escherichia coli and function of
xanthine dehydrogenase in purine salvage. *J. Bacteriol.* **182**, (2000).
28. Tack, D. S. *et al.* The genotype-phenotype landscape of an allosteric protein. *Mol. Syst.
Biol.* **17**, (2021).
29. Kleina, L. G. & Miller, J. H. Genetic studies of the lac repressor. XIII. Extensive amino
acid replacements generated by the use of natural and synthetic nonsense suppressors.
J. Mol. Biol. **212**, (1990).
30. McClelland, M. *et al.* Complete genome sequence of Salmonella enterica serovar
Typhimurium LT2. *Nature* **413**, (2001).
31. Stabler, R. A. *et al.* Comparative genome and phenotypic analysis of Clostridium difficile
027 strains provides insight into the evolution of a hypervirulent bacterium. *Genome Biol.*
10, (2009).
32. Soni, I., Chakrapani, H. & Chopra, S. Draft genome sequence of methicillin-sensitive
Staphylococcus aureus ATCC 29213. *Genome Announc.* **3**, (2015).
33. Tucker, S. C. & Galán, J. E. Complex function for SicA, a Salmonella enterica serovar
typhimurium type III secretion-associated chaperone. *J. Bacteriol.* **182**, (2000).

34. Lobo, S. A. L. *et al.* Staphylococcus aureus haem biosynthesis: Characterisation of the
enzymes involved in final steps of the pathway. *Mol. Microbiol.* **97**, (2015).
35. Beyer, D. *et al.* New Class of Bacterial Phenylalanyl-tRNA Synthetase Inhibitors with High
Potency and Broad-Spectrum Activity. *Antimicrob. Agents Chemother.* **48**, (2004).
36. Matsui, K., Ishii, N. & Kawabata, Z. Release of extracellular transformable plasmid DNA
from Escherichia coli cocultivated with algae. *Appl. Environ. Microbiol.* **69**, 2399–2404
(2003).
37. Lorenz, M. G., Gerjets, D. & Wackernagel, W. Release of transforming plasmid and
chromosomal DNA from two cultured soil bacteria. *Arch. Microbiol.* **156**, 319–326 (1991).
38. Zou, Y. *et al.* Nucleic acid purification from plants, animals and microbes in under 30
seconds. *PLoS Biol.* **15**, (2017).
39. Ibáñez de Aldecoa, A. L., Zafra, O. & González-Pastor, J. E. Mechanisms and regulation
of extracellular DNA release and its biological roles in microbial communities. *Front.*
Microbiol. **8**, 1390 (2017).
40. Zhang, X. *et al.* Stress-induced, highly efficient, donor cell-dependent cell-to-cell natural
transformation in Bacillus subtilis. *J. Bacteriol.* **200**, e00267-18 (2018).
41. McBride, M. T. *et al.* Autonomous Detection of Aerosolized Bacillus anthracis and
Yersinia pestis. *Anal. Chem.* **75**, (2003).
42. Bustin, S. & Huggett, J. qPCR primer design revisited. *Biomolecular Detection and*
Quantification **14**, (2017).
43. Cheng, Y.-Y., Papadopoulos, J. M., Falbel, T., Burton, B. M. & Venturelli, O. S. Donor-
recipient interactions drive dynamics of horizontal gene transfer via natural competence.
bioRxiv 2020.10.19.342733 (2020). doi:10.1101/2020.10.19.342733
44. Ragheb, M. N., Merrikh, C., Browning, K. & Merrikh, H. Mfd regulates RNA polymerase
association with hard-to-transcribe regions in vivo, especially those with structured RNAs.
Proc. Natl. Acad. Sci. U. S. A. **118**, (2020).
45. Chan, C. T. Y., Lee, J. W., Cameron, D. E., Bashor, C. J. & Collins, J. J. 'Deadman' and
'Passcode' microbial kill switches for bacterial containment. *Nat. Chem. Biol.* **12**, (2016).
46. Marchand, N. & Collins, C. H. Synthetic Quorum Sensing and Cell-Cell Communication in
Gram-Positive Bacillus megaterium. *ACS Synth. Biol.* **5**, (2016).
47. Kurushima, J. *et al.* Unbiased homologous recombination during pneumococcal
transformation allows for multiple chromosomal integration events. *Elife* **9**, (2020).
48. Cooper, R. M. *et al.* Engineered bacteria detect tumor DNA in vivo. *bioRxiv*
2021.09.10.459858 (2021). doi:10.1101/2021.09.10.459858
49. Overkamp, W. *et al.* Benchmarking various green fluorescent protein variants in Bacillus
subtilis, Streptococcus pneumoniae, and Lactococcus lactis for live cell imaging. *Appl.*
Environ. Microbiol. **79**, 6481–6490 (2013).
50. Konkol, M. A., Blair, K. M. & Kearns, D. B. Plasmid-encoded comI inhibits competence in
the ancestral 3610 strain of Bacillus subtilis. *J. Bacteriol.* **195**, (2013).
51. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods*
16, 1226–1232 (2019).
52. Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nature*
Methods **9**, (2012).
53. Forootan, A. *et al.* Methods to determine limit of detection and limit of quantification in
quantitative real-time PCR (qPCR). *Biomol. Detect. Quantif.* **12**, (2017).
514
515

Figure 1. Construction and characterization of a cell-based DNA sensor (Cell-Sens). (A)
Schematic of a synthetic genetic circuit in *Bacillus subtilis* that can specifically recognize
programmable target DNA sequences. The gene expression of the toxin-antitoxin system *txpA-ratA* and a fluorescent reporter *gfp* integrated onto the genome in a different position are regulated
by the IPTG-inducible P$_{hyperspank}$ promoter. In the presence of the target DNA sequence, DNA
uptake and homologous recombination via natural competence can eliminate the toxin and the
LacI repressor. Consequently, growth of the GFP expressing transformed sub-population is
permitted in the presence of IPTG and the target DNA sequence. **(B)** Schematic of the purified
DNA detection procedure for Cell-Sens. The DNA sensor is inoculated into a liquid culture
containing the target DNA sequence. Xylose controls the competence state of *B. subtilis* and
enhances transformation efficiency by inducing the expression of the master regulator of
competence *comK*. The fluorescence output of the DNA sensor can be quantified in liquid culture
or solid media supplemented with IPTG, which induces expression of the kill switch and GFP and
thus selects for the transformed sub-population. **(C)** Line plot of homology length versus
transformation frequency for the *E. coli* DNA sensor strain targeting sequences in *xdhABC* in the
presence (solid lines) and absence (dashed lines) of 100 ng/mL purified *E. coli* genomic DNA. Kill
switch escape mutants that can grow in the presence of IPTG and absence of gDNA either
express GFP (triangles) or are not fluorescent (circles). Data points represent biological replicates
and lines are the average of three biological replicates. **(D)** Line plot of time versus transformation

  frequency for the *E. coli* DNA sensor strain. Data points represent biological replicates and lines
  are the average of three biological replicates.

Figure 2. Cell-Sens can detect diverse bacterial species including major human pathogens.
(A) Transformation frequencies of DNA sensor strains that can detect *E. coli* (EC sensor), *S. typhimurium* (ST sensor), *S. aureus* (SA sensor), or *C. difficile* (CD sensor) in the presence of 100 ng/mL purified gDNA from each target bacteria (triangles) or no gDNA (circles) at 10 hr. Bar height represents the mean of biological replicates. Data points represent biological replicates (n = 3). **(B)** GFP expression of each DNA sensor strain at 12 hr in response to 100 ng/mL purified

gDNA of a given donor species or no gDNA representing the negative control (NC). DNA sensor
strains were cultured in LB media supplemented with 2 mM IPTG, 5 µg/mL chloramphenicol and
MLS (1 µg/mL erythromycin and 25 µg/mL lincomycin) for 12 h. Data are the average of four
biological replicates. GFP fold changes were 2.6, 2.6, 2.1, and 1.5 for the EC, ST, SA, or CD
sensor in the presence of *E. coli*, *S. typhimurium*, *S. aureus* or *C. difficile* purified gDNA,
respectively compared to NC with *p*-values of 8.5×10^{-3}, 4.5×10^{-5}, 1.3×10^{-4}, and 3.5×10^{-4},
respectively based on an unpaired *t*-test. Expression of GFP as a function of time for the **(C)** EC
sensor, **(D)** ST sensor, **(E)** SA sensor, or **(F)** CD sensor transformed with different concentrations
of target bacterial gDNA. Lines represent the mean of biological replicates (n=4). Shaded region
denotes 1 s.d. from the mean. Expression of GFP as a function of gDNA concentration at 12 hr
for the **(G)** EC sensor, **(H)** ST sensor, **(I)** SA sensor, and **(J)** CD sensor. Unpaired *t*-test was
performed on GFP fluorescence in the presence and absence of gDNA where *, **, or *** denotes
p-values < 0.05, 0.01 or 0.001, respectively. Horizontal line denotes the mean GFP fluorescence
in the absence of gDNA (n=4). **(K)** Horizontal bar plot denoting the ability to detect gDNA for each
DNA sensor strain based on data in **(G)**-**(J)** or based on quantitative real-time PCR. Solid bars
indicate gDNA concentrations where the given DNA sensor strain displayed higher GFP
fluorescence than the no gDNA condition based on the unpaired t-test in **(G)**-**(J)**. For the qPCR
data, dashed bars represent gDNA concentration conditions where Ct values were determined in
each of the three technical replicates.

Figure 3. Cell-Sens can perform multiplexed detection of diverse bacterial genomic DNA. **(A)** Schematic of the experimental design for multiplexed detection of genomic DNA (gDNA) of *E. coli*, *S. typhimurium*, and *S. aureus*. The *E. coli*, *S. typhimurium*, and *S. aureus* DNA sensor strains were labeled GFP (EC-G), RFP (ST-R) and BFP (SA-B), respectively. EC-G, ST-R, and SA-B sensors were combined into a mixed culture and exposed to all combinations of gDNA from each species (100 ng/mL each). The mixed culture of DNA sensor strains supplemented with combinations of species' gDNA was incubated for 10 hours and then plated onto 6-well LB agar plates containing 2 mM IPTG, 5 μg/mL chloramphenicol and MLS (1 μg/mL erythromycin and 25 μg/mL lincomycin). Fluorescent imaging was used to quantify the number of GFP, RFP and BFP expressing *B. subtilis* colonies in each condition. **(B)** Fluorescent microscopy images of GFP, RFP and BFP in the presence of different combinations of gDNA from *E. coli*, *S. typhimurium*, and/or *S. aureus*. Bar plots of the number of colonies expressing each fluorescent reporter in each condition for the **(C)** EC-G sensor, **(D)** ST-R sensor, and **(E)** SA-B sensor. Bar height represents the mean of biological replicates (n = 3). Data points denote biological replicates (n=3).

583
Figure 4. Cell-Sens can detect diverse bacteria in co-culture. **(A)** Schematic of the
experimental design for co-culturing each DNA sensor strain with a given donor species in 96-
well plates with an initial OD600 of 0.1 for each species. The pairwise communities were co-
cultured 10 hours in LB containing xylose (50 mM) and spectinomycin (100 μg/mL) and then
plated onto LB plates containing 2 mM IPTG, 5 μg/mL chloramphenicol and MLS (1 μg/mL
erythromycin and 25 μg/mL lincomycin). Antibiotics inhibit the growth of the donor species.
Fluorescence microscopy image (right) of *B. subtilis* DNA sensor strain colonies on the agar plate
for each co-culture condition with three biological replicates (n=3). Pixel average fluorescence
intensity of colonies on agar plates were analyzed for the **(B)** EC sensor, **(C)** ST sensor, **(D)** SA
sensor, or **(E)** CD sensor. A positive control of 100 ng/mL target bacterial gDNA was included
(PC) and no gDNA representing the negative control (NC). Unpaired *t*-test was performed in the
presence and absence (NC) of the donor species where ** and *** denote *p*-values < 0.01 or
0.001, respectively.