Influenza A Virus Exacerbates Group A Streptococcus Infection and Thwarts Anti-bacterial Inflammatory Responses in Murine Macrophages

Johann Aleith[1,*], Maria Brendel[1], Erik Weipert[1], Michael Müller[1], Daniel Schultz[2], Ko-Infekt Study Group[2,3,4] , Brigitte Müller-Hilke[1]

[1]Core Facility for Cell Sorting and Cell Analysis, Rostock University Medical Center, Rostock, Germany

[2]Institute of Biochemistry, University of Greifswald, Greifswald, Germany

[3]Institute of Immunology, Friedrich-Loeffler-Institut, Greifswald-Insel Riems, Germany

[4]Institute of Medical Microbiology, Virology and Hygiene, Rostock University Medical Center, Rostock, Germany

*Correspondence: johann.aleith@med.uni-rostock.de

Abstract

Seasonal influenza epidemics pose a considerable hazard for global health. In the past decades, accumulating evidence revealed that influenza A virus (IAV) renders the host vulnerable to bacterial superinfections which in turn are a major cause for morbidity and mortality. However, whether the impact of influenza on anti-bacterial innate immunity is restricted to the vicinity of the lung or systemically extends to remote sites is underexplored. We therefore sought to investigate intranasal infection of adult C57BL/6J mice with IAV H1N1 in combination with bacteremia elicited by intravenous application of Group A Streptococcus (GAS). Co-infection *in vivo* was supplemented *in vitro* by challenging murine bone marrow derived macrophages and exploring gene expression and cytokine secretion. Our results show that viral infection of mice caused mild disease and induced the depletion of CCL2 in the periphery. Influenza preceding GAS infection promoted the occurrence of paw edemas and was accompanied by exacerbated disease scores. *In vitro* co-infection of macrophages led to significantly elevated expression of TLR2 and CD80 compared to bacterial mono-infection, whereas CD163 and CD206 were downregulated. The GAS-inducible upregulation of inflammatory genes, such as *Nos2*, as well as the secretion of TNFα and IL-1β were notably reduced or even abrogated following co-infection. Our results indicate that IAV primes an innate immune layout that is inadequately equipped for bacterial clearance.

Keywords: influenza A virus, Group A Streptococcus, co-infection, inflammation, sepsis, macrophage, innate immunity

1   Introduction

Seasonal influenza is a major cause of respiratory disease that affects 5 – 10% of the global population annually with an estimated death toll of up to 500,000 [1,2]. The segmented genome of influenza A virus (IAV) combined with an error-prone RNA polymerase enables the periodical emergence of new strains with elevated pandemic capacities, which annually challenge humankind yet devoid of adequate adaptive immunity [3,4]. The most prominent paradigm for the dramatic consequences of an influenza pandemic is the 1918/1919 flu that caused roughly 50 Million casualties [5]. Notably, the vast majority of fatal cases were attributed to secondary bacterial infections predominantly caused by pneumococci and hemolytic streptococci [6,7]. Along these lines, excess morbidity due to bacterial superinfection with the nasopharyngeal colonizers *S. pneumoniae*, *S. aureus* and *S. pyogenes* (Group A Streptococcus, GAS) was confirmed for the most recent influenza pandemic in 2009 [8]. As of yet, there is neither a licensed vaccine against S. *aureus* nor against *S. pyogenes* that would help contain invasive infections with these pathogens during future influenza pandemics [9–11].

Several modes by which an immune response against IAV supports viral clearance yet fails to oppose bacterial pathogens have been suggested [1]. For instance, Okamoto and colleagues demonstrated that IAV infection led to the presentation of hemagglutinin (HA) by epithelial cells, which is utilized by GAS to breach cellular barriers [12,13]. Other groups reported that HA, among other viral proteins, caused the exposure of receptors that act as adhesins for bacterial attachment and invasion [14–16]. Others showed that viral infection caused damage of the respiratory epithelium, expediting initial bacterial adherence [6,17,18]. Moreover, experimental data indicated that IAV paves the way for the dissemination of opportunistic bacterial pathogens by impacting the innate immune response, which is critical for bacterial containment [19,20]. In fact, the virus was shown to induce an increased secretion of anti-inflammatory interleukin (IL-)10 as

57 well as inflammatory type I and type II interferons (IFNs), which was associated with both, impaired phagocytic activity by pulmonary immune cells and diminished production of chemokines [14,19,21–24].

60 Together, these data illustrate some aspects of post-influenza pneumonia and the interplay of viral and bacterial pneumopathogens in life-threatening infections. While the aforementioned studies focused on bacterial superinfections of the respiratory tract, we were intrigued that seasonal or pandemic influenza outbreaks seem to coincide with a broad spectrum of invasive GAS-associated infectious diseases like necrotizing fasciitis, pneumonia and bacteremia [8,25–30]. We therefore asked whether pulmonary IAV can also alter systemic innate immunity and facilitate secondary bacterial insults at remote sites. We were particularly interested in the impact IAV exerts on the response of macrophages – immune cells that are indispensable for initial anti-streptococcal resistance [19,31,32]. We established co-infection models that *in vivo* combined respiratory IAV infection with GAS bacteremia and *in vitro* investigated primary macrophages for their potential to respond to both pathogens simultaneously.

2 Materials and Methods

2.1 Pathogens

Pandemic influenza A virus (IAV) A/Germany-BY/74/2009 (H1N1pdm09) propagation and titer determination was performed as previously described [33]. In brief, IAV was replicated in Mardin-Darby canine kidney II (MDCKII) cells using minimal essential medium supplemented with 0.2% bovine serum albumin and 2 μg/mL N-Tosyl-L-phenylalanin-chlormethylketon (Sigma). For the determination of the tissue culture infectious dose 50 ($TCID_{50}$), virus suspensions were serially diluted and applied to MDCKII cultures. Cells were then incubated for three days at 37°C and 5% CO_2 followed by examination of cytopathogenicity.

Streptococcus pyogenes (Group A Streptococcus, GAS) strain AP1 of the *emm1* (M1) serotype was originally acquired from the World Health Organization Collaborating Center for Reference and Research on Streptococci (Prague, Czech Republic). Bacteria were thawed onto Colombia agar plates containing 5% sheep blood (Becton Dickinson) and were cultured overnight followed by storage at 4°C for up to three weeks. Colonies were picked from the plate, suspended into Todd-Hewitt broth (THB, Becton Dickinson) and cultured overnight at 37°C and 5% CO_2. The suspension was diluted 20-fold in THB and bacteria were incubated until exponential phase of growth was reached. Subsequently, bacteria where washed thrice with PBS (Thermo Fisher) prior to their application in mice and *in vitro* infection models, respectively. The determination of colony forming units (CFU) was performed the following day by counting of serially diluted suspensions.

2.2 Animals

C57BL/6J mice were initially purchased from Charles River. Mice were bred in the animal core facility under specific germ-free conditions. Animals were transferred to individually ventilated cages prior to infection experiments and were housed at a 12-hour light/dark cycle, an am-

bient temperature of $22 \pm 2°C$ and $50 \pm 20\%$ humidity. Food and water were provided *ad libitum.*

Animal experiments were reviewed and approved by the ethics committee of the State Depart-

ment for Agriculture, Food Safety and Fishery in Mecklenburg-Western Pomerania under the file

reference number 7221.3-1-017/19.

2.3   In vivo infection models and clinical scoring

For the induction of viral infections, 20 µL of a suspension containing 1.5×10^5 TCID$_{50}$

IAV were applied to both nostrils of 20- to 22-week old male mice under anesthesia by isoflurane

inhalation. This volume was chosen in order to guarantee an infection of both, the upper and low-

er respiratory tracts [34]. Applying the same volume of PBS only served as the negative (healthy)

control. Mice were subsequently monitored daily for 16 days for alterations in body weight rela-

tive to the day of infection (day 0). On days 2, 4 and 7, a maximum of 80 µL of anti-coagulated

blood was drawn by saphenous venipuncture using a 25G needle followed by centrifugation and

collection of plasma. On day 16, mice were anesthetized with 75 mg of Ketamine (Pharmanovo)

and 5 mg Xylazin (Bayer) per kg bodyweight. Subsequently, mice were exsanguinated by cardiac

puncture. Mice were then sacrificed by cervical dislocation and lungs were excised, snap frozen

and stored at -80°C for later analyses.

In order to induce bacteremia, GAS was diluted in PBS and 1×10^5 CFU were applied in a

100 µL volume by injection into the lateral tail vein. Intravenous injection of PBS served as a

control. For co-infection, IAV was applied as described above either two days prior or subse-

quent to bacterial infection. Mice were given tramadol (Ratiopharm) in drinking water for anal-

gesia. Animals were monitored following bacterial infection for a maximum of 14 days or until

humane endpoints were reached. Sepsis severity was assessed by a scoring system that incorpo-

rated the assessment of macroscopic signs of burden as previously described [35,36]. In brief,

scores of four categories were added together to provide an estimate for overall sepsis activity: i)

weight loss of $\geq$ 5% (Score 5), $\geq$ 10% (Score 10), $\geq$ 20% (Score 20, humane endpoint); ii) ap-

pearance deviations such as piloerection (Score 5), high myotonicity or scruffy orifices (Score

10), convulsions or paralysis (Score 20, humane endpoint); iii) impairment of consciousness such

as suppressed activity or limited reaction to stimuli (Score 5), self-isolation or lethargy (Score

10), perpetual pain vocalization or apathy (Score 20, humane endpoint) and iv) signs of impaired

respiratory quality or inflammation such as edemas on small body areas (Score 5), disseminated

edemas or labored breathing (Score 10), open wounds or gasping (Score 20, humane endpoint).

Mice were sacrificed as described above upon reaching the end of the observation period, at

any humane endpoint or when reaching an overall sepsis score of $\geq$ 20. Cardiac blood samples

were plated on blood agar and medial arthrotomy on both knee joints was performed under a ste-

reo microscope followed by plating of the synovial fluid on blood agar. Agar plates were subse-

quently incubated overnight and examined for the presence of β-hemolytic bacteria. Hind paws

were extracted, snap frozen and stored at -80°C for the analysis of eicosanoids.

2.4   Eicosanoid extraction and analysis

Lipidomics analyses were performed as previously described [35]. In brief, paw samples

were chilled in liquid nitrogen, pulverized and 50 mg of the resulting powder was immersed in

500 μL cold methanol containing 0.1% butylated hydroxytoluene and 500 μL ice cold water. 100

μL deuterated internal standards containing 12-HETE-d_8, 13-HODE-d_4, PGE_2-d_4 and Resolvin

D1-d_5 (each 100 ng/mL, Cayman Chemicals) were subsequently added followed by an additional

lysis step with matrix B at 6 m/s for 45 s on a FastPrep (MP Biomedicals). Following this, 300

μL sodium acetate (1 M) was added on ice and 10 M acetic acid was added until pH 6 was

reached. Solid phase extraction was performed on methanol and sodium acetate conditioned

Bond Elut Certify II cartridges (Agilent). After loading the samples, cartridges were washed with

50% methanol. Elution of eicosanoids was carried out by addition of hexane/ethyl acetate (75/25)

containing 1% acetic acid.

For measurements, paw extracts were dried under nitrogen flow using a TurboVap (Bio-

tage) and reconstituted in 70 µL 25% acetonitrile. Separation was done on a Gemini NX-C18

column (3 µm, 100 × 2 mm) utilizing an Agilent 1200 series HPLC systems. Dynamics multiple

reaction monitoring MS/MS was executed using a 6460 series triple quadrupole tandem mass

spectrometer (Agilent) with electrospray ionization in negative mode. Calibration by internal and

external standards was performed as previously described [35]. Agilent Mass Hunter Qualitative

Analysis software and Agilent Mass Hunter Quantitative Analysis software (both version

B.07.00) were used for MS data analysis. Quantities of individual eicosanoids were standardized

to a mean of 0 and a standard deviation of 1 for data visualization.

2.5 Isolation of RNA and DNA from lung samples

Lung samples were submerged in liquid nitrogen, slightly fragmented and weighed. Sixty

to one hundred twenty milligrams were transferred to lysis tubes containing bashing beads (Zymo

Research) and 1 mL TRIzol (Thermo Fisher). Lung fragments were subsequently homogenized at

4,000 rpm for 4×20 s using a FastPrep. Samples were then centrifuged at $10,000 \times g$ for 7 min at

4°C and transferred into new tubes. Apart from centrifugation at 4°C, the following steps were

conducted at room temperature. After resting for 5 min, 200 µL chloroform (Sigma) was added

and samples were extracted for 3 min. Subsequently, samples were centrifuged for 15 min at

$12,000 \times g$. The RNA-enriched upper phase was mixed with 500 µL 2-propanol, incubated for 10

min and centrifuged at $12,000 \times g$ for 10 min. RNA pellets were suspended in 75% Ethanol fol-

lowed by centrifugation at $7,500 \times g$ for 5 min. Supernatants were subsequently discarded, pellets

were dried and dissolved in 40 µL RNAse-free water by incubation at 60°C for 15 min. RNA

contents were then determined photometrically on a NanoDrop (Thermo Fisher). DNA was iso-

lated by precipitation of the appropriate phase upon addition of 300 µL ethanol, incubation for 3

min and centrifugation for 5 min at $2,000 \times g$. The resulting pellet was then washed twice by 30

min incubation with 0.1 M sodium citrate (pH 8.5) in 10% ethanol. DNA samples were subse-

quently suspended in 75% ethanol and incubated for 20 min. After centrifugation, supernatants

were discarded, pellets were dried and dissolved by incubation in 8 mM NaOH for 10 min. DNA

contents were determined fluorometrically using the Qubit 1X dsDNA Assay Kit to the manufac-

turer's instructions (Thermo Fisher).

2.6   Lung pathogen burden and gene expression

Primer pairs were designed for the detection of IAV H1N1 matrix protein, nucleoprotein

and hemagglutinin in murine lung extracts according to the strain specific sequences found at

https://www.fludb.org/brc/fluStrainDetails.spg?strainName=A%2FGermany-

BY%2F74%2F2009%28H1N1%29&decorator=influenza (supplementary Table I). For this,

RNA was isolated as described above and 500 ng were reverse transcribed using the High Capac-

ity cDNA Reverse Transcription Kit (Thermo Fisher) according to the manufacturer's instruc-

tions. Twenty-five nanograms of the resulting cDNA together with 500 nM of the primer pairs

were submitted to qPCR using the PowerUP SYBR Green Mastermix (Thermo Fisher). The am-

plification reaction was monitored on the ViiA 7 Real-Time PCR System running on the

QuantStudio Real Time PCR Software V1.3 (Thermo Fisher). The size of the respective ampli-

cons was confirmed by 2% agarose gel and ethidium bromide staining. Primer pairs for the detec-

tion of GAS strain AP1 specific genomic DNA were designed according to sequence information

found at https://www.ncbi.nlm.nih.gov/nuccore/CP007537?report=genbank (supplementary Ta-

ble II). A total of 20 ng DNA from lung extracts were used together with 500 nM of the primer

pairs for qPCR as described above, followed by confirmation of amplicon sizes on agarose gels.

Gene expression analyses were performed on 25 ng cDNA that was obtained from reverse tran-

scribed lung RNA. For qPCR analysis, TaqMan primer pairs and probes (Thremo Fisher) were

used for *Ccl2* (assay ID: Mm00441242_m1) and *Ifnb1* (Mm00439552_s1) utilizing *Gapdh*

(Mm05724508_g1) as a reference gene. All reactions were amplified using the TaqMan Gene

Expression Master Mix (Thermo Fisher).

2.7   Bone marrow derived macrophage infection model

C57BL/6J mice used for bone marrow isolation had a median age of 10 weeks (range 7 –

42 weeks) and 30% were female. Bone marrow was obtained from long bones by centrifugation

as previously described [37]. The resulting pellet was subsequently suspended in Dulbecco's

Modified Eagle's Medium (DMEM) supplemented with 10% fetal calf serum (FCS), 5 IE/mL

Penicillin, 5 µg/mL Streptomycin, 2 mM L-Glutamine (Thermo Fisher), 10 mM HEPES and 1

mM sodium pyruvate (PAN Biotech). After determination of vital cells using a hemocytometer

and trypan blue (Thermo Fisher), cells were seeded into 6-well culture plates (Greiner) at a densi-

ty of 3×10^5 cells per cm^2 in $2 - 5$ mL supplemented DMEM. The differentiation to macrophages

was initiated at day 0 by the addition of 20 ng/mL macrophage colony-stimulating factor (M-

CSF, R&D Systems). Cells were cultured afterwards at 37°C and 5% CO_2 for 7 days including

the replacement of supplemented DMEM and replenishment of M-CSF at days 1 and 4. For viral

infection (day 7, t_0), supplemented DMEM was refreshed and 4×10^5 $TCID_{50}$ IAV were added.

Following this, infected or uninfected macrophages were incubated for 48 h upon which the cells

were either collected for downstream analyses or submitted to bacterial (super-)infection (day 9).

In case of the latter, supplemented DMEM was removed, the cells were washed thrice with PBS

and Minimal Essential Medium α containing additional nucleosides and 10% FCS (Thermo Fish-

er) was added. GAS was then applied at 4.5×10^6 CFU. Subsequently, macrophages were incu-

bated for 6 h followed by sample collection.

2.8   Single cell analysis by flow cytometry

Gentle detachment of macrophages from culture plates was carried out by washing with

PBS and subsequently incubating with 5 mL PBS containing 10 mM EDTA for 10 min. Culture

plates were tapped multiple times and suspensions were collected afterwards. For increased

yields, 0.7 mL accutase (Pan Biotech) was added for 10 – 15 min followed by alternately tapping

and pipetting. Subsequently, another 0.7 mL accutase were added for an additional 10 – 15 min,

tapping and pipetting were repeated and suspensions collected and pooled with the PBS/EDTA

fraction. Finally, 1 mL supplemented DMEM was added and the remaining cells were obtained

using a cell scraper (Sarstedt). Suspensions were centrifuged at $400 \times g$ and 4°C for 5 min and

cells were suspended in autoMACS Running Buffer (RB, Miltenyi Biotec) followed by counting.

Antibody binding to CD16 and CD32 was prevented by incubation of macrophages with 0.5 µg

Trustain FcX (Biolegend) in RB supplemented with 10% FCS for 10 min on ice. Subsequently,

an antibody mixture containing 0.13 µg (anti-)F4/80:FITC (clone BM8), 0.5 µg CD163:APC

(S150491), 0.25 µg CD206:BV605 (C068C2), 0.25 µg CD80:BV421 (16-10A1, Biolegend), 0.22

µg CD86:APC/Vio770 (PO3.3), 4.5 µL TLR2:PE (REA109) and 0.15 µg MHCII:PerCP/Vio770

(REA813, Miltenyi Biotec) was added and incubated for 20 min on ice in the dark. Cells were

washed afterwards, suspended in RB and 7-Aminoactinomycin (7-AAD, Biolegend) was added at

a concentration of 1.25 µg/mL for at least 5 min prior to measurement.

Data acquisition was performed on the Aurora spectral flow cytometer running on the

SpectroFlo software v2.2.0.3 (Cytek Biosciences). Data analysis was conducted using the FlowJo

software v10.7.1. Supplementary Fig. 1 illustrates the gating strategy. Live macrophages were

identified as 7-AAD⁻F4/80⁺ singlets. This population was used for the subsequent determination

of expression levels based on median fluorescence intensity (MFI) values and as a parent for

measuring the proportions of subpopulations expressing different combination of the above-listed

surface antigens. For dimension reduction, 10,000 macrophage events were down-sampled, con-

catenated and submitted to the algorithm t-distributed stochastic neighbor embedding (t-SNE)

using an automated learning configuration (opt-SNE combined with the exact KNN algorithm

and the Barnes-Hut gradient algorithm) with a perplexity of 50 and a maximum of 1000 iterations

[38]. Unsupervised clustering of subpopulations expressing any combinations of the analyzed

surface proteins was conducted by FlowSOM [39].

2.9   Macrophage gene expression

After aspirating cell culture supernatants, 700 µL of a chaotropic agent solution (Qiagen)

was added to individual wells and cells were lysed by scraping and vigorous shaking. RNA was

subsequently isolated using the RNeasy Plus Mini Kit (Qiagen) after the manufacturer's instruc-

tions. Quantification of RNA contents were determined photometrically and 200 ng RNA was

submitted to reverse transcription as described above. Amplification of cDNA was then per-

formed by TaqMan Gene Expression Master Mix, primer pairs and probes for the relative quanti-

fication of *Ccl2*, *Cxcl2* (assay ID: Mm00436450_m1), *Ifnb1*, *Il1b* (Mm00434228_m1), *Il6*

(Mm00446190_m1), *Il10* (Mm00439614_m1), *Mgl2* (Mm00460844_m1), *Nos2*

(Mm00440502_m1), *Tgfb1* (Mm01178820_m1) and *Tnf* (Mm00443258_m1) using *Gapdh* as a

reference gene. Polymerase chain reactions were performed on the Viia 7 System. In detail, sam-

ples were first incubated for 2 min at 50°C followed by 10 min at 95°C for polymerase activation.

Subsequently, 40 automated cycles of PCR were performed that incorporated denaturation at

95°C for 15 sec and annealing and elongation at 60°C for 1 min. After each cycle the fluorescein

amidite fluorescence signal was measured. Ct values were obtained when fluorescence intensities

reached data-dependent and automatically defined thresholds. Quantification of gene expression

was then performed by the $2^{-\Delta\Delta Ct}$ Method that incorporated normalization of the target gene Ct

values to the reference gene (ΔCt) as well the difference between ΔCt values from uninfected and

infected cells ($\Delta\Delta Ct$).

2.10 Cytokine analysis

Cytokine concentration in mouse plasma samples were quantified by a 3-plex LEGEND-

plex assay (Biolegend) that contained capture beads and detection antibodies for CCL2 (mono-

cyte chemoattractant protein-1, MCP1), Interferon (IFN)γ and tumor necrosis factor (TNF)α. For

the quantification of CCL2, Interleukin (IL-)1β, IL-6, IL-10 and TNFα in cell culture superna-

tants, a 5-plex LEGENDplex assay was used following the manufacturer's guidelines. Data ac-

quisition was performed on the Cytek Aurora flow cytometer. Cell culture supernatant concentra-

tions of CXCL2 (macrophage inflammatory protein 2-α, MIP2-α) were determined by the

CXCL2/MIP-2 DuoSet enzyme linked immunosorbent assay (ELISA) kit to the manufacturer's

instructions (R&D Systems). Horseradish peroxidase-catalyzed color reaction were initiated by

addition of the TMB Substrate Kit (Biolegend) and quenched by 0.5 M sulfuric acid (Merck).

The absorbance at 450 nm was measured on the Infinite M200 spectral photometer (Tecan).

2.11 Statistical analysis

Data analysis and visualization were performed using RStudio v1.2.5033 that ran R v3.5.1.

Normalization was either performed by division of individual values from infection groups and

control groups, respectively, or by feature scaling into a 0 – 1 range by the formula $x_i' = (x_i -$

$x_{min})/(x_{max} - x_{min})$. Heatmaps and hierarchical clusters were generated with the "pheatmap" pack-

age that incorporated feature scaling by standardization applying the formula $z_i = (x_i - \bar{x})/\sigma$. Cali-

bration curves were fitted and samples values were estimated by n-parameter logistic regression

using the "nplr" package. Two-sided statistical tests were used for the comparisons of group me-

dians or means. Repeated measures (body weight trajectories) were compared by one-way and

two-way analyses of variance (ANOVA), respectively. Probabilities of survival and incidences

were compared by the logrank test. Bivariate interdependencies were evaluated by the Pearson

product-moment correlation coefficient (r). Data sets were tested for normality by the Shapiro-

Wilk test. Normal distribution of within-group raw or normalized variables was rejected when the

test resulted in a p-value of < 0.05. Depending on the outcome of this test, univariate statistical

analyses on variables that were normalized to respective controls were performed by the one-

sample Wilcoxon signed rank test and the one-sample t-test, respectively. Two independent sam-

ples were compared with the Mann-Whitney U test or the t-test. Comparisons of variables be-

tween multiple groups were performed with the Dunn's test and the Tukey HSD test, respective-

ly, in combination with type I error correction using the Bonferroni-Holm method. A p-value of $<$

0.05 was considered statistically significant.

3   Results

3.1   Infection with influenza A virus H1N1 caused mild symptoms and induced a persistent immune reaction in the lung

In order to examine clinical manifestations of influenza, we used a model of intranasal infection with 2009 pandemic H1N1 IAV in adult mice (Fig. 1A). Intranasal application of PBS served as a control. Mice were monitored for relative weight loss post infection as a proxy for disease severity and indeed, exhibited minor reductions in body weight as early as two days after virus application (Fig. 1B). This trend continued until day seven after infection and resulted in a maximum weight loss of 5.5% ± 2.1% (mean ± SEM). Thereafter, body weight continuously increased and returned to starting values by day 14 suggesting robust recovery from infection. When comparing weight trajectories over the entire observation period using two-way analysis of variance (ANOVA), we found a statistically significant difference between infected mice and uninfected controls ($p < 0.001$). In accordance with the observed mild disease courses, we did not measure quantifiable amounts of the inflammatory cytokines TNFα and IFNγ in plasma samples from IAV infected animals (not shown). We did, however detect significant reductions of plasma CCL2 concentrations by 12.8% and 13.6% at days two and four after infection, respectively, relative to uninfected controls (Fig. 1C). By day seven, CCL2 plasma levels equalized between both groups ($p = 0.65$, not shown).

In order to examine immune responses in the lower respiratory tract, we further performed gene expression analyses on whole lung homogenates that were obtained 16 days after IAV application. For this, we focused on mRNA expression levels of *Ccl2* and *Ifnb1*, as the former was altered in the periphery and the latter can be indicative of an anti-viral response. Protein data were not collected because of limited sample quantities. We found no meaningful differences in the expression of *Ccl2* between the IAV and control groups (Fig. 1D, left panel). Interestingly,

Ifnb1 expression was found to be significantly increased in lungs from infected mice (Fig. 1D,

right panel). Given this prolonged upregulation of *Ifnb1*, we consequently utilized primer pairs

for the detection of viral genes in lung samples that code for hemagglutinin, matrix protein and

nucleoprotein (supplementary Fig. 2). We indeed detected IAV-specific RNA in 38% (3/8) of

infected animals by quantitative PCR (Fig. 1E). False positive detection of unspecific targets was

ruled out by confirming the expected amplicon melting temperatures (supplementary Fig. 3).

However, the quantities of all three viral genes were generally low ($C_t > 32$) and might rather

indicate residual viral antigen. Interestingly, two of the three samples that were positive for viral

genes were also among the samples that expressed the highest amounts of *Infb1* indicating some

correlation between both parameters.

In summary, we here show that an infection with IAV H1N1 in mice induced minor clinical

manifestations that were accompanied by an early and reversible reduction of plasma CCL2 lev-

els. Our data further show that barely detectable genetic material of the virus persisted in the

lungs of some animals, which was accompanied by an ongoing type I IFN production.

3.2   Group A Streptococcal infection was aggravated following Influenza A virus infection

As CCL2 is integral to bacterial control [40–42], yet reduced during respiratory tract infec-

tion with IAV, we sought to investigate the clinical features of IAV superimposed bacteremia. To

this end, we compared infection with bacteria only to co-infection models combining intranasal

virus application with intravenous GAS infection in alternating succession (Fig. 2A). By monitor-

ing for macroscopic symptoms following bacterial infection, we observed the occurrence of lo-

calized paw inflammation (Fig. 2B). Of note, the emergence of these edemas was accelerated and

more frequent in post-influenza bacteremia (IAV+GAS) compared to bacterial infection only

(GAS, p = 0.01) and pre-influenza bacteremia (GAS+IAV, p = 0.045), respectively (Fig. 2C). In

detail, 80% (8/10) of mice in the IAV+GAS group exhibited signs of paw inflammation already

one day after bacterial infection. In contrast, the incidence of paw edemas was increased to only

40% (4/10) in the GAS+IAV group as opposed to 20% (2/10) in the GAS only group, and this

difference did not reach statistical significance (p = 0.33). We further analyzed eicosanoids from

paw extracts and found that these immunologically active lipid metabolites were upregulated in

some animals irrespective of the (co-)infection regimen (supplementary Fig. 4).

Additionally, more blood smears and knee joint capsule swabs were positive for β-

hemolytic bacteria in co-infected mice from the IAV+GAS group (Table I), which suggested that

preceding influenza promoted bacterial dissemination and invasion into synovial tissues. By as-

sessing macroscopic signs of burden as a proxy for sepsis severity (see Materials and Methods),

we found a significantly increased median disease score when comparing post-influenza bacte-

remia with monocausal GAS infection (Fig. 2D). Interestingly, when correlating sepsis scores

with eicosanoids from paw homogenates, we found a significant relationship between the indi-

vidual disease severity and the corresponding amounts of prostaglandins D_2 and E_2 as well as 5-

and 12-Hydroxyeicosatetraenoic acid (supplementary Table III). Furthermore, elevated disease

severity in the IAV+GAS group was paralleled by a reduction in survival probability to 40%

compared to 80% in the GAS only group (Fig. 2E). In contrast, mice from the GAS+IAV group

had an only marginally decreased survival chance of 70%. However, the overall probability for a

fatal outcome was, according to logrank statistics, not significantly different between groups (p =

0.13) and this was likely due to low sample sizes and a high degree of uncertainty.

For our further analyses, we focused on the IAV+GAS co-infection sequence because our

data suggested that the clinical outcome was not different between the GAS+IAV and GAS

groups. We next aimed to investigate whether post-influenza GAS infection impacted on the im-

mune activation in the lower respiratory tract. To this extent, we analyzed lung homogenates for

the expression of *Ccl2* and *Ifnb1*, and compared the data from co-infected mice to GAS mono-

infection or uninfected controls. We found that neither GAS nor IAV+GAS infection resulted in

a meaningful alteration of the *Ccl2* expression in the lung (Fig. 2F, left panel). Of note, lungs

from both mono- and co-infected mice exhibited a median 2-fold upregulation of *Ifnb1* relative to

lungs from uninfected animals (p = 0.008 for GAS and p = 0.039 for IAV+GAS; Fig. 2F, right

panel). Yet, when comparing the infection regimens with each other, we found that *Ifnb1* overex-

pression was comparable between both infection groups (p = 0.93).We were curious whether the

bacteria are capable of disseminating from the blood into the lower respiratory tract and therefore

analyzed lung homogenates for the presence of GAS specific genes using quantitative PCR (sup-

plementary Fig. 5). Indeed, we detected genomic speB in four out of nine lungs from the

IAV+GAS group whereas only one out of nine lungs from the GAS group was positive for this

bacterial gene (Fig. 2G, left panel). However, when analyzing for spy2158, only two lung ex-

tracts from the IAV+GAS group were positive (Fig. 2G, right panel). Specific amplification was

again confirmed by melting curves (supplementary Fig. 6). As whole lungs were submitted to

chaotropic agent assisted homogenization and PCR analysis, we were not able to confirm wheth-

er there were any vital bacteria present in these samples.

Collectively, our *in vivo* data demonstrated that a preceding IAV infection of the respiratory

tract aggravated intravenous GAS infection by promoting localized inflammation and a dysregu-

lated host response as shown by an elevated sepsis scores. In contrast, application of the virus

following an already established bacteremia did not influence on disease progression and out-

come.

3.3 *Preceding influenza A virus infection impacted on the Group A Streptococcus induced di-*

versification of macrophage surface expression profiles

As our *in vivo* co-infection model implicated a preceding IAV infection to cause impaired

control of the bacterial challenge following a superimposed GAS infection, we sought to explore

any modification of anti-bacterial innate immunity. Macrophages are considered as first line de-

fense immune cells that are substantial to contain pathogens in early phases of infection [43].

These cells take part in numerous bacterial infectious diseases and are especially crucial for a

competent innate immune response during invasive GAS infection [31,44–46]. Hence, we chose

in vitro (co-)infection models of primary murine macrophages in order to investigate whether

IAV influences on the GAS-induced immune landscape. In detail, murine macrophages were

differentiated from bone marrow cells by M-CSF stimulation and were subsequently infected

with IAV, GAS or IAV and GAS (Fig. 3A). We then analyzed the expression patterns of immu-

nologically relevant surface antigens by flow cytometry. In order to gain insight into differential-

ly expressed macrophage markers, we performed dimension reductions on our multiparametric

data sets by t-distributed stochastic neighbor embedding (t-SNE). Fig. 3B demonstrates for the

topological distribution of surface marker expression levels distinct allocations of cells that were

obtained from the different infection models. For instance, macrophage subsets overexpressing

CD80 and CD86 were seemingly enriched in IAV+GAS co-infected cultures, whereas mono-

infection with GAS resulted in the accumulation of CD206 overexpressing macrophages. Unsu-

pervised clustering of macrophage populations on the basis of their respective expression patterns

by flowSOM further indicated that co-infection triggered a different response than viral or bacte-

rial mono-infections (supplementary Fig. 7).

In an effort to obtain a more detailed picture of IAV- and GAS-induced immune responses,

we next focused on the individual expressions of macrophage surface antigens. Given the inter-

experimental variance of macrophage cultures, median fluorescence intensities (MFI) of (co)-

infected cells were normalized to their corresponding uninfected controls that were acquired from

the same donor animal (supplementary Fig. 8). Notably, expression patterns were similar within

each group, which resulted in a robust hierarchical clustering for IAV, GAS and IAV+GAS in-

fected macrophages (Fig. 3C). In detail, apart from a significant upregulation of CD163 com-

pared to both bacterial infection and co-infection, IAV had hardly any impact on the expression

of the investigated surface proteins (Fig. 3C, 3D). Conversely, GAS infection induced the over-

expression of TLR2, which was even amplified following co-infection (Fig. 3D). Both the appli-

cations of GAS and IAV+GAS comparably prompted an elevated production of MHCII. Alt-

hough not statistically significant, GAS infection led to a slight downregulation of CD80, which

was reversed to an upregulated expression in the IAV+GAS group. Similarly, co-infection trig-

gered a minor overexpression of CD86 that was short of reaching statistical significance due to a

high within-group variance ($p = 0.067$, compared to uninfected). The downregulation of CD163

as well as the attenuation of the GAS-induced CD206 upregulation in the IAV+GAS group fur-

ther supports the notion that a preceding IAV infection led to a distinct immune response in mac-

rophages during co-infection (Fig. 3D).

As a result of differentially affected expression landscapes, the proportions of distinctive

macrophage subpopulations shifted depending on the infection regimen (Fig. 3E). We found a

minor depletion of CD80$^+$CD86$^+$ cells following GAS infection ($p = 0.1$), whereas co-infection

caused a significantly increased proportion of this population when compared to uninfected con-

trols (Fig. 3F). Both bacterial mono-infection and co-infection induced an enrichment of MHCII$^+$

macrophages, suggesting a retained ability of these immune cells to inform and coordinate an

adaptive immune response. In accordance with the altered expression profiles shown in Fig. 3D,

the proportions of CD163$^+$ and CD206$^+$ cells, respectively, were decreased upon co-infection

relative to GAS infection only (Fig. 3F).

Collectively, our data on the diversification of surface antigen expression demonstrated that

the immune response of macrophages towards co-infection with IAV and GAS was considerably

distinct from the effects that were induced by either mono-infection. Although we encountered

some similarities between the GAS and IAV+GAS groups, the preceding viral infection seeming-

ly manipulated or obliterated the macrophages´ reaction towards the bacterial pathogen.

3.4   A preceding influenza A virus infection impaired the inflammatory capacity of macrophag-

es during co-infection

In order to gain further insights into the immune response triggered by infected macrophag-

es, we next performed analyses on mRNA expression of immune mediators by quantitative PCR

and measured cytokine secretion by ELISA or bead-based multiplex analysis (Fig. 4A). As illus-

trated in Fig. 4B, the different (co-)infection regimens triggered distinct expression patterns of

immunomodulatory agents that resulted in strong within-group associations as shown via hierar-

chical clustering. IAV infection was specifically characterized by a relatively higher expression

of *Mgl2* and *Tgfb1* (Fig 4C, supplementary Fig. 9). Bacterial infection, on the other hand, com-

prehensively stimulated the overexpression of several genes that mediate an inflammatory re-

sponse (Fig. 4B). Strikingly, co-infected macrophages mostly failed to induce a similar magni-

tude of GAS inducible overexpression, yet upregulated *Arg1* (Fig. 4B, 4C).

By further examining individual expressions, we found that *Mgl2* was significantly reduced

following bacterial mono-infection and co-infection by 1.8- and 3.9-fold, respectively, compared

to uninfected controls (Fig. 4C). Remarkably, GAS application induced an approximately 3,000

fold overexpression of *Nos2* that was impeded during co-infection to a mere, yet statistically sig-

nificant, 10-fold overexpression. Furthermore, co-infection triggered the upregulation of *Ccl2*,

Cxcl2 and *Tnf*, which were significantly less pronounced in comparison to GAS infection only

(Fig. 4D). Secretion of these cytokines was mostly comparable between these groups, however

TNFα production by co-infected macrophages was reduced (Fig. 4E). Although both *Il6* and *Il10*

were increased in the GAS and IAV+GAS group, respectively, only co-infection caused a signifi-

cant secretion of the protein products (supplementary Fig. 9). While *Ifnb1* was only upregulated

following GAS infection, *Tgfb1* was downregulated after GAS infection as well as co-infection

(supplementary Fig. 9A). Of note, although the GAS-induced overexpression of *Il1b* was also

observed in the IAV+GAS group, co-infection entirely abrogated the secretion of mature IL-1β,

which suggests that a preceding IAV infection compromised innate immune sensing of Strepto-

cocci (Fig. 4D, 4E).

In summary, the expression patterns of immunologically active mediators were noticeably

different between GAS mono-infection and IAV+GAS co-infection, implying that prior virus

infection modifies anti-streptococcal immunity.

4   Discussion

In this study we demonstrated that influenza promoted subsequent intravenous GAS infection and allowed for dissemination of the bacterial pathogen within the blood and its migration into lungs as well as synovial tissues. Although we did not assess any alterations in bone or cartilage morphology, we would like to argue that an invasion of articular tissue by GAS is reminiscent of septic arthritis [47]. Indeed, we previously demonstrated that the occurrence of paw edemas, which in the present study was more likely during IAV and GAS co-infection, was due to bacterial colonization of both, subcutaneous and periarticular tissues and was paralleled by immune cell infiltration [35]. Hence, we here show for the first time, that a preceding IAV infection predisposes the host to severe complications during GAS blood infection. Conversely, IAV infection elicited subsequent to intravenous GAS infection did not aggravate disease severity, suggesting that immune priming events in response to a prior viral encounter mitigate an otherwise competent anti-bacterial immune response.

Influenza in humans is usually characterized by mild-to-moderate disease that is rarely lethal and resolves shortly after infection [48], which was also shown in our animal model of IAV inoculation. Upon entry into nasopharyngeal cavities, the virus trespasses into the mucus, invades the epithelium and spreads to immune cells [49,50]. The host then recognizes parts of the viral RNA genome by intracellular pattern recognitions receptors, which triggers the production of several inflammatory cytokines, among them type I IFNs, that establish an anti-viral immune state [51–53]. We have demonstrated that residual viral genes persisted for 16 days in the lungs of some infected mice, which was paralleled by a continuous upregulation of *Ifnb1*. However, we believe it to be unlikely that replicative viral particles were still present in the lungs up to this point because IAV is typically cleared within a couple days following infection and the quantities of viral genes were barely detectable in our samples [54–56]. Type I IFN can have beneficial ef-

fects during bacterial infection by promoting host resilience and by preventing systemic hyperin-

flammation [57–60]. However, several studies advocated that the consequences of type I IFN

expression are detrimental for the containment of a secondary bacterial insult subsequent to influ-

enza [20,61,62].

By using a mouse strain that lacks the common IFNα/β receptor (IFNAR) in a model of

pneumococcal superinfection, Shahangian and colleagues demonstrated that the IAV-induced

IFNAR signaling led to an impaired production of the neutrophil attractants CXCL1 and CXCL2

[22]. They argued that, in agreement with a complementary study by Didierlaurent *et al*., type I

IFNs desensitize subsequent TLR-mediated recognition of bacterial components by macrophages,

which are major producers for these chemokines [22,23]. Another work on IFNAR[-/-] mice by

Nakamura and colleagues had some contrasting results concerning the impact of type I IFN sig-

naling on pneumococcal superinfection [24]. In their study, they found that the virus and the bac-

teria were capable of synergistically inducing an overproduction of type I IFNs, which led to an

impaired production of CCL2 while CXCL1/2 production was unaltered [24,63]. CCL2 supports

bacterial clearance by the attraction of CCR2[+] monocytes to the infected tissue [64,65]. Along

these lines, we found in our study that CCL2 was significantly reduced in the plasma of IAV-

infected mice and that both, monocausal bacterial infection and co-infection featured *Ifnb1* over-

expression in the lung. Hence, although the role of CCL2 during GAS infections is not yet fully

elucidated, we find it possible that a preceding influenza restricts anti-bacterial immunity by lim-

iting monocyte homing and their differentiation to macrophages not only in pulmonary tissues

but also in remote host compartments that would be affected during GAS blood infection. How-

ever, the cellular source of this chemokine was not identified in our study. Furthermore, the ex-

pression of *Ccl2* in lung samples that were taken at endpoints was comparable between bacterial

infection and co-infection, which challenges the idea that IAV-induced alterations in immune cell

recruitment continues after bacterial superinfection. In order to delineate the progression of co-

infection in more detail, future studies should therefore focus on observations that are performed

at specific time points rather than taking samples at endpoints that might be difficult to compare.

Apart from the ramifications due to an impaired chemokinogenesis, we suspected other

means by which IAV dampens innate immune sensing of GAS. We hence focused on macro-

phage immunobiology in the context of co-infection and found that the virus comprehensively

altered GAS-induced gene expression patterns and cytokine layout. In detail, we detected that the

immune sensors CD163 and CD206 were markedly downregulated in co-infected compared to

GAS only infected macrophages. CD163 is an acute phase-regulated scavenger receptor that is

exclusively expressed by cells of the monocyte lineage and aids in the removal of potentially tox-

ic iron complexes during intravascular hemolysis [66–69]. Due to the fact that CD163 also medi-

ates tissue repair [70], host resilience [66,68], immune resolution and is able to sense gram-

positive bacteria [71,72], we speculate that this receptor might confer a protective immune state

during hemolytic bacteremia, even though its role in GAS infection is yet underexplored. Similar-

ly, the mannose receptor CD206 might support pathogen sensing during co-infection [73–77],

however mice that lack this sensor molecule are not more susceptible to infection [78,79].

Strikingly, a preceding IAV inoculation notably reduced the GAS-induced upregulation of

Nos2 while boosting *Arg1* expression. Both genes code for enzymes that compete for the sub-

strate L-Arginine, yet induce opposed immune mechanisms [80–82]. While nitric oxide synthase

2 (NOS2) provides inflammatory and bactericidal metabolites [83–85], arginase (ARG1) supports

tissue repair and immune resolution [83]. Thus, our data hint at a distortion of anti-bacterial pro-

cesses due to a prior IAV infection. This is further corroborated by an inadequate sensing of the

bacterial pathogen indicated by the reduced and abolished production of TNFα and IL-1β, respec-

tively, which was similarly shown in a model of pneumococcal superinfection [19]. Interestingly,

we detected for both, GAS mono-infection and superinfection an upregulation of *Il1b*, which

suggests that the incapacity of co-infected macrophages to process and secrete IL-1β is due to a

failure in the GAS-inducible activation of the NLRP3 inflammasome [86–89]. In fact, it was

shown that different variants of IAV, including a 2009 pandemic strain, were capable of thwart-

ing IL-1β maturation by interfering with NLRP3 inflammasome assembly [90–92], which is cru-

cial for innate immune sensing and coordination [93]. An IAV-mediated nullification of IL-1β

secretion would be of dramatic consequences during streptococcal superinfections. The absence

of signaling via the IL-1 receptor (IL-1R) was in fact associated with an increased susceptibility

to systemic GAS infection in both mice and humans [86,94,95]. Remarkably, rheumatoid arthritis

patients that received the IL-1R antagonist Anakinra exhibited a roughly 330-fold increased rate

of invasive GAS infections which included an elevated likelihood of life-threatening complica-

tions such as necrotizing fasciitis and sepsis [95].

Although our study yielded several findings that are of interest to the research about influ-

enza and bacterial co-infections, it has some limitations. Apart from the fact that our observations

remained purely phenomenological and no experiments on underlying mechanisms were per-

formed, we were unable to link the gap between our *in vivo* and *in vitro* models. For instance, the

IAV-induced reduction of CCL2 in mice was not observed in macrophages that were challenged

with the virus. Furthermore, it would have been interesting to test whether IAV impedes the ca-

pacity of macrophages to phagocytize GAS. Nevertheless, the multitude of IAV-inducible altera-

tions to macrophage immunobiology in the context of GAS superinfection was surprising to us

and warrant further investigations.

In summary, we here describe in complementary *in vivo* and *in vitro* co-infection models

that IAV infection thwarts anti-streptococcal innate immunity. This finding warrants further in-

vestigations on the mechanisms underlying this phenomenon that sets the stage for post-influenza

superinfection. As an important side issue, our work underscores the importance of regular vac-

cinations against influenza in order to avert bacterial superinfection and prevent fatal invasive

GAS complications [10,96–100].

References

References

1. Siemens, N.; Oehmcke-Hecht, S.; Mettenleiter, T.C.; Kreikemeyer, B.; Valentin-Weigand, P.;

Hammerschmidt, S. Port d'Entrée for Respiratory Infections – Does the Influenza A Virus Pave the Way for

Bacteria? *Front. Microbiol.* **2017**, *8*, 2602, doi:10.3389/fmicb.2017.02602.

2. Tjon-Kon-Fat, R.; Meerhoff, T.; Nikisins, S.; Pires, J.; Pereyaslov, D.; Gross, D.; Brown, C.; Drishti, A.;

Hasibra, I.; Kota, M.; et al. The Potential Risks and Impact of the Start of the 2015–2016 Influenza Season in

the WHO European Region: A Rapid Risk Assessment. *Influenza Other Respi. Viruses* **2016**, *10*, 236–246,

doi:10.1111/irv.12381.

3. McHardy, A.C.; Adams, B. The Role of Genomics in Tracking the Evolution of Influenza A Virus. *PLoS*

Pathog. **2009**, *5*, e1000566, doi:10.1371/journal.ppat.1000566.

4. Neumann, G.; Noda, T.; Kawaoka, Y. Emergence and Pandemic Potential of Swine-Origin H1N1 Influenza

Virus. *Nature* **2009**, *459*, 931–939, doi:10.1038/nature08157.

5. Taubenberger, J.K.; Morens, D.M. 1918 Influenza: The Mother of All Pandemics. *Emerg. Infect. Dis.* **2006**,

12, 15–22, doi:10.3201/eid1201.050979.

6. Morens, D.M.; Taubenberger, J.K.; Fauci, A.S. Predominant Role of Bacterial Pneumonia as a Cause of

Death in Pandemic Influenza: Implications for Pandemic Influenza Preparedness. *J. Infect. Dis.* **2008**, *198*,

962–970, doi:10.1086/591708.

7. Brundage, J.F.; Shanks, G.D. Deaths from Bacterial Pneumonia during 1918–19 Influenza Pandemic. *Emerg.*

Infect. Dis. **2008**, *14*, 1193–1199, doi:10.3201/eid1408.071313.

8. Tasher, D.; Stein, M.; Simões, E.A.F.; Shohat, T.; Bromberg, M.; Somekh, E. Invasive Bacterial Infections

in Relation to Influenza Outbreaks, 2006-2010. *Clin. Infect. Dis.* **2011**, *53*, 1199–1207,

doi:10.1093/cid/cir726.

9. McKenna, S.; Malito, E.; Rouse, S.L.; Abate, F.; Bensi, G.; Chiarot, E.; Micoli, F.; Mancini, F.; Gomes

Moriel, D.; Grandi, G.; et al. Structure, Dynamics and Immunogenicity of a Catalytically Inactive CXC

Chemokine-Degrading Protease SpyCEP from Streptococcus Pyogenes. *Comput. Struct. Biotechnol. J.* **2020**,

18, 650–660, doi:10.1016/j.csbj.2020.03.004.

10. Chaussee, M.S.; Sandbulte, H.R.; Schuneman, M.J.; DePaula, F.P.; Addengast, L.A.; Schlenker, E.H.;

Huber, V.C. Inactivated and Live, Attenuated Influenza Vaccines Protect Mice against

Influenza:Streptococcus Pyogenes Super-Infections. *Vaccine* **2011**, *29*, 3773–3781,

doi:10.1016/j.vaccine.2011.03.031.

11. Teymournejad, O.; Montgomery, C.P. Evasion of Immunological Memory by S. Aureus Infection:

Implications for Vaccine Design. *Front. Immunol.* **2021**, *12*, doi:10.3389/fimmu.2021.633672.

12. Okamoto, S.; Kawabata, S.; Nakagawa, I.; Okuno, Y.; Goto, T.; Sano, K.; Hamada, S. Influenza A Virus-

Infected Hosts Boost an Invasive Type of Streptococcus Pyogenes Infection in Mice. *J. Virol.* **2003**, *77*,

4104–4112, doi:10.1128/JVI.77.7.4104-4112.2003.

13. Okamoto, S.; Kawabata, S.; Terao, Y.; Fujitaka, H.; Okuno, Y.; Hamada, S. The Streptococcus Pyogenes

Capsule Is Required for Adhesion of Bacteria to Virus-Infected Alveolar Epithelial Cells and Lethal

Bacterial-Viral Superinfection. *Infect. Immun.* **2004**, *72*, 6068–6075, doi:10.1128/IAI.72.10.6068-6075.2004.

14. van der Sluijs, K.F.; Nijhuis, M.; Levels, J.H.M.; Florquin, S.; Mellor, A.L.; Jansen, H.M.; van der Poll, T.;

Lutter, R. Influenza-Induced Expression of Indoleamine 2,3-Dioxygenase Enhances Interleukin-10

Production and Bacterial Outgrowth during Secondary Pneumococcal Pneumonia. *J. Infect. Dis.* **2006**, *193*,

214–222, doi:10.1086/498911.

15. Wang, B.; Li, S.; Southern, P.J.; Cleary, P.P. Streptococcal Modulation of Cellular Invasion via TGF-B1

Signaling. *Proc. Natl. Acad. Sci.* **2006**, *103*, 2380–2385, doi:10.1073/pnas.0506668103.

16. Herrera, A.L.; Suso, K.; Allison, S.; Simon, A.; Schlenker, E.; Huber, V.C.; Chaussee, M.S. Binding Host

Proteins to the M Protein Contributes to the Mortality Associated with Influenza–Streptococcus Pyogenes

Superinfections. *Microbiol. (United Kingdom)* **2017**, *163*, 1445–1456, doi:10.1099/mic.0.000532.

17. Korteweg, C.; Gu, J. Pathology, Molecular Biology, and Pathogenesis of Avian Influenza A (H5N1)

Infection in Humans. *Am. J. Pathol.* **2008**, *172*, 1155–1170, doi:10.2353/ajpath.2008.070791.

18. Plotkowski, M.C.; Bajolet-Laudinat, O.; Puchelle, E. Cellular and Molecular Mechanisms of Bacterial

Adhesion to Respiratory Mucosa. *Eur. Respir. J.* **1993**, *6*, 903–916.

19. Sun, K.; Metzger, D.W. Inhibition of Pulmonary Antibacterial Defense by Interferon-γ during Recovery

from Influenza Infection. *Nat. Med.* **2008**, *14*, 558–564, doi:10.1038/nm1765.

20. Navarini, A.A.; Recher, M.; Lang, K.S.; Georgiev, P.; Meury, S.; Bergthaler, A.; Flatz, L.; Bille, J.;

Landmann, R.; Odermatt, B.; et al. Increased Susceptibility to Bacterial Superinfection as a Consequence of

Innate Antiviral Responses. *Proc. Natl. Acad. Sci.* **2006**, *103*, 15535–15539, doi:10.1073/pnas.0607325103.

21. Metzger, D.W.; Sun, K. Immune Dysfunction and Bacterial Coinfections Following Influenza. *J. Immunol.*

2013, *191*, 2047–2052, doi:10.4049/jimmunol.1301152.

22.   Shahangian, A.; Chow, E.K.; Tian, X.; Kang, J.R.; Ghaffari, A.; Liu, S.Y.; Belperio, J.A.; Cheng, G.; Deng,

J.C. Type I IFNs Mediate Development of Postinfluenza Bacterial Pneumonia in Mice. *J. Clin. Invest.* **2009**,

119, 1910–1920, doi:10.1172/JCI35412.

23.   Didierlaurent, A.; Goulding, J.; Patel, S.; Snelgrove, R.; Low, L.; Bebien, M.; Lawrence, T.; Van Rijt, L.S.;

Lambrecht, B.N.; Sirard, J.C.; et al. Sustained Desensitization to Bacterial Toll-like Receptor Ligands after

Resolution of Respiratory Influenza Infection. *J. Exp. Med.* **2008**, *205*, 323–329, doi:10.1084/jem.20070891.

24.   Nakamura, S.; Davis, K.M.; Weiser, J.N. Synergistic Stimulation of Type I Interferons during Influenza

Virus Coinfection Promotes Streptococcus Pneumoniae Colonization in Mice. *J. Clin. Invest.* **2011**, *121*,

3657–3665, doi:10.1172/JCI57762.

25.   Cunningham, M.W. Pathogenesis of Group A Streptococcal Infections. *Clin. Microbiol. Rev.* **2000**, *13*, 470–

511, doi:10.1128/CMR.13.3.470-511.2000.

26.   Musher, D. Trends in Bacteremic Infection Due to Streptococcus Pyogenes (Group A Streptococcus), 1986-

1995. *Emerg. Infect. Dis.* **1996**, *2*, 54–56, doi:10.3201/eid0201.960107.

27.   Herrera, A.L.; Huber, V.C.; Chaussee, M.S. The Association between Invasive Group A Streptococcal

Diseases and Viral Respiratory Tract Infections. *Front. Microbiol.* **2016**, *7*, 1–7,

doi:10.3389/fmicb.2016.00342.

28.   Jean, C.; Louie, J.K.; Glaser, C.A.; Harriman, K.; Hacker, J.K.; Aranki, F.; Bancroft, E.; Farley, S.;

Ginsberg, M.; Hernandez, L.B.; et al. Invasive Group a Streptococcal Infection Concurrent with 2009 H1n1

Influenza. *Clin. Infect. Dis.* **2010**, *50*, 59–62, doi:10.1086/652291.

29.   Zakikhany, K.; Degail, M.A.; Lamagni, T.; Waight, P.; Guy, R.; Zhao, H.; Efstratiou, A.; Pebody, R.;

George, R.; Ramsay, M. Increase in Invasive Streptococcus Pyogenes and Streptococcus Pneumoniae

Infections in England, December 2010 to January 2011. *Eurosurveillance* **2011**, *16*, 1–4,

doi:10.2807/ese.16.05.19785-en.

30.   Okamoto, S.; Nagase, S. Pathogenic Mechanisms of Invasive Group A Streptococcus Infections by Influenza

Virus–Group A Streptococcus Superinfection. *Microbiol. Immunol.* **2018**, *62*, 141–149, doi:10.1111/1348-

0421.12577.

31.   Goldmann, O.; Rohde, M.; Chhatwal, G.S.; Medina, E. Role of Macrophages in Host Resistance to Group A

Streptococci. *Infect. Immun.* **2004**, *72*, 2956–2963, doi:10.1128/IAI.72.5.2956-2963.2004.

32.   Ghoneim, H.E.; Thomas, P.G.; McCullers, J.A. Depletion of Alveolar Macrophages during Influenza

Infection Facilitates Bacterial Superinfections. *J. Immunol.* **2013**, *191*, 1250–1259,

doi:10.4049/jimmunol.1300014.

33.   Schultz, D.; Methling, K.; Rothe, M.; Lalk, M. Eicosanoid Profile of Influenza A Virus Infected Pigs.

Metabolites **2019**, *9*, 130, doi:10.3390/metabo9070130.

34.   Miller, M.A.; Stabenow, J.M.; Parvathareddy, J.; Wodowski, A.J.; Fabrizio, T.P.; Bina, X.R.; Zalduondo, L.;

Bina, J.E. Visualization of Murine Intranasal Dosing Efficiency Using Luminescent Francisella Tularensis:

Effect of Instillation Volume and Form of Anesthesia. *PLoS One* **2012**, *7*, e31359,

doi:10.1371/journal.pone.0031359.

35.   Volzke, J.; Schultz, D.; Kordt, M.; Müller, M.; Bergmann, W.; Methling, K.; Kreikemeyer, B.; Müller-Hilke,

B. Inflammatory Joint Disease Is a Risk Factor for Streptococcal Sepsis and Septic Arthritis in Mice. *Front.*

Immunol. **2020**, *11*, 1–15, doi:10.3389/fimmu.2020.579475.

36.   Shrum, B.; Anantha, R. V.; Xu, S.X.; Donnelly, M.; Haeryfar, S.; McCormick, J.K.; Mele, T. A Robust

Scoring System to Evaluate Sepsis Severity in an Animal Model. *BMC Res. Notes* **2014**, *7*, 233,

doi:10.1186/1756-0500-7-233.

37.   Amend, S.R.; Valkenburg, K.C.; Pienta, K.J. Murine Hind Limb Long Bone Dissection and Bone Marrow

Isolation. *J. Vis. Exp.* **2016**, *110*, doi:10.3791/53936.

38.   Belkina, A.C.; Ciccolella, C.O.; Anno, R.; Halpert, R.; Spidlen, J.; Snyder-Cappione, J.E. Automated

Optimized Parameters for T-Distributed Stochastic Neighbor Embedding Improve Visualization and

Analysis of Large Datasets. *Nat. Commun.* **2019**, *10*, 1–26, doi:10.1038/s41467-019-13055-y.

39.   Van Gassen, S.; Callebaut, B.; Van Helden, M.J.; Lambrecht, B.N.; Demeester, P.; Dhaene, T.; Saeys, Y.

FlowSOM: Using Self-Organizing Maps for Visualization and Interpretation of Cytometry Data. *Cytom. Part*

A **2015**, *87*, 636–645, doi:10.1002/cyto.a.22625.

40.   Gomes, R.N.; Teixeira-Cunha, M.G.A.; Figueiredo, R.T.; Almeida, P.E.; Alves, S.C.; Bozza, P.T.; Bozza,

F.A.; Bozza, M.T.; Zimmerman, G.A.; Castro-Faria-Neto, H.C. Bacterial Clearance in Septic Mice Is

Modulated by MCP-1/CCL2 and Nitric Oxide. *Shock* **2013**, *39*, 63–69,

doi:10.1097/SHK.0b013e31827802b5.

41.   Winter, C.; Taut, K.; Srivastava, M.; Länger, F.; Mack, M.; Briles, D.E.; Paton, J.C.; Maus, R.; Welte, T.;

Gunn, M.D.; et al. Lung-Specific Overexpression of CC Chemokine Ligand (CCL) 2 Enhances the Host

Defense to Streptococcus Pneumoniae Infection in Mice: Role of the CCL2-CCR2 Axis. *J. Immunol.* **2007**,

178, 5828–5838, doi:10.4049/jimmunol.178.9.5828.

42.　Nakamura, S.; Davis, K.M.; Weiser, J.N. Synergistic Stimulation of Type I Interferons during Influenza

Virus Coinfection Promotes Streptococcus Pneumoniae Colonization in Mice. *J. Clin. Invest.* **2011**, *121*,

3657–3665, doi:10.1172/JCI57762.

43.　Valderrama, J.A.; Nizet, V. Group A Streptococcus Encounters with Host Macrophages. *Future Microbiol.*

2018, *13*, 119–134, doi:10.2217/fmb-2017-0142.

44.　Carreno, D.; Wanford, J.J.; Jasiunaite, Z.; Hames, R.G.; Chung, W.Y.; Dennison, A.R.; Straatman, K.;

Martinez-Pomares, L.; Pareek, M.; Orihuela, C.J.; et al. Splenic Macrophages as the Source of Bacteraemia

during Pneumococcal Pneumonia. *EBioMedicine* **2021**, *72*, 103601, doi:10.1016/j.ebiom.2021.103601.

45.　Ercoli, G.; Fernandes, V.E.; Chung, W.Y.; Wanford, J.J.; Thomson, S.; Bayliss, C.D.; Straatman, K.;

Crocker, P.R.; Dennison, A.; Martinez-Pomares, L.; et al. Intracellular Replication of Streptococcus

Pneumoniae inside Splenic Macrophages Serves as a Reservoir for Septicaemia. *Nat. Microbiol.* **2018**, *3*,

600–610, doi:10.1038/s41564-018-0147-1.

46.　Horino, T.; Matsumoto, T.; Ishikawa, H.; Kimura, S.; Uramatsu, M.; Tanabe, M.; Tateda, K.; Miyazaki, S.;

Aramaki, Y.; Iwakura, Y.; et al. Interleukin-1 Deficiency in Combination with Macrophage Depletion

Increases Susceptibility to Pseudomonas Aeruginosa Bacteremia. *Microbiol. Immunol.* **2009**, *53*, 502–511,

doi:10.1111/j.1348-0421.2009.00143.x.

47.　Newman, J.H. Review of Septic Arthritis throughout the Antibiotic Era. *Ann. Rheum. Dis.* **1976**, *35*, 198–

205, doi:10.1136/ard.35.3.198.

48.　Layne, S.P.; Beugelsdijk, T.J.; Patel, C.K.N.; Taubenberger, J.K.; Cox, N.J.; Gust, I.D.; Hay, A.J.; Tashiro,

M.; Lavanchy, D. A Global Lab Against Influenza. *Science* **2001**, *293*, 1729–1729,

doi:10.1126/science.293.5536.1729.

49.　Manicassamy, B.; Manicassamy, S.; Belicha-Villanueva, A.; Pisanelli, G.; Pulendran, B.; Garcia-Sastre, A.

Analysis of in Vivo Dynamics of Influenza Virus Infection in Mice Using a GFP Reporter Virus. *Proc. Natl.*

Acad. Sci. **2010**, *107*, 11531–11536, doi:10.1073/pnas.0914994107.

50.　Perrone, L.A.; Plowden, J.K.; García-Sastre, A.; Katz, J.M.; Tumpey, T.M. H5N1 and 1918 Pandemic

Influenza Virus Infection Results in Early and Excessive Infiltration of Macrophages and Neutrophils in the

Lungs of Mice. *PLoS Pathog.* **2008**, *4*, e1000115, doi:10.1371/journal.ppat.1000115.

51.   Iwasaki, A.; Pillai, P.S. Innate Immunity to Influenza Virus Infection. *Nat. Rev. Immunol.* **2014**, *14*, 315–
328, doi:10.1038/nri3665.

52.   Tough, D.F.; Borrow, P.; Sprent, J. Induction of Bystander T Cell Proliferation by Viruses and Type I
Interferon in Vivo. *Science* **1996**, *272*, 1947–1950, doi:10.1126/science.272.5270.1947.

53.   Matikainen, S.; Pirhonen, J.; Miettinen, M.; Lehtonen, A.; Govenius-Vintola, C.; Sareneva, T.; Julkunen, I.
Influenza A and Sendai Viruses Induce Differential Chemokine Gene Expression and Transcription Factor
Activation in Human Macrophages. *Virology* **2000**, *276*, 138–147, doi:10.1006/viro.2000.0542.

54.   Ip, D.K.M.; Lau, L.L.H.; Leung, N.H.L.; Fang, V.J.; Chan, K.-H.; Chu, D.K.W.; Leung, G.M.; Peiris,
J.S.M.; Uyeki, T.M.; Cowling, B.J. Viral Shedding and Transmission Potential of Asymptomatic and Pauci-
Symptomatic Influenza Virus Infections in the Community. *Clin. Infect. Dis.* **2016**, ciw841,
doi:10.1093/cid/ciw841.

55.   Schwaiger, T.; Sehl, J.; Karte, C.; Schäfer, A.; Hühr, J.; Mettenleiter, T.C.; Schröder, C.; Köllner, B.; Ulrich,
R.; Blohm, U. Experimental H1N1pdm09 Infection in Pigs Mimics Human Seasonal Influenza Infections.
PLoS One **2019**, *14*, 1–21, doi:10.1371/journal.pone.0222943.

56.   Cuypers, F.; Schäfer, A.; Skorka, S.B.; Surabhi, S.; Tölken, L.A.; Paulikat, A.D.; Kohler, T.P.; Otto, S.A.;
Mettenleiter, T.C.; Hammerschmidt, S.; et al. Innate Immune Responses at the Asymptomatic Stage of
Influenza A Viral Infections of Streptococcus Pneumoniae Colonized and Non-Colonized Mice. *Sci. Rep.*
2021, *11*, 1–14, doi:10.1038/s41598-021-00211-y.

57.   Kovarik, P.; Castiglia, V.; Ivin, M.; Ebner, F. Type I Interferons in Bacterial Infections: A Balancing Act.
Front. Immunol. **2016**, *7*, 1–8, doi:10.3389/fimmu.2016.00652.

58.   Castiglia, V.; Piersigilli, A.; Ebner, F.; Janos, M.; Goldmann, O.; Damböck, U.; Kröger, A.; Weiss, S.;
Knapp, S.; Jamieson, A.M.; et al. Type I Interferon Signaling Prevents IL-1β-Driven Lethal Systemic
Hyperinflammation during Invasive Bacterial Infection of Soft Tissue. *Cell Host Microbe* **2016**, *19*, 375–
387, doi:10.1016/j.chom.2016.02.003.

59.   LeMessurier, K.S.; Häcker, H.; Chi, L.; Tuomanen, E.; Redecke, V. Type I Interferon Protects against
Pneumococcal Invasive Disease by Inhibiting Bacterial Transmigration across the Lung. *PLoS Pathog.* **2013**,
9, e1003727, doi:10.1371/journal.ppat.1003727.

60.   Maier, B.B.; Hladik, A.; Lakovits, K.; Korosec, A.; Martins, R.; Kral, J.B.; Mesteri, I.; Strobl, B.; Müller,
M.; Kalinke, U.; et al. Type I Interferon Promotes Alveolar Epithelial Type II Cell Survival during

Pulmonary Streptococcus Pneumoniae Infection and Sterile Lung Injury in Mice. *Eur. J. Immunol.* **2016**, *46*,

2175–2186, doi:10.1002/eji.201546201.

61. Antonelli, L.R.V.; Gigliotti Rothfuchs, A.; Gonçalves, R.; Roffê, E.; Cheever, A.W.; Bafica, A.; Salazar, A.M.; Feng, C.G.; Sher, A. Intranasal Poly-IC Treatment Exacerbates Tuberculosis in Mice through the Pulmonary Recruitment of a Pathogen-Permissive Monocyte/Macrophage Population. *J. Clin. Invest.* **2010**, *120*, 1674–1682, doi:10.1172/JCI40817.

62. Jia, T.; Leiner, I.; Dorothee, G.; Brandl, K.; Pamer, E.G. MyD88 and Type I Interferon Receptor-Mediated Chemokine Induction and Monocyte Recruitment during Listeria Monocytogenes Infection. *J. Immunol.* **2009**, *183*, 1271–1278, doi:10.4049/jimmunol.0900460.

63. Zimmerer, J.M.; Lesinski, G.B.; Radmacher, M.D.; Ruppert, A.; Carson, W.E. STAT1-Dependent and STAT1-Independent Gene Expression in Murine Immune Cells Following Stimulation with Interferon-Alpha. *Cancer Immunol. Immunother.* **2007**, *56*, 1845–1852, doi:10.1007/s00262-007-0329-9.

64. Zhang, Z.; Clarke, T.B.; Weiser, J.N. Cellular Effectors Mediating Th17-Dependent Clearance of Pneumococcal Colonization in Mice. *J. Clin. Invest.* **2009**, doi:10.1172/JCI36731.

65. Davis, K.M.; Nakamura, S.; Weiser, J.N. Nod2 Sensing of Lysozyme-Digested Peptidoglycan Promotes Macrophage Recruitment and Clearance of S. Pneumoniae Colonization in Mice. *J. Clin. Invest.* **2011**, *121*, 3666–3676, doi:10.1172/JCI57761.

66. Kristiansen, M.; Graversen, J.H.; Jacobsen, C.; Sonne, O.; Hoffman, H.-J.; Law, S.K.A.; Moestrup, S.K. Identification of the Haemoglobin Scavenger Receptor. *Nature* **2001**, *409*, 198–201, doi:10.1038/35051594.

67. Pulford, K.; Micklem, K.; McCarthy, S.; Cordell, J.; Jones, M.; Mason, D.Y. A Monocyte/Macrophage Antigen Recognized by the Four Antibodies GHI/61, Ber-MAC3, Ki-M8 and SM4. *Immunology* **1992**, *75*, 588–595.

68. Buechler, C.; Ritter, M.; Orsó, E.; Langmann, T.; Klucken, J.; Schmitz, G. Regulation of Scavenger Receptor CD163 Expression in Human Monocytes and Macrophages by Pro- and Antiinflammatory Stimuli. *J. Leukoc. Biol.* **2000**, *67*, 97–103.

69. Oliviero, S.; Cortese, R. The Human Haptoglobin Gene Promoter: Interleukin-6-Responsive Elements Interact with a DNA-Binding Protein Induced by Interleukin-6. *EMBO J.* **1989**, *8*, 1145–1151.

70. van den Heuvel, M.M.; Tensen, C.P.; van As, J.H.; van den Berg, T.K.; Fluitsma, D.M.; Dijkstra, C.D.; Döpp, E.A.; Droste, A.; van Gaalen, F.A.; Sorg, C.; et al. Regulation of CD163 on Human Macrophages:

Cross-Linking of CD163 Induces Signaling and Activation. *J. Leukoc. Biol.* **1999**, *66*, 858–866,

doi:10.1002/jlb.66.5.858.

71.   Fabriek, B.O.; van Bruggen, R.; Deng, D.M.; Ligtenberg, A.J.M.; Nazmi, K.; Schornagel, K.; Vloet, R.P.M.;

Dijkstra, C.D.; van den Berg, T.K. The Macrophage Scavenger Receptor CD163 Functions as an Innate

Immune Sensor for Bacteria. *Blood* **2009**, *113*, 887–892, doi:10.1182/blood-2008-07-167064.

72.   Kneidl, J.; Löffler, B.; Erat, M.C.; Kalinka, J.; Peters, G.; Roth, J.; Barczyk, K. Soluble CD163 Promotes

Recognition, Phagocytosis and Killing of Staphylococcus Aureus via Binding of Specific Fibronectin

Peptides. *Cell. Microbiol.* **2012**, *14*, 914–936, doi:10.1111/j.1462-5822.2012.01766.x.

73.   Reading, P.C.; Miller, J.L.; Anders, E.M. Involvement of the Mannose Receptor in Infection of Macrophages

by Influenza Virus. *J. Virol.* **2000**, *74*, 5190–5197, doi:10.1128/JVI.74.11.5190-5197.2000.

74.   Pontow, S.E.; Kery, V.; Stahl, P.D. Mannose Receptor. *Int. Rev. Cytol.* **1993**, *137*, 221–244,

doi:10.1016/S0074-7696(08)62606-6.

75.   Sallusto, F.; Cella, M.; Danieli, C.; Lanzavecchia, A. Dendritic Cells Use Macropinocytosis and the Mannose

Receptor to Concentrate Macromolecules in the Major Histocompatibility Complex Class II Compartment:

Downregulation by Cytokines and Bacterial Products. *J. Exp. Med.* **1995**, *182*, 389–400,

doi:10.1084/jem.182.2.389.

76.   Stahl, P.D. The Macrophage Mannose Receptor: Current Status. *Am. J. Respir. Cell Mol. Biol.* **1990**, *2*, 317–

318, doi:10.1165/ajrcmb/2.4.317.

77.   Upham, J.P.; Pickett, D.; Irimura, T.; Anders, E.M.; Reading, P.C. Macrophage Receptors for Influenza A

Virus: Role of the Macrophage Galactose-Type Lectin and Mannose Receptor in Viral Entry. *J. Virol.* **2010**,

84, 3730–3737, doi:10.1128/JVI.02148-09.

78.   Lee, S.J.; Zheng, N.-Y.; Clavijo, M.; Nussenzweig, M.C. Normal Host Defense during Systemic Candidiasis

in Mannose Receptor-Deficient Mice. *Infect. Immun.* **2003**, *71*, 437–445, doi:10.1128/IAI.71.1.437-

445.2003.

79.   Swain, S.D.; Lee, S.J.; Nussenzweig, M.C.; Harmsen, A.G. Absence of the Macrophage Mannose Receptor

in Mice Does Not Increase Susceptibility to Pneumocystis Carinii Infection In Vivo. *Infect. Immun.* **2003**, *71*,

6213–6221, doi:10.1128/IAI.71.11.6213-6221.2003.

80.   Gordon, S. Alternative Activation of Macrophages. *Nat. Rev. Immunol.* **2003**, *3*, 23–35, doi:10.1038/nri978.

81.   Wu, G.; Morris, S.M. Arginine Metabolism: Nitric Oxide and Beyond. *Biochem. J.* **1998**, *336*, 1–17,

doi:10.1042/bj3360001.

82. Starikova, E.A.; Sokolov, A. V.; Burova, L.A.; Golovin, A.S.; Lebedeva, A.M.; Vasilyev, V.B.; Freidlin, I.S.

The Role of Arginine Deaminase from Streptococcus Pyogenes in Inhibition Macrophages Nitrogen

Monooxide (NO) Synthesis. *Russ. J. Infect. Immun.* **2018**, *8*, 211–218, doi:10.15789/2220-7619-2018-2-211-

218.

83. Hesse, M.; Modolell, M.; La Flamme, A.C.; Schito, M.; Fuentes, J.M.; Cheever, A.W.; Pearce, E.J.; Wynn,

T.A. Differential Regulation of Nitric Oxide Synthase-2 and Arginase-1 by Type 1/Type 2 Cytokines In

Vivo: Granulomatous Pathology Is Shaped by the Pattern of l-Arginine Metabolism. *J. Immunol.* **2001**, *167*,

6533–6544, doi:10.4049/jimmunol.167.11.6533.

84. Serbina, N. V; Salazar-Mather, T.P.; Biron, C.A.; Kuziel, W.A.; Pamer, E.G. TNF/INOS-Producing

Dendritic Cells Mediate Innate Immune Defense against Bacterial Infection. *Immunity* **2003**, *19*, 59–70,

doi:10.1016/S1074-7613(03)00171-7.

85. Tavares, L.P.; Teixeira, M.M.; Garcia, C.C. The Inflammatory Response Triggered by Influenza Virus: A

Two Edged Sword. *Inflamm. Res.* **2017**, *66*, 283–302, doi:10.1007/s00011-016-0996-0.

86. Richter, J.; Brouwer, S.; Schroder, K.; Walker, M.J. Inflammasome Activation and IL-1β Signalling in

Group A Streptococcus Disease. *Cell. Microbiol.* **2021**, *23*, 1–9, doi:10.1111/cmi.13373.

87. Harder, J.; Franchi, L.; Munoz-Planillo, R.; Park, J.-H.; Reimer, T.; Nunez, G. Activation of the Nlrp3

Inflammasome by Streptococcus Pyogenes Requires Streptolysin O and NF- B Activation but Proceeds

Independently of TLR Signaling and P2X7 Receptor. *J. Immunol.* **2009**, *183*, 5823–5829,

doi:10.4049/jimmunol.0900444.

88. Lin, A.E.; Beasley, F.C.; Keller, N.; Hollands, A.; Urbano, R.; Troemel, E.R.; Hoffman, H.M.; Nizet, V. A

Group a Streptococcus ADP-Ribosyltransferase Toxin Stimulates a Protective Interleukin 1beta-Dependent

Macrophage Immune Response. *MBio* **2015**, *6*, 1–12, doi:10.1128/mBio.00133-15.

89. Valderrama, J.A.; Riestra, A.M.; Gao, N.J.; LaRock, C.N.; Gupta, N.; Ali, S.R.; Hoffman, H.M.; Ghosh, P.;

Nizet, V. Group A Streptococcal M Protein Activates the NLRP3 Inflammasome. *Nat. Microbiol.* **2017**, *2*,

1425–1434, doi:10.1038/s41564-017-0005-6.

90. Stasakova, J.; Ferko, B.; Kittel, C.; Sereinig, S.; Romanova, J.; Katinger, H.; Egorov, A. Influenza A Mutant

Viruses with Altered NS1 Protein Function Provoke Caspase-1 Activation in Primary Human Macrophages,

Resulting in Fast Apoptosis and Release of High Levels of Interleukins 1β and 18. *J. Gen. Virol.* **2005**, *86*,

185–195, doi:10.1099/vir.0.80422-0.

91. Park, H.S.; Lu, Y.; Pandey, K.; Liu, G.Q.; Zhou, Y. NLRP3 Inflammasome Activation Enhanced by

TRIM25 Is Targeted by the NS1 Protein of 2009 Pandemic Influenza A Virus. *Front. Microbiol.* **2021**, *12*,

doi:10.3389/fmicb.2021.778950.

92. Pothlichet, J.; Meunier, I.; Davis, B.K.; Ting, J.P.-Y.; Skamene, E.; von Messling, V.; Vidal, S.M. Type I

IFN Triggers RIG-I/TLR3/NLRP3-Dependent Inflammasome Activation in Influenza A Virus Infected

Cells. *PLoS Pathog.* **2013**, *9*, e1003256, doi:10.1371/journal.ppat.1003256.

93. Lamotte, L.A.; Tafforeau, L. How Influenza a Virus Ns1 Deals with the Ubiquitin System to Evade Innate

Immunity. *Viruses* **2021**, *13*, 1–26, doi:10.3390/v13112309.

94. Midiri, A.; Mancuso, G.; Beninati, C.; Gerace, E.; Biondo, C. The Relevance of Il-1-Signaling in the

Protection against Gram-Positive Bacteria. *Pathogens* **2021**, *10*, 1–13, doi:10.3390/pathogens10020132.

95. LaRock, C.N.; Todd, J.; LaRock, D.L.; Olson, J.; O'Donoghue, A.J.; Robertson, A.A.B.; Cooper, M.A.;

Hoffman, H.M.; Nizet, V. IL-1β Is an Innate Immune Sensor of Microbial Proteolysis. *Sci. Immunol.* **2016**,

1, eaah3539–eaah3539, doi:10.1126/sciimmunol.aah3539.

96. Ely, C.F. Influenza as Seen at the Puget Sound Navy Yard. *J. Am. Med. Assoc.* **1919**, *72*, 24,

doi:10.1001/jama.1919.26110010003009.

97. Okamoto, S.; Kawabata, S.; Fujitaka, H.; Uehira, T.; Okuno, Y.; Hamada, S. Vaccination with Formalin-

Inactivated Influenza Vaccine Protects Mice against Lethal Influenza Streptococcus Pyogenes

Superinfection. *Vaccine* **2004**, *22*, 2887–2893, doi:10.1016/j.vaccine.2003.12.024.

98. Grabenstein, J.D. Immunization to Protect the US Armed Forces: Heritage, Current Practice, and Prospects.

Epidemiol. Rev. **2006**, *28*, 3–26, doi:10.1093/epirev/mxj003.

99. Ozgur, S.K.; Beyazova, U.; Kemaloglu, Y.K.; Maral, I.; Sahin, F.; Camurdan, A.D.; Kizil, Y.; Dinc, E.;

Tuzun, H. Effectiveness of Inactivated Influenza Vaccine for Prevention of Otitis Media in Children.

Pediatr. Infect. Dis. J. **2006**, *25*, 401–404, doi:10.1097/01.inf.0000217370.83948.51.

100. Clements, D.A. Influenza A Vaccine Decreases the Incidence of Otitis Media in 6- to 30-Month-Old

Children in Day Care. *Arch. Pediatr. Adolesc. Med.* **1995**, *149*, 1113,

doi:10.1001/archpedi.1995.02170230067009.

Acknowledgment

The authors would like to thank Wendy Bergman for her excellent support during the work for this study. We also thank Dirk Koczan and Ildiko Toth for their help with the transcription analyses. Moreover, we like to express our gratitude to Karin Gerber and Chantal von Hörsten for their participation in the animal breeding and care at the animal core facility. The authors sincerely give their thanks to all staff and students that took part in the scientific discussions and experimental conduct.

Funding

This research was funded by Federal Excellence Initiative of Mecklenburg-Western Pomerania and European Social Fund (ESF) Grant KoInfekt (ESF/14-BM-A55-0011/16 and ESF/14-BM-A55-0005/16).

Ko-Infekt Study Group

Karen Methling[2], Michael Lalk[2], Ulrike Blohm[3], Alexander Schäfer[3], Bernd Kreikemeyer[4] (Affiliations are listed on the title page)

Author Contributions

J.V. and B.M-H. designed the study. J.V., M.B., E.W., M.M., D.S., K.M., U.B. and A.S. designed and performed the experiments and collected the data. J.V., M.B. and D.S. performed the statistical analyses. J.V. wrote the first draft of the manuscript. All authors contributed to the article and approved the submitted version.

Disclosures

The authors have no financial conflict of interest.

Abbreviations Used in This Article

In order of appearance:

IAV	Influenza A Virus	
GAS	Group A Streptococcus	
HA	Hemagglutinin	
IL	Interleukin	
IFN	Interferon	
MDCKII	Mardin-Darby Canine Kidney II (cells)	
TCID50	Tissue Culture Infectious Dose 50	
THB	Todd-Hewitt Broth	
CFU	Colony-forming Units	
20-HETE	20-Hydroxyeicosatetraenoic acid	
13-HODE	13-Hydroxyoctadecadienoic acid	
PGE$_2$	Prostanglandin E$_2$	
AA	Arachidonic Acid	
DMEM	Dulbecco's Modified Eagle's Medium	
FCS	Fetal Calf Serum	
M-CSF	Macrophage Colony-stimulating Factor	
RB	Running Buffer	
7-AAD	7-Aminoactinomycin	
MFI	Median Fluorescence Intensity	
t-SNE	t-distributed stochastic neighbor embedding	
CCL2/MCP-1	Chemokine CC-Motif Ligand 2/Monocyte Chemoattractant Protein-1	
TNFα	Tumor Necrosis Factor α	
CXCL2/MIP2-α	Chemokine CXC-Motif Ligand 2/Macrophage Inflammatory Protein 2-α	
ELISA	Enzyme-linked Immunosorbent Assay	
ANOVA	Analysis of Variance	
r	Pearson Product-moment Correlation Coefficient	

918	**SEM**	Standard Error of the Mean
919	**IFNAR**	Interferon-α/β Receptor
920	**NOS2**	Nitric Oxide Synthase 2
921	**ARG1**	Arginase
922	**NLRP3**	NLR Family Pyrin Domain Containing 3
923	**IL-1R**	Interleukin-1 Receptor

924 # Supplementary Material

925 Contents:

Figures

Figure 1. Influenza A virus infection induced minor weight loss and inhibited the production of CCL2. (**A**) Experimental Design. Mice were intranasally infected with influenza A virus (IAV, n = 8). PBS was adminstered as a control (n = 3). Blood samples were drawn on days 2 and 4 following infection. Lungs were excised at day 16. (**B**) Mean weight changes relative to day 0. Weight loss was confirmed by one-way ANOVA (p < 0.0001) in the IAV group and by two-way ANOVA (***p < 0.001) comparing the IAV group to PBS controls. Error bars depict the SEM. (**C**) Boxplots display CCL2 concentrations in plasma samples from uninfected controls (n = 13) and IAV infected mice (n = 9). The differences in samples sizes between days 2 and 4 are due to the fact that some animals were later subjected to the bacterial and co-infection experiments that are shown in Figure 2. Day 2 and Day 4 p-values result from Mann-Whitney U test and t-test, respectively. (**D**) Dotplots show normalized mRNA expression of *Ccl2* and *Ifnb1* in lung homogenates based on ΔCt values. p-value results from t-test. (**E**) Normalized viral loads based on Ct values for IAV specific genes in day 16 lung homogenates. HA: hemagglutinin, M: matrix protein, N: nucleoprotein.

Figure 2. Preceding IAV infection promotes bacterial dissemination and sepsis severity during co-infection.
(**A**) Experimental design. For monocausal bacterial infection, Group A Streptococcus was administered intravenously (GAS, n = 10). For co-infection, mice were either infected with GAS followed by intranasal IAV administration (GAS+IAV, n = 10) or infected with IAV followed by infection with GAS (IAV+GAS, n = 10). (**B**) Representative photographies of a healthy compared to an edematous paw after IAV+GAS co-infection. (**C**) Kaplan-Meier curves display the incidences of paw edemas. *p < 0.05, log-rank test with p-value adjustment for multiple comparisons (Bonferroni-Holm method). (**D**) Dotplot shows sepsis scores at endpoints (day 14 or humane endpoint). p-value results from Mann-Whitney U test. The dashed line indicates the minimum score for humane endpoints. (**E**) Kaplan-Meier curves display survival probabilities. (**F**) Dotplots show endpoint bulk lung mRNA gene expressions of *Ccl2* and *Ifnb1*that were normalized to *Gapdh* and lungs from uninfected mice (dashed line) by the $2^{-\Delta\Delta Ct}$ method. *p < 0.05, **p < 0.01, one-sample Wilcoxon signed-rank test (μ = 1). (**G**) Normalized bacterial loads based on Ct values for GAS specific genes in endpoint lung homogenates.

Figure 3. Activation of bone-marrow derived murine macrophages during *in vitro* co-infection was distinct from bacterial mono-infection. (**A**) Experimental Design. Macrophages were differentiated from bone marrow cells and then infected with either IAV for 48 h (n = 4) or GAS for 6 h (n = 4). For co-infection, IAV was first applied for 48 h followed by GAS infection for 6 h (IAV+GAS, n = 7). Each sample was obtained from individual mice to obtain biological replicates. (**B**) t-distributed stochastic neighbor embedding (tSNE) on flow cytometry data from the three different macrophage infection models and topology of surface antigen expression levels. 10,000 events from each sample were integrated into the dimension reduction analysis. MFI: median fluorescence intensity. (**C**) Heatmap and hierarchical clustering on standardized fold changes of surface antigen expression levels based on their MFI. Fold changes were generated by normalization of MFI data from infected macrophages to their respective paired uninfected controls. (**D**) Dotplots depict the alteration of surface antigen expression levels due to (co-)infection. (**E**) Representative pseudocolor plots illustrate the alteration of proportions of macrophages expressing CD80 and CD86 (top) or CD206 (bottom) after (co-)infection. (**F**) Dotplots demonstrate the shift of macrophage subpopulation fractions after (co-)infection relative to uninfected controls (dashed lines). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Dunn's test or Tukey HSD test with p-value adjustments for multiple comparisons (Bonferroni-Holm method). [#]$p < 0.05$, [##]$p < 0.01$, [###]$p < 0.001$, Wilcoxon signed-rank test or one-sample t-test for the comparison to uninfected cultures ($\mu = 1$).

Figure 4. Preceeding influenza A virus infection impedes pro-inflammatory immunological features of

macrophages during co-infection. (**A**) Experimental Design. Bone-marrow derived macrophages were infected

with IAV (n = 4), GAS (n = 4) or IAV+GAS (n = 7). Each sample was obtained from individual mice to obtain

biological replicates. (**B**) Heatmap and hierarchical clustering on standardized relative mRNA expression levels from

quantitative PCR analyses using the $2^{-\Delta\Delta Ct}$ method. Data from infected cultures were normalized to *Gapdh* and their

respective paired uninfected controls. (**C**) Dotplots show the alterations of *Mgl2*, *Nos2* and *Arg1* mRNA expression

levels due to (co-)infection. (**D**) Dotplots illustrate distinct patterns of chemokine and cytokine mRNA production by

macrophages after (co-)infection. (**E**) Dotplots demonstrate (co-)infection induced protein production of chemokines

and cytokines that were measured in cell culture supernatants. Dashed lines represent control cultures. *p < 0.05, **p

< 0.01, ***p < 0.001, Dunn's test or Tukey HSD test with p-value adjustments for multiple comparisons (Bonferro-

ni-Holm method). #p < 0.05, ##p < 0.01, ###p < 0.001, Wilcoxon signed-rank test or one-sample t-test for the compari-

son to uninfected cultures (μ = 1).

Tables

Table I. Frequencies of blood agar cultures from endpoint blood smears and synovial knee joint swabs positive for

β-hemolytic bacteria.

positive cultures	GAS	GAS+IAV	IAV+GAS
blood	20% (2/10)	30% (3/10)	50% (5/10)
knee joint capsule	10% (1/10)	30% (3/10)	50% (5/10)

998 $p = 0.25$, Fisher's exact test