Elevated human footprint on islands promotes both introduction and extinction probability of insular reptiles at opposite ends of geographic, evolutionary and ecological continua.

Jesse, W.A.M.[1]

Ellers, J. [1]

Behm, J.E.[2]

Costa, Gabriel C.[3]

Hedges, S. Blair [2]

Helmus, M.R. [2]

[1] Department of Ecological Science, Vrije Universiteit Amsterdam, De Boelelaan 1085
Amsterdam, NL 1081 HV

[2] Department of Biology, Temple University, 1925 N. 12th Street Suite 502, Philadelphia, PA
19122

[3] Department of Biology and Environmental Sciences, Auburn University at Montgomery,
Montgomery, AL, USA

Abstract

The Anthropocene is marked by unprecedented changes to species ranges worldwide. Introduced

species expand their native range into new geographic regions while extinction-prone species

experience severe native range contraction. Introductions and extinctions are both caused by how

species respond to human influence within their native ranges. The question arises if the opposite

response in distributional change is due to introduced and extinction-prone species falling at

opposite ends of geographic, evolutionary and ecological trait continua. Here, we assess the factors

that explain the introduction and extinction-prone status for 3111 squamate reptile species native

to the western hemisphere, of which 142 species have been introduced elsewhere and 483 species

are threatened with extinction. We first asked what geographic regions contained the most

introduced and extinction-prone species and if these species are evolutionarily related. We then

tested the opposite ends hypothesis for several species characteristics. Our results show that the

native ranges of introduced and extinction-prone reptiles are skewed towards island regions and

that significant phylogenetic signal exists for these two groups, indicating they are not a random

selection of the reptiles that have evolved in the western hemisphere. Species whose native ranges

were more insular and were more influenced by humans were more likely to be introduced or

extinction-prone. Evolutionary factors did not explain species status, but multiple geographic and

ecological factors supported the opposite ends hypothesis. Introduced species were positively

associated with herbivory and omnivory and had large and complex ranges, whereas extinction-

prone species were generally carnivorous with small and simple ranges. In addition, introduced

species produced larger clutches and extinction-prone species had relatively small body sizes. In

the Anthropocene, the naive ranges of introduced and extinction-prone species are situated in the

same human-impacted regions, but species range characteristics and functional traits determine

whether species ranges expand or contract in the continuing face of global change.

Introduction

In the Anthropocene, humans have altered species distributions and reassembled global biogeographic patterns that arose naturally across eons of evolution (Alroy 2015, Capinha et al. 2015, Ceballos et al. 2017). The "winners" of the Anthropocene are introduced species, which expand into new geographic regions through human-vectored dispersal and establishment (Colautti and MacIsaac 2004). The "losers" are the extinct and threatened species that have experienced severe native range retraction often because of human-associated stress factors such as overharvesting, pollution, and land-use change (Vitousek et al. 1997, Sax and Gaines 2008, Böhm et al. 2013, Young et al. 2016). Species that are introduced to other regions benefit from having native ranges that encompass high human populations because of increased likelihood of intentional (e.g. via pet trade) or unintentional (e.g., as stowaway) human-aided dispersal (Latella et al. 2011, Liu et al. 2014, Su et al. 2016). After introduction, species with suitable functional traits are better able to utilize novel food sources and endure local climatic regimes (Mahoney et al. 2015, Monaco et al. 2020). In contrast, extinction-prone species generally are ecological specialists, characterized by narrow dietary and climatic niches and poor dispersal abilities due to their geographic restriction to mountain tops, islands, or isolated nature reserves in a matrix of anthropogenic land use (Kotiaho et al. 2005, Böhm et al. 2016). Hence, introduced and extinction-prone species appear to be each other's antipodes in many characteristics that cause opposite distributional responses in the Anthropocene. This raises the question if these winners and losers are at opposite ends of the same geographic, evolutionary and ecological trait continua (Jeschke and Strayer 2008, Blackburn and Jeschke 2009). Testing the opposite ends hypothesis requires identification of functional traits, native geographic range characteristics, and phylogenetic

clustering associated with introduced and extinction-prone species, followed by an analysis to

determine the (dis)similarity between these two groups.

The few studies that test the opposite ends hypothesis show variable results. For instance,

extinction-prone and successfully introduced plants (Bradshaw et al. 2008, Pandit et al. 2011,

Schmidt et al. 2012) and crayfish (Larson and Olden 2010) were on opposite extremes of the same

trait axes associated with body size, fecundity, genetic diversity and generalist versus specialist

habitat use. Conversely, studies on birds, fishes (Jeschke and Strayer 2008, Blackburn and Jeschke

2009) and reptiles (Tingley et al. 2016) found different set of traits to characterize either extinction-

prone or introduced species. Furthermore, such comparative studies generally lack an (in-depth)

phylogenetic and geographic perspective on species introduction and extinction probabilities,

possibly obscuring or exaggerating observed effects by not accounting for phylogenetic

interdependence of species.

Nonoverlapping phylogenetic clustering of introduced and extinct species is expected under

the opposite ends hypothesis because many species characteristics exhibit phylogenetic signal.

Phylogenetic clustering of extinction-prone species has been detected in plants (Davies et al. 2011,

Loza et al. 2017), fish (Adeoba et al. 2019), bumblebees (Arbetman et al. 2017), birds, and

mammals (Fritz and Purvis 2010). In addition, moderate to strong levels of phylogenetic signal

have been found in traits associated with introduction and/or invasion success (Cadotte et al. 2009),

dispersal capacity (Park and Potter 2015), native range size (Pigot et al. 2018), and fecundity (Allen

et al. 2017) for plants (Cadotte et al. 2009, Yessoufou et al. 2016, Pyšek et al. 2017), fish (Alcaraz

et al. 2005), and birds (Su et al. 2016). However, the proposed reasons for this phylogenetic signal

vary considerably between studies. For instance, Fritz and Purvis (2010) ascribe phylogenetic

signal to the distribution of life-history traits associated with attractiveness, making species

sensitive to over-harvesting. In contrast, Davies and colleagues (2011) detected a phylogenetic

signal for extinction risk that is unrelated to the distribution of specific traits, but associated with

young, recently diversified lineages within a single geographic region. Overall, there is compelling

evidence that the predisposition to becoming introduced or extinct is phylogenetically clustered

and should be considered when testing for opposite ends of the trait continua.

Native geographic range characteristics associated with range expansions and contractions

should underlie how species respond to human influence. In particular, island-living may be an

important characteristic determining the likeliness of introduction and extinction. Islands generally

have high population densities and are subjected to disproportionate levels of human influence

(Kier et al. 2009), thus insular species are much more likely to be introduced or go extinct than

mainland species. Islands are prime locations of species loss (Myers et al. 2000, Mittermeier et al.

2011) and also provide unique environments that select for traits that naturally determine major

range expansions or contractions. Insular species have evolved through processes of oversea

colonization, followed by often rapid adaptive diversification (Cowie and Holland 2006, Hedges

2006, 2011). Thus, insular species are selected to be efficient dispersers and quick adapters to

available niche space, which could favor survival of human-vectored dispersal events (Poe et al.

2011). Indeed, post-introduction ecological niche shifts occur often for introduced species that

originate from oceanic islands (Liu et al. 2014). In contrast, adaptive radiations cause island biota

to be highly specialized, extremely suited to exploit a narrow ecological niche within island

environments (Losos 2009, Mahler et al. 2010, 2013). This level of specialization leaves island

endemics sensitive to human-caused (a)biotic change and thus prone to extinction (Jantz et al.

2015).

In this study, we used a phylogenetic comparative approach to identify (dis)similarities in

introduced and extinction-prone species characteristics. We used a comprehensive dataset of 13

characteristics of 3111 squamate reptiles (lizards and snakes) native to the western hemisphere

(Figure 1). Of these, 142 species have been introduced to at least one location somewhere in the

world and 483 species are threatened with extinction according to the International Union for the

Conservation of Nature (IUCN). Reptiles are particularly suitable to test opposite responses to

human influence, as reptiles are species-rich in the western hemisphere and are greatly impacted

by human activities (Young et al. 2016, Jesse et al. 2018). We analyzed this data set to ask: 1) Are

islands of the western hemisphere disproportionate sources of introduced and sinks of extinction-

prone species? 2) Can we detect phylogenetic signal among introduced and extinction-prone

species? 3) For which geographic, evolutionary and ecological trait continua are introduced and

extinction-prone species positioned at opposite ends?

Methods

Data compilation

<u>Squamate phylogeny</u>

To robustly test for (dis)similarities in introduced and extinction-prone species characteristics

across western hemisphere squamates, a phylogenetic tree of all species was needed. Thus, we

began our data compilation by updating a global, smoothed, and interpolated phylogeny of

squamates from the TimeTree of life project (Kumar et al. 2017), previously used in (Marin et al.

2018, Rapacciuolo et al. 2019) with species synonyms from The Reptile Database (Uetz, P., Freed,

P. & Hošek 2012) to match up with the spatial range dataset that we obtained on western

hemisphere squamates (see "Geographic data" below). Any species not in the phylogeny and not

in the spatial dataset was removed from further data compilation and not included in our analyses.

This phylogeny of 3111 species (Figure 2) was used to calculate phylogenetic signal among

introduced and extinction-prone species and included in multivariable phylogenetic generalized

linear models (PGLMs).

<u>Species introduction and extinction-prone status</u>

The response variables in our analyses were based on species statuses of squamates across the

western hemisphere. We constructed a binomial variable to indicate whether a species was

introduced into some area of earth or not and a response variable that indicated if a species was

extinction-prone in its native range or not. To determine introduction status, we cross-referenced

3111 species from the phylogeny for which we had spatial native range data (see "Geographic data"

below) with published invasive species databases, such as the Global Register of Introduced and

Invasive Species, the Global Invasive Species Database and the Invasive Species Compendium, as

well as several multispecies scientific studies (e.g., Kraus 2009, Powell et al. 2011, 2012, Mahoney

et al. 2015, Krysko et al. 2016, Tingley et al. 2016, Helmus et al. 2017; see Supplement S1 for the

full list of references). For all species that did not match with these databases, we performed a

Google query for each species using the search term "*Species name* AND introduced OR exotic

OR invasive OR alien OR nonnative OR non-indigenous" to look for species-specific literature

that indicates it as introduced regardless of whether an exotic population is currently extant. We

excluded species for which introduction records were not scientifically published, uncertain or

debated. To estimate extinction risk, we reviewed IUCN records for all species with spatial native

range data (accessed August 2018). Species that were "vulnerable", "endangered", "critically

endangered", "extinct in the wild", and "extinct" were all categorized as extinction-prone, and

species with "near threatened" and "least concern" status were considered to be not threatened with

extinction, consistent with the categorization by IUCN (IUCN Standards and Petitions

Subcommittee 2019). Of the 3111 species, 349 of category "data deficient" and 494 of category

"not assessed" were excluded from the analyses (see "Analyses" below).

<u>Geographic data</u>

Distributional range maps for 3463 reptile species were compiled from the IUCN (IUCN Redlist

2021) and the Caribherp Caribbean island databases (Hedges 2021) and edited to only include

polygon shapes of native ranges of squamates in the western hemisphere which also appeared in

the phylogenetic tree. The resultant dataset included 3111 species. This difference in the number

of species between datasets was caused by the mismatch between phylogeny and geographic ranges

and removal of duplicate ranges, taxonomic synonyms, exotic ranges, and non-squamate species.

We used these range polygons to calculate independent covariates (italicized subheadings below)

for our statistical analyses.

Range size and complexity

Species range characteristics are thought to influence introduced and extinction-prone status. For

each species, we calculated native range size and complexity. First, range area (m^2) and range

perimeter (m) were estimated with the "areaPolygon" and "perimeter" functions, respectively, from

the geosphere R package (Hijmans 2019). Size was estimated as range area and complexity was

estimated as a normalized perimeter where range perimeter values are standardized for circularity

by dividing the measured perimeter of species range by a hypothetical perimeter of a circle with

the same surface area (variation of method proposed by Cox (1927) and Polsby and Popper (2017):

$$normalized\ perimeter_i = \frac{(2\pi r)_i}{2\sqrt{\pi * (\pi * r^2)_i}}$$

Low values indicate more circular ranges and high values indicate deviations from a circular range.

Species with high range complexity values thus have ranges that are unevenly spread across a

landscape and/or include many geographic boundaries (e.g., multiple islands).

Species insularity

For each species we calculated its degree of insularity as the portion of a species range located on

oceanic islands. We intersected species range polygons with a map of the continental mainland of

North and South America (GADM 2018), enabling us to identify the mainland continental ranges

for all species. Based on the area difference between the total species range and the continental

range, we were able to infer the insular fraction of a species range. An insularity value of 1 indicates

exclusive island living (i.e., island endemic) and values of 0 indicated that species only occur on

the continental mainland.

Evolutionary data

Genus age

Genus age has been associated with adaptability of species, and can be indicative of whether

species has been part of a recent radiation or has been able to persist for a long time (Davies et al

2011, Title & Burns 2015). Genus age was determined by calculating the branching-time from the

most recent common ancestor of the genus (i.e., the crown age) with the 'AssessMonophyly' and

'branching.time' functions from the 'ape' R package. Of 280 western hemisphere genera, 212 were

monophyletic, 65 were monotypic (a genus containing 1 species) and 3 were paraphyletic

(*Epicrates*, *Leposoma*, *Homonota*). For monotypic genera, we took the branching-time from the

most recent node connecting the focal tree tip as genus age. The intruder or outlier clades in

paraphyletic genera were all monophyletic or monotypic for which we could obtain a reliable age.

After close examination of the paraphyletic genera, we concluded that division into smaller clades

would not benefit our analysis nor significantly change our outcomes. We thus used the most recent

common ancestor of paraphyletic genera for further analysis.

Evolutionary range expansion rate

The rate at which a clade has expanded its range in the past has been suggested to be a proxy for

the potential of its species to do so in the present (Title & Burns 2015). We summed the species

range areas for all the species within the same genus to represent the cumulative range area that a

genus has spread into since the time of speciation from the most recent common ancestor.

Cumulative range area was divided by the focal genus age to calculate the average evolutionary

range expansion rate of geographic spread for all congeners.

Ecological data

Functional traits

We obtained data on functional traits thought to be related to opposite distributional responses to

human influence (Jeschke and Strayer 2008, Blackburn and Jeschke 2009, Tingley et al. 2016).

Natural log-transformed values of clutch size and maximum body size, as well as two categorical

variables for diet (herbivorous, carnivorous, and omnivorous) and reproductive mode (binomial:

oviparous and (partially) viviparous), were included as independent variables in PGLMs. These

trait data were from Rapacciuolo et al. 2019.

Seasonality and elevation

The environment that a species has adapted to in their native range may be associated with

distributional changes in the Anthropocene. We used Principal Component Analysis (PCA) to

extract composite variables that explain the climatic variation experienced among species. We

calculated the median of all 37 climatic variables in WorldClim and ENVIREM raster layers per

species range (Fick and Hijmans 2017, Title and Bemmels 2018). All range medians were

subsequently included in a scaled PCA in R, of which the first two principal components were

selected, and explained a cumulative proportion of 70% of the median environmental variation

among all species ranges. The first axis (PC1) aligned with climatic seasonality variables such as

annual temperature range, diurnal temperature range, and seasonal variation in potential

evapotranspiration (PET), and the second axis (PC2) aligned with two elevation-derived variables

from the ENVIREM dataset: terrain roughness and topographic wetness (indicative for the

catchment of water in a watershed) (Title and Bemmels 2018; See Supplement S2 for biplot).

Climatic niche differentiation

Species with little ability to adapt to a changing climate should be more extinction-prone, while

those with high adaptive potential should be more likely to establish in a new environment once

introduced. To estimate a species' adaptive capacity, we assessed the climatic variability in species'

ranges as indication of the various climatic regimes a species has adapted to. We calculated the

standard deviations of the 37 WorldClim and ENVIREM climatic variables per species range and

included these as new variables in a scaled PCA. PC1 of this PCA (explaining 63% of all variation)

was positively correlated to variability for all climatic variables (See supplement S3). Thus, PC1

score per species was taken as a value for climatic variability in a species range, which was

positively correlated with log-transformed range area ($r = 0.65$, $t_{3054} = 47.45$, $P < 0.001$), i.e.,

species with large ranges experience high levels of climatic variation. We used the residuals of this

linear relationship to reflect climatic niche differentiation relative to a species range size, indicating

disproportionate climatic niche diversification/uniformity as a measure of adaptive capacity

(hereafter referred to as climatic niche differentiation).

<u>Anthropogenic data</u>

<u>Human footprint and number of ports</u>

Unlike the previous species traits and characteristics described above where we expect introduced

and extinction-prone species to be at opposite ends of the continua, anthropogenic metrics are

expected to show positive relationships with both introduced and extinction-prone status, because

both processes are (likely) human-mediated. We determined the number of ports (incl. harbors,

seaports, and major terminals) per native range area (World Port Index; National Geospatial-

Intelligence Agency 2017) as a proxy for the level of seaborne trade to and from the species native

range. While trade clearly increases introductions directly, it also should increase extinctions

indirectly by indicating increased potential of introduced predators, pathogens and competitors.

We also calculated the median human footprint index (Venter et al. 2016) for every species range,

which varies between 0 to 50 and is a compilation of several human pressures, including the amount

of built environment, population density, agriculture, and several types of terrestrial and

waterborne infrastructure. Like the number of ports, we expected human footprint to increase both

introductions and extinctions. The number of ports and human footprint index were not

significantly correlated ($r=-0.02$, $t_{2988}=-0.98$, $P=0.33$), and therefore both were included in the

statistical analyses.

Analyses

We first asked if islands are sources of introduced and sinks of extinction-prone species.

We made maps of squamate species richness and proportions of introduced and extinction-prone

richness in relation to total species richness using 1 km^2 grid cells. This fine grain represents the

relevant scale for the occurrence of introduction and extinction particularly on islands, because

many islands are small. Further, the Caribherp island database has explicitly made ranges at this

fine scale resolution (Hedges 2021). We then calculated the proportion of species with introduced

or extinction-prone status whose ranges were found only on the mainland, mainland and islands,

or only islands; and tested if our species insularity covariate explained introduced and extinction-

prone status with chi-squared tests.

Second, we asked if there was phylogenetic signal among introduced and extinction-prone

species. The level of phylogenetic signal among introduced and extinction-prone species was

determined with the phylo.d-function for binomial data ('caper' package). The observed

distributions of introduced and extinction-prone species was compared to 1000 permutations of a

random trait distribution and a clumped trait distribution assuming Brownian trait evolution.

Finally, we tested the opposite ends hypothesis by asking which ecological, geographic and

evolutionary characteristics predict introduction and extinction-prone species status. To do this, we

used multivariate phylogenetic generalized linear models (PGLMs) to test the dichotomous

variables of introduction and extinction status against all predictor variables (see text above) with

logistic error term and Firth's penalized likelihood correction ('phyloglm' function, 'phylolm'

package). Two separate PGLMs with introduction and extinction status as dependent variables

were performed with range insularity as the sole independent factor, because of collinearity issues

between insularity and most other predictors (Supplement S4). All other predictor variables had

relatively low levels of collinearity ($r \leq |0.60|$ and VIF < 3.0). Prior to all analyses, we scaled the

continuous variables (mean = 0, sd = 1), so model estimates represented effect sizes. All analyses

were executed in R statistics, version 4.0.3 (R Development Core Team 2021) within the R studio

environment (R Studio Team 2021).

Results

The geographic distributions of introduced and extinction-prone species were strikingly different from the distribution of all reptiles in the western hemisphere (Figure 1). For instance, the native ranges of the 142 introduced species in our dataset spanned most of the western hemisphere and were more prevalent in North America and the Caribbean than South America, opposite to the cumulative distribution of all reptiles (Figure 1A,B). In contrast, the ranges of the 483 extinction-prone species in our dataset were much more localized and concentrated to specific endangered ecosystems and mountainous regions such as the Andes, the Pampas, Mesoamerican cloud forests and the Caribbean (Figure 1C).

The native ranges of introduced and extinction-prone species were similar to each other in that these species were often found on islands. Of all species in our dataset, 16% were island endemics; 32% had ranges encompassing both island and mainland regions, and a majority of 52% had native ranges only on the mainland, primarily Amazonia (Figure 1A). In contrast, 91% of the introduced reptiles had an association with island-living, either as island endemics (44%) or with oceanic islands in their native range (47%, Figure 1B). On many islands, >50% of the native species have been introduced elsewhere, making islands key sources of introduced reptiles globally. Similarly, island endemics constituted 41% of the extinction-prone species, and on many islands >30% of species are threatened with extinction (Figure 1C). This clear overrepresentation of island endemics in introduced ($\chi^2 = 47.83$, df = 1, $P < 0.001$) and extinction-prone reptile groups ($\chi^2 = 164.67$, df = 1, $P < 0.001$) indicates that islands are simultaneous sources and sinks of biodiversity in the Anthropocene.

Figure 1. The geographic distribution of reptile species richness in the western hemisphere, and identification of sources and sinks of reptile biodiversity in the Anthropocene. A) The total of overlapping native species ranges per 1km² grid cell. Proportions of introduced (B) and extinction-prone (C) richness in relation to total species richness were calculated for all 1 km² grid cells inhabited by >1 species. Points in all maps represent mean values per island and were plotted as such to increase visibility. Points follow the same color scheme as the gridded layers. Pie charts represent the geographic context of the species ranges depicted in the respective maps, showing the proportions of species that are restricted to the continental mainland of North and South America (white), partially insular species that inhabit island and mainland areas (light grey), and species that are restricted to oceanic islands (dark grey). Percentages within pie charts are (clockwise, i.e., *mainland only, island and mainland, island only*) (A) 52%, 32%, 16%, (B) 9%,

47%, 44% , (C) 53%, 6%, 41%. All maps were projected as Eckert IV spherical world projection

in ESRI ArcMap 10.6.1.

Figure 2. Significant phylogenetic signal among introduced and extinction-prone reptiles.

Introduced (142 spp) and extinction-prone reptiles (483 spp) plotted across the tree of life (3111

spp total). Silhouettes are of genera containing proportionally high numbers of introduced and/or

extinction-prone   species   (clockwise: *Boa,   Sphaerodactylus,   Ameiva,   Scincus,   Iguana,*

Cnemidophorus, Anolis, Typhlops).

Introduced and extinction-prone species were non-randomly distributed across the
phylogeny of Western Hemisphere reptiles (Figure 2). The signal for introduced (D = 0.63, *P* <
0.001) and extinction-prone (D = 0.71, *P* < 0.001) species was intermediate in strength and
significantly different from random and clumped null models based on 1000 permutations. Further
multivariate testing was done using phylogenetic generalized linear models to account for
phylogenetic non-randomness of observations. While introduced and extinction-prone species
exhibited phylogenetic clustering, the composition of the two groups showed little overlap. Of the
142 introduced and 483 extinction-prone squamates, only 15 species were included in both
categories. Of these 15 species, eight are popular species in the pet trade and threatened with
extinction in their native range (e.g., iguanas of the genus *Cyclura*).

Range insularity (the proportion of species range overlap with oceanic islands) was
positively related to both introduction (0.39 ± 0.08, $z = 4.45$, *P* < 0.001) and extinction probabilities
(0.54 ± 0.05, $z = 10.52$, *P* < 0.001; Figure 3A). Range insularity showed strong collinearity with
other variables in our analyses, most prominently with range area ($r = -0.39$, $t = -23.3$, *P* < 0.001),
diet type (Kruskal-Wallis: $\chi^2 = 18.2$, df = 2, *P* < 0.001; bias towards omnivory), and human
footprint ($r = 0.46$, $t = 29.1$, df = 3056, *P* < 0.001) as well as significant correlations with range
complexity (negative), parity (association with oviposition), clutch size (negative), climatic niche
differentiation (negative), genus age (positive), and evolutionary range expansion rate (negative)
(see Supplement S4 for pairwise correlations between all variables). This indicates that island-
living can be seen as a trait "syndrome": a set of correlations among individual phenotypic traits.
Therefore, in our PGLM analyses range insularity was excluded from the set of trait continua used
to test if introduced and extinction-prone species are positioned at opposite ends. After exclusion
of range insularity from the models, human footprint was the only other explanatory variable that
was positively related to introduction and extinction probability (Figure 3B). The underlying cause

of this is likely to be the significantly higher human footprint in species ranges located on oceanic

islands than on the mainland (t = 22.03, df = 653.79, $P < 0.001$; Figure 3C), as well as in the ranges

of introduced and extinction-prone species compared to other western hemisphere reptiles (t = 3.76,

df = 145.41, $P < 0.001$ and t = 11.94, df = 579.02, $P < 0.001$, respectively; Figure 3C).

Several explanatory variables had significant opposing effects on introduced and

extinction-prone status consistent with the opposite ends hypothesis (i.e. diet, range area, and range

perimeter in Figure 3B). Native ranges of introduced species were larger (0.78 ± 0.14, z = 5.58, P

< 0.001) and more complex (0.74 ± 0. 12, z = 6.32, $P < 0.001$) than the ranges of other western-

hemisphere reptiles, while extinction-prone ranges were smaller (-0.86 ± 0.07, z = -12.64, $P <$

0.001) and relatively circular (-0.15 ± 0.05, z = -2.83, $P = 0.005$). Extinction-prone and introduced

species also had divergent diets, with extinction-prone species adhering to a carnivorous, rather

than omnivorous (-1.70 ± 0.22, z = -7.62, $P < 0.001$) or herbivorous diet (-2.19 ± 0.41, z = -5.37,

$P < 0.001$), while introduced species were positively related to herbivory (1.27 ± 0.47, z = 2.70, P

= 0.007) and omnivory (0.68 ± 0.28, z = 2.39, $P = 0.02$; carnivory being the contrast level in the

PGLM analysis, not shown). Other predictor variables also showed opposite effects for introduced

and extinction-prone species, but did not differ significantly. In total, seven of the predictor

variables supported the opposite ends hypothesis (Figure 3B). Finally, clutch size had relatively

small single effect with introduced species having larger clutches (0.25 ± 0.12, z = -2.10, $P = 0.04$).

Figure 3. Species functional traits and characteristics as predictors for introduced and extinction-prone status. A) Range insularity is positively related to introduction (blue) and extinction (red) probabilities of western hemisphere reptiles. Bars are standardized logistic PGLM effect sizes (i.e., model estimate/standard error). B) Associations between range and species characteristics with introduction and extinction probabilities. A total of 2936 species, of which 128 were introduced, and 2148 of which 435 extinction-prone, were included in separate PGLM analyses. Bars are ordered according to the level of congruent responses between introduced and extinction-prone species. C) Human footprint is significantly higher within exclusively insular species ranges (range insularity = 1) compared to (partially) mainland ranges (range insularity < 1) and significantly higher for introduced and extinction-prone species compared to other reptiles. $P < 0.05$ (*), $P < 0.01$ (**), $P < 0.001$ (***).

Discussion

Range insularity and human footprint are important shared characteristics of introduced and extinction-prone species, indicating highly impacted, oceanic islands as simultaneous sources and sinks of biodiversity in the Anthropocene. In addition, we found significant phylogenetic signal among introduced and extinction-prone species, illustrating that these species are not a random draw from the tree of life, but evolutionarily predisposed to success or failure when faced with global change. In contrast to the parallel effects of insularity and human footprint, several ecological and geographic variables affected introduction probability and extinction risk in opposite directions. Our analysis thus shows that while winners and losers of the Anthropocene can inhabit the same geographic areas, their persistence is largely determined by their native range size, range shape, and ecological characteristics that lie on opposite ends of the same trait continua.

Phylogeographic signature of introduced and extinction-prone species

We detected significant phylogenetic signal for both introduced and extinction-prone species groups, indicating that a species' fate in the Anthropocene is partially related to its phylogenetic history. This phylogenetic signal is likely due to the functional traits that clades have evolved and where clades diversified geographically. Previous studies demonstrated that native ranges of exotic fish (Alcaraz et al. 2005) and birds (Blackburn and Duncan 2001) span a wide latitudinal range, and the source distributions of exotic birds are skewed towards northern latitudes (Blackburn and Duncan 2001). However, we found a strong link between insularity of species ranges and introduction status, which to our knowledge has not been established before this study. Reasons for high introduction potential among insular reptiles can be evolutionary, in that insular species are selected to be naturally good dispersers and strong competitors for the limited resources on

islands. Evidence for this has been found in Caribbean tree lizards of genus *Anolis* for which a so-

called "colonizer body type" has been identified: a set of morphological traits shared by species

that recently colonized the islands through human agency and congeners that naturally colonized

the islands millions of years ago (Poe et al. 2011). Alternatively, the positive association between

insularity and introduction status can also have a contemporary, purely ecological reason, as the

high human impact levels on islands increase the probability of being exported elsewhere (Latella

et al. 2011, Liu et al. 2014).

The association between island-living and extinction-prone status has already received

much attention. For instance, all island regions of the world are considered "biodiversity hotspots",

characterized by simultaneous high levels of endemism and extinction risk (Myers et al. 2000,

Mittermeier et al. 2012). In addition, islands harbor 58% of all critically endangered reptiles (Spatz

et al. 2017) and most of the recorded species extinctions occurred on oceanic islands (Sax and

Gaines 2008). Potential causes of this correlation between insularity and extinction risk can be that

the many endemic species with range-restricted populations are prone to random stochastic

extinction events (Fordham and Brook 2010, Smith and Almeida 2020) or insular species may be

innately predisposed to fast rates of species turnover (i.e., diversification and extinction). The flora

of the South African biodiversity hotspot provides evidence for such predisposition, as young,

recently speciated lineages are particularly at risk of extinction, irrespective of human stress factors

(Davies et al. 2011); however, we did not find that to be the case for western hemisphere squamates.

Hence the location of a species' native range and how heavily it is impacted by humans is a key

determinant of species fate in the Anthropocene.

Introduced and extinction-prone species characteristics

Like Kier et al. (2009) who looked at islands worldwide, our human footprint analysis showed that

insular species are subjected to disproportionate levels of human impact. However, we did not

detect an association with the number of ports in the native range as a predictor for exotic

introductions, while travel movements are a central condition for species introductions (Mahoney

et al. 2015). This could be because aerial travel was not taken into account in the number of ports

and likely better reflected by human footprint, because air travel volume strongly correlates with

population size and infrastructure (Mao et al. 2015).

We found that introduced species generally have large and complex ranges, indicative of

an uneven spread across the landscape. However, we did not find indications of exceptional

adaptive climatic capacity among introduced species compared to resident native species; a trait

that has previously been associated with historic range expansion of western hemisphere passerines

(Title and Burns 2015) and rapid climatic niche shift in amphibians and reptiles (Wiens et al. 2019).

A potential reason for the high levels of native range complexity (i.e., deviation from a circular

range) of introduced species might be that these species follow human impact, which is

characterized by a network of irregularly shaped roads and dense settlements (Herold et al. 2003)

or that large ranges are more likely to encounter erratic geographic boundaries such as coastlines

and mountain ranges. In contrast, extinction-prone species have significantly smaller and simpler

range shapes than non-threatened species, often being confined to single point populations. This

finding serves as a positive control of our analysis, as range size is one of the main criteria for

IUCN to consider a species endangered with extinction (IUCN Standards and Petitions

Subcommittee 2019).

Herbivorous and omnivorous diets are overrepresented among introduced species, while

carnivory is predominant among extinction-prone species. These results contrast a recent global

study across several taxa including reptiles that indicates herbivores to be most at risk of extinction

(Atwood et al. 2020). The positive association between herbivory and introduction success are

congruent with a modeling study that shows a lower trophic niche level and generalist herbivorous

diet to be the most important predictors of introduction success (Romanuk et al. 2009), and with a

global study on reptiles that identifies herbivores as most likely to establish exotic populations

(Mahoney et al. 2015). However, we expected a positive association between introduction status

and carnivory rather than herbivory as most well-known reptile introductions feature carnivorous

species, notorious for their impact on native prey populations (e.g., *Boiga irregularis, Boa*

constrictor) and responsible for most exotic-facilitated species extinctions (Sax and Gaines 2008).

Our analysis shows that such introductions do not make up the majority of reptile introductions, at

least for those species that originate in the western hemisphere.

Introduced species exhibit a single positive association with clutch size, a trait that is

associated with fast life-history "r strategies" (i.e., species adhering to r strategies are highly fecund

and reach sexual maturity early). Having "fast traits" greatly benefits species establishment and

invasive spread in reptiles as these traits facilitate rapid population growth from a small population

size and buffer against vulnerability to stochastic events (Mahoney et al. 2015, Allen et al. 2017).

In addition, humans may also favor species with large clutches to breed for the pet trade, a major

introduction pathway for exotic reptiles (Stringham & Lockwood 2018).

We detected a single, negative association between extinction-prone species and maximum

body size, contradicting earlier studies that indicated large-bodied reptiles as most threatened, and

introduced reptiles as generally small (Mahoney et al. 2015, Tingley et al. 2016). However, these

studies included crocodilians and tortoises in their datasets, which are relatively large-bodied

compared to squamates (snakes and lizards). A possible reason for the negative association between

body size and species endangerment in our study is that small reptiles may be particularly sensitive

to temperature shifts (Herczeg et al. 2007) making them vulnerable to human-caused microhabitat

and climate change (Jesse et al. 2018).

Other possibly important variables for species introduction probability include behavioral

traits that correspond to cognitive function. Brain size (Amiel et al. 2011) and behavioral plasticity

(Perry et al. 1999; Salazar et al. 2019) have been associated with introduction success, but could

not be taken into account in this large-scale and multi-species analysis. Also, reproductive traits,

such as the ability of parthenogenesis and long-term sperm storage, could greatly facilitate species

establishment in an exotic range (Mahoney et al. 2015), as theoretically only a single pregnant

female can establish a viable exotic population.

<u>Opposite ends of the same trait continua</u>

Introduced and extinction-prone reptiles showed opposite species characteristics for seven out of

fourteen (50%) of our model variables. Other studies also report opposite effect sizes for 50-73%

of species-level predictors in animal taxa (Jetschke & Strayer 2008, Blackburn & Jetschke 2009,

Tingley et al. 2016). The support for the opposite ends hypothesis seems to find even stronger

support in plant studies (Bradshaw et al. 2008, Pandit et al. 2011), although due to their different

methodology, the strength of support for the hypothesis is difficult to compare among studies. In

our study, there were strong positive associations of both introduced and extinction-prone species

with insularity and human foot print index. Therefore, our results imply that (dis)similarities in

species characteristics alone do not provide a full understanding of what makes a species introduced

or threatened. To answer this question, we may need to adopt a tiered model, in which the level of

human activities in the native area of a species determines the likelihood of distributional change

in the species' range, while its eco-evolutionary and geographic characteristics then determine if

the species will expand or contract in range.

<u>Implications for (insular) biodiversity</u>

Islands have always served as important stepping stones for overwater dispersal and locations from

which many colonizer species originated (MacArthur & Wilson, 1967, Harbaugh et al. 2009).

However, colonization and extinction rates greatly exceed historic background rates of species loss

and gain (Ricciardi 2007, Loehle & Eschenbach, 2012). For instance, in Hawaii, the colonization

rate before human settlement has been estimated at 0.03 species per 1000 years, increasing to 20

species per 1000 years with the arrival of Polynesians and 20000 species per 1000 years over the

last two centuries (Smith 1985, Loope et al. 1988, Ricciardi 2007). While oceanic islands are

generally more susceptible to exotic invasion than mainland regions (Dawson et al. 2017), the level

of colonization pressure on Hawaii in the last centuries is likely an extreme example.

The current rate of species introductions currently surpasses the rate of species extinctions

(Ellis et al. 2012). In our dataset, only eight of 483 extinction-prone species were already

considered extinct, while a total of 21% of species (483 out of 2268) were threatened with

extinction based on IUCN status. This threat level approximates results for all vertebrates, of which

18% are at risk of extinction (Atwood et al. 2020) and 32% of terrestrial vertebrates experience

range reduction or population decline (also including low concern species) (Ceballos et al. 2017).

The discrepancy between extinct and threatened species numbers indicates the existence of an

extinction debt: the surplus of species that will go extinct in the foreseeable future, but still

contributes to the species richness we see today.

The swift influx of colonizers and prospective loss of endemic species leads to significant

taxonomic and functional homogenization (Olden et al. 2006, Clavel et al. 2010, Capinha et al.

2015), especially within island biota (Longman et al. 2018). This threatens biodiversity and

ecosystem functioning on a small scale, but also poses a threat to global biodiversity because of

the profound risk to lose a large portion of diversity on earth. To mitigate this process, this study

provides applicable knowledge that can be implemented to prevent species introductions and

extinctions via screening and biosecurity measures as well as targeted protection policies.

Acknowledgement

We thank the Netherlands Organization for Scientific Research (grant number: 858.14.041) and

the National Science Foundation program "Dimensions of Biodiversity" (DEB-1136586) for

financial support. We thank students and staff from the iEco and Hedges labs at Temple University

in Philadelphia for their hospitality and helping hand (especially Camille Lynch and Tyler Tran)

during a very inspiring and productive visit to conceptualize this chapter. We thank Catherine H.

Graham for a friendly review of the work.

References

Adeoba, M., S. G. Tesfamichael, and K. Yessoufou. 2019. Preserving the tree of life of the fish

family Cyprinidae in Africa in the face of the ongoing extinction crisis. Genome 62:170–

182.

Alcaraz, C., A. Vila-Gispert, and E. García-Berthou. 2005. Profiling invasive fish species: The

importance of phylogeny and human use. Diversity and Distributions 11:289–298.

Allen, W. L., S. E. Street, and I. Capellini. 2017. Fast life history traits promote invasion success

in amphibians and reptiles.

Alroy, J. 2015. Current extinction rates of reptiles and amphibians. Proceedings of the National

Academy of Sciences of the United States of America 112:13003–13008.

Amiel, J. J., R. Tingley, and R. Shine. 2011. Smart moves: Effects of relative Brain size on

Establishment success of invasive amphibians and reptiles. PLoS ONE 6.

Arbetman, M. P., G. Gleiser, C. L. Morales, P. Williams, and M. A. Aizen. 2017. Global decline of bumblebees is phylogenetically structured and inversely related to species range size and pathogen incidence. Proceedings of the Royal Society B: Biological Sciences 284.

Atwood, T. B., S. A. Valentine, E. Hammill, D. J. McCauley, E. M. P. Madin, K. H. Beard, and W. D. Pearse. 2020. Herbivores at the highest risk of extinction among mammals, birds, and reptiles. Science Advances 6.

Blackburn, T. M., and R. P. Duncan. 2001. Establishment patterns of exotic birds are constrained by non-random patterns in introduction. Journal of Biogeography 28.

Blackburn, T. M., and J. M. Jeschke. 2009. Invasion success and threat status: Two sides of a different coin? Ecography 32:83–88.

Böhm, M., B. Collen, J. E. M. Baillie, P. Bowles, J. Chanson, N. Cox, G. Hammerson, M. Hoffmann, S. R. Livingstone, M. Ram, A. G. J. Rhodin, S. N. Stuart, P. P. van Dijk, B. E. Young, L. E. Afuang, A. Aghasyan, A. García, C. Aguilar, R. Ajtic, F. Akarsu, L. R. V. Alencar, A. Allison, N. Ananjeva, S. Anderson, C. Andrén, D. Ariano-Sánchez, J. C. Arredondo, M. Auliya, C. C. Austin, A. Avci, P. J. Baker, A. F. Barreto-Lima, C. L. Barrio-Amorós, D. Basu, M. F. Bates, A. Batistella, A. Bauer, D. Bennett, W. Böhme, D. Broadley, R. Brown, J. Burgess, A. Captain, S. Carreira, M. del R. Castañeda, F. Castro, A. Catenazzi, J. R. Cedeño-Vázquez, D. G. Chapple, M. Cheylan, D. F. Cisneros-Heredia, D. Cogalniceanu, H. Cogger, C. Corti, G. C. Costa, P. J. Couper, T. Courtney, J. Crnobrnja-Isailovic, P. A. Crochet, B. Crother, F. Cruz, J. C. Daltry, R. J. R. Daniels, I. Das, A. de Silva, A. C. Diesmos, L. Dirksen, T. M. Doan, C. K. Dodd, J. S. Doody, M. E. Dorcas, J. Duarte de Barros Filho, V. T. Egan, E. H. El Mouden, D. Embert, R. E. Espinoza, A. Fallabrino, X. Feng, Z. J. Feng, L. Fitzgerald, O. Flores-Villela, F. G. R. França, D. Frost, H. Gadsden, T. Gamble, S. R. Ganesh, M. A. Garcia, J. E. García-Pérez, J. Gatus, M. Gaulke,

587 P. Geniez, A. Georges, J. Gerlach, S. Goldberg, J. C. T. Gonzalez, D. J. Gower, T. Grant, E.

588 Greenbaum, C. Grieco, P. Guo, A. M. Hamilton, K. Hare, S. B. Hedges, N. Heideman, C.

589 Hilton-Taylor, R. Hitchmough, B. Hollingsworth, M. Hutchinson, I. Ineich, J. Iverson, F. M.

590 Jaksic, R. Jenkins, U. Joger, R. Jose, Y. Kaska, U. Kaya, J. S. Keogh, G. Köhler, G.

591 Kuchling, Y. Kumlutaş, A. Kwet, E. La Marca, W. Lamar, A. Lane, B. Lardner, C. Latta, G.

592 Latta, M. Lau, P. Lavin, D. Lawson, M. LeBreton, E. Lehr, D. Limpus, N. Lipczynski, A. S.

593 Lobo, M. A. López-Luna, L. Luiselli, V. Lukoschek, M. Lundberg, P. Lymberakis, R.

594 Macey, W. E. Magnusson, D. L. Mahler, A. Malhotra, J. Mariaux, B. Maritz, O. A. V.

595 Marques, R. Márquez, M. Martins, G. Masterson, J. A. Mateo, R. Mathew, N. Mathews, G.

596 Mayer, J. R. McCranie, G. J. Measey, F. Mendoza-Quijano, M. Menegon, S. Métrailler, D.

597 A. Milton, C. Montgomery, S. A. A. Morato, T. Mott, A. Muñoz-Alonso, J. Murphy, T. Q.

598 Nguyen, G. Nilson, C. Nogueira, H. Núñez, N. Orlov, H. Ota, J. Ottenwalder, T. Papenfuss,

599 S. Pasachnik, P. Passos, O. S. G. Pauwels, N. Pérez-Buitrago, V. Pérez-Mellado, E. R.

600 Pianka, J. Pleguezuelos, C. Pollock, P. Ponce-Campos, R. Powell, F. Pupin, G. E. Quintero

601 Díaz, R. Radder, J. Ramer, A. R. Rasmussen, C. Raxworthy, R. Reynolds, N. Richman, E.

602 L. Rico, E. Riservato, G. Rivas, P. L. B. da Rocha, M. O. Rödel, L. Rodríguez Schettino, W.

603 M. Roosenburg, J. P. Ross, R. Sadek, K. Sanders, G. Santos-Barrera, H. H. Schleich, B. R.

604 Schmidt, A. Schmitz, M. Sharifi, G. Shea, H. T. Shi, R. Shine, R. Sindaco, T. Slimani, R.

605 Somaweera, S. Spawls, P. Stafford, R. Stuebing, S. Sweet, E. Sy, H. J. Temple, M. F.

606 Tognelli, K. Tolley, P. J. Tolson, B. Tuniyev, S. Tuniyev, N. üzüm, G. van Buurt, M. Van

607 Sluys, A. Velasco, M. Vences, M. Veselý, S. Vinke, T. Vinke, G. Vogel, M. Vogrin, R. C.

608 Vogt, O. R. Wearn, Y. L. Werner, M. J. Whiting, T. Wiewandt, J. Wilkinson, B. Wilson, S.

609 Wren, T. Zamin, K. Zhou, and G. Zug. 2013. The conservation status of the world's reptiles.

610 Biological Conservation 157:372–385.

Böhm, M., R. Williams, H. R. Bramhall, K. M. Mcmillan, A. D. Davidson, A. Garcia, L. M.

Bland, J. Bielby, and B. Collen. 2016. Correlates of extinction risk in squamate reptiles: The

relative importance of biology, geography, threat and range size. Global Ecology and

Biogeography 25:391–405.

Bradshaw, C. J. A., X. Giam, H. T. W. Tan, B. W. Brook, and N. S. Sodhi. 2008. Threat or

invasive status in legumes is related to opposite extremes of the same ecological and life-

history attributes. Journal of Ecology 96:869–883.

Cadotte, M. W., M. A. Hamilton, and B. R. Murray. 2009. Phylogenetic relatedness and plant

invader success across two spatial scales. Diversity and Distributions 15.

Capinha, C., F. Essl, H. Seebens, D. Moser, and H. M. Pereira. 2015. The dispersal of alien

species redefines biogeography in the Anthropocene. Science 348:1248–1251.

Ceballos, G., P. R. Ehrlich, and R. Dirzo. 2017. Biological annihilation via the ongoing sixth

mass extinction signaled by vertebrate population losses and declines. Proceedings of the

National Academy of Sciences of the United States of America 114:E6089–E6096.

Clavel, J., R. Julliard, and V. Devictor. 2011. Worldwide decline of specialist species: toward a

global functional homogenization? Frontiers in Ecology and the Environment 9: 222-228.

Colautti, R. I., and H. I. MacIsaac. 2004. A neutral terminology to define "invasive" species.

Diversity and Distributions 10:135–141.

Cowie, R. H., and B. S. Holland. 2006. Dispersal is fundamental to biogeography and the

evolution of biodiversity on oceanic islands.

Cox, E. P. 1927. A Method of Assigning Numerical and Percentage Values to the Degree of

Roundness of Sand Grains. Journal of Paleontology 1.

Davies, T. J., G. F. Smith, D. U. Bellstedt, J. S. Boatwright, B. Bytebier, R. M. Cowling, F.

Forest, L. J. Harmon, A. M. Muasya, B. D. Schrire, Y. Steenkamp, M. van der Bank, and V.

Savolainen. 2011. Extinction risk and diversification are linked in a plant biodiversity

hotspot. PLoS Biology 9.

Dawson, W., Moser, D., Van Kleunen, M., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Winter, M.,

Lenzner, B., Blackburn, T.M. and Dyer, E.E., 2017. Global hotspots and correlates of alien

species richness across taxonomic groups. Nature Ecology & Evolution 1:1-7.

Ellis, E. C., E. C. Antill, and H. Kreft. 2012. All is not loss: plant biodiversity in the

Anthropocene. PloS ONE 7: e30535.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. International Journal of Climatology 37.

Fordham, D. A., and B. W. Brook. 2010. Why tropical island endemics are acutely susceptible to

global change. Biodiversity and Conservation 19.

Fritz, S. A., and A. Purvis. 2010. Selectivity in mammalian extinction risk and threat types: A

new measure of phylogenetic signal strength in binary traits. Conservation Biology

24:1042–1051.

GADM. 2018. Database of Global Administrative Areas, version 3.6. The University of

California.

Harbaugh, D. T., W. L. Wagner, G. J. Allan, and E. A. Zimmer. 2009. The Hawaiian Archipelago

is a stepping stone for dispersal in the Pacific: an example from the plant genus Melicope

(Rutaceae). Journal of Biogeography 36: 230-241.

Hedges, S. B. 2006. An overview of the evolution and conservation of West Indian amphibians

and reptiles. Applied Herpetology 3.

Hedges, S. B. 2011. An overview of the evolution and conservation of West Indian amphibians

and reptiles. Pages 25–37 Conservation of Caribbean Island Herpetofaunas.

Hedges, S.B. 2021. Caribherp: amphibians and reptiles of Caribbean Islands. Available online at

http://www.caribherp.org/ (accessed 1 July 2021). Philadelphia, Pennsylvania: Temple

University.

Helmus, M. R., J. E. Behm, A. M. Wendy, J. J. Kolbe, J. Ellers, and J. B. Losos. 2017. Exotics

Exhibit More Evolutionary History Than Natives : a Comparison of the Ecology and

Evolution of Exotic and Native Anole Lizards. Invasion Genetics: The Baker and Stebbins

Legacy.

Herczeg, G., J. Török, and Z. Korsós. 2007. Size-dependent heating rates determine the spatial

and temporal distribution of small-bodied lizards. Amphibia Reptilia 28:347–356.

Herold, M., X. H. Liu, and K. C. Clarke. 2003. Spatial metrics and image texture for mapping

urban land use.

Hijmans, R. J. 2019. geosphere: Spherical Trigonometry. R package version 1.5-10.

IUCN Standards and Petitions Subcommittee. 2019. Guidelines for Using the IUCN Red List

Categories and Criteria. IUCN 14.

IUCN Redlist. 2021. IUCN Redlist of Threatened Species. Available from

http://www.iucnredlist.org/ (accessed 1 July 2021). Gland, Switzerland: International Union

for the Conservation of Nature.

Jantz, S. M., B. Barker, T. M. Brooks, L. P. Chini, Q. Huang, R. M. Moore, J. Noel, and G. C.

Hurtt. 2015. Future habitat loss and extinctions driven by land-use change in biodiversity

hotspots under four scenarios of climate-change mitigation. Conservation Biology 29:1122–

1131.

Jeschke, J. M., and D. L. Strayer. 2008. Are threat status and invasion success two sides of the

same coin? Ecography 31:124–130.

Jesse, W. A. M., J. E. Behm, M. R. Helmus, and J. Ellers. 2018. Human land use promotes the

abundance and diversity of exotic species on Caribbean islands. Global Change Biology

24:4784–4796.

Kier, G., H. Kreft, T. M. Lee, W. Jetz, P. L. Ibisch, C. Nowicki, J. Mutke. and W. Barthlott.

2009. A global assessment of endemism and species richness across island and mainland

regions. Proceedings of the National Academy of Sciences, 106: 9322-9327.

Kotiaho, J. S., V. Kaitala, A. Komonen, and J. Päivinen. 2005. Predicting the risk of extinction

from shared ecological characteristics. Proceedings of the National Academy of Sciences of

the United States of America 102:1963–1967.

Kraus, F. 2009. Appendix A : Database of Introductions. Alien Reptiles and Amphibians: A

Scientific Compendium and Analysis:133–369.

Krysko, K. L., L. A. Somma, D. C. Smith, C. R. Gillette, D. Cueva, J. A. Wasilewski, K. M.

Enge, S. A. Johnson, T. S. Campbell, J. R. Edwards, M. R. Rochford, R. Tompkins, J. L.

Fobb, S. Mullin, C. Lechowicz, D. Hazelton, and A. Warren. 2016. New verified

nonindigenous amphibians and reptiles in Florida, 1976 through 2015, with a summary of

over 152 years of introductions. Reptiles & Amphibians 23.

Kumar, S., G. Stecher, M. Suleski, and S. B. Hedges. 2017. TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. Molecular biology and evolution 34.

Larson, E. R., and J. D. Olden. 2010. Latent extinction and invasion risk of crayfishes in the

southeastern United States. Conservation Biology 24:1099–1110.

Latella, I. M., S. Poe, and J. T. Giermakowski. 2011. Traits associated with naturalization in

Anolis lizards: Comparison of morphological, distributional, anthropogenic, and

phylogenetic models. Biological Invasions 13:845–856.

Liu, X., X. Li, Z. Liu, R. Tingley, F. Kraus, Z. Guo, and Y. Li. 2014. Congener diversity,

topographic heterogeneity and human-assisted dispersal predict spread rates of alien

herpetofauna at a global scale. Ecology Letters 17:821–829.

Loehle, C. and W. Eschenbach. 2012. Historical bird and terrestrial mammal extinction rates and causes. Diversity and Distributions 18: 84-91.

Longman, E. K., K. Rosenblad, and D. F. Sax. 2018. Extreme homogenization: The past, present and future of mammal assemblages on islands. Global Ecology and Biogeography 27:77–95.

Loope, L. L., O. Hamann, and C. P. Stone. 1988. Comparative Conservation Biology of Oceanic Archipelagoes: Hawaii and the Galápagos. BioScience 38:272–282.

Losos, J. B. 2009. Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles. Page Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles.

Loza, M. I., I. Jiménez, P. M. Jørgensen, G. Arellano, M. J. Macía, V. W. Torrez, and R. E. Ricklefs. 2017. Phylogenetic patterns of rarity in a regional species pool of tropical woody plants. Global Ecology and Biogeography 26.

Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the macroevolutionary landscape in island lizard radiations. Science 341:292–295.

Mahler, D. L., L. J. Revell, R. E. Glor, and J. B. Losos. 2010. Ecological opportunity and the rate of morphological evolution in the diversification of greater Antillean anoles. Evolution 64:2731–2745.

Mahoney, P. J., K. H. Beard, A. M. Durso, A. G. Tallian, A. L. Long, R. J. Kindermann, N. E. Nolan, D. Kinka, and H. E. Mohn. 2015. Introduction effort, climate matching and species traits as predictors of global establishment success in non-native reptiles. Diversity and Distributions 21:64–74.

Mao, L., X. Wu, Z. Huang, and A. J. Tatem. 2015. Modeling monthly flows of global air travel passengers: An open-access data resource. Journal of Transport Geography 48.

MacArthur, R. H. and E.O. Wilson. 1967. Island biogeography. Princeton, USA.

Marin, J., G. Rapacciuolo, G. C. Costa, C. H. Graham, T. M. Brooks, B. E. Young, V. C.

Radeloff, J. E. Behm, M. R. Helmus, and S. B. Hedges. 2018. Evolutionary time drives

global tetrapod diversity. Proceedings of the Royal Society B: Biological Sciences 285.

Mittermeier, R. A., W. R. Turner, F. W. Larsen, T. M. Brooks, and C. Gascon. 2011. Global

Biodiversity Conservation: The Critical Role of Hotspots. Pages 3–22 Biodiversity

Hotspots.

Monaco, C. J., C. J. A. Bradshaw, D. J. Booth, B. M. Gillanders, D. S. Schoeman, and I.

Nagelkerken. 2020. Dietary generalism accelerates arrival and persistence of coral-reef

fishes in their novel ranges under climate change. Global Change Biology 26.

Myers, N., R. A. Mittermeler, C. G. Mittermeler, G. A. B. Da Fonseca, and J. Kent. 2000.

Biodiversity hotspots for conservation priorities. Nature 403:853–858.

National Geospatial-Intelligence Agency. 2017. World Port Index.

https://msi.nga.mil/Publications/WPI.

Olden, J. D., N. L. Poff, and M. L. McKinney. 2006. Forecasting faunal and floral

homogenization associated with human population geography in North America. Biological

Conservation 127: 261–271.

Pandit, M. K., M. J. O. Pocock, and W. E. Kunin. 2011. Ploidy influences rarity and invasiveness

in plants. Journal of Ecology 99.

Park, D. S., and D. Potter. 2015. Why close relatives make bad neighbours: Phylogenetic

conservatism in niche preferences and dispersal disproves Darwin's naturalization

hypothesis in the thistle tribe. Molecular Ecology 24:3181–3193.

Perry, G., Razi Dmi'el, & Lazell, J. 1999. Evaporative Water Loss in Insular Populations of the

Anolis cristatellus Group (Reptilia: Sauria) in the British Virgin Islands II: The Effects of

Drought. Biotropica 31: 337–343.

Pigot, A. L., E. E. Dyer, D. W. Redding, P. Cassey, G. H. Thomas, and T. M. Blackburn. 2018.

Species invasions and the phylogenetic signal in geographical range size.

Poe, S., J. T. Giermakowski, I. Latella, E. W. Schaad, E. P. Hulebak, and M. J. Ryan. 2011.

Ancient colonization predicts recent naturalization in anolis lizards. Evolution 65:1195–

1202.

Polsby, D. D., and R. Popper. 2017. The Third Criterion: Compactness as a Procedural Safeguard

Against Partisan Gerrymandering. SSRN Electronic Journal.

Powell, R., R. W. Henderson, and (Eds.). 2012. Island lists of West Indian amphibians and

reptiles. Bulletin of the Florida Museum of Natural History 51:85–166.

Powell, R., R. W. Henderson, M. C. Farmer, M. Breuil, A. C. Echternacht, G. van Buurt, C. M.

Romagosa, and G. Perry. 2011. Introduced amphibians and reptiles in the Greate Caribbean:

Patterns and conservation implications. Page Conservation of Caribbean Island

Herpetofaunas.

Pyšek, P., J. Pergl, F. Essl, B. Lenzner, W. Dawson, H. Kreft, P. Weigelt, M. Winter, J. Kartesz,

M. Nishino, L. A. Antonova, J. F. Barcelona, F. J. Cabezas, D. Cárdenas, J. Cárdenas-Toro,

N. Castaño, E. Chacón, C. Chatelain, S. Dullinger, A. L. Ebel, E. Figueiredo, N. Fuentes, P.

Genovesi, Q. J. Groom, L. Henderson, Inderjit, A. Kupriyanov, S. Masciadri, N. Maurel, J.

Meerman, O. Morozova, D. Moser, D. Nickrent, P. M. Nowak, S. Pagad, A. Patzelt, P. B.

Pelser, H. Seebens, W. S. Shu, J. Thomas, M. Velayos, E. Weber, J. J. Wieringa, M. P.

Baptiste, and M. Van Kleunen. 2017. Naturalized alien flora of the world: Species diversity,

taxonomic and phylogenetic patterns, geographic distribution and global hotspots of plant

invasion. Preslia 89.

R Development Core Team, R. 2021. R: A Language and Environment for Statistical Computing.

R Foundation for Statistical Computing.

R Studio Team. 2021. RStudio: Integrated Development for R. PBC, Boston, MA.

779 Rapacciuolo, G., C. H. Graham, J. Marin, J. E. Behm, G. C. Costa, S. B. Hedges, M. R. Helmus,

780   V. C. Radeloff, B. E. Young, and T. M. Brooks. 2019. Species diversity as a surrogate for

781   conservation of phylogenetic and functional diversity in terrestrial vertebrates across the

782   Americas. Nature Ecology and Evolution 3:53–61.

783 Ricciardi, A. 2007. Are modern biological invasions an unprecedented form of global change?

784 Romanuk, T. N., Y. Zhou, U. Brose, E. L. Berlow, R. J. Williams, and N. D. Martinez. 2009.

785   Predicting invasion success in complex ecological networks. Philosophical Transactions of

786   the Royal Society B: Biological Sciences 364.

787 Salazar JC, Del Rosario Castañeda M, Londoño GA, Bodensteiner BL, Muñoz MM. 2019.

788   Physiological evolution during adaptive radiation: A test of the island effect in Anolis

789   lizards. Evolution 73:1241-1252.

790 Sax, D. F., and S. D. Gaines. 2008. Species invasions and extinction: The future of native

791   biodiversity on islands. Proceedings of the National Academy of Sciences of the United

792   States of America 105:11490–11497.

793 Schmidt, J. P., P. R. Stephens, and J. M. Drake. 2012. Two sides of the same coin Rare and pest

794   plants native to the United States and Canada. Ecological Applications 22:1512–1525.

795 Smith, C. W. 1985. Impact of alien plants on Hawa'i's native biota. Hawai'i's Terrestrial

796   Ecosystems. Proc. Symposium, 1984, Hawai'i.

797 Smith, K. G., and R. J. Almeida. 2020. When are extinctions simply bad luck? Rarefaction as a

798   framework for disentangling selective and stochastic extinctions. Journal of Applied

799   Ecology 57.

800 Spatz, D. R., K. M. Zilliacus, N. D. Holmes, S. H. M. Butchart, P. Genovesi, G. Ceballos, B. R.

801   Tershy, and D. A. Croll. 2017. Globally threatened vertebrates on islands with invasive

802   species. Science Advances 3.

Stringham, O. C. and J. L. Lockwood. 2018). Pet problems: biological and economic factors that influence the release of alien reptiles and amphibians by pet owners. Journal of Applied Ecology 55: 2632-2640.

Su, S., P. Cassey, and T. M. Blackburn. 2016. The wildlife pet trade as a driver of introduction
and establishment in alien birds in Taiwan. Biological Invasions 18.

Tingley, R., P. J. Mahoney, A. M. Durso, A. G. Tallian, A. Morán-Ordóñez, and K. H. Beard.
2016. Threatened and invasive reptiles are not two sides of the same coin. Global Ecology
and Biogeography 25.

Title, P. O., and J. B. Bemmels. 2018. ENVIREM: an expanded set of bioclimatic and
topographic variables increases flexibility and improves performance of ecological niche
modeling. Ecography 41.

Title, P. O., and K. J. Burns. 2015. Rates of climatic niche evolution are correlated with species
richness in a large and ecologically diverse radiation of songbirds. Ecology Letters 18:433–
440.

Uetz, P., Freed, P. & Hošek, J. (eds. . 2012. The Reptile Database.

Venter, O., E. W. Sanderson, A. Magrach, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham,
W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. M. Watson. 2016. Global
terrestrial Human Footprint maps for 1993 and 2009. Scientific Data 3.

Vitousek, P. M., H. A. Mooney, J. Lubchenco, and J. M. Melillo. 1997. Human domination of
Earth's ecosystems. Science 277:494–499.

Wiens, J. J., Y. Litvinenko, L. Harris, and T. Jezkova. 2019. Rapid niche shifts in introduced
species can be a million times faster than changes among native species and ten times faster
than climate change. Journal of Biogeography 46.

Yessoufou, K., K. Mearns, H. O. Elansary, and G. H. Stoffberg. 2016. Assessing the

phylogenetic dimension of Australian Acacia species introduced outside their native ranges.

Botany Letters 163:33–39.

Young, H. S., D. J. McCauley, M. Galetti, and R. Dirzo. 2016. Patterns, Causes, and

Consequences of Anthropocene Defaunation. Annual Review of Ecology, Evolution, and

Systematics 47:333–358.

829