The extinction of threatened species by anthropogenic threats endangers the

functional diversity of turtles and crocodiles

Rodríguez-Caro, R.C.[1,2*], Graciá, E.[2], Blomberg, S. P.[3], Cayuela, H.[4], Grace, M.[1],

Carmona, C.P.[5], Pérez-Mendoza, H.A.[6], Giménez, A.[2], Salguero-Gómez, R.[1,7,*]

[1] Department of Zoology, Oxford University, 11a Mansfield Road, OX1 3SZ, Oxford, UK.

[2] Departamento de Biología Aplicada, Universidad Miguel Hernández, Elche, Alicante, Spain.

[3] School of Biological Sciences, The University of Queensland, Brisbane, QLD, Australia.

[4] Laboratoire de Biométrie et Biologie Evolutive, Université Lyon 1, CNRS, UMR 5558, Villeurbanne F-

769622, France

[5] Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

[6] Facultad de Estudios Superiores Iztacala, Universidad Autónoma de México, Tlalnepantla, México.

[7] Max Plank Institute for Demographic Research, Konrad-Zuße straße, Rostock, Germany.

* Corresponding authors: r.rodriguez@umh.es

17   rob.salguero@zoo.ox.ac.uk

Abstract: 200 words

Body text (intro, results, and discussion): 2719 words

Methods: 2890 words

Figure: 4

Tables: 0

ABSTRACT (200/200 words)

The Anthropocene is tightly associated with a drastic loss of species worldwide, and

consequently, with the disappearance of key ecosystem functions. The on-going reduction

in species functionality is driven by global and local threats. The orders Testudines

(turtles and tortoises) and Crocodilia (crocodilians and alligators) contain numerous

threatened, long-lived species for which their functional diversity and potential erosion

by anthropogenic impacts remains unknown. Here, we examine 259 of the existing 375

species of Testudines and Crocodilia quantifying their life history strategies (*i.e.*,

schedules of survival, development, and reproduction) from open-access data on their

demography, ancestry, and threats. We find that the loss of functional diversity in

simulated extinction scenarios of threatened species is greater than expected by chance.

Moreover, the effects of unsustainable local consumption, diseases, and pollution are

associated to specific functional strategies. Meanwhile climate change, habitat

disturbance, and global trade affect all species independent of their life history strategy.

Importantly, the loss of functional diversity for threatened species by habitat disturbance

is twice greater than for all other threats. Our findings highlight the key importance of

conservation programmes focused on preserving the functional diversity of life history

strategies jointly with the phylogenetic representativity of these highly threatened groups.

Keywords: animal disease, biodiversity, comparative demography, climate change,

habitat disturbance, life history trait, phylogenetic analyses, pollution, wildlife trade.

INTRODUCTION

Human impacts, such as habitat loss, climate change, pollution, poaching, and unsustainable trade, impose major threats to the persistence of species worldwide (Maxwell et al., 2016). Modern human activities have drastically accelerated species extinctions by three orders of magnitude (Pimm et al., 1995). As such, effective conservation measures are urgently needed to bend the curve of biodiversity loss (Díaz et al., 2019).

The on-going biodiversity loss is resulting in the erosion of species functionality, key to ecological function (Oliver et al., 2015; Carmona et al., 2021; Toussaint et al., 2021). A robust way to characterise species functionality is via quantifying life history strategies, *i.e*. trade-offs between an individual's investment in survival, development, and reproduction driven by limited resources and physiological constraints (Stearns, 1999; Salguero-Gómez et al., 2016a; Cernansky, 2017; Healy et al., 2019; Capdevila et al., 2020). In this context, different species can possess similar life history trait values, resulting in functional redundancy (Naeem, 1998; *e.g*., Texas tortoises, *Gopherus berlandieri,* and Common snapping turtle, *Chelydra serpentina,* achieve similar longevities; Germano, 1994), or display dissimilar functionality, giving rise to functional dispersion (Elmqvist et al., 2003; *e.g*. loggerhead sea turtle, *Caretta caretta,* reproduces later than other Testudines of similar size, such as Aldabra giant tortoise, *Aldabrachelys gigantea*; Scott et al., 2012). Thus, understanding how redundant or dispersed life history traits are and how sensitive these are to anthropogenic threats worldwide is critical for conservation (Chichorro et al. 2019).

Evaluating the risk of extinction of long-lived species is important for species functionality. The extinction of long-lived species can be delayed due to their long generation times (Gaillard et al., 1989; Jackson and Sax, 2010; Salguero-Gómez et al.,

2016a; Capdevila et al. 2020). Thus, environmental effects typically manifest at much longer time scales in them than in short-lived species (Gibbs and Amato, 2000; Kuussaari et al., 2009). In this respect, two of the most threatened and long-lived groups are the order Testudines (tortoises, turtles, and sea turtles) and Crocodilia (crocodilians and alligators) (Rhodin et al., 2018; Colston et al., 2020). In fact, Testudines show one of the highest proportions of threatened species across vertebrates (65%; Rhodin et al., 2018; UICN, 2020). Importantly, the conservation status of Testudines and Crocodilia could be even worse because 25% of them lack data to accurately identify their IUCN status (Rhodin et al., 2018).

Here, we examine the life history strategies of 69% of Testudines and Crocodilia species –259 out of 375 extant species– to identify how the impacts of different human-led global (such as climate change, global wildlife trade and emergent diseases) and local threats (habitat disturbances, local overconsumption, and pollution) may alter their functional diversity. Using open-access data on their demography, phylogenetic relationships, and threats, we quantify life history strategies to estimate their functional diversity. We then simulate different extinction scenarios according to their extinction risk and the main threats that affect each species. We hypothesise that: (H1) life history strategies across testudines and crocodiles will show high functional redundancy because most of them are long-lived species with delayed maturity (Healy et al., 2019). As such, the loss of functional diversity due to the extinction of threatened species will be lower than expected by chance; (H2) human threats (*e.g.,* unsustainable global trade, unsustainable local consumption, disease) will differentially erode the functional diversity of these species (Ripple et al., 2017). Tortoises and turtles are mostly declining by habitat disturbances, local overconsumption, and collection for the international pet trade (Stanford et al., 2020), so global threats like climate change are expected to have

less impact in the functional diversity than other local threats such as habitat disturbances

because knowledge about the consequences of climate change in long-lived species is still

emerging.

RESULTS

High functional redundancy

Two dominant axes of life history traits describe most of the functional diversity of

Testudines and Crocodilia. Using life history trait data from 236 species of turtles and

tortoises (67% of the extant species), and 23 species of crocodilians and alligators (85%)

from multiple sources we characterise their life history strategies via a phylogenetic

principal component analysis (*p*PCA; Table S1) where adult body mass is accounted for

(Table S2). To address gaps in the dataset (See Appendix 1), we performed a multiple

phylogenetic-trait imputation for missing values (see Materials and Methods). The two

dominant axes of life history strategies encompass 62% of the total functional variation

(Fig. 1). These axes correspond to (i) the 'fast–slow continuum' (Sæther, 1987; Stearns,

1999; Oli, 2004; Bielby et al 2007), and it separates species with short maximum lifespan,

like some freshwater turtles (*e.g.*, 4.3 years for the Alabama red-bellied turtle, *Pseudemys*

alabamensis), to species with high maximum lifespan such as terrestrial tortoises (*e.g.*

177 years for the Pinzon Island giant tortoise, *Chelonoidis ephippium*); and (ii) the

'reproductive strategy' (Gaillard et al., 1989; Salguero-Gómez et al., 2016a; Hughes,

2017; Paniw et al. 2018; Capdevila et al. 2020), defined by a trade-off between clutch

frequency *vs.* reproductive output, with species with high number of clutches per year but

lower clutch size per clutch at one end (*e.g.*, 1.2 eggs per clutch and 3 clutches per year

as in the Madagascan flat-tailed tortoise, *Pyxis planicauda*), and extremely reproductive

species at the other end, having high clutch sizes but few clutches per year (*e.g.,* 91.5 eggs

per clutch in 1 clutches per year as in the South American River Turtle, *Podocnemis*

expansa).

Examining the functional spectra of the life history traits of testudines and

crocodilians reveals a high degree of functional redundancy. We used a trait probability

density (TPD) approach considering the two-dimensional functional space (Fig. 1) to

identify areas with hotspots and valleys of species diversity (Carmona et al., 2016;

Toussaint et al., 2021). The examined reptile species tend to congregate around a single

hotspot with median values of maximum lifespan and high number of clutches per year

with lower clutch sizes. This hotspot of life history strategies includes 50% of the 259

species and represents 13.2% of the total spectrum (Fig. 1). Freshwater turtles such as the

eastern long-necked turtle, *Chelodina longicollis,* (max. lifespan 37 years, 13.9

eggs/clutch, 2 clutches/yr) and crocodiles such as the mugger crocodile *Crocodylus*

palustris (max. lifespan 31.5 years, 28.75 eggs/clutch, 2 clutches/yr) are represented in

that hotspot of functional diversity. Sea turtles are represented in a valley of functional

diversity (bottom right of Fig. 1), which is characterised by species with higher clutch

sizes, few clutches per year, and higher maximum lifespan (*e.g. Caretta caretta:* max.

lifespan 76 years, 115 eggs/clutch, 1.29 clutches/yr). In contrast, terrestrial tortoises are

represented at the top right of Fig. 1, with their typically shorter clutch sizes, high clutches

per year, and longer maximum lifespans (*e.g.*, the Hood Island giant tortoise *Chelonoidis*

hoodensis: max. lifespan 177 years, 10 eggs/clutch, 3 clutches/yr).

Unique strategies may become extinct

The extinction of threatened species would result in the loss of a quarter of functional

diversity of testudines and crocodiles. To estimate the effect of the potential extinctions

of threatened species on the existing amount of functional diversity in these groups, we

used the IUCN Red List categories of our species and simulated an accumulative loss of

threatened species (*i.e.*, in the first scenario "-CR", we removed the *Critically*

Endangered species only; in the "-EN" scenario, we removed Endangered and Critically

Endangered species, and so on; see Materials and Methods). If all threatened species

disappeared (*Critically Endangered*, *Endangered,* and *Vulnerable* species*;* "-VU"

scenario, Fig. 2A), the functional diversity would decrease by 26.8%.

The loss of functional diversity due to the extinction of threatened species is

greater than expected by chance. We compared the potential loss of functional diversity,

considering the number of threatened species in each extinction scenario, with a simulated

loss of functional diversity where the identity of the threatened species was randomised

within the pool of species. The resulting loss of functional diversity due to the extinction

of threatened species is greater than 95% of randomisations for most scenarios (Fig. 2A).

In the "-CR" scenario, the loss of functional diversity (-12.67%) is four times greater than

in randomisations (-3.82% [-0.4%, -9.69%] 5% and 95% CI). This finding suggests that

most of the Critically Endangered species of testudines and crocodilians show unique life

history strategies, such as the pancake tortoise (*Malacochersus tornieri* max. lifespan 25.9

years, 1 eggs/clutch, 1 clutches/yr).

Unique or less redundant species strategies of testudines and crocodilians are more

vulnerable to vanish following the extinction of threatened species. To identify

differences in extinction risk between the life history strategies, we mapped extinction

risk in the functional trait spectra via generalized additive models (GAMs), using the

IUCN Red List category of each species (Carmona et al. 2021). The position of each

species along the trait spectra is significantly correlated with its extinction risk when

considering high-risk species as Critically Endangered and Endangered (p = 0.039; Fig.

2B). Using this grouping, high-risk species show distinct positions in the life history

functional space, with species with lower functional redundancy located around the

periphery and likely being more threatened. Only species located at the centre of the

functional space and those with short maximum lifespan and smaller clutch sizes but with

greater number of clutches per year show a lower probability of being threatened (Fig.

2B).

Habitat disturbance is the main threat

The life history strategies of Testudines and Crocodilia can predict vulnerability to

extinction in three of the six analysed threats (local consumption, disease, and pollution;

Fig. 3). To explore the differential effects of global and local threats on the functional

spectra of testudines and crocodilians, we analysed the probability of being affected by

each threat separately using GAMs (more results in Fig. S1). Local consumption (Fig.

3A) is associated with increased extinction risk in species with higher clutch size such as

sea turtles (*e.g., Caretta caretta*), and higher maximum lifespans, such as the radiated

tortoise (*Astrochelys radiata*) and the Orinoco crocodile (*Crocodylus intermedius*). In

contrast, diseases are associated to threaten species with slow life histories (Fig. 3B), such

as the desert tortoise (*Gopherus agassizii*). Finally, pollution primarily is associated to

species with higher reproductive output (*i.e.,* higher clutch sizes), such as freshwater

turtles like *Chelydra serpentina* or crocodilians like the saltwater crocodile, *Crocodylus*

porosus, (Fig. 3C).

Among all examined threats, the extinction of species affected by habitat

disturbance results in the worst loss of functional diversity, especially in the northern

hemisphere (Fig 4A). To estimate the effect of specific threats on the loss of functional

diversity, we simulated the extinction of species affected by each threat to calculate the

percentage of loss of functional diversity (Fig. 4B). We compared the potential loss of

functional diversity by each threat, with a simulated scenario where the identity of the

species affected by each threat was randomised within the pool of species. Most of the

effects of loss of functional diversity due to the six threats considered are greater than

expected by random extinctions, especially for climate change, whose effect is beyond

the 95% percentile for random simulations (Fig. 4B). Only habitat disturbance and

diseases show similar or lower values of loss of functional diversity than expected by

random simulations (Fig 4B).

DISCUSSION

Here, we have shown that threatened species of turtles and crocodilians show great deal

of functional dispersion. Toussaint et al. (2021), exploring 5,686 species of reptiles, found

a loss of functional diversity linked to the extinction of the threatened species

corresponding to less than 10%. Here, focussing on Testudines and Crocodilia, we find

twice the loss of functional diversity. By linking this information to their conservation

status, we demonstrate that global and local threats are exposing the functional diversity

of these already highly threatened taxonomic groups to great peril. Using the largest

collection of trait data to date for turtles and crocodiles, we find that the threatened species

show unique life history strategies. If we lost these species due to extinction, this would

translate into a greater reduction of functional diversity and potentially severe ecological

consequences (Violle et al., 2017) due to the ecosystem services they provide, such as the

high values of biomass contribution of turtles (*e.g.*, red-eared slider turtle, *Trachemys*

scripta, Lovich et al., 2018), mineral cycling and bioaccumulation (*e.g.*, *Chelydra*

serpentine; Hebert et al., 1993), or seed dispersal (*e.g.*, Galapagos tortoises, *Chelonoidis*

spp; Blake et al., 2012).

Local disturbances such as habitat disturbance and unsustainable local consumption are

the main threats for the loss of functional diversity in the studied species. Habitat

disturbances affect most species of turtles and crocodilians, especially in North America

and Europe (Fig. 4A), independent of their life history strategies. Therefore, habitat

protection for these species is key to preserve the functional diversity (Lourenço-de-

Moraes et al., 2021). In fact, Mittermeier et al. (2015) highlighted that the protection of

only 16 hotspot of turtle biodiversity would successfully encompass 83% of these species.

On the other hand, we find that the second local threat for the functional diversity of

Testudines and Crocodiles is unsustainable local consumption. Our results predict the

extinction of species with higher rates of clutch size (*e.g*., such as sea turtles, whose eggs

are consumed in high quantities; Madrigal-Ballestero et al., 2013) and long-lived species

(linked with bigger size, which typically provides more meat; Donlan et al., 2010).

Unsustainable local consumption of turtles is one of the most important threats in Asia,

South America, and Africa (Fig. 4A). For example, in China, one million turtle farms are

the primary purchasers of turtles caught in the wild, including several threatened species

(Shi et al., 2007)

Global threats such as unsustainable or illegal international trade and climate

change can also have important effects in reducing the functional diversity of Testudines

and Crocodilia. The global trade of these species is the third most important threat for

their functional diversity, affecting tortoises and turtles mainly for their use as pets

(Luiselli et al., 2016) and crocodiles for their skin (Thorbjarnarson, 1999; Platt and

Thorbjarnarson, 2000). On the other hand, although climate change affects fewer species

than other threats, turtles and crocodilians with unique life history strategies could

disappear according to our projections (Azanza-Ricardo et al., 2017). Indeed, the

consequences of climate change for these species are found worldwide, including extreme

droughts by compromising the fitness of the populations (Poiani and Johnson, 1991;

Rodríguez-Caro et al., 2021), flood nesting sites by rising sea levels (Pike et al., 2015),

coral bleaching due to increasing in temperatures, affecting feeding areas of sea turtles

(Chaloupka et al., 2008) and the unbalance of sex-ratios in populations of temperature

sex-dependent species (Janzen, 1994), such as the threatened leatherback turtle

(*Dermochelys coriacea*; Tomillo et al., 2014).

Our study does not capture all human activities with impacts on species'

ecological strategies. As such, we deem our findings to be conservative. For instance,

although we have summarised the breadth of testudines and crocodilians' threats to

extinction, threat interactions, which remain largely unknown in these taxonomic groups,

have been reported to have synergistic effects on the loss of species, as was described for

mammals (Romero-Muñoz et al., 2020) and birds (Chen et al., 2015). Moreover,

examining the ecological functions of each species in higher resolution, by examining

their diets or habitat breadth (*i.e.*, generalist–specialist), could improve the available

information of ecological function (Cooke et al., 2019).

Protecting threatened species with unique life history strategies is key to conserve

their functional diversity (Violle et al., 2016) and ecosystem services they provide

(Lovich et al., 2018). Our findings support previous studies that suggest that conservation

policies should consider together the taxonomic and functional dimensions of biodiversity

(Carmona et al., 2021) to implement targeted and more effective conservation actions in

the context of a global biodiversity crisis (Toussaint et al., 2021). The IUCN Red List of

Threatened Species may include information about population trends, distribution area,

threats, and abundance. The inclusion of functional diversity could greatly aid

international, national, and local efforts to preserve functionally unique species, and

recently, new indexes have been developed to include some aspects of species

functionality. The IUCN Green Status of Species (Grace et al. 2021; IUCN 2021), a new

part of the Red List assessment, quantifies species recovery and explicitly includes

ecological function (Akçakaya et al., 2020). Incorporating the functional diversity of life

history strategies in conservation assessments constitutes a promising approach to help

prioritise conservation efforts to maintain high levels of functional diversity against

current and future threats.

MATERIALS AND METHODS

Life history traits

To describe the life cycle of species in the orders Testudines and Crocodilia, we used life

history trait data. Life history traits quantify how the life cycle of a species and its

underlying vital rates —survival, development, reproduction— have been shaped by

natural selection to optimise its performance (Roof, 1992). We obtained such traits from

published literature, open access databases, and via direct researcher contributions

(Cayuela et al. 2019). In some cases, life history trait information was derived from the

demographic information of different databases and datasets. These data were obtained

from the COMADRE Animal Matrix Database (Salguero-Gómez et al. 2016b), DATLife

Database (2021), Amniote Life History Database (Myhrvold et al. 2015), and the reviews

by Allen et al. (2017), Pfaller et al. (2018) and Reinke et al. (under review) (more

information about data selection in Appendix 2).

To quantify species' life history strategies, we selected six life history traits that

encompass detailed information regarding the timing, intensity, frequency, and duration

of demographic processes across the life cycle of any species (Stearns, 1992). These traits

are well-established metrics in comparative demographic analyses, and their emerging

syndromes typically explain high variance (~75%) in life history strategies of

multicellular organisms (Bielby et al., 2007; Carmona et al., 2021; Gaillard et al. 1989,

2005; Paniw et al., 2018; Salguero-Gómez et al. 2016a; Stearns 1992). The life history

traits chosen were adult survival (Sa), juvenile survival (Sj), maximum lifespan (ML), age

at sexual maturity ($L\alpha$), mean number of clutches per year (CN), and clutch size (CS;

Table S1). The data obtained from the aforementioned sources encompass 259 species

with at least one of these six traits (Fig. S2).

To calculate the life history strategies of Testudines and Crocodilia, we
collected/estimated the life history traits according to the different datasets:

- Mean number of clutches per year and clutch size (*CN* and *CS*, respectively) were

obtained from Amniote (Myhrvold et al. 2015) and Allen et al. (2017). Moreover,

Allen et al. (2017) includes several data (including Amniote) and, when they found

multiple records of the same trait for a species, the average of the species' trait was

estimated by taking the mean of unique records per species. When the same species

was present in both datasets, we selected the value in Allen et al. (2017).

- Adult and juvenile survival (*Sa* and *Sj*, respectively) were calculated from the

MPMs in COMADRE (Salguero-Gómez et al., 2016a), life tables in DATLife

(2021), and from direct estimates of capture-recapture published studies (such as

Pfaller et al., 2018; Cayuela et al. 2019). Adult survival (*Sa*) from MPMs was

estimated as the column sum of the matrix U of the stages representing reproductive

individuals represented in the sub-matrix F (considering adults, all ages/stages after

the first reproduction). We used arithmetic averages for the resulting stage-specific

survival values, rather than weighting them by the stable stage distribution (*e.g.*,

Roper et al., 2021) because we collected juvenile and adult data from other sources

where calculating this weighted mean would not feasible, thus rendering

comparisons impossible. In DATLife, we used age-specific fertility to determine

the reproductive ages, and then estimated adult survival as the arithmetic average

of age-specific survival rates after the first reproduction. We estimated juvenile

survival (*Sj*) using a similar approach: as the average of the sum of the columns that

represent juveniles (all the stages prior to the first reproduction) in the pertinent

sub-matrix U in COMADRE. In DATLife, we estimated the average of non-

reproductive age- or stage-specific survival (prior to the first reproduction) of the

life tables.

- Age at maturity ($L\alpha$) and maximum lifespan (ML) were estimated with the MPMs

from COMADRE or obtained directly from the databases with this information

available (such as Amniote or DATLife). In COMADRE, $L\alpha$ was calculated using

age-from-stage decompositions (Caswell 2001, p. 124). Briefly, we defined the

reproductive stages as those columns of the F sub-matrix that contain values greater

than zero. We estimated the mean time between birth and the first entry into the

reproductive stage. This conditional mean is obtained by creating an absorbing state

corresponding to the event of reproducing at least once before death, with a Markov

chain, and calculating the mean time to absorption (Caswell, 2001). ML was

estimated from the MPMs of COMADRE projecting 100 individuals in the first

state and iterating up to 1000 years to identify the first year with fewer than one

individual in the virtual cohort. We also used the databases Amniote and DATLife

because they contain information about $L\alpha$ and ML of most of the examined species.

When several values from different datasets for one species and life history trait

were available, we selected the lowest value for $L\alpha$ and the highest values for ML.

The rationale behind this choice is that the lowest value of $L\alpha$ identifies the most

likely first age at maturity reported in the life cycle of this species, whereas the

highest values of ML approximate the maximum values of this trait reported for the

species of interest in wildlife populations.

Phylogeny and data imputation

To identify and quantify the potential role of evolution in shaping our species' life history

strategies, we performed phylogenetic comparative analyses. We used a species-level

phylogenetic tree recently published for Testudines and Crocodilia (Colston et al. 2020)

to both impute the missing data and account for the effect of evolutionary constraints in

the observed life history strategies. Briefly, to build the phylogenetic tree, 14

mitochondrial loci and six nuclear loci were sampled for 357 species of extant or recently

extinct turtle and tortoises, and the 27 species described crocodilian species (Rhodin et

al. 2017). Colston et al. (2020) computed Maximum Likelihood trees with RaxML

(Stamatakis 2014) and used taxonomic imputation to fill the gaps of 17 out of 384 (<5%)

species in the tree. Further details on the methods for tree construction are available in

Colston et al. (2020). We selected the demographic information of species when

information about the phylogeny is available, to perform phylogenetic comparative

analyses (Revell, 2010; Revell, 2012), the final dataset included 259 species of Crocodilia

(23) and Testudines (236) with this resource

Demographic data of reptiles are scarce and contain missing values (Conde et al.

2019; Fig S2). As principal component analyses require full data (*i.e.* non-gappy data;

Rubin 1976, Nakagawa 2004), we carried out phylogenetic imputations to fill in the gaps

in the life history traits for the studied species (see Appendix 2). To impute the missing

life history traits in our dataset, we used the R function *phylomice* from the package *mice*

(Van Buuren & Groothuis-Oudshoorn, 2011), which use multiple imputation with

phylogenetic information. This approach has been shown recently to correctly impute

missing data in similar datasets better than other available approaches (Murray, 2018).

Briefly, this approach uses a bootstrap expectation–maximization algorithm to impute

missing data (Honaker et al., 2011). As suggested by previous studies (Janssen et al.,

2010), we included two more traits in the imputation analyses in order to have more robust

estimated of the imputed data (we included body weight and body size, which is available

for all the individuals, and incubation time available for 16% of the species from Amniote

Database). We then created 40 imputed datasets and ran analyses on each separately. This

approach also is based on using the position of each species in the phylogenetic tree to

impute new values under the assumption of strong phylogenetic signal (Goolsby et al.

2017). To check this assumption, we first estimated the phylogenetic signal for each trait

separately using Pagel's λ, which describes the strength of phylogenetic relationships on

trait evolution under a Brownian motion model (Freckleton 2000). Pagel's λ ranges

between 0, when the patterns in the traits cannot be explained by the employed phylogeny,

and 1 when the observed patterns in traits are tightly correlated with the placement of

species in the phylogeny. Most of the selected traits showed a strong phylogenetic signal

(> 0.8; see Table S3), so we used the imputed values for traits in further analyses.

Exploring the relation between the axes of life history strategies

To identify potential differences between the patterns of association among life history

traits for Testudines and Crocodilia species while simultaneously assessing non-

independence of lineages, we used a phylogenetically informed PCA (*p*PCA; Revell,

2009). *p*PCA is a multivariate analysis that reduces the number of variables of interest

due to their plausible correlation. The *p*PCA considers the correlation matrix of species'

traits while accounting for phylogenetic relationships and estimate Pagel's λ. The *p*PCA

was estimated using the R package *phytools* (Revell, 2012), assuming a Brownian motion

model of evolution (Revell, 2010). Life history trait data were log-transformed to fulfil

normality assumptions of PCA and z-transformed to mean $= 0$, and SD $= 1$ (Legendre &

Legendre, 2021). We used the Kaiser criterion (Kaiser, 1960) after optimisation through

varimax rotation to determine the number of axes necessary to explain a substantial

amount of variation. Namely, the axes we retained had an associated eigenvalue >1.

As body weight is strongly correlated with life history traits (Gaillard et al., 1989; Bielby

et al. 2007; Capdevila et al., 2020), we accounted for this potential effect in our

multivariate analyses too. Out of the multiple possible ways to account for body size or

weight in life history analyses, we used the residuals of phylogenetic regression between

body size and each trait (Revell, 2009). We collected most of adult body mass (g) data

from the studies by Myhrvold et al. (2015) and Colston et al. (2020), moreover, we

collected the remained data from previous literature for all our 259 species.

Estimating the functional spectra

To describe the probabilistic distribution of the species within the functional spaces, we

used the main axes of functional trait variation by pPCA corrected by size (*i.e.,* residuals

of regressions). We performed multivariate kernel density using the '*TPD'* and '*ks'* R

packages (Duong, 2007, 2014; Carmona, 2019; Carmona et al., 2019) to estimate the

functional spectra. The kernel was a multivariate normal distribution for each species

centered in the location of the species in pPCA and bandwidth chosen using unconstrained

bandwidth selectors from the '*Hpi'* function in the '*ks'* package (Duong & Hazelton,

2003). The grouped kernels for all species drive into the continuous TPD function

(Carmona et al., 2017; 2019). We divided the continuous functional space into a two-

dimensional grid composed of 200 equal sized cells per dimension because it was

suggested by previous authors (Carmona et al., 2021; Toussaint et al., 2021). We

estimated the value of the TPD function for the 40,000 cells. The value of the TPD

function represent the density of species in that particular region of the functional space

(i.e., species with similar life history strategy). Results of TPD were represented

graphically with the contours containing 50, 60, 70, 80, 90, and 99% of the total density

of species.

Threats to the studied species

To identify causes of recent declines in Testudines and Crocodilia species we reviewed published information. The main threats described for these species were i) habitat loss, fragmentation, and degradation (Luiselli, 2009), ii) over-collection of individuals and their eggs for food consumption (Gong et al., 2017), iii) unsustainable or illegal international trade, as well as over-collection for the trade in medicines (Sung & Fong, 2018), iv) climate change (Gibbons et al., 2000); v) emerging diseases (Jacobson et al., 2014; Tompkins et al., 2015). We collected the information on the threats to each species from Stanford et al. (2020), Bonin et al. (2006) and IUCN Red List (IUCN, 2020). We did dichotomous variables when one of these threats were mentioned. We were not able to quantify the effect of the threat, because effect size is only available for the Critically Endangered species of Stanford et al., (2020). To convert the threat descriptions of the Red List into our broader categories, we used the following process: for habitat disturbances we considered the Red List threat classifications "residential & commercial development", "agriculture & aquaculture" and "natural system modifications"; for climate change the classification "climate change and extreme weather"; and for disease the classification "invasive and other problematic species, genes & diseases".

Effects of extinctions of threatened species on global functional diversity

To identify the loss of functional diversity by threatened species extinctions, we simulated the disappearance of species according to their Red List category (IUCN, 2020). The species classified among the threatened species categories (i.e. Critically Endangered (CR), Endangered (EN) and Vulnerable (VU) have a higher risk of extinction than the

rest of species: Near Threatened (NT), Least Concern (LC); the extinction risk of Data

Deficient (DD) species is unknown. In our analysis, we removed threatened species and

estimated the resulting shifts in functional diversity. Firstly, we removed the species with

a higher risk of extinction (-CR), then we removed successively the species with lower

risks (-EN includes the extinction of species classified as a CR and EN, -VU includes CR,

EN and VU, and -NT includes CR, EN, VU and NT). These extinction scenarios

represented a gradient of extinction risk from the persistence of all species to a more

dramatic scenario where all threatened species (including the NT species) went extinct.

We compared the TPD function considering all the species assessed by IUCN (current

spectra of functional diversity), and TPD function after removing the species in the

different scenarios. We can compare TPD functions because probability density functions

can be integrated to 1 across the whole functional space (Carmona et al. 2016). To reduce

the potential effect of outliers in the functional space, we applied a quantile threshold of

99%. To quantify how much the functional spectra is lost after the extinction scenarios,

we estimated which functional space cells become empty after extinctions. This space

(TPD function of the current spectra – TPD function after removing the threatened

species) represented the loss of functional diversity for Testudines and Crocodilia. We

calculated this loss in the functional space for all the extinction scenarios (-CR, -EN, -

VU, and -NT).

To assess if the impacts of each extinction scenario were different from what would be

expected if extinction risk is not related to life history traits, we also compared the

observed changes in functional diversity to a null model where the extinct species were

randomly selected within pool of species. For each scenario of extinction risk, we

compared the functional diversity to 999 losses of functional diversity where the same

number of threatened species were randomly selected among the pool of species. This

strategy allowed us to understand whether the extinction in the different scenarios reduces

more or less than expected the functional diversity. For each scenario, we created 999

TPD functions simulating cases in which the same number of species were lost at random

from the total set of IUCN-assessed species. We compared the 5% and 95% percentile of

the random simulation with the value of loss of functional diversity per each extinction

scenario to estimate if the values were significant different.

After examining the loss of functional diversity, we plotted the extinction risk of species

within the functional spaces, using the species assessed by IUCN. Using binomial

smoother-based GAM (Wood, 2006) with the R package *mgcv* (Wood, 2011), we

analysed the relationship between extinction risk (1: high-risk and 0: nonthreatened or

low-risk) and the position in the functional space (PC axes). We then represented the

predictions of the models (including the 95% confidence intervals of the means) to

visually examine how life history strategies affect the probability of species being

threatened.

Effects of anthropic threats on global functional diversity

To identify the effect of the different threats in the functional spectra, we simulated

scenarios where extinctions were based upon species reported as affected by specific

threats. We did two comparisons: 1) the TPD function considering all the species (current

spectra of functional diversity) and TPD function after removing the species affected by

specific threat (habitat disturbance, trade, local consumption, climate change or disease)

and, 2) the TPD function considering all the IUCN-assessed species, and the TPD

function after removing the threatened species (CR, EN, or VU) affected by specific

threat. With the comparison between TPD functions, we estimated the potential loss of

functional diversity attributable to each threat and evaluated the differences between

them. We also compared the observed changes in functional diversity using a null model

to assess if the impacts of each extinction scenario were different from what would be

expected by chance. For each threat, we compared the loss of functional diversity with

the 999 losses of functional diversity where the same number of species were randomly

selected among the pool of the species. We did the comparison for the extinction of all

the species affected by each threat and also for threatened species.

To visually the relation between life history strategies and the threats, we mapped the

probability of a species being affected by a specific threat within the functional spaces.

We used binomial smoother-based GAM (Wood, 2006) with the R package *mgcv* (Wood,

2011) to identify the relationship between being affected by each threat (1: affected and

0: nonaffected) and the position in the functional space (PCA axes). We represented the

predictions of the models (including the 95% confidence intervals of the means) when

the relation between threat and life history strategy was significant.

ACKNOWLEDGEMENTS

RCRC was supported by a post-doctoral grant funded by the Regional Valencian

Government (APOSTD/2020/090) hosted by RSG. RSG was supported by a NERC IRF

grant (NE/M018458/1). The Spanish Ministry of Science and European Regional

Development Fund funded this work through Project PID2019-105682RA-

I00/AEI/10.13039/501100011033 (MINECO).

REFERENCES

Allen, W. L., Street, S. E., & Capellini, I. (2017). Fast life history traits promote invasion

success in amphibians and reptiles. Ecology Letters, 20(2), 222-230.

Akçakaya et al. (2020). Assessing ecological function in the context of species recovery.

Conservation Biology, 34(3), 561-571.

Azanza-Ricardo, J., Martín, M. E. I., Sansón, G. G., Harrison, E., Cruz, Y. M., & Bretos,

F. (2017). Possible effect of global climate change on *Caretta caretta* (Testudines,

Cheloniidae) nesting ecology at Guanahacabibes Peninsula, Cuba. Chelonian

conservation and Biology, 16(1), 12-19.

Bielby, J., Mace, G. M., Bininda-Emonds, O. R. P., Cardillo, M., Gittleman, J. L., Jones,

K. E., … Purvis, A. (2007). The fast-slow continuum in mammalian life history:

An empirical reevaluation. The American Naturalist, 169(6), 748–757.

Blake, S., Wikelski, M., Cabrera, F., Guezou, A., Silva, M., Sadeghayobi, E., ... &

Jaramillo, P. (2012). Seed dispersal by Galápagos tortoises. Journal of

Biogeography, 39(11), 1961-1972.

Bonin, F., Devaux, B., & Dupré, A. (2006). Turtles of the World. JHU Press.

Capdevila, P., Beger, M., Blomberg, S. P., Hereu, B., Linares, C., & Salguero-Gómez, R.

(2020). Longevity, body dimension and reproductive mode drive differences in

aquatic versus terrestrial life-history strategies. Functional Ecology, 34(8), 1613-

1625.

Carmona, C. P., De Bello, F., Mason, N. W., & Lepš, J. (2016). Traits without borders:

integrating functional diversity across scales. Trends in ecology & evolution,

31(5), 382-394.

Carmona, C. P. (2019). TPD: Methods for Measuring Functional Diversity Based on Trait

Probability Density. https://cran.r-project.org/web/packages/TPD/index.html.

Carmona, C. P., de Bello, F., Mason, N. W., & Lepš, J. (2019). Trait probability density
(TPD): measuring functional diversity across scales based on TPD with R.
Ecology, 100(12), e02876.

Carmona, C. P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., ... &
Toussaint, A. (2021). Erosion of global functional diversity across the tree of life.
Science Advances, 7(13), eabf2675.

Caswell, H. (2001). Matrix population models: Construction, analysis, and interpretation
(2nd ed.). Sunderland, MA: Sinauer Associates.

Cayuela, H., Akani, G. C., Hema, E. M., Eniang, E. A., Amadi, N., Ajong, S. N., ... &
Luiselli, L. (2019). Life history and age-dependent mortality processes in tropical
reptiles. Biological Journal of the Linnean Society, 128(2), 251-262.

Cernansky, R. (2017). The biodiversity revolution. Nature 546, 22–24.

Chaloupka, M., Kamezaki, N., & Limpus, C. (2008). Is climate change affecting the
population dynamics of the endangered Pacific loggerhead sea turtle?. Journal of
Experimental Marine Biology and Ecology, 356(1-2), 136-143.

Chen, S., Fan, Z., Roby, D. D., Lu, Y., Chen, C., Huang, Q., ... & Zhu, J. (2015). Human
harvest, climate change and their synergistic effects drove the Chinese Crested
Tern to the brink of extinction. Global Ecology and Conservation, 4, 137-145.

Chichorro, T., Juslén, A., Cardoso, P. (2019). A review of the relation between species
traits and extinction risk. Biol. Conserv. 237, 220–229.

Colston, T. J., Kulkarni, P., Jetz, W., & Pyron, R. A. (2020). Phylogenetic and spatial
distribution of evolutionary diversification, isolation, and threat in turtles and
crocodilians (non-avian archosauromorphs). BMC Evolutionary Biology, 20(1),
1-16.

Conde, D. A., Staerk, J., Colchero, F., da Silva, R., Schöley, J., Baden, H. M., ... &

Vaupel, J. W. (2019). Data gaps and opportunities for comparative and

conservation biology. Proceedings of the National Academy of Sciences, 116(19),

9658-9664.

Cooke, R. S., Eigenbrod, F., & Bates, A. E. (2019). Projected losses of global mammal

and bird ecological strategies. Nature Communications, 10(1), 1-8.

DATLife Database. (2021). Max-Planck Institute for Demographic Research (Germany).

Available at www.datlife.org (data downloaded on [25-02-2021]

Díaz, S., Settele, J., Brondízio, E. S., Ngo, H. T., Agard, J., Arneth, A., ... & Zayas, C. N.

(2019). Pervasive human-driven decline of life on Earth points to the need for

transformative change. Science, 366(6471).

Donlan, C. J., Wingfield, D. K., Crowder, L. B., & Wilcox, C. (2010). Using expert

opinion surveys to rank threats to endangered species: a case study with sea turtles.

Conservation Biology, 24(6), 1586-1595.

Duong, T. (2007). Ks: Kernel density estimation and kernel discriminant analysis for

multivariate data in R. Journal of Statistical Software, 21(7), 1-16.

Duong, T. (2014). ks: Kernel Smoothing. R Package Version 1.9.2 (R package version

1.11.4). https://CRAN.R-project.org/package=ks.

Duong, T., & Hazelton, M. (2003). Plug-in bandwidth matrices for bivariate kernel

density estimation. Journal of Nonparametric Statistics, 15(1), 17-30.

Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., & Norberg,

J. (2003). Response diversity, ecosystem change, and resilience. Frontiers in

Ecology and the Environment, 1(9), 488-494.

Freckleton, R. P. (2000). Phylogenetic tests of ecological and evolutionary hypotheses:

Checking for phylogenetic independence. Functional Ecology, 14(1), 129–134.

Gaillard, J. M., Pontier, D., Allaine, D., Lebreton, J. D., Trouvilliez, J., & Clobert, J.

(1989). An analysis of demographic tactics in birds and mammals. Oikos, 59-76.

Gaillard, J.-M., Yoccoz, N G, Lebreton, J.-D., Bonenfant, C., Devillard, S., Loison, A.,

… Allaine, D. (2005). Generation time: A reliable metric to measure life-history

variation among mammalian populations. The American Naturalist, 166(1), 119–

123. https://doi.org/10.1086/430330

Germano, D. J. (1994). Comparative life histories of North American tortoises. Biology

of North American Tortoises, 175-185.

Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S.,

Greene, J.L., Mills, T., Leiden, Y., Poppy, S., and Winne, C.T. (2000). The global

decline of reptiles, déja vu amphibians. Bioscience 50, 653–666.

Gibbs, J. P., & Amato, G. D. (2000). Genetics and demography in turtle conservation.

Turtle conservation, 207-217.

Gong, S., Shi, H., Jiang, A., Fong, J.J., Gaillard, D., and Wang, J. (2017). Disappearance

of endangered turtles within China's nature reserves. Curr. Biol. 27, 170–171.

Goolsby, E. W., Bruggeman, J., & Ane, C. (2017). R phylopars: fast multivariate

phylogenetic comparative methods for missing data and within-species

variation. Methods in Ecology and Evolution, 8(1), 22-27.

Grace, M. K., Akçakaya, H. R., Bennett, E. L., Brooks, T. M., Heath, A., Hedges, S., ...

& Quintana, M. G. (2021). Testing a global standard for quantifying species

recovery and assessing conservation impact. Conservation Biology, 35(6), 1833-

1849.

Healy, K., Ezard, T. H., Jones, O. R., Salguero-Gómez, R., & Buckley, Y. M. (2019).

Animal life history is shaped by the pace of life and the distribution of age-specific

mortality and reproduction. Nature ecology & evolution, 3(8), 1217-1224.

Hebert, C. E., Glooschenko, V., Haffner, G. D., & Lazar, R. (1993). Organic contaminants

in snapping turtle (*Chelydra serpentina*) populations from southern Ontario,

Canada. Archives of Environmental contamination and Toxicology, 24(1), 35-43.

Honaker, J., King, G., & Blackwell, M. (2011). Amelia II: A program for missing data.

Journal of statistical software, 45(1), 1-47.

Hughes, P. W. (2017). Between semelparity and iteroparity: empirical evidence for a

continuum of modes of parity. Ecology and evolution, 7(20), 8232-8261.

IUCN. (2020). The IUCN Red List of Threatened Species. Version 2020–3; 2020.

IUCN (2021). IUCN Green Status of Species: a global standard for measuring species

recovery and assessing conservation impact. IUCN, Gland, Switzerland.

https://portals.iucn.org/library/node/49511

Jackson, S. T., & Sax, D. F. (2010). Balancing biodiversity in a changing environment:

extinction debt, immigration credit and species turnover. Trends in ecology &

evolution, 25(3), 153-160.

Jacobson, E.R., Brown, M.B., Wendland, L.D., Brown, D.R., Klein, P.A., Christopher,

M.M., and Berry, K.H. (2014). Mycoplasmosis and upper respiratory tract disease

of tortoises: a review and update. Vet. J. 201, 257–264.

Janssen, K. J., Donders, A. R. T., Harrell Jr, F. E., Vergouwe, Y., Chen, Q., Grobbee, D.

E., & Moons, K. G. (2010). Missing covariate data in medical research: to impute

is better than to ignore. Journal of clinical epidemiology, 63(7), 721-727.

Janzen, F. J. (1994). Climate change and temperature-dependent sex determination in

reptiles. Proceedings of the National Academy of Sciences, 91(16), 7487-7490.

Kaiser, H. F. (1960). The varimax criterion for analytic rotation in factor analysis.

Psychometrika, 23, 187–200.

Kuussaari, M., Bommarco, R., Heikkinen, R. K., Helm, A., Krauss, J., Lindborg, R., ...

& Steffan-Dewenter, I. (2009). Extinction debt: a challenge for biodiversity

conservation. Trends in ecology & evolution, 24(10), 564-571.

Legendre, P., & Legendre, L. (2012). Numerical ecology (3rd ed.). Amsterdam, the

Netherlands: Elsevier Science.

Lovich, J. E., Ennen, J. R., Agha, M., & Gibbons, J. W. (2018). Where have all the turtles

gone, and why does it matter?. BioScience, 68(10), 771-781.

Lourenço-de-Moraes, R., Campos, F. S., Carnaval, A. C., Otani, M., França, F. G.,

Cabral, P., & Benedito, E. (2021). No more trouble: An economic strategy to

protect taxonomic, functional and phylogenetic diversity of continental turtles.

Biological Conservation, 261, 109241.

Luiselli, L. (2009). A model assessing the conservation threats to freshwater turtles of

Sub-Saharan Africa predicts urgent need for continental conservation planning.

Biodivers. Conserv. 18, 1349–1360.

Luiselli, L., Starita, A., Carpaneto, G. M., Segniagbeto, G. H., & Amori, G. (2016). A

short review of the international trade of wild tortoises and freshwater turtles

across the world and throughout two decades. Chelonian Conservation and

Biology, 15(2), 167-172.

Madrigal-Ballestero, R., Schlüter, A., & Lopez, M. C. (2013). What makes them follow

the rules? Empirical evidence from turtle egg harvesters in Costa Rica. Marine

Policy, 37, 270-277.

Maxwell, S. L., Fuller, R. A., Brooks, T. M., & Watson, J. E. (2016). Biodiversity: The

ravages of guns, nets and bulldozers. Nature News, 536(7615), 143.

Mittermeier, R. A., van Dijk, P. P., Rhodin, A. G., & Nash, S. D. (2015). Turtle hotspots:

an analysis of the occurrence of tortoises and freshwater turtles in biodiversity

673   hotspots, high-biodiversity wilderness areas, and turtle priority areas. Chelonian

674   Conservation and Biology, 14(1), 2-10.

675 Murray, J. S. (2018). Multiple imputation: a review of practical and theoretical findings.

676   Statistical Science, 33(2), 142-159.

677 Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. M.

678   (2015). An amniote life-history database to perform comparative analyses with

679   birds, mammals, and reptiles: Ecological Archives E096-269. Ecology, 96(11),

680   3109-3109.

681 Naeem, S. (1998). Species redundancy and ecosystem reliability. Conservation biology,

682   12(1), 39-45.

683 Nakagawa S (2004) A farewell to Bonferroni: the problems of low statistical power and

684   publication bias. Behav Ecol 15:1044–1045

685 Oli, M. K. (2004). The fast-slow continuum and mammalian life-history patterns: An

686   Empirical evaluation. Basic Appl Ecol 5(5):449–463.

687 Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., … &

688   Bullock, J. M. (2015). Biodiversity and resilience of ecosystem functions. Trends

689   in ecology & evolution, 30(11), 673-684.

690 Paniw, M., Ozgul, A., & Salguero-Gómez, R. (2018). Interactive life-history traits predict

691   sensitivity of plants and animals to temporal autocorrelation. Ecology Letters,

692   21(2), 275–286.

693 Pfaller, J. B., Chaloupka, M., Bolten, A. B., & Bjorndal, K. A. (2018). Phylogeny,

694   biogeography and methodology: a meta-analytic perspective on heterogeneity in

695   adult marine turtle survival rates. Scientific reports, 8(1), 1-10.

696 Pike, D. A., Roznik, E. A., & Bell, I. (2015). Nest inundation from sea-level rise threatens

697   sea turtle population viability. Royal Society Open Science, 2(7), 150127.

Pimm, S. L., Russell, G. J., Gittleman, J. L., & Brooks, T. M. (1995). The future of
biodiversity. Science, 269(5222), 347-350.

Platt, S. G., & Thorbjarnarson, J. B. (2000). Population status and conservation of
More'et's crocodile, *Crocodylus moreletii*, in northern Belize. Biological
Conservation, 96(1), 21-29.

Poiani, K. A., & Johnson, W. C. (1991). Global warming and prairie wetlands.
BioScience, 41(9), 611-618.

Reinke, B. A., Cayuela, H., Janzen F. J., Lemaitre, J. F., Gaillard, J. M., … & Miller, D.
Under review. A. W. Under review. Diverse aging rates in ectothermic tetrapods
provide insights for the evolution of aging and longevity.

Revell, L. J. (2009). Size-correction and principal components for interspecific
comparative studies. Evolution, 63(12),

Revell, L J. (2010). Phylogenetic signal and linear regression on species data. Methods
in Ecology and Evolution, 1(4), 319–329.

Revell, L. J. (2012). *phytools*: An R package for phylogenetic comparative biology (and
other things). Methods in Ecology and Evolution, 3(2), 217–223.

Rhodin AGJ, Iverson JB, Bour R, Fritz U, Georges A, Shaffer HB, et al. (2017). Turtles
of the world. Annotated checklist and atlas of taxonomy, synonymy, distribution,
and conservation status. 7th ed.

Rhodin, A. G., Stanford, C. B., Van Dijk, P. P., Eisemberg, C., Luiselli, L., Mittermeier,
R. A., ... & Vogt, R. C. (2018). Global conservation status of turtles and tortoises
(order Testudines). Chelonian Conservation and Biology, 17(2), 135-161.

Ripple, W. J., Wolf, C., Newsome, T. M., Hoffmann, M., Wirsing, A. J., & McCauley,
D. J. (2017). Extinction risk is most acute for the world's largest and smallest

vertebrates. Proceedings of the National Academy of Sciences, 114(40), 10678-
10683.

Rodríguez-Caro, R. C., Capdevila, P., Graciá, E., Barbosa, J. M., Giménez, A., &
Salguero-Gómez, R. (2021). The limits of demographic buffering in coping with
environmental variation. Oikos, 130: 1346–1358

Roff, D. (1993). Evolution of life histories: theory and analysis. Springer Science &
Business Media.

Romero-Muñoz, A., Benítez-López, A., Zurell, D., Baumann, M., Camino, M., Decarre,
J., ... & Kuemmerle, T. (2020). Increasing synergistic effects of habitat destruction
and hunting on mammals over three decades in the Gran Chaco. Ecography, 43(7),
954-966.

Roper, M., Capdevila, P., & Salguero-Gomez, R. (2021). Senescence: why and where
selection gradients might not decline with age. Proceedings of the Royal Society
B: Biological Sciences.

Rubin DB (1976) Inference and missing data. Biometrika 63:581–590

Sæther. B. E. (1987). The influence of body weight on the covariation between
reproductive traits in European birds. Oikos 48(1):79–88.

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Bein, C., deBuhr, H., Farack, C.,
… Vaupel, J. W. (2016b). COMADRE: A global data base of animal
demography. Journal of Animal Ecology, 85(2), 371–
384. https://doi.org/10.1101/027821

Salguero-Gómez, R., Jones, O. R., Jongejans, E., Blomberg, S. P., Hodgson, D.
J., Mbeau-Ache, C., … Buckley, Y. M. (2016a). Fast–slow continuum and
reproductive strategies structure plant life-history variation

worldwide. Proceedings of the National Academy of Sciences of the United States of America, 113(1), 230–235.

Scott, R., Marsh, R., & Hays, G. C. (2012). Life in the really slow lane: loggerhead sea turtles mature late relative to other reptiles. Functional Ecology, 26(1), 227-235.

Shi, H., Parham, J. F., Lau, M., & Tien-Hsi, C. (2007). Farming endangered turtles to extinction in China. Conservation Biology, 21(1), 5-6.

Stanford, C. B., Iverson, J. B., Rhodin, A. G., van Dijk, P. P., Mittermeier, R. A., Kuchling, G., ... & Walde, A. D. (2020). Turtles and tortoises are in trouble. Current Biology, 30(12), R721-R735.

Stamatakis A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics, 30:1312–3.

Stearns, S. C. (1999). The evolution of life histories. New York, NY: Oxford University Press.

Sung, Y.-H., and Fong, J.J. (2018). Assessing conservation trends and illegal activity by monitoring the online wildlife trade. Biological Conservation. 227, 219–225.

Thorbjarnarson, J. (1999). Crocodile tears and skins: international trade, economic constraints, and limits to the sustainable use of crocodilians. Conservation Biology, 13(3), 465-470.

Tomillo, P. S., Oro, D., Paladino, F. V., Piedra, R., Sieg, A. E., & Spotila, J. R. (2014). High beach temperatures increased female-biased primary sex ratios but reduced output of female hatchlings in the leatherback turtle. Biological Conservation, 176, 71-79.

Tompkins, D.M., Carver, S., Jones, M.E., Krosek, M., and Skerrat, L.F. (2015). Emerging infectious diseases of wildlife: a critical perspective. Trends in Parasitology. 31, 149–159.

Toussaint, A., Brosse, S., Bueno, C. G., Pärtel, M., Tamme, R., & Carmona, C. P. (2021).

Extinction of threatened vertebrates will lead to idiosyncratic changes in

functional diversity across the world. Nature Communications, 12(1), 1-12.

Van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by

chained equations in R. Journal of statistical software, 45(1), 1-67.

Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N. J., Cadotte, M. W.,

Livingstone, D. & Mouillot, D. (2017). Functional rarity: the ecology of outliers.

Trends in Ecology & Evolution, 32(5), 356-367.

Wood, S. (2006). Generalized Additive Models: An Introduction with R (CRC Press);

https://books.google.co.uk/books/about/Generalized_Additive_Models.html?id=

hr17lZC-3jQC&pgis=1.

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. J. R. Stat. Soc. 73, 3–36.

Figures

Fig. 1. The global functional spectra of the life history strategies of Testudines and Crocodilia is described by two dominant axes of life history variation: (PC1) fast-slow continuum and (PC2) reproductive strategies. The probabilistic species distribution in this space is defined by the two first principal components axes (PC1 = 39.1% and PC2 = 23.1% of variance explained) of a phylogenetically-corrected principal component analysis (*p*PCA), where adult body mass has also been corrected. The phylogenetic signal from this analysis as per Pagel's λ is 0.674 ± 0.030. The life history traits are: adult survival (*Sa*), juvenile survival (*Sj*), maximum lifespan (*ML*), age at sexual maturity (*La*), mean of number of clutches per year (*CN*), clutch size (*CS*). Arrows indicate the direction and weighting of each trait in the *p*PCA. The colour gradient (red, yellow, and white) depicts the density of species in the defined space, where red corresponds to more densely populated areas). Thick contour lines indicate the 0.5 and 0.99 quantiles, and thinner ones indicate quantiles 0.6, 0.7, 0.8, and 0.9. The species silhouettes correspond to (starting at the top left and moving anticlockwise): painted turtle (*Chrysemys picta*), American alligator (*Alligator mississippiensis*), desert tortoise (*Gopherus agassizii*), Hood Island giant tortoise (*Chelonoidis hoodensis*), Orinoco crocodile (*Crocodylus intermedius*), loggerhead sea turtle (*Caretta caretta*), leatherback sea turtle (*Dermochelys coriacea*), northern map turtle (*Graptemys geographica*), and Western Caspian turtle (*Mauremys rivulata*). Source: PhyloPic, www.phylopic.org.

Fig. 2. The life history strategies of testudines and crocodilians predict their

vulnerability to extinction due to human threats. A) Losses of functional diversity

under extinction scenarios of threatened species. CR: Critically Endangered, EN:

Endangered, VU: Vulnerable; NT: Near Threatened. The loss of functional diversity is

expressed as a percentage of the current functional diversity in the life history strategies

of species of the order Testudines and Crocodilia (Fig. 1). We simulate the loss of

functional diversity by removing all species from the IUCN category in a progressive

manner, by degree of "endangerement", removing first the species with a higher risk of

extinction (*i.e.* -CR), then continuing progressively to remove species from the categories

with lower threatened risks (-EN scenario: removing CR and EN species; -VU scenario:

removing CR, EN & VU; -NT scenario: removing CR, EN, VU & NT). This loss of

diversity is represented by a black line. For each scenario, we compared the loss of

functional diversity with 999 iterations of a null model where threatened species are

randomly selected among the 259 species in the analyses. The 999 randomisations are

represented by a blue polygon defined by the 5^{th} and 95^{th} percentiles of functional

diversity loss, with the blue line representing the 50^{th} percentile; B) Risk of extinction in

the functional space of Testudines and Crocodilia species. Probability of species being

classified as high-risk of extinction (*i.e.*, CR & EN IUCN status) according to GAMs

(with binomial distribution) using the position of each species in the two-dimensional

functional space as predictors. Yellow tones indicate lower risk of extinction, whereas

purple tones indicate higher risk of extinction. *P* value of the GAM model is 0.03 and χ^2

is 28.59. The red contour lines indicate the average threat probability (proportion of

species classified as threatened in the group). Here, we consider only the 214 out of the

259 available species of testudines and crocodilians for which the threat status is known.

Fig. 3. The life history strategies of Testudines and Crocodilia predict the

vulnerability to extinction due to local consumption, disease, and pollution. Other

risks, such as climate change, global trade, or habitat disturbance, do not show a

significant relation with the species' life history strategies (Fig. S1). Probability of

species being affected by (A), local consumption (B), disease, and (C), pollution

according to GAMs (with binomial distribution) using the position of species in the

two-dimensional functional space (Fig. 1) as predictors. Yellow tones indicate lower

risk of extinction due to each threat, whereas purple tones indicate higher risk. *P* values

associated with each GAM are shown at the top-right corner of each panel. The red

contour lines indicate the average threat probability. We consider only species whose

threats are known (n = 251 species).

Fig. 4. Habitat disturbance is the main threat for the functional diversity, especially in the north hemisphere. a) Ratio of species affected by each threat per continent. The ratio is calculated for species whose threats are known in each continent (North America, n = 65; Europe, n = 4; Asia, n = 66; South America, n = 57; Africa, n = 39; Oceania, n = 14). b) Simulated losses of functional diversity of Testudines and Crocodilia under extinction scenarios by threat. The loss of functional diversity is expressed as a percentage of the total spectra of functional diversity of species, as detailed in Fig. 1. We simulated the loss of functional diversity by removing all species affected by specific threats, in salmon colour removing only threatened species (*i.e.* Critically Endangered [CR], Endangered [EN] & Vulnerable [VU] as per the IUCN Red List) and in cyan removing all the species affected (threatened or not). For each scenario, we compared the loss of functional diversity with 999 iterations of a null model where the same number of species were randomly selected among all 251 species. The 999 randomisations are represented for each threat as a grey dot for the 50th percentile, with grey whiskers representing the 5th and 95th percentiles.