Anthropogenic impacts on threatened species erode functional diversity in

turtles and crocodilians

Rodríguez-Caro, R.C.[1,2*], Graciá, E.[2], Blomberg, S. P.[3], Cayuela, H.[4], Grace, M.[1],

Carmona, C.P.[5], Pérez-Mendoza, H.A.[6], Giménez, A.[2], Salguero-Gómez, R.[1,7,*]

[1] Department of Zoology, Oxford University, 11a Mansfield Road, OX1 3SZ, Oxford, UK.

[2] Departamento de Biología Aplicada, Universidad Miguel Hernández, Elche, Alicante, Spain.

[3] School of Biological Sciences, The University of Queensland, Brisbane, QLD, Australia.

[4] Laboratoire de Biométrie et Biologie Evolutive, Université Lyon 1, CNRS, UMR 5558, Villeurbanne F-

769622, France

[5] Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

[6] Facultad de Estudios Superiores Iztacala, Universidad Autónoma de México, Tlalnepantla, México.

[7] Max Plank Institute for Demographic Research, Konrad-Zuße straße, Rostock, Germany.

* Corresponding authors: r.rodriguez@umh.es

17   rob.salguero@zoo.ox.ac.uk

Abstract: 200 words

Body text (intro, results, and discussion): 2786 words

Methods: 2474 words

Figures: 4

Tables: 0

ABSTRACT (200/200 words)

The Anthropocene is tightly associated with a drastic loss of species worldwide and the

disappearance of their key ecosystem functions. The on-going reduction in ecosystem

functionality is driven by global and local threats. The orders Testudines (turtles and

tortoises) and Crocodilia (crocodiles, alligators, and gharials) contain numerous

threatened, long-lived species for which their functional diversity and potential erosion

by anthropogenic impacts remains unknown. Here, we examine 259 (69%) of the

existing 375 species of Testudines and Crocodilia, quantifying their life history

strategies (*i.e.*, schedules of survival, development, and reproduction) from open-access

data on their demography, ancestry, and threats. We find that the loss of functional

diversity in simulated extinction scenarios of threatened species is greater than expected

by chance. Moreover, the effects of unsustainable local consumption, diseases, and

pollution are associated with specific functional strategies. In contrast, climate change,

habitat disturbance, and global trade affect all species independent of their life history

strategy. Importantly, the loss of functional diversity for threatened species by habitat

disturbance is twice that for all other threats. Our findings highlight the importance of

conservation programmes focused on preserving the functional diversity of life history

strategies jointly with the phylogenetic representativity of these highly threatened

groups.

Keywords: animal disease, biodiversity, comparative demography, climate change,

habitat disturbance, life history trait, phylogenetic analyses, pollution, wildlife trade.

INTRODUCTION

Human impacts, such as habitat loss, climate change, pollution, poaching, and unsustainable trade, impose major threats to the persistence of species worldwide (Maxwell et al., 2016). Indeed, modern human activities have drastically accelerated species extinctions by three orders of magnitude (Pimm et al., 1995). As such, effective conservation measures are urgently needed to bend the curve of biodiversity loss (Díaz et al., 2019).

The on-going biodiversity loss is resulting in the erosion of functional diversity, key to ecological function (Oliver et al., 2015; Carmona et al., 2021; Toussaint et al., 2021). A robust way to characterise diversity in functional traits across a specific clade is via quantifying life history strategies, *i.e.* trade-offs between an individual's investment in survival, development, and reproduction driven by limited resources and physiological constraints (Stearns, 1999; Salguero-Gómez et al., 2016a; Cernansky, 2017; Healy et al., 2019; Capdevila et al., 2020). In this context, different species can possess similar life history trait values, resulting in functional redundancy (Naeem, 1998; *e.g.*, the Texas tortoises, *Gopherus berlandieri,* and the common snapping turtle, *Chelydra serpentina,* achieve similar longevities; Germano, 1994), or display dissimilar functionality, giving rise to functional dispersion (Elmqvist et al., 2003; *e.g.*, the loggerhead sea turtle, *Caretta caretta,* reproduces later than other Testudines of similar size, such as the Aldabra giant tortoise, *Aldabrachelys gigantea*; Scott et al., 2012). Understanding how redundant or dispersed life history traits are and how sensitive these are to anthropogenic threats worldwide is critical for conservation (Chichorro et al. 2019).

Evaluating the risk of extinction of long-lived species is important for an accurate understanding and preservation of their functional diversity. Critically, the

extinction of long-lived species can be delayed due to their long generation times (Gaillard et al., 1989; Jackson and Sax, 2010; Salguero-Gómez et al., 2016a; Capdevila et al. 2020). Thus, environmental effects typically manifest at much longer time scales in long-lived species than in short-lived ones (Gibbs and Amato, 2000; Kuussaari et al., 2009). Perhaps not surprisingly, two of the most threatened and long-lived groups are the order Testudines (tortoises, turtles, and sea turtles) and Crocodilia (crocodiles, alligators, and gharials) (Rhodin et al., 2018; Colston et al., 2020; Cox et al., 2022). In fact, Testudines and Crocodilia contain some of the highest proportions of threatened species across vertebrates (57.9% and 50%, respectively; Cox et al., 2022; UICN, 2020). Moreover, the conservation status of Testudines and Crocodilia could be even worse, since 25% of them lack reliable data to accurately identify their IUCN status (Rhodin et al., 2018).

Here, we characterise the life history strategies of 69% of Testudines and Crocodilia species –259 out of 375 extant species– to identify how the impacts of different human-led global (climate change, global wildlife trade, and emergent diseases) and local threats (habitat disturbances, unsustainable local consumption, and pollution) may alter their functional diversity. Using demographic and phylogenetic open-access data, we quantify life history strategies to estimate their functional diversity. We then simulate different extinction scenarios according to their extinction risk and the main threats that affect each species. We hypothesise that: (H1) the loss of functional diversity due to the extinction of threatened species will be lower than expected by chance. This is so because the life history strategies across testudines and crocodilians are expected to show high functional redundancy, since most of them are long-lived species with delayed maturity (Healy et al., 2019); (H2) human threats (*e.g.,* global trade, unsustainable local consumption, disease) will differentially erode the

functional diversity of these species (Ripple et al., 2017). For instance, some evidence

exists that tortoise and turtle populations may be declining due to habitat disturbance,

unsustainable local consumption, and international pet trade collection (Stanford et al.,

2020). In contrast, threats to the functional diversity of these species due to global

climate change may *a priori* not seem more intense than local threats (Cox et al., 2022).

However, a fundamental knowledge about the consequences of climate change in long-

lived species, such as Testudines and Crocodilia, is still to emerge. Filling in this

knowledge gap is the primary motivation of this research.

RESULTS

High functional redundancy

Two dominant axes of life history traits describe most of the functional diversity of

Testudines and Crocodilia. Using life history trait data from 236 species of turtles and

tortoises (67% of the extant species), and 23 species of crocodilians (85%) from

multiple sources, we characterise their life history strategies via a phylogenetic principal

component analysis (*p*PCA; Table S1) accounting for adult body mass (Table S2). To

address gaps in the dataset (See Appendix 1), we performed a multiple phylogenetic-

trait imputation for missing values (see Materials and Methods). The two dominant axes

of life history strategies encompass 62% of the total functional variation (Fig. 1). These

axes correspond to (i) the 'fast–slow continuum' (Sæther, 1987; Stearns, 1999; Oli,

2004; Bielby et al 2007), which separates species with short maximum lifespans, like

some freshwater turtles (*e.g.*, 4.3 years for the Alabama red-bellied turtle, *Pseudemys*

alabamensis), to species with long maximum lifespans such as terrestrial tortoises (*e.g.*

177 years for the Pinzon Island giant tortoise, *Chelonoidis ephippium*); and (ii) the

'reproductive strategy' (Gaillard et al., 1989; Salguero-Gómez et al., 2016a; Hughes,

2017; Paniw et al. 2018; Capdevila et al. 2020), defined by a trade-off between clutch

frequency *vs.* reproductive output, with species with high numbers of clutches per year

but lower clutch size per clutch at one end (*e.g.*, 1.2 eggs per clutch and three clutches

per year, as in the Madagascan flat-tailed tortoise, *Pyxis planicauda*), and extremely

fecund species at the other end, having high clutch sizes but few clutches per year (*e.g.,*

91.5 eggs per clutch and one clutch per year, as in the South American River Turtle,

Podocnemis expansa).

Examining the functional spectra of the life history traits of testudines and

crocodilians reveals a high degree of functional redundancy. We use a trait probability

density (TPD) approach considering the two-dimensional functional space (Fig. 1) to

identify peaks (areas of high density) and valleys (low density) of species diversity

(Carmona et al., 2016; Toussaint et al., 2021). The examined reptile species tend to

congregate around a single hotspot with median values of maximum lifespan and high

number of clutches per year with lower clutch sizes. This hotspot of life history

strategies includes 50% of the 259 species and represents 13.2% of the total spectrum

(Fig. 1), in agreement with a recent analyses examining 6,567 species of reptiles (12.3%

of the spectrum; Carmona et al., 2021). Freshwater turtles, such as the eastern long-

necked turtle, *Chelodina longicollis,* (max. lifespan 37 years, 13.9 eggs/clutch, two

clutches/yr) and crocodiles, such as the mugger crocodile *Crocodylus palustris* (max.

lifespan 31.5 years, 28.75 eggs/clutch, two clutches/yr), are located in that hotspot of

functional diversity. Sea turtles are represented in a valley of functional diversity

(bottom right of Fig. 1), which is characterised by species with higher clutch sizes, few

clutches per year, and higher maximum lifespan (*e.g. Caretta caretta:* max. lifespan 76

years, 115 eggs/clutch, 1.29 clutches/yr). In contrast, terrestrial tortoises are represented

at the top right of Fig. 1, with their typically shorter clutch sizes, high clutches per year,

and longer maximum lifespans (*e.g.*, the Hood Island giant tortoise *Chelonoidis*

hoodensis: max. lifespan 177 years, 10 eggs/clutch, three clutches/yr).

Unique strategies may become extinct

The extinction of threatened species of testudines and crocodilians would result in the

loss of a quarter of the global functional diversity of these taxa. To estimate the effect of

the potential extinctions of threatened species on the existing amount of functional

diversity in these groups, we use the IUCN Red List categories of our species and

simulate an accumulative loss of threatened species (*i.e.*, in the first scenario "-CR", we

remove the *Critically Endangered* species only; in the "-EN" scenario, we remove

Endangered and Critically Endangered species, and so on; see Materials and Methods).

If all threatened species disappeared (*Critically Endangered*, *Endangered,* and

Vulnerable species*;* "-VU" scenario, Fig. 2A), the functional diversity would decrease

by 26.8%.

The loss of functional diversity due to the extinction of threatened species is

greater than expected by chance. We compare the potential loss of functional diversity,

considering the number of threatened species in each extinction scenario, with a

simulated loss of functional diversity where the identity of the threatened species is

randomised within the pool of species. The resulting loss of functional diversity due to

the extinction of threatened species is greater than 95% of randomisations for most

scenarios (Fig. 2A). In the "-CR" scenario, the loss of functional diversity (-12.67%) is

four times greater than in randomisations (-3.82% [-0.4%, -9.69%] 5% and 95% CI).

This finding suggests that most of the Critically Endangered species of testudines and

crocodilians show unique life history strategies, such as the pancake tortoise

(*Malacochersus tornieri* max. lifespan 25.9 years, one egg/clutch, one clutch/yr).

Unique or less redundant species strategies of testudines and crocodilians are

more vulnerable to vanish following the extinction of threatened species. To identify

differences in extinction risk between the life history strategies, we map extinction risk

in the functional trait spectra via generalized additive models (GAMs), using the IUCN

Red List category of each species (Carmona et al. 2021). The position of each species

along the trait spectra is significantly correlated with its extinction risk when

considering high-risk species as Critically Endangered and Endangered ($p = 0.039$; Fig.

2B). Using this grouping, high-risk species show distinct positions in the life history

functional space, with species with lower functional redundancy located around the

periphery and likely being more threatened. Only species located at the centre of the

functional space and those with short maximum lifespan and smaller clutch sizes but

with greater number of clutches per year show a lower probability of being threatened

(Fig. 2B).

Habitat disturbance as the main threat

The life history strategies of Testudines and Crocodilia can predict vulnerability to

extinction due to three –local consumption, disease, and pollution– of the six analysed

threats (Fig. 3). To explore the differential effects of global and local threats on the

functional spectra of testudines and crocodilians, we analyse the probability of being

affected by each threat separately using GAMs (more results in Fig. S1). Local

consumption (Fig. 3A) is associated with increased extinction risk in species with

higher clutch size such as sea turtles (*e.g., Caretta caretta*), and higher maximum

lifespans, such as the radiated tortoise (*Astrochelys radiata*) or the Orinoco crocodile

(*Crocodylus intermedius*). In contrast, diseases are associated to threaten species with

slow life histories (Fig. 3B), such as the desert tortoise (*Gopherus agassizii*). Finally,

pollution primarily is associated to species with higher reproductive output (*i.e.,* higher

clutch sizes), such as freshwater turtles like *Chelydra serpentina* or crocodilians like the

saltwater crocodile, *Crocodylus porosus,* (Fig. 3C).

Among all examined threats, the extinction of species affected by habitat

disturbance results in the worst loss of functional diversity, especially in the northern

hemisphere (Fig 4A). To estimate the effect of specific threats on the loss of functional

diversity, we simulate the extinction of species affected by each threat to calculate the

percentage of loss of functional diversity (Fig. 4B). We compare the potential loss of

functional diversity by each threat, with a simulated scenario where the identity of the

species affected by each threat is randomised within the pool of species. Most of the

effects of loss of functional diversity due to the six threats considered are greater than

expected by random extinctions, especially for climate change, whose effect is beyond

the 95% percentile for random simulations (Fig. 4B). Only habitat disturbance and

diseases show similar or lower values of loss of functional diversity than expected by

random simulations (Fig 4B).

DISCUSSION

Here, we demonstrate that threatened species of turtles and crocodilians show great deal

of functional dispersion. Toussaint et al. (2021), exploring 5,686 species of reptiles,

found a loss of functional diversity linked to the extinction of the threatened species

corresponding to less than 10%. Here, focussing on Testudines and Crocodilia, for

which much information has been missing to date (Conde et al., 2019), we find twice

the loss of functional diversity. By linking this information to their conservation status,

we demonstrate that global and local threats are exposing the functional diversity of

these already highly threatened taxonomic groups to great peril. Using the largest

collection of trait data to date for turtles and crocodilians, we find that the threatened

species show unique life history strategies. Losing these species due to extinction will

result in a greater reduction of functional diversity and potentially severe ecological

consequences (Violle et al., 2017) due to the ecosystem services they provide, such as

the high values of biomass contribution of turtles (*e.g.*, red-eared slider turtle,

Trachemys scripta, Lovich et al., 2018), mineral cycling and bioaccumulation (*e.g*.,

Chelydra serpentine; Hebert et al., 1993), or seed dispersal (*e.g.*, Galapagos tortoises,

Chelonoidis spp; Blake et al., 2012).

Local disturbances such as habitat disturbance and unsustainable local consumption are

the main threats for the functional diversity of the studied species. Habitat disturbances

affect most species of turtles and crocodilians, especially in North America and Europe

(Fig. 4A), independent of their life history strategies. Therefore, habitat protection for

these species is key to preserve the functional diversity (Lourenço-de-Moraes et al.,

2021). In fact, Mittermeier et al. (2015) highlighted that the protection of only 16

hotspot of turtle biodiversity would successfully encompass 83% of these species. On

the other hand, we find that the second local threat for the functional diversity of

Testudines and Crocodiles is unsustainable local consumption. Our results predict the

extinction of species with higher rates of clutch size (*e.g*., such as sea turtles, whose

eggs are consumed in high quantities; Madrigal-Ballestero et al., 2013) and long-lived

species (linked with bigger size, which typically provides more meat; Donlan et al.,

2010). Unsustainable local consumption of turtles is one of the most important threats in

Asia, South America, and Africa (Fig. 4A). For example, in China, one million turtle

farms are the primary purchasers of turtles caught in the wild, including several

threatened species (Shi et al., 2007)

Global threats such as unsustainable or illegal international trade and climate change can also have important effects in reducing the functional diversity of Testudines and Crocodilia. The global trade of these species is the third most important threat for their functional diversity, affecting tortoises and turtles mainly for their use as pets (Luiselli et al., 2016) and crocodilians for their skin (Thorbjarnarson, 1999; Platt and Thorbjarnarson, 2000). On the other hand, although climate change affects fewer species than other threats, turtles and crocodilians with unique life history strategies could disappear according to our projections (Azanza-Ricardo et al., 2017). Indeed, the consequences of climate change for these species are found worldwide, including extreme droughts by compromising the fitness of the populations (Poiani and Johnson, 1991; Rodríguez-Caro et al., 2021), flood nesting sites by rising sea levels (Pike et al., 2015), coral bleaching due to increasing in temperatures, affecting feeding areas of sea turtles (Chaloupka et al., 2008) and the unbalance of sex-ratios in populations of temperature sex-dependent species (Janzen, 1994), such as the threatened leatherback turtle (*Dermochelys coriacea*; Tomillo et al., 2014).

Our study does not capture all human activities with impacts on species' ecological strategies. As such, we deem our findings to be conservative. For instance, although we have summarised the breadth of testudines and crocodilians' threats to extinction, threat interactions, which remain largely unknown in these taxonomic groups, have been reported to have synergistic effects on the loss of species, as was described for mammals (Romero-Muñoz et al., 2020) and birds (Chen et al., 2015). Moreover, examining the ecological functions of each species in higher resolution, by examining their diets or habitat breadth (*i.e.*, generalist–specialist), could improve the available information of ecological function (Cooke et al., 2019).

Protecting threatened species with unique life history strategies is key to

conserve their functional diversity (Violle et al., 2016) and ecosystem services they

provide (Lovich et al., 2018). Our findings support previous studies that suggest that

conservation policies should consider together the taxonomic and functional dimensions

of biodiversity (Carmona et al., 2021) to implement targeted and more effective

conservation actions in the context of a global biodiversity crisis (Toussaint et al.,

2021). The IUCN Red List of Threatened Species may include information about

population trends, distribution area, threats, and abundance. The inclusion of functional

diversity could greatly aid international, national, and local efforts to preserve

functionally unique species, and recently, new indexes have been developed to include

some aspects of functional diversity. The IUCN Green Status of Species (Grace et al.

2021; IUCN 2021), a new part of the Red List assessment, quantifies species recovery

and explicitly includes ecological function (Akçakaya et al., 2020). Incorporating the

functional diversity of life history strategies in conservation assessments constitutes a

promising approach to help prioritise conservation efforts to maintain high levels of

functional diversity against current and future threats.

MATERIALS AND METHODS

Life history traits

To describe the life cycle of species in the orders Testudines and Crocodilia, we used

life history trait data. Life history traits quantify how the life cycle of a species and its

underlying vital rates —survival, development, reproduction— have been shaped by

natural selection to optimise its performance (Roff, 1992). We obtained such traits from

published literature, open access databases, and via direct researcher contributions

(Cayuela et al. 2019). In some cases, life history trait information was derived from the

demographic information of different databases and datasets. These data were obtained

from the COMADRE Animal Matrix Database (Salguero□Gómez et al. 2016b),

DATLife Database (2021), Amniote Life History Database (Myhrvold et al. 2015), and

the reviews by Allen et al. (2017), Pfaller et al. (2018) and Reinke et al. (under review)

(more information about data selection in Appendix S2).

To quantify species' life history strategies, we selected six life history traits that

encompass detailed information regarding the timing, intensity, frequency, and duration

of demographic processes across the life cycle of any species (Stearns, 1992). The life

history traits chosen were adult survival (Sa), juvenile survival (Sj), maximum lifespan

(ML), age at sexual maturity ($L\alpha$), mean number of clutches per year (CN), and clutch

size (CS; Table S1). These traits are well-established metrics in comparative

demographic analyses, and their emerging syndromes typically explain high variance

(~75%) in life history strategies of multicellular organisms (Bielby et al., 2007;

Carmona et al., 2021; Gaillard et al. 1989, 2005; Paniw et al., 2018; Salguero□Gómez

et al. 2016a; Stearns 1992). The data obtained from the aforementioned sources

encompass 259 species with at least one of these six traits (more information about data

collection/estimation in Appendix S2).

Phylogeny and data imputation

To identify and quantify the potential role of evolution in shaping our species' life history strategies, we performed phylogenetic comparative analyses. We used a species‐level phylogenetic tree recently published for Testudines and Crocodilia (Colston et al. 2020) to both impute the missing data and account for the effect of evolutionary constraints in the observed life history strategies. Briefly, to build the phylogenetic tree, 14 mitochondrial loci and six nuclear loci were sampled for 357 species of extant or recently extinct turtle and tortoises, and the 27 species described crocodilian species (Rhodin et al. 2017). Colston et al. (2020) computed Maximum Likelihood trees with RaxML (Stamatakis 2014) and used taxonomic imputation to fill the gaps of 17 out of 384 (<5%) species in the tree. Further details on the methods for tree construction are available in Colston et al. (2020). We selected the demographic information of species when information about the phylogeny is available, to perform phylogenetic comparative analyses (Revell, 2010; Revell, 2012), the final dataset included 259 species of Crocodilia (23) and Testudines (236) with this resource

Demographic data of reptiles are scarce and contain missing values (Conde et al. 2019; Fig S2). As principal component analyses require full data (*i.e.* non-gappy data; Rubin 1976, Nakagawa 2004), we carried out phylogenetic imputations to fill in the gaps in the life history traits for the studied species (see Appendix S2. Missing data). To impute the missing life history traits in our dataset, we used the R package mice (Van Buuren & Groothuis-Oudshoorn, 2011), which uses multiple imputation and the addon-phylomice to include phylogenetic information. This approach has been shown recently to correctly impute missing data in similar datasets better than other available approaches (Murray, 2018). Briefly, this approach uses Fully Conditional Specification

(FCS) of the imputation model, which specifies the multivariate imputation model on a

variable-by-variable basis by a set of conditional densities, one for each incomplete

variable (Van Buuren & Groothuis-Oudshoorn, 2011). As suggested by previous studies

(Janssen et al., 2010), we included three more traits in the imputation analyses in order

to have more robust estimations of the imputed data (we included body weight and body

size, which is available for all the species, and incubation time available for 16% of the

species from Amniote Database). Such as there are some stochasticity in the imputation

method, we then created 40 imputed datasets (the percentage of gaps in the data were

38%) and ran analyses on each separately. Like this approach used the phylogenetic

information, we evaluated the phylogenetic signal. We first estimated the phylogenetic

signal for each trait separately using Pagel's λ, which describes the strength of

phylogenetic relationships on trait evolution under a Brownian motion model

(Freckleton 2000). Pagel's λ ranges between 0, when the patterns in the traits cannot be

explained by the employed phylogeny, and 1 when the observed patterns in traits are

tightly correlated with the placement of species in the phylogeny. Most of the selected

traits showed a strong phylogenetic signal (> 0.8; see Table S3), so we used the imputed

values for traits in further analyses. We evaluated the similarity between imputed and

non-imputed data using density plots for each trait. Moreover, we analysed the position

in the functional spectra using Procrustres analyses between imputed data and a subset

where a maximum of two traits were imputed (see Appendix S2. Imputation validation).

Exploring the relation between the axes of life history strategies

To identify potential differences between the patterns of association among life history

traits for Testudines and Crocodilia species while simultaneously assessing non-

independence of lineages, we used a phylogenetically informed PCA (*p*PCA; Revell,

2009). *p*PCA is a multivariate analysis that reduces the number of variables of interest

due to their plausible correlation. The *p*PCA considers the correlation matrix of species'

traits while accounting for phylogenetic relationships and estimate Pagel's λ. The *p*PCA

was estimated using the R package *phytools* (Revell, 2012), assuming a Brownian

motion model of evolution (Revell, 2010). Life history trait data were log‑transformed

to fulfil normality assumptions of PCA and z‑transformed to mean $= 0$, and $SD = 1$

(Legendre & Legendre, 2021). We used the Kaiser criterion (Kaiser, 1960) after

optimisation through varimax rotation to determine the number of axes necessary to

explain a substantial amount of variation. Namely, the axes we retained had an

associated eigenvalue >1.

As body weight is strongly correlated with life history traits (Gaillard et al., 1989;

Bielby et al. 2007; Capdevila et al., 2020), we accounted for this potential effect in our

multivariate analyses too. Out of the multiple possible ways to account for body size or

weight in life history analyses, we used the residuals of phylogenetic regression

between body size and each trait (Revell, 2009). We collected most of adult body mass

(g) data from the studies by Myhrvold et al. (2015) and Colston et al. (2020), moreover,

we collected the remained data from previous literature for all our 259 species.

Estimating the functional spectra

To describe the probabilistic distribution of the species within the functional spaces, we

used the main axes of functional trait variation by *p*PCA corrected by size (*i.e.,*

residuals of regressions). We performed multivariate kernel density using the '*TPD'* and

'*ks'* R packages (Duong, 2007, 2014; Carmona, 2019; Carmona et al., 2019) to estimate

the functional spectra. The kernel was a multivariate normal distribution for each

species centered in the location of the species in *p*PCA and bandwidth chosen using

unconstrained bandwidth selectors from the '*Hpi*' function in the '*ks*' package (Duong

& Hazelton, 2003). The grouped kernels for all species drive into the continuous TPD

function (Carmona et al., 2017; 2019). According to Carmona et al. (2021) and

Toussaint et al. (2021), we divided the continuous functional space into a two-

dimensional grid composed of 200 equal sized cells per dimension. We estimated the

value of the TPD function for the 40,000 cells. The value of the TPD function represent

the density of species in that particular region of the functional space (i.e., species with

similar life history strategy). Results of TPD were represented graphically with the

contours containing 50, 60, 70, 80, 90, and 99% of the total density of species.

Threats to the studied species

To identify causes of recent declines in Testudines and Crocodilia species we reviewed

published information. The main threats described for these species were i) habitat loss,

fragmentation, and degradation (Luiselli, 2009), ii) over-collection of individuals and

their eggs for food consumption (Gong et al., 2017), iii) unsustainable or illegal

international trade, as well as over-collection for the trade in medicines (Sung & Fong,

2018), iv) climate change (Gibbons et al., 2000); v) emerging diseases (Jacobson et al.,

2014; Tompkins et al., 2015). We collected the information on the threats to each

species from Stanford et al. (2020), Bonin et al. (2006) and IUCN Red List (IUCN,

2020). We did dichotomous variables when one of these threats were mentioned. We

were not able to quantify the effect of the threat, because effect size is only available for

the Critically Endangered species of Stanford et al., (2020). To convert the threat

descriptions of the Red List into our broader categories, we used the following process:

for habitat disturbances we considered the Red List threat classifications "residential &

commercial development", "agriculture & aquaculture" and "natural system

modifications"; for climate change the classification "climate change and extreme

weather"; and for disease the classification "invasive and other problematic species,

genes & diseases".

Effects of extinctions of threatened species on global functional diversity

To identify the loss of functional diversity by threatened species extinctions, we

simulated the disappearance of species according to their Red List category (IUCN,

2020). The species classified among the threatened species categories (i.e. Critically

Endangered (CR), Endangered (EN) and Vulnerable (VU) have a higher risk of

extinction than the rest of species: Near Threatened (NT), Least Concern (LC); the

extinction risk of Data Deficient (DD) species is unknown. In our analysis, we removed

threatened species and estimated the resulting shifts in functional diversity (Toussaint et

al., 2021). Firstly, we removed the species with a higher risk of extinction (-CR), then

we removed successively the species with lower risks (-EN includes the extinction of

species classified as a CR and EN, -VU includes CR, EN and VU, and -NT includes

CR, EN, VU and NT). These extinction scenarios represented a gradient of extinction

risk from the persistence of all species to a more dramatic scenario where all threatened

species (including the NT species) went extinct. We compared the TPD function

considering all the species assessed by IUCN (current spectra of functional diversity),

and the TPD function after removing the species in the different scenarios. We can

compare TPD functions because they are probability density functions, which means

that they integrate to 1 across the whole functional space, regardless of the number of

species considered (Carmona et al. 2016). To reduce the potential effect of outliers in

the functional space, we applied a quantile threshold of 99%. We quantify how much of

the functional spectra is lost after the extinction scenarios by estimating which

functional space cells become empty after extinctions. The difference in the amount of

space occupied before and after extinctions (TPD function of the current spectra – TPD

function after removing the threatened species) represented the loss of functional

diversity for Testudines and Crocodilia. We calculated this loss in the functional space

for all the extinction scenarios (-CR, -EN, -VU, and -NT).

To assess if the impacts of each extinction scenario were different from what would be

expected if extinction risk is not related to life history traits, we also compared the

observed changes in functional diversity to a null model where the extinct species were

randomly selected within the pool of species. For each scenario of extinction risk, we

compared the functional diversity to 999 losses of functional diversity where the same

number of threatened species were randomly selected among the pool of species. This

strategy allowed us to understand whether the extinction in the different scenarios

reduces more or less than expected the functional diversity. For each scenario, we

created 999 TPD functions simulating cases in which the same number of species were

lost at random from the total set of IUCN-assessed species. We compared the 5% and

95% percentile of the random simulation with the value of loss of functional diversity

per each extinction scenario to estimate if the values were significantly different. In this

case, higher than expected reductions in functional diversity would mean that the

species that are going extinct in the considered scenario are relatively unique in terms of

their functional traits, whereas lower than expected reductions would imply that the

species going extinct are mostly functionally redundant.

After examining the loss of functional diversity, we plotted the extinction risk of species

within the functional spaces, using the species assessed by IUCN. Using binomial

smoother-based GAM (Wood, 2006) with the R package *mgcv* (Wood, 2011), we

analysed the relationship between extinction risk (1: high-risk and 0: nonthreatened or

low-risk) and the position in the functional space (PC axes). We then represented the

predictions of the models (including the 95% confidence intervals of the means) to

visually examine how life history strategies affect the probability of species being

threatened (Carmona et al., 2021).

Effects of anthropic threats on global functional diversity

To identify the effect of the different threats in the functional spectra, we simulated

scenarios where extinctions were based upon species reported as affected by specific

threats. We did two comparisons: 1) the TPD function considering all the species

(current spectra of functional diversity) and TPD function after removing the species

affected by specific threat (habitat disturbance, trade, local consumption, climate change

or disease) and, 2) the TPD function considering all the IUCN-assessed species, and the

TPD function after removing the threatened species (CR, EN, or VU) affected by each

specific threat. Using a comparison between TPD functions similar to the one explained

above, we estimated the potential loss of functional diversity attributable to each threat

and evaluated the differences between them. We also compared the observed changes in

functional diversity using a null model to assess if the impacts of each extinction

scenario were different from what would be expected by chance. For each threat, we

compared the loss of functional diversity with the 999 losses of functional diversity

where the same number of species were randomly selected among the pool the species.

We did these comparisons both considering the extinction of all the species affected by

each threat and considering only the extinction of threatened species.

To visualise the relationship between life history strategies and threats, we mapped the

probability of a species being affected by a specific threat within the functional spaces.

We used binomial smoother-based GAM (Wood, 2006) with the R package *mgcv*

(Wood, 2011) to identify the relationship between being affected by each threat (1:

affected and 0: nonaffected) and the position in the functional space (PCA axes). We

represented the predictions of the models (including the 95% confidence intervals of the

means) when the relation between threat and life history strategy was significant.

ACKNOWLEDGEMENTS

RCRC was supported by a post☐doctoral grant funded by the Regional Valencian

Government (APOSTD/2020/090) hosted by RSG. RSG was supported by a NERC IRF

grant (NE/M018458/1). RCRC, EG, and AG were supported by the Spanish Ministry of

Science and European Regional Development Fund (PID2019-105682RA-

I00/AEI/10.13039/501100011033). CPC was supported by the Estonian Research

Council (PSG293). Illustrations by Carmen Cañizares (@canitailustradora)

REFERENCES

Allen, W. L., Street, S. E., & Capellini, I. (2017). Fast life history traits promote
invasion success in amphibians and reptiles. Ecology Letters, 20(2), 222-230.

Akçakaya et al. (2020). Assessing ecological function in the context of species
recovery. Conservation Biology, 34(3), 561-571.

Azanza-Ricardo, J., Martín, M. E. I., Sansón, G. G., Harrison, E., Cruz, Y. M., &
Bretos, F. (2017). Possible effect of global climate change on *Caretta caretta*

(Testudines, Cheloniidae) nesting ecology at Guanahacabibes Peninsula, Cuba.

Chelonian conservation and Biology, 16(1), 12-19.

Bielby, J., Mace, G. M., Bininda‑Emonds, O. R. P., Cardillo, M., Gittleman, J.

L., Jones, K. E., … Purvis, A. (2007). The fast‑slow continuum in mammalian

life history: An empirical reevaluation. The American Naturalist, 169(6), 748–

757.

Blake, S., Wikelski, M., Cabrera, F., Guezou, A., Silva, M., Sadeghayobi, E., ... &

Jaramillo, P. (2012). Seed dispersal by Galápagos tortoises. Journal of

Biogeography, 39(11), 1961-1972.

Bonin, F., Devaux, B., & Dupré, A. (2006). Turtles of the World. JHU Press.

Capdevila, P., Beger, M., Blomberg, S. P., Hereu, B., Linares, C., & Salguero‑Gómez,

R. (2020). Longevity, body dimension and reproductive mode drive differences

in aquatic versus terrestrial life‑history strategies. Functional Ecology, 34(8),

1613-1625.

Carmona, C. P., De Bello, F., Mason, N. W., & Lepš, J. (2016). Traits without borders:

integrating functional diversity across scales. Trends in ecology & evolution,

31(5), 382-394.

Carmona, C. P. (2019). TPD: Methods for Measuring Functional Diversity Based on

Trait   Probability   Density.   https://cran.r-

project.org/web/packages/TPD/index.html.

Carmona, C. P., de Bello, F., Mason, N. W., & Lepš, J. (2019). Trait probability density

(TPD): measuring functional diversity across scales based on TPD with R.

Ecology, 100(12), e02876.

Carmona, C. P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., ... & Toussaint, A. (2021). Erosion of global functional diversity across the tree of life. Science Advances, 7(13), eabf2675.

Caswell, H. (2001). Matrix population models: Construction, analysis, and interpretation (2nd ed.). Sunderland, MA: Sinauer Associates.

Cayuela, H., Akani, G. C., Hema, E. M., Eniang, E. A., Amadi, N., Ajong, S. N., ... & Luiselli, L. (2019). Life history and age-dependent mortality processes in tropical reptiles. Biological Journal of the Linnean Society, 128(2), 251-262.

Cernansky, R. (2017). The biodiversity revolution. Nature 546, 22–24.

Chaloupka, M., Kamezaki, N., & Limpus, C. (2008). Is climate change affecting the population dynamics of the endangered Pacific loggerhead sea turtle?. Journal of Experimental Marine Biology and Ecology, 356(1-2), 136-143.

Chen, S., Fan, Z., Roby, D. D., Lu, Y., Chen, C., Huang, Q., ... & Zhu, J. (2015). Human harvest, climate change and their synergistic effects drove the Chinese Crested Tern to the brink of extinction. Global Ecology and Conservation, 4, 137-145.

Chichorro, T., Juslén, A., Cardoso, P. (2019). A review of the relation between species traits and extinction risk. Biol. Conserv. 237, 220–229.

Colston, T. J., Kulkarni, P., Jetz, W., & Pyron, R. A. (2020). Phylogenetic and spatial distribution of evolutionary diversification, isolation, and threat in turtles and crocodilians (non-avian archosauromorphs). BMC Evolutionary Biology, 20(1), 1-16.

Conde, D. A., Staerk, J., Colchero, F., da Silva, R., Schöley, J., Baden, H. M., ... & Vaupel, J. W. (2019). Data gaps and opportunities for comparative and

conservation biology. Proceedings of the National Academy of Sciences, 116(19), 9658-9664.

Cooke, R. S., Eigenbrod, F., & Bates, A. E. (2019). Projected losses of global mammal and bird ecological strategies. Nature Communications, 10(1), 1-8.

DATLife Database. (2021). Max-Planck Institute for Demographic Research (Germany). Available at www.datlife.org (data downloaded on [25-02-2021]

Díaz, S., Settele, J., Brondízio, E. S., Ngo, H. T., Agard, J., Arneth, A., ... & Zayas, C. N. (2019). Pervasive human-driven decline of life on Earth points to the need for transformative change. Science, 366(6471).

Donlan, C. J., Wingfield, D. K., Crowder, L. B., & Wilcox, C. (2010). Using expert opinion surveys to rank threats to endangered species: a case study with sea turtles. Conservation Biology, 24(6), 1586-1595.

Duong, T. (2007). Ks: Kernel density estimation and kernel discriminant analysis for multivariate data in R. Journal of Statistical Software, 21(7), 1-16.

Duong, T. (2014). ks: Kernel Smoothing. R Package Version 1.9.2 (R package version 1.11.4). https://CRAN.R-project.org/package=ks.

Duong, T., & Hazelton, M. (2003). Plug-in bandwidth matrices for bivariate kernel density estimation. Journal of Nonparametric Statistics, 15(1), 17-30.

Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., & Norberg, J. (2003). Response diversity, ecosystem change, and resilience. Frontiers in Ecology and the Environment, 1(9), 488-494.

Freckleton, R. P. (2000). Phylogenetic tests of ecological and evolutionary hypotheses: Checking for phylogenetic independence. Functional Ecology, 14(1), 129–134.

Gaillard, J. M., Pontier, D., Allaine, D., Lebreton, J. D., Trouvilliez, J., & Clobert, J. (1989). An analysis of demographic tactics in birds and mammals. Oikos, 59-76.

Gaillard, J. M., Yoccoz, N G, Lebreton, J. D., Bonenfant, C., Devillard, S., Loison,

A., … Allaine, D. (2005). Generation time: A reliable metric to measure

life history variation among mammalian populations. The American

Naturalist, 166(1), 119–123. https://doi.org/10.1086/430330

Germano, D. J. (1994). Comparative life histories of North American tortoises. Biology

of North American Tortoises, 175-185.

Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S.,

Greene, J.L., Mills, T., Leiden, Y., Poppy, S., and Winne, C.T. (2000). The

global decline of reptiles, déja vu amphibians. Bioscience 50, 653–666.

Gibbs, J. P., & Amato, G. D. (2000). Genetics and demography in turtle conservation.

Turtle conservation, 207-217.

Gong, S., Shi, H., Jiang, A., Fong, J.J., Gaillard, D., and Wang, J. (2017).

Disappearance of endangered turtles within China's nature reserves. Curr. Biol.

27, 170–171.

Grace, M. K., Akçakaya, H. R., Bennett, E. L., Brooks, T. M., Heath, A., Hedges, S., ...

& Quintana, M. G. (2021). Testing a global standard for quantifying species

recovery and assessing conservation impact. Conservation Biology, 35(6), 1833-

1849.

Healy, K., Ezard, T. H., Jones, O. R., Salguero-Gómez, R., & Buckley, Y. M. (2019).

Animal life history is shaped by the pace of life and the distribution of age-

specific mortality and reproduction. Nature ecology & evolution, 3(8), 1217-

1224.

Hebert, C. E., Glooschenko, V., Haffner, G. D., & Lazar, R. (1993). Organic

contaminants in snapping turtle (*Chelydra serpentina*) populations from

southern Ontario, Canada. Archives of Environmental contamination and
Toxicology, 24(1), 35-43.

Honaker, J., King, G., & Blackwell, M. (2011). Amelia II: A program for missing data.
Journal of statistical software, 45(1), 1-47.

Hughes, P. W. (2017). Between semelparity and iteroparity: empirical evidence for a
continuum of modes of parity. Ecology and evolution, 7(20), 8232-8261.

IUCN. (2020). The IUCN Red List of Threatened Species. Version 2020–3; 2020.

IUCN (2021). IUCN Green Status of Species: a global standard for measuring species
recovery and assessing conservation impact. IUCN, Gland, Switzerland.
https://portals.iucn.org/library/node/49511

Jackson, S. T., & Sax, D. F. (2010). Balancing biodiversity in a changing environment:
extinction debt, immigration credit and species turnover. Trends in ecology &
evolution, 25(3), 153-160.

Jacobson, E.R., Brown, M.B., Wendland, L.D., Brown, D.R., Klein, P.A., Christopher,
M.M., and Berry, K.H. (2014). Mycoplasmosis and upper respiratory tract
disease of tortoises: a review and update. Vet. J. 201, 257–264.

Janssen, K. J., Donders, A. R. T., Harrell Jr, F. E., Vergouwe, Y., Chen, Q., Grobbee, D.
E., & Moons, K. G. (2010). Missing covariate data in medical research: to
impute is better than to ignore. Journal of clinical epidemiology, 63(7), 721-727.

Janzen, F. J. (1994). Climate change and temperature-dependent sex determination in
reptiles. Proceedings of the National Academy of Sciences, 91(16), 7487-7490.

Kaiser, H. F. (1960). The varimax criterion for analytic rotation in factor analysis.
Psychometrika, 23, 187–200.

Kuussaari, M., Bommarco, R., Heikkinen, R. K., Helm, A., Krauss, J., Lindborg, R., ...

634     & Steffan-Dewenter, I. (2009). Extinction debt: a challenge for biodiversity

635     conservation. Trends in ecology & evolution, 24(10), 564-571.

636 Legendre, P., & Legendre, L. (2012). Numerical ecology (3rd ed.). Amsterdam, the

637     Netherlands: Elsevier Science.

638 Lovich, J. E., Ennen, J. R., Agha, M., & Gibbons, J. W. (2018). Where have all the

639     turtles gone, and why does it matter?. BioScience, 68(10), 771-781.

640 Lourenço-de-Moraes, R., Campos, F. S., Carnaval, A. C., Otani, M., França, F. G.,

641     Cabral, P., & Benedito, E. (2021). No more trouble: An economic strategy to

642     protect taxonomic, functional and phylogenetic diversity of continental turtles.

643     Biological Conservation, 261, 109241.

644 Luiselli, L. (2009). A model assessing the conservation threats to freshwater turtles of

645     Sub-Saharan Africa predicts urgent need for continental conservation planning.

646     Biodivers. Conserv. 18, 1349–1360.

647 Luiselli, L., Starita, A., Carpaneto, G. M., Segniagbeto, G. H., & Amori, G. (2016). A

648     short review of the international trade of wild tortoises and freshwater turtles

649     across the world and throughout two decades. Chelonian Conservation and

650     Biology, 15(2), 167-172.

651 Madrigal-Ballestero, R., Schlüter, A., & Lopez, M. C. (2013). What makes them follow

652     the rules? Empirical evidence from turtle egg harvesters in Costa Rica. Marine

653     Policy, 37, 270-277.

654 Maxwell, S. L., Fuller, R. A., Brooks, T. M., & Watson, J. E. (2016). Biodiversity: The

655     ravages of guns, nets and bulldozers. Nature News, 536(7615), 143.

656 Mittermeier, R. A., van Dijk, P. P., Rhodin, A. G., & Nash, S. D. (2015). Turtle

657     hotspots: an analysis of the occurrence of tortoises and freshwater turtles in

biodiversity hotspots, high-biodiversity wilderness areas, and turtle priority

areas. Chelonian Conservation and Biology, 14(1), 2-10.

Murray, J. S. (2018). Multiple imputation: a review of practical and theoretical findings.

Statistical Science, 33(2), 142-159.

Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. M.

(2015). An amniote life history database to perform comparative analyses with

birds, mammals, and reptiles: Ecological Archives E096 269. Ecology, 96(11),

3109-3109.

Naeem, S. (1998). Species redundancy and ecosystem reliability. Conservation biology,

12(1), 39-45.

Nakagawa S (2004) A farewell to Bonferroni: the problems of low statistical power and

publication bias. Behav Ecol 15:1044–1045

Oli, M. K. (2004). The fast-slow continuum and mammalian life-history patterns: An

Empirical evaluation. Basic Appl Ecol 5(5):449–463.

Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., … &

Bullock, J. M. (2015). Biodiversity and resilience of ecosystem functions.

Trends in ecology & evolution, 30(11), 673-684.

Paniw, M., Ozgul, A., & Salguero-Gómez, R. (2018). Interactive life-history traits

predict sensitivity of plants and animals to temporal autocorrelation. Ecology

Letters, 21(2), 275–286.

Pfaller, J. B., Chaloupka, M., Bolten, A. B., & Bjorndal, K. A. (2018). Phylogeny,

biogeography and methodology: a meta-analytic perspective on heterogeneity in

adult marine turtle survival rates. Scientific reports, 8(1), 1-10.

Pike, D. A., Roznik, E. A., & Bell, I. (2015). Nest inundation from sea-level rise threatens sea turtle population viability. Royal Society Open Science, 2(7), 150127.

Pimm, S. L., Russell, G. J., Gittleman, J. L., & Brooks, T. M. (1995). The future of biodiversity. Science, 269(5222), 347-350.

Platt, S. G., & Thorbjarnarson, J. B. (2000). Population status and conservation of More'et's crocodile, *Crocodylus moreletii*, in northern Belize. Biological Conservation, 96(1), 21-29.

Poiani, K. A., & Johnson, W. C. (1991). Global warming and prairie wetlands. BioScience, 41(9), 611-618.

Reinke, B. A., Cayuela, H., Janzen F. J., Lemaitre, J. F., Gaillard, J. M., … & Miller, D. Under review. A. W. Under review. Diverse aging rates in ectothermic tetrapods provide insights for the evolution of aging and longevity.

Revell, L. J. (2009). Size correction and principal components for interspecific comparative studies. Evolution, 63(12),

Revell, L J. (2010). Phylogenetic signal and linear regression on species data. Methods in Ecology and Evolution, 1(4), 319–329.

Revell, L. J. (2012). *phytools*: An R package for phylogenetic comparative biology (and other things). Methods in Ecology and Evolution, 3(2), 217–223.

Rhodin AGJ, Iverson JB, Bour R, Fritz U, Georges A, Shaffer HB, et al. (2017). Turtles of the world. Annotated checklist and atlas of taxonomy, synonymy, distribution, and conservation status. 7th ed.

Rhodin, A. G., Stanford, C. B., Van Dijk, P. P., Eisemberg, C., Luiselli, L., Mittermeier, R. A., ... & Vogt, R. C. (2018). Global conservation status of turtles and tortoises (order Testudines). Chelonian Conservation and Biology, 17(2), 135-161.

Ripple, W. J., Wolf, C., Newsome, T. M., Hoffmann, M., Wirsing, A. J., & McCauley,
D. J. (2017). Extinction risk is most acute for the world's largest and smallest
vertebrates. Proceedings of the National Academy of Sciences, 114(40), 10678-
10683.

Rodríguez☐Caro, R. C., Capdevila, P., Graciá, E., Barbosa, J. M., Giménez, A., &
Salguero☐Gómez, R. (2021). The limits of demographic buffering in coping
with environmental variation. Oikos, 130: 1346–1358

Roff, D. (1993). Evolution of life histories: theory and analysis. Springer Science &
Business Media.

Romero☐Muñoz, A., Benítez☐López, A., Zurell, D., Baumann, M., Camino, M.,
Decarre, J., ... & Kuemmerle, T. (2020). Increasing synergistic effects of habitat
destruction and hunting on mammals over three decades in the Gran Chaco.
Ecography, 43(7), 954-966.

Roper, M., Capdevila, P., & Salguero-Gomez, R. (2021). Senescence: why and where
selection gradients might not decline with age. Proceedings of the Royal Society
B: Biological Sciences.

Rubin DB (1976) Inference and missing data. Biometrika 63:581–590

Sæther. B. E. (1987). The influence of body weight on the covariation between
reproductive traits in European birds. Oikos 48(1):79–88.

Salguero☐Gómez, R., Jones, O. R., Archer, C. R., Bein, C., deBuhr, H., Farack, C.,
… Vaupel, J. W. (2016b). COMADRE: A global data base of animal
demography. Journal of Animal Ecology, 85(2), 371–
384. https://doi.org/10.1101/027821

Salguero☐Gómez, R., Jones, O. R., Jongejans, E., Blomberg, S. P., Hodgson, D.
J., Mbeau☐Ache, C., … Buckley, Y. M. (2016a). Fast–slow continuum and

reproductive strategies structure plant life￼history variation worldwide. Proceedings of the National Academy of Sciences of the United States of America, 113(1), 230–235.

Scott, R., Marsh, R., & Hays, G. C. (2012). Life in the really slow lane: loggerhead sea turtles mature late relative to other reptiles. Functional Ecology, 26(1), 227-235.

Shi, H., Parham, J. F., Lau, M., & Tien-Hsi, C. (2007). Farming endangered turtles to extinction in China. Conservation Biology, 21(1), 5-6.

Stanford, C. B., Iverson, J. B., Rhodin, A. G., van Dijk, P. P., Mittermeier, R. A., Kuchling, G., ... & Walde, A. D. (2020). Turtles and tortoises are in trouble. Current Biology, 30(12), R721-R735.

Stamatakis A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics, 30:1312–3.

Stearns, S. C. (1999). The evolution of life histories. New York, NY: Oxford University Press.

Sung, Y.-H., and Fong, J.J. (2018). Assessing conservation trends and illegal activity by monitoring the online wildlife trade. Biological Conservation. 227, 219–225.

Thorbjarnarson, J. (1999). Crocodile tears and skins: international trade, economic constraints, and limits to the sustainable use of crocodilians. Conservation Biology, 13(3), 465-470.

Tomillo, P. S., Oro, D., Paladino, F. V., Piedra, R., Sieg, A. E., & Spotila, J. R. (2014). High beach temperatures increased female-biased primary sex ratios but reduced output of female hatchlings in the leatherback turtle. Biological Conservation, 176, 71-79.

Tompkins, D.M., Carver, S., Jones, M.E., Krosek, M., and Skerrat, L.F. (2015).
Emerging infectious diseases of wildlife: a critical perspective. Trends in
Parasitology. 31, 149–159.

Toussaint, A., Brosse, S., Bueno, C. G., Pärtel, M., Tamme, R., & Carmona, C. P.
(2021). Extinction of threatened vertebrates will lead to idiosyncratic changes in
functional diversity across the world. Nature Communications, 12(1), 1-12.

Van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by
chained equations in R. Journal of statistical software, 45(1), 1-67.

Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N. J., Cadotte, M. W.,
Livingstone, D. & Mouillot, D. (2017). Functional rarity: the ecology of outliers.
Trends in Ecology & Evolution, 32(5), 356-367.

Wood, S. (2006). Generalized Additive Models: An Introduction with R (CRC Press);
https://books.google.co.uk/books/about/Generalized_Additive_Models.html?id=
hr17lZC-3jQC&pgis=1.

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. J. R. Stat. Soc. 73, 3–
36.

 Figures

Fig. 1. The global functional spectra of the life history strategies of Testudines and

Crocodilia is described by two dominant axes of life history variation: (PC1) fast-

slow continuum and (PC2) reproductive strategies. The probabilistic species

distribution in this space is defined by the two first principal components axes (PC1 =

39.1% and PC2 = 23.1% of variance explained) of a phylogenetically-corrected

principal component analysis (*p*PCA), where adult body mass has also been corrected.

The phylogenetic signal is Pagel's $\lambda = 0.674 \pm 0.030$ (sd). The life history traits are:

adult survival (*Sa*), juvenile survival (*Sj*), maximum lifespan (*ML*), age at sexual

maturity (*Lα*), mean of number of clutches per year (*CN*), clutch size (*CS*). Arrows

indicate the direction and weighting of each trait in the *p*PCA. The colour gradient

(orange, yellow, and white) depicts the density of species in the defined space, where

orange corresponds to more densely populated areas). Thick contour lines indicate the

0.5 and 0.99 quantiles, and thinner ones indicate quantiles 0.6, 0.7, 0.8, and 0.9. The

species silhouettes correspond to (starting at the top left and moving clockwise): painted

turtle (*Chrysemys picta*), American alligator (*Alligator mississippiensis*), desert tortoise

(*Gopherus agassizii*), Hood Island giant tortoise (*Chelonoidis hoodensis*), Orinoco

crocodile (*Crocodylus intermedius*), loggerhead sea turtle (*Caretta caretta*), leatherback

sea turtle (*Dermochelys coriacea*), northern map turtle (*Graptemys geographica*), and

Western Caspian turtle (*Mauremys rivulata*).

Fig. 2. The life history strategies of testudines and crocodilians predict their vulnerability to extinction due to human threats. A) Loss of functional diversity under extinction scenarios of threatened species. CR: Critically Endangered, EN: Endangered, VU: Vulnerable; NT: Near Threatened. The loss of functional diversity is expressed as a percentage of the current functional diversity in the life history strategies of species of the order Testudines and Crocodilia (Fig. 1). We simulate the loss of functional diversity by removing all species from the IUCN category in a progressive manner, by degree of "endangerment", removing first the species with a higher risk of extinction (*i.e.* -CR), then continuing progressively to remove species from the categories with lower threatened risks (-EN scenario: removing CR and EN species; -VU scenario: removing CR, EN & VU; -NT scenario: removing CR, EN, VU & NT). This loss of diversity is represented by a black line. For each scenario, we compare the loss of functional diversity with 999 iterations of a null model where threatened species are randomly selected among the 259 species in the analyses. The 999 randomisations are represented by a blue polygon defined by the 5th and 95th percentiles of functional diversity loss, with the blue line representing the 50th percentile; **B**) Risk of extinction in the functional space of Testudines and Crocodilia species. Probability of species being classified as high-risk of extinction (*i.e.*, CR & EN IUCN status) according to GAMs (with binomial distribution) using the position of each species in the two-dimensional functional space as predictors. Green tones indicate lower risk of extinction, whereas red tones indicate higher risk of extinction. *P* value of the GAM model is 0.03 and χ^2 is 28.59. The red contour lines indicate the average threat probability (proportion of species classified as threatened in the group). Here, we consider only the 214 out of the 259 available species of testudines and crocodilians for which the threat status is known.

Fig. 3. The life history strategies of Testudines and Crocodilia predict the

vulnerability to extinction due to local consumption, disease, and pollution. Other

risks, such as climate change, global trade, or habitat disturbance, do not show a

significant relation with the examined species' life history strategies (Fig. S1).

Probability of species being affected by: **A**) local consumption, **B**) disease, and **C**)

pollution, according to GAMs (with binomial distribution) using the position of species

in the two-dimensional functional space (Fig. 1) as predictors. Light tones indicate

lower risk of extinction due to each threat, whereas dark tones indicate higher risk. *P*

values associated with each GAM are shown at the top-right corner of each panel. The

red contour lines indicate the average threat probability. Here, we consider only species

whose threats are known (n = 251 species).

Fig. 4. Habitat disturbance is the main threat for the functional diversity,

especially in the north hemisphere. A) Ratio of species affected by each threat type

per continent. The ratio is only estimated for species whose threats are known at the

continent level (North America, n = 65; Europe, n = 4; Asia, n = 66; South America, n =

57; Africa, n = 39; Oceania, n = 14). **B**) Simulated loss of functional diversity of

Testudines and Crocodilia under extinction scenarios by threat. The loss of functional

diversity is expressed as a percentage of the total spectra of functional diversity of

species, as detailed in Fig. 1. We simulate the loss of functional diversity by removing

all species affected by specific threats, in light blue colour removing only threatened

species (*i.e.* Critically Endangered [CR], Endangered [EN] & Vulnerable [VU] as per

the IUCN Red List) and in dark blue removing all the species affected (threatened or

not). For each scenario, we compare the loss of functional diversity with 999 iterations

of a null model where the same number of species were randomly selected among all

251 species. The 999 randomisations are represented for each threat as a grey dot for the

50[th] percentile, with grey whiskers representing the 5[th] and 95[th] percentiles.