In vitro susceptibility to beta-lactam antibiotics and viability of *Neisseria gonorrhoeae* strains producing plasmid-mediated broad- and extended-spectrum beta-lactamases

Ilya Kandinov[1*], Dmitry Gryadunov[1], Alexandra Vinokurova[1], Olga Antonova[1], Alexey
Kubanov[2], Victoria Solomka[2], Julia Shagabieva[2], Dmitry Deryabin[2] and Boris Shaskolskiy[1]

[1]Center for Precision Genome Editing and Genetic Technologies for Biomedicine, Engelhardt
Institute of Molecular Biology, Russian Academy of Sciences, 119991 Moscow, Russia

[2]State Research Center of Dermatovenerology and Cosmetology, Russian Ministry of Health, 107076
Moscow, Russia

* Correspondence:
Ilya Kandinov
ilya9622@gmail.com

Keywords: *Neisseria gonorrhoeae*, beta-lactamase-producing plasmids, *N. gonorrhoeae*
viability, extended-spectrum beta-lactamase

ABSTRACT

N. gonorrhoeae strains producing plasmid-mediated broad- and extended-spectrum beta-lactamases
(ESBLs) have been obtained *in vitro* and studied for their viability and susceptibility to beta-lactam
antibiotics. The artificial $pbla_{TEM-1}$ and $pbla_{TEM-20}$ plasmids were constructed by site-directed
mutagenesis from a $pbla_{TEM-135}$ plasmid of the Toronto/Rio type, which was extracted from the
clinical isolate. MIC values were determined for a series of beta-lactam antibiotics, including
benzylpenicillin, ampicillin, cefuroxime, ceftriaxone, cefixime, cefotaxime, cefepime, meropenem,
imipenem, and doripenem. The *N. gonorrhoeae* strain carrying the $pbla_{TEM-20}$ plasmid exhibited a
high level of resistance to penicillins and II-IV generation cephalosporins (MIC $\geq$ 2 mg/L) but not to
carbapenems (MIC $\leq$ 0.008 mg/L). However, this strain stopped growing after 6 hours of cultivation.
The reduced viability was not associated with the plasmid loss but can be explained by the following
factors: the presence of the plasmid itself, which requires additional costs for its reproduction, and
the expression of ESBL, which can affect the structure of the peptidoglycan layer in the cell
membrane. The cell growth was mathematically modeled using the generalized Verhulst equation.
The parameter, characterizing the maximum possible number of cells grown under given conditions,
decreased within the wild type (without plasmids) - $pbla_{TEM-135}$ - $pbla_{TEM-1}$ - $pbla_{TEM-20}$ series, *i.e.*, the
plasmid-bearing strains had reduced viability compared to that of the wild-type strain. The kinetics
for the cell death of *N. gonorrhoeae* strains without the $pbla_{TEM-20}$ plasmid in the presence of
ceftriaxone can be described by a modified Chick-Watson law. For the *N. gonorrhoeae* strain that
harbors the $pbla_{TEM-20}$ plasmid, the kinetics for cell number changes with time reflected several
processes: the hydrolysis of ceftriaxone by TEM-20 beta-lactamase, growth and gradual death of
cells. The demonstrated reduced viability of *N. gonorrhoeae* strains with the $pbla_{TEM-20}$ plasmid
probably explains the absence of clinical isolates of *N. gonorrhoeae* that produce ESBL.

INTRODUCTION

The development of multidrug resistance in the *N. gonorrhoeae* pathogen is a major problem worldwide. According to the World Health Organization (WHO), gonorrhea may become incurable due to the ineffectiveness of old antimicrobial preparations (AMPs) and the lack of new AMPs for its treatment (Tacconelli et al., 2018; Golparian and Unemo, 2021).

Currently, and possibly in the future, antibiotics of the beta-lactam group are the preferred drugs for treating gonococcal infection (Workowski and Bolan, 2015; WHO Guidelines, 2016; Tacconelli et al., 2018; Unemo et al., 2020). The group includes penicillin, cephalosporin, carbapenem, and monobactam subgroups. The mechanism of action of beta-lactam antibiotics is to disrupt the synthesis of the peptidoglycan layer in bacterial cell walls, leading to bacterial death. In the mid-40s of the 20th century, penicillins started a revolution in the treatment of gonococcal infection and became the gold standard of treatment — a cure after a single injection. However, by the end of the 20th century, most clinical isolates of *N. gonorrhoeae* showed reduced susceptibility or resistance to penicillin, and for this reason, the use of penicillin was discontinued (Unemo and Shafer, 2014; Unemo and Jensen, 2017).

Now, for the treatment of gonococcal infections, the drugs of choice include another group of beta-lactams, that is, third-generation cephalosporins, involving ceftriaxone, cefotaxime and cefixime, which are used as a monotherapy or in combination with the macrolide antibiotic azithromycin (Workowski and Bolan, 2015; WHO Guidelines, 2016; Tacconelli et al., 2018; Unemo et al., 2020). However, the levels of gonococcal resistance to cephalosporins are increasing every year, and these drug treatments were already reported to fail in several countries (Unemo et al., 2012; Allen et al., 2013; Golparian et al., 2014; Attram et al, 2019). The next group of beta-lactams that could replace cephalosporins if cephalosporin resistance spreads is the carbapenem group (Unemo et al., 2020).

The resistance of *N. gonorrhoeae* to beta-lactam antibiotics is associated with both chromosomal and plasmid determinants. Chromosomal mutations include substitutions in penicillin-binding proteins 1 and 2 (PBP1 and PBP2), mutations in the porin protein PorB that lead to a change in cell membrane permeability, and mutations that cause an increase in the expression level of MtrCDE efflux pump (Unemo and Shafer, 2014; Unemo and Jensen, 2017; Unemo et al., 2019). In this resistance phenomenon, the presence of the beta-lactamase enzyme, which hydrolyzes the C-N bond in the beta-lactam ring of the antibiotic and inactivates the action of AMP, is particularly important. The proportion of *N. gonorrhoeae* clinical isolates carrying pbla_{TEM} plasmids is not very high; for example, in the Russian population, the proportion remains at ~5% of the total number of isolates (Kubanov et al., 2019). However, the presence of beta-lactamase in gonococci causes a significant increase in the level of resistance to penicillins ($MIC_{pen} \geq 16$ mg/L) compared to chromosomal mutations ($MIC_{pen} = 0.12$-1.0 mg/L) (Shaskolskiy et al., 2019). To date, clinical isolates of *N. gonorrhoeae* have produced only enzyme variants that belong to broad-spectrum beta-lactamases (penicillinases), which are unable to hydrolyze cephalosporins or carbapenems.

The following types of pbla_{TEM} plasmids are known for gonococci: Asian (7426 bp), African (5599 bp), Toronto/Rio (5154 bp), Nimes (6798 bp), New Zealand (9309 bp), Johannesburg (4865 bp) and Australian (3269 bp) (Müller et al., 2011). The plasmid *bla* gene has a length of 861 bp. Most penicillinase-producing isolates of *N. gonorrhoeae* contain the plasmid with the bla_{TEM-1} gene, but recently, the $bla_{TEM-135}$ variant has been increasingly found in the worldwide population of *N. gonorrhoeae* (Ohnishi et al., 2010; Nakayama et al., 2012 Muhammad et al., 2014; Cole et al., 2015; Tanaka et al., 2021). The bla_{TEM-1} gene was found in both African and Asian plasmids, while $bla_{TEM-135}$ was present mainly in Toronto/Rio and Asian type plasmids (Cole et al., 2015; Yan et al., 2019). $bla_{TEM-135}$ differs from bla_{TEM-1} in one thymine-to-cytosine substitution at codon 182 (ATG→ACG). The resulting Met182Thr mutation is located far from the enzymatic active site (17 Å) in the hinge

region between two beta-lactamase domains and leads to stabilization of the beta-lactamase structure (Orencia et al., 2001). As previously noted (Cole, 2015; Yan et al., 2019), the increased enzymatic stability may have contributed to the fixation of the TEM-135 beta-lactamase variant in the *N. gonorrhoeae* population.

There is a large group of enzymes that are related to bacterial beta-lactamases. TEM-1 and TEM-135 belong to class A serine beta-lactamases according to the Ambler classification (Bradford, 2001; Bush and Jacoby, 2010; Tooke et al., 2019). This class also includes extended-spectrum beta-lactamases (ESBLs), such as TEM-20, which are capable of hydrolyzing both penicillins and cephalosporins. Just one GGT→AGT nucleotide substitution in the $bla_{TEM-135}$ gene, which results in the Gly238Ser mutation, enables gonococci to express TEM-20 type ESBLs (Arlet et al., 1999; Bradford, 2001). The TEM-20 beta-lactamase variant already exists in nature. For example, in *Escherichia coli* (Adator et al., 2020), this variant resulted in increased MIC levels of cephalosporins (Jeong et al., 2004). The Gly238Ser substitution is associated with a change in the beta-lactamase conformation, which increases the flexibility of the enzyme active site, and as a result, beta-lactamase gains the ability to bind cephalosporins without losing its penicillin-hydrolyzing character (Orencia et al., 2001; Yan et al., 2019). Due to the increasing use of carbapenems in medicine, beta-lactamases have emerged that can hydrolyze carbapenems; for example, Ambler class A *Klebsiella pneumoniae* carbapenemase (KPC) and Guiana extended-spectrum beta-lactamase (GES) can be both chromosomal- and plasmid-localized (Tooke et al., 2019; Sawa et al., 2020). Thus, there is a risk that ESBLs may appear in gonococci, which could end the use of third-generation cephalosporins as treatments.

The goal of this work was to construct pbla_{TEM} plasmids that contain different variants of the *bla* gene, to prepare genetically engineered *N. gonorrhoeae* strains that contain pbla_{TEM} plasmids, to study the properties of *N. gonorrhoeae* strains that are transformed with the plasmids, and to assess the viability and resistance of these strains to beta-lactam antibiotics (penicillins, cephalosporins and carbapenems). Particular attention was placed on constructing the pbla_{TEM-20} plasmid and studying the properties of the *N. gonorrhoeae* strain carrying this plasmid, which was not previously found in nature. This research will allow for the assessment of risks that are associated with the emergence and spread of third-generation cephalosporin-resistant strains and the possibility of further using beta-lactam drugs for the treatment of gonococcal infection.

MATERIALS AND METHODS

All experiments with *N. gonorrhoeae*, including transforming the cells, isolating plasmid DNA, electroporating the cells, cultivating the cells on solid media, washing the cells from Petri dishes and passaging them to other dishes, counting the grown colonies and assessing the number of viable bacteria (colony-forming units, CFU), were performed as described previously (Spence et al. al., 2008; Dillard, 2011).

Isolation of the p$bla_{TEM-135}$ plasmid from a clinical strain of *N. gonorrhoeae* with natural resistance to penicillins

We used an *N. gonorrhoeae* clinical isolate from the collection of the State Scientific Center of Dermatovenerology and Cosmetology of the Ministry of Health of the Russian Federation, obtained in 2017 from Chuvash Republic, and this isolate contained a Toronto/Rio type p$bla_{TEM-135}$ plasmid, $MIC_{pen} \geq 32$ mg/L (Shaskolskiy et al., 2019). The clinical isolate was seeded on GC chocolate agar (Thermo Fisher Scientific, USA) that was supplemented with IsoVitaleX Enrichment (Becton-Dickinson, USA) and contained 16 mg/L benzylpenicillin (Sigma–Aldrich, USA). The dish was incubated overnight at 37°C and 5% CO_2. Grown colonies were harvested using a culture loop in 100 μl of PBS buffer. Isolation of plasmid DNA was performed with a Monarch Plasmid Miniprep

129 Kit/T1010 (New England Biolabs, UK). The concentration and purity of DNA preparations were
130 determined using a Nanodrop 2000 instrument (Thermo Fisher Scientific, USA). Plasmid DNA was
131 analyzed by PCR as described previously (Palmer et al., 2000; Shaskolskiy et al., 2019).

Construction of plasmid vectors with different variants of the *bla* gene on the basis of p*bla*$_{TEM-135}$ from a clinical *N. gonorrhoeae* strain

Amplification of the *bla* gene fragment with simultaneous site-directed mutagenesis, which ensures
that the mutations were introduced in the gene, was carried out. Using the primers 5'-
TTACTTCTGACAACGATCGGAGGACCGAAGG-3'-FOR and 5'-AATGATACCGCGAGAC
CCACGCTCACTGGCT-3'-REV, a PCR fragment of the *bla* gene was obtained, and this fragment
contained the GGT→AGT mutation that resulted in Gly238Ser and is present in TEM-20 ESBL
(Figure 1). A PCR fragment corresponding to the part of the *bla* gene that encodes TEM-1 broad-
spectrum beta-lactamase was obtained in a similar way.
The PCR fragments and the p*bla*$_{TEM-135}$ plasmid were treated with the restriction
endonucleases PvuI and BsaI. After that, the linearized p*bla*$_{TEM-135}$ plasmid and the PCR fragments
with sticky ends were treated with a ligation mixture to obtain the final p*bla*$_{TEM-1}$ и p*bla*$_{TEM-20}$
plasmids. The map of the resulting p*bla*$_{TEM-20}$ plasmid vector is shown in Figure 1.

Transformation of *E. coli* cells with plasmid vectors

The ready-to-use 10-beta competent *E. coli* strain DH10B (New England Biolabs, UK) was used for
transformation with the plasmid vectors. The puc19 (K+) vector was used as a transformation
control, and a sample without DNA was used as a negative control. The cells were grown and
subsequently reseeded on selective media. Media containing 256-512 mg/L benzylpenicillin was
used for *E. coli* with p*bla*$_{TEM-135}$ и p*bla*$_{TEM-1}$ plasmids; for cultures with p*bla*$_{TEM-20}$ plasmid, the media
contained ceftriaxone (Sigma–Aldrich, USA) at a concentration of 64 mg/L.
E. coli colonies grown on the selective media were collected with a culture loop in sodium
phosphate buffer (PBS), and plasmid DNA was isolated using the Plasmid Miniprep/BC021S kit
(Evrogen, Russia). The *bla* gene sequence and p*bla*$_{TEM}$ plasmid type were confirmed by PCR and
Sanger sequencing.

Transformation of *N. gonorrhoeae* cells with plasmid vectors

We used *N. gonorrhoeae* cells, strain ATCC 49226 (F-18, CDC 10, 001, P935), which is a reference
strain for antimicrobial susceptibility testing (https://www.culturecollections.org.uk/products/bacteria
/detail.jsp?refId=NCTC+12700&collection=nctc). Transformation of *N. gonorrhoeae* with p*bla*$_{TEM-1}$,
p*bla*$_{TEM-135}$ and p*bla*$_{TEM-20}$ plasmids, which were extracted from *E. coli* cells, was performed by
electroporation.
The optimal condition for transformation was resuspension of *N. gonorrhoeae* cells from the
overnight culture inoculum, collected from the dish with a sterile culture loop, in 0.3 M sucrose. The
cells were successively precipitated at 12000 RPM and resuspended in 0.3 M sucrose 2-3 times. For
electroporation, 100-200 µl of purified cell suspension and 50-100 ng of the plasmid vector were put
into 2 mm electroporation cells. A test tube without the addition of plasmid DNA was used as a
negative control. Electroporation was performed by applying single pulsed discharges of 2.5 kV, 200
Ω, and 25 µF. The cooled cell suspensions were plated on chocolate agar dishes, washed and
subcultured on dishes with a selective antibiotic; for strains with the p*bla*$_{TEM-1}$ and p*bla*$_{TEM-135}$
plasmids penicillin was used at a concentration of 2 mg/L, and for strains with the p*bla*$_{TEM-20}$ plasmid
ceftriaxone was used at a concentration of 0.5 mg/L.
The wild-type *N. gonorrhoeae* strain and strains with the p*bla*$_{TEM-1}$ and p*bla*$_{TEM-135}$ plasmids
were stored at −80°C in cryo-medium containing trypticase-soy broth and glycerol at a ratio of 4:1.
The presence of beta-lactamases in transformed cells was checked using nitrocefin discs

(Remel, USA), which qualitatively detect the presence of enzymes, including penicillinases and cephalosporinases that are capable of destroying the beta-lactam ring in the substrate (nitrocefin).

Estimating the susceptibility of *N. gonorrhoeae* to beta-lactam antibiotics

The following beta-lactam antibiotics were used in this work: benzylpenicillin; ampicillin (aminopenicillin with the extended spectrum of action); cefuroxime (2-nd generation cephalosporin); ceftriaxone, cefixime and cefotaxime (3-rd generation cephalosporins); cefepime (4-th generation cephalosporin); and meropenem, imipenem, and doripenem (carbapenems), which were all from Sigma–Aldrich, USA.

The MICs of antibiotics for *N. gonorrhoeae* strains were measured by a serial dilution method. *N. gonorrhoeae* was cultivated on chocolate agar GC supplemented with IsoVitaleX Enrichment. An inoculum (0.5 McFarland) was prepared for the wild-type *N. gonorrhoeae* strain (ATCC 49226) and strains with the p*bla*$_{TEM}$ plasmids. Microorganisms were seeded on Petri dishes containing selective medium with different concentrations of antimicrobial drugs. After incubation at 37°C with the addition of 5% CO_2, the presence/absence of bacterial growth on Petri dishes was assessed. According to the measurement results, each strain was characterized in accordance with the established EUCAST criteria (https://www.eucast.org/clinical_breakpoints): S – susceptible, R – resistant.

Assessment of *N. gonorrhoeae* cell viability: studies of cell growth and death

To estimate the effects of plasmids on the survival of gonococcus in the presence of antibiotics, the viability of *N. gonorrhoeae* cells that contain p*bla*$_{TEM}$ plasmids was studied by obtaining cell growth curves in the absence of antibiotics and cell death curves in the presence of ceftriaxone. The change in the number of viable cells (colony-forming units, CFU) versus time was determined by counting the number of colonies grown on solid media.

All experiments during the studies of the growth and death of *N. gonorrhoeae* cells were performed immediately after transformation that allowed us to avoid the cell storage stage. The absence of storage was important because of the low viability of *N. gonorrhoeae* cells carrying the p*bla*$_{TEM-20}$ plasmid.

To study cell growth, Petri dishes with chocolate agar (10 ml) without the addition of antimicrobials were used. An inoculum of cells (0.5 McFarland) was diluted to obtain ~50 CFU/ml, and 1 ml of cell suspension was plated on each dish, yielding 9 dishes that contained ~50 cells/dish for each *N. gonorrhoeae* strain. Every hour, the grown cells were washed off from every dish as follows: 1 ml of a sterile 0.3 M sucrose solution was poured onto the agar surface, the cells were resuspended on the dish with a sterile plastic loop without disturbing the agar surface layer, and the resulting cell suspension was transferred to a new dish. Thus, the cells grew on dish No. 1 for one hour, on dish No. 2 for two hours, and so on. The dishes with the washed-off cells were incubated for 24-48 hours at 37°C in the presence of 5% CO_2, and the grown colonies were counted.

To study cell death in the presence of ceftriaxone, an inoculum of cells (0.5 McFarland) was diluted to obtain ~1000 CFU/ml, and 1 ml of cell suspension was plated on each dish with ceftriaxone, yielding 9 dishes that contained ~1000 cells/dish for each *N. gonorrhoeae* strain. Concentrations of ceftriaxone were 0.03, 0.125, and 2 mg/L. Every hour, the grown cells were washed off from every dish, and the resulting cell suspension was transferred to a new dish without the addition of ceftriaxone. The dishes were incubated as described above, and the grown colonies were counted.

Mathematical models for cell growth and death

The generalized Verhulst equation was used to model cell growth curves for all strains under study (Peleg and Normand, 2007; Peleg and Corradini, 2011). To obtain cell death curves in the presence

of ceftriaxone for the strains that did not contain the pbla_{TEM-20} plasmid, a modified Chick-Watson
model (Jensen, 2010; Peleg, 2021) was used. The equation parameters were fit by enumerating the
solutions of the Cauchy problem using MATLAB 2021b software for different r, N_{assymp}, and α for
the Verhulst equation and for different k_{obs} values for the Chick-Watson equation. The optimization
of the obtained numerical solutions was carried out by the least squares method when compared with
the experimental results. The statistical significance of the difference between the mathematical
models, which describe experimental results belonging to different data series, was evaluated by
Fisher's test (F test) in accordance with a procedure described previously (Motulsky and Ransnas,
1987).

RESULTS

Obtaining pbla_{TEM-1}, p$bla_{TEM-135}$, and pbla_{TEM-20} vector constructs

The p$bla_{TEM-135}$ plasmid obtained from the clinical isolate of *N. gonorrhoeae* was used to construct
artificial pbla_{TEM-1} and pbla_{TEM-20} vectors. As a result, three 5154 bp plasmids were obtained, which
differed in the *bla* gene structure (Figure 1) as follows:
pbla_{TEM-1}, no mutations in codons 182 and 238;
p$bla_{TEM-135}$, ATG→ACG substitution at codon 182 leading to Met182Thr substitution, no
mutation at codon 238;
pbla_{TEM-20}, ATG→ACG substitution at codon 182 leading to Met182Thr substitution, and
GGT→AGT substitution at codon 238 leading to the Gly238Ser substitution.
Plasmids could replicate in both *E. coli* and *N. gonorrhoeae* due to the presence of replication origins
that were compatible with gram-negative bacteria and replication initiation proteins.

Preparation of *E. coli* and *N. gonorrhoeae* strains containing pbla_{TEM} plasmids

E. coli and *N. gonorrhoeae* strains containing pbla_{TEM-1}, p$bla_{TEM-135}$, and pbla_{TEM-20} plasmids were
obtained. *E. coli* strains with pbla_{TEM} plasmids were grown on selective media with antibiotics and
stored on Petri dishes at 4°C.
The *N. gonorrhoeae* strains with pbla_{TEM-1} and p$bla_{TEM-135}$ plasmids retained their viability
after several (4-5) passages from dishes and after storage in cryo-medium. However, the *N.
gonorrhoeae* strain with the pbla_{TEM-20} plasmid could not be stored in the cryo-medium, and the cells
lost their viability after 12 hours of incubation on the plates; colonies did not grow after undergoing a
subculture on the media both with and without antibiotics (penicillin, ceftriaxone). The loss of
viability cannot be explained by the loss of plasmids, since the result of the test on nitrocephin disks
was positive, indicating the presence of beta-lactamases. PCR and Sanger sequencing also confirmed
the presence of pbla_{TEM} plasmids with introduced mutations in the beta-lactamase gene.

Antimicrobial susceptibility testing of the *N. gonorrhoeae* strains containing different pbla_{TEM}
variants

The susceptibility of *N. gonorrhoeae* strains that contain various pbla_{TEM} plasmids to several groups
of beta-lactam antibiotics was studied. The antibiotics included penicillins, which were previously
used to treat gonococcal infection, cephalosporins, which are currently used for gonorrhea treatment,
and carbapenems, which may replace cephalosporins if cephalosporin resistance spreads (Unemo et
al., 2020). The MICs measured for these drugs are shown in Table 1.
After transformation of the wild-type *N. gonorrhoeae* strain with the pbla_{TEM-1} and pbla_{TEM-}
$_{135}$ plasmid vectors, which contain genes encoding the broad-spectrum beta-lactamase (penicillinase),
the bacteria exhibited a sharp decrease in susceptibility to antibiotics from the penicillin subgroup
(MIC$_{pen}$ = 16-32 mg/L, MIC$_{amp}$ = 8 mg/L). The MICs measured on the solid media confirmed that the
TEM-1 and TEM-135 enzymes were unable to hydrolyze cephalosporins and carbapenems.

The MICs of the strain containing the p*bla*TEM-20 plasmid confirmed that the expressed TEM-
20 beta-lactamase is an ESBL that can hydrolyze both penicillins and cephalosporins of different
generations. The cephalosporin MICs were above the EUCAST threshold for susceptible/resistant
strains established to be 0.125 mg/L (Table 1). For example, the MICs of 3-rd and 4-th-generation
cephalosporins were equal to or higher than 4 mg/L. Thus, the level of resistance of the strain with
the p*bla*TEM-20 plasmid was higher than the threshold value established for 3-rd and 4-th-generation
cephalosporins by more than a fivefold dilution of the antibiotic.
The measured MICs of carbapenems in all *N. gonorrhoeae* strains did not exceed 0.008 mg/L,
i.e., all strains were susceptible to carbapenems, proving that the TEM-1, TEM-135, and TEM-20
beta-lactamase variants are unable to hydrolyze carbapenems (the EUCAST threshold for
carbapenems is absent).

Growth curves of the *N. gonorrhoeae* strains harboring different p*bla*TEM plasmids, and the
mathematical model for cell growth

Figure 2 shows the growth curves of the *N. gonorrhoeae* cells harboring the p*bla*TEM-1, p*bla*TEM-135,
and p*bla*TEM-20 plasmids compared to that of the wild-type cells in the absence of antimicrobials. It
should be noted that in our experiments we determined the number of viable cells calculated as the
number of colonies grown on a dish (CFU). The wild-type strain and strains with p*bla*TEM-1 and
p*bla*TEM-135 plasmids demonstrated bacterial growth during all 8 hours of cultivation. At the same
time, the number of viable *N. gonorrhoeae* cells with the p*bla*TEM-20 plasmid began to decrease after 6
hours of cultivation.

To compare the *N. gonorrhoeae* strains that contain different plasmids, we performed
mathematical modeling for the kinetic curves of cell growth. The generalized Verhulst equation
(Peleg, 2007; Peleg, 2011) (eq. (1)) was chosen for modeling since it is the simplest and most
universal equation for describing the dynamics of population growth up to the transition to the
stationary phase.

$$\frac{dN(t)}{dt} = rN(t)^{\alpha}\left(1 - \frac{N(t)}{N_{assymp}}\right) \qquad eq. (1)$$

where $dN(t)/dt$ is the growth rate at a given time;
r is the proportionality coefficient;
$N(t)$ is the number of organisms at a given time;
N_{assymp} is the maximum possible number of grown cells of a given type under given experimental
conditions;
α is the constant characterizing the growth peculiarity; α < 1 means that the organism does not
achieve its potential for exponential growth; and α > 1 means that the organism exceeds this ability
(Peleg, 2007).

Fitting of the parameters for equation (1) by enumerating the solutions of the Cauchy
problems for different r, N_{assymp}, and α resulted in equations (2)-(5), which describe the growth curves
of cells with and without plasmids (Figure 2).
N. gonorrhoeae without plasmids (wild-type):

$$\frac{dN(t)}{dt} = 0.188\,N(t)^{1.093}\left(1 - \frac{N(t)}{9300}\right) \qquad eq. (2)$$

303　　　　*N. gonorrhoeae* with p*bla*TEM-1:

$$\frac{dN(t)}{dt} = 0.018\,N(t)^{1.593}\left(1 - \frac{N(t)}{1100}\right) \qquad eq.\,(3)$$

304   *N. gonorrhoeae* with p*bla*$_{TEM-135}$:

$$\frac{dN(t)}{dt} = 0.21\,N(t)^{1.071}\left(1 - \frac{N(t)}{2600}\right) \qquad eq.\,(4)$$

305   *N. gonorrhoeae* with p*bla*$_{TEM-20}$ (up to 6 hours of growth):

$$\frac{dN(t)}{dt} = 0.012\,N(t)^{1.819}\left(1 - \frac{N(t)}{235}\right) \qquad eq.\,(5)$$

The R-squared coefficients were 0.9916 for the wild-type, 0.9958 for p*bla*$_{TEM-1}$, 0.9801 for p*bla*$_{TEM-}$
$_{135}$, and 0.9778 for p*bla*$_{TEM-20}$.
The statistical significance of the difference between the obtained models for describing the
data (*p* value) was calculated using the Fisher criterion. The wild-type model outperformed the model
for TEM-1 with $p = 0.02$ and TEM-135 with $p = 0.02$. The model for describing TEM-1 was superior
to that of the wild-type and TEM-135 models, with $p = 0.01$. The model for describing TEM-135 was
superior to that of the wild-type model with $p = 0.05$ and the model for TEM-1 with $p = 0.06$. This
indicates that the resulting models adequately describe cell growth for the corresponding strain.
The calculated value of N_{assymp} was different for the wild-type *N. gonorrhoeae* cells and for
the cells with the p*bla*$_{TEM}$ plasmids as follows: a maximum N_{assymp} of 9300 CFU was observed for the
wild-type *N. gonorrhoeae*, lower values were observed for *N. gonorrhoeae* with the p*bla*$_{TEM-135}$ and
p*bla*$_{TEM-1}$ plasmids, which were 2600 and 1100 CFU, respectively, and a very low CFU of 235 was
observed for *N. gonorrhoeae* with the p*bla*$_{TEM-20}$ plasmid. This means that the cells containing
p*bla*$_{TEM}$ plasmids had a reduced capacity of growth compared to that of the cells without plasmids,
which can be explained by additional energy and metabolic costs for plasmid reproduction. At the
same time, the strain with the p*bla*$_{TEM-20}$ plasmid that expressed beta-lactamase with the Gly238Ser
substitution showed significantly reduced growth ability compared to that of the wild-type strain and
strains harboring the other plasmids. The reduced viability of the strain with the p*bla*$_{TEM-20}$ plasmid
was also confirmed by the previously noted strain characteristics: the cells survived for no more than
12 hours when stored on solid media and did not survive when stored in a cryogenic medium.

Change in the number of viable *N. gonorrhoeae* cells containing different p*bla*$_{TEM}$ plasmids in
the presence of ceftriaxone

Changes in the number of viable cells over time in the presence of various concentrations of
ceftriaxone (0.03, 0.125 and 2 mg/L) are shown in Figure 3. Similar to the studies of bacterial
growth, the CFU number on the solid media (chocolate agar) under various cultivation conditions
was determined rather than the total number of cells. As shown in Figure 3A-C, the strains that did
not contain ceftriaxone resistance determinants rapidly lost viability in the presence of this
antimicrobial agent (the ceftriaxone concentrations in this experiment were equal to or greater than
the MIC$_{cro}$ for these strains, which were 0.015-0.03 mg/L (Table 1)). After 2-8 hours of incubation,
the bacteria were completely extinct.
Under our experimental conditions, the change in the concentration of ceftriaxone over time
can be neglected, since this concentration is much higher than that of the cells in the sample; [CRO]
= 0.03-2.0 mg/L or $(0.045\text{-}3.030)\cdot 10^{-6}$ M, N_0 = 1000 cells (cells at zero time point). The decrease in
CFU versus time for the curves in Figure 3A-C (strains without plasmids and with p*bla*$_{TEM-1}$ and

p*bla*TEM-135) followed a logarithmic law, *i.e.,* kinetic curves can be considered applicable for chemical
reactions of the first (pseudo-first) order (Scheme 1).

$$\text{Viable cells} + \text{CRO} \xrightarrow{k_{obs}} \text{Nonviable cells} \qquad \text{(Scheme 1)}$$

To describe the kinetic curves, a modified Chick-Watson model, which was developed for
disinfection curves (cell death under the action of a disinfectant agent), can be applied. According to
Chick's law, the dependence of cell survival on time is described by the equations for the first-order
chemical reaction (Jensen, 2010; Peleg, 2021) (equations (6) and (7)).

$$\frac{dN(t)}{dt} = -k_{obs}\, N(t) \qquad\qquad eq.\,(6)$$

$$\ln \frac{N_0}{N} = k_{obs}\, t \qquad\qquad eq.\,(7)$$

where $dN(t)/dt$ is the growth rate at a given time;
$N(t)$ is the number of cells at a given time;
k_{obs} is the observed first-order rate constant.

Fitting the cell death curves (Figure 3A-C) using equation (7) made it possible to obtain the
values of the observed first-order rate constants k_{obs} for all strains at different concentrations of the
disinfectant agent, which in our case was ceftriaxone (pseudo-first order rate constants). The
dependences of k_{obs} on the concentration of ceftriaxone are shown in Figure 4.
According to the modified Chick-Watson model, the dependence of the cell death rate on the
concentration of ceftriaxone is determined by the following equation:

$$\frac{dN(t)}{dt} = -k_d\, [CRO]^n N(t) \qquad\qquad eq.\,(8)$$

356 where $[CRO]$ is the ceftriaxone concentration;
357 n is the fitting coefficient, which is also called the coefficient of dilution;
358 and k_d is the true rate constant of cell death.

359   Thus, the observed death rate constant is related to the true rate constant and ceftriaxone
360 concentration as follows:

$$k_{obs} = k_d\, [CRO]^n \qquad\qquad eq.\,(9)$$

361 The dependences of k_{obs} on the concentration of ceftriaxone, shown in Figure 4, were approximated
362 by the following power functions (R^2 – approximation confidence value):
363   *N. gonorrhoeae* without plasmids (wild-type):

$$k_{obs} = 2.71\, [CRO]^{0.44} \qquad\qquad eq.\,(10) \quad (R^2 = 0.9677)$$

365   *N. gonorrhoeae* with p*bla*TEM-1:

$$k_{obs} = 1.86\, [CRO]^{0.31} \qquad\qquad eq.\,(11) \quad (R^2 = 0.8461)$$

367   *N. gonorrhoeae* with p*bla*TEM-135:

$$368 \qquad k_{obs} = 1.44\,[CRO]^{0.22} \qquad\qquad eq.(12) \quad (R^2 = 0.9626)$$

369   Thus, the equations describing the decrease in *N. gonorrhoeae* CFU versus time under the influence
370   of ceftriaxone were as follows:
371   *N. gonorrhoeae* without plasmids (wild-type):

$$\frac{dN(t)}{dt} = -2.71\,[CRO]^{0.44}\,N(t) \qquad\qquad eq.(13)$$

372   *N. gonorrhoeae* with $pbla_{TEM-1}$:

$$\frac{dN(t)}{dt} = -1.86\,[CRO]^{0.31}\,N(t) \qquad\qquad eq.(14)$$

373   *N. gonorrhoeae* with $pbla_{TEM-135}$:

$$\frac{dN(t)}{dt} = -1.44\,[CRO]^{0.22}\,N(t) \qquad\qquad eq.(15)$$

The results showed that the death of *N. gonorrhoeae* cells in the presence of ceftriaxone
occurred differently in various strains. In the wild-type strain without $pbla_{TEM}$, the cells died
somewhat faster than in the strains carrying $pbla_{TEM-1}$ and $pbla_{TEM-135}$, although the beta-lactamases
expressed by these plasmids were not capable of destroying cephalosporins.
For the *N. gonorrhoeae* strain carrying the $pbla_{TEM-20}$ plasmid that expresses ESBL, the CFU
reduction versus time curves looked completely different (Figure 3D). The MIC of ceftriaxone for
this strain was 4 mg/L (Table 1), and at a ceftriaxone concentration of 0.03-0.125 mg/L, some
decrease in the number of live cells was observed in the first 3 hours. Then, a short-term increase in
CFU was observed over a period of 3-6 hours, and this was changed to a reduction in CFU that was
similar to the reduction observed in the *N. gonorrhoeae* strain with the $pbla_{TEM-20}$ plasmid when it
was cultivated in the absence of the antibiotic (Figure 2, curve for the strain with the $pbla_{TEM-20}$
plasmid). At ceftriaxone concentration of 0.03 mg/L, *i.e.*, much less than the MIC_{cro} for the strain
with $pbla_{TEM-20}$, a significant amount of the antibiotic seemed to be hydrolyzed in the first 3 hours,
and the number of cells reached its initial value during the next 3 hours. It should also be noted that
the CFU versus time curves for this strain did not obey the logarithmic law, and in this case, the
Chick-Watson model was not applicable to describe the kinetic curves.
Obviously, the kinetic curves for cells with $pbla_{TEM-20}$ reflected several of the following
processes: hydrolysis of ceftriaxone by TEM-20 beta-lactamase, growth, and gradual cell death.
Therefore, these curves could not be described by a simple Chick-Watson model for cell death in the
presence of a disinfectant agent, which in our case was ceftriaxone.

DISCUSSION

In this work, *N. gonorrhoeae* strains containing different types of beta-lactamase plasmids were
studied for their susceptibility to beta-lactam antibiotics and for their viability. The artificial $pbla_{TEM-1}$
and $pbla_{TEM-20}$ plasmids were obtained by site-directed mutagenesis from the natural $pbla_{TEM-135}$
plasmid of the Toronto/Rio type. Efficient plasmid amplification occurred in *E. coli* cells due to the
presence in the vector's structure of specific replication origins (ori1, ori2, ori3) and *repA* and *repB*
genes encoding replication initiation proteins. Transformation of the *N. gonorrhoeae* strain ATCC

49226 with plasmid vectors was carried out using electroporation to obtain individual strains, for
which the presence of $pbla_{TEM}$ plasmids was confirmed. Among the strains, the *N. gonorrhoeae*
strain was obtained with the $pbla_{TEM-20}$ plasmid producing ESBL, which has not yet been found
among clinical isolates.

Indeed, a single Gly238Ser amino acid substitution in the beta-lactamase plasmid gene was
sufficient to provide this enzyme with the ability to hydrolyze cephalosporins, while retaining the
ability to destroy penicillins. However, the *N. gonorrhoeae* strain that contained the $pbla_{TEM-20}$
plasmid showed substantially reduced viability; cell growth reached a plateau after 6 hours of
cultivation both in the absence and in the presence of the antibiotic. In addition, the cells did not
grow after incubation on plates for more than 12 hours and after freezing. Our results intersect with
the work of Cole et al., 2018, in which genetically engineered strains of *N. gonorrhoeae* of the
widespread NG-MAST type 1407, carrying African-type penicillinase-producing plasmids, were
obtained. However, the strains described in the work of Cole et al. were unable to retain plasmids in
the absence of antibiotic selection, and the cells lost plasmids even after one passage in the absence
of penicillin. In our work, the reduced viability of the *N. gonorrhoeae* cells carrying the $pbla_{TEM-20}$
plasmid was not associated with the loss of the plasmid during cultivation.

The reduced viability of the strain carrying the $pbla_{TEM-20}$ plasmid that was demonstrated in
this work can be explained by the following reasons: (a) the presence of the plasmid itself, which
requires additional expenses from cells for its reproduction, and (b) the expression of ESBL. The
action of cephalosporins, like all beta-lactam antimicrobials, is aimed at inhibiting the synthesis of
the bacterial cell wall by covalently inhibiting the action of transpeptidases (penicillin-binding
proteins). The main component of the cell wall is peptidoglycan, which is a macromolecular structure
that consists of peptide and sugar components. To protect themselves against beta-lactam
antimicrobials, bacteria express beta-lactamases, which are located in the periplasmic space and
hydrolyze the C-N bond in the beta-lactam ring of the antibiotic, thereby inactivating the AMP.
Because peptidoglycan in gram-negative bacteria possesses a C-terminal motif in the acyl-D-Ala-D-
Ala peptide chain that is similar to the structural fragment of beta-lactam preparations, beta-
lactamases can cause changes in the composition of peptidoglycan, thereby reducing the viability of
bacteria. A change in the structure of the cell membrane leads to the suppression of cell division
(bacteriostatic effect) or to the rupture of bacteria due to osmotic pressure (bactericidal activity)
(Sawa et al., 2020). In the work of Fernández et al., 2012, it was shown that *E. coli*, which expressed
certain beta-lactamase variants, underwent a change in their peptidoglycan structures, in particular, a
decrease in the level of cross-linked muropeptides that negatively affected the viability of such
strains.

Evaluation of the resistance of the *N. gonorrhoeae* strains with $pbla_{TEM}$ plasmids to beta-
lactam antibiotics also showed that none of the obtained beta-lactamase variants, including ESBL
TEM-20, were capable of hydrolyzing carbapenems. Thus, carbapenems remain among the drugs of
the beta-lactam series that can resist hydrolysis by *N. gonorrhoeae* bacterial lactamases, which
confirms that they are potential drugs for the treatment of gonococcal infection.

The kinetic curves of cell growth were mathematically modeled using the generalized
Verhulst equation (Peleg, 2007; Peleg, 2011). The value of N_{assymp}, a parameter that characterizes the
maximum possible number of cells grown under given conditions, was lower for the strains carrying
the $pbla_{TEM}$ plasmids compared with the cells without plasmids, *i.e.,* plasmid-bearing strains had
reduced viability compared to that of the wild-type *N. gonorrhoeae*. N_{assymp} decreased within the
series WT – $pbla_{TEM-135}$ – $pbla_{TEM-1}$ – $pbla_{TEM-20}$. The reduced viability of the strains with $pbla_{TEM}$
plasmids may explain why these strains have a lower frequency of occurrence in the *N. gonorrhoeae*
population compared to that of the strains without plasmids. The higher viability of *N. gonorrhoeae*
with $pbla_{TEM-135}$ compared to that of *N. gonorrhoeae* with $pbla_{TEM-1}$ may be associated with the
presence of the Met182Thr mutation in beta-lactamase, which leads to the stabilization of the

structure of the enzyme. As noted previously (Orencia et al., 2001; Cole et al., 2015; Yan et al.,
2019), increased enzyme stability may have contributed to the fixation and spread of the TEM-135
beta-lactamase variant in the *N. gonorrhoeae* population.
The decrease in the number of viable cells with cultivation time in the presence of ceftriaxone
has been studied. The kinetics of *N. gonorrhoeae* cell death without plasmids and with the p*bla*$_{TEM-1}$
and p*bla*$_{TEM-135}$ plasmids that lack ceftriaxone resistance determinants could be described by a
modified Chick-Watson law for cell death modeling in the presence of a disinfectant (Jensen, 2010;
Peleg, 2021). The death rate for the wild-type *N. gonorrhoeae* strain in the presence of ceftriaxone
was higher than that of the strains with the p*bla*$_{TEM-1}$ and p*bla*$_{TEM-135}$ plasmids, although the strains
carrying these plasmids were susceptible to ceftriaxone according to the EUCAST criteria.
Curves of the CFU versus time for the *N. gonorrhoeae* strain carrying the p*bla*$_{TEM-20}$ plasmid
that expresses ESBL cannot be described by a simple Chick-Watson model. The kinetic death curves
in this case reflected several processes: hydrolysis of ceftriaxone by TEM-20 beta-lactamase, growth,
and gradual cell death.
The obtained data on the reduced viability of *N. gonorrhoeae* strains with the p*bla*$_{TEM-20}$
plasmid may explain the absence of *N. gonorrhoeae* clinical isolates producing ESBL that hydrolyze
cephalosporins of various generations.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

ETHICS STATEMENT

Ethical approval/written informed consent was not required for the study of animals/human
participants in accordance with the local legislation and institutional requirements.

AUTHOR CONTRIBUTIONS

I.K. performed the experiments, analyzed results, and wrote the manuscript; D.G. designed and
supervised the project, wrote the manuscript, A.V., O.A. and J.S. carried out antimicrobial
susceptibility testing; A.K. and V.S. supervised work with cell cultures; D.D. wrote the manuscript;,
B.S. directed the project, performed mathematical modeling and wrote the manuscript. All authors
contributed to the article and approved the submitted version.

FUNDING

This work was supported by the Russian Science Foundation, grant number 17-75-20039 (plasmid
construction, susceptibility testing, mathematical modeling, cell growth characteristics) and by the
Ministry of Science and Higher Education of the Russian Federation to the EIMB Center for
Precision Genome Editing and Genetic Technologies for Biomedicine under the Federal Research
Program for Genetic Technologies Development for 2019-27, agreement number 075-15-2019-1660
(gene sequencing and analysis of sequence data). Work with cell cultures was performed according to
the Ministry of Health of the Russian Federation assignment number 056-03-2021-124.

REFERENCES

Adator, E. H., Narvaez-Bravo, C., Zaheer, R., Cook, S. R., Tymensen, L., Hannon, S.J., et al. (2020).
A One Health Comparative Assessment of Antimicrobial Resistance in Generic and Extended-

Spectrum Cephalosporin-Resistant *Escherichia coli* from Beef Production, Sewage and Clinical
Settings. *Microorganisms* 8 (6), 885. doi: 10.3390/microorganisms8060885.
Allen, V. G., Mitterni, L., Seah, C., Rebbapragada, A., Martin, I.E., Lee, C., et al. (2013). *Neisseria*
gonorrhoeae Treatment Failure and Susceptibility to Cefixime in Toronto, Canada. *JAMA* 309
(2), 163-170. doi: 10.1001/jama.2012.176575.
Arlet, G., Goussard, S., Courvalin, P., and Philippon, A. (1999). Sequences of the Genes for the
TEM-20, TEM-21, TEM-22, and TEM-29 Extended-Spectrum Beta-Lactamases. *Antimicrob.*
Agents Chemother. 43 (4), 969-971. doi: 10.1128/AAC.43.4.969.
Attram, N., Agbodzi, B., Dela, H., Behene, E., Nyarko, E. O., Kyei, N. N. A., et al. (2019).
Antimicrobial Resistance (AMR) and Molecular Characterization of *Neisseria gonorrhoeae* in
Ghana, 2012-2015. *PLoS One* 14 (10), e0223598. doi: 10.1371/journal.pone.0223598.
Bradford, P. A. (2001). Extended-Spectrum Beta-Lactamases in the 21st Century: Characterization,
Epidemiology, and Detection of This Important Resistance Threat. *Clin. Microbiol. Rev.* 14 (4),
933-951. doi: 10.1128/CMR.14.4.933-951.2001.
Bush, K., and Jacoby, G. A. (2010). Updated Functional Classification of Beta-Lactamases.
Antimicrob. Agents Chemother. 54 (3), 969-976. doi: 10.1128/AAC.01009-09.
Cole, M. J., Unemo, M., Grigorjev, V., Quaye, N., and Woodford, N. (2015). Genetic Diversity of
blaTEM Alleles, Antimicrobial Susceptibility and Molecular Epidemiological Characteristics of
Penicillinase-Producing *Neisseria gonorrhoeae* from England and Wales. *J. Antimicrob.*
Chemother. 70 (12), 3238-3243. doi: 10.1093/jac/dkv260.
Cole, M. J., Ison, C., and Woodford, N. (2018). Transfer of a Gonococcal beta-Lactamase Plasmid
into *Neisseria gonorrhoeae* Belonging to the Globally Distributed ST1407 Lineage. *J*
Antimicrob. Chemother. 73 (9), 2576-2577. doi: 10.1093/jac/dky194.
Dillard, J. P. (2011). Genetic Manipulation of *Neisseria gonorrhoeae. Curr. Protoc. Microbiol.*
Chapter 4, Unit4A.2. doi: 10.1002/9780471729259.mc04a02s23.
EUCAST (2022) The European Committee on Antimicrobial Susceptibility Testing. Breakpoint
Tables for Interpretation of MICs and Zone Diameters. Version 12.0. http://www.eucast.org.
Fernández, A., Pérez, A., Ayala, J. A., Mallo, S., Rumbo-Feal, S., Tomás, M., et al. (2012).
Expression of OXA-Type and SFO-1 beta-Lactamases Induces Changes in Peptidoglycan
Composition and Affects Bacterial Fitness. *Antimicrob. Agents Chemother.* 56 (4), 1877-1884.
doi: 10.1128/AAC.05402-11.
Golparian, D., Ohlsson, A., Janson, H., Lidbrink, P., Richtner, T., Ekelund, O., et al. (2014). Four
Treatment Failures of Pharyngeal Gonorrhoea with Ceftriaxone (500 mg) or Cefotaxime (500
mg), Sweden, 2013 and 2014. *Euro Surveill.* 19 (30), 20862. doi: 10.2807/1560-
7917.es2014.19.30.20862.
Golparian, D., and Unemo, M. (2022). Antimicrobial Resistance Prediction in *Neisseria*
gonorrhoeae: Current Status and Future Prospects. *Expert Rev. Mol. Diagn.* 22 (1), 29-48. doi:
10.1080/14737159.2022.2015329.
Jensen, J. N. (2010). Disinfection Model Based on Excess Inactivation Sites: Implications For Linear
Disinfection Curves and the Chick-Watson Dilution Coefficient. *Environ. Sci. Technol.* 44 (21),
8162-8168. doi: 10.1021/es101818z.
Jeong, S. H., Bae, I. K., Lee, J. H., Sohn, S. G., Kang, G. H., Jeon, G. J., et al. (2004). Molecular
Characterization of Extended-Spectrum Beta-Lactamases Produced by Clinical Isolates of
Klebsiella pneumoniae and *Escherichia coli* from a Korean Nationwide Survey. *J Clin*
Microbiol. 42 (7), 2902-2906. doi: 10.1128/JCM.42.7.2902-2906.2004.
Kubanov, A., Solomka, V., Plakhova, X., Chestkov, A., Petrova, N., Shaskolskiy, B. et al. (2019).
Summary and Trends of the Russian Gonococcal Antimicrobial Surveillance Programme, 2005-
2016. *J. Clin. Microbiol.* 5 (6), e02024-18. doi: 10.1128/JCM.02024-18.
Motulsky, H. J., and Ransnas, L. A. (1987). Fitting Curves to Data Using Nonlinear Regression: a

Practical and Nonmathematical Review. *FASEB J.* 1 (5), 365-374. PMID: 3315805.

Müller, E.E., Fayemiwo, S. A., and Lewis, D. A. (2011). Characterization of a Novel beta-
Lactamase-Producing Plasmid in *Neisseria gonorrhoeae*: Sequence Analysis and Molecular
Typing of Host Gonococci. *J. Antimicrob. Chemother.* 66 (7), 1514-1517. doi:
10.1093/jac/dkr162.

Muhammad, I., Golparian, D., Dillon, J. A., Johansson, A., Ohnishi, M., Sethi, S., et al. (2014).
Characterisation of blaTEM Genes and Types of beta-Lactamase Plasmids in *Neisseria*
gonorrhoeae - the Prevalent and Conserved blaTEM-135 Has Not Recently Evolved and Existed
in the Toronto Plasmid from the Origin. *BMC Infect. Dis.*14, 454. doi: 10.1186/1471-2334-14-
454.

Nakayama, S., Tribuddharat, C., Prombhul, S., Shimuta, K., Srifuengfung, S., Unemo, M., et al.
(2012). Molecular Analyses of TEM Genes and Their Corresponding Penicillinase-Producing
Neisseria gonorrhoeae Isolates in Bangkok, Thailand. *Antimicrob. Agents Chemother.* 56 (2),
916-920. doi: 10.1128/AAC.05665-11.

Ohnishi, M., Ono, E., Shimuta, K., Watanabe, H., and Okamura, N. (2010) Identification of TEM-
135 beta-Lactamase in Penicillinase-Producing *Neisseria gonorrhoeae* Strains in Japan.
Antimicrob. Agents Chemother. 54 (7), 3021-3023. doi: 10.1128/AAC.00245-10.

Orencia, M. C., Yoon, J.S., Ness, J. E., Stemmer, W. P., and Stevens, R. C. (2001). Predicting the
Emergence of Antibiotic Resistance by Directed Evolution and Structural Analysis. *Nat. Struct.*
Biol. 8 (3), 238-242. doi: 10.1038/84981.

Palmer, H. M., Leeming, J. P., and Turner, A. (2000). A Multiplex Polymerase Chain Reaction to
Differentiate beta-Lactamase Plasmids of *Neisseria gonorrhoeae*. *J. Antimicrob. Chemother.* 45
(6), 777-782. doi: 10.1093/jac/45.6.777.

Peleg, M., Corradini, M. G., and Normand, M. D. (2007). The Logistic (Verhulst) Model for Sigmoid
Microbial Growth Curves Revisited. *Food Research International* 40 (7), 808-818.

Peleg, M., and Corradini, M. G. (2011). Microbial Growth Curves: What the Models Tell Us and
What They Cannot. *Crit. Rev. Food Sci. Nutr.* 51 (10), 917-945. doi:
10.1080/10408398.2011.570463.

Peleg, M. (2021). Modeling the Dynamic Kinetics of Microbial Disinfection with Dissipating
Chemical Agents -a Theoretical Investigation. *Appl. Microbiol. Biotechnol.* 105 (2), 539-549.
doi: 10.1007/s00253-020-11042-8.

Sawa, T., Kooguchi, K., and Moriyama, K. (2020). Molecular Diversity of Extended-Spectrum beta-
Lactamases and Carbapenemases, and Antimicrobial Resistance. *J. Intensive Care* 8, 13. doi:
10.1186/s40560-020-0429-6.

Shaskolskiy, B., Dementieva, E., Kandinov, I., Filippova, M., Petrova, N., Plakhova, X., et al.
(2019). Resistance of *Neisseria gonorrhoeae* Isolates to Beta-Lactam Antibiotics
(Benzylpenicillin and Ceftriaxone) in Russia, 2015-2017. *PLoS One* 14 (7), e0220339. doi:
10.1371/journal.pone.0220339.

Spence, J. M., Wright, L., and Clark, V. L. (2008). Laboratory Maintenance of *Neisseria*
gonorrhoeae. *Curr. Protoc. Microbiol.* Chapter 4, Unit4A.1. doi:
10.1002/9780471729259.mc04a01s8.

Tacconelli, E., Carrara, E., Savoldi, A., Harbarth, S., Mendelson, M., Monnet, D. L., et al. (2018).
Discovery, Research, and Development of New Antibiotics: the WHO Priority List of
Antibiotic-Resistant Bacteria and Tuberculosis. *Lancet Infect. Dis.* 18 (3), 318-327. doi:
10.1016/S1473-3099(17)30753-3.

Tanaka, M., Furuya, R., Kobayashi, I., Ohno, A., and Kanesaka, I. (2021). Molecular Characteristics
and Antimicrobial Susceptibility of Penicillinase-Producing *Neisseria gonorrhoeae* Isolates in
Fukuoka, Japan, 1996-2018. *J. Glob. Antimicrob. Resist.* 26, 45-51. doi:
10.1016/j.jgar.2021.04.014.

Tooke, C. L,, Hinchliffe, P., Bragginton, E. C., Colenso, C. K., Hirvonen, V. H. A., et al. (2019).
beta-Lactamases and beta-Lactamase Inhibitors in the 21st Century. *J. Mol. Biol.* 431 (18), 3472-
3500. doi: 10.1016/j.jmb.2019.04.002.
Unemo, M., Golparian, D., Nicholas, R., Ohnishi, M., Gallay, A., and Sednaoui, P. (2012). High-
level Cefixime- and Ceftriaxone-Resistant *Neisseria gonorrhoeae* in France: Novel *penA* Mosaic
Allele in a Successful International Clone Causes Treatment Failure. *Antimicrob. Agents*
Chemother. 56 (3), 1273-1280. doi: 10.1128/AAC.05760-11.
Unemo, M., and Shafer, W. M. (2014). Antimicrobial Resistance in *Neisseria gonorrhoeae* in the
21st Century: Past, Evolution, and Future. *Clin. Microbiol. Rev.* 27 (3), 587-613. doi:
10.1128/CMR.00010-14.
Unemo, M., and Jensen, J. S. (2017). Antimicrobial-Resistant Sexually Transmitted Infections:
Gonorrhoea and *Mycoplasma genitalium*. *Nat. Rev. Urol.* 14 (3), 139-152. doi:
10.1038/nrurol.2016.268.
Unemo, M., Seifert, H. S., Hook, E. W. 3rd, Hawkes, S., Ndowa, F., and Dillon, J. R. (2019).
Gonorrhoea.. *Nat. Rev. Dis. Primers* 5 (1), 79. doi: 10.1038/s41572-019-0128-6.
Unemo, M., Ross, J., Serwin, A. B., Gomberg, M., Cusini, M., and Jensen, J. S. (2020). 2020
European Guideline for the Diagnosis and Treatment of Gonorrhoea in Adults. *Int. J. STD AIDS*.
956462420949126. doi: 10.1177/0956462420949126.
WHO (2016). WHO Guidelines for the Treatment of *Neisseria gonorrhoeae*.
https://www.ncbi.nlm.nih.gov/books/NBK379221/.
Workowski, K. A., and Bolan, G. A. (2015). Sexually Transmitted Diseases Treatment Guidelines,
2015. *MMWR Recomm. Rep.* 5, 1-137.
Yan, J., Zhang, J., and van der Veen, S. (2019). High Prevalence of TEM-135 Expression from the
Asian Plasmid in Penicillinase-Producing *Neisseria gonorrhoeae* from Hangzhou, China. *Int. J.*
Antimicrob. Agents. 54 (3), 361-366. doi: 10.1016/j.ijantimicag.2019.06.012.

FIGURE CAPTIONS

FIGURE 1. Map of the p*bla*$_{TEM-20}$ plasmid vector, 5154 base pairs long, containing the *bla*$_{TEM-20}$ gene associated with the resistance to beta-lactam antibiotics, including penicillins and cephalosporins. The vector contained three replication origins (ori1, ori2, ori3) compatible with Gram-negative bacteria and the *repA* and *repB* genes encoding replication initiation proteins. Below is a nucleotide alignment of a 329 bp insert that contained a part of the *bla* gene with the GGT→AGT mutation, which resulted in a Gly→Ser substitution at codon 238, and the ATG→ACG mutation, which resulted in a Met→Thr substitution at codon 182.

FIGURE 2. Growth curves of *N. gonorrhoeae* (change in CFU over time), strain ATCC 49226, without plasmids (WT) and with p*bla*$_{TEM}$ plasmids, in the absence of antimicrobials (37°C, 5% CO_2). Dashed lines and dots show the experimental results, and solid lines indicate the theoretical curves constructed using equations (2)-(5). The inset shows the theoretical curves up to 24 hours of incubation to reach N_{assymp}.

FIGURE 3. Change in the CFU number of *N. gonorrhoeae* in the presence of ceftriaxone 0.03; 0.125; 2 mg/L. *N. gonorrhoeae* strain ATCC 49226 without the plasmid (A), with p*bla*$_{TEM-1}$ (B), p*bla*$_{TEM-135}$ (C), and p*bla*$_{TEM-20}$ (D) plasmids (37°C, 5% CO_2).

FIGURE 4. Dependences of the cell death rate constant k_{obs} on ceftriaxone concentration for *N. gonorrhoeae* strains without the plasmids and with p*bla*$_{TEM-1}$ and p*bla*$_{TEM-135}$ plasmids. The solid lines show the power-approximating curves constructed according to equations (10)-(12).

632 TABLE 1. Susceptibility of *N. gonorrhoeae* strains containing different p*bla*$_{TEM}$ plasmids to beta-
633 lactam antibiotics. S – susceptible, R – resistant, according to the EUCAST criteria (EUCAST, 2022)

Antibiotic	EUCAST criteria	MIC, mg/L			
		WT*	p*bla*$_{TEM-1}$	*pbla*$_{TEM-135}$	*pbla*$_{TEM-20}$
Benzylpenicillin (PEN)	S: $MIC_{pen} \leq 0.06$ R: $MIC_{pen} > 1$	0.25	16 (R)	32 (R)	16 (R)
Ampicillin (AMP)	–	0.125	8	8	16
Cefuroxime (CXM)	–	0.008	0.004	0.004	2
Ceftriaxone (CRO)	S: $MIC_{cro} \leq 0.125$ R: $MIC_{cro} > 0.125$	0.03 (S)	0.015 (S)	0.03 (S)	4 (R)
Cefixime (CFM)	S: $MIC_{cfm} \leq 0.125$ R: $MIC_{cfm} > 0.125$	0.015 (S)	0.015 (S)	0.015 (S)	16 (R)
Cefotaxime (CTX)	S: $MIC_{ctx} \leq 0.125$ R: $MIC_{ctx} > 0.125$	0.015 (S)	0.008 (S)	0.015 (S)	8 (R)
Cefepime (FEP)	–	0.03	0.015	0.015	16
Meropenem (MEM)	–	0.002	0.002	0.002	0.004
Imipenem (IPM)	–	0.004	0.008	0.004	0.008
Doripenem (DOR)	–	0.002	0.002	0.002	0.002

634 *WT – wild-type, *N. gonorrhoeae*, ATCC 49226, without p*bla*$_{TEM}$ plasmid

Insertion (329 b.p)

| *pvuI* | M182T | G238S | *bsaI* |

5'-TTACTCTGACAACGATCGGAGGACCGAAGGAG...ACCACGACGCCTGCA...TCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCA-3'
3'-AATGAGACTGTTGCTAGCCTCCTGGCTTCCTC...TGGTGCTGCGGACGT...AGACCTCGGCCACTCGCACCCAGAGCGCCATAGTAACGTCGT-5'

For-5'-TTACTCTGACAACGATCGGAGGACCG-3'   3'-TCGGTCACTCGCACCCAGAGCGCCATAGTAA-5'-Rev