Combining two Genetic Sexing Strains allows sorting of non-transgenic males for *Aedes* genetic control

Authors

Célia Lutrat [1,2,3,4,*], Myriam Burckbuchler [4], Roenick Proveti Olmo [4], Rémy Beugnon [5,6], Albin Fontaine [7], Omar S. Akbari [8], Rafael Argilés-Herrero [9], Thierry Baldet [1,10], Jérémy Bouyer [1,11,12,†], Eric Marois [4,†,*]

Affiliations

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France.

[2] ASTRE, CIRAD, INRA, Univ. Montpellier, Montpellier, France.

[3] Université de Montpellier, Montpellier, France.

[4] CNRS UPR9022, INSERM U1257, Université de Strasbourg, France

[5] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Puschstrasse 4, 04103 Leipzig, Germany

[6] Institute of Biology, Leipzig University, Puschstrasse 4, 04103 Leipzig, Germany

[7] Unité Parasitologie et Entomologie, Département Microbiologie et maladies infectieuses, Institut de Recherche Biomédicale des Armées (IRBA), Marseille, France.

[8] School of Biological Sciences, Department of Cell and Developmental Biology, University of California, San Diego, 92093, USA.

[9] Cicindela Ltd, Valencia, Spain

[10] CIRAD, UMR ASTRE, Sainte-Clotilde F-97490, Reunion, France

[11] CIRAD, UMR ASTRE, Saint-Pierre F-97410, Reunion, France

[12] Insect Pest Control Sub-Programme, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International Atomic Energy Agency (IAEA), Vienna, Austria

[†] These authors jointly supervised this work

[*] To whom correspondence should be addressed: celia.lutrat@outlook.com, e.marois@unistra.fr

Summary

Chemical control of disease vectoring mosquitoes *Aedes albopictus* and *Aedes aegypti* is costly, unsustainable, and increasingly ineffective due to the spread of insecticide resistance. The Sterile Insect Technique is a valuable alternative but is limited by the slow, error-prone, and wasteful sex-separation requirement. Here, we present four Genetic Sexing Strains (two for each *Aedes* species) based on fluorescence markers linked to the m and M sex loci, allowing for the isolation of transgenic males. Furthermore, we demonstrate how combining these sexing strains enables the production of non-transgenic males. Scaling-up would allow the sorting of 100,000 first instar male larvae in under 1.5 hours with an estimated 0.01-0.1% female contamination. Cost-efficiency analyses revealed that using these strains could result in important savings while setting up and running a mass-rearing facility. Altogether, these Genetic Sexing Strains should enable a major upscaling in control programmes against these important vectors.

Keywords

Graphical abstract

Introduction

Introduction

Aedes aegypti and *Aedes albopictus* are invasive mosquito species responsible for the transmission of

many pathogens including dengue (DENV), chikungunya (CHIKV), Zika (ZIKV) and yellow fever

(YFV) viruses (*1, 2*). Driven by climate change and worldwide trade, both vectors are spreading

rapidly and it is predicted that 49% of the world population will be at risk of *Aedes*-borne diseases by

2050 in the absence of effective control measures (*3, 4*).

Suppression of mosquito populations using genetic control is one of the most effective, sustainable

and environmentally friendly alternatives to insecticide use. It relies on repeated mass releases of non-

biting male mosquitoes – either sterile (the Sterile Insect Technique, SIT (*5*), and its derivatives

including pgSIT (*6*)), *Wolbachia* infected (the Incompatible Insect Technique, IIT(*7–9*)), both (*10,*

11), or carrying a lethal transgene (Release of Insects carrying a Dominant Lethal, RIDL) (*12*). For all

these interventions, an efficient sex separation method is required. Currently, *Aedes* mosquitoes are

sexed based on natural pupal size dimorphism using a Hoch's sorter, which can be either fully manual

(*13, 14*) or partly automated (*11*). This method, which requires homogenous pupal size and therefore

density optimised larval rearing conditions, suffers from female contamination rates between 0.8 and

1% and high daily and user-to-user variability (*15*). Notably, a multi-step pupal and adult sorter was

recently described which allowed 1.13×10^{-7}% female contamination (*9*). However, this system like

all other existing methods share the drawback of sorting at a late stage and recovering less than half of

the reared males (*14*), meaning that > 75% of total pupae are reared and fed in vain.

In *Anopheles* mosquitoes, transgenic genetic sexing strains (GSSs) allowing automated sex separation

of young larvae have been described (*16–19*). Fluorescent markers display male-specific expression,

either by using male-specific regulatory sequences or by linking markers to the Y chromosome. One

Anopheles coluzzii sexing strain is X-linked (*20*), rendering females more fluorescent than males. Sex

separation is accomplished using a Complex Object Parametric Analyzer and Sorter (COPAS) device,

which functions as a flow cytometer, sorting large particles according to their fluorescence.

Additionally, as an alternative to the release of transgenic males, a crossing scheme generating non-

transgenic male-only populations using COPAS was proposed (*21*). This scheme requires a strain

carrying a fluorescence marker on the Y chromosome and another carrying a fluorescence marker on

the X chromosome. Crossing non-transgenic (X-/X-) females from the first strain to transgenic (X+/Y-

) males from the second results in progeny consisting of transgenic (X+/X-) females and non-

transgenic (X-/Y-) males.

In *Aedes* mosquitoes, GSSs allowing sex-sorting of transgenic males exist. Two studies have proposed

the use of a repressible female-specific flightless phenotype (*22, 23*). In this system, transgenic males

and females are released together but adult females do not survive long as they are unable to fly. A

method combining such sex-sorting and sterilization has been developed and adapted to *Ae. aegypti*

for SIT (*6, 24*). However, this system requires perfect sex-sorting of the parental strains to be crossed,

otherwise fertile transgenic mosquitoes (some of them homozygous for a *Cas9* transgene) could be

released. In Ae. Aegypti, a red-eye GSS has also been developed and proven efficient for sorting at the

pupal stage but requires automation (*25*). Recently, we developed a GSS for *Ae. albopictus* carrying a

fluorescence marker within a masculinizing transgenic cassette, allowing automated separation of

transgenic males from non-transgenic females (*26*).

Here, we examined if a crossing scheme similar to the one for *Anopheles* could be designed in *Aedes*

mosquitoes despite the absence of heteromorphic sex chromosomes. In *Aedes*, sex is encoded by non-

homologous sex loci located on the first pair of autosomes. The male locus is called "M", while the

common locus is called "m"; thus, females are m/m and males are m/M. These sex loci are about 1.18

Mbp long in *Ae. aegypti* (*27*), and are delimited by antagonistic factors that protect them from

recombination (*28, 29*). They are embedded within a 63 Mbp region with strong male-female genetic

differentiation, in which recombination remains suppressed (*30*). Consequently, linking a fluorescence

marker transgene to the *Aedes* sex loci or to the surrounding non-recombining region might allow

automated sorting of either transgenic males directly from an M-linked strain, or of non-transgenic

males after crossing hemizygous m-transgenic males with wild-type females. In this work, we

developed two GSSs for both *Aedes* vector species, one linked to the M-locus and another to the m-

locus, using genome editing and transgenesis. We show that these GSSs can be used separately to

release transgenic mosquitoes, or combined to purify large populations of wild-type males. We discuss

the suitability of the method for mass-rearing and inundative releases.

Results

Obtaining four Genetic Sexing Strains

In *Ae. aegypti*, linkage of an e*GFP* marker transgene to the m and M loci was achieved by CRISPR-

Cas9 knock-in targeting the AAEL019619 gene located in the non-recombining region encompassing

the sex-loci (*30*) (**Figure 1a-b,** see **Methods** for further details). We isolated an *Ae. aegypti* M-linked

strain that we termed Aaeg-M, and an *Ae. aegypti* m-linked strain that we termed Aaeg-m. *GFP*

fluorescence expressed from a ubiquitous promoter allows sex separation at the first larval stage: in

Aaeg-M, males express *GFP* while females are non-transgenic (**Figure 1c**) and in Aaeg-m, males

express one copy of the *GFP* transgene, while females express two copies resulting in brighter

fluorescence (**Figure 1d**). Both lines were backcrossed 7 times to a Brazilian genetic background

(Bra). In *Ae. albopictus*, M-linkage of fluorescence markers was achieved by *piggyBac* homing

stimulated by sequences derived from the *Nix* gene and cloned in the *piggyBac* transgenesis plasmid

(**Figure 1e,** see **Methods**). This approach yielded at least eight lines with tight M-linkage out of about

60 screened *piggyBac* insertions (strikingly, no or weak M-linkage was obtained with these *piggyBac*

constructs devoid of *Nix* sequence). Among the tightly M-linked *Ae. albopictus* strains, we selected

one expressing YFP, that we termed Aal-M. Sequencing revealed that the transposon had landed in a

non-coding sequence in scaffold 16, located on 1q12 according to the latest genome assembly ((*31*).

To obtain a second sexing strain with m linkage in the absence of knowledge of the *Ae. albopictus* m

locus sequence, we screened >120 *piggyBac* random insertions (**Figure 1f**). The best m-linked

insertion line that we obtained showed a recombination frequency of 0.1%. Its transgenic cassette

harboured a *Cas9* transgene associated with a *DsRed* fluorescence marker and was flanked by lox

sites. A *Cas9* transgene being undesirable in a sexing strain, we excised it using CRE recombinase and

replaced it with an *eGFP* transgene. This *Ae. albopictus* m-linked strain was termed Aal-m.

Sequencing of the transposon's flanking genomic sequence indicated that it had landed into a highly

repeated region; hence, its exact genomic location could not be identified but matched several loci in

1q31 (see **Methods** and **Supplementary Data 1**). Both *Ae. albopictus* transgenic lines show a clear

sex-separation pattern in COPAS analyses (**Figure 1g-h**). All four lines were fluorescence-sorted and

screened at each generation. In the Aaeg-M line, a single recombination event was observed after 15

generations (>10,000 individuals screened in total). Consequently, we estimated that the Aaeg-M and

Aaeg-m lines recombine at approximately 0.01%. In Aal-M, four recombinant individuals were

observed in the tenth generation after successively screening a total of >50,000 individuals. These

recombinations are likely to have arisen from a single event, suggesting that recombination is

extremely rare in the Aal-M strain as well. Of note, all tested recombinants in Aal-M appeared to be

sterile.

Figure 1: Schematic of the procedures used to obtain linkage of a fluorescence marker to the M and m-loci in *Ae. aegypti* and *Ae. albopictus*. In. *Ae. aegypti*, a single CRISPR-Cas9 target was used to obtain an M-linked strain (**a**) and an m-linked strain (**b**) within 2- 4 generations. Both lines allow sex separation using a COPAS flow cytometer (**c-d**). In *Ae. albopictus*, M-linkage was obtained by *piggyBac* homing near the M-locus, requiring screening and testing for 4 generations (**e**). "pB Trp" = helper plasmid expressing piggyBac transposase. In *Ae. albopictus*, m-linkage was obtained by *piggyBac* random integration through intensive screening over 6 generations (**f**). The selected m-linked line was then modified to remove undesirable transgenes using CRE-recombinase and a *GFP* expressing plasmid docking into the attP site left-over after lox cassette excision. Both Aal-M and Aal-m allow sex separation using a COPAS flow cytometer (**g-h**).

Combining M and m-linked Genetic Sexing Strains for purifying non-transgenic males

A pilot experiment to obtain non-transgenic males using two GSS was performed in both *Ae. aegypti* and *Ae. albopictus*. In *Ae. aegypti*, 1,000 negative females from Aaeg-M were COPAS-sorted (**Figure 2a**) and crossed with 300 COPAS sorted hemizygous males from Aaeg-m (**Figure 2b**). Their progeny (Aaeg-CS) comprised 65,839 larvae, of which 32,960 were negative and supposedly males (**Figure 2c**). We COPAS-extracted 2,000 of those, all of which emerged as non-transgenic males. In *Ae. albopictus*, we applied the same crossing scheme using 850 negative females from the Aal-M strain crossed with 250 hemizygous males from the Aal-m strain (**Figure 2d-e**). In their progeny, we obtained 24,239 Aal-CS larvae, of which 12,173 were negative and presumably males (**Figure 2f**). Of

the 2,000 negative COPAS-extracted larvae, 1,468 pupae were recovered and visually verified,

revealing the presence of one contaminant non-transgenic female, which is consistent with the

estimated 0.1% recombination rate of the Aal-m strain.

Figure 2: Successive sortings leading to purification of non-transgenic males. Panels show the fluorescence

graphs log(Green) = f(log(Red)) displayed by the COPAS software. Gated regions were defined for sorting

populations of interest. **a.** Sorting of non-transgenic females from Aaeg-M strain. Total larvae = 6,079. **b.**

Sorting of hemizygous transgenic males from Aaeg-m strain. Total larvae = 711. **c.** Sorting of non-transgenic

males in the progeny of 1,000 non-transgenic females from Aaeg-M strain crossed with 300 hemizygous

transgenic males from Aaeg-m strain. Total larvae = 65,839. **d.** Sorting of non-transgenic females from Aal-M

strain. Total larvae = 6,797. **e.** Sorting of hemizygous transgenic males from Aal-m strain. Total larvae = 1,960.

f. Sorting of non-transgenic males in the progeny of 850 non-transgenic females from Aal-M strain crossed with

250 hemizygous transgenic males from Aal-m strain. Total larvae = 24,239.

Assessing the fitness of the Genetic Sexing Strains

The fitness of the selected M and m-linked lines, as well as of the intermediary colony from the sexing

scheme (non-transgenic females from the M-linked line crossed with hemizygous transgenic males

from the m-linked line), was analysed by examining the following parameters: sex ratio, fecundity

(number of eggs per female), egg hatching rate, and survival from larva to adulthood. In *Ae. aegypti*,

lines Aaeg-M, Aaeg-m and the intermediary colony from the crossing scheme (Aaeg-CS) showed

similar sex ratios to the Bra WT line (**Figure 3a, Supplementary Table 1 and Data 2**). The fecundity

of Aaeg-M and Aaeg-m was not different from that of Bra (WT), while Aaeg-CS had a higher

fecundity (**Figure 3b**). Aaeg-M eggs hatched similarly to Bra eggs, while Aaeg-m eggs and Aaeg-CS

eggs hatched significantly better (**Figure 3c**). Survival from first instar larvae to adult stages was not

significantly different in Aaeg-M, Aaeg-m and Aaeg-CS compared to Bra (**Figure 3d**). In *Ae.*

albopictus, the sex ratios of the Aal-M, Aal-m and Aal-CS lines were not significantly different from

that of the wild-type line BiA (**Figure 3a**). Aal-M had significantly higher fecundity than BiA (WT),

while Aal-m and Aal-CS had similar fecundities to WT (**Figure 3b**). Eggs from the Aal-M and Aal-

CS lines had a slightly higher hatching rate than eggs from BiA (WT), while Aal-m eggs hatched

similarly (**Figure 3c**). Survival from L1 larvae to adults was marginally increased in Aal-M and Aal-m

compared to BiA (WT), while it was not significantly different in the progeny of the crossing scheme

(Aal-CS, **Figure 3c**).

Figure 3: Fitness of the genetic sexing strains and of the intermediary colony yielding non-transgenic

males. In all panels, the black dots with vertical bars represent the mean value and the 95% CI in the *Ae. aegypti*

and *Ae. albopictus* assays, respectively. The thin coloured dots in the squares above and below estimates

represent individual data points (n), while the large coloured dots around estimates represent the mean values of

each replicate (N). Significant differences between the lines are specified in the figure (n.s.: no significant

difference. *: *p*-value < 0.1, **: *p*-value < 0.01, ***: *p*-value < 0.001). The percentages of males were compared

using a generalised linear model with a binomial distribution in the *Ae. aegypti* (**a**) and *Ae. albopictus* (**e**) lines.

Comparisons of the mean number of eggs laid by an individual were analysed using a generalised linear model

with a Poisson distribution in the *Ae. aegypti* (**b**) and *Ae. albopictus* (**f**) lines. Comparisons of the percentage of

egg hatch were analysed using a generalised linear model with a binomial distribution in the *Ae. aegypti* (**c**) and

Ae. albopictus (**g**) lines. Comparison of survival from first instar larvae to adult stages were analysed using a

linear model with the assumption of residual normality in the *Ae. aegypti* (**d**) and *Ae. albopictus* (**h**) lines.

Assessing the fitness of the males to be released

In an operational SIT approach, transgenic males from the M-linked lines or non-transgenic males

from the crossing scheme would be produced for release. With this aim in mind, we assessed their

fitness through competitiveness assays, flight tests and two-week survival tests under laboratory

conditions. In the competitiveness assay between transgenic and non-transgenic males in *Ae. aegypti*,

Aaeg-M performed equally to Bra (WT) and Aaeg-CS (**Figure 4a, Supplementary Table 1**). In *Ae.*

albopictus, given the higher hatching rate and fecundity of Aal-M compared to BiA (WT) and Aal-CS,

equal competitiveness would result in higher than expected percentages of Aal-M larvae in both

competitiveness assays. Aal-M showed a 0.59 competitiveness compared to BiA (WT) and a similar

competitiveness compared to Aal-CS (**Figure 4b**), which means that both Aal-M and Aal-CS had a

reduced competitiveness as compared to WT. In flight tests, *Ae. aegypti* transgenic males and non-

transgenic males from the crossing scheme performed as well as control males, while in *Ae.*

albopictus, the escape rates of Aal-M males were marginally lower and those of Aal-CS males

marginally higher than those of WT, with no significant difference (**Figure 4c-d**). Adult survival was

monitored for two weeks and no significant differences were observed between transgenic males, non-

transgenic males from the crossing scheme, and males from their respective WT lines (**Figure 4e-f**).

Figure 4: Fitness of transgenic and non-transgenic males compared to wild type. Black dots with vertical

bars represent the mean value and 95% CI in the competitiveness and male flight ability assays. The thin

coloured dots in the squares above and below the estimates represent individual data points (n), while the large

coloured dots around the estimates represent the mean values of each replicate (N). Significant differences

between the lines are indicated in the figure (n.s.: no significant difference. *: p-value < 0.1, **: p-value < 0.01,

***: p-value < 0.001). The competitiveness of males of the Aaeg-M line relatively to Bra (WT) or Aaeg-CS in

Ae. aegypti (**a**) and the competitiveness of males of the Aal-M line relatively to BiA (WT) and Aal-CS in *Ae.*

albopictus (**b**) were compared by measuring the number of transgenic and non-transgenic L1 larvae in the

progeny of each of the N = 5 replicates. Results were compared to the percentage expected if transgenic males

were as competitive as non-transgenic males (dashed line) and were analysed using a generalised mixed-effects

model with a binomial distribution and replicates as a random effect. The flight ability of transgenic and non-

transgenic males compared to the wild-type in *Ae. aegypti* (**c**), and *Ae. albopictus* (**d**) was tested by N = 3

replicates in a reference flight test device ("FTD"). The results were analysed using a generalised mixed-effect

model with a binomial distribution and replicates as a random effect. Survival of adult males 14 days after

emergence was monitored in *Ae. aegypti* (**e**) and *Ae. albopictus* (**f**) lines. N=4 replicates were performed for each

line. All values being above 90%, the y-axis is discontinuous between 0 and 90% for better visualisation. Male

survival at 7 and 14 days was compared using the log-rank test and neither comparison was significantly

different.

Determining the ideal COPAS-sorting speed

To assess the reliability of COPAS-sorting for high-throughput mass production of males in a

mosquito production facility, we evaluated the contamination rate as a function of sorting speed. A

pool of 10,000 Aaeg-CS first instar larvae (fluorescent females + non-fluorescent males) was

repeatedly COPAS-sorted and we measured the male recovery rate and the female contamination rate

at different sorting speeds. Sorting at a given speed was repeated N = 3 times. We observed that the

percentage of recovery decreased from 91.5 ±1.4% to 26.9 ±1.9% when increasing the sorting speed

from 6 to 300 larvae / sec, following a degree-two polynomial regression model ($R^2 = 0.983$), while

contamination remained absent below 300 larvae/sec (**Figure 5a**). We estimate that an operational

speed for sex sorting would be around 60 larvae/sec (which corresponds to 200 larvae/mL in the

reservoir water with our instrument settings) with a mean recovery rate of 69.6 ±1.1% and no female

contamination detected among more than 6,500 sorted males. At this speed, a small COPAS reservoir

(250mL) holds 50,000 larvae that are sorted in <14 minutes, yielding approximately 17,400 ± 270

males (considering a 50:50 sex ratio). Consequently, 100,000 males could be sorted in less than 1.5

hours. The same test run was repeated with *Ae. albopictus* larvae and gave comparable results (**Figure**

5a).

Figure 5: Evaluation of the feasibility of using GSS in *Aedes* SIT, and suggested improvement. a.
Evolution of male recovery and female contamination rates as a function of COPAS sorting speed. 10,000 *Ae.*
aegypti L1 larvae from the crossing scheme were COPAS-sorted at different concentrations, resulting in
different sorting speeds. Percentage of recovery (black dots) and percentage of contamination (red dots) were
measured on three replicates for each sorting speed. Recovery values were fitted using a degree-two polynomial
regression ($R^2 = 0.983$) and contamination values were fitted using a linear regression ($R^2 = 0.556$). The model
predictions were plotted by black and red lines, respectively, with 95% confidence intervals ribbons. The same
experiment was repeated with 4,000 *Ae. albopictus* L1 larvae from the crossing scheme and is shown on the
same graph. In this species, percentage of recovery (black triangles) and percentage of contamination (red
triangles) were measured on a single replicate for each sorting speed. **b-g.** Estimate of the cost-efficiency of
Genetic Sexing Strains for *Aedes* SIT by simulating the construction of a mass rearing and a release facility with
different sexing systems. Included sexing systems were the manual sorting of pupae using glass plates, "Size
manual" (dashed black lines), the automated sorting of pupae with robotic glass plates, "Size auto" (dashed grey
lines), the automated sorting of L1 larvae from a GSS using COPAS, "GSS" (light green lines), and the
combination of two GSS for automated sorting of non-transgenic L1 larvae, "GSS-CS" (dark green lines).
Different production objectives were considered: release of 5, 10, 20, 50 and 100 million males per week for 52
weeks per year. **b.** Comparison of the number of larvae present at any time in the rearing facility. **c.** Comparison
of the construction cost in US dollars for both the mass rearing and the release facilities. **d.** Comparison of the
equipment cost in US dollars for both the mass rearing and the release facilities (yearly cost estimated by
dividing the cost of each equipment by its lifespan). **e.** Comparison of the diet and consumable cost per year in
US dollars for both the mass rearing and the release facilities. **f.** Sum of the yearly costs for construction
(estimated lifespan = 20 years), equipment, diet and consumables in US dollar. **g.** Comparison of the number of
staff (labourers, managers etc.) needed for both the mass rearing and the release facilities. **h.** Proposed
chromosomal inversion that could prevent recombination between the fluorescence marker and the sex-locus.

Assessing the cost-efficiency of using transgenic GSS as compared to standard sex sorting in SIT programs

By allowing the separation of males from females at the first larval stage (L1), the numbers of subsequent larval stages and pupae to be reared in a mass-rearing facility is greatly reduced. However, filter colonies are required, as well as the acquisition of expensive fluorescence-sorting devices such as COPAS. Adapting the 'FAO/IAEA INTERACTIVE SPREADSHEET FOR DESIGNING MOSQUITO MASS REARING AND MALE HANDLING FACILITIES' (*32*) (see **Methods** and **Supplementary Data 3**), we estimated the number of insects to be reared, the facility sizes, the workforce, the construction, equipment and consumable costs *etc*. These estimates were compared between standard rearing and sorting procedures (hereafter referred to as 'Size manual' for manual sorting of pupae based on their size dimorphism (*13*), and 'Size auto' for automated sorting of pupae developed for the IIT-SIT in Guangzhou, China (*11*)) as well as the use of a single GSS (sorting transgenic males, referred to as 'GSS') and the use of two GSS combined for sorting non-transgenic males (referred to as 'GSS-CS'). Different release scales (namely 5, 10, 20, 50 and 100 million males to be released per week) were tested according to what is currently being done in genetic control field trials against *Aedes* (5 and 10M/week) and what would be operationally relevant for a broader *Aedes* control. We observed that both GSS and GSS-CS allow important reductions in the number of insects to be reared in comparison to the standard, no matter the release scale and including the filter colonies (GSS *vs.* Size manual = -65% larvae and -25-30% adults, GSS-CS *vs.* Size manual = -46% larvae and -8-17% adults, see **Fig. 5b** and **Supplementary Data 3**). These reductions translate into decreases in the mass-rearing facility size and therefore in the construction cost: facility size and construction cost are estimated to be reduced by 22-43% in GSS and 12-28% in GSS-CS compared to Size manual (**Fig. 5c**). Given the currently high costs of COPAS devices (from recent quotes, we considered $40k for a refurbished device), equipment costs are increased in GSS-CS compared to the manual sorting of pupae (+9 to +18% yearly, **Fig. 5d**) while GSS alone is estimated to have the lowest equipment cost from 10M males/week (-5 to -16% yearly in comparison to Size manual, **Fig. 5d**). However, the consumable costs, including insect diet costs, are also reduced and allow important savings starting at 20M males/week and above (GSS *vs.* Size manual = -54%, GSS-CS *vs.* Size manual = -37%, with >$50,000 savings at 20M males/week and above, **Fig. 5e**). In total, considering construction, equipment, diet and consumable costs, GSS is predicted to be the most economical option at all tested throughputs, while GSS-CS starts to be more economical compared to manual sorting of pupae from a 10M males/week throughput (**Fig. 5f**). Moreover, the workload, the cost of which can only be estimated on a country-by-country basis, is reduced by 29-38% with GSS and 18-27% with GSS-CS (**Fig. 5g**). For all comparisons, including equipment cost, the most expensive option is the automated sorting of pupae, as the devices are expensive (e.g. $40k for the automated sorter from(*11*), J. Bouyer) and the number of insects to be reared is high (**Fig. 5b-g**).

Discussion

Here we present the first genetic sexing strains (GSSs) for *Ae. aegypti* and *Ae. albopictus* based on m/M-linked transgenic fluorescent markers. We demonstrate that the M-linked Aaeg-M and Aal-M strains can be used alone to sex transgenic first instar larvae and that the M- and m-linked strains of each species can also be combined to produce a non-transgenic male population by two rounds of sex-sorting (of the parental strains and of the cross' offspring). In *Ae. aegypti*, transgenic males from the M-linked strain and non-transgenic males from the crossing scheme were found to be at least as fit and competitive as wild-type males of the same laboratory strain. In *Ae. albopictus*, a mild competitiveness reduction was observed in Aal-M transgenic males as well as in Aal-CS non-transgenic males. The reason for this is unclear, especially since Aal-CS males are non-transgenic. At a speed of 60 larvae/second using a COPAS device, we estimate that 100,000 males can be obtained in under 1.5 hour with the current settings, recovering 70% males and with a female contamination of 0.01-0.1%.

Unexpectedly, we observed rare recombination events between the transgene and the M-locus in the Aaeg-M line after screening tens of thousands of larvae. In this line, the transgene is located approximately 9 Mbp from *Nix*, which remains well within the presumed non-recombining region (*30*). Compared to the probability of recombination between two markers separated by the same distance on ordinary loci (3.4-4.7% according to the latest genomic information (*27*)), these results confirm that recombination is strongly reduced in this region, but not totally suppressed. Although not yet observed, the Aaeg-m line is expected to recombine at the same frequency as Aaeg-M, as both transgenes are inserted in the same target gene at a similar distance from their respective sex loci. Unfortunately, the recombination events imply that these GSSs will require regular purification of the parental colonies to avoid a gradual accumulation of recombinant individuals. Of note, such recombination events are easy to spot in single-sex batches of pupae due to size dimorphism and protandry, which will facilitate regular quality control of each strain.

Although our attempts at knocking-in a GFP marker into the large intron of *Nix* failed (not shown), improved GSSs may be constructed by targeting regions closer to or within the sex loci, especially in *Ae. aegypti* which benefits from a fully assembled genome. In *Ae. albopictus*, rare recombination events were also found in Aal-M amongst more than 10,000 individuals screened over 10 generations. For more accurate genome editing, better assembly of the first chromosome would be necessary in this species. The possibility of genetic instability under mass-rearing conditions and the appearance of aberrant sex recombinants have already been reported in the development of GSSs in other insect model organisms (*33*). Alternatively, the creation of viable chromosomal inversions spanning the sex-determining locus and the fluorescent transgene, akin to a balancer chromosome, could help suppress crossing-over (**Fig. 5h**) (*34, 35*). Finally, we have recently reported that transgenic *Ae. albopictus* pseudo-males masculinised by ectopic expression of *Nix* can be used as the Aal-M GSS in which the fluorescent marker will show an absolute linkage to sex (*26*). This type of GSS is particularly

interesting for the production of non-transgenic males, since in this case the wild-type females derived

from the Aal-M line and used in the crossing scheme are both non-transgenic and devoid of any

recombinant contamination. Compared to current sex-separation methods in *Aedes* mosquitoes (*9, 11,*

13, 36, 37), these GSSs allow a higher male recovery than any other sorting method based on pupal

size (about 70% *vs.* 30%). The female contamination rate (0.01 to 0.1% depending on recombination

events) is higher than with the Verily's multi-step sorter (*9*), but much lower than with manual [13] or

automated [11] glass sorters. Our system offers a sorting speed similar to that of the automated glass

sorter, which is higher than that of manual size sorters and multi-step sorters. Compared to all of these

currently available methods, the main advantage of our GSSs is to allow sorting at the earliest stage of

development, which results in significant savings in space, time and cost otherwise spent on rearing

unnecessary individuals. Moreover, reducing the number of females reaching the pupal stage in the

mass-rearing facility decreases the biting nuisance for the workers, a problem frequently observed

with pupal sorting methods, due to early adult emergences.

Additionally, we propose a crossing strategy that proved to be effective in obtaining non-transgenic

males. Compared to sorting pupae by size using glass plates or metal sieves, it has the advantage of

automating all sorting steps and achieving much lower female contamination rates. Although it

requires the rearing of two strains plus the sorting of males in the mass-rearing facility, this still

represents a reduction of approximately 46% in the number of mosquito larvae raised for current size-

based manual sortings. Despite the high cost of the sorting devices and the higher insect numbers than

for GSS, this method is predicted to be more cost-effective than the automated sorting of pupae at all

throughputs, and than manual sorting of pupae when exceeding a production throughput of 10M

males/week.

Although this work has focussed on the production of *Aedes* males for SIT, the same procedures for

the production of transgenic or non-transgenic male could be directly applied in other mosquito

control approaches such as IIT, RIDL, pg-SIT or other genetic control methods currently under

development.

Of note, GSSs with an m-linked fluorescence marker show less obvious sex separation than M-linked

GSSs. However, contamination of the sexing cross with females homozygous for the transgene yields

transgenic progeny that is subsequently eliminated when sorting the non-transgenic males. The release

of non-transgenic males from crossbreeding schemes may in fact be more appropriate than the release

of transgenic males in many countries, since GMO mosquito releases face public opposition and

regulatory constraints (*38*). Therefore, costs related to public information and consultation campaigns,

stakeholder engagement and compliance with regulatory processes may be reduced. These aspects

need to be considered as in the past, lack of information and consultation has led to such strong public

opposition that some release programmes have had to be cancelled (*38*). Of note, recombination in our

GSSs would result in contamination of released males by non-transgenic irradiated females at a rate equal to the recombination rate; however, this release of sterile females would still be much lower compared to current methods.

COPAS sorting errors during the purification of non-transgenic males were extremely rare in our pilot experiments. However, such contaminations may occur due to occasional technical problems, and would result in the release of a small percentage of transgenic mosquitoes amidst the non-transgenic. This should be absolutely avoided in settings where GMOs are banned. An additional quality control step to eliminate any residual GFP fluorescent larvae from the sorted larvae could be achieved by a second COPAS run. It could also be achieved by visual verification of the sorted larvae collectively under a fluorescence microscope, as GFP positives are easy to spot and eliminate, even among many wild-type larvae. Also, the irradiation of all mosquitoes before release will prevent leaking of the transgenes into the target population.

In conclusion, the four GSSs developed in this work represent very promising tools for improving the cost-efficiency of *Aedes* mass-rearing in genetic control programmes. According to our laboratory pilot tests, a facility harbouring two COPAS machines could produce 1,000,000 male larvae per day in a 7.5-hours regime. Considering that no females are reared, space constraints would be eased. Moreover, this technique would also reduce the production costs associated with the need to reach low or no female contamination, facilitating compliance with local regulations. Our results suggest that COPAS sorting of our GSSs would allow a rapid, cost-effective, and safe mass-production of male mosquitoes for SIT and other genetic control methods, with flexible up-scaling possibilities according to local needs, resources and regulations.

Acknowledgements

The authors thank Dr. H. Maiga and Dr. W. Mamai from the International Atomic Energy Agency (Vienna, Austria) for the revision of the SIT-related costs, Nathalie Schallon and Amandine Gautier for insectary operation, and Dr. Stéphanie Blandin for continuous support and scientific discussion.

This study was funded by EU ERC grant CoG—682387 REVOLINC to J.B. The contents of this publication are the sole responsibility of the authors and do not necessarily reflect the views of the European Commission. Mosquito production and insectarium operation were supported by Agence Nationale de la Recherche grant #ANR-11-EQPX-0022 and contrat triennal "Strasbourg capitale européenne" 2019-2021. Part of the mosquito experiments have been performed on the Baillarguet insectarium platform, member of the National Infrastructure EMERG'IN and of the Vectopole Sud network (http://www.vectopole-sud.fr/). The Baillarguet insectarium platform is led by the joint units Intertryp (IRD, Cirad) and ASTRE (Cirad, INRAE). E.M. received funding from Agence Nationale de la Recherche through grants #ANR-19-CE35-0007 GDaMO and # 18-CE35-0003-02 BAKOUMBA.

381 R.B. was supported by the International Research Training Group TreeDì funded by the Deutsche
382 Forschungsgemeinschaft (DFG, German Research Foundation)—319936945/GRK2324. O.S.A was
383 supported by NIH awards (R01AI151004, 5R21AI156018, 5RO1AI148300).

384 All figures were designed on BioRender.com under paid license.

Author contributions

387 C.L., T.B., J.B. and E.M. designed research. A.F. contributed to genomic data. C.L., M.B., R.P.O. and
388 E.M. performed research. C.L., R.P.O., R.B., A.F., O.S.A., R.A.H. and E.M. analysed data. R.B. and
389 R.A.H. contributed analytic tools. C.L. wrote the paper with inputs from all authors.

Declaration of interest

392 O.S.A. is a co-founder of Agragene, Inc. and Synvect, Inc. with an equity interest. The terms of this
393 arrangement have been reviewed and approved by the University of California, San Diego in accord-
394 ance with its conflict of interest policies. All remaining authors declare no competing interests.

Inclusion and diversity statement

397 We support inclusive, diverse, and equitable conduct of research.

 Figure titles and legends

Figure 1: Schematic of the procedures used to obtain linkage of a fluorescence marker to the M and m-loci

in *Ae. aegypti* **and** *Ae. albopictus*. In. *Ae. aegypti*, a single CRISPR-Cas9 target was used to obtain an M-linked

strain (**a**) and an m-linked strain (**b**) within 2- 4 generations. Both lines allow sex separation using a COPAS

flow cytometer (**c-d**). In *Ae. albopictus*, M-linkage was obtained by *piggyBac* homing near the M-locus,

requiring screening and testing for 4 generations (**e**). "pB Trp" = helper plasmid expressing piggyBac

transposase. In *Ae. albopictus*, m-linkage was obtained by *piggyBac* random integration through intensive

screening over 6 generations (**f**). The selected m-linked line was then modified to remove undesirable transgenes

using CRE-recombinase and a *GFP* expressing plasmid docking into the attP site left-over after lox cassette

excision. Both Aal-M and Aal-m allow sex separation using a COPAS flow cytometer (**g-h**).

Figure 2: Successive sortings leading to purification of non-transgenic males. Panels show the fluorescence

graphs log(Green) = f(log(Red)) displayed by the COPAS software. Gated regions were defined for sorting

populations of interest. **a.** Sorting of non-transgenic females from Aaeg-M strain. Total larvae = 6,079. **b.**

Sorting of hemizygous transgenic males from Aaeg-m strain. Total larvae = 711. **c.** Sorting of non-transgenic

males in the progeny of 1,000 non-transgenic females from Aaeg-M strain crossed with 300 hemizygous

transgenic males from Aaeg-m strain. Total larvae = 65,839. **d.** Sorting of non-transgenic females from Aal-M

strain. Total larvae = 6,797. **e.** Sorting of hemizygous transgenic males from Aal-m strain. Total larvae = 1,960.

f. Sorting of non-transgenic males in the progeny of 850 non-transgenic females from Aal-M strain crossed with

250 hemizygous transgenic males from Aal-m strain. Total larvae = 24,239.

Figure 3: Fitness of the genetic sexing strains and of the intermediary colony yielding non-transgenic

males. In all panels, the black dots with vertical bars represent the mean value and the 95% CI in the *Ae. aegypti*

and *Ae. albopictus* assays, respectively. The thin coloured dots in the squares above and below estimates

represent individual data points (n), while the large coloured dots around estimates represent the mean values of

each replicate (N). Significant differences between the lines are specified in the figure (n.s.: no significant

difference. *: *p*-value < 0.1, **: *p*-value < 0.01, ***: *p*-value < 0.001). The percentages of males were compared

using a generalised linear model with a binomial distribution in the *Ae. aegypti* (**a**) and *Ae. albopictus* (**e**) lines.

Comparisons of the mean number of eggs laid by an individual were analysed using a generalised linear model

with a Poisson distribution in the *Ae. aegypti* (**b**) and *Ae. albopictus* (**f**) lines. Comparisons of the percentage of

egg hatch were analysed using a generalised linear model with a binomial distribution in the *Ae. aegypti* (**c**) and

Ae. albopictus (**g**) lines. Comparison of survival from first instar larvae to adult stages were analysed using a

linear model with the assumption of residual normality in the *Ae. aegypti* (**d**) and *Ae. albopictus* (**h**) lines.

Figure 4: Fitness of transgenic and non-transgenic males compared to wild type. Black dots with vertical

bars represent the mean value and 95% CI in the competitiveness and male flight ability assays. The thin

coloured dots in the squares above and below the estimates represent individual data points (n), while the large

coloured dots around the estimates represent the mean values of each replicate (N). Significant differences between the lines are indicated in the figure (n.s.: no significant difference. *: p-value < 0.1, **: p-value < 0.01, ***: p-value < 0.001). The competitiveness of males of the Aaeg-M line relatively to Bra (WT) or Aaeg-CS in *Ae. aegypti* (**a**) and the competitiveness of males of the Aal-M line relatively to BiA (WT) and Aal-CS in *Ae. albopictus* (**b**) were compared by measuring the number of transgenic and non-transgenic L1 larvae in the progeny of each of the N = 5 replicates. Results were compared to the percentage expected if transgenic males were as competitive as non-transgenic males (dashed line) and were analysed using a generalised mixed-effects model with a binomial distribution and replicates as a random effect. The flight ability of transgenic and non-transgenic males compared to the wild-type in *Ae. aegypti* (**c**), and *Ae. albopictus* (**d**) was tested by N = 3 replicates in a reference flight test device ("FTD"). The results were analysed using a generalised mixed-effect model with a binomial distribution and replicates as a random effect. Survival of adult males 14 days after emergence was monitored in *Ae. aegypti* (**e**) and *Ae. albopictus* (**f**) lines. N=4 replicates were performed for each line. All values being above 90%, the y-axis is discontinuous between 0 and 90% for better visualisation. Male survival at 7 and 14 days was compared using the log-rank test and neither comparison was significantly different.

Figure 5: Evaluation of the feasibility of using GSS in *Aedes* SIT, and suggested improvement. a. Evolution of male recovery and female contamination rates as a function of COPAS sorting speed. 10,000 *Ae. aegypti* L1 larvae from the crossing scheme were COPAS-sorted at different concentrations, resulting in different sorting speeds. Percentage of recovery (black dots) and percentage of contamination (red dots) were measured on three replicates for each sorting speed. Recovery values were fitted using a degree-two polynomial regression ($R^2 = 0.983$) and contamination values were fitted using a linear regression ($R^2 = 0.556$). The model predictions were plotted by black and red lines, respectively, with 95% confidence intervals ribbons. The same experiment was repeated with 4,000 *Ae. albopictus* L1 larvae from the crossing scheme and is shown on the same graph. In this species, percentage of recovery (black triangles) and percentage of contamination (red triangles) were measured on a single replicate for each sorting speed. **b-g.** Estimate of the cost-efficiency of Genetic Sexing Strains for *Aedes* SIT by simulating the construction of a mass rearing and a release facility with different sexing systems. Included sexing systems were the manual sorting of pupae using glass plates, "Size manual" (dashed black lines), the automated sorting of pupae with robotic glass plates, "Size auto" (dashed grey lines), the automated sorting of L1 larvae from a GSS using COPAS, "GSS" (light green lines), and the combination of two GSS for automated sorting of non-transgenic L1 larvae, "GSS-CS" (dark green lines). Different production objectives were considered: release of 5, 10, 20, 50 and 100 million males per week for 52 weeks per year. **b.** Comparison of the number of larvae present at any time in the rearing facility. **c.** Comparison of the construction cost in US dollars for both the mass rearing and the release facilities. **d.** Comparison of the equipment cost in US dollars for both the mass rearing and the release facilities (yearly cost estimated by dividing the cost of each equipment by its lifespan). **e.** Comparison of the diet and consumable cost per year in US dollars for both the mass rearing and the release facilities. **f.** Sum of the yearly costs for construction (estimated lifespan = 20 years), equipment, diet and consumables in US dollar. **g.** Comparison of the number of staff (labourers, managers etc.) needed for both the mass rearing and the release facilities. **h.** Proposed chromosomal inversion that could prevent recombination between the fluorescence marker and the sex-locus.

Methods

Mosquito rearing

The *Ae. albopictus* wild-type colony (called BiA) was established from larvae collected in Bischheim (France, 48.36°N 07.45°E) in 2018. *Ae. aegypti* from the Bangkok genetic background obtained from MR4-BEI were used for molecular cloning and genetic engineering. *Ae. aegypti* strain BgR9, derived from Bangkok, and expressing *Cas9* under the control of the Exuperentia promoter and a *DsRed* reporter gene under the *Ae*PUb promoter was used for CRISPR/Cas9 knock-in experiments (plasmid sequence provided in **Supplementary Data 4**). The Bra *Ae. aegypti* strain originated from Juazeiro, Brazil and was provided to the IPCL in 2012 by Biofabrica Moscamed, a collaborative centre of the IAEA on the development of the SIT against mosquitoes. It was used for backcrossing the sexing transgenes.

Mosquitoes were maintained in standard insectary conditions (25-28°C, 75-80% relative humidity, 14-hr/10-hr light/dark cycle). Larvae were reared in 35 cm square pans in 1 L distilled water and daily fed ground TetraMin fish food. Adult mosquitoes were kept in 16x16x16-cm cages and provided a 10% sugar solution. Females were blood-fed on anesthetised mice. Eggs were laid on humid kraft paper and hatched or dried after 3 days.

Plasmid construction

Plasmid construction employed standard molecular biology procedures including PCR amplification of genomic DNA using Phusion™ High-Fidelity DNA Polymerase (F530, Thermo Scientific, France), cloning with NEBuilder® HiFi DNA Assembly (New England Biolabs, France), GoldenGate cloning in destination backbones using Anza 36 Eco31I restriction enzyme (Thermo Fisher Scientific, France), transformation in competent *Escherichia coli* bacteria, and Miniprep or Endo-Free Midiprep plasmid purifications (Macherey Nagel, France). For CRISPR/Cas9 knock-in in the *Ae. aegypti* M and m loci, we used plasmid pX4 and pX3 as repair donors, respectively (**Supplementary Data 4,** Addgene #183903 and #183904). For *piggyBac* integration near the *Ae. albopictus* M locus yielding the Aal-M strain, we used plasmid ppBAalbNixE1E3E4 PUb-YFP (Addgene #173666 described in (*26*)). For random *piggyBac* insertion near the m locus, we used plasmid ppBExu-Cas9-sv40, the lox cassette of which (encompassing *Cas9* and PUb-*DsRed*) was subsequently excised from the genome by injection of a plasmid expressing Cre recombinase under the control of the PUb promoter, and replaced by inserting plasmid pENTR-attB-PUb-GFP (Addgene #183911) into the *attP* site by co-injecting it with a plasmid expressing integrase from phage PhiC31 (Addgene #183966).

Microinjection

Injection mixes were composed of 400ng/µL of DNA in 0.5x PBS. *piggyBac* injection mixes were prepared as described in (*20*). For CRISPR-Cas knock-ins, the initial mix injected into the BgR9 *Ae.*

aegypti Cas9-expressing line comprised 84 µM gRNAs, 100 ng/µL repair plasmid and 2µM Scr7.

Given the very low number of transgenics obtained (1 individual out of >20,000 screened larvae), we

later injected the repair donor plasmid (190 ng/µL) with 5µM Scr7 and three plasmids expressing

different gRNAs under the control of different *Ae*U6 promoters (70ng/µL each plasmid). This method

gave significantly higher knock-in rates (25 GFP positive larvae out of about 4000 screened).

Embryo microinjection was performed essentially as published using a Nikon Eclipse TE2000-S

inverted microscope, an Eppendorf Femtojet injector and a TransferMan NK2 micromanipulator [18]

with the following modifications: eggs were hatched 3-8 days post-injection and first instar larvae

were screened under a Nikon SMZ18 fluorescence microscope, only G0 larvae showing transient

expression of the reporter gene were retained for subsequent outcross.

Ae. aegypti **transgenesis**

For targeting the sex loci, we exploited the work of Fontaine *et al.* showing a cluster of markers inside

a non-recombining 63 Mbp region encompassing the sex loci with high male-female genetic

differentiation and male heterozygosity consistent with Y chromosome-like null alleles (*30*)

(**Supplementary Table 3**). DNA sequences of these markers with putative Y-linked-like null alleles

were blasted on the assembled AegL5 genome (*27*) to identify a 4.9 Mb region extending from

160,092,339 bp to 165,014,624 bp on chromosome 1 in the AegL5 assembly. Within this non-

recombining region, we picked the gene AAEL019619, located ~9Mbp away from *Nix*. We used the

CRISPR/Cas9 system to knock-in a 2.3 kb fluorescence marker cassette (*eGFP* marker under the *Ae.*

aegypti poly-Ubiquitin promoter - PUb) flanked by 850-960 bp homology arms either within the fifth,

9 kb-intron of this gene (pX3 construct), or to create a synthetic intron within exon 5 (pX4 construct).

We obtained a first M-linked strain for *Ae. aegypti* following the injection of 200 eggs of the BgR9

strain, which expresses *Cas9* under control of the *Exuperentia* promoter, with the pX4 repair plasmid

and two synthetic gRNAs (GATGAATCATGGGGCGCCT[GGG] and

GATTCATCAATCAACGGAG[CGG], brackets indicate PAM). About 30 G0 larvae showing

transient GFP expression were crossed *en masse* to an excess of wild-type mosquitoes. Out of >

20,000 G1 larvae screened, a single transgenic male larva was found, raised to adulthood, and crossed

to WT females. Its progeny was composed of 100% eGFP positive sons and 100% eGFP negative

daughters, indicative of M linkage. We termed this *Ae. aegypti* M-linked strain Aaeg-M. Upon

injection of 600 BgR9 eggs with the pX3 repair plasmid and a mixture of three plasmids expressing

gRNAs (GTGGCATAGCGCCGTGTGGA[GGG], GTCTTAAATGAAAGAGGCG[AGG],

GTATCATGCGTATTGCGAG[AGG]) under the control of three different U6 promoters (gRNA

cloning vectors: **Supplementary Data 4**), 43 larvae with transient eGFP expression were recovered in

G0. Resulting 20 males and 20 females were crossed *en masse* to adults of the opposite sex. 25

transgenic G1 larvae were obtained, 10 of which were males carrying an M-linked insertion, 4 were

males carrying an m-linked insertion and 11 were females carrying an m-linked insertion. Proper

insertion site in AAEL019619 was confirmed by PCR in all tested individuals, which revealed that in

some of these mosquitoes the whole repair plasmid had integrated. From a single male devoid of the

plasmid backbone, an *Ae. aegypti* m-linked strain named Aaeg-m was established and further

characterized. Both Aaeg-M and Aaeg-m lines were backcrossed into a Brazilian (Bra) genetic

background for 7 generations.

556 *Ae. albopictus* transgenesis: obtaining M-locus linked strains

Following several unsuccessful attempts to knock-in a fluorescence marker near the *Ae. albopictus Nix*

gene using the CRISPR/Cas9 system, we took advantage of the frequent spontaneous insertion near

the M locus of *piggyBac* transposon constructs carrying *Nix* DNA sequences that we previously

reported (*26*). In the course of several transgenesis experiments, using *piggyBac* constructs carrying

Nix sequences, we obtained 12 independent transgenic insertions with obvious M-linkage of the

transgenesis reporter (100% fluorescent males) in addition to several *Nix*-expressing masculinized

female lines and to non-M linked, non-masculinizing insertions. From the third generation on, these

lines were amplified to confirm their M-linkage and estimate their recombination rate. Six lines

showed no recombination out of >700 individuals screened in each line until the sixth generation

(**Supplementary Table 4**). Most other M-linked lines also showed 100% fluorescent males but

contained additional, non-M-linked multiple insertions and were discarded. Based on line fitness and

COPAS profiles, we selected a YFP line termed Aal-M for further work as an *Ae. albopictus* M-linked

strain. Additional M-linked lines marked with OpIE2-GFP or Pub-DsRed showing no recombination

to date, carrying a docking *attP* site for subsequent transgenesis, and representing additional GSSs are

also being maintained to date but were not further characterized in detail.

573 *Ae. albopictus* transgenesis: obtaining an m-locus linked strain

We first attempted to obtain an m-linked fluorescence marker by screening through >120 random

piggyBac insertions. For this, we screened our existing collection of *Ae. albopictus* transgenic lines for

m/M-linkage and constructed an additional library of *piggyBac* constructs expressing various

fluorescence proteins (eGFP, mTurquoise2, YFP or DsRed) under the control of promoters showing

distinct expression patterns (*3xP3, OpIE2*, *Ae. aegypti PUb*, *Drosophila melanogaster actin5C*). By

performing multiplex injections of these *piggyBac* plasmids together, we obtained lines carrying up to

six distinct insertions as indicated by their colours and patterns of fluorescence. Out of 40 independent

founder transgenic individuals screened (most of which carrying multiple insertions), only two

insertions appeared m-linked: one carried an OpIE2-mTurquoise2 marker gene with a recombination

rate of about 3%, and the other, called m-albR9, carried a *Cas9* transgene with a *DsRed* reporter gene

and showed a recombination rate of 0.1%. We exploited the latter, in which transgenes are flanked by

two *lox* sites adjacent to an *attP* docking site. We injected m-albR9 eggs with a plasmid encoding Cre

recombinase to excise *Cas9* and *DsRed* transgenes, as *Cas9* is undesirable in a GSS. From

successfully excised individuals, we established an m-linked *attP* docking line, mX1, carrying no

further transgene. In a second step, an *attB*-containing plasmid carrying a PUb-eGFP marker cassette

was integrated into the *attP* site. The m-linkage of this new strain was verified by crossing eGFP

males to WT females and screening their offspring. The new *Ae. albopictus* m-linked line, named Aal-

m was made hemi/homozygous and amplified.

Automated larva sorting

Automated sorting of L1 larvae depending on their fluorescence level was performed using a large-

object flow cytometer named Complex Object Parametric Analyzer and Sorter (COPAS SELECT;

Union Biometrica, Belgium) as published [16], with the provided Biosort software. Sorting accuracy was

controlled under a Nikon SMZ18 fluorescence microscope.

Analysis of COPAS outputs

To determine the exact number of larvae in COPAS clusters, we built a clustering algorithm based on

manual outlier screening of particle size and fluorescence distribution, and an automated clustering

algorithm. All data handling and statistical calculations were performed using the R statistical software

version 4.1.0 (*39*). All R scripts used for this project can be found in our GitHub repository

(**https://github.com/Celia-Lutrat/Lutrat-et-al_2022_GSS**). Briefly, COPAS raw data were filtered

by the user via the visual examination of the individuals' size measurements (i.e., "log(EXT)" and

"log(TOF)") to remove outliers (e.g., egg debris and dust). A second filtering was performed on the

two first axes of a Primary Component Analyses (PCA) applied to the size variables (EXT and TOF).

A third manual filter was applied by examination of the individuals' fluorescence (i.e., "log(first

fluorescence)" and "log(second fluorescence)") . A final filtering was performed on the two first axes

of a PCA applied to fluorescence variables. PCA analyses were performed using the R function

'prcomp' from the package 'stat' (*39*) and the PCA outputs were extracted using the R package

'factoextra' (*40*). Filtered data were clustered using the R function 'kmeans' from the R package 'stat'

(*39*). Data were plotted using the R packages 'ggplot2' and 'ggpubr' (*41, 42*).

Targeted sequencing method

Genomic DNA was extracted using NEB Monarch HMW DNA extraction kit (T3060, New England

Biolabs, France) according to manufacturer instructions with the following modifications: after

addition of isopropanol, precipitated DNA was caught directly on a pipette tip and transferred to a new

tube containing 550 µl wash buffer and then to a final tube containing 70% ethanol. DNA was stored

in 70% ethanol at 4°C until use. On the day of use, ethanol was removed and DNA was resuspended in

water at 37°C with 500rpm agitation and gentle pipetting. DNA concentration was then measured

using a Nanodrop One device (Ozyme, France).

Sequencing method was adapted from the Nanopore Cas9 Targeted Sequencing (nCATS) (*43*).

gRNAs were selected in the known transgene region, with their PAMs towards the unknown genomic

sequence to be determined. This orientation ensured preferential ligation of Nanopore adapters on

DNA strands of interest.

Adapter ligation and library preparation were performed using the LSK109 kit (Oxford Nanopore

Technologies, UK) according to manufacturer instructions. Library was loaded on Flongle flow cells.

Sequencing, basecalling and alignment of reads to the transgenesis plasmid were performed using the

provided MinKNOW v. 21.06.10 software (Oxford Nanopore Technologies, UK). Reads that showed

homology to the transposon ends were then aligned to the transposon sequence using ClustalW on

Geneious software v9.1.8 (*44*) and one or several consensus sequences were determined. Finally, the

consensus sequences were compared to genomic data using the online NCBI BLAST service (*45*).

This method allowed us to determine the insertion site of the Aal-m transgene. We obtained only two

low-quality reads extending by about 20 kb into the genomic sequence adjacent to the transgene, from

which we designed a series of PCR primers. Sanger sequencing of PCR products spanning the

transposon/genomic junction allowed us to refine 1647 bp of the flanking sequence, provided in

Supplementary Data 1. Analysis of this sequence using NCBI BLAST showed only partial alignment

to sequences of the AalbF2 genome (*31*), presumably corresponding to a repetitive element,

suggesting that the m-linked region where the transgene inserted is not represented in the available

genome assemblies.

Assessment of lines' fitness

In transgenic lines, the sex ratio was measured by COPAS counting of the larvae of each fluorescence.

In non-transgenic lines (BiA and Bra), we reared samples to the pupal stage and counted males and

females under a binocular microscope based on genitalia examination. N=3 replicates were made for

all lines.

For fecundity, males and females were allowed to mate for several days. Cages were then offered a

blood meal and engorged females were selected. After 2 days, engorged females were shortly

anesthetised using CO_2 and placed individually in 24-well plates coated with wet filter paper for 2

days. Females were then released and each well was photographed under a Zeiss SteREO binocular

microscope. Eggs laid by each female were counted using the ImageJ software (*46*).

Egg hatching rate was measured on egg samples collected from N=3-5 cages of each line,

photographed under a Zeiss SteREO binocular microscope and counted using the ImageJ software

(*46*). Eggs were immersed in water, placed into a vacuum chamber for 20 to 30 minutes and allowed

to hatch overnight. The following morning, larvae were counted by COPAS.

Larva to adult survival was measured by sorting N=4-8 samples of 100-200 L1 larvae by COPAS and

rearing them to adulthood. The number of adults emerging from each batch was counted manually.

Assessment of males' fitness

Males' competitiveness was assessed by placing 30 transgenic males with 30 non-transgenic males in

a cage with 30 females and measuring the proportion of transgenic progeny in their offspring. If both

types of males are of similar fitness, it is expected that the progeny would be composed of ~25%

transgenic individuals (transgenic fathers being homozygous), corrected by the line sex ratio. Relative

competitiveness was estimated by comparing the observed proportion of transgenic progeny to the

theoretical value in N=4 or N=5 replicates.

Males' flight ability was assessed using a flight test device (FTD) as described (*47*). Tests were

performed on pools of about 100 males with N = 3 replicates for each condition.

Males' 2-week survival was measured by counting the daily number of deaths in N=4-8 cages of ~100

males of each line for 14 days.

Cost-efficiency analysis

The cost efficiency analyses were performed by comparing the outputs of the 'FAO/IAEA

INTERACTIVE SPREADSHEET FOR DESIGNING MOSQUITO MASS REARING AND MALE

HANDLING FACILITIES' (*32*) for different weekly production targets with that of its adapted

versions taking into account the use of a GSS or two GSS combined for sorting non-transgenic males

(see scheme in **Supplementary Data 3**). Briefly, these new spreadsheets were designed in a similar

way to the original one, but incorporate the GSS-specific structure previously developed in fruit flies

that includes separate filter and rearing colonies (*33*). In the filter colonies, first instar larvae are sorted

using COPAS, and sorted once more at the pupal stage using a classical pupal sorter. These colonies

produce the eggs feeding the rearing colonies at each generation. In a single GSS scenario, the rearing

colony is sorted as L1s for producing the males to be released. In a GSS-CS scenario, two separate

filter colonies produce the eggs for their respective rearing colony, which in turn are sorted and mated

to produce the eggs that will be hatched and sex-sorted as L1s for male-only releases. In these

scenarios, filter colonies are used at each generation, which is more careful than what is being done in

fruit flies despite a higher recombination rate (*33*). Input parameters can be read in **Supplementary**

Data 3. Of note, the costs mentioned in the spreadsheet have been implemented in euros (€) and

updated in July 2022. These costs are average observations from European SIT assays and were

revised by IAEA researchers. To facilitate downstream analyses, these Excel spreadsheets were

translated into an R (*39*) function. This function uses common R-base (*39*)functions as well as

functions from the 'purrr' and 'tidyverse' packages (*48*). It was deposited on our GitHub repository at

https://github.com/Celia-Lutrat/Lutrat-et-al_2022_GSS. The output costs were converted in US dollars

($).

Statistical analysis

All data statistical analyses were performed using the R statistical software version 4.1.0 (*39*), the R

scripts used for this study can be found in our GitHub repository (https://github.com/Celia-

Lutrat/Lutrat-et-al_2022_GSS).

The difference of sex ratio and egg hatching rate between the lines of each species was modelled using

the generalised model with binomial distribution and residuals distribution normality assumption (R

function 'glm' (*39*)), and pairwise comparison of the lines was tested using a Tukey HSD test (R

function 'glht' (*49*)). The difference in fecundity (number of eggs laid per female) between the lines of

each species was modelled using a generalised model with Poisson distribution and residuals

distribution normality assumption (R function 'glm'), and a pairwise comparison of the lines was

tested using a Tukey HSD test (R function 'glht'). The differences in larval survival between the lines

of each species was modelled using linear models with residuals distribution normality assumption (R

function 'lm' (*39*)), and tested using a Tukey HSD test (R function 'TukeyHSD' (*39*)). The difference

in male competitiveness and flight ability between the lines of each species was modelled using the

generalised mixed-effect model with binomial distribution and residuals distribution normality

assumption (R function 'glmer' from the package 'lme4' (*50*)), where technical replicates were set as

random factor. Pairwise comparisons between lines were tested using a Tukey HSD test (R function

'glht'). Male adult survival during the first 14 days was modelled using Cox regressions and Kaplan

Meir formula (function from the R packages 'survminer', 'survival' (*51–53*)). Differences in survival

at 7 and 14 days were modelled using linear models with residuals distribution normality assumption

(R function 'lm') and tested using a Tukey HSD test (R function 'TukeyHSD'). Model performances

were assessed using the R function 'check_model' from the package 'performance' (*54*). The

packages used, data handling procedures, model structure, fit and performances can be found in

Supplementary Data 2.

Data availability

All plasmid sequences are available in **Supplementary Data 4.** Plasmids are available from Addgene

under reference numbers #173666, #183903, #183904, #183911, #183912, #183913, #183914 and

#183966. Mosquito strains are available upon request from EM. All data needed to evaluate the

conclusions in the paper are present in the paper and/or in the **Supplementary Information.**

Code availability

The R function replicating all four spreadsheets is available at https://github.com/Celia-Lutrat/Lutrat-

et-al_2022_GSS. All other code used in this study is available from the authors upon reasonable

request.

Animal care

Animal care was in accordance with CNRS guidelines and approved by the CREMEAS ethics

committee.

Supplementary Information titles and legends
738

Supplementary Table 1: Summary and significance of statistical tests performed. More details
can be found in **Supplementary Data 2**.

Supplementary Table 2: Sequences of markers marking out and within the 63Mbp with high
male-female genetic differentiation in *Ae. aegypti*.

Supplementary Table 3: Cumulative number of fluorescent and negative males and females
between generations 3 and 6 in 11 *Ae. albopictus* lines carrying an M-linked transgene.
Underlined numbers are observations that may correspond to recombination events or to the presence
of additional, less sex-linked transgene insertions. Lines showing such events were thus discarded. The
selected M-linked line (Aal-M) is the one named "4.4Y" in the table.

Supplementary Data 1: 1239bp *Ae. albopictus* genomic sequence flanking the Aal-M transposon
and 1647 bp *Ae. albopictus* genomic sequence flanking the Aal-m transposon. Complete flanking
sequences and detailed explanations can be found in Supplementary Data 4.

Supplementary Data 2: Detailed statistical outputs

Supplementary Data 3: Production costs and initial parameters

Supplementary Data 4: Plasmid and genomic integration nucleotide sequences

1. pX4 plasmid used for CRISPR/Cas9 knock-in of a Pub-GFP cassette into exon 6 of the *Aedes aegypti* M-linked gene AAEL019619 (Addgene #183903)

2. Nucleotide sequence of three gRNA expression cassettes, each under the control of a different *Aedes aegypti* U6 promoter, used to knock-in the pX4 GFP cassette in the *Ae. aegypti* M locus (Addgene #183912, #183913, #183914)

3. pX3 plasmid used for CRISPR/Cas9 knock-in into the *Aedes aegypti* m-linked gene AAEL019619 (Addgene #183904)

4. ppBAalbNixE1E3E4 PUb-YFP plasmid that produced M-linked *piggyBac* insertion Aalb-M on *Aedes albopictus* chromosome 1 (Addgene #173666)

5. *piggyBac* plasmid containing *Cas9* under control of the *Exuperentia* promoter, used to obtain the m-linked m-albR9 *Ae. albopictus* line (Addgene #183905)

6. pENTRPUb-OptpBTransposase helper plasmid expressing a hyper-active, codon-optimized and NLS-modified piggyBac transposase under the control of the Ae. aegypti Polyubiquitin promoter

7. pattBPUb-GFP, attB site-containing plasmid expressing *GFP* under control of the *Ae. aegypti* PUb promoter, inserted in the m-linked attP site of *Ae. albopictus* strain mX1 derived from m-albR9 (Addgene #183911)

8. Helper plasmid (*piggyBac* construct) used to express PhiC31 integrase under control of the PUb promoter (Addgene #183966)

9. Genomic sequence flanking the Aalb-M insertion on the 3' piggyBac side.

10. Genomic sequence flanking the Aalb-m insertion on the 3' piggyBac side.

References

1.　N. J. Kassebaum *et al.,* Global, regional, and national disability-adjusted life-years (DALYs) for

315 diseases and injuries and healthy life expectancy (HALE), 1990Global Burden of Disease

Study 2015. *The Lancet*. **388**, 1603–1658 (2016), doi:10.1016/S0140-6736(16)31460-X.

2.　D. L. Vanlandingham, S. Higgs, Y.-J. S. Huang, Aedes albopictus (Diptera: Culicidae) and

Mosquito-Borne Viruses in the United States. *Journal of Medical Entomology*. **53**, 1024–1028

(2016), doi:10.1093/jme/tjw025.

3.　M. U. G. Kraemer *et al.,* Past and future spread of the arbovirus vectors Aedes aegypti and Aedes

albopictus. *Nature Microbiology*. **4**, 854 (2019), doi:10.1038/s41564-019-0376-y.

4.　S. Leta *et al.,* Global risk mapping for major diseases transmitted by Aedes aegypti and Aedes

albopictus. *International journal of infectious diseases*. **67**, 25–35 (2018),

doi:10.1016/j.ijid.2017.11.026.

5.　R. Gato *et al.,* Sterile Insect Technique: Successful Suppression of an Aedes aegypti Field

Population in Cuba. *Insects*. **12** (2021), doi:10.3390/insects12050469.

6.　M. Li *et al.,* Suppressing mosquito populations with precision guided sterile males. *Nature

Communications*. **12**, 5374 (2021), doi:10.1038/s41467-021-25421-w.

7.　N. A. Ahmad *et al.,* Wolbachia strain wAlbB maintains high density and dengue inhibition

following introduction into a field population of Aedes aegypti. *Philosophical transactions of the

Royal Society of London. Series B, Biological sciences*. **376**, 20190809 (2021),

doi:10.1098/rstb.2019.0809.

8.　A. Utarini *et al.,* Efficacy of Wolbachia-Infected Mosquito Deployments for the Control of

Dengue. *N Engl J Med*. **384**, 2177–2186 (2021), doi:10.1056/NEJMoa2030243.

9.　J. E. Crawford *et al.,* Efficient production of male Wolbachia-infected Aedes aegypti mosquitoes

enables large-scale suppression of wild populations. *Nature Biotechnology*. **38**, 482–492 (2020),

doi:10.1038/s41587-020-0471-x.

10.　The Project Wolbachia – Singapore Consortium, N. L. Ching, Wolbachia-mediated sterility

suppresses Aedes aegypti populations in the urban tropics. *medRxiv* (2021),

doi:10.1101/2021.06.16.21257922.

11. X. Zheng *et al.,* Incompatible and sterile insect techniques combined eliminate mosquitoes.

Nature. **572**, 56–61 (2019), doi:10.1038/s41586-019-1407-9.

12. D. O. Carvalho *et al.,* Suppression of a Field Population of Aedes aegypti in Brazil by Sustained

Release of Transgenic Male Mosquitoes. *PLOS Neglected Tropical Diseases.* **9**, e0003864 (2015),

doi:10.1371/journal.pntd.0003864.

13. R. W. Fay, H. B. Morlan, A mechanical device for separating the developmental stages, sexes and

species of mosquitoes. *Mosquito News.* **19**, 144–147 (1959).

14. P. A. Papathanos *et al.,* A perspective on the need and current status of efficient sex separation

methods for mosquito genetic control. *Parasites & Vectors.* **11**, 654 (2018), doi:10.1186/s13071-

018-3222-9.

15. W. Mamai *et al.,* Aedes aegypti larval development and pupal production in the FAO/IAEA mass-

rearing rack and factors influencing sex sorting efficiency. *Parasite (Paris, France).* **27**, 43 (2020),

doi:10.1051/parasite/2020041.

16. F. Bernardini *et al.,* Site-specific genetic engineering of the Anopheles gambiae Y chromosome.

Proceedings of the National Academy of Sciences. **111**, 7600–7605 (2014),

doi:10.1073/pnas.1404996111.

17. E. Marois *et al.,* High-throughput sorting of mosquito larvae for laboratory studies and for future

vector control interventions. *Malaria Journal.* **11**, 302 (2012), doi:10.1186/1475-2875-11-302.

18. F. Catteruccia, J. P. Benton, A. Crisanti, An Anopheles transgenic sexing strain for vector control.

Nature Biotechnology. **23**, 1414–1417 (2005), doi:10.1038/nbt1152.

19. N. L. Ntoyi *et al.,* Life-history traits of a fluorescent Anopheles arabiensis genetic sexing strain

introgressed into South African genomic background. *Malaria Journal.* **21**, 254 (2022),

doi:10.1186/s12936-022-04276-6.

20. G. Volohonsky *et al.,* Tools for Anopheles gambiae Transgenesis. *G3: Genes, Genomes, Genetics.*

5, 1151–1163 (2015), doi:10.1534/g3.115.016808.

21. C. Lutrat *et al.,* Sex Sorting for Pest Control: It's Raining Men! *Trends in Parasitology.* **35**, 649–

662 (2019), doi:10.1016/j.pt.2019.06.001.

22. G. M. C. Labbé, S. Scaife, S. A. Morgan, Z. H. Curtis, L. S. Alphey, Female-Specific Flightless

(fsRIDL) Phenotype for Control of Aedes albopictus. *PLOS Neglected Tropical Diseases*. **6**,

e1724 (2012), doi:10.1371/journal.pntd.0001724.

23. G. Fu *et al.,* Female-specific flightless phenotype for mosquito control. *Proceedings of the*

National Academy of Sciences. **107**, 4550–4554 (2010), doi:10.1073/pnas.1000251107.

24. N. P. Kandul *et al.,* Transforming insect population control with precision guided sterile males

with demonstration in flies. *Nature Communications*. **10**, 84 (2019), doi:10.1038/s41467-018-

07964-7.

25. P. Koskinioti *et al.,* Genetic sexing strains for the population suppression of the mosquito vector

Aedes aegypti. *Philosophical transactions of the Royal Society of London. Series B, Biological*

sciences. **376**, 20190808 (2021), doi:10.1098/rstb.2019.0808.

26. C. Lutrat, R. P. Olmo, T. Baldet, J. Bouyer, E. Marois, Transgenic expression of Nix converts

genetic females into males and allows automated sex sorting in Aedes albopictus. *Communications*

biology. **5**, 210 (2022), doi:10.1038/s42003-022-03165-7.

27. B. J. Matthews *et al.,* Improved reference genome of Aedes aegypti informs arbovirus vector

control. *Nature*. **563**, 501 (2018), doi:10.1038/s41586-018-0692-z.

28. C. L. Campbell *et al.,* Alternative patterns of sex chromosome differentiation in Aedes aegypti (L).

BMC Genomics. **18**, 1–11 (2017), doi:10.1186/s12864-017-4348-4.

29. E. Krzywinska *et al.,* The sex locus is tightly linked to factors conferring sex-specific lethal effects

in the mosquito Aedes aegypti. *Heredity*. **117**, 408–416 (2016), doi:10.1038/hdy.2016.57.

30. A. Fontaine *et al.,* Extensive Genetic Differentiation between Homomorphic Sex Chromosomes in

the Mosquito Vector, Aedes aegypti. *Genome Biology and Evolution*. **9**, 2322–2335 (2017),

doi:10.1093/gbe/evx171.

31. U. Palatini *et al.,* Improved reference genome of the arboviral vector Aedes albopictus. *Genome*

biology. **21**, 215 (2020), doi:10.1186/s13059-020-02141-w.

32. R. Argiles-Herrero, C. Cáceres, J. Bouyer, Eds., *Spreadsheet for Dimensioning Aedes Mosquito*

Mass-rearing and Release Facilities (Vienna, Austria, 2019).

33. J. S. Meza *et al.,* Comparison of classical and transgenic genetic sexing strains of Mediterranean

fruit fly (Diptera: Tephritidae) for application of the sterile insect technique. *PLOS ONE*. **13**,

e0208880 (2018), doi:10.1371/journal.pone.0208880.

34. D. L. Stern, Transgenic tools for targeted chromosome rearrangements allow construction of balancer chromosomes in non-melanogaster Drosophila species. *G3: Genes, Genomes, Genetics*. **12** (2022), doi:10.1093/g3journal/jkac030.

35. A. A. Augustinos *et al.,* Irradiation induced inversions suppress recombination between the M locus and morphological markers in Aedes aegypti. *BMC Genetics*. **21**, 142 (2020), doi:10.1186/s12863-020-00949-w.

36. J. Bouyer, H. Maiga, M. J. B. Vreysen, Assessing the efficiency of Verily's automated process for production and release of male Wolbachia-infected mosquitoes. *Nature Biotechnology* (2022), doi:10.1038/s41587-022-01324-z.

37. J. E. Crawford *et al.,* Reply to: Assessing the efficiency of Verily's automated process for production and release of male Wolbachia-infected mosquitoes. *Nature Biotechnology* (2022), doi:10.1038/s41587-022-01325-y.

38. A. Panjwani, A. Wilson, What Is Stopping the Use of Genetically Modified Insects for Disease Control? *PLOS Pathogens*. **12**, e1005830 (2016), doi:10.1371/journal.ppat.1005830.

39. R Core Team, R: A Language and Environment for Statistical Computing (2021).

40. A. Kassambara, F. Mundt, *factoextra: Extract and Visualize the Results of Multivariate Data Analyses* (2020).

41. A. Kassambara, *ggpubr: 'ggplot2' Based Publication Ready Plots* (2020).

42. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New York, 2016).

43. T. Gilpatrick *et al.,* Targeted nanopore sequencing with Cas9-guided adapter ligation. *Nature Biotechnology*, 433–438 (2020), doi:10.1038/s41587-020-0407-5.

44. M. Kearse *et al.,* Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)*. **28**, 1647–1649 (2012), doi:10.1093/bioinformatics/bts199.

45. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *Journal of Molecular Biology*. **215**, 403–410 (1990), doi:10.1016/S0022-2836(05)80360-2.

46. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image analysis.

Nature methods. **9**, 671–675 (2012), doi:10.1038/nmeth.2089.

47. N. J. Culbert *et al.,* A rapid quality control test to foster the development of genetic control in

mosquitoes. *Scientific Reports.* **8**, 1–9 (2018), doi:10.1038/s41598-018-34469-6.

48. H. Wickham *et al.,* Welcome to the Tidyverse. *JOSS.* **4**, 1686 (2019), doi:10.21105/joss.01686.

49. T. Hothorn, F. Bretz, P. Westfall, Simultaneous Inference in General Parametric Models.

Biometrical Journal. **50**, 346–363 (2008).

50. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using lme4.

Journal of Statistical Software. **67**, 1–48 (2015), doi:10.18637/jss.v067.i01.

51. T. Therneau, *A Package for Survival Analysis in R* (2021).

52. T. M. Therneau, P. M. Grambsch, Modeling Survival Data: Extending the Cox Model. *Statistics in*

Medicine. **20**, 2053–2054 (2001), doi:10.1002/sim.956.

53. A. Kassambara, M. Kosinski, P. Biecek, *survminer: Drawing Survival Curves using 'ggplot2'*

(2021) (available at https://CRAN.R-project.org/package=survminer).

54. D. Lüdecke, D. Makowski, P. Waggoner, I. Patil, performance: Assessment of Regression Models

Performance. *CRAN* (2020), doi:10.5281/zenodo.3952174.

Aedes aegypti

Aaeg-M

a.

CRISPR-Cas9

M-locus
m-locus — AAEL019619

Cas9

GFP   pU6 gRNA

G0 Transient expression (TE)

x

G1 Transgenic

x

G2 GSS

m m m

m m M

c.

log(Green) vs log(Red)

♂
♀

Aaeg-m

b.

CRISPR-Cas9

M-locus AAEL019619
m-locus

Cas9

G0 Transient expression (TE)

x

G1 Transgenic

x

G2-G3 Make homo/ hemizygous

x

G4 GSS

m m m

m m M

d.

log(Green) vs log(Red)

♀
♂

Aedes albopictus

Aal-M

e.

piggyBac homing near *Nix*

M-locus
Nix
Nix GFP

piggyBac transposon

pB int   Nix GFP
Nix YFP   Nix DsRed

G0 ~100 TE individuals

5 pools

G1 >100 transgenic ♂ Isolate 60 insertions

20 cages x 3 insertions

G2 Follow insertions of interest

4 indiv. males x 20 insertions

G3 Check presence of endogenous *Nix*

PCR

8 interesting lines => isolate and amplify

G4 GSS

m m m

m m M

g.

log(Green) vs log(Red)

♂
♀

Aal-m

f.

piggyBac random integration

DsRed

piggyBac transposon

pB Trp   GFP
mTurq   DsRed

G0 ~80 TE individuals

5 pools

G1 >100 transgenic ♂ Isolate 120 insertions

40 cages x 3 insertions

G2-G4 Look for insertions of interest

100s of individual crosses => Found 1 m-linkage

G5-G6 Make homo/ hemizygous

x

Cre recombinase + integrase, GFP

G15 GSS

m m m

m m M

h.

log(Green) vs log(Red)

♀
♂

d. Aal-M

e. Aal-m

f. Aal-CS

Mating competitiveness

a. 30 ♂ Bra (WT) / 30 ♂ Aaeg-M / 30 ♀ WT · 30 ♂ Aaeg-CS / 30 ♂ Aaeg-M / 30 ♀ WT — expected

b. 30 ♂ BiA (WT) / 30 ♂ Aal-M / 30 ♀ WT · 30 ♂ Aal-CS / 30 ♂ Aal-M / 30 ♀ WT — expected

Male flight ability

c. d.

Male survival (14 days)

e. f.

COPAS sorting efficiency

a.

Percentage of recovery and contamination (%) vs Sorting speed (larvae/sec) [larval concentration (larvae/mL)]

$R^2 = 98.3\%$

$R^2 = 55.6\%$

Measurement
- Recovery
- Contamination

Species
- Ae. aegypti
- Ae. albopictus

Cost-efficiency simulation per sexing method

b. Number of larvae in the mass-rearing facility vs Flying male production (M males / week)

c. Construction cost for both facilities ($) vs Flying male production (M males / week)

d. Equipment cost for both facilities (yearly cost, $/year) vs Flying male production (M males / week)

e. Diet and consumable yearly cost ($/year) vs Flying male production (M males / week)

f. Estimate of total yearly costs excluding staff ($/year) vs Flying male production (M males / week)

g. Total number of staff needed for both facilities vs Flying male production (M males / week)

method
- Size.manual
- Size.auto
- GSS
- GSS-CS

Balancer chromosome

h. Induce chromosomal inversion