A fungal endophyte-generated nucleoside signal regulates host cell death and
promotes root colonization

Nick Dunken[1], Heidi Widmer[1], Gerd U. Balcke[2], Henryk Straube[,3], Gregor Langen[1,4],
Nyasha M. Charura[1], Pia Saake[1,4], Lisa Leson[1], Hanna Rövenich[1,4], Stephan Wawra[1,4],
Armin Djamei[5], Alain Tissier[2], Claus-Peter Witte[3] and Alga Zuccaro[1,4*]

[1] University of Cologne, Institute for Plant Sciences, Cologne, Germany

[2] Department of Cell and Metabolic Biology, Institute of Plant Biochemistry, Halle (Saale),
Germany

[3] Department of Molecular Nutrition and Biochemistry of Plants, Institute of Plant Nutrition,
Leibniz University Hannover, Hannover, Germany

[4] Cluster of Excellence on Plant Sciences (CEPLAS)

[5] Department of Plant Pathology, Institute of Crop Science and Resource Conservation
(INRES), University of Bonn, Bonn, Germany

[*] Corresponding author: azuccaro@uni-koeln.de

Summary

Intracellular colonization of plant roots by the beneficial fungal endophyte *Serendipita indica*
follows a biphasic strategy. After an early biotrophic phase, the interaction switches to a
host cell death phase restricted to the root epidermis and cortex layers. Host cell death
contributes to successful fungal accommodation during beneficial interaction in *Arabidopsis*
thaliana. How the host cell death is initiated and controlled is largely unknown. Here we
show that two fungal enzymes, the ecto-5'-nucleotidase *Si*E5NT and the nuclease *Si*NucA,
act synergistically in the plant apoplast at the onset of cell death to produce deoxyadenosine
(dAdo), a potent cell death inducer in animal systems. Uptake of extracellular dAdo, but not
of the structurally related adenosine (Ado), activates a previously undescribed cell death
mechanism in *A. thaliana* and the basal land plant *Marchantia polymorpha*, suggesting a
conserved cell death response to dAdo across land plant lineages. Mutation of the *A.*
thaliana equilibrative nucleoside transporter *ENT3* confers resistance to extracellular dAdo-
triggered cell death and leads to decreased fungal-mediated cell death during root
colonization. A library screen of *A. thaliana* T-DNA insertion lines identified a nucleotide-
binding leucine-rich repeat protein (NLR) as an additional intracellular component in dAdo-
triggered cell death. Our data show that the combined activity of two apoplast-resident
fungal enzymes leads to the production of a metabolite, which induces an NLR-modulated
plant host cell death upon uptake.

In brief

Regulated host cell death is part of a plant defense strategy against pathogens but it is also
involved in accommodating certain beneficial root microbes. We have identified extracellular
metabolites and intracellular metabolic signals that contribute to beneficial root fungal
endophyte colonization, and uncovered a conserved cell death mechanism likely co-opted
for establishing plant-endophyte symbiosis.

Introduction

Regulated cell death occurs in plants as part of normal growth and development and in
response to abiotic and biotic stimuli. In plant-microbe interactions, host cell death programs
can mediate either resistance or successful infection. Depending on the type of microbial
lifestyle (biotrophic or necrotrophic), host cell death can benefit the plant by stopping growth
of biotrophs, or the microbe by promoting growth of necrotrophs. Thus, control of plant host
cell death is vitally important for determining the outcome of an interaction. Several research
lines indicate that controlled host cell death have a crucial role in certain beneficial
interactions, challenging the paradigm that cell death in plant-microbe interactions implies
pathogenesis or host-microbe incompatibility. Symbiosis with beneficial microbes, as well
as infection by pathogens, requires a sophisticated control of host defence and nutrient
fluxes. Certain features of beneficial microbes, such as the manipulation of host immunity,
metabolism and host cell death, are reminiscent of pathogen infections. In *Rhizobium*-
legume symbioses, root nodules are formed to provide a niche for bacterial nitrogen fixation.
The formation of infection pockets in some of these symbioses is associated with host cell
death and production of hydrogen peroxide (D'Haeze et al., 2003). Restricted cell death is
also observed in ectomycorrhizal symbioses (Mucha et al., 2014; Ragnelli et al., 2014), and
is required for the establishment of symbiotic interactions with the widely distributed
beneficial fungi of the order Sebacinales (Deshmukh et al., 2006; Lahrmann et al., 2013;
Qiang et al., 2012). Molecular environmental studies revealed that some of the most
common taxa of this order display little, if any, host-specificity and interact with a large
variety of plant species. Sebacinales isolates, including *Serendipita indica* and *S. vermifera*,
exhibit beneficial effects such as growth promotion, increased seed production and
pathogen protection, thereby playing a significant role in natural and managed ecosystems
(Oberwinkler et al., 2013; Tedersoo et al., 2014; Weiss et al., 2016). A requirement for
restricted host cell death in accommodation of certain beneficial microorganisms leads to
the hypothesis that the activation of cell death mechanisms in roots have a more important
ecological function than previously recognized.

In the hosts *Hordeum vulgare* (hereafter barley) and *Arabidopsis thaliana* (hereafter
Arabidopsis), fungi of the order Sebacinales initially colonize living cells which die as the
colonization proceeds (Deshmukh et al., 2006; Lahrmann et al., 2013; Lahrmann et al.,
2015; Lahrmann and Zuccaro, 2012; Qiang et al., 2012; Zuccaro et al., 2011). This
symbiosis-induced cell death is thought to contribute to niche differentiation during microbial
competition for space and nutrients in the root and is restricted to colonized host cells in the
epidermis and outer cortex. The host pathways controlling induction and execution of plant
root cell death and the fungal elicitors/effectors that initiate this process in the roots remain

largely unknown (Lahrmann et al., 2013; Nizam et al., 2019; Qiang et al., 2012; Schneider and Lynch, 2018). Pathogenic and beneficial fungi have large repertoires of secreted effectors that can manipulate host cell physiology and suppress plant defence which promotes fungal colonization. In fungi, effectors are mainly described from biotrophic and hemibiotrophic leaf pathogens (Lo Presti et al., 2015). By contrast, only few effectors have been functionally characterized from root symbiotic fungi (Kloppholz et al., 2011; Nizam et al., 2019; Nostadt et al., 2020; Plett et al., 2014; Voß et al., 2018; Wawra et al., 2016). Therefore, the modes of action of mutualistic fungal effectors remain poorly understood.

Using genomics, transcriptomics and proteomics we identified *Serendipita indica* secreted proteins present in the root apoplast, the space outside the plasma membrane, including the cell walls, within which material can diffuse freely (Nizam et al., 2019). One secreted fungal protein consistently found at different symbiotic stages is the ecto-5'-nucleotidase *Si*E5NT (PIIN_01005) (Nizam et al., 2019). Expression of *E5NT* is induced during colonization of barley and Arabidopsis hosts, but not in fungal axenic culture (Nizam et al., 2019). Animal ecto-5'-nucleotidases play a key role in the conversion of AMP to adenosine, counteracting, the immunogenic effects of extracellular adenosine-5'-triphosphate (eATP) released from stimulated host cells (Antonioli et al., 2013). Extracellular ATP is another essential signalling molecule in plants where it regulates development and responses to biotic and abiotic stresses. In Arabidopsis, eATP mediates different cellular processes through its binding to the purinergic membrane-associated receptor proteins DORN1/P2K1 and recently described P2K2 (Pham et al., 2020). In the apoplast, ATP accumulation increases cytoplasmic calcium and induces a defence response against invading microbes. We have recently shown that perception of extracellular nucleotides, such as eATP, is important in plant-fungus interactions via colonization assays of the Arabidopsis *dorn1* knock out mutant that is better colonized by *S. indica*. *Si*E5NT is capable of hydrolysing ATP, ADP and AMP to adenosine and phosphate, altering eATP levels in the apoplast and plant responses to fungal colonization. Expression of extracellular *Si*E5NT in Arabidopsis leads to increased *S. indica* colonization, confirming its role as an apoplastic effector protein. Considering the important role *Si*E5NT plays in fungal accommodation at early symbiotic stages, we proposed that modulation of extracellular nucleotide levels and their perception play a key role in compatibility during early plant-fungus interactions in roots (Nizam et al., 2019). Secreted *Si*E5NT homologues are present also in fungal pathogens as *Colletotrichium incanum* and *Fusarium oxysporum* and other endophytes of Arabidopsis such as *C. tofieldiae* where their expression is induced during colonization, suggesting that extracellular purines-based biomolecules play a role also in other plant-fungus interactions.

At the onset of root cell death a small fungal endonuclease, which we named *Si*NucA (PIIN_02121), is secreted together with *Si*E5NT during barley and Arabidopsis colonization by *S. indica* (Nizam et al., 2019; Thürich et al., 2018). In plants, mechanisms linking immune sensing of DNA danger signals in the extracellular environment to innate pathways in the cytosol are poorly understood as the role of (deoxy)nucleotide metabolism in root colonization and cell-death. Here we show that the synergistic activity of *Si*NucA and *Si*E5NT leads to the production of deoxyadenosine (dAdo) and, in smaller amounts, other

deoxynucleosides from extracellular DNA (eDNA) that is released by the roots upon
microbial colonization in a process named root extracellular trap (RET) formation or by
microorganisms during biofilm development (Chambard et al., 2021; Driouich et al., 2019;
Tran et al., 2016; Wen et al., 2009). The production of dAdo by extracellular fungal enzymes
bears resemblance to *Staphylococcus aureus* dAdo-mediated immune cell death in
animals, which appears to ensure bacterial survival in host tissues (Thammavongsa et al.,
2013; Winstel et al., 2018). Staphylococcal nuclease and AdsA (a homologue of *S. indica*
E5NT) are both required to release dAdo from neutrophil extracellular traps (NETs) which
has a potent cytotoxic effect on macrophages and other immune cells (Thammavongsa et
al., 2013). Here we demonstrate that extracellular dAdo, but not the structurally similar Ado,
activates a previously unidentified cell death mechanism in Arabidopsis which induces cell
death marker gene expression, activates the 26S proteasome, increases electrolyte
leakage and decreases photosynthetic activity (F_V/F_M). Metabolomic analyses of exudates
of dAdo-treated or *S. indica*-colonized Arabidopsis roots showed extracellular accumulation
of the plant metabolite methylerythritol cyclodiphosphate (MEcPP), a precursor of plastidial
isoprenoids and a stress-specific retrograde signalling metabolite (Xiao et al., 2012).
Consistently, expression of either extracellular *Si*NucA or *Si*E5NT *in planta* increased
colonization by *S. indica* and host cell death. We found that mutation in the Arabidopsis
equilibrative nucleoside transporter 3 (ENT3) leads to a strong resistance phenotype to
dAdo-induced cell death but not to exposure to methyl jasmonate, a cell death inducer of
senescence. Congruently, the *ent3* KO Arabidopsis line displays less fungal-mediated cell
death in the root compared to *S. indica* colonized WT and accumulates less of the
extracellular signalling metabolite MEcPP in response to fungal colonization or dAdo
treatment. Finally, in a mutant screening of Arabidopsis T-DNA insertion lines, we found
that a previously uncharacterized Toll-like Interleukin 1 receptor (TIR) domain type NLR is
involved in dAdo-triggered cell death. We propose that modification of extracellular
metabolites by the activity of the fungal derived enzymes *Si*NucA and *Si*E5NT provides a
direct link between purine salvage, immunity and cell death in the roots.

Results

Fading of host nuclei and vacuolar collapse are hallmarks of symbiotic cell death
during *S. indica* root colonization

S. indica induces cell death in colonized root cells of Arabidopsis and barley resulting in
characteristic cytological features at later colonization stages. In these two hosts the timing
and extent of cell death differ (Lahrmann et al., 2013; Qiang et al., 2012). In Arabidopsis the
cell death phenotype is less prominent than in barley but cytological analyses revealed
fading of host nuclei, vacuolar collapse (Figure 1A-M) and endoplasmic reticulum (ER)
swelling in colonized cells (Qiang et al., 2012), indicative of ER-stress and host cell death
during root colonization. To characterize the timing of fungal-mediated root cell death, the
presence and shape of plant nuclei during the progression of *S. indica* colonization were
monitored by confocal laser scanning microscopy either by the use of the nucleic acid stain
DAPI (4',6-Diamidino-2'-phenylindole dihydrochloride) and the cell death stain SYTOX

Orange (Figure 1C-I) or by the use of an Arabidopsis line expressing the nuclear marker
H2B:mCherry (Figure 1A, B). Nuclei stained with SYTOX Orange were visible 7 to 8 days
post inoculation (dpi), indicating that at this time point Arabidopsis plasma and nuclear
membranes were permeable to the dye, a hallmark for cell death (Figure 1F, G). In the
Arabidopsis H2B:mCherry line the plant nuclei in the epidermal layer were often stretched,
faded and finally disappeared around 8 to 10 dpi in heavily colonized areas of the root
(Figure 1A, B). *S. indica* hyphae embedded within the plant nuclei in dying cells were
frequently observed, suggesting that the fungus can digest and feed on host nuclear DNA
(Figure 1C-E, H, I). The cytological analyses show that Arabidopsis root cell death starts
around 7 dpi and at 10 dpi most host nuclei have faded or disappeared in *S. indica* colonized
cells.

170 *Si*NucA and *Si*E5NT act synergistically to produce deoxynucleosides

Although the interaction between *S. indica* and plant roots has been studied in detail,
comparatively little is known about the contribution of apoplastic effectors to fungal
accommodation or the mechanism of cell death in this system (Nizam et al., 2019; Nostadt
et al., 2020; Rafiqi et al., 2013; Wawra et al., 2016). We therefore analysed soluble
apoplastic proteins in barley at different stages of *S. indica* colonization. We found that
*Si*E5NT was one of the predominant fungal proteins consistently present in this extracellular
compartment (Nizam et al., 2019). We previously showed that *Si*E5NT functions as a
membrane bound nucleotidase that is released to the apoplast during host colonization.
*Si*E5NT is capable of liberating phosphate and adenosine from ATP, ADP and AMP (Nizam
et al., 2019). Additionally, the small secreted protein *Si*NucA with predicted endonuclease
activity (Figure S1), was found in the apoplastic fluid of colonised barley roots at the onset
of cell death (5 dpi) (Nizam et al., 2019). *Si*NucA is also secreted during root colonization in
Arabidopsis (Thürich et al., 2018). Its expression is transiently induced during cell death in
barley and Arabidopsis compared to axenic fungal growth according to transcriptomic data
(Lahrmann et al., 2013) and quantitative PCR analyses (Figure 2A). This prompted us to
examine the involvement of this small secreted protein in fungal colonization and host cell
death in more detail. Secretion of *Si*NucA and its enzymatic activity were assessed by
overexpression of a *Si*NucA:HA:His construct in *S. indica* (Figure 2B, S2). Supernatants of
fungal OE strains, but not of the empty vector control strains, as well as the affinity purified
*Si*NucA were capable of degrading single and double-stranded DNA and RNA derived from
fungal and plant material, suggesting a non-specific nuclease activity (Figure 2C). Addition
of magnesium and calcium increased activity of *Si*NucA in *in vitro* assays whereas EDTA
inhibited it (Figure S3).

To explore the effects of *Si*NucA on colonization, we tested independent homozygous T3
lines heterologously expressing a native version of *Si*NucA in Arabidopsis under control of
the 35S promoter. As controls we used WT and segregating lines without *Si*NucA gene.
Expression of *Si*NucA in Arabidopsis led to higher fungal colonisation at 7 dpi, which
correlated with its expression level (Figure 2D, E), demonstrating its importance in fungal
accommodation. Localization studies by confocal microscopy imaging of Arabidopsis roots

expressing either full length *Si*NucA or a version without the N-terminal signal peptide (SP)
fused to mCherry confirmed secretion of the full-length mCherry-tagged *Si*NucA fusion
protein to the apoplast and functionality of the SP. After plasmolysis with 1M NaCl or
sorbitol, the mCherry fluorescence signal was visible at the cell walls for the full-length
*Si*NucA fusion protein and not for the cytoplasmic version without SP (Figure S4).
Remarkably, the presence of *S. indica* led to a re-localization of the full-length mCherry-
tagged *Si*NucA to the host nuclei only in cells colonized by the fungus and not in bystander
cells where the mCherry signal remained at the cell periphery also after plasmolysis (Figure
2F and S4, S5). The presence of *Si*NucA in the apoplast (Nizam et al., 2019) and in the
nucleus of colonized host cells (Figure 2F) suggests that this small secreted protein can be
redirected to the nucleus when produced by the fungus and can act on both eDNA and
nuclear DNA.

The co-occurrence of *Si*NucA and *Si*E5NT in the apoplast at 5 dpi led us to speculate that
these two enzymes might cooperate in promoting fungal colonization of roots. Using *Si*E5NT
affinity purified from leaves of *Nicotiana benthamiana*, which transiently expressed the
enzyme as secreted C-terminal Strep-tagged variant, we found activity with dAMP and AMP
whereas no activity was detected for any other deoxynucleotides (dGMP, dCMP, dTMP) or
ribonucleotides (GMP, UMP, 3′,5′-cAMP) tested. Moreover, dATP and the general
phosphatase substrate *para*-nitrophenyl pyrophosphate (pNPP) were not *Si*E5NT
substrates. Interestingly, only dAMP was hydrolysed at a constant rate whereas the AMP
hydrolysis rate gradually declined under the chosen reaction conditions (Figure S6A, B).
The Km for AMP (15.9 μM) was ~20-fold lower than for dAMP (361.6 μM) but the k_{cat} for
dAMP (11.9 s^{-1}) exceeded that of AMP (1.1 s^{-1}) by a factor of ~10 (Figure 3A, B). These
data suggest that *Si*E5NT hydrolyses AMP slightly better at low substrate concentrations
(below 20 μM) but is by far more efficient for dAMP at higher substrate concentrations which
might prevail in a nucleus undergoing degradation and in case of eDNA degradation. To
test the ability of *Si*NucA and *Si*E5NT to work together, we incubated DNA with *Si*E5NT and
*Si*NucA alone and in combination. As expected, *Si*NucA alone degraded DNA (Figure 3C)
but neither deoxynucleotides nor deoxynucleosides were detected as reaction products by
LC-MS analysis indicating that *Si*NucA degrades DNA to oligonucleotides. *Si*E5NT alone
also did not release deoxynucleotides from DNA but released small amounts of
deoxynucleosides (Figure 3D). However, in combination *Si*NucA and *Si*E5NT released
deoxynucleosides from DNA with a strong preference for dAdo, a potent cell death inducer
in animal systems. This demonstrates the ability of *Si*E5NT to use oligonucleotides released
by *Si*NucA as substrate and to act synergistically with *Si*NucA to preferentially release dAdo
from DNA (Figure S6C).

dAdo induces cell death *in planta*

The production of dAdo and additional deoxynucleosides by the synergistic activity of the
two secreted fungal enzymes bears resemblance to *Staphylococcus aureus* dAdo-mediated
immune cell death in animal cells, which ensure the exclusion of macrophages from the
center of abscesses within which the bacteria survive (Thammavongsa et al., 2013; Winstel

et al., 2018). This motivated us to test the effect of dAdo *in planta*. Incubation of Arabidopsis
H2B-mCherry line with extracellular dAdo but not with Ado or buffer led to fading and
disappearance of nuclei in the roots within 48 hours (Figure 4A). This root cell death
phenotype was quantified via the use of the azo dye Evans Blue (Figure 4B, C).
Furthermore, extracellular dAdo triggered hallmarks of cell death such as increased
electrolyte leakage, induction of cell death marker gene expression as well as activation of
the 26S proteasome and decreased photosynthetic activity (F_V/F_M) (Figure 4D-H, S7). The
effects were concentration dependent and the removal of dAdo from the culture supernatant
24 hours after treatment led to recovery of Arabidopsis seedlings, suggesting that the
activation of the cell death program in Arabidopsis is still reversible at this stage (Figure S8,
S9).

A cell death phenotype was also observed in *N. benthamiana* young leaves during
expression via *Agrobacterium tumefaciens* infiltration of a *Si*E5NT construct with an N-
terminus signal peptide for secretion into the apoplastic space (Figure S10, S11) and in
seedlings incubated with dAdo (Figure S12). Cell death was not visible in older leaves or in
leaves of *N. benthamiana* expressing the control vector p19 or *Si*NucA alone. The observed
phenotype suggests that the presence of *Si*E5NT is sufficient to induce cell death in this
plant host upon wounding/damage by agroinfiltration which could release DNA and DNAses
into the apoplast or activate a response to the presence of the bacterium or its proteins
(Figure S10, S11). To test if dAdo-triggered cell death is conserved in basal plant lineages
we additionally tested its activity in the liverwort *Marchantia polymorpha*. Incubation with
dAdo led to cell death also in this plant species (Figure S13). Taken together, these data
indicate that the dAdo-mediated cell death mechanism is conserved across plant lineages.

dAdo induces the MEP pathway and accumulation of the stress signal metabolite MEcPP

To investigate the mechanism by which dAdo induces and regulates cell death in plants and
if it activates stress signalling pathways, we analysed the transcriptional response of
Arabidopsis to this metabolite at 0, 3 and 12 hours, compared to a buffer control by RNA-
seq. Most marker genes for regulated cell death (Olvera-Carrillo et al., 2015) were
upregulated at 12 hours (Figure 4I & Table S1-3). Additionally, the plastidial 2-C-methyl-D-
erythritol 4-phosphate (MEP) pathway was induced upon incubation with dAdo at 12 hours
in Arabidopsis seedlings (Figure 4J & Table S4). Congruently, MEcPP a precursor of
plastidial isoprenoids and a stress-specific retrograde signalling metabolite produced by the
MEP pathway, accumulated extracellularly at 3 days post treatment (dpt) measured via LC-
MS/MS (Figure 4J). Abiotic stresses and wounding elevate the levels of cytoplasmic
MEcPP, which coordinates stress-response pathways *in planta* (Xiao et al., 2012).
Detection of this isoprenoid intermediate in the extracellular environment upon dAdo
treatment suggests that this metabolite may serve as a stress signal in bystander cells, a
putatively important function that needs to be further investigated. Incubation of Arabidopsis
with extracellular MEcPP did not lead to cell death, demonstrating that accumulation of this

metabolite is not sufficient to trigger cell death and therefore it is not the cause of the
observed dAdo-mediated cell death (Figure S14).

dAdo-triggered signalling and cell death are not mediated by canonical pattern-
triggered immune responses

To test if dAdo-triggered cell death is mediated by signalling generated by an extracellular
immune receptor, we tested the ability of this metabolite to induce a fast apoplastic response
by monitoring calcium influxes and ROS production. Both responses are part of PTI
(pattern-triggered immunity), a process activated by the recognition of MAMPs (microbe-
associated molecular patterns) or DAMPs (damage-associated molecular patterns) by
pattern-recognition receptors (PRRs) at the plasma membrane (Ngou et al., 2021; Yuan et
al., 2021; Zipfel and Oldroyd, 2017).

Incubation of Arabidopsis seedlings with dAdo did not elicit a calcium influx whereas
treatment with ATP or ADP, which have been previously described as DAMPs (Choi et al.
2014; Tanaka et al. 2014), or with dATP or dADP elicited a fast calcium influx dependent
on the extracellular ATP receptor DORN1 (Figure 5 A-C). Treatment with dAdo or with any
of the other purine derivatives did not trigger a ROS burst (Figure S15). On the other hand,
incubation with dAdo and to a lesser extent with ATP or dAMP, but not Ado, led to
accumulation of the marker metabolite MEcPP and the induction of the DAMP/MAMP and
fungal-responsive At1g58420 gene at 3 dpt (Figure 5D, E & Table S5) (Choi et al., 2014;
Kilian et al., 2007; Nizam et al., 2019). Collectively, these data indicate that dAdo does not
act as a typical extracellular DAMP or MAMP but induces a signalling pathway independent
of calcium influxes and ROS production.

The *Arabidopsis* transporter ENT3 is required for dAdo-mediated signalling and cell
death

The fact that we did not observe a canonical PTI response to dAdo let us speculate that
uptake of this metabolite is necessary to promote plant signalling and induce cell death. In
the animal system, treatment with extracellular dAdo causes accumulation of intracellular
dATP, which affects DNA synthesis and triggers apoptosis via caspase 3 activation (Winstel
et al., 2018). Rapidly dividing cells are particularly susceptible to dAdo-induced apoptosis
(Figure 4 A,B) and the toxic activity of dAdo in the animal system was shown to depend on
dAdo uptake by the human equilibrative nucleoside transporter 1 (hENT1) (Winstel et al.,
2018). *S. indica* was not sensitive to dAdo and displayed normal growth also at high
concentrations (Figure S16). In Arabidopsis, eight potential members of the ENT family are
annotated in the genome. Two are expressed in roots, the plasma membrane localized
ENT3 and the tonoplast localized ENT1. The *Arabidopsis ent3* knockout (KO) line displayed
a strong resistance phenotype to dAdo-induced cell death compared to WT and *ent1* KO,
but is not affected by methyl jasmonate-induced cell death (positive control) (Figure 6 A-C;
S17). *At*ENT3 was shown to transport adenosine and uridine with high affinity and to be
inhibited in competition assays by a number of different purine and pyrimidine nucleosides
and 2'-deoxynucleosides, including dAdo (Li et al., 2003). This indicates that *At*ENT3 has a

broad substrate specificity and is a strong candidate for the uptake of extracellular dAdo. In
a competition assay, addition of extracellular Ado reduced the cell death phenotype induced
by dAdo, suggesting that these two extracellular metabolites compete for the same
transporter at the cell membrane (Figure 6D). Interestingly, *ENT3* is more strongly
expressed in the epidermis compared to the rest of the root (Rich-Griffin et al., 2020),
suggesting that dAdo-induced cell death could display a certain level of cell type specificity.
This is in line with the root cell death phenotype observed after dAdo treatment with the
Evans Blue staining (Figure 4B, C). Together, these data demonstrate that a functional
ENT3 plays an important role in dAdo-triggered cell death in Arabidopsis most likely by
importing dAdo into the cytoplasm where it activates signalling resulting in cell death.
Congruently, the *ent3* KO line accumulates lower amounts of extracellular signalling
metabolites including MEcPP, GSSG and 3′,5′-cAMP in response to dAdo treatment at 3
dpt compared to the WT line (Figure 6E ,Table S6).

Mutation of *At*ENT3 affects *S. indica*-mediated cell death and colonization

Next, we determined whether *At*ENT3 plays a role in fungal accommodation and *S. indica*-
mediated cell death in roots. The *ent3* KO Arabidopsis line displayed significantly less cell
death upon colonization by *S. indica* compared to the WT line at 7, 10 and 14 dpi (Figure
6G, H). Furthermore, the *ent3* KO line was significantly less colonized by *S. indica* at 8 dpi
compared to the WT control as determined by quantitative reverse transcription PCR
(Figure 6I).

Upon *S. indica* colonization we observed a transient accumulation of the extracellular
signalling metabolites 3′,5′-cAMP at 3 dpi (early biotrophic phase) and MEcPP at 6 dpi
(onset of cell death). Additionally, a higher level of free phosphate at 10 dpi was observed
(Figure 6F, Table S6). In accordance with the decreased cell death phenotype in the *ent3*
KO line, the amount of these metabolites was lower upon *S. indica* colonization compared
to the WT line, indicating that the *Arabidopsis* ENT3 is important for fungal-mediated
signalling and cell death in this host. Modification of extracellular metabolites by the activity
of the fungal derived enzymes *Si*NucA and *Si*E5NT and the host transporter ENT3 provides,
for the first time, a direct link between purine salvage, immunity and cell death in roots. How
the metabolic status of the host affects fungal-induced cell death needs to be thoroughly
elucidated.

An *At*NLR is involved in dAdo-mediated cell death

To identify downstream genetic determinants associated with dAdo-mediated cell death *in*
planta, we performed a mutant screening of 6868 SALK Arabidopsis T-DNA insertion lines
(Figure S18). Sensitivity to dAdo was tested using Arabidopsis Col-0 WT as control and the
SALK mutant lines grown on solid media (1/2 MS) in 24-well plates with and without 500
µM dAdo for 14 days. The dAdo-insensitive lines (survivors) from the screening were
subsequently analysed using the PAM fluorometry in three independent biological
repetitions together with the Col-0 WT line as control. Thirteen lines with different degrees
of dAdo-resistance were identified (Table S7). One of the survivor SALK lines had a

mutation at the locus At5g45240 (SALK_034517) encoding a nucleotide binding, leucine-
rich repeat receptor (NLR). Mutation in this gene led to a reproducible and significant high
resistance to dAdo-induced cell death resulting in decreased electrolyte leakage and higher
photosynthetic activity compared to the wild type Col-0 upon incubation with dAdo (Figure
7 A-E). The NLR locus At5g45240 is directly adjacent to *RPS4* and *RRS1* and is part of a
bigger locus with several TIR-NLR (TNL) genes, which are not functionally characterized
(Figure S19). The involvement of an NLR suggests that dAdo-induced cell death is
regulated by the host immune response in plants rather than a consequence of cytoplasmic
toxicity as suggested for the animal cells.

Discussion

Symbiotic cell death: an evolutionary partially conserved mechanism?

Root colonization by *S. indica* is associated with restricted host cell death. The mechanisms
behind the induction and regulation of this symbiotic cell death are poorly understood.
Earlier work showed that *S. indica* secretes E5NT, a ubiquitous fungal apoplastic enzyme
that catalyzes the conversion of adenosine nucleotides released by immune activated or
damaged host tissues to adenosine (Nizam et al., 2019). The implication of a secreted ecto-
5′-nucleotidase in host colonization resembles the mechanism evolved by *S. aureus* and
other animal pathogens to escape the host immune response by the secretion of AdsA that
converts eATP to Ado, a suppressor of immunity in animal cells (Thammavongsa et al.,
2009). In addition to the release of eATP by damaged animal cells, activated neutrophils
release NETs, extracellular matrices composed of nuclear and mitochondrial DNA armed
with granular proteins, cell-specific proteases, and antimicrobial peptides (Brinkmann et al.,
2004). NETs rapidly immobilize and kill bacterial pathogens (Driouich et al., 2019). *S.*
aureus escapes NETs by the secretion of proteases and the nuclease Nuc which degrade
antimicrobial peptides and DNA. Nuclease-mediated degradation of neutrophil NETs leads
to the formation of dAMP, which is converted by *S. aureus* AdsA into dAdo. Macrophages
and other immune cells are highly sensitive to dAdo intoxication. This cellular intoxication is
mediated by uptake via equilibrative nucleoside transporters (Thammavongsa et al., 2013;
Winstel et al., 2018). The *S. aureus* escape mechanism is based on the cooperative work
of two extracellular microbial enzymes and enables staphylococcus to block phagocyte
infiltration of abscess lesions and bacterial clearance. Similar to neutrophils, plant roots
secret RETs, root extracellular traps made of extracellular DNA and a variety of
antimicrobial compounds and polysaccharides. RETs and NETs exhibit similar
compositional and functional properties but unlike the NETs, RETs production is not solely
induced upon microbial infection but RETs are released continuously during root
development to form a large mucilage-rich network in the rhizosphere (Chambard et al.,
2021; Driouich et al., 2019; Hawes et al., 2016; Tran et al., 2016). *S. indica* is able to grow
within RETs and along the root tip (Figure S20) and to digest extracellular DNA via the
secretion of the nuclease NucA which accumulates in the root apoplast at the onset of cell
death (Nizam et al., 2019). Here, we show that the combined enzymatic activity of *Si*E5NT
and *Si*NucA, concomitantly present in the apoplast, catalyses the conversion of

deoxyadenosine nucleotides derived by nuclease digestion of extracellular DNA to
deoxyadenosine. Most importantly, we demonstrate that uptake via *At*ENT3 of
deoxyadenosine is required for host cell death in the roots. Mutation of ENT3 leads to
decreased cell death and impaired colonization by *S. indica*, substantiating the importance
of extracellular nucleosides uptake in modulating fungal accommodation and cell death. The
identification of a previously unknown immune-metabolic axis by which cells respond to
extracellular purine nucleosides and trigger cell death in plants suggests a certain
conservation or functional convergence between the immune evasion and escape
mechanisms evolved by *S. aureus* and other bacterial pathogens of animals and dAdo-
triggered cell death during plant-fungal endophyte interaction in the roots.

To ward off infections by microbial pathogens, plants have evolved immune receptors that
are essential for mounting a successful defense response. Plasma membrane localized
PRRs are able to perceive conserved MAMPs/DAMPs and induce a relatively mild PTI
immune response. Successful plant-associated microbes can deliver suites of effectors to
dampen PTI for successful host colonization. In turn, plants have evolved polymorphic
intracellular resistance (R) proteins to detect the presence and/or activity of the effectors,
which results in a robust defense response termed effector-triggered immunity (ETI) which
potentiate PTI (Ngou et al., 2021; Yuan et al., 2021). The majority of R proteins belong to
the nucleotide-binding (NB) leucine-rich repeat (LRR) (NLR) family, which is further
classified as Toll-like Interleukin 1 receptor (TIR) type NLRs (TNLs) or coiled-coil type NLRs
(CNLs) based on their N-terminal domains (Wu et al., 2019). TNLs generally act as sensors
of microbial effectors, while several CNLs are called helper NLRs and signal downstream
of many sensor NLRs in Arabidopsis (Saur et al., 2021; Wu et al., 2019). The implication of
a TNL in resistance to dAdo-mediated cell death in Arabidopsis suggest that this NLR could
be guarding a protein that is targeted by dAdo. Alternatively, it is possible that dAdo converts
into a plausible substrate of TNLs related to either nicotinamide adenine dinucleotide (NAD)
or to 2′,3′-cAMP, a noncanonical cyclic nucleotide monophosphate (cNMP) (Horsefield et
al., 2019; Wan et al., 2019; Yu et al., 2021). Plant TIR domains of NLRs are enzymes
capable of degrading NAD in its oxidized form (NAD^+). The TIR domain NADase function is
necessary but insufficient to activate plant immune responses. Recently, it was shown that
plant TIR proteins, besides being NADases, act as 2′,3′-cAMP/cGMP synthetases by
hydrolyzing RNA/DNA. Mutations specifically disrupting the synthetase activity abrogate
TIR-mediated cell death in *N. benthamiana*, supporting an important role for these cNMPs
in TIR signalling (Yu et al., 2021). Accumulation of extracellular 3′,5′-cAMP upon *S. indica*
colonization and dAdo treatment establish a link with cyclic nucleotide monophosphates.
Further analysis needs to be done in order to characterize the metabolism of dAdo inside
plant cells and how intracellular dAdo leads to activation of the TNL in Arabidopsis. As TIR
domain type NLRs are also present in animals, our findings open opportunities for further
study the implication of TNLs in dAdo-triggered cell death in plants and beyond.

In the animal systems, it was shown that after import of dAdo into macrophages, dAdo-
mediated toxicity involves conversion of dAdo to dAMP by the activity of deoxycytidine
kinase (DCK) and adenosine kinase (ADK) and signalling via subsequent conversion into

the corresponding di- and tri-phosphates by nucleotide kinases and activation of caspase-
3-induced apoptosis (Winstel et al., 2018). The absence of caspases in plants and the
implication of an NLR in dAdo-mediated cell death in Arabidopsis, strongly suggest that this
part of the pathway is not conserved between plants and animals and rely on different
regulation and execution mechanisms that need to be further studied.

In conclusion, we uncovered a cellular signalling pathway that responds to extracellular
DNA–derived metabolites, coupling extracellular nucleoside transport by an equilibrative
nucleoside transporter to cellular activation of the MEP pathway and cell death regulated
by an NLR. The fact that dAdo triggers cell death in different plant species, including a basal
one, suggests that this pathway is likely conserved and may represent an ancient cell death
mechanism that is co-opted for establishing plant-endophyte symbiosis. This paves the way
for a better understanding of immunometabolism in plant-microbe interactions.

Material and Methods

Fungal strains and culturing techniques

458 *S. indica* strain DSM11827 (Deutsche Sammlung von Mikroorganismen und Zellkulturen,
459 Braunschweig, Germany) was grown on complete medium (CM) with 2% (w/v) glucose and
460 1.5 % (w/v) agar as described before in (Hilbert et al., 2012). Liquid (CM) cultures were
461 incubated from spores at 28 °C & 120 rpm.

Plant lines and Arabidopsis transformation

Seeds of the *A. thaliana* ecotype Columbia 0 (Col-0), Col-0^{AEQ} and Col-0^{AEQ} *dorn1* (Choi et
al. 2014), SALK lines SALK_034517 (*nlr*) and SALK_204257(*ent3*) were used in the context
of the experiments. Additionally, Col-0 expressing histone H2B fused to mCherry
(H2B:2xmCherry) under the UBQ10 promoter in root cells (Marquès-Bueno et al., 2016)
were used in this study. Col-0 transformation was performed by the floral dip method as
described in (Nizam et al., 2019) to create the transgenic lines *35S::SiNucA* and
35S::SiNucA:mCherry (with and without signal peptide).

For *Marchantia polymorpha*, gemmae of male Tak1 and female Tak2 gametophyte were
used.

Plant growth conditions and fungal inoculation

Surface-sterilized seeds were germinated and grown on ½ MS medium (Murashige-Skoog-
Medium, with vitamins, pH 5.7) with 1 % (w/v) sucrose and 0.4 % (w/v) Gelrite under short
day conditions (8 h light, 16 h dark) with 130 µmol m^{-2} s^{-1} of light and 22 °C/18 °C.
For fungal inoculation, 7-day-old seedlings were transferred to ½ MS or ⅒ PNM (Plant
Nutrion Medium, pH 5.7) plates without sucrose (15-20 seedlings per plate). 1 ml water
containing 5x10^5 *S. indica* chlamydospores was pipetted on the root and the surrounding
area. Control plants were inoculated with sterile water.

For harvesting, single roots were thoroughly washed with water, a 4 cm long root part
starting 0.5 cm below the shoot cut and immediately frozen in liquid nitrogen. Two to three
plates of 20 seedlings each were pooled per replicate.

Confocal microscopy

Colonized Arabidopsis roots were stained with 5 µg/ml Wheat Germ Agglutinin Alexa Fluor
488 conjugate (WGA-AF 488, Life Technologies, Thermo Fisher Scientific, Schwerte,
Germany) or 1 µM FGB1-FITC from (Wawra et al., 2019) to visualize the fungal cell wall.
250 ng/ml 4',6-Diamidino-2'-phenylindole dihydrochloride (DAPI) or 500 nM SYTOX Orange
(Life Technologies, Thermo Fisher Scientific, Schwerte, Germany) were used as nucleic
acid stains.

For live cell confocal laser scanning microscopy, a TCS SP8 confocal microscope (Leica,
Wetzlar, Germany) was used. Excitation of AF 488 and FITC was done with an argon laser
at 488 nm and the emitted light was detected with a hybrid detector at 500-550 nm. mCherry
and SYTOX orange were excited with a DPSS laser at 561 nm and the signal was detected
with a hybrid detector at 590-660 nm. DAPI was excited with a diode laser at 405 nm and
the emitted light was detected with a hybrid detector at 415-460 nm.

DNA & RNA extraction

DNA was extracted from frozen root or fungal material as described in Wawra et al. 2016.
In short, ca. 500 mg ground frozen material was dissolved in 1 ml CTAB extraction buffer
(100 mM TrisHCl pH 7.5, 50 mM EDTA pH 8, 1.5 M NaCl, 2 % (w/v)
Cetyltrimethylammonium bromide, 0.05 % (v/v) ß- mercaptoethanol) and homogenized for
10 min. 500 µl of chloroform:isoamylalcohol mixture (24:1) was added and the tubes were
mixed for 5 min and centrifuged. Ethanol and 1 volume chloroform:isoamylalcohol mixture
(24:1) were added to the upper phase and centrifuged again. The DNA in the upper phase
was precipitated with 1 volume isopropanol at 4 °C for 1 h.

RNA was extracted with TRIzol (Invitrogen, Thermo Fisher Scientific, Schwerte, Germany)
and DNA digested with DNase I (Thermo Fisher Scientific, Schwerte, Germany) according
to manufacturer's instructions. cDNA was synthesized using the Fermentas First Strand
cDNA Synthesis Kit (Thermo Fisher Scientific, Schwerte, Germany).

Quantitative RT-PCR analysis

For quantitative real-time PCR the 2x GoTaq qPCR master mix (Promega, Mannheim,
Germany) was used. 500 nM primer forward and reverse each and 10-20 ng template cDNA
or gDNA were added. The reaction was running in a CFX connect real time system (BioRad,
Munich, Germany) with the following program: 95°C 3min, 95°C 15s, 59°C 20s, 72°C 30s,
40 cycles and melting curve analysis. Relative expression was calculated using the $2^{-\Delta\Delta CT}$
method (Livak and Schmittgen 2001). All used Oligonucleotides can be found in Table S8.

Serendipita indica transformation

S. indica protoplasts were transformed by the PEG-mediated transformation system as described in (Wawra et al., 2019).

Nuclease activity assay

To test nuclease activity of the purified protein, 10 nM *Si*NUCA were mixed with 100 ng linearized plasmid, 1 µg gDNA or 1 µg RNA in buffer (5 mM Tris pH 8, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 0.1 % microelement solution (from CM medium)). The mixture was incubated for 1 to 30 min at RT, loading dye was added and the samples run on a 1-2 % agarose gel.

To test nuclease activity in *S. indica* culture filtrate, a 5-day-old CM culture was crushed and incubated for another 3 days. The culture was filtered through Miracloth (Millipore Merck, Darmstadt, Germany) and 50 µl culture filtrate were added to linearized plasmid DNA, gDNA or RNA.

*Si*NucA-HA-His purification

A 7-day-old liquid culture of *S. indica* grown in CM medium was filtered through Miracloth. The mycelium was washed with 0.9 % NaCl and crushed in a blender (MicrotronR MB550 homogenizer (Kinematica, Luzern, Switzerland)) in fresh CM medium. The culture was regenerated for two days. Subsequently, the culture was filtered with Miracloth and through a 0.45 µm membrane filter. To this cell-free culture filtrate 1 mM Phenylmethanesulfonyl fluoride (PMSF) was added and the pH was adjusted to pH 7 with 1 M Tris pH 8. The proteins were precipitated using 80 % ammonium sulfate. The protein pellet was resuspended in 20 mM Tris pH 8. The proteins were separated by size exclusion chromatography (Sephadex G 200 column, Hiload 6/600) usinga 20 mM Tris pH 8 / 150 mM NaCl buffer. Fractions containing *Si*NucA-HA-His were desalted by dialysis and checked by SDS-PAGE and anti-HA western blot. The protein was stored in 20 mM Tris pH 8. Identity of the protein was checked by LC-MS/MS.

Protein purification and measurement of enzymatic kinetics of *Si*E5NTs

*Si*E5NT fused to a C-terminal haemaglutinin (HA) and Strep tag (pXCScpmv-HAStrep, V69) was transiently expressed in *Nicotiana benthaminana* and purified by affinity chromatography as described by (Myrach et al., 2017; Werner et al., 2008). Enzyme concentrations were determined using bovine serum albumin standards after SDS-PAGE and Coomassie staining employing an Odyssey Fc dual mode imaging system (Li-cor Biosciences, Germany).

To determine the substrate specificity, different substrates were tested. Briefly, 10 µl purified enzyme (1.9×10^{-4} mg protein) were added to 70 µl reaction buffer (10 mM MES buffer, pH 6.0; 1 mM $CaCl_2$ and 1 mM $MgCl_2$) and pre-incubated for 5 min. Substrates were added (20 µl of a 5 mM stock solution) reaching a final reaction volume of 100 µl. The reaction was quenched with 400 µl MeOH at the indicated time points and centrifuged for 10 min at

21.000*g* at 4°C. The supernatant was evaporated in a vacuum concentrator and the dry
pellet resuspended in 100 µl HPLC mobile phase A.

The samples were analyzed on an Agilent 1200 SL HPLC system equipped with a diode
array detector. 10 µl of the sample was injected onto a Supelcosil LC–18–T column (Sigma-
Aldrich) with a flowrate of 0.8 ml min^{-1} and a column temperature of 25°C. The analytes
were separated by the following gradient: 0 min, 100% A; 9 min, 100% A; 15 min, 75% A;
17.5 min, 10% A; 19 min, 0% A; 23 min, 0% A; 24 min, 100% A; 30 min, 100% A. The mobile
phase A consisted of 100 mM KH_2PO_4, pH 6.0, in deionized water and the mobile phase B
consisted of 90% 100 mM KH_2PO_4, pH 6.0, in deionized water and 10% MeOH. Respective
standard solutions were used for quantification. The data were analyzed using the Agilent
Chemstation software.

The kinetic constants for deoxyadenosine monophosphate and adenosine monophosphate
were determined using the EnzCheck (Thermo Fisher Scientific, Waltham, USA)
phosphatase assay kit according to specifications but using only one tenth of the
recommended reaction volume.

Release of deoxynucleosides from salmon sperm DNA by *Si*NucA and *Si*E5NT

10 µg Salmon sperm DNA were incubated with 4 µl *Si*NucA, 10 µl *Si*E5NT or a combination
of both in a final volume of 40 µl containing 10 mM HEPES, pH 7.2; 100 mM $MgCl_2$ and 1
mM DTT. Before adding the DNA, the enzymes were incubated for 5 min in the buffer
mixture. The reaction was incubated at 22°C for 1 h and inactivated at 95°C for 5 min
followed by centrifugation for 20 min at 40000*g* and 4°C. 20 µl of the supernatant were
analyzed on a 2% agarose gel. The remaining 20 µl of the supernatant were mixed with 50
µl water and used for HPLC MS/MS analysis. Samples and standards were analyzed using
an Agilent 1290 Infinity II HPLC System coupled with an Agilent 6470 triple quadrupole
mass spectrometer. Analytes (10 µl samples) were separated on a 50 x 4.6 mm Polaris
C18A column (Agilent) using the following gradient: 0 min, 96% A; 8 min, 35% A; 8.2 min,
0% A; 10 min, 0% A; 10.1 min, 96% A. Mobile phase A consisted of 10 mM ammonium
acetate, pH 7.5, and mobile phase B was pure MeOH. The flow rate was 0.6 ml min^{-1} and
the column temperature was 30°C. The in-source parameters were set as described
previously for deoxyribonucleoside analysis (Straube et al., 2021). All analytes were
measured in positive mode. Transitions (precursor ions and product ions), collision energies
and fragmentor energies are listed in Table S9.

PAM fluorometric measurements

The measurements of PAM fluorometry were performed by transferring 9-day-old
Arabidopsis seedlings into 24 well plates containing 2 ml of 2.5 mM MES buffer (pH 5.6). 3
seedlings were pooled into one well. After 24 h of regeneration, the seedlings were treated
with solutions of 5' deoxyadenosine (dAdo), adenosine (Ado) or methyljasmonate (all
Sigma-Aldrich, Taufkirchen, Germany), adjusting to an end concentration of 500 µM. The
treated seedlings were incubated for 20 minutes in total darkness to reach a dark-adapted

state. The photosynthetic activity of the plants was measured using the M-Series PAM
fluorometer (Heinz Walz GmbH, Effeltrich, Germany). The data was evaluated using the
ImagingWin software (v.2.41a; Walz, Germany). If further analysis of the development of
photosynthetic area was required, it was evaluated with Fiji (ImageJ).

Cell death staining with Evans Blue

A modified protocol as described in (Vijayaraghavareddy et al., 2017)was used.To quantify
S. indica induced cell death in *Arabidopsis*, plants were used for cell death staining at three
time points, 7, 10 and 14 days post fungal spore inoculation, using five plants per treatment.
For dAdo (or chemically induced cell death) induced cell death, plants were used for
microscopy after 4 days of cell death treatment. To remove external fungal growth or
chemical treatment solutions, plants were washed three times in ddH$_2$O before cell death
staining in a 2.5mM Evans Blue solution (Sigma-Aldrich, Lot #MKCH7958), dissolved in
0.1M CaCl$_2$ pH 5.6 for 15 minutes. After extensive washing for one hour with ddH$_2$O, images
were obtained using a Leica M165 FC microscope.

Activity-based protein profiling

After they were treated with 500 µM dAdo, 500 µM Ado or MES buffer, 9-day-old
Arabidopsis seedlings were frozen in liquid nitrogen, ground and resolved in 50 mM Tris-
HCl buffer (pH 7). After centrifugation, the supernatant was split and either treated with the
proteasome inhibitor MG132 (end concentration: 50 µM, Company) or DMSO for 30
minutes. The samples were then incubated with the 26S proteasome probe MVB072 (end
concentration: 1 µM, (Kolodziejek et al., 2010) for 2 h. Afterwards, the samples were
denatured at 95°C in SDS loading dye and separated on 12 % SDS gels. The probe was
visualized using the rhodamine settings (excitiation: 532 nm, emission: 580 nm) on a
ChemiDoc (BioRad, CA, USA). The protein content of the samples were visualized by
staining the gel with SYPRO^{TM}Ruby (Invitrogen, Carlsbad, CA,USA) according to the
manufacturer's instructions.

RNA seq

7-day-old seedlings were transferred from plates to individual wells of 24-well plates
containing 0.5 mL liquid ½ MS medium supplemented with 0.5% (w/v) sucrose. Following 5
additional days in liquid culture, the medium was replaced with 1 mL 500 µM dAdo in 2.5
mM MES buffer pH 5.6, or 2.5 mM MES buffer pH 5.6 alone as negative control. Three
replicates of four seedlings each were harvested for each treatment at 0, 3, and 12 hours
post treatment (hpt), snap-frozen in liquid nitrogen and stored at -80°C until processing for
RNA extraction. Total RNA was extracted using the TRIZOL reagent as described above.
RNA integrity was confirmed by gel electrophoresis and quantity and purity were determined
using a NanoDrop 2000. Stranded mRNA-seq Libraries were prepared according to the
manufacturer's instructions (Vazyme Biotech Co., Nanjing, China). Qualified libraries were
sequenced on a HiSeq 3000 system instrument at Genomics & Transcriptomics Laboratory,
Heinrich-Heine University, Germany, to generate >100 million reads with a 150 bp read

length from three biological replicates. Trimmomatic (v. 0.36,(Bolger et al., 2014)) was used
for quality trimming and adapter clipping and reads were then mapped to the *Arabidopsis*
thaliana TAIR10 CDS assembly and quantified using kallisto (v. 0.46.2,(Bray et al., 2016)),
which resulted in estimated counts and transcripts per million (TPM) values. The log2 fold
difference of the gene expression between conditions was calculated using R packages
tximport (Soneson et al., 2015) and DESeq2 (Love et al., 2014). Genes with statistical
significance were selected (FDR adjusted p value < 0.05). Data are deposited to the NCBI
under the BioProject accession number xxxx.

Collection of extracellular fluid

640 7-day-old Arabidopsis seedlings, germinated on ½ MS agar with 1 % sucrose were
641 transferred into 24 well plates. Each well contained 3 seedlings in 1.5 ml 2.5 mM MES and
642 the specific treatment. After transfer the plants were returned to the growth chamber for
643 three days. The seedlings were then removed and the liquid was centrifuged at 4000 g for
644 15 minutes. The supernatant was lyophilized and sent for metabolite analysis.

Metabolite analysis

5 µl of apoplastic fluid were injected to an Acquity UPLC (Waters Inc.) equipped with a
Nucleoshell RP18 column (Macherey & Nagel, 150mm x 2 mm x 2.1µm) using
tributylammonium as ion pairing agent. Solvent A: 10 mM tributylamine (aqueous) acidified
with glacial acetic acid to pH 6.2; solvent B acetonitrile. Gradient: 0-2 min: 2% B, 2-18 min
2-36% B, 18-21 min 36-95% B, 21-22.5 min 95% B, 22.51-24 min 2% B. The column flow
was 0.4 ml min^{-1} throughout. The column temperature was 40 °C. Scheduled multiple
reaction monitoring (MRM)-based metabolite detection was performed in negative mode
electrospray ionization (ESI) on a QTrap 6500 (AB-Sciex GmbH, Darmstadt, Germany: ion
source gas 1: 60 psi, ion source gas 2: 70 psi, curtain gas: 35 psi, temperature: 450 °C, Ion
spray voltage floating: and -4500V). MRM transitions were of 189 metabolites covering the
central carbon and energy metabolism were previously signal-optimized and retention times
were determined (Table S10).

Ca^{2+} influx quantification

Calcium influx assays were performed as previously described in (Wanke et al. 2020). In
brief, single 7-day-old Arabidopsis seedlings were placed into white 96-well plates filled with
200 µl reconstitution buffer (2.5 mM MES pH 5.7 [Sigma-Aldrich, Taufkirchen, Germany],
10 mM CaCl$_2$ [Roth, Karlsruhe, Germany]). Before overnight incubation in the dark, the
solution was replaced with 133 µl reconstitution buffer containing 10 mM coelenterazine
(Roth, Karlsruhe, Germany). On the following day, chemiluminescence was measured
using a TECAN SPARK 10M microplate plate reader. After the baseline measurement, 67
µl of three-fold concentrated elicitor solutions (or Milli-Q-water as a mock control) was
manually added. Cytosolic calcium influx upon elicitor addition was constantly measured for
30 min. In order to assess undischarged aequorin for treatment normalization, 100 µl of 3

M CaCl$_2$ (in 30% EtOH) was injected into each well, followed by constant measurement for
1 min. All steps were performed with an integration time of 450 msec.

Heterologous protein production in *Nicotiana benthamiana* and protein purification

For heterologous protein production in *Nicotiana benthamiana, l*eaves of 4-week-old plants
were infiltrated with *Agrobacterium tumefaciens* GV3101 strains. *A. tumefaciens* were
grown for 2 days in LB liquid medium with the appropriate antibiotics at 28°C, 180 rpm until
an OD$_{600}$ of 1 was reached. The cultures were centrifuged at 3500 rpm for 15 min at RT,
supernatant was discarded and resuspended in 1 ml infiltration buffer (10 mM MES pH 5.5,
10 mM MgCl$_2$, 200 µM Acetosyringone) and incubated for 1 h in the dark at 28°C, 180 rpm.
All strains were diluted to OD$_{600}$ of 1 using the infiltration buffer. Each strain was mixed with
P19-expressing strains in a 1:1 ratio. 2-3 leaves were infiltrated per plant using a needleless
syringe. At 4 dpi, the leaves were detached from the plant and ground in liquid nitrogen.
The protein purification was performed according to (Werner et al., 2008) including minor
modifications: a 15 ml tube was filled up to the 2 ml mark with grinded leaf material and 2
ml cold extraction buffer (100 mM Tris pH 8.0, 100 mM NaCl, 5 mM EDTA, 0.5 % Triton
X100, 10 mM DTT, 100 µg/ml Avidin) was added. The powder was re-suspended by
vortexing and subsequently centrifuged at 12000 rpm and 4°C for 10 min. Supernatant was
transferred to a new tube and 100 µl Strep-Tactin® Macroprep (50% slurry) was added and
incubated for 60 minutes at 4°C in a rotation wheel. The mixture was centrifuged for 30 s at
700xg at RT and the supernatant was removed completely from the beads. The beads were
washed once with 4 ml and 2 more times with 2 ml washing buffer (50 mM Tris pH 8.0, 100
mM NaCl, 0.5 mM EDTA, 0.005 % Triton X-100, 2 mM DTT) by centrifugation for 30 s at
700xg, and discarding the supernatant. During the last wash, the beads were transferred
into a 1.5 ml low-binding tube. The beads were either directly boiled for 5 min at 95°C with
6x SDS loading dye to run on a SDS PAGE or the proteins were eluted from the beads by
adding 100 µl elution buffer (wash buffer + 2.5 – 10 mM biotin) and incubation at 25 °C,
>800 rpm for 5 min. The samples were centrifuged for 20 s at 700xg and elution was
repeated. Elution fractions were pooled, SDS loading dye was added and samples were run
on a SDS PAGE followed by a Western Blot.

Ion leakage measurements

700   The measurements of ion leakage were prepared identically like the seedlings for PAM
701   fluorometry measurements. The ion leakage was measured using a conductivity meter
702   (LAQUAtwin EC-11; Horiba, Newhampton,UK).

Oxidative burst assay

Individual 7-day-old Arabidopsis seedlings were transferred into white 96-well plates
containing 200 µl reconstitution buffer (2.5 mM MES pH 5.7 (Sigma-Aldrich, Taufkirchen,
Germany), 10 mM CaCl$_2$ (Roth, Karlsruhe, Germany) and incubated overnight in the growth
chamber. The following day, the buffer was exchanged with 133 µl of reconstitution
supplemented with 15 µg ml^{-1} horseradish peroxidase (Sigma-Aldrich, Taufkirchen,

Germany) and 15 µM L-O12 (Wako Chemicals, Neuss, Germany). Following 10 min
incubation, 67 µL of three-fold concentrated elicitor solutions (or Milli-Q water as a mock
control) was manually added to the wells. Measurements were started immediately and
chemiluminescence was constantly measured with an integration time of 450 msec using a
TECAN SPARK 10M microplate reader (Tecan, Männedorf, Switzerland).

Acknowledgments

We would like to thanks Prof. Jane Parker for discussions and for reading the manuscript
prior submission. ND & AZ also want to thank Lisa Madhi and Lucia Schmitz for their
excellent support in the laboratory. ND was supported by the Max-Planck-Gesellschaft
through the International Max Planck Research School (IMPRS) on 'Understanding
Complex Plant Traits using Computational and Evolutionary Approaches' and the University
of Cologne. AZ acknowledges support from the Cluster of Excellence on Plant Sciences
(CEPLAS) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) under Germany's Excellence Strategy – EXC 2048/1 – Project ID: 390686111
and the SFB project ID: 1403–414786233. CPW acknowledges support from DFG grant
WI3411/8-1. Graphical illustrations were designed with the BioRender online tool.

References
Antonioli, L., Pacher, P., Vizi, E.S., and Hasko, G. (2013). CD39 and CD73 in immunity and
inflammation. Trends in Molecular Medicine *19*, 355-367.
Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics *30*, 2114-2120.
Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. Nature biotechnology *34*, 525-527.
Brinkmann, V., Reichard, U., Goosmann, C., Fauler, B., Uhlemann, Y., Weiss, D.S., Weinrauch,
Y., and Zychlinsky, A. (2004). Neutrophil extracellular traps kill bacteria. Science *303*, 1532-1535.
Chambard, M., Plasson, C., Derambure, C., Coutant, S., Tournier, I., Lefranc, B., Leprince, J.M.,
Kiefer-Meyer, M.C., Driouich, A., Follet-Gueye, M.L.*, et al.* (2021). New Insights into Plant
Extracellular DNA. A Study in Soybean Root Extracellular Trap. Cells *10*.
Choi, J., Tanaka, K., Cao, Y., Qi, Y., Qiu, J., Liang, Y., Lee, S.Y., and Stacey, G. (2014).
Identification of a plant receptor for extracellular ATP. Science *343*, 290-294.
D'Haeze, W., De Rycke, R., Mathis, R., Goormachtig, S., Pagnotta, S., Verplancke, C., Capoen,
W., and Holsters, M. (2003). Reactive oxygen species and ethylene play a positive role in lateral
root base nodulation of a semiaquatic legume. Proceedings of the National Academy of Sciences
100, 11789-11794.
Deshmukh, S., Huckelhoven, R., Schafer, P., Imani, J., Sharma, M., Weiss, M., Waller, F., and
Kogel, K.H. (2006). The root endophytic fungus Piriformospora indica requires host cell death for
proliferation during mutualistic symbiosis with barley. Proc Natl Acad Sci U S A *103*, 18450-18457.
Driouich, A., Smith, C., Ropitaux, M., Chambard, M., Boulogne, I., Bernard, S., Follet-Gueye, M.L.,
Vicre, M., and Moore, J. (2019). Root extracellular traps versus neutrophil extracellular traps in
host defence, a case of functional convergence? Biol Rev Camb Philos Soc *94*, 1685-1700.
Hawes, M., Allen, C., Turgeon, B.G., Curlango-Rivera, G., Tran, T.M., Huskey, D.A., and Xiong,
Z.G. (2016). Root Border Cells and Their Role in Plant Defense. Annu Rev Phytopathol *54*, 143-
161.
Hilbert, M., Voll, L.M., Ding, Y., Hofmann, J., Sharma, M., and Zuccaro, A. (2012). Indole
derivative production by the root endophyte Piriformospora indica is not required for growth
promotion but for biotrophic colonization of barley roots. New Phytologist *196*, 520-534.

Horsefield, S., Burdett, H., Zhang, X., Manik, M.K., Shi, Y., Chen, J., Qi, T., Gilley, J., Lai, J.S.,
Rank, M.X.*, et al.* (2019). NAD(+) cleavage activity by animal and plant TIR domains in cell death
pathways. Science *365*, 793-799.
Kilian, J., Whitehead, D., Horak, J., Wanke, D., Weinl, S., Batistic, O., D'Angelo, C., Bornberg-
Bauer, E., Kudla, J., and Harter, K. (2007). The AtGenExpress global stress expression data set:
protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses.
The Plant Journal *50*, 347-363.
Kloppholz, S., Kuhn, H., and Requena, N. (2011). A Secreted Fungal Effector of Glomus
intraradices Promotes Symbiotic Biotrophy. Curr Biol *21*, 1204-1209.
Kolodziejek, I., Misas-Villamil, J.C., Kaschani, F., Clerc, J., Gu, C., Krahn, D., Niessen, S.,
Verdoes, M., Willems, L.I., Overkleeft, H.S.*, et al.* (2010). Proteasome Activity Imaging and
Profiling Characterizes Bacterial Effector Syringolin A Plant Physiology *155*, 477-489.
Lahrmann, U., Ding, Y., Banhara, A., Rath, M., Hajirezaei, M.R., Doehlemann, S., von Wiren, N.,
Parniske, M., and Zuccaro, A. (2013). Host-related metabolic cues affect colonization strategies of
a root endophyte. Proceedings of the National Academy of Sciences of the United States of
America *110*, 13965-13970.
Lahrmann, U., Strehmel, N., Langen, G., Frerigmann, H., Leson, L., Ding, Y., Scheel, D., Herklotz,
S., Hilbert, M., and Zuccaro, A. (2015). Mutualistic root endophytism is not associated with the
reduction of saprotrophic traits and requires a noncompromised plant innate immunity. New
Phytologist *207*, 841-857.
Lahrmann, U., and Zuccaro, A. (2012). Opprimo ergo sum-Evasion and Suppression in the Root
Endophytic Fungus Piriformospora indica. Molecular Plant-Microbe Interactions *25*, 727-737.
Li, G., Liu, K., Baldwin, S.A., and Wang, D. (2003). Equilibrative nucleoside transporters of
Arabidopsis thaliana. cDNA cloning, expression pattern, and analysis of transport activities. The
Journal of biological chemistry *278*, 35732-35742.
Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A., Reissmann,
S., Kahmann, R., and Merchant, S. (2015). Fungal Effectors and Plant Susceptibility. In Annual
Review of Plant Biology, Vol 66, pp. 513-545.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome biology *15*, 550.
Marquès-Bueno, M.M., Morao, A.K., Cayrel, A., Platre, M.P., Barberon, M., Caillieux, E., Colot, V.,
Jaillais, Y., Roudier, F., and Vert, G. (2016). A versatile Multisite Gateway-compatible promoter
and transgenic line collection for cell type-specific functional genomics in Arabidopsis. The Plant
Journal *85*, 320-333.
Mucha, J., Guzicka, M., Ratajczak, E., and Zadworny, M. (2014). Strategies utilized by trophically
diverse fungal species for Pinus sylvestris root colonization. Tree Physiol *34*, 73-86.
Myrach, T., Zhu, A., and Witte, C.P. (2017). The assembly of the plant urease activation complex
and the essential role of the urease accessory protein G (UreG) in delivery of nickel to urease. The
Journal of biological chemistry *292*, 14556-14565.
Ngou, B.P.M., Ahn, H.K., Ding, P., and Jones, J.D.G. (2021). Mutual potentiation of plant immunity
by cell-surface and intracellular receptors. Nature *592*, 110-115.
Nizam, S., Qiang, X., Wawra, S., Nostadt, R., Getzke, F., Schwanke, F., Dreyer, I., Langen, G.,
and Zuccaro, A. (2019). Serendipita indica E5'NT modulates extracellular nucleotide levels in the
plant apoplast and affects fungal colonization. EMBO Rep *20*.
Nostadt, R., Hilbert, M., Nizam, S., Rovenich, H., Wawra, S., Martin, J., Kupper, H., Mijovilovich,
A., Ursinus, A., Langen, G.*, et al.* (2020). A secreted fungal histidine- and alanine-rich protein
regulates metal ion homeostasis and oxidative stress. New Phytol.
Oberwinkler, F., Riess, K., Bauer, R., Selosse, M.-A., Weiß, M., Garnica, S., and Zuccaro, A.
(2013). Enigmatic Sebacinales. Mycological Progress *12*, 1-27.
Olvera-Carrillo, Y., Van Bel, M., Van Hautegem, T., Fendrych, M., Huysmans, M., Simaskova, M.,
van Durme, M., Buscaill, P., Rivas, S., Coll, N.S.*, et al.* (2015). A Conserved Core of Programmed
Cell Death Indicator Genes Discriminates Developmentally and Environmentally Induced
Programmed Cell Death in Plants. Plant Physiol *169*, 2684-2699.

Pham, A.Q., Cho, S.-H., Nguyen, C.T., and Stacey, G. (2020). Arabidopsis Lectin Receptor Kinase
P2K2 Is a Second Plant Receptor for Extracellular ATP and Contributes to Innate
Immunity1 [OPEN]. Plant Physiology *183*, 1364-1375.
Plett, J.M., Daguerre, Y., Wittulsky, S., Vayssieres, A., Deveau, A., Melton, S.J., Kohler, A.,
Morrell-Falvey, J.L., Brun, A., Veneault-Fourrey, C.*, et al.* (2014). Effector MiSSP7 of the
mutualistic fungus Laccaria bicolor stabilizes the Populus JAZ6 protein and represses jasmonic
acid (JA) responsive genes. Proc Natl Acad Sci U S A *111*, 8299-8304.
Qiang, X., Zechmann, B., Reitz, M.U., Kogel, K.H., and Schafer, P. (2012). The mutualistic fungus
Piriformospora indica colonizes Arabidopsis roots by inducing an endoplasmic reticulum stress-
triggered caspase-dependent cell death. Plant Cell *24*, 794-809.
Rafiqi, M., Jelonek, L., Akum, N.F., Zhang, F., and Kogel, K.H. (2013). Effector candidates in the
secretome of Piriformospora indica, a ubiquitous plant-associated fungus. Front Plant Sci *4*, 228.
Ragnelli, A.M., Aimola, P., Maione, M., Zarivi, O., Leonardi, M., and Pacioni, G. (2014). The cell
death phenomenon during Tuber ectomycorrhiza morphogenesis. Plant Biosystems - An
International Journal Dealing with all Aspects of Plant Biology *148*, 473-482.
Rich-Griffin, C., Stechemesser, A., Finch, J., Lucas, E., Ott, S., and Schäfer, P. (2020). Single-Cell
Transcriptomics: A High-Resolution Avenue for Plant Functional Genomics. Trends in plant
science *25*, 186-197.
Saur, I.M.L., Panstruga, R., and Schulze-Lefert, P. (2021). NOD-like receptor-mediated plant
immunity: from structure to cell death. Nat Rev Immunol *21*, 305-318.
Schneider, H.M., and Lynch, J.P. (2018). Functional implications of root cortical senescence for
soil resource capture. Plant and Soil *423*, 13-26.
Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. F1000Research *4*, 1521.
Straube, H., Niehaus, M., Zwittian, S., Witte, C.P., and Herde, M. (2021). Enhanced nucleotide
analysis enables the quantification of deoxynucleotides in plants and algae revealing connections
between nucleoside and deoxynucleoside metabolism. Plant Cell *33*, 270-289.
Tedersoo, L., Bahram, M., Ryberg, M., Otsing, E., Koljalg, U., and Abarenkov, K. (2014). Global
biogeography of the ectomycorrhizal /sebacina lineage (Fungi, Sebacinales) as revealed from
comparative phylogenetics analyses. Mol Ecol.
Thammavongsa, V., Kern, J.W., Missiakas, D.M., and Schneewind, O. (2009). Staphylococcus
aureus synthesizes adenosine to escape host immune responses. J Exp Med *206*, 2417-2427.
Thammavongsa, V., Missiakas, D.M., and Schneewind, O. (2013). Staphylococcus aureus
degrades neutrophil extracellular traps to promote immune cell death. Science *342*, 863-866.
Thürich, J., Meichsner, D., Furch, A.C.U., Pfalz, J., Krüger, T., Kniemeyer, O., Brakhage, A., and
Oelmüller, R. (2018). Arabidopsis thaliana responds to colonisation of Piriformospora indica by
secretion of symbiosis-specific proteins. PLOS ONE *13*, e0209658.
Tran, T.M., MacIntyre, A., Hawes, M., and Allen, C. (2016). Escaping Underground Nets:
Extracellular DNases Degrade Plant Extracellular Traps and Contribute to Virulence of the Plant
Pathogenic Bacterium Ralstonia solanacearum. PLoS Pathog *12*, e1005686.
Vijayaraghavareddy, P., Adhinarayanreddy, V., Vemanna, R.S., Sreeman, S., and Makarla, U.
(2017). Quantification of Membrane Damage/Cell Death Using Evan's Blue Staining Technique.
Bio-protocol *7*, e2519.
Voß, S., Betz, R., Heidt, S., Corradi, N., and Requena, N. (2018). RiCRN1, a Crinkler Effector
From the Arbuscular Mycorrhizal Fungus Rhizophagus irregularis, Functions in Arbuscule
Development. Frontiers in Microbiology *9*.
Wan, L., Essuman, K., Anderson, R.G., Sasaki, Y., Monteiro, F., Chung, E.H., Osborne Nishimura,
E., DiAntonio, A., Milbrandt, J., Dangl, J.L.*, et al.* (2019). TIR domains of plant immune receptors
are NAD(+)-cleaving enzymes that promote cell death. Science *365*, 799-803.
Wawra, S., Fesel, P., Widmer, H., Neumann, U., Lahrmann, U., Becker, S., Hehemann, J.H.,
Langen, G., and Zuccaro, A. (2019). FGB1 and WSC3 are in planta-induced beta-glucan-binding
fungal lectins with different functions. New Phytol.
Wawra, S., Fesel, P., Widmer, H., Timm, M., Seibel, J., Leson, L., Kesseler, L., Nostadt, R.,
Hilbert, M., Langen, G.*, et al.* (2016). The fungal-specific beta-glucan-binding lectin FGB1 alters
cell-wall composition and suppresses glucan-triggered immunity in plants. Nat Commun *7*, 13188.

Weiss, M., Waller, F., Zuccaro, A., and Selosse, M.A. (2016). Sebacinales - one thousand and one
interactions with land plants. New Phytologist *211*, 20-40.
Wen, F.S., White, G.J., VanEtten, H.D., Xiong, Z.G., and Hawes, M.C. (2009). Extracellular DNA
Is Required for Root Tip Resistance to Fungal Infection. Plant Physiology *151*, 820-829.
Werner, A.K., Sparkes, I.A., Romeis, T., and Witte, C.P. (2008). Identification, biochemical
characterization, and subcellular localization of allantoate amidohydrolases from Arabidopsis and
soybean. Plant Physiol *146*, 418-430.
Winstel, V., Missiakas, D., and Schneewind, O. (2018). Staphylococcus aureus targets the purine
salvage pathway to kill phagocytes. Proc Natl Acad Sci U S A *115*, 6846-6851.
Wu, Z., Li, M., Dong, O.X., Xia, S., Liang, W., Bao, Y., Wasteneys, G., and Li, X. (2019).
Differential regulation of TNL-mediated immune signaling by redundant helper CNLs. New Phytol
222, 938-953.
Xiao, Y., Savchenko, T., Baidoo, E.E., Chehab, W.E., Hayden, D.M., Tolstikov, V., Corwin, J.A.,
Kliebenstein, D.J., Keasling, J.D., and Dehesh, K. (2012). Retrograde signaling by the plastidial
metabolite MEcPP regulates expression of nuclear stress-response genes. Cell *149*, 1525-1535.
Yu, D., Song, W., Tan, E.Y.J., Liu, L., Cao, Y., Jirschitzka, J., Li, E., Logemann, E., Xu, C., Huang,
S.*, et al.* (2021). TIR domains of plant immune receptors are 2′,3′-cAMP/cGMP synthetases
mediating cell death. bioRxiv, 2021.2011.2009.467869.
Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., Wang, Y., Cai, B., Zhou, J.M., He, S.Y., and Xin,
X.F. (2021). Pattern-recognition receptors are required for NLR-mediated plant immunity. Nature
592, 105-109.
Zipfel, C., and Oldroyd, G.E.D. (2017). Plant signalling in symbiosis and immunity. Nature *543*,
328-336.
Zuccaro, A., Lahrmann, U., Gueldener, U., Langen, G., Pfiffi, S., Biedenkopf, D., Wong, P.,
Samans, B., Grimm, C., Basiewicz, M.*, et al.* (2011). Endophytic Life Strategies Decoded by
Genome and Transcriptome Analyses of the Mutualistic Root Symbiont Piriformospora indica. Plos
Pathogens *7*.

Figure 1. Host cell death during *S. indica* root colonization

(A and B) Arabidopsis roots expressing the fluorescent nuclei marker
UBQ10::H2B:mCherry (magenta) and stained with the fungal cell wall marker Alexa Fluor
dye WGA-AF 488 (green) at 10 dpi.
(C) Plant nuclei stained with DAPI (magenta) and fungal cell wall and matrix stained with
the β-glucan binding lectin FGB1-FITC 488 (green) at 6dpi. With progression of *S. indica*
colonization host nuclei often get stretched and fade.
(D and E) Close up of *S. indica* hyphae embedded in a host nucleus from (C).
(F and G) Staining of nucleic acids of *S. indica*-colonized roots with the dead-cell indicator
(membrane integrity marker) SYTOX Orange (magenta) and fungal hyphae using WGA-
AF 488 (green) at 10 dpi.
(H and I) *S. indica* hyphae fluorescently labelled with the β-glucan binding lectin FGB1-
FITC 488 (green), embedded in a DAPI-stained host nucleus (magenta) at 6dpi.
(J-M) Progression of vacuolar collapse of colonized root cells. Fungal hyphae are stained
using WGA-AF 488 (green) while membranes are stained with FM4-64 (red).
(J and K) Initial biotrophic colonization.
(L and M) vacuolar collapse in a dying host cell.
CLSM was repeated at least five times using 3 to 4 *S. indica* colonized plants. Fading of
nuclei at the onset of cell death during fungal colonization was observed regularly.

Figure 2. *Si*NucA is a small secreted nuclease involved in fungal accommodation

(A) *Si*NucA expression in *S. indica*-colonized Arabidopsis and barley roots and axenic
cultures, measured using the $2^{-\Delta CT}$ method. Error bars: SD, n = 3 (biological replicates).
(B) *Si*NucA:HA:His protein enrichment from culture filtrate precipitated with 80% ammonium
sulfate and separated by size exclusion chromatography. The resulting fraction was
separated by SDS-PAGE and *Si*NucA:HA:His was stained with Coomassie Blue.
(C) The purified *Si*NucA:HA:His protein was incubated with fungal (*S. indica*) or plant
(Arabidopsis) RNA or DNA in 5 mM Tris buffer (pH = 8) with 1 mM $MgCl_2$, and 1 mM $CaCl_2$
for 4 (RNA) or 10 (DNA) minutes and visualized after gel electrophoresis.
(D) Arabidopsis lines expressing *Si*NucA driven by the 35S promoter (lines 18, 53 and 56)
compared to control lines (4 and 18: segregating from T2 generation and Col-0 WT). Roots
of plants grown on ½ MS medium were inoculated with *S. indica* and analyzed at 7 dpi. Dots
represent independent biological replicates, while lines represent the mean. Different letters
indicate significant different groups determined with a one-way ANOVA with post-hoc Tukey
HSD test (p<0.05).

(E) Root colonization by *S. indica* in transgenic 35S::*Si*NucA Arabidopsis lines at 7 dpi was
assessed by RT-qPCR comparing the expression of the fungal housekeeping gene *Si*TEF
and the plant gene *At*Ubi and the $2^{-\Delta\Delta CT}$ method, normalizing to the Col-0 colonization. Dots
represent independent biological replicates, while lines represent the mean. Different letters
indicate significant different groups determined with a one-way ANOVA with post-hoc Tukey
HSD test ($p<0.05$).
(F) CLSM live cell images of an Arabidopsis root expressing *Si*NucA:mCherry (yellow)
inoculated with *S. indica*. Fungal cell walls are stained with WGA-AF 488 (green) and nuclei
with DAPI (magenta). *Si*NucA:mCherry accumulates in the apoplast in non-colonized root
cells (asterisk) and (re-)localizes to the host nuclei in colonized cells (arrow).

Figure 3. *Si*NucA and *Si*E5NT act synergistically to produce deoxy-nucleosides
from DNA

(A) Enzymatic activity and kinetic constants of *Si*E5NT. Left axis, initial catalytic velocity (v)
for phosphate production at different dAMP concentrations fitted using the Michaelis-
Menten equation. Right axis, ratio of dAMP concentrations and velocities (s/v) plotted
against dAMP concentrations (s) fitted by linear regression (Hanes plot). Error bars: SD, n
= 3.
(B) Same as in (A) but using AMP as substrate (n = 3).
(C) Degradation of DNA by *Si*NucA and/or *Si*E5NT. 2% agarose gel loaded with the
products of an incubation of 10 µg salmon sperm DNA for one hour at 25° C with different

enzyme combinations.
(D) Relative quantification by HPLC-MS/MS of dAdo (deoxyadenosine), dC (deoxycytidine),
dG (deoxyguanosine) and dT (deoxythymidine) released during the incubation shown in
(C). Deoxynucleotides could not be detected in any of the reactions.

A P_{UBQ10}::H2B:mcherry

Ado

dAdo

Ado
100 μm

dAdo
100 μm

B Evans Blue staining at 4 dpt

Mock   dAdo

C Quantification

Evans Blue staining [intensity]

Col-0 + Mock   Col-0 + dAdo

***

D dAdo induces ion leakage in Arabidopsis seedlings

conductivity [μS/cm]

dpt

- Mock
- Ado
- dAdo

E 3 dpt

conductivity [μS/cm]

Mock   Ado   dAdo

a   a   b

- Mock
- Ado
- dAdo

F dAdo activates the 26S proteasome

Duration   3 hpt   12 hpt   24 hpt
Treatment M   dAdo   M   dAdo   M   dAdo
Inhibitor - + - + - + - + - + - + NPC

25 kDa   26S Proteasome

SyproRuby

G Effect of dAdo on photosynthesis

[F_V/F_M]

dpt

- Mock
- Ado
- dAdo

H 7 dpt

[F_V/F_M]

Mock   Ado   dAdo

a   a   b

- Mock
- Ado
- dAdo

I

relative
row min → row max

Associated PCD | ens_gene
genotoxic | AT2G45460
osmotic | AT1G69260
genotoxic | AT3G45730
osmotic | AT1G07430
osmotic | AT1G16850
dev. | AT5G13820
dev. | AT1G26820
dev. | AT3G21550
genotoxic | AT2G18600
biotic | AT1G13340
biotic | AT1G51890
biotic | AT2G18690
dev. | AT5G18310
biotic | AT3G26830
osmotic | AT4G24960
osmotic | AT2G47770
genotoxic | AT1G07500
genotoxic | AT3G07800
genotoxic | AT4G22960
genotoxic | AT4G21070
genotoxic | AT5G20850
genotoxic | AT4G29170
osmotic | AT1G52690
osmotic | AT3G02480
dev. | AT4G04460
dev. | AT1G45145
osmotic | AT1G60190
osmotic | AT5G06760
osmotic | AT5G59220
genotoxic | AT3G42860

J MEP pathway 12 hpt   extracellular metabolites 3 dpt

Plastid

Pyr + G3P

3.536 *DXPS1* → CO2

DXP

DXR NADPH/H⁺ / NADP⁺

MEP

0.825 *CMS* CTP / PP_i

CDP-ME

0.556 *CMK* ATP / ADP

CDP-MEP

0.652 *MCS* CMP

MEcPP

0.721 *HDS* 2 Fd_{red} / 2 Fd_{ox} / H2O

HMB-PP

0.264 *HDR* NAD(P)H/H⁺ / NAD(P)⁺ / H2O

DMAPP   IPP

DXP — Mock / dAdo

MEP — Mock / dAdo *

CDP-ME — Mock / dAdo

CDP-MEP — Mock / dAdo

MEcPP — Mock / dAdo *

DMAPP — Mock / dAdo ***

Figure 4. dAdo induces cell death *in planta*

(A) CLSM images of seven-day-old Arabidopsis roots expressing the nuclei marker UBQ10::H2B:mCherry. Incubation with dAdo but not Ado (500 µM) leads to disorganization of Arabidopsis cells in the root tip and to disappearance or fading of the nuclear material.

(B) Bright field microscopy of the root tip and differentiation zone of Arabidopsis seedlings treated with Mock/dAdo (500 µM) and stained with the cell death dye Evans Blue.

(C) Quantification of root cell death in Col-0 root tips at 4 days after dAdo-treatment. Cell death was evaluated using Evans Blue staining. Boxplots represent the data obtained from 5 biological replicates. Asterisks represent significant difference to the Mock-treated sample analyzed by Student's t-test ($p < 0.005$ ***). The experiment was repeated independently 3 times with similar results.

(D) Electrolyte leakage of Col-0 seedlings (9-day-old) in MES buffer after Mock-, Ado- or dAdo-treatment. Error bars depict the standard error of the mean (SEM) obtained from 6 biological replicates. The experiment was repeated independently at least 3 times with similar results. (E) Electrolyte leakage at 3 dpt of the seedlings from (D). Error bars depict the standard error of the mean (SEM) obtained from 6 biological replicates. Different letters indicate significant different groups determined with a one-way ANOVA with post-hoc Tukey HSD test ($p<0.05$). The experiment was repeated independently at least 3 times with similar results.

(F) Activity of the Arabidopsis 26S proteasome. Total protein extracts of 14-day-old seedlings treated with 500 µM dAdo or 2.5 mM MES buffer (M) were incubated with 1 µM of the probe MVB072 (Kolodziejek et al. 2011). Before labeling, samples were incubated with 50 µM of the proteasome inhibitor Epoximycin (+) or DMSO (-). Afterwards, the samples were labeled for 2 hours and separated by SDS-PAGE. The 26S proteasome activity was visualized by fluorescence scanning. SYPRO™Ruby staining was performed to compare sample loads. A non-probe control (NPC) consisting of a mix of all samples incubated with DMSO was used as an additional control. The experiment was repeated twice with similar results.

(G) Photosynthetic activity (F_V/F_M) of 7-day-old Col-0 seedlings incubated with 500 µM Ado or dAdo. Error bars depict the standard error of the mean (SEM) obtained from twelve biological replicates. The experiment was repeated at least 3 times with similar results.

(H) Photosynthetic activity (F_V/F_M) of 7-day-old Col-0 seedlings incubated with 500 µM Ado or dAdo at 7 dpt from (G). Error bars depict the standard error of the mean (SEM) obtained from twelve technical replicates. Different letters indicate significant different groups determined with a one-way ANOVA with post-hoc Tukey HSD test ($p<0.05$).

(I) Heatmap of core regulated cell death (RCD) marker genes expression (selected from Olvera-Carrillo et al., 2015) of 7-day-old Arabidopsis seedlings after dAdo-treatment at 0, 3 and 12 hours. RNA-Seq data are shown as $\log_2(tpm)$ (Table S6). Heatmaps and hierarchical clustering (one-minus Pearson correlation) were generated using Morpheus, https://software.broadinstitute.org/morpheus.

(J) Graphical representation of the methyl erythritol 4-phosphate (MEP) pathway. Colored numbers left to key enyzmes represent the $\log_2$ fold-change after 12 hours of dAdo-treatment (Padj < 0.05), measured by RNA-Seq. Boxplots on the right side represent the

corresponding extracellular metabolite levels at 3 dpt. Asterisks represent significant differences to the Mock-treated sample analyzed by Student's t-test (p < 0.05 * ; p < 0.001 ***).

Figure 5. dAdo-triggered signalling and cell death are not mediated by canonical pattern-triggered immune responses

(A) Ca²⁺-influx after treatment of 7-day-old Arabidopsis Col-0^{AEQ} seedlings with 500 μM purine derivatives solved in 2.5 mM MES buffer (pH 5.7). Buffer and 33 μM flg22 were used as positive and negative controls, respectively. Ca²⁺-influx was monitored via a coelenterazine-based chemiluminescence assay and measured as relative light units (RLU). After discharging the remaining aequorin by adding CaCl₂, discharge kinetics were integrated and normalized to the maximum Ca²⁺ level. The discharge integral was then used to normalize Ca²⁺ kinetics in response to elicitor treatment. Curves represent eight biological replicates and the experiments were additionally repeated independently three times with similar results.

(B) Boxplots represent total Ca^{2+}-influx over the measured time period. Calculations were
performed as described in (A). Values represent means ± SEM from eight biological
replicates, each consisting of one seedling. Different letters indicate significant differences
determined by a Kruskal-Wallis test and post-hoc Dunn's test ($p < 0.05$). The experiment
was repeated three times with similar results.
(C) Comparison of total Ca^{2+}-influx in 7-day-old Arabidopsis Col-0^{AEQ} and Arabidopsis Col-
0^{AEQ} *dorn1* seedlings after treatment with 500 µM ATP, dATP, 33 µM flg22 or 2.5 mM MES
pH 5.7 (Mock). Data represent values from eight biological replicates. Calculations were
performed as described in (A). Asterisks represent significant differences between the two
genotypes analyzed by Student's t-test ($p < 0.005$ ***). The experiment was repeated three
times with similar results.
(D) Expression of the marker gene At1g58420, responsive to wounding and *S. indica*
colonization, in treated 7-day-old Col-0 seedlings. Fold change expression was calculated
relative to housekeeping gene *At*Ubi and Mock treatment using the $2^{-\Delta\Delta Ct}$ method. Data
represents 6 independent biological replicates. Different letters indicate significant different
groups determined by Kruskal-Wallis test and post-hoc Wilcox BH adjustment ($p < 0.05$).
(E) Metabolic analysis of the supernatant of purine-treated 7-day-old Col-0 seedlings at 3
dpt. Dots represent data of at least 6 biologically independent replicates normalized to the
average of the corresponding Mock samples. Different letters indicate significant different
groups determined by a Kruskal-Wallis test and post-hoc Wilcox BH adjustment ($p < 0.05$).

Figure 6. The Arabidopsis equilibrative nucleoside transporter ENT3 is required for dAdo-mediated signalling and cell death

(A) Photosynthetic activity (F_V/F_M) of 9-day-old Mock- and dAdo-treated (500 µM) Col-0, *ent3* and *ent1* seedlings. Measurements were performed 3 days post treatment (dpt), every 24 hours. Data show the mean and error bars depict the standard error of the mean (SEM)

obtained from 12 technical replicates consisting of 3 seedlings each. The experiment was
repeated more than three times independently, yielding similar results.
(B) F_V/F_M of the seedlings from (A) at 3 dpt. Boxplots represent the data of 12 technical
replicates consisting of 3 seedlings each. Different letters indicate significant differences
determined with a one-way ANOVA with post-hoc Tukey HSD test ($p < 0.05$).
(C) Visualization of F_V/F_M measured via PAM fluorometry. The FV/FM value is visualized by
the color scale shown below. Depicted are 12 wells containing 9-day-old seedlings,
screened at 0, 3 and 7 dpt with either mock treatment (2.5 mM MES buffer pH 5.6), 500 µM
dAdo or 500 µM MeJA. The experiment was repeated more than three times independently,
yielding similar results.
(D) Photosynthetic activity (F_V/F_M x photosynthetically active area) of 9-day-old Col-0
seedlings incubated with different concentrations of Ado and dAdo at 7 dpt. Dots represent
3 biologically independent replicates consisting of 12 wells containing 3 seedlings each.
Different letters indicate significant differences determined with a one-way ANOVA with
post-hoc Tukey HSD test ($p < 0.05$).
(E) Metabolic analysis of 7-day-old Col-0 supernatants from dAdo-treated and *ent3*
seedlings at 3 dpt. Boxplots represent the data of 6 biologically independent replicates
normalized to the Mock-treated samples. Asterisks represent significant differences
between the two genotypes analyzed by Student's t-test ($p < 0.05$ *,$p < 0.005$ ***).
(F) Metabolic analysis of the supernatant from *S. indica*-inoculated 7-day-old Col-0 and *ent3*
seedlings at different time points after treatment (3′,5′-cAMP: 3 dpi; MEcPP: 6 dpi;
phosphate: 10 dpi). Dots represent the data of 6 biologically independent replicates
normalized to the Mock-inoculated samples. Asterisks represent significant differences
between the two genotypes analyzed by Student's t-test ($p < 0.05$ (*), $p < 0.005$ ***).
(G) Quantification of root cell death in 7-day-old *S. indica*-colonized Col-0 and ent3 roots at
7, 10 and 14 dpi. Cell death was evaluated via Evans Blue staining and normalized to the
Mock-treated roots. Boxplots represent the data obtained from 20 biological replicates.
Asterisks represent significant difference to the Mock-treated sample analyzed by Student's
t-test ($p < 0.05$ *, $p < 0.01$ **, $p < 0.005$ ***). The experiment was repeated three times
independently, yielding similar results.
(H) Evans blue staining of *S. indica*-colonized Col-0 and *ent3* roots at 10 dpi.
(I) *S. indica* abundance in 7-day-old Arabidopsis seedlings. The ratio of fungus (*Si*TEF) to
plant (*At*Ubi) was calculated using cDNA as template and the $2^{-\Delta CT}$ method. Boxplots
represent 6 independent biological replicates. Asterisks indicate significant difference from
Col-0 samples (Student's t-test, $p < 0.05$ *).

Figure 7. An uncharacterized *At*NLR is involved in dAdo-mediated cell death

(A) F_V/F_M of 9-day-old Mock- and dAdo-treated (500 µM) Col-0 and *nlr* knock-out mutant seedlings. Measurements were performed for 3 days. Data show the mean and error bars depict the standard error of the mean (SEM) obtained from 12 technical replicates consisting of 3 seedlings each. Asterisks indicate significant difference from Col-0 samples (Student's t-test, $p < 0.005$ ***). The experiment was repeated three times independently with similar results.

(B) Boxplots of F_V/F_M measurements from (A) at 3 dpt. Different letters indicate significant differences determined with a one-way ANOVA and post-hoc Tukey HSD test ($p < 0.05$).

(C) Visualization of F_V/F_M measured via PAM fluorometry. Blue indicates a high photosynthetic activity while green decreased and yellow strongly decreased photosynthetic activity. Depicted are 12 wells containing 9-day-old seedlings, screened at 0 and 3 days post treatment (dpt) either with Mock or 500 µM dAdo. The experiment was repeated three times independently with similar results.

(D) Conductivity of MES buffer containing 9-day-old Col-0 or *nlr* seedlings 0 – 3 dpt after Mock- or dAdo-treatment (500 µM). Data points depict the mean, while error bars depict the standard error of the mean (SEM) obtained from 12 biological replicates.

(E) Boxplots of conductivity measurements from (D) at 3 dpt. Different letters indicate significant differences determined with a one-way ANOVA with post-hoc Tukey HSD test ($p < 0.05$).

Figure 8. Model for dAdo-triggered cell death during *S. indica* colonization of *A. thaliana* roots

The beneficial endophyte *S. indica* secretes two enzymes into the apoplast, the nuclease NucA and the ecto-nucleotidase E5NT. *Si*E5NT is initially involved in manipulation of the eATP signalling (left side) as described in Nizam et al., 2019. *Si*NucA accumulates at the onset of cell death. The combined activity of *Si*NucA and *Si*E5NT releases deoxynucleosides from DNA with a strong preference for dAdo, a potent cell death inducer in animal systems. dAdo is transported into the cytoplasm *via At*ENT3 where it initiate a cell death process, contributing to successful fungal colonization. dAdo induces production of the retrograde stress signal MEcPP from plastids, activating stress signalling and intercellular communication. dAdo-triggered cell death is modulated by an uncharacterized TR type NLR protein, establishing a link to immunometabolism.