Prokaryotic phospholipid translocation by ubiquitous PplT domain proteins

Janna N. Hauser[1,2,3], Arnaud Kengmo Tchoupa[1,2,3], Susanne Zabel[4], Kay Nieselt[3,4], Christoph M. Ernst[1,5], Christoph J. Slavetinsky[1,2,3,6#], Andreas Peschel[1,2,3#]

1. Interfaculty Institute of Microbiology and Infection Medicine Tübingen, Infection Biology Section, University of Tübingen, Germany.

2. German Center for Infection Research (DZIF), partner site Tübingen, Germany.

3. Cluster of Excellence EXC2124 Controlling Microbes to Fight Infection, University of Tübingen, Germany

4. Institute for Bioinformatics and Medical Informatics, University of Tübingen, Germany

5. Current address: Department of Molecular Biology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

6. Pediatric Surgery and Urology, University Children's Hospital Tübingen, University of Tübingen, Germany.

# Equally contributing corresponding authors. Correspondence to: Christoph J. Slavetinsky and Andreas Peschel, Interfaculty Institute of Microbiology and Infection Medicine, Infection Biology Section, University of Tübingen, Auf der Morgenstelle 28, 72076 Tübingen, Germany.

Abstract

Flippase proteins exchanging phospholipids between the cytoplasmic membrane leaflets have been

identified in Eukaryotes but remained largely unknown in Prokaryotes. Only MprF proteins that

synthesize aminoacyl phospholipids in some bacteria have been found to contain a domain that

translocates the produced lipids. We show here that this domain, which we named 'prokaryotic

phospholipid translocator' (PplT), is widespread in Bacteria and Archaea, encoded as a separate

protein or fused to different types of enzymes. We also demonstrate that the *Escherichia coli* PplT

protein interacts with many phospholipid-synthetic enzymes and deletion of *pplT* impaired bacterial

growth and membrane fluidity. Thus, PplT domain proteins appear to be general prokaryotic lipid

flippases with critical roles in cellular homeostasis.

Introduction

Most of the basic processes and enzyme systems involved in growth and maintenance of prokaryotic

cells have been explored in the past. Major mechanisms of membrane lipid synthesis and turnover

have also been described thoroughly [1, 2]. However, 'flippase' proteins that facilitate the

translocation of phospholipids from the inner leaflet, the site of synthesis, to the outer leaflet of the

cytoplasmic membrane, have remained largely unknown. We speculated that such flippases might be

among the conserved membrane proteins of unknown function found in many prokaryotic taxa.

Phospholipid flippases may not be essential for cell viability, which might explain why they have not

been identified in previous screening approaches for prokaryotic proteins with crucial function for

cellular integrity. Flippases have been characterized in eukaryotic cells [3] but the responsible proteins

do not seem to have obvious homologs in prokaryotes.

Flippase reactions are particularly difficult to study because the flipping process is fast and hard to

monitor as the lipids remain in place and adapt only their orientation. Nevertheless, a prototype

flippase has been identified in prokaryotic MprF proteins that synthesize and then translocate

aminoacyl phosphatidylglycerol (Aa-PG) lipids such as lysyl-phosphatidylglycerol (Lys-PG) or alanyl-

phosphatidylglycerol (Ala-PG) [4, 5]. Those lipids reduce the negative net charge at the outer surface

of cytoplasmic membranes, thereby diminishing the binding of cationic antimicrobial molecules such

as bacterial bacteriocin and host defensin peptides [6]. To achieve protection against such molecules,

it is critical for microorganisms to efficiently translocate Lys-PG or Ala-PG, once these lipids have been

formed by the MprF synthase domain at the inner surface of the cytoplasmic membrane. The synthase

domain cooperates with the flippase domain in a dynamic fashion to deliver the lipid products to the

translocation channel [7]. The two domains function together even if expressed as separate proteins

[8]. The flippase domain of MprF consists of eight transmembrane sections [8] and contains a motif

that is accessible from both sides of the cytoplasmic membrane [9]. It has been proposed that this part

moves during the lipid translocation process or that it is in the center between two larger cavities that

form the lipid translocation channel [9]. The structure of the *Rhizobium tropici* MprF protein with

bound Lys-PG molecules captured in the process of translocation has recently been elucidated by cryo-

electron microscopy [7]. It gives rise to a mechanistic model that proposes a process with Lys-PG

entering an inner protein cavity, which transiently opens a gate to an outer cavity, from which Lys-PG

is subsequent released to the outer membrane leaflet [7]. The extent to which the flippase domain

found in MprF proteins is present in Bacteria and Archaea, and if it may play a more general role in

phospholipid translocation, has remained unknown.

The flippase domain of MprF proteins exhibits a conserved architecture with low sequence identity. It

is referred to as cl04219, pfam03706, or UPF0104 in the NCBI Conserved Domain Database (CDD) [10]

and as IPR022791 in the InterPro protein signatures database [11]. We show here that proteins

consisting of this domain alone or in combination with various enzymatic domains are found in almost

all of the major groups of Bacteria and Archaea, most likely to accomplish the translocation of

membrane lipids. The corresponding *Escherichia coli* protein interacts with most of the phospholipid-

biosynthetic enzymes and it is encoded in the vicinity of a cardiolipin (CL) synthase. Its inactivation is

not lethal but affects membrane fluidity and the fitness of *E. coli*. We suggest naming the

corresponding flippase domain 'prokaryotic phospholipid translocator' (PplT), which is likely to fulfill a

general membrane homeostatic function in prokaryotes.

Results

1. PplT domain proteins are widespread in most groups of prokaryotes. MprF-related proteins are

found in many different bacterial taxa [12]. Notably, we found that the two functional domains of MprF

occur in different combinations. The Lys-PG synthase domain, for instance, occurs sometimes without

the Lys-PG flippase domain [12]. On the other hand, the flippase domain is often encoded without a

Lys-PG synthase or in combination with other enzymatic domains. Systematic evaluation of microbial

genomes for the presence of this domain, referred to as PplT (cl04219), revealed an almost universal

occurrence in most phyla of Bacteria (Fig. 1A) and in several Classes of Archaea (Fig. 1B). Indeed, blastp

searches uncovered that the PplT domain is even found in bacterial species, which do not produce Aa-

PGs, such as *E. coli* and many other Enterobacteriaceae [13, 14]. Thus, PplT may have a broader role in

prokaryotes than previously thought. It should be noted that most prokaryotes with a *pplT* homolog

encode only one protein with a PplT domain in their genome (Fig. S1).

PplT occurs as a separate protein in many different bacterial phyla including Proteobacteria,

Actinobacteria, Firmicutes, Bacteroidetes, Planctomycetes, and Thermotogae as well as archaeal phyla

including Euryarchaeota and Crenarchaeota. Moreover, PplT is found connected to other types of

proteins in addition to Lys-PG or Ala-PG synthases (Fig. 2). The Conserved Domain Architecture

Retrieval Tool [15] lists 148 different architectures of PplT domain-containing proteins. The most

prevalent architectures include combinations with Aa-PG synthase domain (cl41273) of MprF proteins,

glycosyl transferase domains (family A, cl11394; or family B, cl10013), or protein kinase domains

(cl21453) (Fig. 2). MprF proteins were most often found in genomes of Cyanobacteria, Actinobacteria,

Firmicutes, and Proteobacteria. A C-terminal or N-terminal family-B glycosyltransferase domain was

found attached to PplT proteins from Actinobacteria, Nitrospirae, and Chloroflexi or Acidobacteria,

respectively, while a family-A glycosyltransferase domain was found at the N-terminus of PplT in

Archaea of the Euryarchaeota phylum. Several Actinobacteria encode a protein kinase domain, N-

terminally fused to PplT. Further, less frequently occurring combinations were found with enzymatic

domains such as dehalogenase-like hydrolases (cl21460), O-antigen ligases (cl04850), S-

adenosylmethionine-dependent methyltransferase (cl17173), or phosphotransferases (cl37506), to

name a few (Fig. 2). These findings suggest that PplT-containing proteins may have multiple roles,

presumably in the synthesis, modification, and translocation of different types of membrane lipids or

in membrane-linked signal transduction processes. In several bacterial and archaeal taxa, only a

minority of the genera appear to encode PplT proteins in their genomes (Fig. 1) suggesting that PplT

proteins may be important only in certain habitats or that some bacteria may use other transport

systems, that could translocate phospholipids along with other types of cargo molecules.

2. The _E. coli_ PplT is encoded together and interacts with lipid-synthetic enzymes. In order to study

potential functions of PplT proteins not linked to other protein domains, we focused on the _E. coli_

homolog, a protein of unknown function with the generic name YbhN, we suggest to rename PplT.

pplT is not essential in _E. coli_, mutations in this gene are included in the Keio collection, a transposon

mutant library [16], albeit without a reported phenotype. _pplT_ is encoded in an operon together with

the CL synthase gene _clsB_ in _E. coli_ and other Enterobacteriaceae including the genera _Shigella_,

Klebsiella and _Pantoea_ (Fig. 3A), which is in agreement with the role of PplT in lipid synthesis and

translocation.

Phospholipid-biosynthetic proteins have been found to form large multicomponent complexes in

bacteria such as _S. aureus_ [17, 18]. To analyze a role of PplT in such complexes, its potential interaction

with several phospholipid-biosynthetic proteins was studied in the bacterial two-hybrid system [8, 19,

20]. The genes to be analyzed were cloned in _E. coli_ expression plasmids, resulting in C- and N-terminal

fusions of potential interaction partners or PplT with the complementary adenylate cyclase fragment

T18 or T25, respectively. Interaction of T18 with T25 results in the production of cAMP, leading to

expression of β-galactosidase, which was quantified. _E. coli_ does not produce aminoacyl phospholipids

but, in addition to phosphatidylglycerol (PG) and CL, phosphatidylserine (PS) and

phosphatidylethanolamine (PE) [2]. PplT was found to interact with the proteins PgpA, PgpB, and PgpC,

involved in PG synthesis and with the CL synthase ClsB, encoded in the same operon as PplT (Fig. 3A,

B, C). Hence, PplT interacts with the majority of PG and CL biosynthetic enzymes in *E. coli* (Fig. 3C),

which supports a role of PplT in phospholipid metabolism.

3. PplT shapes *E. coli* membrane fluidity and fitness. Although *pplT* is not essential in *E. coli*, its

ubiquitous presence in most of the prokaryotic taxa suggests an important role in basic cellular

functions. *pplT* was deleted in the uropathogenic *E. coli* (UPEC) strain CFT073. The mutant had the

same phospholipid pattern as the wild type (WT) indicating that PplT has no obvious impact on the

overall lipid composition (Fig. 4A, B). However, the mutant showed a significant increase in membrane

fluidity compared to that of the WT strain, as analyzed by monitoring changes in fluorescence

polarization of bacteria stained with 1,6-diphenyl-1,3,5-hexatriene (DPH) (Fig. 4C) [21, 22]. This finding

supports the notion that PplT contributes to membrane homeostasis. The phenotype could be

reverted back to WT level by complementation with a plasmid-encoded *pplT* copy (Fig. 4C). The *pplT*

mutant also exhibited a growth defect compared with the WT in the late logarithmic phase and it did

not reach the cell density of WT cultures (Fig. 5). This phenotype was pronounced at 37°C (Fig. 5A) but

not observed at 42°C (Fig. 5B), which agrees with a role of PplT in membrane fluidity, a process that

depends on temperature. When co-cultivated over several days, the WT grew better than the *pplT*

mutant (Fig. 5C, D), demonstrating a competitive fitness benefit conferred by PplT.

Discussion

Phospholipids are synthesized at the inner leaflet of the cytoplasmic membrane but they need to be

translocated to the outer leaflet to form a stable bilayer [1]. Phospholipids can flip spontaneously but

this process is rather inefficient [23]. Appropriate membrane homeostasis therefore depends on

dedicated flippases that translocate lipids. While such transporters are known for eukaryotic cells, their

presence and nature in prokaryotes has remained largely unclear. ABC transporters translocating the

lipid-bound precursors of peptidoglycan or teichoic acids have been identified [24, 25]. However, these

molecules are usually highly specific and are unlikely to have a major impact on phospholipid

translocation. Nevertheless, the lipid A translocating ABC transporter MsbA of *E. coli* has also a minor

affinity for phospholipids [25]. The only dedicated bacterial phospholipid flippase described to date is

the integral-membrane domain of MprF proteins, which connects the synthesis of aminoacyl

phospholipids with their translocation [4, 5]. We show here that the flippase domain of MprF, named

the PplT domain, can be found as a separate protein or combined in modular proteins with other

domains of diverse function, which supports a general role of PplT in bacterial and archaeal cells.

The PplT protein modulates the membrane fluidity and fitness of *E. coli*. The observed effects were

moderate, which may be based on the fact that the lipid A flippase MsbA has also some basic

phospholipid translocation capacity [25] and may compensate for the lack of PplT to some degree.

msbA is essential though [16, 26], which precludes its simultaneous inactivation with *pplT*. The residual

capacity of MsbA cell wall precursor flippases to translocate phospholipids may also explain why

several bacterial species do not seem to encode obvious PplT orthologs. Membrane fluidity is strongly

affected by temperature, which may explain the temperature-dependent impact of PplT in *E. coli*. It

should be noted though that PplT domain proteins are also found in thermophilic microorganisms such

as Thermotogae and archaeal extremophiles, suggesting that the translocation of certain lipids may

require dedicated flippases even at high temperatures. The frequent combination of PplT with other

enzymatic protein domains also suggests that certain modified lipids, potentially those with bulkier

head groups, may be particularly dependent on cognate translocation machineries. Lipid translocation

is a very dynamic process, which is difficult to monitor directly. Therefore, evidence for the flippase

function of PplT proteins remains in part indirect. However, the fact that PplT interacts directly with

most of the phospholipid-biosynthetic proteins of *E. coli* strongly underscores its central role in

membrane homeostasis. Likewise, MprF proteins have been found to interact with larger

phospholipid-biosynthetic protein complexes in *S. aureus* [17, 18].

It remains unclear how wide or narrow the substrate specificities of PplT domain proteins may be.

Since *E. coli* does not produce aminoacyl phospholipids, PplT may translocate some or all of the *E. coli*

phospholipids PG, CL, PS, and PE. These molecules differ in the net charge of their head groups, which

has been found to be a major determinant for substrate recognition by the PplT domain of the *R. tropici*

MprF [7]. Although the PplT domain of the *S. aureus* MprF is linked to synthesis of the cationic

phospholipid Lys-PG, MprF has been found to also translocate the zwitterionic phospholipid Ala-PG

[27], indicating that PplT proteins may have broader substrate specificities. In agreement with this

assumption, a few point mutations have been shown to alter the function and probably substrate

specificity of the *S. aureus* MprF [28]. It should be noted that most prokaryotic genomes with a PplT

homolog contain only one PplT protein and only a minority encodes two (19.89%) or more (1.60%) of

them, suggesting that one flippase is usually sufficient for most of the membrane-forming

phospholipids. The presence or absence of certain conserved, charged amino acids in trans-membrane

segments of PplT domains may help to predict the substrate specificities of these proteins in the future.

Archaea have different phospholipids compared to bacterial ones, with monolayer-membrane forming

isoprenoid lipids [29]. The broad presence of PplT proteins in Archaea suggest that these proteins can

also flip such lipids, which are much bulkier than bacterial phospholipids and contain two head groups,

one at each end. It remains unclear if the PplT domain only facilitates the exchange of phospholipids

between the inner and outer leaflet of cytoplasmic membranes or if it can translocate phospholipids

in an energy-dependent fashion to generate asymmetric lipid patterns. Some of the transmembrane

domains of MprF seem to be related to the proton-motif force-dependent major facilitator protein

superfamily [30] and some studies found Lys-PG to be unevenly distributed between the two

membrane leaflets [4], suggesting that PplT may use energy to translocate its substrate lipids.

Membrane homeostasis is an important fitness factor for all kinds of organisms. PplT proteins may

therefore become attractive targets for future anti-fitness drugs to combat major human pathogens.

In support of such a strategy, monoclonal antibodies directed against extracellular portions of the PplT

domain of the *S. aureus* MprF have recently been shown by inhibiting the Lys-PG exposure at the outer

leaflet of the membrane and therefore sensitize *S. aureus* to cationic antibiotics [9].

Materials and Methods

Occurrence and architecture of PplT in bacterial and archaeal taxa. In order to determine the

distribution of PplT across the taxonomy a blastp search within NCBI's non-redundant protein database

[31] was performed using the PplT/YbhN protein sequence of *E. coli* (old: NP_752801.1; new:

WP_000045478.1) as a query and an E-value cutoff of 0.05. Blast hits originating from the

superkingdoms Bacteria and Archaea were investigated separately. For each, the subtree of the NCBI

taxonomy rooted at the respective superkingdom was thinned out for those taxa that possess a PplT

homolog. The BLAST search and subsequent taxonomic visualization was performed with BLASTphylo

[https://github.com/Integrative-Transcriptomics/BLASTphylo]. We used NCBI's Conserved Domain

Architecture Retrieval Tool (CDART) [15] to identify different domain architectures that include a PplT

domain. We used MicrobesOnline[32] to identify *pplT*'s operon structure in other Enterobacteriaceae.

Bacterial strains, mutagenesis, and maintenance. *E. coli* UPEC strain CFT073 (DSM103538) (Tab. S1)

was used for *pplT* mutagenesis and analysis [33]. *pplT* was deleted in the chromosome of CFT073 as

described previously [34]. Briefly, *E. coli* CFT073 was transformed with helper plasmid pKD46 encoding

the Lambda-Red recombinase (Tab. S2) and incubated at 30°C on LB agar supplemented with

ampicillin. The kanamycin resistance cassette with chromosomal DNA regions flanking *pplT* was

amplified by PCR with primers pplT_KO_5 and pplt_KO_6 (Tab. S3) using pKD13 as template (Tab. S2).

After arabinose-induced Lambda Red expression in *E. coli* CFT073 containing the helper plasmid, the

PCR fragment was transferred by electroporation and cells were plated on kanamycin-containing LB

agar for primary selection. After verification of the chromosomal *pplT* deletion by PCR (Tab. S3) and

sequencing, mutants were streaked to single colonies on LB agar without antibiotics at 37°C twice for

curing of the temperature-sensitive helper plasmid. The loss of the helper plasmid pKD46 was

confirmed by PCR (Tab. S3). The *pplT* deletion mutant was complemented by a *pplT* gene amplified

from *E. coli* K12 and cloned in the *E. coli/S. aureus* shuttle vector pRB474 [35] (Tab. S2). All bacterial

strains were grown in LB medium (Carl Roth) with appropriate antibiotics unless otherwise noted (Tab.

S1).

Bacterial two-hybrid assay. To identify potential interaction partners of PplT, the commercially

available bacterial two-hybrid kit (BATCH kit, Euromedex) was used [8, 19, 20] and the proteins to be

studied were cloned as described previously [8]. The *pplT* gene was cloned in the high-copy number

plasmid pUT18C, the potential interaction partners in the low copy-number plasmid pKT25 leading to

C-terminal or N-terminal fusion with the adenylate cyclase fragments T25 or T18 of *Bordetella*

pertussis, respectively (Tab. S2). The constructs were used to co-transform chemically competent *E.*

coli BTH101. The resulting transformants were tested in a 96-well format for β-galactosidase activity,

to quantify the protein-protein interaction as described previously [8] with slight modifications. Briefly,

bacterial cells were grown at 30°C overnight in LB, supplemented with 0.5 mM isopropyl-β-D-

thiogalactopyranoside (IPTG), 25 µg/ml kanamycin and 100 µg/ml ampicillin. 100 µl of the overnight

cultures were used to measure the optical density at 600 nm (OD_{600}) in a 96-well microtiter plate (F-

bottom, Falcon), another 100 µl were transferred into a deep 96-well plate (U-bottom, Sarstedt) and

mixed with 1 ml buffer Z (60 mM Na_2HPO_4, 40 mM NaH_2PO_4, 10 mM KCl, 1 mM $MgSO_4$, 50 mM β-

mercaptoethanol). To lyse the cells, 40 µl sodium dodecyl sulfate (0.1%) and 80 µl chloroform were

added and mixed vigorously. After incubation at room temperature to allow phase separation, 100 µl

of the aqueous phase were transferred into a 96-well microtiter plate (F-bottom, Falcon) and mixed

with 20 µl o-nitrophenyl-β-D-galactopyranoside (4 mg/ml) to start the enzymatic reaction. The

reaction was measured continuously for 1 h in a CLARIOStar plate reader (BMG Labtech) at OD_{420} and

OD_{550}. The β-galactosidase activity in Miller Units was calculated by the formula: $1000*((OD_{420}-$

$(1.75*OD_{550}))/(t*v*OD_{600}))$, where t is reaction time in minutes and v the reaction volume in milliliter.

Isolation of polar lipids. Polar lipids were extracted from bacteria grown to exponential or stationary

phase using the Bligh-Dyer procedure [36]. Briefly, phospholipids were extracted by a mixture of

sodium acetate buffer (20mM, pH 4.8), chloroform, and methanol (1:1:1 [by volume]), vacuum dried,

and dissolved in chloroform-methanol (2:1 [by volume]). For detection of phospholipid patterns,

appropriate amounts of polar lipid extracts were spotted onto silica gel high-performance thin-layer

chromatography (HPTLC) plates (silica gel 60 F_{254}, Merck) using a Linomat 5 sample application unit

(CAMAG) and developed with chloroform:methanol:water (65:25:4 [by volume]) in an automatic

developing chamber ADC 2 (CAMAG). Phospholipids were selectively stained with molybdenum blue

spray reagent (1.3% in 4.2 M sulfuric acid, Sigma-Aldrich).

Measurement of the membrane fluidity. To measure the membrane fluidity of *E. coli* CFT073 WT and

mutants, we used the method of fluorescence polarization upon staining with 1,6-diphenyl-1,3,5-

hexatriene (DPH) as described previously [21, 22], with some modifications. Briefly, bacterial cultures

grown to exponential phase were adjusted to an OD_{600} 1 and treated with a subinhibitory

concentration of polymyxin B (30 µg/ml) for 30 min at 37°C for outer membrane permeabilization

purposes. Bacteria were then stained with 8 µM DPH for further 30 min. After washing with PBS, DPH-

stained bacteria were transferred to 96-well microtiter plate (F-bottom, black, Greiner) and

fluorescence polarization was measured at an excitation wavelength of 360-10 nm, and emission

wavelength of 450-10 nm in a CLARIOStar plate reader (BMG Labtech). Polarization values inversely

correlate with membrane fluidity.

Bacterial growth assay. Temperature-dependent growth differences were analyzed in 96-well

microtiter plate format. To this end, overnight cultures grown in LB with appropriate antibiotics were

adjusted to OD_{600} 0.05 in fresh LB and 100 µl of the cell suspensions were transferred to a 96-well

microtiter plate (F-bottom, Falcon) and growth was measured at OD_{600} in 30 min intervals at either

37°C or 42°C in an Epoch2 plate reader (BioTek).

To investigate growth differences of WT and *pplT* mutant in cocultivation, overnight cultures were

adjusted to the same colony forming unit in fresh LB and used to inoculate a coculture, which was

incubated in a shaking incubator at either 37°C or 42°C. Cells were passaged every 24 h in fresh LB for

five days. To measure the numbers of live cells of the two competing strains, appropriate dilutions of

the suspensions were spotted on LB agar plates with or without kanamycin. After incubation of the

agar plates at appropriate temperatures, colonies were counted and ratios were calculated.

Statistics. Statistical analyses were performed with the Prism 8.4.2 package (GraphPad Software) and

Group differences were analyzed for significance with one-way or two-way ANOVA. A P value of ≤0.05

was considered statistically significant.

Acknowledgments

We thank Anne Berscheid for providing plasmids pKD13 and pKD46, Cordula Gekeler and Ulrike Redel

for technical support and Jennifer Müller for her contributions to the BLASTphylo project. This work

was financed by Grants from Deutsche Forschungsgemeinschaft (DFG) SFB766 (to A.P.) and TRR261,

project ID 398967434 (to K.N. and A.P), the German Center of Infection Research (DZIF) (to A.P. and

C.S.). C.S. was also supported by the intramural Experimental Medicine program of the Medical Faculty

at the University of Tübingen. The authors acknowledge project support (C.S.) and infrastructural

support (C.S., K.N., A.P.) by the DFG Cluster of Excellence EXC2124 CMFI, project ID 390838134.

References

1.   Zhang YM, Rock CO. Membrane lipid homeostasis in bacteria. Nat Rev Microbiol. 2008;6(3):222-33. doi: 10.1038/nrmicro1839. PubMed PMID: 18264115.

2.   Dowhan W, Bogdanov M. Eugene P. Kennedy's Legacy: Defining Bacterial Phospholipid Pathways and Function. Front Mol Biosci. 2021;8:666203. Epub 2021/04/13. doi: 10.3389/fmolb.2021.666203. PubMed PMID: 33842554; PubMed Central PMCID: PMCPMC8027125.

3.   López-Marqués RL. Lipid flippases in polarized growth. Curr Genet. 2021;67(2):255-62. Epub 2021/01/04. doi: 10.1007/s00294-020-01145-0. PubMed PMID: 33388852.

4.   Ernst CM, Staubitz P, Mishra NN, Yang SJ, Hornig G, Kalbacher H, et al. The bacterial defensin resistance protein MprF consists of separable domains for lipid lysinylation and antimicrobial peptide repulsion. PLoS Pathog. 2009;5(11):e1000660. Epub 2009/11/17. doi: 10.1371/journal.ppat.1000660. PubMed PMID: 19915718; PubMed Central PMCID: PMCPMC2774229.

5.   Ernst CM, Peschel A. Broad-spectrum antimicrobial peptide resistance by MprF-mediated aminoacylation and flipping of phospholipids. Mol Microbiol. 2011;80(2):290-9. Epub 2011/02/11. doi: 10.1111/j.1365-2958.2011.07576.x. PubMed PMID: 21306448.

6.   Peschel A, Jack RW, Otto M, Collins LV, Staubitz P, Nicholson G, et al. Staphylococcus aureus resistance to human defensins and evasion of neutrophil killing via the novel virulence factor MprF is based on modification of membrane lipids with l-lysine. J Exp Med. 2001;193(9):1067-76. Epub 2001/05/09. doi: 10.1084/jem.193.9.1067. PubMed PMID: 11342591; PubMed Central PMCID: PMCPMC2193429.

7.   Song D, Jiao H, Liu Z. Phospholipid translocation captured in a bifunctional membrane protein MprF. Nat Commun. 2021;12(1):2927. Epub 2021/05/20. doi: 10.1038/s41467-021-23248-z. PubMed PMID: 34006869; PubMed Central PMCID: PMCPMC8131360.

8.   Ernst CM, Kuhn S, Slavetinsky CJ, Krismer B, Heilbronner S, Gekeler C, et al. The lipid-modifying multiple peptide resistance factor is an oligomer consisting of distinct interacting synthase and flippase subunits. mBio. 2015;6(1). Epub 2015/01/30. doi: 10.1128/mBio.02340-14. PubMed PMID: 25626904; PubMed Central PMCID: PMCPMC4324311.

9.   Slavetinsky CJ, Hauser JN, Gekeler C, Slavetinsky J, Geyer A, Kraus A, et al. Sensitizing Staphylococcus aureus to antibacterial agents by decoding and blocking the lipid flippase MprF. Elife. 2022;11. Epub 2022/01/20. doi: 10.7554/eLife.66376. PubMed PMID: 35044295.

10.   Yang M, Derbyshire MK, Yamashita RA, Marchler-Bauer A. NCBI's Conserved Domain Database and Tools for Protein Domain Analysis. Curr Protoc Bioinformatics. 2020;69(1):e90. Epub 2019/12/19. doi: 10.1002/cpbi.90. PubMed PMID: 31851420; PubMed Central PMCID: PMCPMC7378889.

11.   Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The InterPro protein families and domains database: 20 years on. Nucleic Acids Res. 2021;49(D1):D344-d54. Epub 2020/11/07. doi: 10.1093/nar/gkaa977. PubMed PMID: 33156333; PubMed Central PMCID: PMCPMC7778928.

12.   Roy H, Ibba M. Broad range amino acid specificity of RNA-dependent lipid remodeling by multiple peptide resistance factors. J Biol Chem. 2009;284(43):29677-83. doi: 10.1074/jbc.M109.046367. PubMed PMID: 19734140; PubMed Central PMCID: PMCPMC2785599.

13.   Cronan JE, Jr., Rock CO. Biosynthesis of Membrane Lipids. EcoSal Plus. 2008;3(1). Epub 2008/09/01. doi: 10.1128/ecosalplus.3.6.4. PubMed PMID: 26443744.

14.   Furse S, Wienk H, Boelens R, de Kroon AI, Killian JA. E. coli MG1655 modulates its phospholipid composition through the cell cycle. FEBS Lett. 2015;589(19 Pt B):2726-30. Epub 2015/08/15. doi: 10.1016/j.febslet.2015.07.043. PubMed PMID: 26272829.

15.   Geer LY, Domrachev M, Lipman DJ, Bryant SH. CDART: protein homology by domain architecture. Genome Res. 2002;12(10):1619-23. Epub 2002/10/09. doi: 10.1101/gr.278202. PubMed PMID: 12368255; PubMed Central PMCID: PMCPMC187533.

16.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol.

2006;2:2006.0008. Epub 2006/06/02. doi: 10.1038/msb4100050. PubMed PMID: 16738554; PubMed Central PMCID: PMCPMC1681482.

17. García-Lara J, Weihs F, Ma X, Walker L, Chaudhuri RR, Kasturiarachchi J, et al. Supramolecular structure in the membrane of Staphylococcus aureus. Proc Natl Acad Sci U S A. 2015;112(51):15725-30. Epub 2015/12/09. doi: 10.1073/pnas.1509557112. PubMed PMID: 26644587; PubMed Central PMCID: PMCPMC4697411.

18. Weihs F, Wacnik K, Turner RD, Culley S, Henriques R, Foster SJ. Heterogeneous localisation of membrane proteins in Staphylococcus aureus. Sci Rep. 2018;8(1):3657. Epub 2018/02/28. doi: 10.1038/s41598-018-21750-x. PubMed PMID: 29483609; PubMed Central PMCID: PMCPMC5826919.

19. Karimova G, Gauliard E, Davi M, Ouellette SP, Ladant D. Protein-Protein Interaction: Bacterial Two-Hybrid. Methods Mol Biol. 2017;1615:159-76. Epub 2017/07/02. doi: 10.1007/978-1-4939-7033-9_13. PubMed PMID: 28667611.

20. Griffith KL, Wolf RE, Jr. Measuring beta-galactosidase activity in bacteria: cell growth, permeabilization, and enzyme assays in 96-well arrays. Biochem Biophys Res Commun. 2002;290(1):397-402. Epub 2002/01/10. doi: 10.1006/bbrc.2001.6152. PubMed PMID: 11779182.

21. Chen Y, Reinhardt M, Neris N, Kerns L, Mansell TJ, Jarboe LR. Lessons in Membrane Engineering for Octanoic Acid Production from Environmental Escherichia coli Isolates. Appl Environ Microbiol. 2018;84(19). Epub 2018/07/22. doi: 10.1128/aem.01285-18. PubMed PMID: 30030228; PubMed Central PMCID: PMCPMC6146996.

22. Gohrbandt M, Lipski A, Grimshaw JW, Buttress JA, Baig Z, Herkenhoff B, et al. Low membrane fluidity triggers lipid phase separation and protein segregation in living bacteria. Embo j. 2022:e109800. Epub 2022/01/18. doi: 10.15252/embj.2021109800. PubMed PMID: 35037270.

23. Kol MA, de Kroon AI, Killian JA, de Kruijff B. Transbilayer movement of phospholipids in biogenic membranes. Biochemistry. 2004;43(10):2673-81. Epub 2004/03/10. doi: 10.1021/bi036200f. PubMed PMID: 15005602.

24. Swoboda JG, Meredith TC, Campbell J, Brown S, Suzuki T, Bollenbach T, et al. Discovery of a small molecule that blocks wall teichoic acid biosynthesis in Staphylococcus aureus. ACS Chem Biol. 2009;4(10):875-83. Epub 2009/08/20. doi: 10.1021/cb900151k. PubMed PMID: 19689117; PubMed Central PMCID: PMCPMC2787957.

25. Eckford PD, Sharom FJ. The reconstituted Escherichia coli MsbA protein displays lipid flippase activity. Biochem J. 2010;429(1):195-203. Epub 2010/04/24. doi: 10.1042/bj20100144. PubMed PMID: 20412049; PubMed Central PMCID: PMCPMC2888566.

26. Karow M, Georgopoulos C. The essential Escherichia coli msbA gene, a multicopy suppressor of null mutations in the htrB gene, is related to the universally conserved family of ATP-dependent translocators. Mol Microbiol. 1993;7(1):69-79. Epub 1993/01/01. doi: 10.1111/j.1365-2958.1993.tb01098.x. PubMed PMID: 8094880.

27. Slavetinsky CJ, Peschel A, Ernst CM. Alanyl-phosphatidylglycerol and lysyl-phosphatidylglycerol are translocated by the same MprF flippases and have similar capacities to protect against the antibiotic daptomycin in Staphylococcus aureus. Antimicrob Agents Chemother. 2012;56(7):3492-7. Epub 2012/04/12. doi: 10.1128/aac.00370-12. PubMed PMID: 22491694; PubMed Central PMCID: PMCPMC3393434.

28. Ernst CM, Slavetinsky CJ, Kuhn S, Hauser JN, Nega M, Mishra NN, et al. Gain-of-Function Mutations in the Phospholipid Flippase MprF Confer Specific Daptomycin Resistance. MBio. 2018;9(6). Epub 2018/12/20. doi: 10.1128/mBio.01659-18. PubMed PMID: 30563904; PubMed Central PMCID: PMCPMC6299216.

29. Caforio A, Driessen AJM. Archaeal phospholipids: Structural properties and biosynthesis. Biochim Biophys Acta Mol Cell Biol Lipids. 2017;1862(11):1325-39. Epub 2016/12/23. doi: 10.1016/j.bbalip.2016.12.006. PubMed PMID: 28007654.

30. Wang SC, Davejan P, Hendargo KJ, Javadi-Razaz I, Chou A, Yee DC, et al. Expansion of the Major Facilitator Superfamily (MFS) to include novel transporters as well as transmembrane-acting enzymes. Biochim Biophys Acta Biomembr. 2020;1862(9):183277. Epub 2020/03/25. doi:

10.1016/j.bbamem.2020.183277. PubMed PMID: 32205149; PubMed Central PMCID: PMCPMC7939043.

31.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res. 2016;44(D1):D733-45. Epub 2015/11/11. doi: 10.1093/nar/gkv1189. PubMed PMID: 26553804; PubMed Central PMCID: PMCPMC4702849.

32.   Dehal PS, Joachimiak MP, Price MN, Bates JT, Baumohl JK, Chivian D, et al. MicrobesOnline: an integrated portal for comparative and functional genomics. Nucleic Acids Res. 2010;38(Database issue):D396-400. Epub 2009/11/13. doi: 10.1093/nar/gkp919. PubMed PMID: 19906701; PubMed Central PMCID: PMCPMC2808868.

33.   Welch RA, Burland V, Plunkett G, 3rd, Redford P, Roesch P, Rasko D, et al. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic Escherichia coli. Proc Natl Acad Sci U S A. 2002;99(26):17020-4. Epub 2002/12/10. doi: 10.1073/pnas.252529799. PubMed PMID: 12471157; PubMed Central PMCID: PMCPMC139262.

34.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97(12):6640-5. Epub 2000/06/01. doi: 10.1073/pnas.120163297. PubMed PMID: 10829079; PubMed Central PMCID: PMCPMC18686.

35.   Bruckner R. A series of shuttle vectors for Bacillus subtilis and Escherichia coli. Gene. 1992;122(1):187-92. PubMed PMID: 1452028.

36.   Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. Can J Biochem Physiol. 1959;37(8):911-7. Epub 1959/08/01. doi: 10.1139/o59-099. PubMed PMID: 13671378.

Figures

Figure 1: Distribution of the PpIT domain amongst taxa of the Bacteria (A) and Archaea (B) superkingdoms. The taxonomy was collapsed to the phylum rank. Dark grey leaf nodes indicate collapsed nodes. The bar chart indicates the fraction of genera of a particular Phylum (Bacteria) or

Class (Archaea) carrying a PpIT homolog within the subtree rooted at the respective taxon. #, taxa include *pplT* genes but percentages of covered genera could not be calculated due to incomplete assignment of the genome on genus level.

Figure 2: PplT domain protein architectures. Combinations of the PplT domain (cl04219) with other protein domains found in Bacteria and Archaea. The 10 most commonly occurring domain combinations are shown, as revealed by the Conserved Domain Architecture Retrieval Tool [15]. The domains are not drawn to scale.

Figure 3: Genetic context and interaction partners of PplT in *E. coli*. (A) PplT is encoded in an operon together with YbhP, a protein of unknown function, and the cardiolipin synthase ClsB in Enterobacteriaceae, including the genera *Escherichia, Shigella*, *Klebsiella,* and *Pantoea*. The genes of the operon are not drawn to scale. (B) Schematic representation of the phospholipid synthesis pathways in *E. coli*. Cytidine diphosphate diacylglycerol (CDP-DAG), phosphatidylglycerol-3-phosphate (PGP), phosphatidylglycerol (PG), cardiolipin (CL), phosphatidylserine (PS), phosphatidylethanolamine (PE). Enzymes are marked in blue or grey, if they interacted with PplT or not, respectively. (C) Interaction studies with PplT and lipid-biosynthetic enzymes using the bacterial two-hybrid system and β-galactosidase assay for quantification. Data represent means plus SD of five biological replicates. Statistical significances were determined by two-way ANOVA followed by Dunnett's test in comparison to the negative control (ns, not significant; ****, P < 0.0001). Figures A and B were created with BioRender.com.

Figure 4: Impact of PplT on cell membrane lipid pattern and fluidity. Detection of phospholipids of *E. coli* CFT073 WT, *pplT* mutant and complemented mutant at exponential (A) or stationary (B) growth phase. Total lipids were isolated, separated via thin-layer chromatography and stained using molybdenum blue spray reagent. PE and PG spots are indicated, concentrations of the minor lipids PS and CL are too low to be detectable. Three biological replicates are represented. (C) Membrane fluidity analyzed via DPH staining. Bacterial cells were stained during exponential growth with DPH and polarization was measured. Means of three biological replicates with SEM are shown. Data were compared to the WT and statistically analyzed by one-way ANOVA (ns, not significant; *, $P < 0.05$).

Figure 5: The *E. coli pplT* mutant exhibits a growth deficit compared to the WT. Growth curves of *E. coli* CFT073 WT, *pplT* mutant and complemented mutant at 37°C (A) or 42°C (B). Means of three biological replicates plus SEM are shown. The difference in growth between WT and *pplT* mutant was statistically analyzed by two-way ANOVA (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.0005$; ****, $P < 0.0001$). Competition assay of WT and the *pplT* mutant over five days at 37°C (C) or 42°C (D). Means of three biological replicates with SD are shown. Data were statistically analyzed by two-way ANOVA and compared to the WT at the appropriate timepoint (ns, not significant; **, $P < 0.01$; ****, $P < 0.0001$).

Supplemental Data

Table S1: Bacterial strains.

Strain	Characteristics
E. coli CFT073	Uropathogenic wild type strain [33]
E. coli CFT073 Δ*pplT*	*pplT* chromosomal deletion mutant, *pplT* gene was replaced by kanamycin resistance cassette, Kan^R. This study.
E. coli CFT073 Δ*pplT* pRB*pplT*	*pplT* deletion mutant complemented by plasmid-encoded *pplT* version. This study.

Table S2: Plasmids.

Plasmid	Characteristics
pRB474*pplT*	*pplT* gene cloned in *E. coli/S. aureus* shuttle vector pRB474 [35], Amp^R.
pKD13	Template for Kanamycin resistance cassette. Amp^R, Kan^R [34].
pKD46	Lambda Red helper plasmid, Lambda Red expression inducible by L-arabinose. Amp^R [34].
pKT25-*clsA*	*clsA* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*clsB*	*clsB* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*clsC*	*clsC* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*ybhP*	*ybhP* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*pssA*	*pssA* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*psd*	*psd* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*pgsA*	*pgsA* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*pgpA*	*pgpA* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*pgpB*	*pgpB* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-*pgpC*	*pgpC* gene encoded on low copy number plasmid pKT25 (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T25. This study.
pKT25-zip	Leucine zipper of GCN4 fused to the C-terminal end of the adenylate cyclase fragment T25. BATCH system kit, Euromedex.
pUT18C-*pplT*	*pplT* gene encoded on high copy number plasmid pUT18C (Euromedex), fused to the C-terminal end of the adenylate cyclase fragment T18. This study.
pUT18C-zip	Leucine zipper of GCN4 fused to the C-terminal end of the adenylate cyclase fragment T18. BATCH system kit, Euromedex.

Table S3: Primers.

Primer	Sequence 5' → 3'	Usage

pplT_KO_5	ATGCACATACTAAAGAACTTAACTATACTT CACATCGCCGCTTCATTTTTTGTAGGCTGG AGCTGCTTCG	Amplification of kanamycin resistance cassette from pKD13 with overhang homolog to the chromosomal *pplT* surrounding; for construction of the chromosomal *pplT* deletion mutant in *E. coli* CFT073.
pplT_KO_6	GAAACGCAGGATCGGGTAGAAACTGAAA ACACGGGGGTAAAACCCTGATGATTCCGG GGATCCGTCGACC	
pKD46_gone_1	CCCGTGCGTTTGATGACGATG	Primers to confirm the curing of the helper plasmid pKD46.
pKD46_gone_2	GGATTCATTGTCCTGCTCAAAGTCC	
pRB_pplT_BamHI_fw	AAATTATggatccCAGAGGAGGTCGGCTGA TGAGTAAATCAC	Cloning of *pplT* into pRB474 for plasmid-based complementation of the *pplT* deletion mutant.
pRB_pplT_EcoRI_rev	AAATTATgaattcAACTAAACTTCACATCGCC GCTTC	
pplT_up	ggttcgttgtgcttttatagg	Sequencing primers to verify the chromosomal pplT deletion mutant.
pplT_down	cttcccggcgctgg	
ClsA_BamHI_for	TAATCggatccTATGACAACCGTTTATACG	Cloning of *clsA* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
ClsA_EcoRI_rev	ACGTTgaattcTTACAGCAACGGGCTG	
ClsB_BamHI_for	TCATTggatccTATGAAATGTAGCTGGCGCG	Cloning of *clsB* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
ClsB_EcoRI_rev	ACTCAgaattcTCAGGGTTTTACCCCCGTG	
ClsC_BamHI_for	GTCTGTggatccTATGATGAAGAAAACGCCCAC	Cloning of *clsC* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
ClsC_EcoRI_rev	CGCGTgaattcTTACAATAACCATTCCACGG	
YbhP_BamHI_for	GAAAATggatccTATGCCCGATCAAACAC	Cloning of *ybhP* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
YbhP_EcoRI_rev	ACATTgaattcTCATAAATGAATCTCCGC	
PssA_BamHI_for	CATCCggatccTATGAAAATGACAAAACTGG	Cloning of *pssA* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
PssA_EcoRI_rev	CAGGGgaattcTTACTGCGTGGTACCG	
Psd_BamHI_for	CACTGTggatccGATGTTGTCAAAATTTAAGCG	Cloning of *psd* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
Psd_KpnI_rev	TGTGAgaattcTTACAGGATGCGGCTAATTAATC	
PgsA_BamHI_for	GCTACggatccCTTGTTAAATTCATTTAAAC	Cloning of *pgsA* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
PgsA_EcoRI_rev	ATGGAggtaccTTAGACCTGGTCTTTTTTG	
PgpA_BamHI_for	GTCATggatccTATGCAATTTAATATCCCTAC	Cloning of *pgpA* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
PgpA_EcoRI_rev	AACGAgaattcTCACTGATCAAGCAAATCTG	
PgpB_BamHI_for	AAGGAggatccTATGACCATTTTGCCACGCC	Cloning of *pgpB* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
PgpB_EcoRI_rev	CACAAgaattcCTACGACAGAATACCCAGC	
PgpC_BamHI_for	GAGGCggatccCATGCGTTCGATTGCC	Cloning of *pgpC* from *E. coli* CFT073 into pKT25 for bacteria two-hybrid.
PgpC_EcoRI_rev	CAGCGgaattcTTAACTTTCTTGTTCTCGTTG	
pUT18_PplT_BamHI_for	CCCTGggatccGATGAGTAAATCACACCCGCG	Cloning of *pplT* from *E. coli* CFT073 into pUT18C for bacteria two-hybrid.
PplT_EcoRI_rev	ATACTgaattcTCACATCGCCGCTTCATTTTTC	
pKT25 fw_2	CAACTTCCGCGACTCGGCG	Sequencing primer to verify insert of pKT25.
pKT25 rev	gattaagttgggtaacgccag	
pUT18C for	CGTCACCCGGATTGCGGC	Sequencing primer to verify insert of pUT18C.
pUT18C rev	cggggctggcttaactatgc	

Figure S1: Most PplT containing genomes of prokaryotes harbor only one homolog. Number of genomes with one, two or more genes encoding PplT domains are shown.