Manuscript title: **Antecedent effect models as an exploratory tool to link climate drivers to**

herbaceous perennial population dynamics data

Author names: Aldo Compagnoni[1,2,3.], Dylan Childs[4], Tiffany M. Knight[1,2,3], Roberto Salguero-

Gómez[6]

Affiliations: [1] Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale),
Germany.

Deutscher Platz 5e, Leipzig, 04103, Germany; [2] Department of Community Ecology, Helmholtz

Centre for Environmental Research–UFZ, 06120 Halle (Saale), Germany; [3] German Centre for

Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Puschstrasse 4, Leipzig 04103,

Germany; [4] Department of Animal and Plant Sciences, University of Sheffield. Western Bank,

Sheffield S10 2TN, UK; [6] Department of Biology, University of Oxford. South Parks Road,

Oxford, OX1 3SZ, United Kingdom.

Corresponding Author: Aldo Compagnoni; aldo.compagnoni@gmail.com

Open Research statement: Data and code to reproduce the analyses are available on zenodo at

https://dx.doi.org/10.5281/zenodo.7839199.

Keywords: Bayesian moving window, environmental drivers, demography, lag, lagged climate

effect, delay, COMPADRE Plant Matrix Database, cross-validation, forecasting, population

growth rate, regularization, vital rate.

Abstract

Understanding mechanisms and predicting natural population responses to climate is a key goal of Ecology. However, studies explicitly linking climate to population dynamics remain limited. Antecedent effect models are a set of statistical tools that capitalize on the evidence provided by climate and population data to select time windows correlated with a response (*e.g.*, survival, reproduction). Thus, these models can serve as both a predictive and exploratory tool. We compare the predictive performance of antecedent effect models against simpler models, and showcase their exploratory analysis potential by selecting a case study with high predictive power. We fit three antecedent effect models: (1) weighted mean models (WMM), which weigh the importance of monthly anomalies based on a Gaussian curve, (2) stochastic antecedent models (SAM), which weigh the importance of monthly anomalies using a Dirichlet process, and (3) regularized regressions using the Finnish Horseshoe prior (FHM), which estimate a separate effect size for each monthly anomaly. We compare these approaches to a linear model using a yearly climatic predictor and a null model with no predictors. We use demographic data from 77 natural populations of 34 plant species ranging between seven and 36 years of length. We then fit models to the asymptotic population growth rate (λ) and its underlying vital rates: survival, development, and reproduction. We find that models including climate do not consistently outperform null models. We hypothesize that the effect of yearly climate is too complex, weak, and confounded by other factors to be easily predicted using monthly precipitation and temperature data. On the other hand, in our case study, antecedent effect models show biologically sensible correlations between two precipitation anomalies and multiple vital rates. We conclude that, in temporal datasets with limited sample sizes, antecedent effect models are better suited as exploratory tools for hypothesis generation.

Introduction

Understanding and predicting population dynamics is a key objective of Ecology (Sutherland et al. 2013). In population ecology, the last decades have witnessed the development and application of many statistical methods aimed precisely at this objective (Caswell 2001, Ellner et al. 2016). Increasingly, emphasis is being devoted to understanding how climatic drivers such as temperature and precipitation affect vital rates like survival, development (*i.e.,* changes in stage along the life cycle), and reproduction (Ehrlén et al. 2016; Teller et al. 2016). The ultimate goal of this ambitious research agenda is to achieve a mechanistic understanding of what makes species vulnerable to climate change worldwide (Sutherland et al. 2013). However, progress in this understanding is underpinned by synthetic studies that explicitly link vital rates to climatic drivers using standardized methods (Ehrlén et al. 2016, Compagnoni et al. 2021).

To understand and predict the effects of climate drivers on populations, ecologists need to estimate the predictability of different response variables. While ecologists are ultimately interested in linking population performance metrics (*e.g.*, population growth rate) to climatic drivers, this link often has poor predictive ability (Knape and de Valpine 2011, Tredennick et al. 2016). The low predictive performance might occur because *a priori* knowledge of the system is often poor, and/or because the number of potential covariates is orders of magnitude larger than the number of data points available in population studies (Tredennick et al. 2021). In addition, poor predictive performance might reflect how variation in population growth rates might be buffered against environmental stochasticity, in that responses of single vital rates might not fully translate into population growth rate responses (Hilde et al. 2020). Thus, we might have a better understanding and ability to predict the effects of climate on vital rates (*e.g.*, Clark et al., 2016, 2021; Schulze et al., 2019). Alternatively, the low predictive ability of our models might

simply reflect how the approaches to link populations dynamics to climate implemented so far

are ineffective. Currently, we lack a key piece of the puzzle: the relative ability of existing

approaches to predict vital rates and population growth rate, and how this predictability varies

across species. Establishing this predictive ability in synthesis study requires addressing the

technical hurdles associated with linking climatic drivers to population dynamics.

Studies that explicitly link population responses to climate drivers typically identify

climatic predictors *a priori*. Most population studies are short-term (Salguero-Gómez et al. 2015,

2016, Römer et al. 2021, Levin et al. 2022), thus preventing investigators from effectively using

exploratory or model selection approaches. As a result, ecologist have historically considered

seasonal or, more frequently, annual climatic effects (reviewed in Evers et al., 2021). This choice

has practical reasons: first, monthly or annual climate data are widely available (Daly et al. 1994,

Karger and Zimmermann 2018). Second, annual anomalies are a logical predictor of annual

demographic variation, and seasonal anomalies can be justified by physiological or behavioural

considerations (White et al. 1999, Catchpole et al. 2000). However, no consensus on these *a*

priori choices exists. For example, expectations based on physiology, such as the importance of

the growing season in plants, or the importance of exceeding minimum thermal limits in insects

(Angilletta, 2009), are often met with opposing evidence (Kreyling 2010, Czachura and Miller

2020), or evidence of lagged effects from previous years (Hacket□Pain et al. 2018, Tenhumberg

et al. 2018, Evers et al. 2021). At worst, models that link population responses to monthly or

annual climatic variables could be an excessive oversimplification of the processes that drive

plant physiology. The effect of climate on plants unfolds via the dynamic, nonlinear interaction

of temperature, soil moisture, relative humidity, and radiative fluxes at sub-hourly time scales

(Lambers et al. 2008). While these factors correlate with annual or seasonal temperature and

precipitation, such correlation might be too weak to make annual or seasonal climate effective predictors of population dynamics at a yearly resolution.

If the most common *a priori* choices on the climate predictors of population fluctuations are suboptimal, then more complex models could increase predictive power. This would be the case for models that simultaneously identify the critical time window and magnitude of climatic effects that drive their system dynamics. Several such models have been proposed that automatically select the timing most relevant to population processes (van de Pol and Cockburn 2011, Ogle et al. 2015). Henceforth, we refer to these models as "antecedent effect models". These models control for multiple comparisons, and weigh the time windows when drivers are most influential for population dynamics, dividing the climate (or any other relevant factor) of a time window (*e.g.*, a year) into sub-periods of equal lengths – usually months. Such weighing can occur by estimating the importance of each monthly anomaly using weights constrained to follow a specific function (van de Pol and Cockburn 2011, Ogle et al. 2015). Alternatively, the weighing can be performed by estimating a different effect size for each monthly climate anomaly, using regularization or informative priors to minimize overfitting (Hoerl & Kennard, 1970). In cases where the timing of climatic predictors is not crucial in explaining population responses to climate, antecedent effect models would have low predictive power when compared to simpler models. Moreover, it has been suggested that, when the climatic effects on populations are small (e.g. due to their ability to buffer the effects of climate; Hilde et al. 2020), antecedent effect models would be most useful when 20 or more years of climatic data are available (Teller et al. 2016, van de Pol et al. 2016). Still, despite these potential issues, antecedent effect models provide a key advantage of being tools for exploratory analysis: the

automatic estimation of the timing and magnitude of climatic effects can help generate new

hypotheses.

Here, we provide an overview on antecedent effect models, we evaluate their predictive

performance, and then showcase their potential for exploratory data analysis. We systematically

evaluate our performance to predict the out-of-sample variation of the vital rates of survival,

development, and reproduction and the overall population growth rate (λ) of 77 natural

populations of 34 plant species. We compare and contrast the predictive performance of three

types of antecedent effect models to that of models based on annual climatic anomalies, and null

models. Here, we include null models to test the hypothesis that, in relatively small datasets,

monthly or annual climatic values are inadequate predictors of variation in demographic data.

We use monthly temperature and precipitation as climatic drivers, and demographic information

ranging from 7-36 years from the COMPADRE Plant Matrix Database (Salguero-Gómez *et al.*

2015). Antecedent effect models were previously employed with great success on tree rings

datasets (*e.g.*, Peltier et al. 2018). Here, we test these models on plant datasets for which the

interest in climate effects has been steadily growing (Ehrlén et al. 2016). Our analysis goes from

small, relatively common datasets to rare, long-term data that should maximize the predictive

potential of antecedent effect models. We evaluate the predictive performance of these five

models to address four objectives: (1) evaluate the predictive performance of each statistical

model; (2) compare the ability of climate to predict among vital rates and population growth rate;

(3) determine whether the predictive ability of antecedent effect models increase with temporal

replication; and (4) choose a model as a case study to show the potential of antecedent effect

models as exploratory data analysis tools.

Methods

Data and response variables

To compare the predictive performance of antecedent effect models to that of simpler

models, we used long-term time series of vital rate data obtained from matrix population models

(MPMs, hereafter) contained in the COMPADRE Plant Matrix Database (Salguero-Gómez et al.

2015). In this study, we ignored similar data on animals because it lacks taxonomic, temporal,

and spatial replication (Salguero-Gómez et al. 2016). COMPADRE contains MPMs of natural

populations where individuals have been categorized into discrete stages and their vital rates

(*e.g.*, survival, development, reproduction) tracked in discrete time steps (Caswell 2001). We

selected datasets from COMPADRE with at least seven contiguous years (*i.e.*, six MPMs) for a

single natural population. We chose seven years as the shortest study length that allowed

comparing a sufficient number of species: only half of plant demographic studies exceed four

years in duration (Salguero-Gómez et al. 2015, Römer et al. 2021). This criterion resulted in

1,057 MPMs originating from 77 populations across 34 species, with an average length of 15

years (median = 10; range = 7-36), and an average number of two populations (median = 1,

range=1-6). In our dataset, each study corresponded to a separate species or set of species, with

the exception of *Cirsium pitcheri*. This species appeared in three separate studies, conducted on

three geographically separate populations (Supplementary Online Materials Table S1). The

characteristics of our species pool reflected the well-known bias of plant demographic data for

temperate environments and herbaceous species (Supplementary Online Materials Table S2

(Salguero-Gómez et al. 2015, Compagnoni et al. 2021, Römer et al. 2021). Importantly, 86% our

populations are eudicot, 86% herbaceous perennials, and 78% of populations are from temperate

environments. We only had three shrubs, two succulent species, four species found in

Mediterranean environments, and four found in deserts.

We used these MPMs to calculate the expectation of survival, development, reproduction,

and the population growth rate (λ) associated with each annual MPM. By "development", we

refer to changes in life cycle stage. These four year-specific metrics refer to a population at its

stable stage distribution: the distribution of discrete stages that would result if each yearly

MPM's vital rates were kept constant. We did not divide these data in vital rates of non-

reproductive and reproductive plants because their year-to-year fluctuations are strongly

correlated (Tredennick et al. 2018). Accordingly, our preliminary analyses on such split vital rate

data showed inconspicuous differences between classes results, motivating us to lump data

across size classes. We calculated survival, development, and reproduction using standard

methods (Franco & Silvertown, 2004), and λ as the dominant eigenvalue of each MPM (Caswell

2001) with the R package popbio (Stubben and Milligan 2007). Here, $\lambda >$ (or $<$) 1 corresponds to

a population that is expected to grow (decline) at the stable stage distribution (Caswell, 2001).

Note that we used λ instead of a population growth rate derived from year-to-year changes in

population size which cannot be derived from matrix projection models. We calculated net

annual vital rates by taking a weighted mean of the stage-specific vital rates, where the weights

were derived from the stable stage distribution.

We used gridded and weather station climatic data to generate the climatic predictors for

our statistical models. The climatic predictors in our models include mean monthly air

temperature and total monthly precipitation at the location of each of our 77 populations. We

obtained these data primarily from the CHELSA (Karger et al. 2017), but we also used PRISM

(Daly et al. 1994) and a single Global Historical Climatology Network (GHCN) daily weather

station (Menne et al. 2012) in the temporal range not covered by CHELSA. We used CHELSA

data for all populations sampled between 1979 and 2013 (Supplementary Online Materials Table

S1). We used PRISM data for the two populations of *Astragalus scaphoides* that extend to 2014

(Tenhumberg *et al.* 2018). We used data from weather station number USC00143527 of the

GHCN for the 10 populations collected between 1938 and 1972 in Hays, Kansas, USA

(Supplementary Online Materials Table S1). We used data from this weather station instead of

PRISM because the quality of gridded climatic data tends to decrease with time since present.

We note that CHELSA and PRISM provide gridded climatic data at a 30 arcsec (~1 km^2)

resolution which approximate weather station climate averages well (Behnke et al. 2016). We

consider this spatial resolution more than sufficient, because climate anomalies are strongly

correlated spatially. For example, in North America, to observe correlations of 0.5 between the

annual anomalies of two sites, one needs to travel at least 300Km, for precipitation anomalies,

and 1,400 Km, for temperature anomalies (Compagnoni et al. 2024).

Our climatic predictors consisted of the monthly climatic anomalies observed in the 36

months preceding each vital rate observation (*i.e.*, preceding the end month of a given

demographic projection interval). Specifically, vital rate observations describe a transition from

year *t* to year *t* + 1: our models use the 36 monthly anomalies preceding year *t* + 1 as predictors.

These anomalies were *z*-scores (mean = 0, SD = 1) with respect to the long-term monthly mean.

For example, the anomalies for the month of December were calculated with respect to the long-

term mean and standard deviation of December. We calculated long-term means and standard

deviations exceeding the recommended minimum of 30 years of data (World Meteorological

Organization 2017). For CHELSA and PRISM datasets, we calculated the long-term monthly

mean using a 35-year period between 1979 and 2013, and 1980 and 2014, respectively. We

calculated the long-term monthly mean for the 10 datasets collected before 1979 using 50 years

of climate data. The difference between anomalies computed using the 50- versus 35-year mean

was negligible, as their correlations were 0.996 and 0.995 for temperature and precipitation,

respectively.

Modeling overview

Using vital rates and $\log(\lambda)$ as response variables, we compared and contrasted the

predictive performance of five models (Supplementary Online Materials: Fig. S1): the (1) Null

Model (NM), with no climatic covariates, the (2) Climate Summary Model (CSM, Dalgleish et

al., 2011), using yearly climate as a predictor, and three types of antecedent effect models: the

(3) Weighted Means Model (WMM, van de Pol & Cockburn, 2011), the (4) Stochastic

Antecedent Model (SAM, Ogle et al., 2015), and the (5) Finnish Horseshoe Model (FHM, Hastie

et al., 2009). Models 1-5 are ranked by increasing flexibility, by progressively relaxing key

assumptions (Supplementary Online Materials: Fig. S1). NMs are a "model of the mean" that do

not account for the effect of climatic drivers. CSMs assume that the monthly anomalies observed

within a specific period have all equal weight and that these anomalies affect population

dynamics in the same direction. Therefore, this model uses as a predictor the mean of 12 monthly

anomalies. WMM's weigh the contribution of the 12 monthly anomalies based on a Gaussian

function. The mean of this function will identify the month with the highest contribution, and all

other months will have progressively lower weight (Supplementary Online Materials: Fig. S1).

Like in WMMs, SAMs assume that monthly anomalies influence population dynamics in the

same direction. SAMs estimate the relative weights of monthly anomalies by assigning a

Dirichlet distribution prior to the weights (Supplementary Online Materials: Fig. S1; Ogle et al.,

2015). Finally, FHMs fit a separate coefficient for each of the 12 monthly anomalies, shrinking these toward zero through their prior distribution. In regularized horseshoe regressions, monthly anomalies can have positive or negative effects. FHMs were not developed to model climatic effects (Piironen and Vehtari 2017), though regularization approaches have been previously employed to estimate climatic effects on vital rate information (e.g. Tredennick et al., 2016).

We chose these three antecedent effect models as they were the most useful exploratory models for our dataset. There are four types of exploratory models to estimate climate sensitivity (van de Pol and Bailey 2019): sliding windows (van de Pol et al. 2016), weighted means, regularization, and machine learning (Teller et al. 2016). We discarded sliding windows and machine learning because not easily implemented in a Bayesian framework, and because they were too complex for our small datasets. We also excluded the use of splines (Teller et al. 2016), as they are a form of regularization, and we discarded three alternatives to our WMM because of convergence issues. We provide further details on these decisions in the Supplementary Online Materials.

We fit each model type using climate data from three years: one from the year of the census, one prior to the census, and one two years before the census. We used these three years of data to account for the potential of lagged climatic effects. There is convincing evidence of lagged climatic effects in several case studies, including for three populations analyzed in this study: one population of *Cryptantha flava* (Evers et al. 2021) and two of *Astragalus scaphoides* (Tenhumberg et al., 2018). We re-fit each model using climate data from three separate years: each year corresponding to a subset of the 36 monthly anomalies preceding each demographic observation. We subdivided these 36 anomalies as belonging to the year (year t), one year (t-1), and two years (t-2) preceding each demographic observation. Hence, we fit five model types for

each response variable. Because four of these models included climatic predictors, we refit these

four models on the three years of climate data.

Below, we describe the equations of the models we tested assuming a normal response

variable, $\log(\lambda)$. To facilitate model comparison and model convergence, we used weakly

informative priors for the intercept and variance terms (Lemoine 2019). We document the priors

for all of the five (Eq. 1-5) models presented below, and the prior predictive checks we ran to

choose them, in the section of the Supplementary Online Materials on prior predictive checks.

Our null is a model of the mean, such that

256
$$y_{ti} \sim Normal(\hat{y}, \sigma) \qquad (1a)$$

257
$$\hat{y} = \alpha, \qquad (1b)$$

258
$$\alpha \sim Normal(0,0.5), \qquad (1c)$$

259
$$\sigma \sim Gamma(0.01,0.01), \qquad (1d)$$

260   where y_{ti} refers to a $\log(\lambda)$ value, t refers to the year of observation, i to a replicate within years (i

261   exceeds one in datasets that are spatially replicated), $\hat{y}$ is the mean model prediction, σ is the

262   standard deviation, and α is a model intercept, the mean. In the CSM, we model the response

263   variable as a function of $\bar{x}_t$, the mean of the 12 monthly anomalies (z-scores) over year t, as

264
$$\hat{y}_t = \alpha + \beta \bar{x}_t, \qquad (2a)$$

265
$$\beta \sim Normal(0,1), \qquad (2b)$$

266   where β is the coefficient for the effect of $\bar{x}_t$, and $\hat{y}$ is the mean model prediction referred to year

267   t.

268   The first antecedent effect model is the WMM, which models the response as a function

269   of x_{tk}, the monthly temperature deviations for year t and month k:

270
$$\hat{y}_t = \alpha + \beta \sum_k w_k x_{kt} \qquad (3a)$$

271
$$w_k = g_k / \sum_k g_k \qquad (3b)$$

272
$$g_k = \frac{1}{\sigma_w \sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{k-\mu}{\sigma_w}\right)^2} \qquad (3c)$$

273
$$\mu \sim Normal(6.5,12)[0,12], \quad (3d)$$

274
$$\sigma_w \sim Normal(1,12)[1,\infty), \quad (3e)$$

where the 12 x_k monthly anomalies prior to the end of the relevant demographic year, are each

weighted by w_k. In turn, w_k is a normalized vector of Gaussian weights, g_k, with mean μ and

scale σ_w. These priors on the μ and σ_w parameters allowed consistent convergence to these

models. The prior in eq. 3d provides equal probability to μ values between 1 and 12. The prior in

Eq. 3e allows values of σ_w so large that the weights tend to be equal. The square brackets in Eq.

3d and 3e show the limits beyond which the potential values of these parameters are truncated.

While values of μ do not extend beyond 0 and 12, σ_w has no upper bound.

The SAM (Ogle et al. 2015) adds flexibility to the WMM above,

283
$$\hat{y}_t = \alpha + \beta \sum_k w_k x_{kt}, \qquad (4a)$$

284
$$\boldsymbol{w} = \text{Dirichlet}\left(\theta, \overbrace{\ldots, \theta}^{n=12}\right), \quad (4b)$$

where vector $\boldsymbol{w}$ contains 12 weights that sum to 1, and w_k is a single weight that refer to monthly

anomaly x_{kt} observed in month k and year t. We set the concentration parameter (θ) to 1,

corresponding to a uniform distribution on the 12-dimensional simplex of weights. The added

flexibility originates because the values in vector $\boldsymbol{w}$ do not follow a functional form (albeit they

are correlated). Henceforth, we refer to the term $\sum_k w_k x_{kt}$, in both Eq. 3a and 4a, as the

"antecedent" of year t.

Finally, the FHM is a multiple regression whose coefficients are shrunk via the horseshoe

prior proposed by Piironen & Vehtari (2017). These models estimate a separate slope value, β_k,

for each k month. The Finnish horseshoe prior prevents overfitting by shrinking estimates

towards zero. This model is fit as follows:

$$\hat{y}_t = \alpha + \sum_k \beta_k x_{kt}, \qquad (5a)$$

$$\beta_k \sim Normal\left(0, \tau\tilde{\theta}_k\right), \qquad (5b)$$

$$\tilde{\theta}_k = \frac{c\theta_k}{\sqrt{c^2 + \tau^2\theta_k^2}}, \qquad (5c)$$

$$\theta_k \sim HalfCauchy(0, 0.1), \qquad (5d)$$

$$c^2 \sim InvGamma\left(\frac{v}{2}, \frac{v}{2}s^2\right), \qquad (5e)$$

$$\tau \sim HalfCauchy(0, \tau_0), \qquad (5f)$$

$$\tau_0 = \frac{m_0}{M - m_0}\frac{\sigma}{\sqrt{N}}. \qquad (5g)$$

Here, 12 β_k values depend on a normal distribution whose standard deviation is controlled

by τ, the "global scale", and 12 $\tilde{\theta}_k$ "local scales". The global scale τ tends to shrink every

monthly estimate β_k toward zero. On the other hand, the $\tilde{\theta}_k$ values allow the β_k monthly estimates

to become relatively large. In the FHM, the combination of τ and $\tilde{\theta}_k$ shrinks all β_k coefficients

toward zero, while controlling the magnitude of β_k values for which there is evidence for a

substantial departure from zero. We modified these models to fit survival, development, and

reproduction data, which are non-normally distributed. We describe these modifications in the

Supplementary Online Material.

Model fitting

We fit models using a Hamiltonian Monte Carlo sampler via the *R* package *rstan* (Stan

Development Team 2018). We fit models running four parallel chains for 16,000 iterations, with

a burn-in of 4,000 iterations, and thinning by saving one out of every eight iterations. Because

we had 13 alternative models, four demographic responses ($\log(\lambda)$ and three vital rates), two

climatic predictors, and 36 datasets, we fit a total of 3,744 models. Fitting these many models

made it unfeasible to check models one at a time. Therefore, we opted to perform model checks

across model types, response variables, and dataset length. We checked for model convergence

using four metrics. First, we flagged models with more than nine divergent transitions, models

with a single parameter whose Gelman and Rubin convergence diagnostic was above 1.1, models

where at least one parameter had a ratio of the Monte Carlo standard error to the standard

deviation of the posterior above 0.1, and a single parameter whose effective sample size was

lower than 10% of the posterior sample.

In the Supplementary Online Material, we provide the methods and results to assess the

convergence of the Hamiltonian Monte Carlo samplers and assess model fit (Bayesian model

checks).

Model comparison

We compared model performance for each vital rate, climate predictor, and species

combination. We performed comparisons based on the log pointwise predictive density (LPPD)

(Bernardo and Smith 2009, Hooten and Hobbs 2015) calculated through a leave-one-year-out

(LOYO) cross-validation (e.g. Tredennick et al., 2016). A LOYO cross-validation creates a

training set by leaving out an entire year of data rather than a single data point. The model is then

fit for as many years the dataset is composed of, each time leaving a different year out of the

training set. The higher the value of LPPD of the held-out data, the better the predictive

performance of the model. Because we refit models, each time leaving out a single year of data,

we fit a total of 54,048 models. Our model selection focused on LOYO cross-validation, because

we aim to predict year-level variation in vital rates. We did not perform k-fold, holdout, or

exhaustive cross-validations because, despite being more conservative cross-validations, our

datasets were sufficiently small as to generate convergence issues. We performed this procedure

because the climate is strongly correlated among different sites in our spatially replicated

datasets. In particular, 16 of our 36 datasets are spatially replicated, and their mean correlation is

0.98 (range = 0.74-1).

We compared model performance by plotting the difference in ΔLPPD — the difference

in LPPD with the best model. The $\Delta LPPD_i$ of model i is calculated as $LPPD_i$ – max(LPPD). For

each species, we plotted heat maps of the $\Delta LPPD_i$ values for each model, flagging the best three

models. We then ordered heat maps by data length along the y-axis, and by the year of climate

data (t, t-1, and t-2) on the x-axis.

We addressed our three objectives by testing for differences in the relative support of our

models using Pseudo-Bayesian Model Averaging. We calculated model weights using Pseudo-

Bayesian Model Averaging for each combination of species, response variable, and predictor

(temperature or precipitation). We calculated model weights following equations 8 and 9 in Yao

et al. (2018). LPPD relative weights using the same formula used for AIC weights, so that

354
$$p_k = \frac{\exp\left(LPPD_k - \frac{1}{2}\mathrm{se}(LPPD_k)\right)}{\sum_{k=1}^{K}\exp\left(LPPD_k - \frac{1}{2}se(LPPD_k)\right)} \qquad (8)$$

Where p is the LPPD relative weight, k refers to each model, se is the standard error, and K is 13,

the total number of models tested for each combination of species, response variable, and

climatic predictor. We arcsine transformed these LPPD relative weights to make them normally

distributed. We then used these arcsine transformed LPPD relative weights as the response

variable to address our three objectives. We addressed these objectives using graphs and average

statistics. We plotted boxplots of model weights across all datasets, grouping them based on

model types. This evaluated (1) the predictive performance of each of the five models. We

plotted model weights of NMs to compare (2) the ability of climate to predict among vital rates

and population growth rate. We also used these results to explore the role of plant functional

type, ecoregion, and generation time in explaining our results. Then, we linked relative model

weights against the temporal replication of each dataset to (3) verify whether such replication

increased the predictive ability of antecedent effect models. We expected that replication would

increase either the predictive ability of either the worst or best models. Therefore, to test this

hypothesis we fit quantile regressions on these plots using the R package quantreg (Koenker

2022), from the 10^{th} to the 90^{th} quantile in increments of 10. Finally, used our model selection

results to (4) select a model that showcases the potential of antecedent effect models as

exploratory data analysis tools. To select the dataset of our case study, we first visually selected

models with high predictive performance. Finally, we plotted these models to choose a

particularly striking and biologically sensible example.

Sensitivity analysis

We assessed the solidity of our results by running three alternative analyses: one using only the

longest datasets, another fitting antecedent effect models on 36 months of climatic data, and an

analysis fitting models that included both precipitation and temperature. First, we re-run analyses

including only datasets that were at least 20 years long. This analysis could have increased

predictive ability, because 20 years of data are considered the minimum sample size to detect a

climatic signal (Teller et al. 2016, van de Pol et al. 2016a). Second, we fit our three antecedent

effect models using all 36 months of climatic data. Third, we re-run all analyses including year-

models using precipitation only, and models including both precipitation and temperature as

predictors (using both 12 months and 36 months of data).

Case study and exploratory analysis

We present a case study to showcase the potential of antecedent effect models as exploratory

tools. We selected models with high predictive ability because they were the only ones whose

model weights were unequal (for WMMs and SAMs), or whose effect sizes clearly differed from

zero (for FHMs). Our case study modeled reproduction in *Astragalus tyghensis*, an endemic

legume plant species with an extremely restricted range encompassing arid lands of Western

North America (Thorpe and Kaye 2008). This dataset has only nine years of data but five sites,

which must have substantially improved the chance to detect climatic signals (Compagnoni et al.

2024). Using this example, we first show how antecedent effect models identify monthly

climatic drivers of demographic responses, and we discuss how these inferences can inform

biological hypotheses. Second, we show how antecedent effect models improve model fit. We

present models fit on the reproduction of *A. tyghensis* using precipitation anomalies one year

prior to the last demographic observation (year t-1). This climatic predictor had high predictive

power in our model comparisons.

We use this case study to show for each model its fit to data and the effect sizes of monthly

climatic anomalies. To show model fit to data, we plotted each raw reproduction data point

against its corresponding prediction according to 200 randomly selected posterior samples. To

show model fit to data, we plotted each raw reproduction data point against 200 randomly

selected posterior samples predictions. To display the monthly effect sizes, we produced

estimates of effect sizes suitable for a graphic comparison among CSMs, WMMs, SAMs, and

FHMs. For the CSM, we divided the regression slope, β (Eq. 2a), by 12, the number of months.

For the WMMs and SAMs, we multiplied the regression slope, β (Eq. 3-4a), by the 12 monthly

weights, w_k (Eq. 3-4b). Finally, the FHMs directly estimate 12 β_k separate regression slopes. We

represented these monthly effect sizes by plotting the median and the 95% credible intervals of

the posterior on the y-axis, against the month number on the x-axis.

Results

Model comparison

Antecedent effect models were not generally useful in predicting the effects of climate,

but performed well in isolated cases. When comparing LPPD weights, NMs were consistently

the best performing models (Fig. 1). The variance in model weights was smaller in NMs models,

while the models including climatic predictors could reach very high or very low model weights

(Fig. 1). This variance in the LPPD weights suggests that estimating climate effects is

advantageous for a subset of populations which are presumably particularly sensitive to climate.

However, estimating climate effects can substantially decrease predictive ability in the other

populations. There were no clear trends in which antecedent effect model had the best predictive

performance, even if FHMs tended to perform better on $\log(\lambda)$ data (Fig. 1; Supplementary

Online Materials: Fig. S10-13). Still, these average differences were small, potentially reflecting

that these three models tend to provide similar information.

Models including climate had similar predictive performance across response variables,

plant functional types, and ecoregions. On average, we found little difference in the predictive

performance of response variables (Fig. 2), albeit FHMs tended to predict $\log(\lambda)$ better than other

climate models (Fig. 1; Supplementary Online Materials: Fig. S10-11). The predictive

performance was also similar across plant functional types (Supplementary Online Materials:

Fig. S14), Ecoregions (Supplementary Online Materials: Fig. S15), and plant species generation

time (Supplementary Online Materials: Fig. S16).

The temporal replication of the studies did not improve average model performance, but

it decreased the likelihood of fitting models with very low LPPD relative weights. The only

quantile regression that shows a significant trend is linked to the 10^{th} quantile, for which

temporal replication increases LPPD relative weights (Supplementary Online Materials: Fig.

S17). This result suggests that sample size only limits the negative effects of fitting models with

large numbers of parameters on predictive ability.

In our sensitivity analysis, the results of our model comparison held independently of the

data and model used. First, selecting datasets longer than 20 years did not change the main

results that null models have the greatest predictive performance, and that other climate models

are roughly equivalent (Supplementary Online Materials: Fig. S18). Moreover, in these long-

term datasets, the predictive performance of climate models was lower than that of null models

across response variables (Supplementary Online Materials: Fig. S19). These results held despite

these data included 36-month models. These 36-month models did not improve the predictive

ability of climatic models (Supplementary Online Materials: Fig. S20-21). Finally, comparing

precipitation-only models with models including precipitation plus temperature indicated that

precipitation-only models tended to have better predictive ability (Supplementary Online

Materials: Fig. S22-26).

Case study and exploratory analysis

The results of antecedent effect models provided biologically interpretable information that can

be used to formulate hypotheses on the demography of *A. tyghensis*. All four models show that

precipitation is negatively associated with reproduction (Fig. 3). According to the WMM, the

monthly anomalies preceding a demographic census tend to be the most important in driving

reproduction. The SAM and FHM provide more precise insights, indicating that the most

important climate windows are located two (May) and six (January) months prior to the

demographic census. Thus, antecedent effect models indicate that the fecundity of this species is

negatively correlated to the precipitation of the growing and dormant seasons. Therefore, in this

case antecedent effect models provided information with theoretical and practical implications,

even if their predictive ability was similar to the CSM (Fig. S10, Fig. 3).

Discussion

Predicting and understanding how climate affects the vital rates (survival, development,

reproduction) of natural populations and their overall performance (*e.g.*, population growth rate,

λ) is a core ecological question. By comparing the predictive performance of antecedent effect

models to simpler linear models (NMs and CSMs), we found that (1) models using climate as

predictor did not reliably increase our ability to predict any of the vital rates and population

performance, (2) dataset length has negligible effects on predictive power, and (3) despite having

similar predictive ability to CSMs, antecedent effect models provide biologically sensible

insights on the timing of climatic effects. First, the hypothesize that the weak predictive ability of

climate models, which is irrespective of dataset length, originates from either small climatic

signals, and from low signal-to-noise ratio. A potential solution to deal with the low signal-to-

noise ratio is to use individual-level data with abundant co-variate information. Second, given

the generally poor predictive ability of our climate models, we conclude that on herbaceous

demographic data, antecedent effect models are best used as exploratory tools for hypothesis

generation. For this exploratory aim, we recommend the Finnish Horseshoe Model (FHM),

because FHM has the largest flexibility.

Our initial expectation was that antecedent effect models would improve predictions,

because they accommodate a variety of potential links between demography and climate

(Kreyling 2010, Evers et al. 2021). This expectation was strong at least for our 12 datasets that

exceeded 20 years of data, as this was the dataset length recommended for simulation studies

focused on antecedent effect models (Teller et al. 2016, van de Pol et al. 2016). The predictive

success of null models is a clear refutation of our initial expectation. We expected dataset length

would positively affect the predictive ability of climatic variables by increasing total sample size,

and by increasing the range of climatic conditions observed during the study (Compagnoni et al.

2024). For the fact that climate had small predictive power even in datasets exceeding 30 years

we propose three possible explanations. First, the true effect of climate could be much smaller

than what was used in previous simulation studies. For example, Teller et al. (2016) defined the

climate signals they simulated as "strong". However, subsequent analysis on the type of data

mimicked by their simulations found weak climatic signals with inconsistent effects on

predictive power (Tredennick et al. 2016, 2021).

Second, the signal-to-noise ratio might be quite low in aggregated data such as what we

employed for this study. For example, the influential article by van de Pol et al. (2016) simulated

a signal strength based on R^2, choosing 0.2 as their lowest R^2. Our experience with a large dataset

of herbaceous plant population growth rates (Compagnoni et al. 2021) indicated a mean in-

sample R^2-squared of about 0.24. If such low signal-to-noise ratios were the norm, then

accounting for confounding factors would be critical to understand and predict the effects of

climate. To account for confounding factors, demographic data on individuals could facilitate

accounting for co-varying factors (e.g. competition and spatial location Chu et al. 2016). Third,

the way in which climate affects plants might be too complex to capture via phenomenological

models applied on aggregated population data. In principle, temperature and precipitation should

correlate with the main factors affecting plant processes: solar radiation, leaf temperature, soil

moisture, and vapor pressure deficit (Lambers et al. 2008). Accordingly, temperature and

precipitation are excellent predictors in optimal scenarios, such as tree ring studies (Fritts 2012).

In these studies, much larger sample sizes belong to individuals selected to be from range

positions and microsite conditions that maximize climate sensitivity (Klesse et al. 2018). On the

other hand, our population dynamics data samples entire populations located at any position

within a species range. Thus, our data comprises individuals and populations of varying climate

sensitivity. With these type of data, unveiling climatic correlations might require sophisticated

models; for example, models that approximate the microclimatic conditions at each site (e.g.

Kearney and Leigh 2024).

The case study presented in this article showcases the potential of antecedent effect

models for exploratory analysis and hypothesis generation. While antecedent effect models have

on average similar predictive performance to CSMs, they provide granular information that is

more useful scientifically. In our *A. tyghensis* case study, both the SAM and FHM showed a

strong negative association between reproduction and the precipitation anomalies that occurred

in the May and January of the preceding year. These results raise three hypotheses. First, such

negative association with precipitation anomalies occurring a year prior to a demographic

observation could result from indirect effects mediated through competition (Suttle et al. 2007,

Adler et al. 2012) or physiological tradeoffs (*e.g.*, costs of reproduction, Crone et al., 2009;

Miller et al., 2012). Second, the correlation with the precipitation in January, a month when the

mean temperature is just above or below 0° C, could reflect the effect of snowfall. This effect is

often distinct from rain (Dalgleish et al. 2011, Compagnoni and Adler 2014) and could arise in

different ways: for example, because snow affects pathogens (Smull et al. 2019), it protects

plants from frost (Inouye, 2000), or because snowmelt provides a large pulse of resources (Noy-

Meir 1973). Third, the large effect of May precipitation could reflect a key developmental phase

occurring either in May or June. All three hypotheses call for further tests to be performed via

analysis of existing data or field experiments. For exploratory studies, we recommend to exploit

the flexibility of FHMs. FHMs usually provide qualitatively similar results to the SAMs, but

with the added benefit of simultaneously identifying positive and negative climatic signals.

While it is unlikely for climate signals to go in opposite directions, we did observe such cases in

some of our model fits. It is perhaps because of this flexibility that FHMs outperform WMMs

and SAMs when predicting $\log(\lambda)$ (Figure 1).

Our experience building and comparing antecedent effect models informs our suggestions

on dataset length, hyperparameter priors, and emphasizes the role of prior predictive checks.

While most of our models had no problems converging, problems occasionally arose when

datasets had less than about 20 years of data (Fig. S5-6). Therefore, in our case the often

recommended dataset length of 20 years (Teller et al. 2016, van de Pol and Bailey 2019)

improved model convergence rather than model predictive ability. The choice of hyperparameter

priors has strong effects on model fitting and performance. In weighted mean models (WMMs),

truncated hyperparameter priors (Eq. 3d-e) prevented the sampler from exploring unrealistic

parameter values, greatly facilitating model convergence. In SAMs, we chose a vector of ones as

the Dirichlet concentration parameter (Eq. 4b) to not favor any particular lag, and to ensure that

prior weights had a uniform distribution. A much lower number than 1 would favor models with

a single non-zero weight. In FHMs, we suggest using a standard deviation of 0.1 for the prior on

monthly effect sizes (Eq. 5d). Using a standard deviation of 1 (*e.g.*, Piironen & Vehtari, 2017)

generated unrealistically high variation in our prior predictive checks (Supplementary Online

Materials: prior predictive checks). More generally, prior predictive checks were essential

because our initial prior choices produced very different variance in predictions among model

types and response variables. Moreover, priors generating unrealistically high variance had

model convergence issues and lower predictive performance.

These results justify the continued data and conceptual efforts to understand and predict

the effects of climatic drivers on the vital rates of herbaceous plants. These effects of climate are

complex, and antecedent effect models could contribute to understand this complexity as a

hypothesis generation tool. However, improvements in predictive ability will rely on datasets

with larger replication, especially temporal replication. Moreover, conceptual breakthroughs

such as those regarding the importance of indirect effects (Clark et al., 2021; Hacket☐Pain et al.,

2018), might also prove useful.

Data availability statement. The data and reproducible code for the analyses connected to this study are available at https://zenodo.org/records/7839199.

Competing Interests Statement. We declare no competing interests.

Acknowledgements. AC and TMK were supported by the Alexander von Humboldt Foundation under the framework of the Humboldt Professorship, and by the Helmholtz Recruitment Initiative of the Helmholtz Association (both to TMK); RSG was supported by NERC (NE/M018458/1 and NE/X103766/1). This work emanated from research discussions during the sAPROPOS (Analysis of PROjections of POpulationS) working group led by RSG and TMK at the German Centre for Integrative Biodiveristy Research (iDiv), which is funded by the German Research Foundation (FZT-118-202548816). We thank the Max Planck Institute for Demographic Research for the support of the COMPADRE Plant Matrix Database and the hundreds of plant population ecologists who have contributed with demographic data to COMPADRE. We thank B. Teller and P. Barks for discussions in the early phases of this project.

Author Contributions. **Aldo Compagnoni**: Conceptualization (Equal); Data curation (Equal); Formal analysis (Lead); Methodology (Lead); Software (Lead); Validation (Lead); Visualization (Lead); Writing – original draft (Lead); Writing – review & editing (Equal). **Dylan Z. Childs**: Conceptualization (Supporting); Data curation (Equal); Formal analysis (Supporting); Methodology (Supporting); Project administration (Supporting); Software (Supporting); Supervision (Supporting); Validation (Supporting); Writing – review & editing (Equal). **Tiffany M Knight**: Data curation (Equal); Funding acquisition (Lead); Resources (Supporting); Supervision (Supporting); Writing – review & editing (Equal). **Roberto Salguero-Gómez**: Conceptualization (Equal); Data curation (Equal); Formal analysis (Supporting); Funding

582   acquisition (Supporting); Methodology (Supporting); Project administration (Lead); Supervision

583   (Lead); Writing – review & editing (Equal).

References

Adler, P. B., H. J. Dalgleish, and S. P. Ellner. 2012. Forecasting plant community impacts of climate

variability and change: when do competitive interactions matter? Journal of Ecology 100:478–

487.

Angilletta, M. J. J. 2009. Thermal Adaptation: A Theoretical and Empirical Synthesis. OUP Oxford.

Behnke, R., S. Vavrus, A. Allstadt, T. Albright, W. E. Thogmartin, and V. C. Radeloff. 2016. Evaluation of

downscaled, gridded climate data for the conterminous United States. Ecological Applications

26:1338–1351.

Bernardo, J. M., and A. F. M. Smith. 2009. Bayesian Theory. John Wiley & Sons.

Caswell, H. 2001. Matrix population models. Construction, analysis, and interpretation. Sinauer

Associates, Sunderland, MA.

Catchpole, E. A., B. J. T. Morgan, T. N. Coulson, S. N. Freeman, and S. D. Albon. 2000. Factors influencing

Soay sheep survival. Journal of the Royal Statistical Society: Series C (Applied Statistics) 49:453–

472.

Chu, C., A. R. Kleinhesselink, K. M. Havstad, M. P. McClaran, D. P. Peters, L. T. Vermeire, H. Wei, and P. B.

Adler. 2016. Direct effects dominate responses to climate perturbations in grassland plant

communities. Nature Communications 7:11766.

Clark, J. S., R. Andrus, M. Aubry-Kientz, Y. Bergeron, M. Bogdziewicz, D. C. Bragg, D. Brockway, N. L.

Cleavitt, S. Cohen, B. Courbaud, R. Daley, A. J. Das, M. Dietze, T. J. Fahey, I. Fer, J. F. Franklin, C.

A. Gehring, G. S. Gilbert, C. H. Greenberg, Q. Guo, J. HilleRisLambers, I. Ibanez, J. Johnstone, C. L.

Kilner, J. Knops, W. D. Koenig, G. Kunstler, J. M. LaMontagne, K. L. Legg, J. Luongo, J. A. Lutz, D.

Macias, E. J. B. McIntire, Y. Messaoud, C. M. Moore, E. Moran, J. A. Myers, O. B. Myers, C.

Nunez, R. Parmenter, S. Pearse, S. Pearson, R. Poulton-Kamakura, E. Ready, M. D. Redmond, C.

D. Reid, K. C. Rodman, C. L. Scher, W. H. Schlesinger, A. M. Schwantes, E. Shanahan, S. Sharma,

M. A. Steele, N. L. Stephenson, S. Sutton, J. J. Swenson, M. Swift, T. T. Veblen, A. V. Whipple, T.

G. Whitham, A. P. Wion, K. Zhu, and R. Zlotin. 2021. Continent-wide tree fecundity driven by

indirect climate effects. Nature Communications 12:1242.

Clark, J. S., L. Iverson, C. W. Woodall, C. D. Allen, D. M. Bell, D. C. Bragg, A. W. D'Amato, F. W. Davis, M.

H. Hersh, I. Ibanez, S. T. Jackson, S. Matthews, N. Pederson, M. Peters, M. W. Schwartz, K. M.

Waring, and N. E. Zimmermann. 2016. The impacts of increasing drought on forest dynamics,

structure, and biodiversity in the United States. Global Change Biology 22:2329–2352.

Compagnoni, A., and P. B. Adler. 2014. Warming, soil moisture, and loss of snow increase Bromus

tectorum's population growth rate. Elementa: Science of the Anthropocene 2:000020.

Compagnoni, A., S. Evers, and T. Knight. 2024. Spatial replication can best advance our understanding of

population responses to climate. Ecography 2024:e06833.

Compagnoni, A., S. Levin, D. Z. Childs, S. Harpole, M. Paniw, G. Römer, J. H. Burns, J. Che-Castaldo, N.

Rüger, G. Kunstler, J. M. Bennett, C. R. Archer, O. R. Jones, R. Salguero-Gómez, and T. M. Knight.

2021. Herbaceous perennial plants with short generation time have stronger responses to

climate anomalies than those with longer generation time. Nature Communications 12:1824.

Crone, E. E., E. Miller, and A. Sala. 2009. How do plants know when other plants are flowering? Resource

depletion, pollen limitation and mast-seeding in a perennial wildflower. Ecology Letters

12:1119–1126.

Czachura, K., and T. E. X. Miller. 2020. Demographic back-casting reveals that subtle dimensions of

climate change have strong effects on population viability. Journal of Ecology 108:2557–2570.

Dalgleish, H. J., D. N. Koons, M. B. Hooten, C. A. Moffet, and P. B. Adler. 2011. Climate influences the

demography of three dominant sagebrush steppe plants. Ecology 92:75–85.

Daly, C., R. P. Neilson, and D. L. Phillips. 1994. A Statistical-Topographic Model for Mapping

Climatological Precipitation over Mountainous Terrain. Journal of Applied Meteorology 33:140–

158.

Ehrlén, J., W. F. Morris, T. von Euler, and J. P. Dahlgren. 2016. Advancing environmentally explicit

structured population models of plants. Journal of Ecology 104:292–305.

Ellner, S. P., D. Z. Childs, and M. Rees. 2016. Data-driven Modelling of Structured Populations: A Practical

Guide to the Integral Projection Model. Springer International Publishing.

Elton, C. S. 1924. Periodic Fluctuations in the Numbers of Animals: Their Causes and Effects. Journal of

Experimental Biology 2:119–163.

Evers, S. M., T. M. Knight, D. W. Inouye, T. E. X. Miller, R. Salguero-Gómez, A. M. Iler, and A.

Compagnoni. 2021. Lagged and dormant season climate better predict plant vital rates than

climate during the growing season. Global Change Biology 27:1927–1941.

Franco, M., and J. Silvertown. 2004. A Comparative Demography of Plants Based upon Elasticities of Vital

Rates. Ecology 85:531–538.

Fritts, H. 2012. Tree Rings and Climate. Elsevier.

Hacket-Pain, A. J., D. Ascoli, G. Vacchiano, F. Biondi, L. Cavin, M. Conedera, I. Drobyshev, I. D. Liñán, A. D.

Friend, M. Grabner, C. Hartl, J. Kreyling, F. Lebourgeois, T. Levanič, A. Menzel, E. van der

Maaten, M. van der Maaten-Theunissen, L. Muffler, R. Motta, C.-C. Roibu, I. Popa, T.

Scharnweber, R. Weigel, M. Wilmking, and C. S. Zang. 2018. Climatically controlled reproduction

drives interannual growth variability in a temperate tree species. Ecology Letters 21:1833–1844.

Harper, J. L. 1967. A Darwinian Approach to Plant Ecology. Journal of Ecology 55:247–270.

Hastie, T., R. Tibshirani, and J. Friedman. 2009. The Elements of Statistical Learning: Data Mining,

Inference, and Prediction, Second Edition. Springer Science & Business Media.

Hilde, C. H., M. Gamelon, B.-E. Sæther, J.-M. Gaillard, N. G. Yoccoz, and C. Pélabon. 2020. The

Demographic Buffering Hypothesis: Evidence and Challenges. Trends in Ecology & Evolution

35:523–538.

Hoerl, A. E., and R. W. Kennard. 1970. Ridge Regression: Biased Estimation for Nonorthogonal Problems.

Technometrics 12:55–67.

Hooten, M. B., and N. T. Hobbs. 2015. A guide to Bayesian model selection for ecologists. Ecological

Monographs 85:3–28.

Inouye, D. w. 2000. The ecological and evolutionary significance of frost in the context of climate

change. Ecology Letters 3:457–463.

Karger, D. N., O. Conrad, J. Böhner, T. Kawohl, H. Kreft, R. W. Soria-Auza, N. E. Zimmermann, H. P.

Linder, and M. Kessler. 2017. Climatologies at high resolution for the earth's land surface areas.

Scientific Data 4:170122.

Karger, D. N., and N. E. Zimmermann. 2018. CHELSAcruts – High resolution temperature and

precipitation timeseries for the 20th century and beyond.

http://dx.doi.org/10.16904/envidat.159.

Kearney, M. R., and A. Leigh. 2024. Fast, accurate and accessible calculations of leaf temperature and its

physiological consequences with NicheMapR. Methods in Ecology and Evolution n/a.

Klesse, S., R. J. DeRose, C. H. Guiterman, A. M. Lynch, C. D. O'Connor, J. D. Shaw, and M. E. K. Evans.

2018. Sampling bias overestimates climate change impacts on forest growth in the

southwestern United States. Nature Communications 9:5336.

Knape, J., and P. de Valpine. 2011. Effects of weather and climate on the dynamics of animal population

time series. Proceedings of the Royal Society B: Biological Sciences 278:985–992.

Koenker, R. 2022, May 2. quantreg: Quantile Regression.

Kreyling, J. 2010. Winter climate change: a critical factor for temperate vegetation performance. Ecology

91:1939–1948.

Lambers, H., F. S. C. III, and T. L. Pons. 2008. Plant Physiological Ecology. Springer Science & Business

Media.

Lemoine, N. P. 2019. Moving beyond noninformative priors: why and how to choose weakly informative

priors in Bayesian analyses. Oikos 128:912–928.

Levin, S. C., S. Evers, T. Potter, M. P. Guerrero, D. Z. Childs, A. Compagnoni, T. M. Knight, and R.

Salguero-Gómez. 2022. Rpadrino: An R package to access and use PADRINO , an open access

database of Integral Projection Models. Methods in Ecology and Evolution:2041–210X.13910.

Menne, M. J., I. Durre, R. S. Vose, B. E. Gleason, and T. G. Houston. 2012. An Overview of the Global

Historical Climatology Network-Daily Database. Journal of Atmospheric and Oceanic Technology

29:897–910.

Miller, T. E. X., J. L. Williams, E. Jongejans, R. Brys, and H. Jacquemyn. 2012. Evolutionary demography of

iteroparous plants: incorporating non-lethal costs of reproduction into integral projection

models. Proceedings of the Royal Society B: Biological Sciences 279:2831–2840.

Noy-Meir, I. 1973. Desert Ecosystems: Environment and Producers. Annual Review of Ecology and

Systematics:25–51.

Ogle, K., J. J. Barber, G. A. Barron-Gafford, L. P. Bentley, J. M. Young, T. E. Huxman, M. E. Loik, and D. T.

Tissue. 2015. Quantifying ecological memory in plant and ecosystem processes. Ecology Letters

18:221–235.

Peltier, D. M. P., J. J. Barber, and K. Ogle. 2018. Quantifying antecedent climatic drivers of tree growth in

the Southwestern US. Journal of Ecology 106:613–624.

Piironen, J., and A. Vehtari. 2017. Sparsity information and regularization in the horseshoe and other

shrinkage priors. Electronic Journal of Statistics 11:5018–5051.

van de Pol, M., and L. Bailey. 2019. Quantifying the climatic sensitivity of individuals, populations and

species. Page Effects of Climate Change on Birds. Oxford University Press.

van de Pol, M., L. D. Bailey, N. McLean, L. Rijsdijk, C. R. Lawson, and L. Brouwer. 2016a. Identifying the

best climatic predictors in ecology and evolution. Methods in Ecology and Evolution 7:1246–

1257.

van de Pol, M., L. D. Bailey, N. McLean, L. Rijsdijk, C. R. Lawson, and L. Brouwer. 2016b. Identifying the

best climatic predictors in ecology and evolution. Methods in Ecology and Evolution 7:1246–

1257.

van de Pol, M., and A. Cockburn. 2011. Identifying the Critical Climatic Time Window That Affects Trait

Expression. The American Naturalist 177:698–707.

Römer, G., J. P. Dahlgren, R. Salguero-Gómez, I. M. Stott, and O. R. Jones. 2021, April 26. Plant

demographic knowledge is biased towards short-term studies of temperate-region herbaceous

perennials. bioRxiv.

Salguero-Gómez, R., O. R. Jones, C. R. Archer, C. Bein, H. de Buhr, C. Farack, F. Gottschalk, A. Hartmann,

A. Henning, G. Hoppe, G. Römer, T. Ruoff, V. Sommer, J. Wille, J. Voigt, S. Zeh, D. Vieregg, Y. M.

Buckley, J. Che-Castaldo, D. Hodgson, A. Scheuerlein, H. Caswell, and J. W. Vaupel. 2016.

COMADRE: a global data base of animal demography. Journal of Animal Ecology 85:371–384.

Salguero-Gómez, R., O. R. Jones, C. R. Archer, Y. M. Buckley, J. Che-Castaldo, H. Caswell, D. Hodgson, A.

Scheuerlein, D. A. Conde, E. Brinks, H. de Buhr, C. Farack, F. Gottschalk, A. Hartmann, A.

Henning, G. Hoppe, G. Römer, J. Runge, T. Ruoff, J. Wille, S. Zeh, R. Davison, D. Vieregg, A.

Baudisch, R. Altwegg, F. Colchero, M. Dong, H. de Kroon, J.-D. Lebreton, C. J. E. Metcalf, M. M.

Neel, I. M. Parker, T. Takada, T. Valverde, L. A. Vélez-Espino, G. M. Wardle, M. Franco, and J. W.

Vaupel. 2015. The COMPADRE Plant Matrix Database: an open online repository for plant

demography. Journal of Ecology 103:202–218.

Schulze, E.-D., E. Beck, N. Buchmann, S. Clemens, K. Müller-Hohenstein, and M. Scherer-Lorenzen. 2019.

Dynamic Global Vegetation Models. Pages 843–863 *in* E.-D. Schulze, E. Beck, N. Buchmann, S.

Clemens, K. Müller-Hohenstein, and M. Scherer-Lorenzen, editors. Plant Ecology. Springer,

Berlin, Heidelberg.

Simmonds, E. G., E. F. Cole, and B. C. Sheldon. 2019. Cue identification in phenology: A case study of the

predictive performance of current statistical tools. Journal of Animal Ecology 88:1428–1440.

Smull, D. M., N. Pendleton, A. R. Kleinhesselink, and P. B. Adler. 2019. Climate change, snow mold and

the Bromus tectorum invasion: mixed evidence for release from cold weather pathogens. AoB

PLANTS 11:plz043.

Stubben, C., and B. Milligan. 2007. Estimating and Analyzing Demographic Models Using the popbio

Package in R. Journal of Statistical Software 22:1–23.

Sutherland, W. J., R. P. Freckleton, H. C. J. Godfray, S. R. Beissinger, T. Benton, D. D. Cameron, Y. Carmel,

D. A. Coomes, T. Coulson, M. C. Emmerson, R. S. Hails, G. C. Hays, D. J. Hodgson, M. J. Hutchings,

D. Johnson, J. P. G. Jones, M. J. Keeling, H. Kokko, W. E. Kunin, X. Lambin, O. T. Lewis, Y. Malhi, N.

Mieszkowska, E. J. Milner-Gulland, K. Norris, A. B. Phillimore, D. W. Purves, J. M. Reid, D. C.

Reuman, K. Thompson, J. M. J. Travis, L. A. Turnbull, D. A. Wardle, and T. Wiegand. 2013.

Identification of 100 fundamental ecological questions. Journal of Ecology 101:58–67.

Suttle, K. B., M. A. Thomsen, and M. E. Power. 2007. Species Interactions Reverse Grassland Responses

to Changing Climate. Science 315:640–642.

Teller, B. J., P. B. Adler, C. B. Edwards, G. Hooker, and S. P. Ellner. 2016. Linking demography with

drivers: climate and competition. Methods in Ecology and Evolution 7:171–183.

Tenhumberg, B., E. E. Crone, S. Ramula, and A. J. Tyre. 2018. Time-lagged effects of weather on plant

demography: drought and Astragalus scaphoides. Ecology 99:915–925.

Thorpe, A. S., and T. N. Kaye. 2008. Astragalus tyghensis: Actual vs. predicted population sizes. Page 18.

748   Institute for Applied Ecology for Prineville District BLM.

749   Tredennick, A. T., G. Hooker, S. P. Ellner, and P. B. Adler. 2021. A practical guide to selecting models for

750   exploration, inference, and prediction in ecology. Ecology 102.

751   Tredennick, A. T., M. B. Hooten, and P. B. Adler. 2016. Do we need demographic data to forecast plant

752   population dynamics? Methods in Ecology and Evolution 8:541–551.

753   White, M. A., S. W. Running, and P. E. Thornton. 1999. The impact of growing-season length variability

754   on carbon assimilation and evapotranspiration over 88 years in the eastern US deciduous forest.

755   International Journal of Biometeorology 42:139–145.

756   World Meteorological Organization. 2017. WMO Guidelines on the Calculation of Climate Normals.

757   World Meteorological Organization. Geneva.

758   Yao, Y., A. Vehtari, D. Simpson, and A. Gelman. 2018. Using Stacking to Average Bayesian Predictive

759   Distributions (with Discussion). Bayesian Analysis 13.

 Figure legends

Figure 1. Null models (NMs) perform better than models using climate as predictor. Box and

whisker plots and point clouds showing the arcsine transformed model weights for our 34

species, and across two climatic predictors, five model types, and four response variables. Panels

separate models fit using precipitation or air temperature as predictor variables. The middle line

of the boxplots shows the median, the upper and lower hinges delimit the first and third quartile,

and the whiskers extend 1.5 times beyond the first and third quartile. Each point refers to one of

the 3,744 model fits, and colors refer to the response variable of each model.

Figure 2. Climate has a similar predictive power across response variables. Box and whisker

plots and point clouds showing the arcsine transformed model weights of the null models by

climate variable, and based on the type of response variable. In this plot, lower values reflect a

higher predictive ability of the models including a climatic predictor. The middle line of the

boxplots shows the median, the upper and lower hinges delimit the first and third quartile, and

the whiskers extend 1.5 times beyond the first and third quartile. Points represent 288 model fits:

one for each species, response variable, and predictor variable (Null models were only one of the

13 models we fit on data).

Figure 3. Antecedent effect models yield biologically meaningful insights. Case study showing

the fits of the climate summary model (CSM), and the three antecedent effect models (WMM,

SAM, FHM) to the reproduction of a natural population of the herbaceous perennial *A. tyghensis*,

using precipitation anomalies two years preceding the demographic census. In each plot,

columns represent, respectively, model fit and climate effect sizes. The rows refer to model type:

respectively the Climate Summary Model (CSM), Weighted Mean Model (WMM), Stochastic

Antecedent Model (SAM), and Finnish Horseshoe Model (FHM). In the first column, we show

model fits using bivariate plots where the x-axis represents the observed reproductive values, and

the y-axis represents, for each observed value, 200 randomly selected predicted values from the

Bayesian posterior. In the second column, we show the effect sizes of each monthly climatic

anomaly. Each monthly effect size is shown as a point range where the black circles represent

the median, and the lines extend from the 2.5th to the 97.5th quantile of each posterior. In the

CSM, this effect sizes correspond to the regression slope, β (Eq. 2a), divided by 12. In the WMM

and SAM, the effect sizes are produced multiplying the regression slope, β (Eq. 3-4a) by the

monthly weights, w_k (Eq. 3-4b). Finally, the FHM estimates a separate slope, β_k, for each

monthly anomaly.

Figure 1.

797   **Figure 2.**

Figure 3.