The effects of climate drivers are more predictable for development than for other plant population vital rates

Aldo Compagnoni[1,2,3.], Dylan Childs[4], Tiffany M. Knight[1,2,3], Roberto Salguero-Gómez[6,7,8]

[1] Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale), Germany.

Deutscher Platz 5e, Leipzig, 04103, Germany

[2] Department of Community Ecology, Helmholtz Centre for Environmental Research–UFZ, 06120 Halle (Saale), Germany.

[3] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Puschstrasse 4, Leipzig 04103, Germany.

[4] Department of Animal and Plant Sciences, University of Sheffield. Western Bank, Sheffield S10 2TN, UK.

[6] Department of Zoology, University of Oxford. 11a Mansfield Road, Oxford, OX1 3SZ, United Kingdom.

[7] Max Planck Institute for Demographic Research, Konrad Zuse Strasse 1, Rostock, DE-18057, Germany

[8] Centre of Excellence in Environmental Decisions. The University of Queensland, St Lucia QLD 4071, Australia

Number of tables: 0

Number of figures: 5

Word count: 8,090

Number of references: 54

Online appendices: 1

Abstract

1. Understanding mechanisms and predicting trends in natural population responses to climate is a key goal of ecology. It is unclear whether the ability to predict the effects of climate on populations changes according to the response variable (e.g. survival, reproduction). Comparative studies that explicitly link climate to population dynamics remain limited, probably because adequate modeling tools are uncommon. Antecedent effect models are statistical tools capitalizing on the evidence provided by climate and population data to select time windows correlated with a population level response.

2. We compare the predictive performance of antecedent effect models against simpler models. We fit three antecedent effect models: (1) weighted mean models (WMM), which weigh the importance of monthly anomalies based on a Gaussian curve, (2) stochastic antecedent models (SAM), which weigh the importance of monthly anomalies using a Dirichlet process, and (3) regularized regressions using the Finnish Horseshoe prior (FHM), which estimate a separate effect size for each monthly anomaly. We compare these approaches to a linear model using a yearly climatic predictor and a null model with no predictors. We use demographic data from 77 natural populations of 3 plant species ranging between 7 and 36 years of length. We fit models to the overall asymptotic population growth rate (λ) and its underlying vital rates: survival, development, and reproduction.

3. Antecedent effect models do not consistently outperform the linear or null models. Average differences in performance are small. Surprisingly, ranked in order of decreasing predictive performance, the best performing approaches are: null model > linear model > FHM > SAM > WMM. Models that include climate as a predictor have a higher ability to predict

development than survival and reproduction and a higher ability to predict vital rates than

population growth rate.

4. *Synthesis*: We find that development is more predictable than other vital rates, and especially

population growth rates. Improving the predictive ability of survival and reproduction could

be inherently more difficult, and might require larger datasets or conceptual advances.

Conceptual advances could be facilitated by antecedent effect models which, despite not

improving predictive ability, are well suited as exploratory tools.

Keywords: Bayesian moving window, COMPADRE Plant Matrix Database, cross-validation,

exploratory data analysis, forecasting, population growth rate, regularization, vital rate.

Introduction

Understanding and predicting population dynamics is a key objective of ecology. In population ecology, the last decades have witnessed the development and application of many statistical methods aimed precisely at this objective (Caswell, 2001; Ellner et al., 2016). Increasingly, emphasis is being devoted to understanding how climatic drivers such as temperature and precipitation affect demographic rates (henceforth "vital rates") such as survival, development (*i.e.,* changes in stage along the life cycle), and reproduction (Ehrlén et al. 2016; Teller et al. 2016). The ultimate goal of this ambitious research agenda is to achieve a mechanistic understanding of what makes species vulnerable to climate change worldwide. Progress in this understanding requires synthetic studies that link vital rates to climatic drivers using standardized methods (Compagnoni et al., 2021).

To understand and predict the effects of climate drivers on populations, ecologists should understand how predictable different response variables are. Despite the interest of ecologists in linking population growth rates to climatic drivers, this link often results in poor predictive ability (Knape & de Valpine, 2011). Poor predictive performance might reflect that variation in population growth rates is buffered when compared to variation in the underlying vital rates (Hilde et al., 2020). One the other hand, we have a better understanding and ability to predict the effects of climate on vital rates (e.g. Clark et al., 2016, 2021; Schulze et al., 2019). Despite these developments, we do not yet know our relative ability to predict population growth rates and each separate vital rate across many species. Establishing this predictive ability in synthesis study requires addressing the technical hurdles associated with linking climatic drivers to population dynamics.

Studies that explicitly link population responses to climate drivers typically have limited

temporal extent, and thus often identify climatic predictors, usually at the annual scale, *a priori*.

Most population studies are short-term (Salguero-Gómez et al., 2015, 2016), and these low sample

sizes have historically pushed investigators to consider annual or seasonal climatic effects

(reviewed in Evers et al., 2021). This choice has practical reasons: first, monthly or annual climate

data are widely available (Daly et al., 1994; Karger & Zimmermann, 2018). Second, annual

anomalies are a logical predictor of annual demographic variation, and seasonal anomalies can be

justified by physiological or behavioural considerations (Catchpole et al., 2000; White et al.,

1999). However, there is no consensus on which season affects vital rates the most. *A priori*

expectations based on physiology, such as the importance of the growing season in plants, or the

importance of exceeding minimum thermal limits in insects (Angilletta, 2009), are often met with

opposing evidence (Czachura & Miller, 2020; Kreyling, 2010), or evidence of lagged effects from

previous years (Evers et al., 2021; Hacket-Pain et al., 2018; Tenhumberg et al., 2018).

Ecologists need to simultaneously identify the critical time window and magnitude of

climatic effects that drive their system dynamics. To address this need, several models have been

proposed to automatically select the timing most relevant to population processes (Ogle et al.,

2015; Simmonds et al., 2019; van de Pol & Cockburn, 2011). Henceforth, we refer to these models

as "antecedent effect models". These models control for multiple comparisons, and weigh the time

windows when drivers are most influential for population dynamics, dividing the climate (or any

other relevant factor) of a time window (*e.g.*, a year) into sub-periods of equal lengths – usually

months. Such weighing can occur by estimating the importance of each monthly anomaly using

weights, which can be constrained to follow a specific function (van de Pol & Cockburn, 2011) or

not (Ogle et al., 2015). Alternatively, the weighing can be performed by estimating a different

effect size for each monthly climate anomaly, using regularization or informative priors to

minimize overfitting (Hoerl & Kennard, 1970). These models have potential as both exploratory

and predictive tools. The automatic estimation of the timing and magnitude of climatic effects can

help generate hypotheses. Moreover, these models could increase predictive ability by reflecting

the temporal processes that link climatic drivers to population dynamics.

Here, we systematically evaluate our performance to predict the out-of-sample variation of

the vital rates of survival, development, and reproduction and the overall population growth rate

(λ) of 77 natural populations of 34 plant species. We compare and contrast the predictive

performance of three types of antecedent effect models to that of null models and models based on

annual climatic anomalies. We use monthly temperature and precipitation as climatic drivers, and

demographic information ranging from 7-36 years from the COMPADRE Plant Matrix Database

(Salguero-Gómez *et al.* 2015). We evaluate the predictive performance of these five models to

address three objectives: (1) evaluate the predictive performance of each different statistical

model; (2) determine whether the predictive ability of antecedent effect models increase with

temporal replication; and (3) compare the predictive ability among vital rates and population

growth rate.

Methods

Methods

Data and response variables

To compare the predictive performance of antecedent effect models to that of simpler models, we

used long-term time series of vital rate data obtained from matrix population models (MPMs,

hereafter) contained in the COMPADRE Plant Matrix Database (Salguero-Gómez et al., 2015).

COMPADRE contains MPMs of natural populations where individuals have been categorized into

discrete stages and their vital rates (*e.g.*, survival, development, reproduction) tracked in discrete

time steps (Caswell, 2001). We selected datasets from COMPADRE with at least seven contiguous

years (*i.e.*, six MPMs) for a single natural population. We chose seven years as the shortest study

length that allowed comparing a sufficient number of species: only half of plant demographic

studies exceed four years in duration (Römer et al., 2021; Salguero-Gómez et al., 2015). This

criterion resulted in 1,057 MPMs originating from 77 populations across 34 species, with an

average length of 15 years (median = 10; range = 7-36), and an average number of two populations

(median = 1, range=1-6). We note that in our analysis one species, *Cirsium pitcheri*, is repeated

three times because each original study examined a different population, or a different set of

populations (Supplementary Online Materials Table S1).

We used these MPMs to calculate the expectation of survival, development, reproduction,

and the population growth rate (λ) associated with each annual MPM. By "development", we refer

to changes in life cycle stage. These four year-specific metrics refer to a population at its stable

stage distribution: the distribution of discrete stages that would result if each yearly MPM's vital

rates were kept constant. We calculated survival, development, and reproduction using standard

methods (Franco & Silvertown, 2004), and λ as the dominant eigenvalue of each MPM (Caswell,

2001) with the R package popbio (Stubben & Milligan, 2007). Here, $\lambda >$ (or $<$) 1 corresponds to a

population that is expected to grow (decline) at the stable stage distribution (Caswell, 2001). We

calculated net annual vital rates by taking a weighted mean of the stage-specific vital rates, where

the weights were derived from the stable stage distribution.

We used gridded and weather station climatic data to generate the climatic predictors for

our statistical models. The climatic predictors in our models include mean monthly air temperature

and precipitation at the location of each of our 77 populations. We obtained these data from the

CHELSA (Karger et al., 2017) and PRISM (Daly et al., 1994) datasets, and a single Global

Historical Climatology Network (GHCN) daily weather station (Menne et al., 2012). Both

CHELSA and PRISM provide gridded climatic data at a 30 arcsec (~1 km^2) resolution which

approximate weather station climate averages well (Behnke et al., 2016), while providing a

standardized source of climatic data. We used CHELSA data for North American populations

sampled between 1979 and 2013 (Supplementary Online Materials Table S1). We used PRISM

data for the two populations of *Astragalus scaphoides* which extend to 2014 (Tenhumberg *et al.*

2018). We used data from weather station number USC00143527 of the GHCN for the 10

populations collected between 1938 and 1972 in Hays, Kansas, USA (Supplementary Online

Materials Table S1).

Our climatic predictors consisted of the monthly climatic anomalies observed in the 36

months preceding each vital rate observation (*i.e.*, preceding the end month of a given demographic

projection interval). Specifically, vital rate observations describe a transition from year *t* to year *t*

+ 1: our models use the 36 monthly anomalies preceding year *t* + 1 as predictors. These anomalies

were *z*-scores (mean = 0, SD = 1) with respect to the long-term monthly mean. We calculated long-

term means exceeding the recommended minimum of 30 years of data (World Meteorological

Organization, 2017). For CHELSA and PRISM datasets, we calculated the long-term monthly

mean using a 35-year period between 1979 and 2013, and 1980 and 2014, respectively. We

calculated the long-term monthly mean for the 10 datasets collected before 1979 using 50 years of

climate data. The difference between anomalies computed using the 50- versus 35-year mean was

negligible, as their correlations were 0.996 and 0.995 for temperature and precipitation,

respectively.

Modeling overview

Using vital rates and $\log(\lambda)$ as response variables, we compared and contrasted the predictive performance of five models (Fig. 1): the (1) Null Model (NM), with no climatic covariates, the (2) Climate Summary Model (CSM, Dalgleish et al., 2011), using yearly climate as a predictor, and three types of antecedent effect models: the (3) Weighted Means Models (WMM, van de Pol & Cockburn, 2011b), the (4) Stochastic Antecedent Model (SAM, Ogle et al., 2015), and the (5) Finnish Horseshoe Model (FHM, Hastie et al., 2009). Models 1-5 are ranked by increasing flexibility, by progressively relaxing key assumptions (Fig. 1). NMs are a "model of the mean" that do not account for the effect of climatic drivers. CSMs assume that the monthly anomalies observed within a specific period have all equal weight and that these anomalies affect population dynamics in the same direction. Therefore, this model uses as a predictor the mean of 12 monthly anomalies. WMM's weigh the contribution of the 12 monthly anomalies based on a Gaussian function. For example, monthly anomalies could decrease in weight depending on their distance to the demographic census (Fig. 1). Like in WMMs, SAMs assume that monthly anomalies influence population dynamics in the same direction. SAMs construct relative weights of monthly anomalies using coefficients drawn from a Dirichlet distribution (Fig. 1; Ogle et al., 2015). Finally, FHMs fit a separate coefficient for each of the 12 monthly anomalies, shrinking these toward zero through their prior distribution. In regularized horseshoe regressions, monthly anomalies can have positive or negative effects. FHMs were not developed to model climatic effects (Piironen & Vehtari, 2017), though regularization approaches have been previously employed to estimate climatic effects on vital rate information (e.g. Tredennick et al., 2016).

We fit data from three separate years to account for the potential of lagged climatic effects.

There is convincing evidence of lagged climatic effects in several case studies, including for three

populations analyzed in this study: one population of *Cryptantha flava* (Evers et al., 2021) and two

of *Astragalus scaphoides* (Tenhumberg et al., 2018). We re-fit each model using climate data from

three separate years: each year corresponding to a subset of the 36 monthly anomalies preceding

each demographic observation. We subdivided these 36 anomalies as belonging to the year (year

t), one year (*t*-1), and two years (*t*-2) preceding each demographic observation. Hence, we fit 13

models for each vital rate or estimation of population growth rate and climate variable

combination.

Below, we describe the equations of the models we tested assuming a normal response

variable, $\log(\lambda)$. To facilitate model comparison and model convergence, we used weakly

informative priors for the intercept and variance terms (Lemoine, 2019). We document the priors

for all models and the prior predictive checks we ran in the section of the Supplementary Online

Materials on prior predictive checks. Our null is a model of the mean, such that

$$y_{ti} \sim Normal(\hat{y}, \sigma) \qquad (1a)$$

$$\hat{y} = \alpha, \qquad (1b)$$

$$\alpha \sim Normal(0, 0.5), \qquad (1c)$$

$$\sigma \sim Gamma(0.01, 0.01), \qquad (1d)$$

where y_{ti} refers to a $\log(\lambda)$ value, *t* refers to the year of observation, *i* to a replicate within years (*i*

exceeds one in datasets that are spatially replicated), $\hat{y}$ is the mean model prediction, and α is a

model intercept, the mean. In the CSM, we model the response variable as a function of $\bar{x}_t$, the

mean of the 12 monthly anomalies (z-scores) over year *t,* as

$$\hat{y}_t = \alpha + \beta \bar{x}_t, \qquad (2a)$$

$$\beta \sim Normal(0,1), \qquad (2b)$$

where β is the coefficient for the effect of $\bar{x}_t$, and $\hat{y}$ is the mean model prediction referred to year

t.

The first antecedent effect model is the WMM, which models the response as a function of

x_{tk}, the monthly temperature deviations for year t and month k:

$$\hat{y}_t = \alpha + \beta \sum_k w_k x_{kt} \qquad (3a)$$

$$w_k = g_k / \sum_k g_k \qquad (3b)$$

$$g_k = \frac{1}{\sigma_w \sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma_w}\right)^2} \qquad (3c)$$

$$\mu \sim Normal(6.5,12)[0,12], \qquad (3d)$$

$$\sigma_w \sim Normal(1,12)[1,\,], \qquad (3e)$$

where the 12 x_k monthly anomalies prior to the end of the relevant demographic year, are each

weighted by w_k. In turn, w_k is a normalized vector of Gaussian weights, g_k, with mean μ and scale

σ_w. The square brackets in Eq. 3d and 3e show the limits beyond which the potential values of

these parameters are truncated.

The SAM (Ogle et al. 2015) adds flexibility to the WMM above,

$$\hat{y}_t = \alpha + \beta \sum_k w_k x_{kt}, \qquad (4a)$$

$$\boldsymbol{w} = \text{Dirichlet}\left(\alpha, \overbrace{\ldots, \alpha}^{n=12}\right), \qquad (4b)$$

where vector $\boldsymbol{w}$ contains 12 weights that sum to 1, and w_k is a single weight that refer to monthly

anomaly x_{kt} observed in month k and year t. We set the concentration parameter (α) to 1,

corresponding to a uniform distribution on the 12-dimensional simplex of weights. The added

flexibility originates because the values in vector $\boldsymbol{w}$ do not follow a functional form (albeit they

are correlated). Henceforth, we refer to the term $\sum_k w_k x_{kt}$, in both Eq. 3a and 4a, as the

"antecedent" of year t.

Finally, the FHM is a multiple regression whose coefficients are shrunk via the "Finnish"

horseshoe prior proposed by Piironen & Vehtari (2017). These models estimate a separate slope

value, β_k, for each k month. The Finnish horseshoe prior prevents overfitting by shrinking estimates

towards zero. This model is fit as follows:

$$\hat{y}_t = \alpha + \sum_k \beta_k x_{kt}, \qquad (5a)$$

$$\beta_k \sim Normal\left(0, \tau \tilde{\theta}_k\right), \qquad (5b)$$

$$\tilde{\theta}_k = \frac{c\theta_k}{\sqrt{c^2 + \tau^2 \theta_k^2}}, \qquad (5c)$$

$$\theta_k \sim HalfCauchy(0,0.1), \qquad (5d)$$

$$c^2 \sim InvGamma\left(\frac{v}{2}, \frac{v}{2}s^2\right), \qquad (5e)$$

$$\tau \sim HalfCauchy(0, \tau_0), \qquad (5f)$$

$$\tau_0 = \frac{m_0}{M - m_0} \frac{\sigma}{\sqrt{N}}. \qquad (5g)$$

Here, 12 β_k values depend on a normal distribution whose standard deviation is controlled

by τ, the "global scale", and 12 $\tilde{\theta}_k$ "local scales". The global scale τ tends to shrink every

monthly estimate β_k toward zero. On the other hand, the $\tilde{\theta}_k$ values allow the β_k monthly estimates

to become relatively large. In the FHM, the combination of τ and $\tilde{\theta}_k$ shrinks all β_k coefficients

toward zero, while controlling the magnitude of β_k values for which there is evidence for a

substantial departure from zero. We modified these models to fit survival, development, and

reproduction data, which are non-normally distributed. We describe these modifications in the

Supplementary Online Material.

Model fitting

We fit models using a Hamiltonian Monte Carlo sampler via the *R* package *rstan* (Stan

Development Team 2018). We fit models running four parallel chains for 4,000 iterations, with a

burn-in of 1,000 iterations, and thinning every other iteration. Because we had 13 alternative

models, four demographic responses ($\log(\lambda)$ and three vital rates), two climatic predictors, and 36

datasets, we fit a total of 3,744 models. Fitting these many models made it unfeasible to check

models one at a time. Therefore, we opted to perform model checks across model types, response

variables, and dataset length. In the Supplementary Online Material, we provide the methods and

results to assess the convergence of the Hamiltonian Monte Carlo samplers and assess model fit

(Bayesian model checks).

Model comparison

We compared model performance for each vital rate, climate predictor, and species

combination. We performed comparisons based on the log pointwise predictive density (LPPD)

(Bernardo & Smith, 2009; Hooten & Hobbs, 2015) calculated through a leave-one-year-out

(LOYO) cross-validation (e.g. Tredennick et al., 2016). A LOYO cross-validation creates a

training set by leaving out an entire year of data rather than a single data point. The higher the

value of LPPD of the held-out data, the better the predictive performance of the model. Our model

selection focused on LOYO cross-validation, because we aim to predict year-level variation in

vital rates. We performed this procedure because the climate is strongly correlated among different

sites in our spatially replicated datasets. In particular, 16 of our 36 datasets are spatially replicated,

and their mean correlation is 0.98 (range = 0.74-1).

We compared model performance by plotting the difference in ΔLPPD — the difference

in LPPD with the best model. The $\Delta LPPD_i$ of model i is calculated as $LPPD_i$ – max(LPPD). For

each species, we plotted heat maps of the $\Delta LPPD_i$ values for each model, flagging the best three

models. We then ordered heat maps by data length along the y-axis, and by the year of climate

data (*t*, *t*-1, and *t*-2) on the x-axis.

We addressed our three objectives by testing for differences in the relative support of our

models using Pseudo-Bayesian Model Averaging. We calculated model weights using Pseudo-

Bayesian Model Averaging for each combination of species, response variable, and predictor

(temperature or precipitation). We calculated model weights following equations 8 and 9 in Yao

et al. (2018). LPPD relative weights using the same formula used for AIC weights, so that

288
$$w_k = \frac{\exp(LPPD_k - \frac{1}{2}se(\text{LPPD}_k))}{\sum_{k=1}^{K} \exp(LPPD_k - \frac{1}{2}se(\text{LPPD}_k))} \qquad (8)$$

where *k* refers to each model, *se* is the standard error, and *K* is 13, the total number of models

tested for each combination of species, response variable, and climatic predictor. We arcsine

transformed these LPPD weights to make them normally distributed. We then used these acrsine

transformed LPPD weights as the response variable to address our three objectives. We addressed

these objectives using graphs and average statistics. We plotted boxplots of model weights across

all datasets, grouping them based on model types. This evaluated (1) whether antecedent effect

models outperformed NMs and CSMs, and which antecedent effect model had the best predictive

performance. Then, we plotted model weights against the temporal replication of each dataset to

(2) verify whether such replication increased the predictive ability of antecedent effect models.

We fit quantile regressions on these plots using the R package quantreg (Koenker, 2022), from the

10[th] to the 90[th] quantile in increments of 10. Finally, we plotted model weights of NMs grouping

them by response variable to verify (3) which response variable climate is a better predictor of.

Note that in this case, the lower the model weight of NMs, the higher the predictive ability of

climate.

Case study and exploratory analysis

We present a case study to showcase the potential of antecedent effect models as exploratory and

predictive tools. First, we show how antecedent effect models identify monthly climatic drivers of

demographic responses, and we discuss how these inferences can inform biological hypotheses.

Second, we show how antecedent effect models improve model fit. Here, we present data from

five natural populations of *Astragalus tyghensis,* an endemic legume plant species with an

extremely restricted range encompassing arid lands of Western North America (Thorpe & Kaye,

2008). We present models fit on the reproduction of *Astragalus tyghensis* using precipitation

anomalies one year prior to the last demographic observation (year *t*-1). This climatic predictor

had high predictive power in our model comparisons.

We use this case study to show for each model its fit to data and the effect sizes of monthly

climatic anomalies. To show model fit to data, we plotted each raw reproduction data point against

200 randomly selected posterior samples predictions. To display the monthly effect sizes, we

produced estimates which were directly comparable among CSMs, WMMs, SAMs, and FHMs.

For the CSM, we divided the regression slope, β (Eq. 2a), by 12, the number of months. For the

WMMs and SAMs, we multiplied the regression slope, β (Eq. 3-4a), by the 12 monthly weights,

w_k (Eq. 3-4b). Finally, the FHMs directly estimate 12 β_k separate regression slopes. We

represented these monthly effect sizes by plotting the median and the 95% credible intervals of the

posterior on the y-axis, against the month number on the x-axis.

Results

Model comparison

We found no evidence that antecedent effect models have higher predictive power than climate summary models. When comparing LPPD weights, the NMs were on average the best performing, followed by CSMs, and then antecedent effect models (Fig. 2). The variance in model weights was smaller in NMs models, while the models including climatic predictors could reach very high or very low model weights (Fig. 2). This variance in the LPPD weights suggests that estimating climate effects is advantageous for a subset of populations which are presumably particularly sensitive to climate. However, estimating climate effects can substantially decrease predictive ability in the other populations. The best performing antecedent effect models were the FHMs, followed by WMMs, and SAMs. However, these average differences were small, potentially reflecting, as shown in our case study, that these three models tend to provide similar information (Fig. 3). The superior performance of FHMs is in part driven by the relatively good performance of these models in $\log(\lambda)$ models (Fig. 4-5; Supplementary Online Materials: Fig. S8).

The temporal replication of the studies did not improve average model performance, but it decreased the likelihood of fitting models with very low LPPD relative weights. The only quantile regression that shows a significant trend is linked to the 10^{th} quantile, for which temporal replication increases LPPD relative weights (Supplementary Online Materials: Fig. S9). This result suggests that sample size only limits the negative effects of fitting models with large numbers of parameters on predictive ability.

Models including climate tended to predict development better than the other response

variables (Supplementary Online Materials: Fig S10). This higher predictive ability was due to the

higher predictive performance of CSM models (Fig. 3, Fig. 4-5; Supplementary Online Materials:

Fig S8). Moreover, models including climate predicted log(λ) worse than the other response

variables (Fig. S10).

Case study and exploratory analysis

The results of antecedent effect models were biologically interpretable, but they did not

substantially improve the ability to predict the reproduction of *Astragaluys tygensis*. All four

models show that precipitation is negatively associated with reproduction. According to the

WMM, the monthly anomalies preceding a demographic census tend to be the most important in

driving reproduction. The SAM and FHM provide more precise insights, indicating that the most

important climate windows are located two (May) and six (January) months prior to the

demographic census. Thus, antecedent effect models indicate that the demography of this species

is correlated to the precipitation of the growing and dormant seasons. However, the model fits of

antecedent effect models and the CSM are similar (Fig. 3). Hence, despite the predictive

performance of antecedent effect models is slightly better in this case, this is a quantitatively small

difference.

Discussion

Understanding how climate affects the vital rates (survival, development, reproduction) of natural populations and their overall performance (*e.g.*, population growth rate, λ) is a core ecological question. Contrary to our expectations, we found that simple climate summary models (CSMs) showed, on average, slightly better predictive ability than the substantially more complex antecedent effect models. Moreover, we found dataset length had little role in modulating these results. Therefore, we advocate using antecedent effects models as exploratory tools to facilitate hypothesizing the windows of time most strongly associated with demographic variation. Finally, we found predicting climatic effects is easier for development and harder to for population growth rate. This last result suggests that development is more responsive to relatively short-term climatic drivers, and that species responses to climate might reflect "demographic buffering".

Our initial expectation was that antecedent effect models would improve predictions because they accommodate a variety of potential links between demography and climate (Evers et al., 2021; Kreyling, 2010). However, our model comparisons indicate that simple linear models have the best predictive ability in our small datasets. Selecting the most accurate model among a candidate set trades off the bias and variance of model predictions (Hastie et al. 2009). Models such as SAM and FHM are expected to show low bias because their flexibility captures complex relationships. However, in small datasets complex models with many parameters are likely estimated with high uncertainty (variance). Bayesian regularization aims to address this problem by limiting the prior variance of the model parameters. Nonetheless, a simple model with few parameters may achieve similar levels of predictive accuracy when its structural bias is sufficiently low. Our case study suggests another reason why antecedent effect models may not outperform simpler models: the monthly precipitation anomaly with the strongest effect, May, had a

correlation with the annual precipitation anomaly of 0.5. In our dataset, the median correlation

between monthly anomalies and yearly anomalies is 0.37. Such correlations will reduce the

predictive performance gained through the additional flexibility of the antecedent effect models.

The case study exemplified the exploratory power of antecedent effect models. In

Astragalus tyghensis, both the SAM and FHM show a strong negative association between

reproduction and the precipitation anomalies that occurred in the May and January of the preceding

year. These results raise three hypotheses. First, such negative association with precipitation

anomalies occurring a year prior to a demographic observation could result from indirect effects

mediated through competition (Adler et al., 2012; Suttle et al., 2007) or physiological tradeoffs

(*e.g.*, costs of reproduction, Crone et al., 2009; Miller et al., 2012). Second, the correlation with

the precipitation in January, a month when the mean temperature is just above or below $0°$ C, could

reflect the effect of snowfall. This effect is often distinct from rain (Compagnoni & Adler, 2014;

Dalgleish et al., 2011) and could arise in different ways: for example, because snow affects

pathogens (Smull et al., 2019), it protects plants from frost (Inouye, 2000), or because snowmelt

provides a large pulse of resources (Noy-Meir, 1973). Third, the large effect of May precipitation

could reflect a key developmental phase occurring either in May or June. All three hypotheses call

for further tests to be performed via analysis of existing data or field experiments.

We hypothesize that the higher predictability of development with respect to other vital

rates is a general pattern in plants. First, predicting $\log(\lambda)$ should be more challenging, because

evolution should favor buffering its variation through time (Hilde et al., 2020; Tuljapurkar, 1982).

Evidence of such buffering is typically tested, across species, using data on the temporal variability

of vital rates (Le Coeur et al., 2022; McDonald et al., 2017). Here we provide the first evidence,

to our knowledge, that across many species, demographic buffering is reflected in the link between

climatic drivers and vital rates. Second, understanding how climate affects plant survival is still a

rapidly developing field (Trugman et al., 2021). For example, the survival of plants to drought is

specific to the species, plant functional type, and a large number of contextual factors (Volaire,

2018). Third, understanding the response of reproduction to climate is not straightforward: seed

production reflects indirect effects of development (Clark et al., 2021), and the transition from

seed to new recruits is a source of uncertainty (Paniw et al., 2017). On the other hand, the strong

dependence of development to climate is posited by physiology and demonstrated by ecosystem

models (Schulze et al., 2019).

Our experience building and comparing antecedent effect models informs our suggestions

on hyperparameter priors and emphasizes the role of prior predictive checks. The choice of

hyperparameter priors has strong effects on model fitting and performance. In weighted mean

models (WMMs), truncated hyperparameter priors (Eq. 3d-e) prevented the sampler from

exploring unrealistic parameter values, greatly facilitating model convergence. In SAMs, we chose

a vector of ones as the Dirichlet concentration parameter (Eq. 4b) to not favor any particular lag,

and to ensure that prior weights had a uniform distribution. A much lower number than 1 would

favor models with a single non-zero weight. In FHMs, we suggest using a standard deviation of

0.1 for the prior on monthly effect sizes (Eq. 5d). Using a a standard deviation of 1 (*e.g.*, Piironen

& Vehtari, 2017) generated unrealistically high variation in our prior predictive checks

(Supplementary Online Materials: prior predictive checks). More generally, prior predictive

checks were essential because our initial prior choices produced very different variance in

predictions among model types and response variables. Moreover, priors generating unrealistically

high variance had model convergence issues and lower predictive performance.

Our results justify the continued data and conceptual efforts to understand and predict the effects

of climatic drivers on survival and reproduction. Compared with development, the

Acknowledgements

AC and TMK were supported by the Alexander von Humboldt Foundation under the framework

of the Humboldt Professorship, and by the Helmholtz Recruitment Initiative of the Helmholtz

Association (both to TMK); RSG was supported by NERC (NE/M018458/1). This work emanated

from research discussions during the sAPROPOS (Analysis of PROjections of POpulationS)

working group led by RSG and TMK at the German Centre for Integrative Biodiveristy Research

(iDiv), which is funded by the German Research Foundation (FZT-118-202548816). We thank the

Max Planck Institute for Demographic Research for the support of the COMPADRE Plant Matrix

Database and the hundreds of plant population ecologists who have contributed with demographic

data to COMPADRE. We thank B. Teller and P. Barks for discussions in the early phases of this

project.

Conflict of Interest statement

We declare no conflict of interest

Author Contributions

RS-G and AC conceived the study; AC, RS-G, and DC designed the research; AC, performed the

research, analyzed the data, and wrote the article; RS-G, DC, and TMK provided critical

revisions of the article, and approved the final version of the article.

Data archiving statement

We will store the code to reproduce these results on Zenodo. The demographic data used in this

article come from the open access COMPADRE Plant Matrix Database (v. 5.0.1; https://compadre-

db.org/Data/Compadre). The datasets in COMPADRE used in this study are identified in the first

column of Supplementary Table S1. The climate data used in this study come from CHELSA

(available at http://dx.doi.org/doi:10.5061/dryad.kd1d4), PRISM (available at

https://ftp.prism.oregonstate.edu/), and the Global Historical Climatology Network daily

(available at https://dx.doi.org/10.7289/V5D21VHZ).

References

Adler, P. B., Dalgleish, H. J., & Ellner, S. P. (2012). Forecasting plant community impacts of climate variability and change: When do competitive interactions matter? *Journal of Ecology*, *100*, 478–487.

Behnke, R., Vavrus, S., Allstadt, A., Albright, T., Thogmartin, W. E., & Radeloff, V. C. (2016). Evaluation of downscaled, gridded climate data for the conterminous United States. *Ecological Applications*, *26*(5), 1338–1351. https://doi.org/10.1002/15-1061

Bell, T. J., Powell, K. I., & Bowles, M. L. (2013). Viability model choice affects projection accuracy and reintroduction decisions. *The Journal of Wildlife Management*, *77*(6), 1104–1113. https://doi.org/10.1002/jwmg.525

Bernardo, J. M., & Smith, A. F. M. (2009). *Bayesian Theory*. John Wiley & Sons.

Caswell, H. (2001). *Matrix population models. Construction, analysis, and interpretation.* Sinauer Associates.

Catchpole, E. A., Morgan, B. J. T., Coulson, T. N., Freeman, S. N., & Albon, S. D. (2000). Factors influencing Soay sheep survival. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, *49*(4), 453–472. https://doi.org/10.1111/1467-9876.00205

Clark, J. S., Andrus, R., Aubry-Kientz, M., Bergeron, Y., Bogdziewicz, M., Bragg, D. C., Brockway, D., Cleavitt, N. L., Cohen, S., Courbaud, B., Daley, R., Das, A. J., Dietze, M., Fahey, T. J., Fer, I., Franklin, J. F., Gehring, C. A., Gilbert, G. S., Greenberg, C. H., … Zlotin, R. (2021). Continent-wide tree fecundity driven by indirect climate effects. *Nature Communications*, *12*(1), 1. https://doi.org/10.1038/s41467-020-20836-3

Clark, J. S., Iverson, L., Woodall, C. W., Allen, C. D., Bell, D. M., Bragg, D. C., D'Amato, A. W., Davis, F. W., Hersh, M. H., Ibanez, I., Jackson, S. T., Matthews, S., Pederson, N., Peters, M., Schwartz, M. W., Waring, K. M., & Zimmermann, N. E. (2016). The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States. *Global Change Biology*, *22*(7), 2329–2352. https://doi.org/10.1111/gcb.13160

Compagnoni, A., & Adler, P. B. (2014). Warming, soil moisture, and loss of snow increase Bromus tectorum's population growth rate. *Elementa: Science of the Anthropocene*, *2*, 000020. https://doi.org/10.12952/journal.elementa.000020

Compagnoni, A., Levin, S., Childs, D. Z., Harpole, S., Paniw, M., Römer, G., Burns, J. H., Che-Castaldo, J., Rüger, N., Kunstler, G., Bennett, J. M., Archer, C. R., Jones, O. R., Salguero-Gómez, R., & Knight, T. M. (2021). Herbaceous perennial plants with short generation time have stronger responses to climate anomalies than those with longer generation time. *Nature Communications*, *12*(1), 1824. https://doi.org/10.1038/s41467-021-21977-9

Crone, E. E., Miller, E., & Sala, A. (2009). How do plants know when other plants are flowering? Resource depletion, pollen limitation and mast-seeding in a perennial wildflower. *Ecology Letters*, *12*(11), 1119–1126. https://doi.org/10.1111/j.1461-0248.2009.01365.x

Czachura, K., & Miller, T. E. X. (2020). Demographic back-casting reveals that subtle dimensions of
climate change have strong effects on population viability. *Journal of Ecology*, *108*(6), 2557–2570.
https://doi.org/10.1111/1365-2745.13471

Dalgleish, H. J., Koons, D. N., Hooten, M. B., Moffet, C. A., & Adler, P. B. (2011). Climate influences the
demography of three dominant sagebrush steppe plants. *Ecology*, *92*(1), 75–85.
https://doi.org/10.1890/10-0780.1

Daly, C., Neilson, R. P., & Phillips, D. L. (1994). A Statistical-Topographic Model for Mapping
Climatological Precipitation over Mountainous Terrain. *Journal of Applied Meteorology*, *33*(2), 140–158.
https://doi.org/10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2

Ehrlén, J., Morris, W. F., von Euler, T., & Dahlgren, J. P. (2016). Advancing environmentally explicit
structured population models of plants. *Journal of Ecology*, *104*(2), 292–305.
https://doi.org/10.1111/1365-2745.12523

Ellis, M. M., Williams, J. L., Lesica, P., Bell, T. J., Bierzychudek, P., Bowles, M., Crone, E. E., Doak, D. F.,
Ehrlén, J., Ellis-Adam, A., McEachern, K., Ganesan, R., Latham, P., Luijten, S., Kaye, T. N., Knight, T. M.,
Menges, E. S., Morris, W. F., Nijs, H. den, … Weekley, C. W. (2012). Matrix population models from 20
studies of perennial plant populations. *Ecology*, *93*(4), 951–951. https://doi.org/10.1890/11-1052.1

Ellner, S. P., Childs, D. Z., & Rees, M. (2016). *Data-driven Modelling of Structured Populations: A Practical*
Guide to the Integral Projection Model. Springer International Publishing. https://doi.org/10.1007/978-
3-319-28893-2

Elton, C. S. (1924). Periodic Fluctuations in the Numbers of Animals: Their Causes and Effects. *Journal of*
Experimental Biology, *2*(1), 119–163. https://doi.org/10.1242/jeb.2.1.119

Evers, S. M., Knight, T. M., Inouye, D. W., Miller, T. E. X., Salguero-Gómez, R., Iler, A. M., & Compagnoni,
A. (2021). Lagged and dormant season climate better predict plant vital rates than climate during the
growing season. *Global Change Biology*, *27*(9), 1927–1941. https://doi.org/10.1111/gcb.15519

Franco, M., & Silvertown, J. (2004). A Comparative Demography of Plants Based upon Elasticities of Vital
Rates. *Ecology*, *85*(2), 531–538.

Hacket-Pain, A. J., Ascoli, D., Vacchiano, G., Biondi, F., Cavin, L., Conedera, M., Drobyshev, I., Liñán, I. D.,
Friend, A. D., Grabner, M., Hartl, C., Kreyling, J., Lebourgeois, F., Levanič, T., Menzel, A., Maaten, E. van
der, Maaten-Theunissen, M. van der, Muffler, L., Motta, R., … Zang, C. S. (2018). Climatically controlled
reproduction drives interannual growth variability in a temperate tree species. *Ecology Letters*, *21*(12),
1833–1844. https://doi.org/10.1111/ele.13158

Harper, J. L. (1967). A Darwinian Approach to Plant Ecology. *Journal of Ecology*, *55*(2), 247–270. JSTOR.
https://doi.org/10.2307/2257876

Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The Elements of Statistical Learning: Data Mining,*
Inference, and Prediction, Second Edition. Springer Science & Business Media.

Hilde, C. H., Gamelon, M., Sæther, B.-E., Gaillard, J.-M., Yoccoz, N. G., & Pélabon, C. (2020). The
Demographic Buffering Hypothesis: Evidence and Challenges. *Trends in Ecology & Evolution*, *35*(6), 523–
538. https://doi.org/10.1016/j.tree.2020.02.004

Hoerl, A. E., & Kennard, R. W. (1970). Ridge Regression: Biased Estimation for Nonorthogonal Problems.
Technometrics, *12*(1), 55–67. https://doi.org/10.1080/00401706.1970.10488634

Hooten, M. B., & Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists. *Ecological*
Monographs, *85*(1), 3–28.

Inouye, D. w. (2000). The ecological and evolutionary significance of frost in the context of climate
change. *Ecology Letters*, *3*(5), 457–463. https://doi.org/10.1046/j.1461-0248.2000.00165.x

Jolls, C. L., Marik, J. E., Hamzé, S. I., & Havens, K. (2015). Population viability analysis and the effects of
light availability and litter on populations of Cirsium pitcheri, a rare, monocarpic perennial of Great
Lakes shorelines. *Biological Conservation*, *187*, 82–90. https://doi.org/10.1016/j.biocon.2015.04.006

Jr, M. J. A., & Angilletta, M. J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis*. OUP
Oxford.

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder,
H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific*
Data, *4*, 170122. https://doi.org/10.1038/sdata.2017.122

Karger, D. N., & Zimmermann, N. E. (2018). *CHELSAcruts – High resolution temperature and precipitation*
timeseries for the 20th century and beyond. http://dx.doi.org/10.16904/envidat.159

Kaye, T. N., & Pyke, D. A. (2003). The Effect of Stochastic Technique on Estimates of Population Viability
from Transition Matrix Models. *Ecology*, *84*(6), 1464–1476. https://doi.org/10.1890/0012-
9658(2003)084[1464:TEOSTO]2.0.CO;2

Knape, J., & de Valpine, P. (2011). Effects of weather and climate on the dynamics of animal population
time series. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1708), 985–992.
https://doi.org/10.1098/rspb.2010.1333

Koenker, R. (2022). *quantreg: Quantile Regression* (5.93) [Computer software]. https://CRAN.R-
project.org/package=quantreg

Kreyling, J. (2010). Winter climate change: A critical factor for temperate vegetation performance.
Ecology, *91*(7), 1939–1948. https://doi.org/10.1890/09-1160.1

Le Coeur, C., Yoccoz, N. G., Salguero-Gómez, R., & Vindenes, Y. (2022). Life history adaptations to
fluctuating environments: Combined effects of demographic buffering and lability. *Ecology Letters*,
25(10), 2107–2119. https://doi.org/10.1111/ele.14071

Lemoine, N. P. (2019). Moving beyond noninformative priors: Why and how to choose weakly
informative priors in Bayesian analyses. *Oikos*, *128*(7), 912–928. https://doi.org/10.1111/oik.05985

McDonald, J. L., Franco, M., Townley, S., Ezard, T. H. G., Jelbert, K., & Hodgson, D. J. (2017). Divergent
demographic strategies of plants in variable environments. *Nature Ecology & Evolution*, *1*(2), 0029.
https://doi.org/10.1038/s41559-016-0029

Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E., & Houston, T. G. (2012). An Overview of the Global
Historical Climatology Network-Daily Database. *Journal of Atmospheric and Oceanic Technology*, *29*(7),
897–910. https://doi.org/10.1175/JTECH-D-11-00103.1

Miller, T. E. X., Williams, J. L., Jongejans, E., Brys, R., & Jacquemyn, H. (2012). Evolutionary demography of iteroparous plants: Incorporating non-lethal costs of reproduction into integral projection models. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1739), 2831–2840. https://doi.org/10.1098/rspb.2012.0326

Noy-Meir, I. (1973). Desert Ecosystems: Environment and Producers. *Annual Review of Ecology and Systematics*, 25–51.

Ogle, K., Barber, J. J., Barron-Gafford, G. A., Bentley, L. P., Young, J. M., Huxman, T. E., Loik, M. E., & Tissue, D. T. (2015). Quantifying ecological memory in plant and ecosystem processes. *Ecology Letters*, *18*(3), 221–235. https://doi.org/10.1111/ele.12399

Paniw, M., Quintana-Ascencio, P. F., Ojeda, F., & Salguero-Gómez, R. (2017). Accounting for uncertainty in dormant life stages in stochastic demographic models. *Oikos*, *126*(6), 900–909. https://doi.org/10.1111/oik.03696

Piironen, J., & Vehtari, A. (2017). Sparsity information and regularization in the horseshoe and other shrinkage priors. *Electronic Journal of Statistics*, *11*(2), 5018–5051. https://doi.org/10.1214/17-EJS1337SI

Römer, G., Dahlgren, J. P., Salguero-Gómez, R., Stott, I. M., & Jones, O. R. (2021). *Plant demographic knowledge is biased towards short-term studies of temperate-region herbaceous perennials* (p. 2021.04.25.441327). bioRxiv. https://doi.org/10.1101/2021.04.25.441327

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Bein, C., de Buhr, H., Farack, C., Gottschalk, F., Hartmann, A., Henning, A., Hoppe, G., Römer, G., Ruoff, T., Sommer, V., Wille, J., Voigt, J., Zeh, S., Vieregg, D., Buckley, Y. M., Che-Castaldo, J., … Vaupel, J. W. (2016). COMADRE: A global data base of animal demography. *Journal of Animal Ecology*, *85*(2), 371–384. https://doi.org/10.1111/1365-2656.12482

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Buckley, Y. M., Che-Castaldo, J., Caswell, H., Hodgson, D., Scheuerlein, A., Conde, D. A., Brinks, E., de Buhr, H., Farack, C., Gottschalk, F., Hartmann, A., Henning, A., Hoppe, G., Römer, G., Runge, J., Ruoff, T., … Vaupel, J. W. (2015). The COMPADRE Plant Matrix Database: An open online repository for plant demography. *Journal of Ecology*, *103*(1), 202–218. https://doi.org/10.1111/1365-2745.12334

Schulze, E.-D., Beck, E., Buchmann, N., Clemens, S., Müller-Hohenstein, K., & Scherer-Lorenzen, M. (2019). Dynamic Global Vegetation Models. In E.-D. Schulze, E. Beck, N. Buchmann, S. Clemens, K. Müller-Hohenstein, & M. Scherer-Lorenzen (Eds.), *Plant Ecology* (pp. 843–863). Springer. https://doi.org/10.1007/978-3-662-56233-8_22

Shripad Tuljapurkar. (1982). Population dynamics in variable environments. III. Evolutionary dynamics of r-selection. *Theoretical Population Biology*, *21*(1), 141–165.

Simmonds, E. G., Cole, E. F., & Sheldon, B. C. (2019). Cue identification in phenology: A case study of the predictive performance of current statistical tools. *Journal of Animal Ecology*, *88*(9), 1428–1440. https://doi.org/10.1111/1365-2656.13038

Smull, D. M., Pendleton, N., Kleinhesselink, A. R., & Adler, P. B. (2019). Climate change, snow mold and
the Bromus tectorum invasion: Mixed evidence for release from cold weather pathogens. *AoB PLANTS*,
11(5), plz043. https://doi.org/10.1093/aobpla/plz043

Stubben, C., & Milligan, B. (2007). Estimating and Analyzing Demographic Models Using the popbio
Package in R. *Journal of Statistical Software*, *22*, 1–23. https://doi.org/10.18637/jss.v022.i11

Suttle, K. B., Thomsen, M. A., & Power, M. E. (2007). Species Interactions Reverse Grassland Responses
to Changing Climate. *Science*, *315*(5812), 640–642. https://doi.org/10.1126/science.1136401

Teller, B. J., Adler, P. B., Edwards, C. B., Hooker, G., & Ellner, S. P. (2016). Linking demography with
drivers: Climate and competition. *Methods in Ecology and Evolution*, *7*(2), 171–183.
https://doi.org/10.1111/2041-210X.12486

Tenhumberg, B., Crone, E. E., Ramula, S., & Tyre, A. J. (2018). Time-lagged effects of weather on plant
demography: Drought and Astragalus scaphoides. *Ecology*, *99*(4), 915–925.
https://doi.org/10.1002/ecy.2163

Thorpe, A. S., & Kaye, T. N. (2008). *Astragalus tyghensis: Actual vs. Predicted population sizes* (p. 18).
Institute for Applied Ecology for Prineville District BLM.

Tredennick, A. T., Hooten, M. B., & Adler, P. B. (2016). Do we need demographic data to forecast plant
population dynamics? *Methods in Ecology and Evolution*, *8*(5), 541–551. https://doi.org/10.1111/2041-
210X.12686

Trugman, A. T., Anderegg, L. D. L., Anderegg, W. R. L., Das, A. J., & Stephenson, N. L. (2021). Why is Tree
Drought Mortality so Hard to Predict? *Trends in Ecology & Evolution*.
https://doi.org/10.1016/j.tree.2021.02.001

van de Pol, M., Bailey, L. D., McLean, N., Rijsdijk, L., Lawson, C. R., & Brouwer, L. (2016). Identifying the
best climatic predictors in ecology and evolution. *Methods in Ecology and Evolution*, *7*(10), 1246–1257.
https://doi.org/10.1111/2041-210X.12590

van de Pol, M., & Cockburn, A. (2011). Identifying the Critical Climatic Time Window That Affects Trait
Expression. *The American Naturalist*, *177*(5), 698–707. https://doi.org/10.1086/659101

Volaire, F. (2018). A unified framework of plant adaptive strategies to drought: Crossing scales and
disciplines. *Global Change Biology*, *24*(7), 2929–2938. https://doi.org/10.1111/gcb.14062

White, M. A., Running, S. W., & Thornton, P. E. (1999). The impact of growing-season length variability
on carbon assimilation and evapotranspiration over 88 years in the eastern US deciduous forest.
International Journal of Biometeorology, *42*(3), 139–145. https://doi.org/10.1007/s004840050097

World Meteorological Organization. (2017). *WMO Guidelines on the Calculation of Climate Normals*
(World Meteorological Organization).

Yao, Y., Vehtari, A., Simpson, D., & Gelman, A. (2018). Using Stacking to Average Bayesian Predictive
Distributions (with Discussion). *Bayesian Analysis*, *13*(3). https://doi.org/10.1214/17-BA1091

Figure 1. Antecedent effect models used to estimate the effect of monthly climate anomalies on

population dynamics. These plots represent hypothetical Bayesian estimates of the A) weight and

B) effect sizes of 12 monthly climate anomalies observed before a demographic outcome (*e.g.*,

change in survival). In both panels, the symbols represent estimate means, and the extremes of the

error bars represent the 95% credible intervals. In panel A), the effect size of climate depends on

the relative weights of monthly anomalies. The Climate Summary Model (CSM) is a linear

regression where every monthly anomaly has the same weight; the climatic predictor in this

regression is thus given by the mean of 12 monthly anomalies. Hence, monthly weights are

identical, and no error bars are shown. In the Weighted Mean Model (WMM), monthly weights

follow a Gaussian curve; in the Stochastic Antecedent Model (SAM) the monthly weights are

determined by a Dirichlet distribution. Finally, panel B) shows how the Finnish Horseshoe Model

(FHM) tends to produce effect sizes centered around zero – except for the few effect sizes, two in

this hypothetical case, for which there is strong evidence of a positive and negative effect.

Figure 2. Null models (NMs) perform better than models using climate as predictor. Box and

whisker plots and point clouds showing the arcsine transformed model weights of the five model

types for 34 species. Panels separate models fit using precipitation or air temperature as predictor

variables. The middle line of the boxplots shows the median, the upper and lower hinges delimit

the first and third quartile, and the whiskers extend 1.5 times beyond the first and third quartile.

Each point refers to one of the 3,744 model fits, and colors refer to the response variable of each

model.

Figure 3. Antecedent effect models yield biologically meaningful insights. Case study showing

the fits of the climate summary model (CSM), and the three antecedent effect models (WMM,

SAM, FHM; see Fig. 1) to the reproduction of a natural population of the herbaceous perennial

Astragalus tyghensis, using precipitation anomalies two years preceding the demographic census.

In each plot, columns represent, respectively, model fit and climate effect sizes. The rows refer to

model type: respectively the Climate Summary Model (CSM), Weighted Mean Model (WMM),

Stochastic Antecedent Model (SAM), and Finnish Horseshoe Model (FHM). In the first column,

we show model fits using bivariate plots where the x-axis represents the observed reproductive

values, and the y-axis represents, for each observed value, 200 randomly selected predicted values

from the Bayesian posterior. In the second column, we show the effect sizes of each monthly

climatic anomaly. Each monthly effect size is shown as a point range where the black circles

represent the median, and the lines extend from the 2.5th to the 97.5th quantile of each posterior.

In the CSM, this effect sizes correspond to the regression slope, β (Eq. 2a), divided by 12. In the

WMM and SAM, the effect sizes are produced multiplying the regression slope, β (Eq. 3-4a) by

the monthly weights, w_k (Eq. 3-4b). Finally, the FHM estimates a separate slope, β_k, for each

monthly anomaly.

Figure 4. Climate Summary Models (CSM) show the best ability to predict vital rates using

precipitation data. Tile plot of predictive power based on differences in log pointwise predictive

density (ΔLPPD) for models using precipitation as predictor. Tiles with yellow colors denote

models with better predictive ability; the best models have a ΔLPPD of 0. Symbols denote best

(circles), second best (triangle), and third best (square) model. Each panel refers to a separate

response variable: A) survival, B) development, C) reproduction, and D) log(λ). Rows refer to each

dataset, and columns to each model. We compare five types of models (eqs. 1-5). We fit these

models using the climate observations from three years prior to a demographic response. Species

are organized by the temporal replication of the dataset, with the longest datasets being on top.

Figure 5. Climate Summary Models (CSMs) show the best ability to predict vital rates using air

temperature as predictor. Tile plot shows predictive power based on differences in log pointwise

predictive density (ΔLPPD) for models using temperature as predictor. The layout of the plot is

the same of Figure 4.

     Figure 1

706   **Figure 3.**

Figure 4.

712 **Figure 5.**

713