Nementin is a Nematode-Selective Small Molecule Agonist of Neurotransmitter Release

Sean Harrington[1,2], Jessica J. Knox[2,3], Andrew R. Burns[2,3], Ken-Loon Choo[4], Aaron Au[2,5], Megan Kitner[6], Cecile Haeberli[7,8], Jacob Pyche[1,2], Cassandra D'Amata[2], Yong-Hyun Kim[4], Jonathan R. Volpatti[9,10], Maximillano Guiliani[2,5], Jamie Snider[2], Victoria Wong[2], Bruna M. Palmeira[11], Elizabeth M. Redman[11], Aditya S. Vaidya[12,13], John S. Gilleard[11], Igor Stagljar[2,3,14,15,16], Sean R. Cutler[12,13], Daniel Kulke[17, †‡], James J. Dowling[9,10], Christopher M. Yip[2,5], Jennifer Keiser[7,8], Inga Zasada[6], Mark Lautens[4], and Peter J. Roy[1,2,3,20]*

Affiliations:

[1]Department of Pharmacology and Toxicology, University of Toronto, Toronto, ON, M5S 1A8, Canada.

[2]The Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, ON, M5S 3E1, Canada.

[3]Department of Molecular Genetics, University of Toronto; Toronto, ON, M5S 1A8, Canada.

[4]The Department of Chemistry, University of Toronto, 80 St. George Street, M5S 3H6, Toronto, Canada.

[5]Institute of Biomaterials and Biomedical Engineering, University of Toronto, Toronto, Ontario, Canada.

[6]USDA-ARS Horticultural Crops Research Laboratory, Corvallis, OR 97330, USA.

[7]Department of Medical Parasitology and Infection Biology; Swiss-Tropical and Public Health Institute, (Swiss TPH), Socinstr. 57, CH–4051 Basel, Switzerland.

[8]Faculty of Science; University of Basel, Petersplatz 1, 4001 Basel.

[9]Program in Developmental and Stem Cell Biology, The Hospital for Sick Children, Toronto, Ontario, Canada M5G 0A4.

[10]Program in Genetics and Genome Biology, The Hospital for Sick Children, Toronto, Ontario, Canada M5G 0A4.

[11]Department of Comparative Biology and Experimental Medicine, Host-Parasite Interactions (HPI) Program, Faculty of Veterinary Medicine, University of Calgary, Calgary, AB, Canada.

[12]Institute for Integrative Genome Biology, University of California, Riverside, Riverside, CA 92521, USA.

[13]Department of Botany and Plant Sciences, University of California, Riverside, Riverside, CA 92521, USA.

[14]Department of Biochemistry, University of Toronto, Toronto, ON, M5S 1A8, Canada.

[15]Mediterranean Institute for Life Sciences, Meštrovićevo Šetalište 45, HR-21000, Split, Croatia.

[16]School of Medicine, University of Split, Šoltanska ul. 2, 21000, Split, Croatia.

[17]Research Parasiticides, Bayer Animal Health GmbH, Monheim, Germany.

[†]Current affiliation, Department of Biomedical Sciences; Iowa State University, Ames, Iowa 50011, USA.

[‡]Current affiliation, Global Innovation, Boehringer Ingelheim Vetmedica GmbH; Binger Str. 173, 55218 Ingelheim am Rhein, Germany.

*Corresponding author.

The Donnelly Centre, Rm 1202, 160 College St., Toronto, ON, M5S 3E1, CANADA

4f16-978-6116, peter.roy@utoronto.ca

Classification: Biological Sciences, Genetics

Keywords: chemical genetics, nematicide, anthelmintic, *C. elegans*, parasitic nematodes*, Strongyloides ratti*, *Meloidogyne hapla*, small molecule screens, early-stage drug discovery, dense core vesicles, convulsions, dgk-1, egl-30, goa-1, unc-13, unc-17, unc-31, unc-43, unc-64

One-Sentence Summary: A *C. elegans*-based screening pipeline identifies a selective nematicide that also potentiates acetylcholinesterase inhibitors.

Abstract: Nematode parasites of humans, livestock and crops pose a significant burden on human health and welfare. Alarmingly, parasitic nematodes of animals have rapidly evolved resistance to anthelmintic drugs, and traditional nematicides that protect crops are facing increasing restrictions because of poor phylogenetic selectivity. Here, we present a pipeline that exploits multiple motor outputs of the model nematode *C. elegans* for nematicide discovery. This pipeline yielded multiple compounds that selectively kill and/or immobilize diverse nematode parasites. We focus on one compound that induces violent convulsions and paralysis that we call Nementin. We find that Nementin agonizes neuronal dense core vesicle release, which in turn agonizes cholinergic signaling. Consequently, Nementin synergistically enhances the potency of widely-used non-selective acetylcholinesterase inhibitors (AChEIs), but in a nematode-selective manner. Nementin therefore has the potential to reduce the environmental impact of toxic AChEI pesticides used to control nematode infections and infestations.

Significance Statement:

Parasitic nematodes pose a considerable burden to human health and food security. Small molecules that have traditionally been used to control these parasites have either been banned because of toxicity concerns or are being rendered ineffective because of the evolution of resistance. Significant gaps in our nematicidal toolkit are therefore becoming an alarming problem. Here, we describe our discovery of Nementin, a small molecule that disrupts the nematode nervous system but is ineffective against non-targeted organisms. We find that Nementin also enhances the activity of non-selective pesticides but does so in a nematode-selective manner. Hence, Nementin is an innovative solution to combat parasitic nematodes in a safe and phylum-selective manner.

Introduction

Parasitic nematodes are a scourge to humanity. Not only do helminths currently infect more than 1.5 billion people (WHO Soil Transmitted Helminth Fact Sheet, 2022) but nematode parasitism also destroys tens of billions of dollars (USD) worth of livestock annually[1]. Alarmingly, the rate of destruction is growing due to the evolution of anthelmintic resistance[2]. Plant-parasitic nematodes (PPNs) are even more destructive as they ruin over 125 billions of dollars (USD) worth of food crops annually[3]. Traditional nematicides that protect crops have been justifiably banned or severely restricted due to a lack of phylum selectivity[4], but these restrictions severely impact food security[5]. Compounding these issues, global food demand is expected to increase by 70% by the year 2050[6]. Hence, the development of new and selective nematicides is essential to our collective welfare.

Here, we present a novel motor-centric screening pipeline to identify novel candidate nematicides. Disrupting a parasite's motor control has repeatedly proven effective in mitigating nematode infection[7,8]. We therefore screened our collection of worm-active (wactive) compounds[9] using two successive behavioural assays of the free-living nematode *Caenorhabditis elegans*. We identified four molecules that fail to elicit phenotypes in off-target systems at concentrations that kill multiple nematode species. We focused on one of these, called Nementin, that induces *C. elegans* hyperactivity within seconds of exposure, followed by whole-body convulsions minutes later. Nementin and its analogs selectively immobilize and/or kill nematode parasites of plants and mammals. Chemical-genetic analyses reveal that Nementin agonizes neuronal dense core vesicle release, which in turn, agonizes synaptic vesicle release. This

114   insight led to the finding that Nementin enhances the paralytic effects of

115   organophosphate and carbamate acetylcholinesterase inhibitors not only in *C. elegans*,

116   but in nematode parasites of mammals and plants. We conclude that Nementin is a

117   novel nematode-selective scaffold that may also improve the selectivity of broad-acting

118   pesticides.

Results

26 Compounds Disrupt Multiple *C. elegans* Motor Activities

To identify small molecules that disrupt nematode neuromuscular activity, we designed

a 96-well plate imaging system that measures the egg-laying rate of adult *C. elegans*

hermaphrodites (Fig. 1a-c). The regulation of *C. elegans* egg-laying relies on well-

characterized cholinergic, serotonergic, GABAergic and peptidergic circuits[10], and is

therefore a paragon to assess neuromuscular perturbation by small molecules. To

identify egg-laying stimulators, we screened small molecules in the background of M9

worm buffer[9]. Animals incubated in M9-vehicle control wells lay 1.1 eggs per hour on

average (Fig. 1d; Supplementary Data 1). To identify egg-laying inhibitors, we screened

small molecules in the background of exogenous serotonin plus nicotine that stimulates

animals to lay an average of 7.4 eggs per hour (Fig. 1d; Supplementary Data 1).

A screen of the 486 molecules from our worm-active library[9] revealed 29 stimulators

and 29 inhibitors of egg-laying (Fig. 1e; Supplementary Data 2). We cross-referenced

the list of acute egg-laying modulators to the list of 247 wactive molecules that are lethal

in six-day viability assays[9]. We found a 1.8-fold enrichment of lethality among the

stimulators ($p<0.001$) (Fig. 1f), suggesting that excessive neuromuscular modulation

may lead to death. A 1.4-fold enrichment of lethality among the inhibitors was also

found ($p<0.05$), which is not unexpected given that egg-laying inhibition is certain to be

a consequence of molecules that kill rapidly. A Tanimoto structural similarity cut-off of

0.55 reveals that the 58 egg-laying modulators can parsed into 37 distinct core

scaffolds, consisting of 11 clusters and 26 singletons[11] (Fig. 1g). 10 of the 11 clusters

are composed of molecules with the identical egg-laying phenotype, which validates the phenotypic assignments.

We reasoned that molecules that disrupt multiple motor circuits are more likely to disrupt the parasitic nematode lifecycle. We therefore surveyed the 58 egg-laying modulators for those that disrupt the normal sinusoidal locomotion of wild type *C. elegans*. 26 of the 58 egg-laying modulators (45%) induce at least one locomotory phenotype such as convulsions, coiling, and paralysis (Supplementary Table 1; Fig. 1g; Supplementary Movies 1-6; Supplementary Data 3). The molecules that induce slow movement and paralysis are dominated (88%) by molecules that inhibit egg-laying (Supplementary Table 1). This is consistent with the idea that the egg-laying inhibitor class may be enriched with acutely lethal molecules. By contrast, the molecules that induce convulsions, shaking, and coiling are dominated (80%) by those that enhance egg-laying (Supplementary Table 1).

Nementin-1 is Effective Against Multiple Parasitic Nematodes

To prioritize the 26 neuromodulatory molecules, we eliminated those from consideration that: i) are cytotoxic to human HEK293 cells; ii) induce developmental defects in zebrafish; iii) have lackluster lethality against the five nematodes previously assayed; iv) induce subtle locomotory phenotypes (reversal-defective), or v) were closely related to the commercial nematicide fluopyram[9] (Supplementary Table 1; Supplementary Data 4). This left worm-active (wact) molecules 10, 13, 15, 55, 120, 128, and 444 for further consideration (Fig. 2a; Supplementary Table 1).

We further assessed the activity of our seven hits against seven nematode species

from three phylogenetic clades (Fig. 2b). Four of these seven species are parasites of

mammals. We also analyzed (or reanalyzed) the molecules' activity against the non-

target models *D. rerio* fish, HEK293 cells, and *Arabidopsis thaliana* plants (Fig. 2b;

Supplementary Fig. 1). We prioritized wact-55 because it arguably demonstrates the

best combination of broad nematode activity and selectivity. SciFinder-based literature

searches failed to reveal prior annotation of nematicidal activity for wact-55's alkyl

phenylpiperidine core scaffold[12].

A detailed temporal analysis showed that within 90 seconds of exposure of 60 μM

wact-55, *C. elegans* exhibits a hyperactive nose movement phenotype (Fig. 2c). The

hyperactive phenotype dissipates at the expense of spastic convulsions over the course

of 2 hours (Fig. 2c). By 24 hours, 100% of the adult animals incubated in wact-55 are

dead (non-moving and/or disintegrating; three trials, n = 18 animals per trial). For

reasons outlined below, we renamed the wact-55 molecule 'Nementin-1' (nematode

enhancer of neurotransmission-1).

To determine whether Nementin-1 might have a canonical mechanism-of-action

(MOA), we tested whether any of the *C. elegans* mutant strains that resist the effects of

seven popular anthelmintics also resist the lethal effects of Nementin-1. All drug-

resistant strains remain sensitive to Nementin-1 (Supplementary Fig. 2). Furthermore,

we previously showed that *C. elegans* cannot be easily mutated to resist Nementin-1's

lethality; 290,000 randomly mutated genomes failed to yield wact-55-resistant mutants[9].

These results suggest that: i) Nementin-1 does not share an MOA with canonical

anthelmintics, ii) Nementin-1's MOA may not be limited to a single protein target, and,

iii) genetic resistance to Nementin-1 may be difficult to achieve in the field. All of these

properties make Nementin-1 an attractive hit for further investigation.

Nementin Convulsions are Dependent on Dense Core Vesicle Release

We reasoned that mutations in the pathway targeted by Nementin-1 might phenocopy

the motor defects induced by the molecule and provide mechanistic insight. A survey of

the literature yielded eight mutants that share at least some of Nementin-1's

phenotypes, including *acr-2, unc-2, unc-43, unc-58, unc-93, sup-9, sup-10*, and *twk-*

18[13-21] (Supplementary Table 2). We tested the hypothesis that Nementin-1 disrupts the

pathway disrupted by each of these mutants. We did this by asking whether known

suppressors of the phenocopying mutants can suppress Nementin-1's effects (see

Supplementary Table 2). Of the potential suppressing mutants tested, only the *unc-43*

gain-of-function (GF) mutant resisted Nementin-1's convulsions (Fig. 3a). We also found

the converse to be true; *unc-43* reduction-of-function (RF) alleles enhance Nementin-1

convulsions (Fig. 3a). Remarkably, the interaction between the *unc-43* GF and

Nementin-1 is mutual in that Nementin-1 can rescue the paralysis of *unc-43* GF (Fig.

3b). This observation argues against the idea that the *unc-43* GF mutant alters

Nementin-1 absorption or metabolism. These results suggest that Nementin-1 may be

disrupting UNC-43 pathway activity.

UNC-43 is the *C. elegans* ortholog of CaMKII (calcium/calmodulin-dependent protein

kinase II) and is a key negative regulator of dense core vesicle (DCV) release in *C.*

elegans neurons[14,22] (Fig. 3c). In animals with reduced UNC-43 function, neuronal DCV

content is released in excess and is likely responsible for the mutant's convulsive and

constitutive egg-laying phenotypes[13,22]. In *unc-43* GF animals, DCVs accumulate, and

their limited release is likely responsible for the mutant's severely lethargic locomotion[22].

To examine DCV behaviour in response to Nementin-1, we exploited the

neuropeptide reporter (INS-22::GFP) that is packaged into DCVs in cholinergic motor

neurons[22]. Like *unc-43* RF mutants[22], worms treated with either Nementin-1 or a second

analog called Nementin-12 have reduced axonal INS-22::GFP signal ($p<0.05$) (Fig. 3d-

f,h; Supplementary Fig. 3c,d). Coincidentally, INS-22::GFP accumulates in the

pseudocoelomic fluid-scavenging cells, called coelomocytes, of Nementin-treated

animals ( ).

We reasoned that if Nementin disrupts motor behavior by agonizing DCV release,

then disruption of UNC-31 (calcium-dependent activator protein for secretion (CAPS)),

which is required for DCV release[22,23], or UNC-64 syntaxin, which is required for all

vesical fusions[24,25], should suppress the Nementin-1-driven locomotory defects. Indeed,

we find that both the weaker temperature-sensitive *unc-31(e169)*, the *unc-31(e928)* null,

and the canonical *unc-64(e246)* RF mutants suppress the convulsions induced by

Nementin-1 (Fig. 4a). Together, these data indicate that Nementin-induced DCV release

is responsible for convulsions.

Collectively, these results beg the question of whether UNC-43 is likely to be the

physiologically relevant target of Nementin-1. Several observations argue against this.

First, we note that the terminal phenotype of Nementin-1 is death, yet presumptive null

alleles of *unc-43* (*ce685* and *n1186*) are viable[14,22]. Second, if Nementin-1 inhibits UNC-

43, then Nementin-1 should not be able to enhance the convulsion phenotype of *unc-43*

null mutants. We find that Nementin-1 does indeed enhance presumptive *unc-43* null

mutants (*ce685* and *n1186*) (Fig. 3a). Third, if UNC-43 was Nementin-1's sole

physiological target, then *unc-43* GF mutations would have been identified in our

genetic suppressor screens[9], but no resistant mutants arose. Finally, the *unc-43* GF

mutant (and *unc-31* and *unc-64* RF mutants) remain sensitive to the lethal effects of

Nementin-1 (Supplementary Fig. 4). Together, these data argue against the idea that

UNC-43 is Nementin-1's sole physiologically relevant target. Instead, the similarities

between Nementin-1-induced phenotypes and the *unc-43* RF mutants suggest that the

compound targets a component(s) that acts with UNC-43 (Fig. 3c, purple glow).

Nementin-1 Agonizes Cholinergic Signaling

We next investigated whether cholinergic signaling is also required for Nementin-1

activity. We tested whether mutants of the UNC-17 vesicular acetylcholine transporter

(VAChT), which loads synaptic vesicles (SVs) with acetylcholine, or mutants of UNC-13,

which is specifically required for SV release[26,27], could suppress Nementin-1-induced

convulsions. Reduction-of-function mutants of UNC-17 and UNC-13 remained sensitive

to Nementin-1 (Fig. 4a). Incidentally, we found that Nementin-1 could partially rescue

the lethargic locomotion of *unc-17* RF mutants (Fig. 4b). These data show that despite

cholinergic signalling being dispensable for Nementin-1-induced convulsions, Nementin-

1 may nevertheless agonize cholinergic signalling.

We further investigated the idea that Nementin-1 agonizes cholinergic signalling. To

do so, we used the acetylcholinesterase (AChE) inhibitor aldicarb, which is a well-

characterized tool used to investigate perturbations of cholinergic signaling in *C.*

elegans[28]. AChE catabolizes acetylcholine at the neuromuscular junction (NMJ) and its

inhibition increases acetylcholine levels that in turn paralyzes the animal[29] (see Fig. 3c).

We found that Nementin-1 sensitizes animals to the paralytic effects of aldicarb (Fig.

4c), suggesting that Nementin-1 agonizes acetylcholine release at the NMJ. The

relationship between Nementin-1 and aldicarb is synergistic in nature, yielding a zero-

interaction potency (ZIP) δ-score of 31.1 with a δ-score of 64.8 over the most

synergistic set of concentrations (ZIP scores >10 are considered synergistic[30,31] (Fig.

4d,e)). Nementin-1 also enhances the paralysis induced by dichlorvos and trichlorfon,

which are two other AChE inhibitors that are also commercial anthelmintics, by 12.7 and

18.7-fold, respectively (Fig. 4c). By contrast, Nementin-1 fails to enhance the effects of

an AChE-inhibitor on the fruit fly *D. melanogaster* (Fig. 4c). These data provide

additional support for the idea that Nementin-1 agonizes cholinergic signalling and does

so in a nematode-selective manner.

Cholinergic Agonism is a Secondary Effect of DCV Release

Neuropeptides released from DCVs are known to interact with G-protein coupled-

receptors (GPCRs) that signal through G-proteins[32]. In *C. elegans* neurons, the Gα$_q$

protein EGL-30 stimulates DAG production via the EGL-8 and PLC-3 phospholipases[33]

(see Fig. 3c). By contrast, the Gα$_o$ protein GOA-1 reduces DAG abundance by

triggering its phosphorylation by diacylglycerol kinase DGK-1[34]. Neuronal DAG primes

both DCV a. nd SV release via interaction with PKC-1(PKC-ε/η)[35] and UNC-1 [36],

respectively. The bioamines and neuropeptides that are released from the DCVs can

have both paracrine and autocrine activity[37,38]. This circuitry raised the possibility that

Nementin-1-induced DCV release may initiate a positive feedback loop that amplifies

DCV release and agonizes SV release as a secondary consequence (see green and

yellow pathways in Fig. 3c).

We tested this hypothesis in three ways. First, we examined the impact of mutant

components of the 'green' circuit depicted in Fig. 3c. We found that *egl-30* RF, which

reduces DAG production, suppresses Nementin-1 convulsions (Fig. 4a). By contrast,

goa-1 RF and *dgk-1* loss-of-function (LF) mutants, which lack regulation of membrane

DAG, enhance Nementin-1 convulsions. We also found that disruption of PKC-1

reduces Nementin-1 convulsions (Fig. 4a). Second, we asked whether the cell

permeable DAG mimetic phorbol myristate acetate (PMA) enhances Nementin-1

convulsions and found that it did (Fig. 4a). Consistent with the lack of convulsions in

goa-1 RF mutants (not treated with Nementin; Figure 4a), PMA treatment (without

Nementin) does not induce convulsions in wild type animals (Figure 4a). This indicates

that PKC-1-agonism alone cannot severely disrupt motor activity. Finally, we asked

whether Nementin-1's agonism of cholinergic signaling is dependent on the machinery

needed for DCV release. Indeed, we found that the *unc-31(e169)* RF mutation and the

unc-43(n498) GF mutation suppress Nementin-1's agonism of cholinergic signaling (Fig.

4c). Together, these observations are consistent with a model whereby Nementin-1

agonizes DCV release whose contents may stimulate autocrine/paracrine G-protein

signaling, which in turn agonizes both more DCV release and cholinergic signaling.

Nementin Disrupts Parasitic Nematodes of Plants and Animals

We carried out a small structure-activity relationship (SAR) analysis of Nementin

analogs assayed against free-living nematodes, parasitic nematodes, and non-target

models (Supplementary Table 3; Supplementary Data 4). We purchased 18

commercially available analogs of Nementin-1 and found that Nementin-12 was the

most potent inducer of acute motor phenotypes (Supplementary Table 3). We expanded

the SAR by synthesizing 22 analogs of Nementin-12 (see Materials and Methods).

Nementins 1, 12-5, 13 and 14 demonstrated broad anthelmintic activity without inducing

obvious phenotype against non-target organisms. In addition, Nementin analogs were

identified with improved activity in each parasitic nematode model tested. These data

suggest that the Nementin scaffold may be further refined to improve selectivity and

potency.

Given that Nementin-1 synergistically enhances the activity of AChE inhibitors in *C.*

elegans, we tested whether it might act similarly against parasitic nematodes. We tested

Nementin-1's interaction with the organophosphate dichlorvos against *Strongyloides*

ratti, a nematode parasite of mammals. Dichlorvos is approved for anthelmintic use in

mammals[39,40]. Indeed, we find that Nementin-1 synergistically enhances dichlorvos'

activity against *S. ratti* (Fig. 4f,g). Nementin-1 was also found to synergistically paralyze

the plant-parasitic nematode *Meloidogyne hapla* with the carbamate AChE inhibitor

oxamyl, which is approved for field use[39,40] (Fig. 4h,i). The Nementins may therefore

have added benefit against parasitic nematodes when used in combination with

pesticides that are AChE inhibitors.

Discussion

Here, we have developed a *C. elegans*-based pipeline to identify novel anthelmintic small molecules with phylum-selectivity. The pipeline was designed to yield broad-acting anthelmintics that acutely affect nematode behaviour. The strategy yielded four small molecules that exhibit phylum-selectivity within the limited range of organisms tested. We focused on one of these molecules, called Nementin-1, which induces *C. elegans* hyperactivity within seconds of exposure, followed by spastic convulsions minutes later. Several Nementin analogs affect parasitic nematodes from distinct clades at concentrations that fail to elicit obvious phenotypes from non-target organisms. Despite significant effort, we were not able to generate *C. elegans* mutants that resist the lethal effects of Nementin, suggesting that resistance in the wild may be difficult to achieve.

Our data support a model whereby Nementin exposure initiates DCV release, which in turn agonizes SV release that promotes cholinergic signaling. Nementin-induced convulsions depend on DCV release but not cholinergic signaling, indicating that agonized DCV release is key to Nementin's ability to disrupt motor behavior. How Nementin agonizes DCV release remains unclear. Despite the phenotypic similarities between Nementin exposure and *unc-43* RF mutants, our data suggests that Nementin is unlikely to inhibit UNC-43 in any canonical fashion. However, because of CaMKII's complexity and its many isoforms[14,41], we cannot rule out the possibility that Nementin interacts with CaMKII in a complex and perhaps tissue-specific manner.

Convulsions in *C. elegans* and other animals are thought to manifest from an

imbalance of excitatory and inhibitory neurotransmitter signals, termed E/I imbalance [42].

Gain-of-function mutations in motor neuron cation channels like the ACR-2

acetylcholine receptor or the UNC-2 voltage-gated calcium channel simultaneously

increase excitatory cholinergic signals and decrease inhibitory GABAergic signals that

are relayed to muscles. This in turn generates an excitation-dominated E/I imbalance

that culminate in convulsions [18,19,38]. Several parallels exist between Nementin-treated

wild-type worms and either the *unc-2* and/or *acr-2* gain-of-function mutants including

hyperactive locomotion, convulsions, a phenotypic dependence on UNC-31 CAPS, and

aldicarb hypersensitivity [18,19,38]. Nementin-induced convulsions are independent of

ACR-2, UNC-2, and robust cholinergic signaling. We therefore deduce that the likely E/I

imbalance induced by Nementin must originate from signals that are released from

neuronal dense core vesicles.

Nementin has several features that make it an attractive anthelmintic for further

development. First, *in vitro* tests demonstrate that it exhibits potentially broad-spectrum

activity while maintaining phylogenetic selectivity. Such a feature is key to the

development of an environmentally safe anthelmintic. Second, Nementin analogs

exhibit an *in vitro* potency that is comparable to several commercial anthelmintics. Third,

the Nementin synthesis route is relatively simple and uses inexpensive starting

materials ('Chemistry' in Supplementary Materials)[43]. Finally, Nementin has the potential

to reduce the amount of non-selective pesticides that are notoriously released into the

environment.

Despite the toxicity concerns over indiscriminate activity, many AChE inhibitors such

as trichlorfon, dichlorvos, coumaphos, aldicarb, fosthiazate, chlorpyrifos and oxamyl

remain approved for nematicidal field use as of 2021 because of limited control

options[39] (see the FDA Approved Animal Drug Products at https://tinyurl.com/by83ks98,

the US Environmental Protection Agency at https://tinyurl.com/3n3kf34n; and the US

Geological Survey pesticide study at https://tinyurl.com/bp5h5em4). Because of

Nementin's selectivity and synergistic effects, using it in combination with AChE

inhibitors has the potential to maintain nematode-selectivity whilst simultaneously

reducing the amount of non-selective pesticide applied to the environment.

373
374   **Materials and Methods**

375
376   *Materials and Methods are provided in the Supplementary Information for this initial

377   submission.

References

Rashid, M. *et al.* A systematic review on modelling approaches for economic losses studies caused by parasites and their associated diseases in cattle. *Parasitology* **146**, 129-141, doi:10.1017/S0031182018001282 (2019).

Geerts, S. & Gryseels, B. Drug resistance in human helminths: current situation and lessons from livestock. *Clin Microbiol Rev* **13**, 207-222, doi:10.1128/CMR.13.2.207 (2000).

Chitwood, D. J. Research on plant-parasitic nematode biology conducted by the United States Department of Agriculture-Agricultural Research Service. *Pest Manag Sci* **59**, 748-753, doi:10.1002/ps.684 (2003).

Affairs, U. N. D. o. E. a. S. Consolidated list of products whose consumption and / or sale have been banned, withdrawn, severely restricted or not approved by Governments. (New York, 2009).

Godfray, H. C. & Garnett, T. Food security and sustainable intensification. *Philos Trans R Soc Lond B Biol Sci* **369**, 20120273, doi:10.1098/rstb.2012.0273 (2014).

Cole, M. B., Augustin, M. A., Robertson, M. J. & Manners, J. M. The science of food security. *NPJ Sci Food* **2**, 14, doi:10.1038/s41538-018-0021-9 (2018).

Geary, T. G., Klein, R. D., Vanover, L., Bowman, J. W. & Thompson, D. P. The nervous systems of helminths as targets for drugs. *J Parasitol* **78**, 215-230 (1992).

Holden-Dye, L. & Walker, R. J. Anthelmintic drugs and nematicides: studies in Caenorhabditis elegans. *WormBook*, 1-29, doi:10.1895/wormbook.1.143.2 (2014).

9   Burns, A. R. *et al.* Caenorhabditis elegans is a useful model for anthelmintic

discovery. *Nat Commun* **6**, 7485, doi:10.1038/ncomms8485 (2015).

10   Schafer, W. F. Genetics of egg-laying in worms. *Annu Rev Genet* **40**, 487-509,

doi:10.1146/annurev.genet.40.110405.090527 (2006).

11   Bajusz, D., Racz, A. & Heberger, K. Why is Tanimoto index an appropriate

choice for fingerprint-based similarity calculations? *J Cheminform* **7**, 20,

doi:10.1186/s13321-015-0069-3 (2015).

12   Somerville, A. N. SciFinder Scholar. *Journal of Chemical Education* **75**, 959-960,

doi:doi.org/10.1021/ed075p959 (1998).

13   Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71-94 (1974).

14   Reiner, D. J., Newton, E. M., Tian, H. & Thomas, J. H. Diverse behavioural

defects caused by mutations in Caenorhabditis elegans unc-43 CaM kinase II.

Nature **402**, 199-203, doi:10.1038/46072 (1999).

15   Wei, A., Jegla, T. & Salkoff, L. Eight potassium channel families revealed by the

C. elegans genome project. *Neuropharmacology* **35**, 805-829, doi:10.1016/0028-

3908(96)00126-8 (1996).

16   de la Cruz, I. P., Levin, J. Z., Cummins, C., Anderson, P. & Horvitz, H. R. sup-9,

sup-10, and unc-93 may encode components of a two-pore K+ channel that

coordinates muscle contraction in Caenorhabditis elegans. *J Neurosci* **23**, 9133-

9145 (2003).

17   Greenwald, I. S. & Horvitz, H. R. unc-93(e1500): A behavioral mutant of

Caenorhabditis elegans that defines a gene with a wild-type null phenotype.

Genetics **96**, 147-164 (1980).

18   Huang, Y. C. *et al.* Gain-of-function mutations in the UNC-2/CaV2alpha channel

425   lead to excitation-dominant synaptic transmission in Caenorhabditis elegans.

426   *Elife* **8**, doi:10.7554/eLife.45905 (2019).

427   19   Jospin, M. *et al.* A neuronal acetylcholine receptor regulates the balance of

428   muscle excitation and inhibition in Caenorhabditis elegans. *PLoS Biol* **7**,

429   e1000265, doi:10.1371/journal.pbio.1000265 (2009).

430   20   Salkoff, L. *et al.* Potassium channels in C. elegans. *WormBook*, 1-15,

431   doi:10.1895/wormbook.1.42.1 (2005).

432   21   Kasap, M. & Dwyer, D. S. Clozapine, nimodipine and endosulfan differentially

433   suppress behavioral defects caused by gain-of-function mutations in a two-pore

434   domain K(+) channel (UNC-58). *Neurosci Res*, doi:10.1016/j.neures.2020.07.006

435   (2020).

436   22   Hoover, C. M. *et al.* A novel CaM kinase II pathway controls the location of

437   neuropeptide release from Caenorhabditis elegans motor neurons. *Genetics* **196**,

438   745-765, doi:10.1534/genetics.113.158568 (2014).

439   23   Speese, S. *et al.* UNC-31 (CAPS) is required for dense-core vesicle but not

440   synaptic vesicle exocytosis in Caenorhabditis elegans. *J Neurosci* **27**, 6150-

441   6162, doi:10.1523/JNEUROSCI.1466-07.2007 (2007).

442   24   Hammarlund, M., Watanabe, S., Schuske, K. & Jorgensen, E. M. CAPS and

443   syntaxin dock dense core vesicles to the plasma membrane in neurons. *J Cell*

444   *Biol* **180**, 483-491, doi:10.1083/jcb.200708018 (2008).

25   Laurent, P. *et al.* Genetic dissection of neuropeptide cell biology at high and low

activity in a defined sensory neuron. *Proc Natl Acad Sci U S A* **115**, E6890-

E6899, doi:10.1073/pnas.1714610115 (2018).

26   Lackner, M. R., Nurrish, S. J. & Kaplan, J. M. Facilitation of synaptic transmission

by EGL-30 Gqalpha and EGL-8 PLCbeta: DAG binding to UNC-13 is required to

stimulate acetylcholine release. *Neuron* **24**, 335-346, doi:10.1016/s0896-

6273(00)80848-x (1999).

27   Alfonso, A., Grundahl, K., Duerr, J. S., Han, H. P. & Rand, J. B. The

Caenorhabditis elegans unc-17 gene: a putative vesicular acetylcholine

transporter. *Science* **261**, 617-619, doi:10.1126/science.8342028 (1993).

28   Nguyen, M., Alfonso, A., Johnson, C. D. & Rand, J. B. Caenorhabditis elegans

mutants resistant to inhibitors of acetylcholinesterase. *Genetics* **140**, 527-535

(1995).

29   Cambon, C., Declume, C. & Derache, R. Effect of the insecticidal carbamate

derivatives (carbofuran, pirimicarb, aldicarb) on the activity of

acetylcholinesterase in tissues from pregnant rats and fetuses. *Toxicol Appl*

Pharmacol **49**, 203-208, doi:10.1016/0041-008x(79)90242-4 (1979).

30   Ianevski, A., Giri, A. K. & Aittokallio, T. SynergyFinder 2.0: visual analytics of

multi-drug combination synergies. *Nucleic Acids Res* **48**, W488-W493,

doi:10.1093/nar/gkaa216 (2020).

31   Yadav, G. & Ganguly, S. Structure activity relationship (SAR) study of

benzimidazole scaffold for different biological activities: A mini-review. *Eur J Med*

Chem **97**, 419-443, doi:10.1016/j.ejmech.2014.11.053 (2015).

32   Koelle, M. R. Neurotransmitter signaling through heterotrimeric G proteins:

insights from studies in C. elegans. *WormBook* **2018**, 1-52,

doi:10.1895/wormbook.1.75.2 (2018).

33   Robatzek, M. & Thomas, J. H. Calcium/calmodulin-dependent protein kinase II

regulates Caenorhabditis elegans locomotion in concert with a G(o)/G(q)

signaling network. *Genetics* **156**, 1069-1082 (2000).

34   Miller, K. G., Emerson, M. D. & Rand, J. B. Goalpha and diacylglycerol kinase

negatively regulate the Gqalpha pathway in C. elegans. *Neuron* **24**, 323-333,

doi:10.1016/s0896-6273(00)80847-8 (1999).

35   Sieburth, D., Madison, J. M. & Kaplan, J. M. PKC-1 regulates secretion of

neuropeptides. *Nat Neurosci* **10**, 49-57, doi:10.1038/nn1810 (2007).

36   Richmond, J. E., Davis, W. S. & Jorgensen, E. M. UNC-13 is required for

synaptic vesicle fusion in C. elegans. *Nat Neurosci* **2**, 959-964,

doi:10.1038/14755 (1999).

37   Steuer Costa, W., Yu, S. C., Liewald, J. F. & Gottschalk, A. Fast cAMP

Modulation of Neurotransmission via Neuropeptide Signals and Vesicle Loading.

Curr Biol **27**, 495-507, doi:10.1016/j.cub.2016.12.055 (2017).

38   Stawicki, T. M., Takayanagi-Kiya, S., Zhou, K. & Jin, Y. Neuropeptides function in

a homeostatic manner to modulate excitation-inhibition imbalance in C. elegans.

PLoS Genet **9**, e1003472, doi:10.1371/journal.pgen.1003472 (2013).

39   Nixon, S. A. *et al.* Where are all the anthelmintics? Challenges and opportunities

on the path to new anthelmintics. *Int J Parasitol Drugs Drug Resist* **14**, 8-16,

doi:10.1016/j.ijpddr.2020.07.001 (2020).

40   Abdel-Sattar, M., Haikal, A. M. & Hammad, S. E. Meloidogyne incognita

population control and nutritional status and productivity of Thompson seedless

grapevines managed with different treatments. *PLoS One* **15**, e0239993,

doi:10.1371/journal.pone.0239993 (2020).

41   Myers, J. B. *et al.* The CaMKII holoenzyme structure in activation-competent

conformations. *Nat Commun* **8**, 15742, doi:10.1038/ncomms15742 (2017).

42   Shao, L. R., Habela, C. W. & Stafstrom, C. E. Pediatric Epilepsy Mechanisms:

Expanding the Paradigm of Excitation/Inhibition Imbalance. *Children (Basel)* **6**,

doi:10.3390/children6020023 (2019).

43   Ertl, P. & Schuffenhauer, A. Estimation of synthetic accessibility score of drug-

like molecules based on molecular complexity and fragment contributions. *J

Cheminform* **1**, 8, doi:10.1186/1758-2946-1-8 (2009).

Acknowledgements:

We are grateful for the strains given to us by the *Caenorhabditis* Genetics Center

(University of Minnesota), Marie-Anne Félix (IBENS, Paris), Ken Miller (Oklahoma

Medical Research Foundation), and Mei Zhen and Wesley Hung (Lunenfeld-

Tanenbaum Research Institute, Toronto). Many thanks to Dr. Doug Colwell and Dawn

Gray at the Lethbridge and Agri-Food Canada Research Centre for supplying faeces

from infected calves for *Cooperia oncophora* egg collection.

Ethics Statement

DK: Experiments on *D. immitis* and *N. brasiliensis* were performed in the laboratories of Bayer Animal Health GmbH (Monheim, Germany) in accordance with the local Animal Care and Use Committee and governmental authorities. JK: The generation of *N. americanis*, *T. muris*, S. ratti, and *H. polygyrus* and in vitro studies were carried out at the Swiss Tropical Institute (Basel, Switzerland), in accordance with both cantonal (license no. 2070) and Swiss national regulations on animal experimentation. JD: All zebrafish experiments were performed in compliance with any relevant ethical regulations, specifically following an institutionally reviewed and approved animal use protocol as well as the policies and guidelines of the Canadian Council on Animal Care and Animals for Research Act of Ontario.

Funding:

Research in the I.Z. lab is partially supported by the United States Department of Agriculture, Agricultural Research Service. P.J.R. is supported by CIHR project grants (313296 and 173448). P.J.R. is a Canada Research Chair (Tier 1) in Chemical Genetics.

Author Contributions

Conceptualization: PJR, SH

Methodology: SH, JJK, K-L C, AA, MK, CH, JP, JRV, MG, JS, VW, BMP, EMR, ASV,

Investigation: SH, JJK, AB, MK, CH, JP, C'DA, Y-H K, JRV, MG, JS, VW, BMP, EMR, ADS

Visualization: PJR, SH

Funding acquisition: PJR

Project administration: PJR

Supervision: PJR, JG, IS, SRC, JJD, CMY, JK, IZ, ML

Writing – original draft: PJR, SH, JJK, K-LC, AA, SRC, DK, JK

Writing – review & editing: PJR, SH, AB, JJK, SC, DK, JK, IZ, ML

Competing interests:

Mention of trade names or commercial products in this publication is solely for the

purpose of providing specific information and does not imply recommendation or

endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity

provider and employer. S.H., J.K., A.R.B., K-L.C., J.P., M.L. and P.J.R. have patents

pending related to Nementin.

Data and materials availability:

Original data for all analyses presented are included in Supplementary Data 1-S4.

Original images for analyses can be made available per request through the

corresponding author P.J.R.

Supplementary Materials

Materials and Methods

Supplementary Figures 1-4

Supplementary Table 1-3

Supplementary Movies 1 to 6 (description)

Supplementary Data 1 to S4 (description)

Fig. 1 The Identification of 58 Small Molecules that Modulate the *C. elegans* Nervous System.

(**a** and **b**) Two perspectives of the '2020 Imager' showing one 96-well plate in the

imager holder (arrow). (**c**) Representative half-well images of vehicle controls in the

basal (M9-only) background (left) and the 'stimulatory cocktail' of 12.3 mM serotonin

creatine sulfate and 7.7 mM nicotine (right) after 1 hour. A blue arrow points to an adult

hermaphrodite and a red arrow points to a pair of embryos. (**d**) The distribution of

counts of eggs/worm/hour for the basal background (blue) and the stimulatory

background (red) from the indicated well-coordinates over 22 experimental replicates

(96-well plates). The mean and SEM is shown. (**e**) A screen of the worm active

(wactive) library yields 29 inhibitors (purple) and 29 stimulators (green) of the *C.*

elegans egg-laying rate. Mean Egl rate and SEM is shown. (**f**) Venn diagram showing

the overlap of the wactive molecules that are lethal (at 60 μM) with those that modulate

egg-laying among the 486 molecules assayed in both assays. (**g**) A structure similarity

network of the 58 egg-laying modulators. Each node represents a numbered wactive

molecule. Connecting lines indicate shared structural similarity. Additional information is

provided in Supplementary Table 1.

Fig. 2 The motor-centric pipeline yields convulsing-inducing Nementin-1. (**a**)

Structures of the 7 prioritized neuromodulators. (**b**) Bioactivity of the seven prioritized

molecules. Data are EC50s except for the *A. thaliana* greening assay, which is the

lowest concentration that yields a discernible difference from control. (**c**) A time-course

analysis of 60 *μ*M Nementin-1-induced locomotory phenotypes. Data are the mean of

biological triplicate measurement of 18 animals per each time point. The standard

deviation is shown.

Fig. 3 Nementin agonizes dense core vesicle release. (a) Convulsions induced by

Nementin-1 (Nem-1) (60 μM) in the indicated genetic background after 80 minutes. The

mean of three biological replicates with 18 animals per trial is shown. (**b**) The mean

percentage of animals of the indicated genotype (three trials with ~150 animals per trial)

that locomote to zone 2 over 180 minutes after being placed in the centre of a plate (see

schematic on right). For A and B, black and purple asterisks represent $p<0.05$ and

$p<0.001$, Chi-square test of independence with Bonferroni correction relative to control

(open circle); the standard error of the mean (SEM) is shown. (**c**) A schematic of

neuronal signaling pathways relevant to Nementin activity. Purple glow indicates area of

possible Nementin action; the green glow indicates how DCV neurotransmitter cargo

may activate $G\alpha_q$ and $G\alpha_o$ signaling pathways; the yellow glow indicates how the

secondary messenger diacylglycerol (DAG) can also promote synaptic vesicle release.

(**d**) Schematic of adult (strain KG4247) expressing INS-22::GFP (green dots; packaged

into DCVs) and secreted mCherry (fuchsia) in cholinergic motor neurons (cell bodies

are oval). mCherry is constitutively secreted and taken up by the coelomocytes (three

pairs of pink objects in the animal). Box indicates the area shown in E and F. Anterior is

the left and dorsal is up. (**e** and **f**) Images of the midbody region of control and 60 μM

Nementin-1-treated KG4247 animals after 4 hours. dc, dorsal cord; vc, ventral cord; co,

coelomocytes; cb, cell bodies. (**g**) Quantification of the midbody coelomocytes (ratio of

measured GFP/RFP) and dorsal cord fluorescence (relative to background tissues) of

the animal. AU, arbitrary units., 1-way ANOVA with Dunnett correction for multiple

606   comparisons; ** p<0.01; *** p<0.001; SEM is shown. Other regions of the animal are

607   quantified in Supplementary Fig. 3.

C

			AChEi EC50 (μM)		
species	genetic background	AChEi	control	+ 15 μM Nem-1	fold change
C. elegans	WT	trichlorfon	63.5	3.4	18.7
C. elegans	WT	dichlorvos	244.8	19.3	12.7
C. elegans	WT	aldicarb	565.5	61.2	9.2
C. elegans	*unc-31(e169)*	aldicarb	839.1	469.8	1.8
C. elegans	*unc-43(n498)* (GF)	aldicarb	926.5	852.6	1.1
D. melanogaster	WT	oxamyl	14.8	17.8	1.2

global ZIP = 31.1; max ZIP = 64.8   global ZIP = 24.3; max ZIP = 42.0   global ZIP = 25.1; max ZIP = 40.1

Fig. 4 Nementin enhances AChE inhibitor activity via agonism of DCV release. (a)

Convulsions induced after 80 minutes of exposure to Nementin-1 (Nem-1) (60 μM) or

vehicle control in the background of the indicated mutant gene, or in the background of

24 hours of preincubation with the DAG memetic PMA. (**b**) The ability of wild type (WT)

or *unc-17* mutant animals to productively locomote to zone 2 over 180 minutes after

being placed in the centre of a plate (see schematic in Fig. 3b) in the indicated

Nementin concentration. For both A and B, black asterisk p<0.01, red asterisk p<0.001,

Bonferroni corrected Chi-square test of independence relative to control (open circle);

SEM is shown. (**c**) Acetylcholinesterase inhibitor (AChEi) EC50s for different species

and genetic backgrounds. Oxamyl is the only AChEi tested that exhibited activity

against flies. Data are the mean of 3 biological replicates scoring 18 animals per

condition. (**d-f**) Double dose-response analyses of Nementin-1 (y-axis) vs the indicated

AChEi (x-axis) in the indicated nematode species. The scale for the percentage of the

population paralyzed applies to d-f. (**g-i**) The corresponding zero interaction potency

(ZIP) synergy score heatmaps for the double dose-response analyses presented in d-f.

Max ZIP-score is defined as the maximal ZIP-score obtained from a combination of

three concentrations of each compound in the double dose response. Shading

highlights the topology of the plot and differs between g-i. N=3 trials for *elegans* with

n=18 within each trial, N=6 trials for *ratti* with n≥30 within each trial, N≥3 trials for *hapla*

with n≥15 within each trial.