Humans prioritize walking efficiency or walking stability based on environmental risk

Ashwini Kulkarni[1&], Chuyi Cui[1&], Shirley Rietdyk[1&], Satyajit Ambike[1¶*]

[1]Department of Health and Kinesiology, Purdue University, West Lafayette, IN

*Corresponding author

E-mail: sambike@purdue.edu (SA)

[¶]These authors contributed equally to this work.

[&]These authors contributed equally to this work.

Abstract

In human gait, the body's mechanical energy at the end of one step is reused to achieve forward progression during the subsequent step, thereby reducing the required muscle work. During the single stance phase, humans rely on the largely uncontrolled passive inverted pendular motion of the body to perpetuate forward motion. These passive body dynamics, while improving walking efficiency, also indicate that lower passive dynamic stability in the anterior direction since the individual will be less able to withstand a forward external perturbation. Here we test the novel hypothesis that humans manipulate passive anterior-posterior (AP) stability via active selection of step length to either achieve energy-efficient gait or to improve stability when it is threatened. We computed the AP margin of stability, which quantifies the passive dynamic stability of gait, for multiple steps as healthy young adults (N=20) walked on a clear and on an obstructed walkway. Participants used passive dynamics to achieve energy-efficient gait for all but one step; when crossing the obstacle with the leading limb, AP margin of stability was increased. This increase indicated caution to offset the greater risk of falling after a potential trip. Furthermore, AP margin of stability increased while approaching the obstacle, indicating that humans proactively manipulate the passive dynamics to meet the demands of the locomotor task. Finally, the step length and the center of mass motion co-varied to maintain the AP margin of stability for all steps in both tasks at the specific values for each step. We conclude that humans actively regulate step length to maintain specific levels of passive dynamic stability for each step during unobstructed and obstructed gait.

1. Introduction

Gymnasts exploit the mechanical properties of their bodies to achieve remarkable feats. While performing a triple axel, for example, an ice-skater holds her arms close to or farther from her body to modulate her angular velocity. High velocity during the flight phase allows for more turns [1], whereas lower velocity close to landing improves safety [2].

The ability and propensity to adapt the body's mechanical properties to facilitate motion is not limited to trained sportspersons; it is a feature of the walking patterns of most adults. On clear and level walkways, adults use a characteristic step length to recycle the kinetic energy at the end of a step to rotate the body over the new stance foot. Part of the forward progression of the body is thus achieved passively, which improves walking efficiency [3, 4]. However, if the environment poses a threat to stability, the passive forward motion can be altered to improve gait stability rather than efficiency. Such adaptation is suggested by the difference in the behaviors of young and older adults. While stepping over an obstacle, older adults rely less on the passive forward motion compared to young adults [5], since age-related declines in strength and coordination make it more difficult to recover from a trip. In other words, since, after a trip, forward passive motion makes a forward fall more likely, older adults proactively improve their passive stability by reducing this motion while crossing obstacles.

This tradeoff between stability and efficiency has been supported by the study of the margin of stability in the anterior-posterior direction (MOS_{AP}; abbreviations in Table 1). MOS_{AP} has been used to identify differences in the recruitment of passive dynamics across age groups [5-9], in pathological populations [10-12], and in dual tasks [13]. MOS_{AP}, based on the inverted pendulum model of gait, and usually computed at heel contact, is the distance from the anterior boundary of the base of support (BOS; the leading heel) to the extrapolated center of mass (XcoM), which reflects the center of mass (CoM) state [14, 15]. The condition XcoM ahead of the BOS boundary at heel contact indicates efficient gait: the body has sufficient kinetic energy to passively rock over the new stance ankle. More anterior location of XcoM indicates greater forward passive motion and greater efficiency. At the same time, a forward perturbation during the swing phase would accentuate the greater forward passive motion,

making a forward fall more likely. Therefore, a more anterior location of the XcoM at heel

contact is also interpreted as lower passive dynamic stability in the anterior direction [5, 13].

Here, we advance in three ways our understanding of the role of passive dynamics

during gait based on MOS_{AP}. First, we demonstrate that adaptation in passive dynamics in

response to environmental threats to stability is apparent in young adults as well. We show

that young adults increase passive dynamic stability while crossing an obstacle compared to

unobstructed walking. Second, we demonstrate that the shift in passive dynamic stability

occurs in the steps leading up to the obstacle. This novel finding is consistent with the notion

that motor behavior usually does not change instantaneously, but evolves over characteristic

times determined in part by the inertia of the body [16]. Third, we demonstrate that humans

modulate step length to control MOS_{AP}. We show that MOS_{AP} is similar for the steps of

unobstructed gait, and it is maintained at different values for specific steps while approaching

and crossing an obstacle. Hof [15] showed that a consistent MOS_{AP} yields a stable walking cycle

in a mathematical model, and hypothesized that humans would benefit from this strategy as

well [17]. It has been suggested that the variability rather than the average value of MOS better

represents passive dynamic stability of human gait [18]. In the anterior-posterior (AP) direction

specifically, indirect evidence for this hypothesis is provided by the fact that persons with

Multiple Sclerosis (MS) exhibit greater MOS_{AP} variability compared to healthy controls [11], and

within the group of persons with MS, fallers show greater MOS_{AP} variability than non-fallers

[10].

We sought direct evidence for the control of MOS_{AP} in the unobstructed and obstructed

gait of healthy young adults. Since the MOS_{AP} is a function of step length and XcoM, we

hypothesized that the central nervous system responds to changes the XcoM with a

corresponding correction in step length so that the MOS_{AP} itself is relatively invariant at each

heel contact. However, the specific values of MOS_{AP} that are maintained via the covariation in

XcoM and step length are different at various steps during obstructed gait.

We employed the uncontrolled manifold (UCM) method [19] to test this hypothesis. For

each heel contact, we computed the synergy index that quantifies the covariation between

XcoM and step length. A positive synergy index indicates that MOS_{AP} was actively controlled,

i.e., stabilized at the specific across-trial mean MOS_{AP} for that step. A higher value indicates a

stronger synergy or higher stability of MOS_{AP}. We emphasize that stability of MOS_{AP} is different

from the stability of gait. MOS_{AP} is the measure of *passive* dynamic gait stability. In contrast, the

synergy index indicates the efficacy of *active* control at stabilizing or maintaining MOS_{AP} at a

specific value for a given step.

We computed MOS_{AP} and the synergy index for multiple steps for unobstructed gait

trials and for trials where participants approached an obstacle, stepped over it, and continued

walking till the end of the walkway. We hypothesized a task by step interaction for MOS_{AP} (H1).

The MOS_{AP} will not be different across tasks (obstructed and unobstructed) for the steps at the

start and end of the walkway, but the MOS_{AP} will be higher for the approach steps 1-2 steps

before the obstacle and for the crossing steps. These changes will indicate prioritization of

safety over efficiency [5, 6, 20]. Next, we hypothesized that the synergy index will be greater

than zero, indicating that the step length and the XcoM co-vary to stabilize MOS_{AP} for all steps

in both tasks (H2). Finally, we hypothesized a task by step interaction for the synergy index

(H3). The synergy index will not be different across tasks for the steps at the start and end of

the walkway, but the synergy index will be lower while crossing an obstacle placed in the

middle of the walkway. The lower synergy index will reflect the greater motor demands

associated with the crossing steps; larger muscle activations required for stepping over the

obstacle [21] will be associated with greater noise [22], which will make stabilization more

difficult.

Table 1. Abbreviations.

Anterior-posterior	AP
Medial-lateral	ML
Foot placements	fp
Center of mass	CoM
Center of mass velocity	V_{CoM}

Leg length	l
Acceleration due to gravity	g
Base of support	BOS
Extrapolated center of mass	XcoM
Margin of stability in anterior-posterior direction	MOS_{AP}
Uncontrolled manifold	UCM
Orthogonal manifold	ORT
Synergy index	ΔV
Synergy index z transformed	ΔV_Z
Variance along the uncontrolled manifold	V_{UCM}
Variance along the orthogonal manifold	V_{ORT}

2. Materials and Methods

2.1 Participants

Twenty-six healthy young adults participated in the study. We excluded six participants due to poor kinematic tracking. Data from 20 participants (14 women, 22.3 ± 3.7 years, 1.7 ± 0.1 m, 66.9 ± 14.6 kg) were used for analysis. All participants walked without aid, had no orthopedic, neuromuscular, or dementia disorders, and were independent in daily activities. Vision was normal or corrected-to-normal. The study was approved by Purdue University's Institutional Review Board, and all participants provided written informed consent (Protocol number: IRB-2021-331).

2.2 Equipment and Procedures

We assessed leg dominance using the Waterloo Footedness Questionnaire— Revised [23]. Participants walked at their self-selected speed on a 6.0 m walkway and stepped over an obstacle when present (Fig. 1A). The obstacle was 100 cm wide $\times$ 0.4 cm deep. The height of the obstacle was scaled to 25% of the participant's leg length. The obstacle was made of black Masonite and designed to tip if contacted. The starting position was determined for each

participant such that they took five steps before reaching the obstacle, crossed the obstacle

naturally with the right leg first and stopped three to four steps later (Fig. 1A).

Participants first performed 20 trials without an obstacle (no obstacle task), followed by

20 trials of walking with an obstacle (obstacle-crossing task). We collected kinematic data at

100 Hz with a motion capture system (Vicon Vero, Oxford, UK) with marker clusters placed

bilaterally on the lower back, thigh, shank, and foot. We digitized the joint centers and

posterior aspect of the heels to identify their locations relative to the marker clusters. We also

digitized the top edge of the obstacle to identify its position.

(A)

(B)

Figure 1. Experimental task and basic definitions. Illustration of foot placements while

approaching (fp_{-4} to fp_{-1}), crossing (fp_{+1} and fp_{+2}) and after crossing (fp_{+3}) the obstacle, and of

steps while approaching ($Step_{-3}$ to $Step_{-1}$), crossing ($Step_0$ and $Step_{+1}$) and after crossing ($Step_{+2}$)

the obstacle **(A)**. Definitions of the extrapolated center of mass (XcoM) and margin of stability,

MOS_{AP}. Step length, XcoM and MOS_{AP} are computed at the moment of lead heel contact in a

coordinate frame located where the rear heel contacted the ground **(B)**.

2.3 Analysis

Across all trials and all participants, the obstacle was contacted and tipped 12 times out of 3360 trials (0.4 % of trials). We did not include the contact trials in our analysis. Furthermore, some trials were discarded due to poor kinematic tracking. To have the same number of trials for all participants and tasks, we selected 15 trials with good kinematic data. Fifteen trials are sufficient for reliable quantification of the synergy variables [24]. We filtered all kinematic data using a zero-lag, 4th order, low-pass Butterworth filter with a cut-off of 7 Hz. We identified seven foot placements (Fig. 1A) using the AP position of the heel [25]. We quantified spatiotemporal gait parameters and margin of stability at heel contact at the seven foot placements (fp_{-4} to fp_{+3}) and six steps ($Step_{-3}$ to $Step_{+2}$; Fig. 1A). Step length was defined as the distance between two consecutive heel contacts. CoM position was computed as the centroid of the triangle formed by the left and right anterior superior iliac spines and the center of the left and right posterior superior iliac spines [26]. CoM velocity was obtained by differentiating the CoM position data. The extrapolated center of mass (XcoM) was calculated in the sagittal plane as [14]:

$$XcoM = CoM + \frac{V_{CoM}}{\sqrt{\frac{g}{l}}},$$

where CoM is the anterior-posterior CoM position, V_{CoM} is the anterior-posterior CoM velocity, g is the acceleration due to gravity, and l is the participant's leg length (Fig. 1B). Leg length was calculated as the sagittal-plane distance between the CoM and the ankle of the limb that contacted the ground. We used the average of the leg length values obtained from the 15 trials for each step to compute the XcoM for that step [18]. We computed MOS_{AP} at the instant of leading heel contact in a coordinate frame fixed at the location of the rear heel contact (Fig. 1B):

$$MOS_{AP} = Step\ length - XcoM, \qquad\qquad (1)$$

i.e., the distance from the anterior boundary of the BOS, defined by the position of the leading heel, to the XcoM. Negative MOS_{AP} (XcoM ahead of the anterior BOS boundary) at heel contact

indicates that the body possesses sufficient energy to passively rotate beyond the upright

position and fall forward (assuming that the energy loss at heel contact and the energy input

during the subsequent push-off are either equal or negligible). Conversely, positive MOS_{AP}

(XcoM is behind the anterior BOS boundary) indicates that the body cannot passively rotate

beyond the upright position, and it will eventually fall backward.

For a passive walker, $MOS_{AP} = 0$ serves as a threshold for detecting unstable gait. Hof

demonstrated that stable gait arises with XcoM ahead of the BOS ($MOS_{AP} < 0$; Eqn. 1) in a

mathematical model of a passive walker [15]. Conversely, $MOS_{AP} > 0$ would mean that the

forward progression of the passive walker will stop, and the gait will be unstable.

In this work, we are interested in the forward loss of stability arising from a trip.

Therefore, we invert the interpretation of the MOS_{AP} from that for the passive walker. We

consider MOS_{AP} directly proportional to gait stability in the anterior direction. That is, lower

MOS_{AP} (more anterior location of the XcoM) indicates less stable gait, since a forward fall is

more likely if a perturbation like a trip occurs. Conversely, higher MOS_{AP} (more posterior

location of XcoM) indicates more stable gait [13, 27]. Furthermore, MOS_{AP} is inversely

proportional to efficiency: lower MOS_{AP} indicates greater passive forward motion, and hence

less active propulsion is required to maintain forward progression, and vice-versa. In this way,

MOS_{AP} reflects the tradeoff between stability and efficiency.

We used the uncontrolled manifold (UCM) analysis to quantify the synergy stabilizing

MOS_{AP} at heel contact. A *synergy* is co-variation in a redundant sets of input body variables that

maintains important output variables that define task performance. The UCM method has been

widely used to quantify the synergistic covariation in body variables in a variety of human

movements including gait ([28, 29]; see [30, 31] for recent reviews). Importantly, in addition to

identifying task-specific covariation, the UCM method identifies the salient task variables

controlled by the nervous system.

Here, we use the UCM method to evaluate the hypothesis that the input variables – step

length and XcoM, co-vary to stabilize the output variable, i.e., MOS_{AP}. We performed the

analysis separately for each step for both tasks. We first obtain from the constraint equation

(Eqn. 1), the Jacobian matrix that relates small changes in the step length and XcoM to changes

in MOS$_{AP}$: J = [1   -1]. The one-dimensional null space of this Jacobian defines the UCM, and its

one-dimensional compliment defines the orthogonal (ORT) manifold. We pool the across-trial

step length and XcoM data for a particular step. The deviation in the step length and XcoM data

for each trial from the across-trial mean is projected onto the UCM and the ORT manifolds. The

variances in these projections are the V$_{UCM}$ and the V$_{ORT}$, respectively. These variance

components yield the synergy index:

207
$$\Delta V = \frac{V_{UCM} - V_{ORT}}{\left(\frac{V_{UCM} + V_{ORT}}{2}\right)}.$$

The synergy index ΔV has a threshold value of zero. When ΔV > 0, V$_{UCM}$ > V$_{ORT}$. In general, this

implies that the control is organized so that most of the variability in the inputs is channeled

along the UCM, and therefore, it does not alter task performance, i.e., the output. Here, ΔV > 0

implies that the step length and XcoM covary to stabilize MOS$_{AP}$. Conversely, when ΔV < 0, V$_{UCM}$

< V$_{ORT}$. This implies that control is organized so that most of the variability in the inputs alters

task performance (i.e., change MOS$_{AP}$). In both these cases, when the synergy index differs from

zero, we would conclude that MOS$_{AP}$ is a controlled variable. Finally, ΔV = 0 indicates that there

is no task-specific co-variation in the step length and XcoM. This result would indicate that

MOS$_{AP}$ is not a controlled variable.

The synergy index ΔV ranges from -2 to 2. Therefore, it was z-transformed for statistical

analysis [29, 32]:

219
$$\Delta Vz = \frac{1}{2} \times \log\left[\frac{2 + \Delta V}{2 - \Delta V}\right].$$

220   Note that ΔV = 0 translates to ΔVz = 0. We report ΔVz values in the Results section and use ΔVz

221   values to draw inferences, consistent with most previous studies [30].

222   2.4 Statistical analysis

To determine whether the gait task and foot placement affected MOS$_{AP}$ (H1), we performed a

two-way (task × step) repeated measures ANOVA. To identify the source of changes in MOS$_{AP}$,

we performed the two-way ANOVA separately on the CoM position, CoM velocity at heel

contact and step length. To determine whether the synergies were present, we performed

separate, one-sample t-tests to test if ΔVz was significantly different from zero for each step in

the two tasks (H2). To determine whether the gait task and foot placement affected the

synergy index (H3), we performed a task × step repeated-measures ANOVA. To identify the

source of changes in the synergy index, we performed the two-way ANOVA separately on the

variance components (V_{UCM}, V_{ORT}). For all ANOVA tests, we fit a generalized linear model with

random effects. Recall that the UCM analysis utilizes data from all 15 trials to yield a single

value each for the synergy index, V_{UCM}, and V_{ORT}. Therefore, participant was the random effect

for these ANOVAs. However, data from all 15 trials was used to analyze all the remaining

variables. Therefore, the trial number within each participant was the nested random effect for

these ANOVAs. Tukey-Kramer adjustments were used to perform the following planned

pairwise comparisons: (1) across-task comparison at each of the six steps (or seven foot

placements), and (2) all across-step (or foot placement) comparisons for the obstacle-crossing

task only. All analyses were performed using the PROC GLIMMIX procedure in SAS 9.4 (Cary,

NC, USA) with significance set at 0.05.

242 3. Results

The Waterloo Footedness Questionnaire scores were 8 ± 5, indicating that all

participants were right-leg dominant.

Next, we use data from one participant (Fig. 2) to outline the overall results. The

detailed statistical results are presented in the following subsections. Figure 2 illustrates (1) the

changes in various outcome variables while approaching, crossing, and resuming gait after

crossing an obstacle, and (2) how the variables compare with unobstructed gait for each step.

All outcome variables showed a task × step interaction (supporting H1 and H3) indicating that

the pattern of changes in the variables across steps was different for the obstacle-crossing task

compared to the no obstacle task. The largest across-task changes were observed in all

outcomes for the two crossing steps ($Step_0$ and $Step_{+1}$), and across-task changes were apparent

in some outcomes in the approach to the obstacle ($Step_{-3}$ to $Step_{-1}$). All variables showed

across-step changes for the obstacle-crossing task. The synergy index was greater than zero,

indicating that the step length and the XcoM co-varied to stabilize MOS_{AP} for all steps in both

tasks (supporting H2). In the detailed results below, we first present the results for MOS_{AP},

followed by results for the variables that constitute MOS_{AP}: CoM position relative to rear heel,

CoM velocity at heel contact, and step length. We then present the results for the UCM

outcome variables quantifying the stability of MOS_{AP}.

Figure 2. Representative data from one participant. Figure illustrates changes in various

outcomes while approaching and crossing an obstacle, and then resuming unobstructed gait

(blue ellipses). Steps during unobstructed gait are included for comparison (red ellipses). The

45° line separates the regions of positive and negative MOS_{AP}; the white region above the line

represents passively unstable behavior, and the gray region below the line represents passively

stable behavior. The 45° line also represents the direction of the uncontrolled manifold (UCM).

Variations in step length and XcoM along this direction do not change MOS_{AP}. The ellipses are
centered at the centroid of the across-trial data for each step, and the major axes are aligned
with the 45° line. However, the flatness of the ellipses is representative. Changes in mean
values of the MOS_{AP} are reflected in the position of the ellipses in each chart. The flatness of
the ellipse reflects the synergy index, with flatter ellipses indicating higher synergy indices that
reflect stronger covariation between the step length and XcoM.

3.1 Margin of stability

MOS_{AP}

We observed a task × step interaction for MOS_{AP} ($F_{6,4167}$ = 886.89, p < 0.001; η_p^2 = 0.9;
Fig. 3A). Post-hoc comparisons across tasks revealed that MOS_{AP} was higher (more stable) for
the obstacle-crossing task than the no obstacle task at all except the first (fp_{-4}) and last two foot
placements (fp_{+2} and fp_{+3}). MOS_{AP} was lower (less stable) for fp_{+2} (after trail foot crossing) for
the obstacle-crossing task. All pair-wise across-step comparisons for the obstacle-crossing task
are depicted in Fig. 3A. For brevity, we describe only some of the significant differences.
Changes in MOS_{AP} occurred before the obstacle was reached; MOS_{AP} was higher (more stable)
for fp_{-1} relative to the two preceding foot placements. For the two crossing steps, MOS_{AP} first
increased to the highest value for fp_{+1}, and then decreased to the lowest value for fp_{+2} across all
foot placements.

CoM position relative to the rear heel

A task × step interaction was observed for the CoM position relative to the rear heel
($F_{6,4167}$ = 391.20, p < 0.001; η_p^2 = 0.9; Fig. 3B). Post-hoc comparisons across tasks revealed that
CoM was closer to the rear heel at the foot placements of all approach steps and the lead
crossing step (fp_{-4} to fp_{+1}) for the obstacle-crossing task than the no obstacle task. Conversely,
CoM was closer to the front heel at the foot placement for the trail crossing step (fp_{+2}) for the
obstacle-crossing task. All pair-wise across-step comparisons for the obstacle-crossing task are
depicted in Fig. 3B. For brevity, we describe only some of the significant differences. CoM

position changed for the last three foot placements. The CoM was closer to the rear foot at the

foot placement for the lead crossing step (fp_{+1}). It shifted forward, so that it was closer to the

front heel at the foot placement for the trail crossing step (fp_{+2}). Finally, for the last step (fp_{+3}),

CoM position was consistent with that for the approach steps.

CoM velocity at heel contact

We observed a task $\times$ step interaction for the CoM velocity at heel contact ($F_{6,4167}$=

161.52, $p < 0.001$; $\eta_p^2 = 0.8$; Fig. 3C). Post-hoc comparisons across tasks revealed that the CoM

velocity was lower for the obstacle-crossing task compared to the no obstacle task at all but the

first and the last foot placements (fp_{-4} and fp_{+3}). Post-hoc comparisons across steps for the

obstacle-crossing task revealed that the CoM velocity was lower at the first (fp_{-4}) and last (fp_{+3})

foot placement compared to other foot placements, except at the lead foot crossing (fp_{+1}),

where the CoM velocity was lower compared to all other foot placements.

Step length

We observed a task $\times$ step interaction for the step length ($F_{5,3569} = 149.09$; $p < 0.001$; η_p^2

$= 0.9$; Fig. 3D). Post-hoc comparisons across tasks revealed that step length was longer during

the obstacle-crossing task compared to the no obstacle task at the lead ($step_0$) and trail crossing

($step_{+1}$) steps. Post-hoc comparisons across steps for the obstacle-crossing task revealed that

step length was higher for step-1 compared to step-3. That is, relative to the early step, the

step length increased one step before the first crossing step. Step length was highest for the

lead crossing step ($step_0$), and then it shortened for the trail crossing step ($step_{+1}$), but

remained longer than all other steps. Finally, step length was the shortest for the last step

($step_{+2}$) compared to all other steps.

Figure 3. MOS_AP and its components. MOS_AP **(A)**, CoM position at heel contact **(B)**, CoM velocity at heel contact **(C)** and step length **(D)**. Data are across-subject means (N=20), and error bars denote standard error. Asterisks denote significant task difference at that step (p < 0.05). Across-step pairwise comparisons for the obstacle-crossing task are indicated by the row of letters above the horizontal axis for each panel. Steps that do not have letters in common are significantly different from each other (i.e., A is different from B, but A is not different from AB).

3.2 UCM variables

Synergy index

The synergy index (ΔVz) was significantly greater than zero at all steps for both tasks (t_{19} < 15.97, p < 0.01; Fig. 4A).

There was a task × step interaction for the synergy index ΔVz ($F_{5,209}$ = 2.89, p = 0.015; η_p^2 = 0.3; Fig. 4A). Post-hoc comparisons across tasks revealed that ΔVz was lower during the obstacle-crossing task compared to the no obstacle task for only the lead ($step_0$) and trail crossing steps ($step_{+1}$). Post-hoc comparisons across steps for the obstacle-crossing task revealed that ΔVz was lower at the lead crossing step ($step_0$) compared to all non-crossing steps except $step_{-2}$, and ΔVz was lower at the trail crossing step ($step_{+1}$) compared to the last step ($step_{+2}$).

Variance components

There was a task × step interaction for V_{UCM} ($F_{5,209}$ = 3.12, p = 0.009; η_p^2 = 0.3; Fig. 4B). Post-hoc comparisons across tasks revealed that V_{UCM} was higher for the obstacle-crossing task compared to the no obstacle task at all but the first two steps ($step_{-3}$ and $step_{-2}$). Post-hoc comparisons across steps for the obstacle-crossing task revealed that V_{UCM} was higher at the lead crossing step ($step_0$) compared to the first two steps ($step_{-3}$ and $step_{-2}$).

There was a task × step interaction for V_{ORT} ($F_{5,209}$ = 16.51, p < 0.001; η_p^2 = 0.3; Fig. 4C). Post-hoc comparisons across task revealed that V_{ORT} was higher during the obstacle-crossing task compared to the no obstacle task at the lead ($step_0$) and trail ($step_{+1}$) crossing steps. Post-hoc comparisons across steps for the obstacle-crossing task revealed that V_{ORT} was higher at the lead crossing step ($step_0$) compared to all other steps, and V_{ORT} was higher at the trail crossing step ($step_{+1}$) compared to all steps except the lead crossing step (lower for $step_{+1}$ compared to $step_0$) and the preceding step ($step_{-1}$; no significant difference).

Figure 4. Outcomes of the synergy analysis. Synergy index **(A)**, V$_{UCM}$ **(B)**, and V$_{ORT}$ **(C)**. Data are across-subject means (N=20), and error bars denote standard error. Asterisks denote significant across-task differences at that step (p < 0.05). Across-step pairwise comparisons for the obstacle-crossing task are indicated by the row of letters above the horizontal axis for each panel. Steps that do not have letters in common are significantly different from each other (i.e., A is different from B, but A is not different from AB).

4. Discussion

Our goals were to establish that young adults proactively change MOS_{AP} while approaching and then crossing an obstacle, and that MOS_{AP} is actively controlled during unobstructed as well as obstructed walking. We had hypothesized a task by step interaction for MOS_{AP} (H1). We observed proactive changes in MOS_{AP} in anticipation of a threat to stability; MOS_{AP} changed across steps for unobstructed versus obstructed gait (supporting H1). Next, we had hypothesized that the synergy index will be greater than zero (H2). We report the novel finding that the central nervous system responds to changes in the body's motion with a corresponding correction in step length so that the MOS_{AP} itself is approximately invariant at each heel contact (supporting H2). Finally, we had hypothesized a task by step interaction for the synergy index (H3). We observed that the synergy index was lower while crossing an obstacle compared to earlier and later steps and compared to the corresponding steps during unobstructed gait (supporting H3). We argue below that (1) the proactive changes in the MOS_{AP} for the obstacle-crossing task reflect a stability-efficiency tradeoff; and (2) the positive synergy indices indicate that the passive dynamic stability is not a byproduct of another process, but that the step length is actively controlled to exploit consistently the passive dynamics to achieve either efficient or more stable gait.

4.1 Proactive changes in MOS_{AP} and the compromise between stability and energy efficiency

The first major finding of this study is that the MOS_{AP} changes while approaching the obstacle. When compared to unobstructed gait, greater stability was apparent three steps before the obstacle (Fig. 3A). Changes in MOS_{AP} during approach and crossing resulted from reduced speed, a more posterior CoM, and, for some steps, increased step length (Figs. 3B, 3C, 3D). Previous work has demonstrated that changes in MOS_{AP} and MOS_{ML} were evident during the swing phase immediately prior to taking unnaturally long or quick steps [33], and in MOS_{ML} one step before reaching an obstacle [34]. Here we extend this finding and demonstrate that

MOS$_{AP}$ changes several steps before reaching the obstacle, indicating that the transition from

unobstructed to obstructed gait occurs over several steps.

Our findings extend the argument that passive dynamic stability is modulated in

response to perceived risk [6, 27]. For example, the substantive increase in MOS$_{AP}$ for the lead

crossing step (Figs. 2, 3A) reflects cautious gait and may be a preemptive strategy against a

potential trip that could lead to a forward fall [12]. Similarly, the even-more-substantive

transition back to the least stable passive dynamics (lowest MOS$_{AP}$; Fig. 3A) following the trail

crossing step may reflect the reduced risk of a forward fall after the trail foot has crossed the

obstacle, and the exploitation of passive dynamics to propel the body forward to regain gait

speed.

These fluctuations in passive dynamic stability increase the energetic cost of

locomotion. The increase in MOS$_{AP}$ over approach steps, and especially the positive MOS$_{AP}$ for

the crossing step, indicates that the trailing leg must push off more so that the body can rotate

about and beyond the stance ankle [35]. Furthermore, the lead crossing step (step$_0$) is a slower

and *longer* step (Figs 3C, 3D), which is inconsistent with the typical combination of a slower and

shorter step during unobstructed gait which reduces energetic cost [4, 36]. Therefore, our

results reflect a tradeoff between stability and energy efficiency, consistent with similar

arguments offered in the context of stair descent [6], and model-based optimization

computations of obstacle crossing behaviors [20].

In sum, inspecting changes in MOS$_{AP}$ across tasks and steps leads to the conclusion that

MOS$_{AP}$ is proactively adjusted (1) during the approach, likely to facilitate the transition from

unobstructed gait to the movements required to clear the obstacle, (2) while crossing an

obstacle, likely to prioritize safety over energy optimality, and (3) when resuming level gait to

regain gait speed. The large fluctuations in MOS$_{AP}$ reflect a tradeoff between stability and

energy efficiency.

4.2 Active control of MOS$_{AP}$ during unobstructed and obstructed gait

The synergy index provides evidence for the control of MOS$_{AP}$

The second major finding of this study is that MOS_{AP} is a controlled variable for

obstructed as well as unobstructed gait ($\Delta Vz > 0$ for all steps; Fig. 4A). In particular, the synergy

index remains significantly larger than zero, even though the input variables that define the

MOS_{AP} change over multiple steps for the obstacle-crossing task (Figs 2 and 3). Thus, the UCM

analysis provides strong quantitative evidence that MOS_{AP} is controlled.

The variance components (Figs 4B, 4C) provide information regarding the underlying

processes that stabilize MOS_{AP}. Higher V_{ORT} indicates higher variability in MOS_{AP}, whereas

higher V_{UCM} indicates greater compensatory covariance between XcoM and step length. We

observed that MOS_{AP} is more variable for the crossing steps (up to 205% increase in V_{ORT}

compared to earlier steps; Fig. 4C). This likely arises from the larger muscle activations and joint

moments required for the crossing steps compared to unobstructed steps [21, 37]. Higher

activations would increase signal-dependent noise [22], which will lead to more variable MOS_{AP}.

However, this increase is offset by an increase in V_{UCM} (up to 103% increase compared to earlier

steps; Fig. 4B). This compensation leads to MOS_{AP} stabilization overall (Fig. 4A).

Does the synergy arise from passive mechanics, or do neural mechanisms

indicate active control?

It is unlikely that passive mechanics of the gait cycle alone can explain the positive

synergy indices that we observed. Rather, both passive mechanics and active control are

responsible for our data. It is indeed likely that mechanics contribute; for example, a greater

push off force would result in a more anterior XcoM at the next heel contact, but it would also

tend to produce longer steps [3], thereby helping to maintain MOS_{AP}. However, the step length

is not entirely determined by a passively swinging leg. Rather, activity in the leg muscles and

power at the hip and knee joints indicate active control of the forward swinging limb, which will

alter the step length from what a passively swinging limb would yield [38]. Spinal stretch reflex

loops have long been implicated in the control of locomotion [39, 40]. Feldman [41] recently

elucidated locomotor control based on his theory of referent configurations which incorporates

spinal reflexes. The central idea in this theory is that to intentionally change a limb's position,

the nervous system modifies the parameter λ, which is the threshold muscle length at which α-

motor neurons are recruited and the stretch reflex is activated. Furthermore, movements are

executed by setting the time courses $\lambda(t)$ for various muscles. Muscle activations arise via the

stretch-reflex loop in relation to their deviations from the corresponding λ to drive the muscles

(and hence the body) towards its referent configuration. It is plausible that variations in muscle

lengths from their reference values, arising from variations in push off forces, will engage spinal

stretch reflexes that will alter step length and contribute to the observed MOS_{AP} synergy [42].

This explanation is also consistent with the view that for steady state, level gait, humans rely on

spinal feedback for control in the AP direction [43, 44].

The contribution of neural mechanisms to the MOS_{AP} synergy may be even greater for

obstructed gait, and in addition to the spinal processes implicated during unobstructed gait,

supraspinal processes may contribute to the synergy. We observed large fluctuations in the

MOS_{AP} over the approach and crossing steps (174% increase for fp_{+1} over fp_{-1}, and a 292%

decline for fp_{+2} over fp_{+1}; Fig. 3A). Nevertheless, the synergy index, although lower (38% decline

for $Step_0$ relative $Step_{-1}$; Fig. 4A), remained positive. The positive synergy index despite large

fluctuations in related variables over consecutive steps suggests the involvement of supraspinal

mechanisms. It is generally thought that supraspinal mechanisms influence synergies. The most

convincing evidence is that synergies in individuals with neurological problems (persons with

Parkinson's disease, stroke survivors, etc.) are altered compared to healthy, age-matched

controls (see [31] for review). Specifically, during obstacle crossing, supra-spinal mechanisms

may influence the synergy index by modulating the gain on spinal reflexes. Indeed, supraspinal

centers are involved in the control of the obstacle crossing steps [21]. Spinal reflex responses in

the stance leg flexors are enhanced, likely due to increased activity in the prefrontal cortex, in

preparation for swinging the leg over the obstacle. Furthermore, the prefrontal cortical activity

remains enhanced (compared to unobstructed walking) while swinging the leg over the

obstacle [21]. It is also known that visual information about the obstacle is gathered during

approach and used to alter gait characteristics [45, 46]. Gathering and using visual information

also implicates higher brain centers in the control of obstructed gait, supporting our view that

the MOS_{AP} synergy arises from spinal and supraspinal neural circuits. We note both spinal and

supraspinal structures are frequently mentioned as candidate neurophysiological bases of

synergies [47], although the specific mechanisms are unknown.

In summary, the synergy values that we observed could arise partially from passive body

mechanics. However, active neurophysiological processes at the spinal and supra-spinal level

likely contribute to this signal as well, indicating active control of MOS_{AP}.

Our ideas parallel previous work indicating that the MOS in the medio-lateral (ML)

direction is controlled during level walking [18, 48-53]. The focus of previous research on ML

stability – versus AP stability – is consistent with the view that level gait requires more control

along the ML direction, and minimal control along the AP direction [43]. The UCM analyses

performed here show that the MOS is also controlled in the AP direction by regulating step

length – not only in gait tasks that require proactive adaptations, but also during unobstructed

gait.

4.3 Limitations

As an indicator of the stability of human gait, MOS_{AP} is challenging to interpret [54]. The

strict interpretation is that XcoM ahead of the BOS boundary indicates that gait cannot be

stopped within a step. This would indicate instability if the task were to stop walking within a

step. However, when the task is to continue walking, Hof's result is more pertinent: the same

condition yields a stable gait in a passive walker by ensuring that the walker does not stall [15].

The challenge in interpreting MOS_{AP} for human gait arise when extending this logic to

(1) different adaptive gait tasks and (2) while considering the relation of the passive dynamic

stability to the overall gait stability. First, authors have interpreted MOS_{AP} depending on the

task. For example, Bosse et al. consider XcoM ahead of the BOS boundary as unstable gait while

descending stairs [6], whereas others (including us) assume roughly the opposite for obstacle

crossing tasks [5, 13]. It seems that these interpretations must be evaluated on a case-by-case

basis. A better approach would be to validate the interpretations by correlating MOS_{AP}

characteristics with fall rates. Although this is difficult to accomplish, and many biomechanical

variables used to quantify stability of human gait lack such validation [54], this is an important

goal for future research.

Second, MOS_{AP} captures only the passive dynamic stability of the body, and the relation

of the passive stability to the overall stability of human gait is not straightforward due to the

contributions of active neuromuscular processes. For example, when MOS_{AP} is positive,

indicating that a passive walker would stall due to insufficient energy, higher active push-off by

the rear leg of a human walker could overcome this deficit. This is reflected in the data.

Although the relationship $MOS_{AP} < 0$ is consistently observed in stable human gait [7-9, 13, 55-

57], the reverse relationship ($MOS_{AP} > 0$) has also been observed in some stable human walking

trials [55]. Even in our data (Fig 3A), we observe positive and negative MOS_{AP} values, and there

were no signs of instability in the gait overall.

Nevertheless, the accumulated evidence regarding the changes in MOS_{AP} across tasks

and populations suggests that the passive dynamic stability might influence the ability of the

person to recover from large perturbations, given that the speed and magnitude of human

neuromuscular responses are bounded. Hence passive stability is enhanced when large

perturbations could occur. This effect will likely be larger in older or patient populations where

maximal capacities are reduced. Therefore, we suggest that studying MOS_{AP} is useful; our study

of MOS_{AP} has revealed information regarding locomotor control, and our findings will serve as a

baseline for identifying potential locomotor issues in various populations.

Another limitation of this study is that we estimated the CoM location using four pelvis

markers [26]. A rigorous whole-body model could have provided slightly different estimates of

the CoM motion. However, these differences would influence MOS_{AP} similarly across all

conditions, and using a different method to obtain CoM kinematics would likely yield similar

qualitative results and overall conclusions, especially given the large effect sizes for all our

outcome variables.

It can be argued that the all the steps of the unobstructed task are similar. Therefore,

the data for the unobstructed task could be collapsed across the steps and compared with the

steps of the obstructed task. We performed this alternate analysis and obtained results that led

to identical conclusions. However, we observed effects of gait initiation and termination in our

data (cf. Fig 3C). Gait was initiated one step before the examined steps, and gait was

terminated one or two steps after the trail crossing step. Therefore, collapsing the data across

all steps of the unobstructed task would be suspect, and the statistical approach we used is

more appropriate. Using a longer walkway – with more steps before and after the obstacle –

may alter the values of our outcome variables. We do not expect that these changes will

influence our key conclusions (proactive changes in and control of MOS_{AP}). However, effects of

gait initiation and termination on MOS_{AP} characteristics may be worth an independent

investigation.

533   5. Conclusion

We demonstrated that the XcoM and the step length covary to maintain MOS_{AP}, indicating that

MOS_{AP} is controlled during unobstructed and obstructed gait. In conjunction with the

consistently low MOS_{AP} values for most of the analyzed steps, we conclude that the MOS_{AP} is

controlled to exploit the passive dynamics and achieve forward progression at low energetic

cost. Furthermore, the value of MOS_{AP} is proactively altered while approaching an obstacle, and

MOS_{AP} shows substantial fluctuations for the two crossing steps. These changes reflect a

tradeoff between stability and energy efficiency. The changes during approach and lead

crossing steps indicate increasingly cautious gait with a growing preference for stability,

whereas the opposite change after the trail crossing step indicates a reversion to improving

efficiency when the risk to stability is reduced. Thus, our results indicate that humans exploit

the passive AP body motion to meet specific ends dictated by the locomotor task. We conclude

that the UCM analysis of MOS_{AP} provides new information regarding the control of stability

during walking, especially for gait tasks requiring proactive adaptations, and our methods could

be valuable in understanding the effects of age and pathology on gait.

REFERENCES

1.   King DL, Arnold AS, Smith SL. A Kinematic Comparison of Single, Double and Triple Axels. Journal
of Applied Biomechanics. 1994;10(1):51-60. doi: DOI 10.1123/jab.10.1.51. PubMed PMID:
WOS:A1994MU82100005.
2.   Knoll K, Hildebrand F. Optimum Movement Coordunation in Multi-Revolution Jumps in Figure
Skating. In: Riehle HJ, Vieten MM, editors. 16 International Symposium on Biomechanics in Sport July 21-
25; Konstanz, Germany1998.
3.   Kuo AD, Donelan JM. Dynamic principles of gait and their clinical implications. Phys Ther.
2010;90(2):157-174. Epub 2009/12/22. doi: 10.2522/ptj.20090125. PubMed PMID: 20023002; PubMed
Central PMCID: PMCPMC2816028.
4.   Kuo AD, Donelan JM, Ruina A. Energetic consequences of walking like an inverted pendulum:
step-to-step transitions. Exerc Sport Sci Rev. 2005;33(2):88-97. Epub 2005/04/12. doi:
10.1097/00003677-200504000-00006. PubMed PMID: 15821430.
5.   Hak L, Hettinga FJ, Duffy KR, Jackson J, Sandercock GRH, Taylor MJD. The concept of margins of
stability can be used to better understand a change in obstacle crossing strategy with an increase in age.
J Biomech. 2019;84:147-152. Epub 2019/01/16. doi: 10.1016/j.jbiomech.2018.12.037. PubMed PMID:
30642664.
6.   Bosse I, Oberlander KD, Savelberg HH, Meijer K, Bruggemann GP, Karamanidis K. Dynamic
stability control in younger and older adults during stair descent. Hum Mov Sci. 2012;31(6):1560-1570.
Epub 2012/08/03. doi: 10.1016/j.humov.2012.05.003. PubMed PMID: 22853941.
7.   Bierbaum S, Peper A, Karamanidis K, Arampatzis A. Adaptive feedback potential in dynamic
stability during disturbed walking in the elderly. J Biomech. 2011;44(10):1921-1926. Epub 2011/05/11.
doi: 10.1016/j.jbiomech.2011.04.027. PubMed PMID: 21555126.
8.   Bierbaum S, Peper A, Karamanidis K, Arampatzis A. Adaptational responses in dynamic stability
during disturbed walking in the elderly. J Biomech. 2010;43(12):2362-2368. Epub 2010/05/18. doi:
10.1016/j.jbiomech.2010.04.025. PubMed PMID: 20472240.
9.   Ohtsu H, Yoshida S, Minamisawa T, Katagiri N, Yamaguchi T, Takahashi T, et al. Does the balance
strategy during walking in elderly persons show an association with fall risk assessment? J Biomech.
2020;103:109657. Epub 2020/02/10. doi: 10.1016/j.jbiomech.2020.109657. PubMed PMID: 32035661.
10.   Peebles AT, Bruetsch AP, Lynch SG, Huisinga JM. Dynamic Balance Is Related to Physiological
Impairments in Persons With Multiple Sclerosis. Arch Phys Med Rehabil. 2018;99(10):2030-2037. Epub
2017/12/25. doi: 10.1016/j.apmr.2017.11.010. PubMed PMID: 29274726; PubMed Central PMCID:
PMCPMC6014868.
11.   Peebles AT, Bruetsch AP, Lynch SG, Huisinga JM. Dynamic balance in persons with multiple
sclerosis who have a falls history is altered compared to non-fallers and to healthy controls. J Biomech.
2017;63:158-163. Epub 2017/09/12. doi: 10.1016/j.jbiomech.2017.08.023. PubMed PMID: 28889946.
12.   Peebles AT, Reinholdt A, Bruetsch AP, Lynch SG, Huisinga JM. Dynamic margin of stability during
gait is altered in persons with multiple sclerosis. J Biomech. 2016;49(16):3949-3955. Epub 2016/11/28.
doi: 10.1016/j.jbiomech.2016.11.009. PubMed PMID: 27889188; PubMed Central PMCID:
PMCPMC5176013.
13.   Raffegeau TE, Brinkerhoff SA, Kellaher GK, Baudendistel S, Terza MJ, Roper JA, et al. Changes to
margins of stability from walking to obstacle crossing in older adults while walking fast and with a dual-
task. Exp Gerontol. 2022;161:111710. Epub 2022/01/30. doi: 10.1016/j.exger.2022.111710. PubMed
PMID: 35090973.
14.   Hof AL, Gazendam MG, Sinke WE. The condition for dynamic stability. J Biomech. 2005;38(1):1-
8. Epub 2004/11/03. doi: 10.1016/j.jbiomech.2004.03.025. PubMed PMID: 15519333.

15. Hof AL. The 'extrapolated center of mass' concept suggests a simple control of balance in
walking. Hum Mov Sci. 2008;27(1):112-125. Epub 2007/10/16. doi: 10.1016/j.humov.2007.08.003.
PubMed PMID: 17935808.
16. Tillman M, Ambike S. The Influence of Recent Actions and Anticipated Actions on the Stability of
Finger Forces During a Tracking Task. Motor Control. 2020;24(3):365-382. Epub 2020/07/15. doi:
10.1123/mc.2019-0124. PubMed PMID: 32663389.
17. Hof AL, Vermerris SM, Gjaltema WA. Balance responses to lateral perturbations in human
treadmill walking. J Exp Biol. 2010;213(Pt 15):2655-2664. Epub 2010/07/20. doi: 10.1242/jeb.042572.
PubMed PMID: 20639427.
18. McAndrew Young PM, Wilken JM, Dingwell JB. Dynamic margins of stability during human
walking in destabilizing environments. J Biomech. 2012;45(6):1053-1059. Epub 2012/02/14. doi:
10.1016/j.jbiomech.2011.12.027. PubMed PMID: 22326059; PubMed Central PMCID: PMCPMC3321251.
19. Scholz JP, Schoner G. The uncontrolled manifold concept: identifying control variables for a
functional task. Exp Brain Res. 1999;126(3):289-306. PubMed PMID: 10382616.
20. Chou LS, Draganich LF, Song SM. Minimum energy trajectories of the swing ankle when stepping
over obstacles of different heights. J Biomech. 1997;30(2):115-120. Epub 1997/02/01. doi:
10.1016/s0021-9290(96)00111-x. PubMed PMID: 9001931.
21. Haefeli J, Vogeli S, Michel J, Dietz V. Preparation and performance of obstacle steps: interaction
between brain and spinal neuronal activity. Eur J Neurosci. 2011;33(2):338-348. Epub 2010/11/13. doi:
10.1111/j.1460-9568.2010.07494.x. PubMed PMID: 21070395.
22. Harris CM, Wolpert DM. Signal-dependent noise determines motor planning. Nature.
1998;394(6695):780-784. Epub 1998/09/02. doi: 10.1038/29528. PubMed PMID: 9723616.
23. Elias LJ, Bryden MP, Bulman-Fleming MB. Footedness is a better predictor than is handedness of
emotional lateralization. Neuropsychologia. 1998;36(1):37-43. Epub 1998/04/09. doi: 10.1016/s0028-
3932(97)00107-3. PubMed PMID: 9533385.
24. Rosenblatt NJ, Hurt CP. Recommendation for the minimum number of steps to analyze when
performing the uncontrolled manifold analysis on walking data. J Biomech. 2019;85:218-223. Epub
2019/02/06. doi: 10.1016/j.jbiomech.2019.01.018. PubMed PMID: 30718066; PubMed Central PMCID:
PMCPMC6905614.
25. Desailly E, Daniel Y, Sardain P, Lacouture P. Foot contact event detection using kinematic data in
cerebral palsy children and normal adults gait. Gait Posture. 2009;29(1):76-80. Epub 2008/08/05. doi:
10.1016/j.gaitpost.2008.06.009. PubMed PMID: 18676147.
26. Whittle MW. Three-dimensional motion of the center of gravity of the body during walking.
Hum Movement Sci. 1997;16(2-3):347-355. doi: Doi 10.1016/S0167-9457(96)00052-8. PubMed PMID:
WOS:A1997WV17500014.
27. Hak L, Houdijk H, Beek PJ, van Dieen JH. Steps to take to enhance gait stability: the effect of
stride frequency, stride length, and walking speed on local dynamic stability and margins of stability.
PLoS One. 2013;8(12):e82842. Epub 2013/12/19. doi: 10.1371/journal.pone.0082842. PubMed PMID:
24349379; PubMed Central PMCID: PMCPMC3862734.
28. Cui C, Kulkarni A, Rietdyk S, Barbieri FA, Ambike S. Synergies in the ground reaction forces and
moments during double support in curb negotiation in young and older adults. J Biomech.
2020;106:109837. Epub 2020/06/11. doi: 10.1016/j.jbiomech.2020.109837. PubMed PMID: 32517974.
29. Kulkarni A, Cho H, Rietdyk S, Ambike S. Step length synergy is weaker in older adults during
obstacle crossing. J Biomech. 2021;118:110311. Epub 2021/02/19. doi:
10.1016/j.jbiomech.2021.110311. PubMed PMID: 33601182.
30. Vaz DV, Pinto VA, Junior RRS, Mattos DJS, Mitra S. Coordination in adults with neurological
impairment - A systematic review of uncontrolled manifold studies. Gait Posture. 2019;69:66-78. Epub
2019/01/25. doi: 10.1016/j.gaitpost.2019.01.003. PubMed PMID: 30677709.

31.   Latash ML, Huang X. Neural control of movement stability: Lessons from studies of neurological
patients. Neuroscience. 2015;301:39-48. doi: 10.1016/j.neuroscience.2015.05.075. PubMed PMID:
26047732; PubMed Central PMCID: PMCPMC4504804.
32.   Ambike S, Penedo T, Kulkarni A, Santinelli FB, Barbieri FA. Step length synergy while crossing
obstacles is weaker in patients with Parkinson's disease. Gait Posture. 2021;84:340-345. Epub
2021/01/18. doi: 10.1016/j.gaitpost.2021.01.002. PubMed PMID: 33454501.
33.   Sivakumaran S, Schinkel-Ivy A, Masani K, Mansfield A. Relationship between margin of stability
and deviations in spatiotemporal gait features in healthy young adults. Hum Mov Sci. 2018;57:366-373.
Epub 2017/10/11. doi: 10.1016/j.humov.2017.09.014. PubMed PMID: 28987772; PubMed Central
PMCID: PMCPMC5770210.
34.   Worden TA, Vallis LA. Stability control during the performance of a simultaneous obstacle
avoidance and auditory Stroop task. Exp Brain Res. 2016;234(2):387-396. Epub 2015/10/22. doi:
10.1007/s00221-015-4461-z. PubMed PMID: 26487180.
35.   Jin J, van Dieen JH, Kistemaker D, Daffertshofer A, Bruijn SM. Does ankle push-off correct for
errors in anterior-posterior foot placement relative to center-of-mass states? BioRxiv. 2022. doi:
https://doi.org/10.1101/2022.03.14.484283.
36.   Kuo AD. A simple model of bipedal walking predicts the preferred speed-step length
relationship. J Biomech Eng. 2001;123(3):264-269. Epub 2001/07/31. doi: 10.1115/1.1372322. PubMed
PMID: 11476370.
37.   Chou LS, Draganich LF. Stepping over an obstacle increases the motions and moments of the
joints of the trailing limb in young adults. J Biomech. 1997;30(4):331-337. Epub 1997/04/01. doi:
10.1016/s0021-9290(96)00161-3. PubMed PMID: 9075000.
38.   Winter DA. The Bionechanics and Motor Control of Human Gait: Normal, Elderly and
Pathological. Second ed. Kitchener, ON: Waterloo Biomechanics; 1991.
39.   Dietz V, Quintern J, Sillem M. Stumbling reactions in man: significance of proprioceptive and
pre-programmed mechanisms. J Physiol. 1987;386:149-163. Epub 1987/05/01. doi:
10.1113/jphysiol.1987.sp016527. PubMed PMID: 3681704; PubMed Central PMCID: PMCPMC1192455.
40.   Takakusaki K. Neurophysiology of gait: from the spinal cord to the frontal lobe. Mov Disord.
2013;28(11):1483-1491. Epub 2013/10/18. doi: 10.1002/mds.25669. PubMed PMID: 24132836.
41.   Feldman AG, Levin MF, Garofolini A, Piscitelli D, Zhang L. Central pattern generator and human
locomotion in the context of referent control of motor actions. Clin Neurophysiol. 2021;132(11):2870-
2889. Epub 2021/10/11. doi: 10.1016/j.clinph.2021.08.016. PubMed PMID: 34628342.
42.   Latash ML. One more time about motor (and non-motor) synergies. Exp Brain Res.
2021;239(10):2951-2967. Epub 2021/08/13. doi: 10.1007/s00221-021-06188-4. PubMed PMID:
34383080.
43.   Bauby CE, Kuo AD. Active control of lateral balance in human walking. J Biomech.
2000;33(11):1433-1440. Epub 2000/08/15. doi: 10.1016/s0021-9290(00)00101-9. PubMed PMID:
10940402.
44.   Collins SH, Kuo AD. Two independent contributions to step variability during over-ground human
walking. PLoS One. 2013;8(8):e73597. Epub 2013/09/10. doi: 10.1371/journal.pone.0073597. PubMed
PMID: 24015308; PubMed Central PMCID: PMCPMC3756042.
45.   Patla AE, Vickers JN. Where and when do we look as we approach and step over an obstacle in
the travel path? Neuroreport. 1997;8(17):3661-3665. Epub 1998/01/14. doi: 10.1097/00001756-
199712010-00002. PubMed PMID: 9427347.
46.   Diaz GJ, Parade MS, Barton SL, Fajen BR. The pickup of visual information about size and location
during approach to an obstacle. PLoS One. 2018;13(2):e0192044. Epub 2018/02/06. doi:
10.1371/journal.pone.0192044. PubMed PMID: 29401511; PubMed Central PMCID: PMCPMC5798835.

47. Tillman M, Ambike S. Cue-induced changes in the stability of finger force-production tasks
revealed by the uncontrolled manifold analysis. J Neurophysiol. 2018;119(1):21-32. Epub 2017/09/22.
doi: 10.1152/jn.00519.2017. PubMed PMID: 28931612.
48. McAndrew Young PM, Dingwell JB. Voluntarily changing step length or step width affects
dynamic stability of human walking. Gait Posture. 2012;35(3):472-477. Epub 2011/12/17. doi:
10.1016/j.gaitpost.2011.11.010. PubMed PMID: 22172233; PubMed Central PMCID: PMCPMC3299923.
49. van Leeuwen AM, van Dieen JH, Daffertshofer A, Bruijn SM. Active foot placement control
ensures stable gait: Effect of constraints on foot placement and ankle moments. PLoS One.
2020;15(12):e0242215. Epub 2020/12/18. doi: 10.1371/journal.pone.0242215. PubMed PMID:
33332421; PubMed Central PMCID: PMCPMC7746185.
50. van Leeuwen AM, van Dieen JH, Daffertshofer A, Bruijn SM. Ankle muscles drive mediolateral
center of pressure control to ensure stable steady state gait. Sci Rep. 2021;11(1):21481. Epub
2021/11/04. doi: 10.1038/s41598-021-00463-8. PubMed PMID: 34728667; PubMed Central PMCID:
PMCPMC8563802.
51. Rosenblatt NJ, Grabiner MD. Measures of frontal plane stability during treadmill and overground
walking. Gait Posture. 2010;31(3):380-384. Epub 2010/02/05. doi: 10.1016/j.gaitpost.2010.01.002.
PubMed PMID: 20129786.
52. Reimann H, Fettrow T, Jeka JJ. Strategies for the Control of Balance During Locomotion.
Kinesiology Review. 2018;7(1):18. doi: 10.1123/kr.2017-0053 10.1123/kr.2017-0053 10.1123/kr.2017-
0053 10.1123/kr.2017-0053.
53. Kazanski ME, Cusumano JP, Dingwell JB. Rethinking margin of stability: Incorporating step-to-
step regulation to resolve the paradox. J Biomech. 2022;144:111334. Epub 2022/10/17. doi:
10.1016/j.jbiomech.2022.111334. PubMed PMID: 36244320.
54. Bruijn SM, Meijer OG, Beek PJ, van Dieen JH. Assessing the stability of human locomotion: a
review of current measures. J R Soc Interface. 2013;10(83):20120999. Epub 2013/03/22. doi:
10.1098/rsif.2012.0999. PubMed PMID: 23516062; PubMed Central PMCID: PMCPMC3645408.
55. Hohne A, Stark C, Bruggemann GP, Arampatzis A. Effects of reduced plantar cutaneous afferent
feedback on locomotor adjustments in dynamic stability during perturbed walking. J Biomech.
2011;44(12):2194-2200. Epub 2011/07/06. doi: 10.1016/j.jbiomech.2011.06.012. PubMed PMID:
21726865.
56. Ohtsu H, Yoshida S, Minamisawa T, Takahashi T, Yomogida SI, Kanzaki H. Investigation of balance
strategy over gait cycle based on margin of stability. J Biomech. 2019;95:109319. Epub 2019/08/31. doi:
10.1016/j.jbiomech.2019.109319. PubMed PMID: 31466715.
57. Watson F, Fino PC, Thornton M, Heracleous C, Loureiro R, Leong JJH. Use of the margin of
stability to quantify stability in pathologic gait - a qualitative systematic review. Bmc Musculoskel Dis.
2021;22(1). doi: 10.1186/s12891-021-04466-4. PubMed PMID: WOS:000669563500004.

(A)

fp$_{-4}$ fp$_{-3}$ fp$_{-2}$ fp$_{-1}$ Obstacle fp$_{+1}$ fp$_{+2}$ fp$_{+3}$

Step$_{-3}$ Step$_{-2}$ Step$_{-1}$ Step$_{0}$ Step$_{+1}$ Step$_{+2}$

Approach Crossing Last

(B)

MOS$_{AP}$ = Step length - XcoM

V$_{CoM}$ × √(l/g)

CoM

leg length l

Coordinate frame

Step length MOS$_{AP}$

XcoM