1 **Title**

2 • Do males know? Evidence-driven rainmaking of male bonobos to meet the fertile window
3 of females
4
5 **Short Title**

6 • Male mating efforts allocation based on female fertility
7

8 **Authors**
9 Heungjin Ryu,[1,2]* Chie Hashimoto,[2] David A. Hill,[3] Keiko Mouri,[2] Keiko Shimizu,[4]
10 Takeshi Furuichi[2]
11
Affiliations

[1]School of Life Sciences, Ulsan National Institute of Science and Technology, Ulsan,
South Korea.

[2]Primate Research Institute, Kyoto University, Aichi, Japan.

[3]Wildlife Research Center, Kyoto University, Kyoto, Japan

[4]Faculty of Science, Okayama University of Science, Okayama, Japan

*Corresponding author. Email: heungjin.ryu.26v@kyoto-u.jp
19

Abstract
Female bonobos exhibit prolonged receptivity. One suggested function of the prolonged

receptivity is it lowers male mating competition. However, it is questionable whether easier

access to receptive females can reduce male-male competition, given the exclusive nature of

male reproductive success. We tested whether males could determine a fertile phase of

females. We found that ovulation probability predicted male mating effort. High-ranking

males copulated with the female with higher fertility, and male-male agonistic interactions

increased when there were fertile females in the party. When there were multiple females with

maximal swelling, males concentrated their mating effort on a female with an older infant

whose maximal swelling started earlier, and they continued mating efforts until detumescence

(rainmaking). These findings suggest that male bonobos distinguish between fertile and non-

fertile phases of females and that having more receptive females in the party does not reduce

male-male competition for fertile females.

Teaser
Males use the rainmaker's rule to meet the periovulatory phase of the female bonobo for
better reproductive success.
36

Introduction

Male-male competition over mates and female choice shapes the behavior, physiology, and social systems of the species (*1, 2*). Bonobos (*Pan paniscus*) are known for their peaceful and mild social characteristics (*3–5*) compared with those of chimpanzees (*Pan troglodytes*). One hypothesis explaining their peaceful nature is their high operational sex ratio that influences male competition over mates – estrus sex ratio hypothesis (*6*). Based on findings that female bonobos are receptive for a longer period than female chimpanzees, this hypothesis proposes that the prolonged receptivity of female bonobo – achieved by prolonged and frequent maximal swelling phases – results in milder mating competition between male bonobos compared to chimpanzees. The main role of prolonged receptivity is, therefore, in this hypothesis, confusing paternity as well as reducing the tension surrounding male reproductive success by allowing males to increase their chance of mating. Prolonged receptivity in bonobo seems to be mainly a reflection of the prolonged maximal swelling periods as female bonobos achieve maximal swelling during the early lactation periods when the infant is younger than 2 years (*7*) and even during pregnancy (*8, 9*), whereas female chimpanzees do not (*10*).

Some researchers suggested that prolonged periods of maximal swelling in bonobos might help females conceal ovulation (*8, 11*). In line with this suggestion, female bonobos are known to have the longest period of maximal sexual swelling among species that exhibit sexual swelling (*6, 8, 11*). Moreover, the probability of ovulation during the maximal swelling phase in bonobos is the least predictable among these species with exaggerated sexual swelling (*11–13*). The lower predictability of ovulation in bonobos might support the idea that the prolonged periods of maximal swelling function to conceal ovulation. However, it is uncertain to what extent mating competition is reduced by easier access to mating due to the greater number of receptive females within a group and low predictability of ovulation. In addition, the estrous sex ratio hypothesis implies that males may not pay much attention to producing offspring if they are able to mate with a female with maximal swelling. Otherwise, male bonobos should be deceived by the signal of ovulation from the sexual swelling, so they do not get much reproductive benefit from male-male competition. However, it is unlikely that male mating competition is not an important factor that determines male reproductive success considering very high reproductive skew to high-ranking males (*14, 15*) and male-male competition over mate which sometimes involves their mothers in bonobos (*16*).

In previous studies on bonobos (*11*, *12*, *17*, *18*), female reproductive events including lactation and pregnancy were not considered thoroughly as an important factor in determining the resumption and continuation of the menstrual and swelling cycle. For example, the maximal swelling phase may less reliably indicate ovulation for females with younger infants, particularly whose infant is younger than 2-years-old. The maximal swelling phase of such females may also vary in length more than it does for females with older infants (*19*). Male bonobos may also respond differently to the sexual swelling of females with young infants, as those females might have a lower probability of getting pregnant, given that the inter-birth interval has a mean of 4.8 years (*20*). In such cases, the sexual swelling of females with young infants can be considered less meaningful for male mating decision-making.

Despite the lower predictability of ovulation from the onset of maximal swelling in bonobos among primates, over half of the ovulation occurs at the second half of maximal swelling periods (*17*, *18*, *11*, *20*); but see also (*12*). We do not have any hormonal evidence, if it ever exists, that explains how ovulation can deviate from sexual swelling given the antagonistic effect of estrogen and progesterone on sexual swelling as well. Therefore, if sexual swelling somewhat reliably signals ovulation, male mating activity should have been selected to occur more frequently at the second half of maximal swelling periods in which the fertile phase is likely situated – fertile phase is called a fertile window (*21*) or peri-ovulatory periods – POP; (*22*, *23*). In chimpanzees, male mating efforts, in particular, those of the alpha male who can exert the most influence on male mating competition, more concentrate their mating efforts on the fertile phase of females (*24*, *25*). These findings are in line with the predictions from the graded signal hypothesis, which implies that sexual swelling advertises a graded signal of the probability of ovulation so that it distributes males' mating attempts with their social dominance (*26*, *27*). Considering the similarity in genetics and mating systems between bonobos and chimpanzees, it is unlikely that bonobo males cannot determine the fertile phase of females unless ovulation is independent of maximal sexual swelling. However given the genetic similarity with chimpanzees and exaggerated color and size of sexual swelling in bonobos, the basic hormonal mechanism that regulates sexual swelling in bonobos may not be very different from other Old World primates (*20*).

In chimpanzees, males adopt a coercive mating strategy or mate-guarding to restrict promiscuous mating of females during maximal swelling periods (*28–31*). The efficiency of male mate-guarding probably depends on the number of males in the group (*32*) and the

degree of synchrony of ovulation (*33*, *34*). Nevertheless, it is highly beneficial for male

chimpanzees if they can determine the fertile phase (peri-ovulatory phase; POP) of females

within the maximal swelling periods, for better reproductive outcomes. In contrast to

chimpanzees, male coercive mating is missing in bonobos (*35*). Alongside which, the high

social status of females (*4*, *36*) might facilitate female choice and prevent males from using a

coercive mating strategy in bonobos as well. Therefore, male bonobos might deploy different

strategies to increase their reproductive success compared with male chimpanzees. However

even if males' behavioral responses to fertile females are different in bonobos, it would be

adaptive for male bonobos if they could judge the probability of ovulation and adjust their

behavior in relation to the ovulation probability. Following previous studies and the above

assumptions, we hypothesize that male bonobos can determine the fertile phase of females

with some precision – "males know" hypothesis. The fertile phase of this study is defined as

4-day of fertile window which includes 3 preceding days to ovulation and the day of ovulation

(*23*). This hypothesis predicts that

P1: Males' mating efforts intensive following (defined in the methods section) will

coincide with the higher reproductive quality of females. Here higher reproductive quality is

determined based on the swelling status that may indicate the probability of ovulation and

one's 'infant age. More specifically, males will focus their mating efforts on a more fertile

female whose infant's age is older, and the onset of the maximal swelling was earlier when

there are several females with maximal swelling.

P2: The number of copulations, interventions, and solicitations of copulation will

increase with the probability of ovulation and infant age.

P3: Male-male agonistic interactions will increase when there is a female who gets close

to ovulation in the party particularly when the female has an old infant. This might also result

in a higher copulation rate between high-ranking males and females with higher reproductive

quality than for low-ranking males.

Our hypothesis will be supported if male bonobos can determine/detect the fertile phase

of female bonobos based on sexual swelling and one's infant age somehow. Therefore, our

study will provide an opportunity to examine whether sexual swelling is a reliable signal

indicating ovulation and male mating efforts will be predicted from the changes in sexual

swelling in bonobos. To test this hypothesis and predictions, we will investigate the male

mating efforts including following over females, copulations, and aggressions in relation to the probability of ovulation (or sexual swelling) and infant age. After we evaluate the relevance of the hypothesis, we will suggest a male decision-making rule for mating effort allocation. We then propose that detumescence of the maximal swelling is a key phenomenon to understanding male decision-making as we assume that the mechanism of detumescence sexual swelling in Old World Monkeys – an increase of serum progesterone concentration after ovulation results in detumescence (*37*, *38*) – is probably the same for bonobos (*11*). To test the effect of detumescence on male mating effort, we will modify the previously proposed calculation of ovulation probability and fertility from the onset of maximal swelling (*23*). Finally, based on our findings and the new calculation of ovulation probability, we will propose and discuss a decision-making model, called a lottery model (evidence-driven rainmaking), that explains how males allocate their mating efforts toward females with different reproductive qualities for the best reproductive outcomes.

Results

We determined 14 ovulatory cycles and 5 anovulatory cycles in 19 maximal swelling phases from 7 females (table S1A) with the daily sexual swelling record and using 660 urine samples collected from September 2013 to April 2015. The 14 ovulatory maximal swelling phases were only found in 5 females whose infants were more than 2.0 years old (fig. S1). The earliest ovulation of females was for a female (Jk) whose infant was 24 months old. There was one possible case of ovulation from a female (Nv) whose infant was 20 months old. However, the exact date of ovulation could not be determined because of a 5-day sampling gap, so this case was not included in data analysis.

Male dominance hierarchy

Three adolescent and 7 adult males were the male subjects of this study (table S1B). Male dominance hierarchy was not stable during the 2 years of study. During Study Period 1, one male named JR temporally occupied the highest rank of the group (table S2A). In this period, however, the h☐ index was 0.84 which indicates unclear linearity of the male dominance hierarchy (*16*, *39*). During Study Periods 2 and 3, JR received many aggressions from the top 3 ranking males and his rank dropped to 5th (table S2B). To take into account this dominance fluctuation, we separated the male dominance hierarchy in Study Period 1 from Study Period 2 and 3.

Male intensive following behavior toward females

163
In total, male intensive following behavior after females – following the target female
over 5 minutes within 10m distance while keeping eyes on the target – was observed on 95
days among 250 observation days. The number of males who intensively followed females
ranged from 0 to 8 (1.4 ± 2.1 males per day). Among these 95 days, there were only 6 days
when two females were the targets of the male intensive following at the same time. To
investigate the effect of one's infant age, sexual swelling status, and ovulation on male
behavior, we used hurdle modeling that divide the data set into two parts to deal with many
zeros values – there were many days that no males intensively followed females (*40*). Further
details are provided in materials and methods.

For the first part of the hurdle model, we ran a generalized linear mixed model (GLMM)
with the data that contained all 250 observation days that included days of no male's intensive
following was observed (GLMM-1A in Table 1). We confirmed that the number of males who
intensively followed a female increased when the target female had an older infant and
maximal swelling (Fig. 2A and 2B). Female age, which probably reflects the parity of females,
did not explain the number of males who intensively followed females which suggests no
preference for older females in bonobos. This result is in contrast to that was found in
chimpanzees (*41*). The number of males and the number of females with maximal swelling
(FMS) in the daily ranging party were not responsible for the increased number of males in the
intensive following. Interaction between infant age and the swelling score was not included in
the GLMM-1A given the large proportion of data from females without maximal swelling but
with an infant.

For the second part of the hurdle model, we ran a GLMM with a reduced data set, which
only consisted of the days when females exhibited maximal swelling (205 days). With this
data set, we investigated the effect of the days from the detumescence of the maximal swelling
and one's infant age on male's intensive following behavior (GLMM-1B in Table 1). We
found that the number of males who intensively followed females increased when maximal
swelling was getting to detumescence. Infant age was also positively related to the number of
males engaged in the intensive following. More males followed females with maximal
swelling of which ovulation was detected during or right after the maximal swelling phase (14
ovulations) compared to females with maximal swelling of which no ovulation was detected
within or after the maximal swelling phase (fig. S2). Such an anovulatory maximal swelling

phase maybe because it was post-conception or anovulatory swelling or we might miss

ovulation due to insufficient data. The other predictor variables were not responsible for the

number of males engaged in the intensive following. Importantly, however, there was a

significant interaction between the days from detumescence and infant age on males' intensive

following over females. This suggests that the increase in the number of males in the intensive

following close to the day of detumescence was more rapid (or the slope of increase was

steeper) with an increase in the age of one's infant – females with older infants attracted more

males when they exhibited maximal swelling (Fig. 3A).

We also investigated how males chose a female with maximal swelling for the intensive

following when there were two or more two females with maximal swelling in the same party

using a binomial GLMM (GLMM-1C in Table 1). The days when there was only one female

with maximal swelling were excluded for this analysis. In total, we found 77 days when there

were more than two females with maximal swelling. Using this data, we tested the effect of

one's infant age, the day from the onset of maximal swelling, female age, and the number of

males in a day on male's choice among females with maximal swelling. Similar to previous

results, male bonobos chose females with older infants, and whose maximal swelling was

started earlier – significant interaction term (Fig 3B). However, the number of males did not

influence males' choice which indicates that males do not alternate their choice even when

there were same-sex competitors in a party. Although female age was significant in this

model, it could be caused by small female age variations in the reduced data set as seen in fig.

S3 – compared with infant age and the onset of maximal swelling (fig. S3A and S3B), female

age data was distributed widely and the two medians in between were almost identical (fig.

S3C).

We finally tested the effect of days from ovulation, not detumescence on male behavior

(GLMM-1D) using a data set that only included 14 ovulatory-detected maximal swelling

phases from 5 females (138 days in total). By doing so, we also tested the reproducibility of

the results from GLMM-1B. There were some sampling gaps within a maximal swelling phase

in the data and the largest gap was 6 consecutive days from Sl (fig. S1). We considered the

female continued to exhibit maximal swelling for these 6 days, so this female considered

starting maximal swelling 18 days before ovulation. As detumescence of maximal swelling

preceded the ovulation day in this cycle, we included the day without maximal swelling until

the day of ovulation in the data set. Similarly, when the maximal swelling phase was

continued after ovulation, these days were also included in the data (*24*). There were also no

sampling gaps over 1 day during POP (4 days of fertile phase) in the data set. GLMM-1D

showed similar results to GLMM-1B except for a non-significant interaction term between the

days from ovulation and infant age (Table 1). The number of males who intensively followed

a given female, increased when females were near to ovulation (Fig. 4), as well as with the

increase of infant age of females. The other predictor variables did not predict the number of

males intensively following the given female. No significant interaction between the

probability of ovulation and infant age may be related to the small size of the data set, or less

likely, that males can indeed detect ovulation precisely, so males follow a female regardless of

her infant's age when ovulation coincides with maximal swelling phase.

Day specific ovulation probability and fertility

To evaluate the effect of detumescence on male behavior, we first calculated ovulation

probability and fertility from the day of onset of maximal swelling. The day-specific

probability of ovulation and fertility were calculated using the 14 ovulatory cycles from 5

females that were used for GLMM-1D. The equation for the calculation of day-specific

ovulation probability was proposed and used in previous studies on chimpanzees and bonobos

(*12*, *23*) and also described in Materials and Methods in detail. The day-specific ovulation

probability from the onset was widely distributed from the 8th to 27th days from the onset of

maximal swelling (closed circle with black lines in Fig. 5A). The peak probability of ovulation

was on the 19th day from the onset of maximal swelling. This pattern is similar to that of the

previous report on wild bonobos from Luikotale (*12*), except that no ovulation was found

before the onset of maximal swelling. The day-specific probability of fertility – the probability

that the egg coincides with sperm – did not exceed 0.3 (open circle with red lines in Fig. 5A),

which also resembles the previous finding (*12*).

To better understand whether ovulation results in detumescent and why males stopped

following females after detumescence, we calculated the day-specific probability of ovulation

and the fertility from the detumescent day using the same equation used for Fig. 5A (see

Materials and Methods). Fig. 5B only differs from Fig. 5A in that day 0 was rearranged to the

detumescent day. Compared with Fig. 5A, the day-specific probability of ovulation was

concentrated to the detumescence day of maximal swelling and showed a clear peak 4 days

before detumescence. This result also suggests that progesterone that is released from a corpus

luteum after ovulation probably result in detumescence of maximal swelling. Due to the high

probability of ovulation before detumescence in our current study, the day-specific fertility exceeded 0.8 from 5 to 4 days before detumescence then decreased rapidly and became zero one day before detumescence. This result indicates that copulation around detumescence would more likely result in pregnancy than copulation on other days.

The distribution of the probability of ovulation and fertility regardless of calculation methods resembles the distribution of the number of males that intensively followed females. The number of males in the intensive following was widely distributed when plotted with the days from the onset of maximal swelling (Fig. 5C). In contrast, the number of males in the intensive following was more concentrated to the day to detumescence (Fig. 5D). Particularly, the median number of males in intensive following increased 5 to 1 day before detumescence and no males followed females when detumescence just started. This result suggests a possibility that males might predict ovulation before detumescence – probably by a large falling of female attractiveness which indicates females are signaling and males take advantage of it.

Male's copulation, intervention, and solicitation of copulation

Ten males copulated with the 9 female subjects 570 times in 250 days (table S1B). The frequency of copulation was 0.066 times/hour/male for males (570/8609 of total presence counted by one-hour party attendance; party presence counts) and 0.070 times/hour/female (570/8137) for females. Among 570 copulations, 492 copulations (86.3%) were observed from females with maximal swelling. We observed 440 solicitations of copulation from a male to a female, regardless of success or failure in 250 days (0.051 times/hour/male; 440 in 8609 party presence counts). Intervention of copulation was observed 80 times in 250 days (0.009 times/hour/male; 80 in 8609 party presence counts). Male-male aggression, including dyadic, tied, and more than 2 individuals, were observed 873 times in 250 days (0.101 times/hour/male; 873 in 8609 party presence counts).

To test the effect of ovulation on male copulation-related behaviors, we also used data from 14 ovulation-detected maximal swelling phases from 5 females as in GLMM-1D. We then limited the data to only contain -14 to 0 days from ovulation to remove the effect of extreme values. Minus 14 to 0 days from the ovulation were selected given the average length of the maximal swelling phase of bonobos is around 14.0 days – based on mean maximal swelling phase length from published studies (Table S3) – and one day after ovulation is still

fertile given the lifespan of an egg. The final data set contained 104 days among 250 total

observation days. In total, 209 copulations within 22 dyads (among possible 48 dyads; except

2 mother-son dyads) were situated in the data set and the frequency of copulation was 0.041

times/hour/dyad (in 5040 dyad-party hours; total presence count of a dyad found in the same

party). We observed 201 instances of solicitation in 5040 dyad-party hours (0.040

times/hour/dyad). The intervention of copulation between males was observed 42 times in

5040 dyad-party hours (0.008 times/hour/dyad). The number of male-male aggressions in the

104 days was divided by the party presences counts of all males. In total, 472 male-male

agonistic interactions were observed in 104 days (4203 party presence counts in total; male-

hours) and the frequency was 0.112 times/hour/male (472/4203).

To investigate further the effect of ovulation, infant age, female age, male age, and rank

(highest: 1, lowest: 10) on the number of copulations, interventions, and solicitations of

copulation, we ran three GLMMs (Table 2). We confirmed in the GLMM-2A that the number

of copulations increased when the ovulation day of a given female was getting close (Fig. 6A).

Infant age was negatively related to the number of copulations – males copulated more with

the females with younger infants (Fig. 6B). Male rank was positively related to the number of

copulations – high-ranking males did more (Fig. 6C), and male age was negatively related to

the number of copulations (Fig. 6D). Female age did not predict the number of copulations

(Fig. 6E). The interaction between infant age and days from ovulation was not significant.

As males did more copulation with females with younger infants whose reproductive

quality is assumed to be lower, we further investigated whether this result came from a greater

number of copulations from low-ranking males with such females. To test this possibility, we

ran a GLMM with the same data set used for GLMM-2A. In this model, we converted male

rank to a categorical variable (1-3: high-ranking, 4-10: low-ranking), and a three-way

interaction between male rank (category), infant age, and the days from ovulation was also

included. Female age was removed as the model failed in converging. The top 3 ranking males

(JR, NB, KT in periods 3 and KT, NB, GC in period 3) were selected as the three males and

their mother Jk (JR's mother) and Ki (NB and KT's mother) competed for alpha position

during the study period. The top 3 ranking males occupied 78.9% of all copulations (165 in

209 copulations). GLMM-2B showed a similar tendency of GLMM-2A in that significance

and the effect of variables were identical in between (Table 2). Importantly, the 3-way

interaction term was significant for high-ranking males. This significant interaction indicates

important contrasts of males' copulation depending on their rank. The ovulation probability

was responsible for the increased number of copulations for the high-ranking males but not for

low-ranking males. Such skewed copulation tendency was mostly because even high-ranking

males could copulate more with a female with younger infants (fig. S4). This result suggests

that females with low reproductive quality also copulate more frequently with high-ranking

males, not with low-ranking males. Low-ranking males, therefore, have less chance to

copulate even with females with low reproductive quality which is different from the

prediction of the estrus sex ratio hypothesis. This result also demonstrated that females are

choosier when their reproductive quality gets high and reduces the copulation rate even with

high-ranking males.

The number of interventions of copulation between males (GLMM-2C in Table 2)

increased when the ovulation day was getting close (Fig. 6F) and increased for females with

older infants (Fig. 6G). High-ranking males intervened more frequently in other males'

copulation attempts (Fig. 6H). It is probably easier for high-ranking males to intervene than it

is for other males. Male age and female age were not related to the number of interventions of

copulation (Fig. 6I, 6J). This result indicates that even though older females have a higher rank

than younger females, the intervention of copulation between males may not be influenced by

the rank of females in copulation attempts.

Males solicited copulation more often when the day of ovulation of a given female was

about to occur (Fig. 6K, GLMM-2D in Table 2). Infant age did not predict the number of

solicitations of copulation by males (Fig. 6L). This finding suggests that the increased number

of copulations in females with younger infants in GLMM-2A was not due to increased

solicitation of copulation by males. Both high-ranking and younger males solicited copulation

more frequently (Fig. 6M, 6N). Female age was not related to the number of solicitations

which suggests that older females are not more attractive for male bonobos.

Finally, we tested whether the number of females with maximal swelling on a given day

was related to the number of male-male aggressions using all 250 observation days using a

generalized linear model (GLM-2D in Table 2). We included the number of females with

maximal swelling and the existence of the male intensive following as explanatory variables.

To control the number of males of the day and observation hours, we include the two variables

in the offset term of the model. We found that the number of male-male aggressions on a

given day increased on days when the male intensive following was observed (Fig. 7A). The number of male aggressions was also increased with the number of females with maximal swelling of the day (Fig. 7B). These results suggest that the availability of receptive/attractive females with maximal swelling in a party does not influence male-male aggression which contrasts with the prediction of the estrus-sex ratio hypothesis.

Discussion

In the current study, we showed that male mating efforts were skewed toward females with older infants and whose maximal swelling was close to detumescence which we defined that the female had higher reproductive quality. Males, regardless of rank, intensively followed females with higher reproductive quality. More specifically, the number of males in intensive following increased for females whose maximal swelling was close to detumescence and whose infants were older. When there were more than two females with maximal swelling, males followed females with older infants whose maximal swelling started earlier. The significant interaction between infant age and sexual swelling makes sense since males did not follow females even with maximal swelling when their infants were younger than 2 years old. It is worth noting, however, that when the data set was reduced to 14 ovulation cycles from 5 females, the significance of the interaction term disappeared. This non-significant interaction suggests that male bonobos, regardless of infant age of females, may be able to determine an ovulatory maximal swelling phase more precisely, though this result could also be due to sampling bias in the reduced data set.

Despite increasing the number of copulations close to ovulation day, high-ranking males copulated more frequently with females with younger infants, contrary to the results of males' intensive following. High-ranking males also intervened in the copulation attempts of other males when the other males tried to copulate with females with older infants whose ovulation day was approaching (higher reproductive quality). These results suggest that male-male competition for access to females with older infants is more intense so the copulation frequency might be lower between them, or females are more selective for mating when they have higher reproductive quality. Males solicited copulation more often from females whose ovulation was approaching. However, solicitation frequency did not change with the infant age of females. Male-male aggression also increased when there were more females with maximal swelling. There were more aggressions between males after controlling the number of males in the party, especially on days when males' intensive following was observed. Contrarily to a

prediction from the estrus sex ratio hypothesis, this result suggests that the availability of

receptive/attractive females is not an important influencing factor determining male-male

agonistic interaction in bonobos.

Overall, our results support the predictions from the males know hypothesis but did not

support the estrus sex ratio hypothesis. The three predictions of the males know hypothesis

regarding male mating were well related to the reproductive quality of females except for the

effect of infant age on females in the ovulatory maximal swelling phase. This exception,

however, makes the hypothesis even stronger as this result implies that males can recognize an

ovulatory swelling cycle. Therefore, our study supports the notion that male bonobos can

determine the fertile phase of females with some precision and they allocate mating efforts to

the females with higher reproductive quality. These results are also in line with previous

findings in other primate species where females have exaggerated sexual swelling, including

chimpanzees and Old World monkeys (*24*, *42*, *43*). We suggest, therefore, that male bonobos

use information from sexual swelling and one's infant for decision-making on mating effort

allocation, and the sexual swelling of females is a reliable indicator of ovulation in bonobos.

Male intensive following and fertility of females

Males intensively followed females with maximal swelling more than females at other sexual

swelling stages. When the female had an older infant and the sexual swelling was near to the

detumescence, a higher number of males intensively followed the female (GLMM-1A and

1B). Therefore, males appear to be able to distinguish females with higher reproductive

quality even when several females exhibit maximal swelling which suggests a possibility that

males use not only sexual swelling but also other factors from females such as lactation status,

olfactory and behavioral cues.

Although males may be able to make a better choice by combining the information from

sexual swelling and reproductive history, there can be two females with similar reproductive

quality at the same time by menstrual synchrony. If there is some degree of the synchronized

menstrual cycle (maximal swelling as well), it would be difficult for high-ranking males to

monopolize copulation attempts against other males. Here, some degree of synchrony

represents loose synchrony in which the ovulation date of each female coincides within a or

two weeks so thus their maximal swelling also overlaps. As seen in the table S1A, in E1

group, two females often gave birth in the same month (within 2 weeks) which suggests a

possibility of loose synchrony of the menstrual cycle. If there is such loose synchrony, it can also allow females to test the physical durability of males as higher-ranking males need to maintain mating efforts for longer periods. It may also reduce female-female competition over mates by differentiating the short fertile phase (within 4 days from ovulation). This loose synchrony in menstrual cycles, therefore, can play a counterstrategy against the male's monopolizing mating strategy and allow females to exert control on males. In such a case, we could recall the estrus hypothesis that the high availability of receptive/attractive females in the bonobo group has some regulative roles in male-male competition. However, a narrow synchrony window may result in a competitive situation between females and so is unlikely beneficial for females.

We confirmed that more males intensively followed the female whose ovulation was about to occur (Fig. 4). Our way of determining ovulation may be strict; ovulation was not determined unless the average of PdG was above the threshold (mean ± 2SD) for 10 days after the ovulation day. This might be why we could not detect ovulation from females with young infants whose age was less than 2, contrary to the previous finding (*12*). However, ovulation in those females might not have much influence on male behavior since males were not interested in females with such young infants, even when the females had maximal swelling (Fig. 3).

As seen in Fig. 5D, the number of males in intensive following seems to decrease one or two days before detumescence. This suggests that male bonobos are better than human observers in evaluating subtle changes in sexual swelling. Or males may have some ability to detect ovulation based on olfactory and other visual cues (*44*). However, such discussion goes beyond the scope of this study. Given that detumescence of maximal swelling is a good predictor of male responses, it is likely that male bonobos also rely on the sexual swelling as an important indicator of ovulation just like males of other primate species with exaggerated sexual swelling such as yellow baboons (*43*), Barbary macaques (*45*), and chimpanzees (*24*). A more important question might then be whether the seemingly unpredictable ovulation of female bonobos, possibly concealed by the maximal swelling phase (*12*, *13*), would be manipulative enough to confuse male mating decision making and to regulate male-male mating competition.

Male intensive following behavior was not directed toward females whose infants were

less than 2 years old, except for 2 days during Nv's maximal swelling phase and that was only

by 2 males (when her infant was 21 months old). Male intensive following sometimes

continued all day, from the bed site in the morning to dusk when group members started

making their beds for the night. In this case, males would even make beds around the target

female. This might be because high-ranking males do not want males other than themselves

sneaking copulations during the night or early next morning. Such all-day-long male intensive

following was most noticeable when the sexual swelling of females with old infants (over 4)

reached its maximal swelling phase. This might be related to the inter-birth interval, which

was 4.8 ± 0.8 years in this study group (*20*). In the future, it will be interesting to see if the

duration of intensive following of a given male varies depending on the reproductive quality

of the female. Females may be also able to test the durability of males that participate in all-

day-long intensive following to make a better mate choice. Together with the possibility of the

loose synchrony of ovulation, such a female mating strategy might be a challenge, particularly

for the highest-ranking male who would have to invest more energy for longer periods than

lower-ranking males (*46*).

In all models (GLMM-1A to 1D), the number of males and females with maximal

swelling in a party did not predict the number of males who intensively followed females. This

indicates that the spatio-temporal distribution of males each day was not responsible for the

increase in the number of males in the intensive following. This strengthens the finding that

male bonobos selectively choose the target of intensive following rather than just following

any female with maximal swelling.

Male mating efforts and competition for access to fertile females

As predicted from the "males know" hypothesis, the number of copulations within a

dyad increased when ovulation was about to occur. However, infant age had a negative

influence on the number of copulations, which is opposite from the prediction of the

hypothesis. Females with older infants may be more selective for their choice of copulation

partners, or that mate guarding between high-ranking males is intense for females with older

infants so they cannot copulate so easily. Male rank was also positively correlated with the

number of copulations. High-ranking males may be chosen more often by females for

copulation or high-ranking males increase their chance of copulation by staying closer to the

female with high reproductive quality (*4*, *47*). Occupying spatio-temporal proximity with the

female with high reproductive quality can also be an effective strategy of male bonobos, given the high possibility of female choice in bonobos because of the high social status of females (*4*, *48*, *49*). A negative binomial GLMM using the proximity data from the individual focal sampling confirmed that the number of females with maximal swelling within 1 meter was negatively related to male age (Z = -3.46, *SE* = 0.21, *P* < 0.001) and positively related to male rank – higher-ranking males had more females around (Z = -2.01, *SE* = 0.18, *P* = 0.044). This result indicates that it is not always the high-ranking males who are closer to more females with maximal swelling and their access to copulation was not simply explained by the availability of females with maximal swelling in proximity. This result is also consistent with GLMM-2A and 2B that younger males copulated more frequently (Table 2). The high frequency of copulations in younger males, therefore, might be due to the close spatio-temporal relationship with females. Young males might get extra benefit from having their mother in the group, since the mother can influence the availability of females around their son directly and indirectly (*4*, *16*, *50*, *51*).

Although copulation in bonobos is assumed to be open for low-ranking individuals (*4*), our results suggest that male-male competition over mates results in a skewed number of copulations toward high-ranking males with highly fertile females. Such skewed mating causes the exclusion of low-ranking males from mating. The estrous sex ratio hypothesis (*4*) predicts that low-ranking males may seek access to females even with lower reproductive quality to maximize chances of copulation and reproductive success. If this is the case, low-ranking males may copulate more with lower reproductive quality such as females with a younger infant but with maximal swelling. However, our results do not support this prediction. Our results instead, demonstrated that high-ranking males occupy most of the copulations during a fertile window and they also copulate more with females with younger infants (GLMM-2B, and fig. S4). It was not because males solicited copulation more to females with younger infants (GLMM-2D). It is possible that females with a young infant are more responsive or females with an old infant are more selective about responding to a copulatory invitation. Females may also prefer high-ranking males to low-ranking males for mating. Investigating female choice will therefore clarify how female mate choice shapes male-male competition and its outcome that results in variations in male reproductive success depending on male rank.

It was suggested that paternity in bonobos seems to be skewed for high-ranking males (*52*) as found in chimpanzees (*53*). Recent studies demonstrated that the paternity of bonobos is indeed skewed to high-ranking males even more than that of chimpanzees. Our current study suggests that male bonobos can discriminate a narrow fertile window using female sexual swelling and infant age and that this ability could explain very skewed paternity in bonobos. Together with the previous finding that male rank is important in bonobos for mating success (*16*), we suggest that male-male competition is also an important characteristic of bonobo society, and male-male competition may cause reproductive skew toward high ranking males who are more durable and can monopolize mating opportunities with females with higher reproductive quality. Such monopolization is probably not solely to the competitive ability of the male, but also to the female preference for the male with high durability in pursuing mating efforts as well.

The number of interventions by high-ranking males increased to the females with older infants when their ovulation was about to occur. Males may also intervene in the copulation of young females more, though this is not statistically significant ($P = 0.053$, GLMM-2c in Table 2). If this was the case, it might be because old females are usually high ranking (*48*, *49*) and their high social status might restrain males from intervening in their copulations. High-ranking and young males solicited copulation from females. Male's solicitation of copulation also increased with ovulation probability. Unlike the results of the result of copulation, solicitation of copulation did not increase with the infant age which denies one of our second predictions (P2). This finding also suggests that males do not discriminate between females for copulation if females exhibit maximal swelling. However, females' willingness to copulate may vary depending on their reproductive quality or probability of conception.

The number of total male aggressions on a given day increased when male intensive following behavior was observed (Fig. 7A). More importantly, the number of females with maximal swelling resulted in a greater number of male-male aggression. The increase in male aggressions might be simply due to the increase in the number of males in the daily ranging party as the number of males increased when there were more females with maximal swelling (Poisson GLM; $\chi^2(1) = 43.18$, $P < 0.001$). However, since the number of males was included as an offset term to control the effect of the number of males in GLM-5E and the number of males did not vary depending on the existence of male's intensive following (Poisson GLM;

$\chi^2(1) = 1.25$, $P = 0.26$), male mating competition over access to ovulating females probably

caused more male-male aggression in the current study.

Male mating strategies to cope with the low predictability of ovulation

We found that the onset and detumescence of maximal swelling were significant

predictors of male mating efforts. Therefore, males seem to rely on the sexual swelling of

female bonobos as a signal to make decisions for mating. However, as shown in Fig. 5A, the

day-specific probability of ovulation was distributed widely from the onset of maximal

swelling. The predictability of ovulation from the onset of the maximal swelling phase is,

therefore, also low in the Wamba population as found in the Luikotale population (*12*).

Although sexual swelling of bonobos was suggested to deviate from that of other primates in

that ovulation does not match well with maximal swelling (*12*), the physiological and

hormonal mechanisms that result in such deviation in bonobos are yet proposed. However, the

role of estrogen and progesterone that regulates sexual swelling in bonobos may have not

changed much from that of chimpanzees (*20*). Moreover, although maximal swelling could

play an important role in female-female interactions (*51*), what is the benefit to keep this

costly signal if it does not indicate ovulation? Therefore, we may have missed some important

keys, or we are stuck with noise or artifacts from the previous model to calculate daily

ovulation probability in evaluating sexual swelling of bonobos.

Bonobo females resume swelling cycles in early lactation periods: 225.4 days on

average after giving birth even when their estrogen level is still low (*20*). In addition, as seen

in fig. S5, some females have very long maximal swelling around 30 days, particularly when

females are with younger infants. Such variation may not be due to the small sample size in

the current study as other studies also reported such variations in the length of maximal

swelling (*11*, *12*). We propose, therefore, that the inter-and intra-individual variations in the

length of the maximal swelling phase are the reason (or the source of noise) why the

probability of ovulation calculated from the onset of the maximal swelling is low in bonobos.

In other words, as the equation calculate the daily probability by summing up all probabilities

of a given day from all defined ovulatory cycles, a great individual variation in the length of

the maximal swelling phase results in poor predictability of ovulation. In such a case, poor

predictability of ovulation is not because ovulation deviates from the maximal swelling phase.

Instead, it is the inter-and intra-individual variations in the length of the maximal swelling

phase that cause such low predictability of ovulation. So even if ovulation is usually situated

last half of the maximal swelling phase as in the previous study (*11*, *17*) and this study, the
day-specific probability of ovulation and fertility is low in bonobos.

The equation to calculate day-specific ovulation probability and fertility in the current
study (see method section) was proposed in a study on chimpanzees (*23*). Although this
equation has provided a solid framework to understand and investigate the relationship
between sexual swelling and ovulation, it does not take into account a greater variation in the
length of the follicular phase compared to the luteal phase between and within an individual
across menstrual cycles (*54*). Such variation in the length of the follicular phase would
inevitably cause low predictability of ovulation when using the equation as seen in the current
study. This in turn may mislead researchers to conclude that ovulation is unpredictable based
on the change in sexual swelling particularly for the species such as bonobos of which the
length of the maximal swelling phase varies much. Nevertheless, our current study does not
deny the notion that the predictability of ovulation from sexual swelling is low in bonobos.
The predictability of ovulation from the onset of maximal swelling is low in the current study
(Fig. 5A). As demonstrated in the current study and paternity data from the wild bonobo
population (*14*, *15*, *55*), males may be just able to cope with well such low predictability of
ovulation for better reproductive outcomes. How would males be able to match their mating
efforts near to ovulation and detumescence of maximal swelling if ovulation is barely
predictable in bonobos? In addition, if ovulation is such difficult to predict, how paternity can
be skewed to high-ranking males in bonobos? Our result implies that sole female choice
cannot explain such paternity skew.

Although sexual swelling is, in general, a reliable signal indicating ovulation probability
of females, the reliability may be lowest in bonobos (*12*, *13*, *20*). Despite such low reliability,
detumescence is a reliable indicator that predicts male behavior in bonobos as seen in Fig. 5B
so thus suggesting that detumescence can be a reliable indicator of ovulation. However, even
if ovulation causes detumescence of maximal swelling in bonobos, male bonobos may not be
able to predict ovulation (or detumescence) of females due to the greater inter-and intra-
individual variation in the length of the maximal swelling phase. We propose a decision-
making model of males to cope with such difficulties in predicting ovulation. This model does
not need to assume that males can pinpoint the detumescent day although such a possibility
exists in bonobos. Instead, males can simply follow the females with older infants and
maximal swelling over days if the detumescence does not happen on that day (today) as the

day-specific probability of ovulation will get higher and higher over days until detumescence.

Our model would be also applicable to understanding male mating effort allocation, not only

in species where sexual swelling is found but also in species that have any graded signals

indicating ovulation (e.g., changes in face colors, odors, behaviors with ovulation probability).

The model assumes that male bonobos, as well as males in the species proposed above, use

three-layered rules to allocate their mating efforts for the best reproductive outcome.

The first rule in the basement layer is called the 'just-do-it' rule. This layer assumes that

males are highly motivated for copulation, which is plausible considering that more mating, in

general, gives males better reproductive outcomes (56). Therefore, male bonobos seek any

copulation opportunities and whenever an opportunity is available, males are willing to engage

in copulation. This rule could be predominant particularly for young males throughout their

puberty as more copulations might allow them to become better at copulation as well.

The second rule in the second layer is that males allocate more mating efforts for access

to females with older infants (called 'check-infant'). This rule requires an assumption that the

attractiveness of females with maximal swelling to male bonobos changes depending on one's

infant growth (infant age). This rule does not necessarily assume that males should recognize

infant age as humans do. It can work if the male can somehow discriminate against females

with different reproductive histories based on some signal or cue related to one's infant

growth. Holding this rule by males does not necessarily result in skewed copulations toward

the females with older infants, since the male-male competition and female choice might

influence the outcomes of copulation.

The final rule in the third layer is more complicated and arguable. This rule is called

'rain-making' as males are following females until the detumescence of maximal swelling. If

maximal swelling (or any indicative signal of ovulation) continued from yesterday, then males

simply increase mating efforts or keep a similar degree of mating efforts toward the female on

that day compared to yesterday. Males, in particular high-ranking males, might skew their

mating efforts more toward females with older infants and whose maximal swelling started

earlier or is close to detumescence. This does not mean that they can predict detumescence or

ovulation. They can make wrong decisions, particularly when the ovulations of several

females are closely synchronized. It is also plausible that males are interested in a female from

the onset of her maximal swelling phase, particularly for a female accompanied by an old and

weaned infant. For such a female with higher reproductive qualities, males maintain their
mating interests and efforts until the day of detumescence. After detumescence, they will not
pay attention, since the probability that the sexual swelling gets detumescent before ovulation
is low (though not zero). Since social-ecological factors, such as the presence of the mother,
party composition that influences male relative rank, vary over time for males, these social
dynamics can cause individual specificity of the rainmaking rule. Therefore, the strategy of
each male for the 3rd layer might vary between individuals as well as within an individual
over his lifetime.

Overall, this male mate decision-making model explains the results of this study that
male mating efforts get more intense around the time of ovulation without an assumption of
male's ability to pinpoint ovulation day. We call this hypothetical model a "lottery" model
since it is somewhat similar to the way how lotteries work. In lotteries, as the prize money
increases when there are no winners for several weeks, more people buy lottery tickets. This
might increase the chance of a winner coming out in the whole game over time. Although the
probability of ovulation does not change due to an increase in male interest and efforts, such
an increase will help females have better reproductive outcomes. It is also possible that males
can predict the fertile phase more precisely using multiple signals (or cues), including female
behavior and olfaction. This might also explain the subtle decrease in the number of males
engaged in intensive following even before detumescence (Fig. 3A, 5D). However, our data is
not enough to test this possibility.

It was reported that the size of sexual swelling gradually increased until the fertile phase
(POP) in chimpanzees (24, 57) and olive baboons (58). Males might be, therefore, able to
discriminate the subtle change in the size of sexual swelling (24). However, an increase in the
size of sexual swelling may simply reflect the gradual increase of serum estrogen level until
ovulation, and males do not need to discriminate precisely the subtle size variation within a
maximal swelling phase (45). In such a case, the gradual increase of size within the maximal
swelling phase can only have significance when it is presented to the males with a continuous
time scale. This possibility will be examined in future studies. With a more precise
measurement of female signals, the future study will clarify whether males can discriminate
subtle increases of size and color (or even olfactory cues) and adjust their mating efforts
precisely to the narrow fertile window. This will provide us with a valuable opportunity to

understand how males have evolved to recognize and respond to females' signaling on
ovulation probability not only in bonobos but in primate species with such graded signals.

An increase of serum progesterone concentration caused by ovulation results in
detumescence in primates with sexual swelling (*37, 59–61*). We do not know yet any
mechanism standing for the physiology, which does not follow this mechanism or that causes
considerable deviation if there is. If there is a mechanism that can explain how an increase of
progesterone would not result in detumescence, it would be important to look deeper into the
reason why. However, it is unlikely that the physiology of sexual hormones has changed much
only in bonobos. Instead, reproductive events of a female, such as lactation, pregnancies, and
early miscarriage of pregnancy – early loss (*62*), might be related to the great variation in the
length of maximal swelling phase as these events add more instability to the follicular phase in
primates including bonobos. For example, as seen in fig. S6A, there could be a short post-
conception swelling (around 1 week) right after implantation within 3 weeks from fertilization
(ovulation). Such post-conception swelling is probably due to a surge of estrogen being earlier
than a surge of progesterone after implantation and such relative concentration difference
between estrogen and progesterone will be a plausible (or the only) mechanism of post-
conception swelling in bonobos and other primate species with sexual swelling. In addition, if
an early loss occurs after the emergence of post-conception maximal swelling as in fig. S6B,
this post-conception maximal swelling can be defined as anovulatory by researcher. Male
bonobos may also be confused with such post-conception swelling as losing mating change is
more costly than allocating mating efforts even for the post-conception swelling. If such early
miscarriage frequency in bonobos is similar to that of humans, over 20% (*62, 63*), this will be
an important reason that causes sexual swelling deviates from ovulation as well. Together with
the proposed possibility of high sensitivity to estrogen of cells in bonobo sex skin (*20*), the
physiological variations from one's reproductive history might be, therefore, important
sources of the noise which causes inter-and intra-individual variations in the length of
maximal swelling. Although this variation in the length of maximal swelling causes the loose
relationship between the probability of ovulation and sexual swelling in bonobos, males do not
have much difficulty meeting their mating efforts to ovulation as suggested in the current
study. Further studies on molecular level hormonal sensitivity of sex skin and hormone–
swelling dynamics induced by fertilization events will elucidate the source of noise and will
help us to understand better the mechanism and evolution of such nosy ovulatory signals in
bonobos.

In this study, we confirmed that male-male competition also exists in bonobos, as found previously (*16*), and male mating competition is important for males to access females with high reproductive quality. This result suggests that male mating efforts and interests are not manipulated by the poor predictability of ovulation from the onset of the maximal swelling. Instead, as we suggested in the three-layered model, males might be able to cope with the poor probability of ovulation and their mating efforts using simple rules based on female sexual swelling and infant growth. Although the high availability of receptive/attractive females achieved by the more frequent and prolonged maximal swelling in bonobos may have some regulative role in male mating competition as suggested in the estrus-sex ratio hypothesis, we could not find evidence to support it. Together with the possibility that females are getting more selective when their infant gets older, future studies should examine and clarify the regulative role of sexual interactions between males and females with low reproductive quality.

Chimpanzee society is characterized as male-centered and aggressive. However, bonobo society is recognized as female-centered and peaceful (*3–5*, *49*, *64*). The prolonged receptivity/attractiveness from the early resumption of maximal swelling and long-lasting maximal swelling phase of bonobos was suggested as the key to understanding the difference in social characteristics between chimpanzees and bonobos (*4*, *9*, *12*, *51*). Although more open chances of mating even for low-ranking males due to higher availability of receptive/attractive females in bonobos may lower male-mating competitions in bonobos (*4*, *6*), it has not yet been tested thoroughly whether more chances of copulation for low-ranking males lower male-mating competition in bonobos. As suggested in the current study, it becomes questionable whether the prolonged period of maximal swelling phase indeed conceals ovulation in bonobos. In addition, we do not know whether the concealed ovulation indeed lowers mating competition in bonobos as when ovulation is concealed, males may need to allocate more efforts toward mate guarding and sexual activity of females for a longer time (*54*, *65*). The main perspective and insight from this study are therefore that the extended periods of maximal swelling through frequent and prolonged maximal swelling phases are not sufficient to explain the social differences between chimpanzees and bonobos.

In contrast to female chimpanzees (*66*), female bonobos make coalitions against male aggression (*49*). The female-female cooperation may contribute to the high social status of female bonobos and the high social status of females in turn may allow more room for female

choice that eases male-male competition over mates. We suggest therefore that the high

operational sex ratio and lower predictability of ovulation may not be the only causation of

less aggressive and more tolerant (namely peaceful) bonobo society compared with that of

chimpanzees. Instead, milder and more tolerant characteristics of bonobo society and less

restricted sexual interactions in bonobos can be rather the outcome from the high social status

of females which is as high or equivalent to that of males (alpha female hypothesis).

Nevertheless, a high operational sex ratio in bonobos because of prolonged sexual swelling

may play some regulative roles in male competition. As a result, we propose that to

comprehend the characteristics of bonobo society, we must investigate the socio-ecological

factors underlying how females achieved high social status in their society, as well as the

mating competition and strategies of females and males.

743
744　**Materials and Methods**
745
Study site and subjects

We conducted the study on a fully habituated free-ranging wild bonobo group, called

E1, at the long-term bonobo field site at Wamba (00˚ 11☐ 07.6☐ N, 022˚ 37☐ 57.5☐ E;

WGS84) in the northern sector of the Luo Scientific Reserve, D. R. Congo (*7*, *67*). The

group's home range consists of primary, secondary, and swamp forests (*68*). Although there is

no clear division of dry and wet seasons, the amount of rainfall and fruits availability are

lower in January and February than in the other months (*69*). E1 group comprised of 32 to 38

individuals, including 9 adult and 3 adolescent immigrant females and 7 adult and 4

adolescent males during the study periods from 2013 to 2015. The age classes of females and

males were defined following a previous study (*70*). Focal data were collected on 9 adult

females and 7 adult and 3 adolescent males. Further information on the study subject is in

table S1A and S1B. Three adolescent females who had no previous birth records were not the

subject of the current study. Although one adolescent male became 8-years-old in October

2014, he was not the subject animal, since he had not shown any sign of dominance

challenges, which can be a good indicator for adolescence (*16*, *71*). All adult females had

immigrated into E1 group and had at least one successful birth after their immigration (*72*).

Behavioral observation

The first author RH collected behavioral data on adult males and females during three

study periods (Period 1; September 2013 to February 2014, Period 2; July 2014 to September

2014, Period 3; November 2014 to April 2015). Bonobos were followed from their bed site, located at around 6 AM, until they made a new night bed, usually after 5 PM. When they split into several parties, the largest party was followed as far as possible. To control for the effect of the number of available group members, party composition was recorded every single hour following the one-hour party method (*73*). Daily ranging party size and membership were defined as the number and identities of all individuals who were found during a given day when the total observation time of the day was longer than 2 hours.

For continuous focal sampling, one of the bonobos in the party was randomly selected based on a random order which was previously generated before field work began. RH and one research assistant searched for the first target focal animal in the random order for at least 30 minutes, after bonobos were located in the morning. If we could not find the first one, we chose the second one in the order, and if we could not find the second one either, we moved to the next one and so on. To compensate for the lack of focal sampling data due to the fission-fusion social system of the species (*74*, *75*), we also gave priority to some individuals with fewer data for focal animal selection. One focal session continued for 20 minutes and was terminated when a focal animal was out of sight for more than 5 minutes within a focal session. Only focal sessions that were conducted for longer than 15 minutes were included in data analysis. After an individual was followed once, it was not followed for another focal sampling before at least 100 minutes. All activities involving the focal animal, including feeding, moving, resting, and social interactions were recorded during a focal session. To estimate neighboring individual patterns, the names, and behaviors of neighboring individuals within 5m of the focal animal were recorded at 5-minute intervals, including a scan at the beginning of the focal session (5 scan points within a focal session). All rare events, such as sexual and agonistic interactions, were also recorded as far as possible during the party following. We recorded the individual ID of the participants of the interactions as well as time and duration. To minimize the sampling error caused by ad-libitum sampling, all copulations, genito-genital rubbings, and agonistic interactions (e.g., bite, hit, displacing, and chase), were recorded whenever observed by RH and at least one research assistant.

Ejaculation during copulation was not always possible to confirm. Therefore, complete heterosexual intercourse without any interruption by others was counted as copulation. The intervention of the copulation included direct aggressions toward the male who solicited copulation from a female or direct intervention of the copulation which targeted the male in

copulation. Intervention directed to females was only 22 cases and most of them caused

counter-aggression of females or were ineffective so only 5 cases were succeeded in stopping

copulation. The solicitation of copulation was counted only when the clear target was

identified. Male bonobos often have an erect penis during feeding while watching females,

and so only the presence of erection was not counted as a solicitation. For solicitation, an

erection should coincide with other behaviors such as directed hand or body gestures (*76*) or

shaking, bending, or dropping branches or twigs to get attention from the target female.

We also recorded agonistic interactions whenever there was a clear target, regardless of

physical contact. Therefore, undirected display behaviors, such as charging and branch

dragging, were not considered agonistic interactions (*77*). When a counterattack was

performed by the target within 5 seconds after the directed aggression to the target, we

considered it as a tied interaction. To evaluate males' interests toward females in the party, we

defined **male intensive following** behavior as when a male keeps close distance, within 10m

in maximum, to the target female for more than 5 minutes while moving, without losing the

target female from his sight. If a male's following behavior fulfilled these criteria, we

considered this male was intensively following the target female.

In total, we followed bonobos for 1462.3 hours on 255 days (5.73 $\pm$ 1.25 hours per day).

The total focal following time for all individuals was 280.22 hours (858 focal sessions: 14.75

$\pm$ 2.01 hours per individual). The average length of a focal session was 19.6 $\pm$1.0 minutes and

each focal session consisted of 4.88 $\pm$ 0.34 scans (4189 scans in 858 focal sessions). Among

255 days we excluded two days on which we could follow bonobos for less than 1 hour. There

was one day with very heavy rain so that no males were confirmed, and two days on which we

could not see any females. These 5 days were removed from the data analysis, except to

calculate the male dominance relationship.

Assessment of sexual swelling

Daily variation in sexual swelling of female bonobos was scored in relation to the

firmness and size of each individual's swelling (*8, 51*). The sexual swelling status was

assigned to one of three categories. Fig. 1 shows an example of changes in sexual swelling of

one individual, named Fk. Swelling 1 (Fig. 1A) is for non-swelling status; swelling 2 (Fig. 1B)

is for intermediate swelling status; swelling 3 (Fig. 1C) represents maximal swelling status.

Unlike female chimpanzees, the sexual swelling of female bonobos is always visible, even in

the non-swelling status. Therefore, the practical definition of non-swelling status (swelling 1)

is that the swelling is within range of its minimum size. The swelling size variation of an

individual between swelling 2 and 3 was sometimes not definitive. Therefore, firmness and

shininess of the surface and movement during locomotion of the sexual swelling were the key

features for distinguishing swelling 2 and 3. The detumescent day was set to the first day when

the maximal swelling started shrinking so lost its firmness and shininess when the swelling

score became 3 to 2. The maximal swelling phase was defined from the onset of the maximal

swelling to the end of maximal swelling. If sexual swelling recovers from 2 to 3 within 3 days,

it was considered a maximal swelling phase continued. Since this study followed a

conventional method (*8*) for scoring sexual swelling stages, there could be subjective bias on

scoring between observers. To reduce observation errors and bias on sexual swelling scoring,

at least one researcher and two research assistants discussed and decided the sexual swelling

status of each female during field observations at the daily meeting every evening.

Urine sample collection and hormonal measurement

Urine was collected using filter paper (Whatman #1 Ø 5.5cm) during a day. After a

female finished urination, RH or one research assistant collected urine on leaves of terrestrial

plants or small trees. We put the tip (around 1/4 of the whole filter paper) of the filter paper

into the urine drops and waited until it absorbed to around 2/3 of the paper. By putting only

the tip of the paper into the urine, we avoided contamination of dirt from leaves. Not to disturb

bonobos, we collected urine after the bonobos left the place so that we could keep at least 5m

distance between the collector and bonobos. When the urine was mixed with their feces or

others' urine or feces, we did not collect the urine. Further information on our sampling

collection and handling is found in a published work (*20*) and the supplementary material

(Urine sample handling after collection). We used enzyme immunoassay (EIA) to measure

urinary metabolites of estrogen (E_1C) and progesterone (PdG). In total 660 urine samples from

9 females were successfully recovered urinary E_1C and PdG and used to determine ovulatory

cycles. The number of urine samples and examined swelling cycles are available in table S1A.

Further details about the EIA analysis and validation are available in the supplementary

material (Enzyme immunoassay and its validation) and a published paper (*20*).

Defining the fertile phase (POP) and probability of ovulation and conception on a

given day within a swelling cycle

The date of ovulation was estimated based on sustained urinary PdG rise above the

baseline (*20*). In brief, the baseline was defined as the mean of urinary PdG concentration of

the 10 days preceding a given day. When the PdG concentration on the given day was more

than two standard deviations above the baseline for 3 successive days within a week, the first

day of the sustained PdG rise (over 2 standard deviations) was considered ovulation date. We

used the 4-day fertile window, which starts from 3 days prior to ovulation, based on findings

that sperm are fertile within three days in the reproductive tracts and eggs are fertile within 24

hours after ovulation (*78, 79*). This criterion has already been used in several studies in several

primate species including wild chimpanzees in Taï (*23*) and wild bonobos in Luikotale (*12*).

To calculate the probability of ovulation on a specific day within a cycle, we used the

following equation proposed previously (*23*). When the onset of maximal swelling phase was

not observed (6 cases) or the detumescent day was not observed (4 cases), the start day and/or

the end day was determined as to when a female was first observed with maximal swelling

and/or the last day when she was observed with maximal swelling.

$$P(T = t) = \frac{n_t}{n}, t = 1,2,3\ldots$$

In this equation, $P(T = t)$ refers the probability of ovulation on a given day; t refers to a

specific day within the maximal swelling phase which starts from the onset of maximal

swelling; n_t is the total number of cycles in which ovulation occurred on day t; n is the total

number of ovulatory cycles (14 in the current study).

We also calculated the daily probability of fertility (the probability that the sperm can

meet the egg; fertilization) using the equation in the same study (*23*).

$$P(X(f) = 1) = \sum_{t=f}^{f+3} P(T = t)$$

$P(X(f) = 1)$ refers to a probability of fertility (conception) on a given day within the

maximal swelling phase. More simply the probability of fertilization on a day is the sum of the

probability of ovulation on the given day f and the following 3 days of from the given day

f that are calculated in the previous equation. By following the same equation used

previously, we can directly compare our results between species, as well as between

populations of the same species.

Estimating male dyadic dominance hierarchy

Male dominance metrics were made based on dyadic agonistic interactions. Agonistic

interactions with physical contact included biting, hitting, and trampling. Agonistic

interactions without physical contact included chasing, charging, and directed display

(sometimes with branch dragging). We only used dyadic agonistic interactions which had a

clear outcome of loser and winner. A loser was defined when the recipient of the aggression

showed submissive behaviors such as grimace, screaming, running away, or retreating. We did

not include tied agnostic interactions for dominance calculation. In total, we observed 844

dyadic agonistic interactions. Only 16 dyadic interactions were followed by counter-

aggressions. Therefore, in total 828 dyadic interactions (p1: 314, p2: 162, p3: 352) were

selected to calculate male dominance hierarchy (table s2A and s2B). We used the 'steepness'

package (*80*) to build dominance hierarchy and David's score. We also checked h☐ index (*81*)

– a modified version of Landau's linearity index (*82*) – using the 'igraph' package (*83*) to

decide whether the male dominance hierarchy is linear or not.

Statistical analysis

We used R 4.0.2 (*84*) with R studio 1.3.1056 (*85*) for all analyses. We used generalized

linear mixed models (GLMM) or generalized linear models (GLM) with negative binomial,

binomial error distributions along with several packages including 'lme4', and 'lmertest' to fit

our models to the data. Mixed models that included random effects were done to control the

repeated measure of the behavior from the same individual. More details of each model will be

described later in this section. We also checked multicollinearity between independent

variables in the model based on variance inflation factors (VIFs) after we ran all mixed

models. In all models, the maximum VIF was less than 3 (only over 2 in GLMM-1C),

therefore the multi-collinearity among the predictor variables is of serious concern (*86*). All

models were also significantly better in explaining the data compared with null models.

In bonobos, it was reported that a male's copulation rate is related to other social factors,

such as the mother's support and relative dominance rank caused by changes in party

composition in a group (*16*). Therefore, we may need to consider those factors as predictor

variables in the models. However, the 4 subject males whose mothers were in the group, were almost always with their mother in the party. There were only 23 out of 744 days (sum of days when the 4 males were observed) that any of the 4 males were found without their mother (only 3.1%). Therefore, we did not include this factor in the models. Fluctuation of the relative rank between days because of change in party composition was not the main predictor, since our question is more generally whether the male rank within the community is an important factor for understanding male mating efforts. In addition, the alpha male was the highest-ranking male on 188 days of the 253 observation days on which we observed males (74.3%). The beta male occupied the highest position on 33 days (13.0%), the 3rd ranking male for 9 days (3.6%) and the 4th for 17 days (6.7%).

We also considered that the reproductive quality of female bonobos has two main components. One is the increased chance of getting pregnant with the growth of infants over time, regardless of the menstrual cycle on a wide time scale. The other is changes in the probability of fertility (the chance of fertilization – a sperm meets an egg) within a menstrual cycle depending on changes in the probability of ovulation over time. In chimpanzees, these two components are closely related to each other probably because energetic constraint from lactation has a regulative role in the menstrual cycle (*10*). This is probably similar to female bonobos, although bonobo female resumes their swelling cycle much earlier (*6*, *20*). To integrate such an effect of one's infant on the female reproductive quality, we integrated infant age, as a proxy for female body condition. To examine the effect of ovulation, we also included the number of days from ovulation or the detumescence or onset of maximal swelling as a predictor variable in each model depending on our question because the chance of fertilization increases until the day of ovulation. The day of ovulation (or detumescence) was coded 0 and one day before the ovulation was -1, two days before was -2, and so on. The days after the ovulation were 1, 2, and so on. The onset of maximal swelling was coded 1, the next day was 2, and so on. In some analyses, we confined the data from -14 days before ovulation to the ovulation day (15 days, -14 to 0) to control large inter-and intra-individual variations in the length of the maximal swelling phase.

Infant age was coded on an ordinal scale by putting together 6 months as 0.5 years. It ranges from -1 to 5.5. From the day of birth to 6 months, the infant age was coded 0.5. From six months before parturition to the day of parturition were therefore coded -0.5. We also coded -1 from the detumescent day of maximal swelling which coincided with ovulation

which resulted in successful deliveries of an infant. Finally, to avoid the high VIF and

convergence errors of the model, we standardized all predictor variables (male rank, days from

ovulation/detumescence/onset, infant age, the number of males and females) to make all

variables has a mean of 0 and standard deviation of 1. To make figures more intuitive and

easier to interpret, they were made with the original scale.

-    **Male intensive following after a female with different reproductive quality**

To investigate factors that influence male intensive following after a certain female, we

ran two GLMMs with negative binomial distribution following the hurdle model approaches

(*40*). In the first model (GLMM-1A), we tested whether maximal swelling and increase in

infant age (explanatory variables) predict the number of males (response variable) who

intensively followed a certain female using all data (250 days). Female age was also included

as a fixed effect to examine whether female age influencing male intensive following. The

number of males and females with maximal swelling were also included as fixed effects to

examine whether the spatio-temporal distribution of males and females with maximal swelling

on the day influences the number of males who intensively follow the female. Female ID was

the random factor in the model. One row of the data set comprised swelling status on a given

day (ordinal 1 to 3), and infant age of a certain female as well as her age. In the same row, the

number of males and females with maximal swelling of the given day was also placed. There

were on average 6.4 ± 2.4 females in a day in the data set.

In the second part of the hurdle model, we took a subset of the data, which only

consisted of the females with maximal swelling (205 days), to examine the effect of

detumescence of maximal swelling and infant age (GLMM-1B). The average number of

females with maximal swelling per day was 2.7 ± 1.3 (range: 1 to 8 per day). In this model,

the days from detumescence of maximal swelling were included as an explanatory variable

instead of swelling scores. The other variables were the same with GLMM-1A, except the

inclusion of the interaction term between detumescent day and infant age, and a categorical

variable that divides maximal swelling phase into ovulation detected or not – 209 ovulation-

detected female-days and 336 ovulation-undetected female-days. Here a female-day represents

a row in the data set that includes other explanatory and response variables of a given female

on the given day. The inclusion of the interaction term was to test whether there would be a

dependency between the two terms on male's intensive following. The categorical variable for

the maximal swelling phase (ovulatory vs anovulatory) was included to test whether males can

discriminate ovulatory cycles between maximal swelling phases. Our division of maximal

swelling into ovulatory or anovulatory is a rough estimation as we could categorize an

ovulatory cycle into anovulatory but not likely vice versa. However, it would provide

therefore a conservative estimation of the effect of ovulation on male behavior.

We also investigated how males chose the target of the intensive following using a

binomial mixed model (GLMM-1C) with the data when there were more than 2 females with

maximal swelling in a daily ranging party (77 observation days). In this model, the number of

days from the onset of maximal swelling and infant age as well as their interaction were

included as explanatory variables. Female age and the number of males of the day were also

included as in the two previous models. There were on average 3.2 ± 1.3 females with

maximal swelling in the data set. Using ovulation-detected menstrual cycle from 14 females

which cover 138 observation days, we finally tested the effect of ovulation on the intensive

following behavior (GLMM-1D). In this model, other explanatory variables were identical

with GLMM-1B except that we changed days from detumescence to days from ovulation

(ovulation day is coded 0). As in other models, it was fitted to a negative binomial distribution.

-   **Copulation, intervention, solicitation of copulation, and male-male aggression**

Male behaviors including copulation and intervention of copulation, and solicitation of

copulation are probably directly related to male mating efforts. To examine the number of

these three events between male-female dyads on a given day, three separated GLMMs with

negative binomial distribution were conducted. For these three analyses, we used the 14

ovulation-detected maximal swelling phases from 5 females and limited the data from -14 to 0

days from ovulation (104 days in total). Within this ovulatory maximal swelling phase, there

were temporal drops from maximal swelling to intermediate swelling (swelling score 2). The

maximal swelling temporally dropped to the intermediate swelling on 11 days from 5 females

(fig. S1). These 11 days were excluded from data analysis. However, within the menstrual

cycle of a female called Sl, 3 days of the fertile phase were outside of the maximal swelling

phase. These 3 days of the intermediate swelling phase remained in the data set as a fertile

window is situated in the days (*24*).

Each model (copulation: GLMM-2A and 2B, intervention: GLMM-2C, solicitation:

GLMM-2D) has the number of copulations, interventions, and solicitations of copulation

observed within a male-female dyad on a given day as a response variable. The number of

events was controlled by the hours that the male and female of the given dyad were observed

together (total counts of a one-hour party on a given day) using the offset function. The

explanatory variables were the number of days from ovulation, the infant age of the female of

the dyad, male rank, male age, and female age. Males with their mothers have better

copulation opportunities and advantages over the male-male competition (*4*, *16*). To include

these social dynamics to some extent male age was included in all three models as the

probability of having mothers in the group will be higher for younger males. In the GLMM-

2B, male rank was converted into two categories, high (1st to 3rd ranking) and low (4th to

10th ranking) to further investigate the effect of 3-way interaction between days from

ovulation, infant age, and the male rank on the number of copulations. The female-male dyad

that was nested within a certain menstrual cycle was integrated into all models as a random

variable (coded as dyad ID: cycle name). We did not use the 'dyad ID*cycle name' term

because of a convergent error.

To test whether the number of females with maximal swelling influences the number of

male-male aggression, we ran a GLM with a negative binomial distribution (GLM-2E). As

male-male aggression might be also related both directly and indirectly with the male mating

competition, we included the existence of male intensive following (O or X) as a fixed effect

along with the number of females with maximal swelling on a given day. Both following

hours of the day and the number of males of the given day were included as an offset term to

control the effect of observation hours and the number of males in a daily ranging party. The

random effect could not be defined as the number of male-male aggression was pooled within

a day. One row of the data included the total number of aggressions of the day, the existence

of intensive following and the number of females with maximal swelling, and so on.

1031
1032
1033
1034 **References**
1035

1. M. Andersson, L. W. Simmons, Sexual selection and mate choice. *Trends in Ecology &*
Evolution. **21**, 296–302 (2006).

2. C. W. Miller, in *The Princeton Guide to Evolution*, J. B. Losos, Ed. (Princeton University
Press, Princeton, 2013; https://doi.org/10.1515/9781400848065-089), pp. 267–299.

3. F. B. M. de Waal, *Peacemaking among primates* (Harvard University Press, Cambridge
Mass, 1989).

4. T. Furuichi, Female contributions to the peaceful nature of bonobo society. *Evolutionary*
Anthropology: Issues, News, and Reviews. **20**, 131–142 (2011).

5. Z. Clay, T. Furuichi, F. B. M. de Waal, Obstacles and catalysts to peaceful coexistence in
chimpanzees and bonobos. *Behaviour*. **153**, 1293–1330 (2016).

6. T. Furuichi, C. Hashimoto, in *Behavioural Diversity in Chimpanzees and Bonobos*, C.
Boesch, G. Hohmann, L. F. Marchant, Eds. (Cambridge University Press, Cambridge, UK,
2002), pp. 156–167.

7. T. Kano, *The last ape: pygmy chimpanzee behavior and ecology* (Stanford University Press,
Stanford, 1992).

8. T. Furuichi, Sexual swelling, receptivity, and grouping of wild pygmy chimpanzee females
at Wamba, Zaire. *Primates*. **28**, 309–318 (1987).

9. F. B. M. de Waal, Bonobo sex and society. *Scientific American*. **272**, 82–88 (1995).

10. M. E. Thompson, M. N. Muller, R. W. Wrangham, The energetics of lactation and the
return to fecundity in wild chimpanzees. *Behavioral Ecology*. **23**, 1234–1241 (2012).

11. K. E. Reichert, M. Heistermann, J. K. Hodges, C. Boesch, G. Hohmann, What females tell
males about their reproductive status: Are morphological and behavioural cues reliable
signals of ovulation in bonobos (*Pan paniscus*)? *Ethology*. **108**, 583–600 (2002).

12. P. H. Douglas, G. Hohmann, R. Murtagh, R. Thiessen-Bock, T. Deschner, Mixed messages:
wild female bonobos show high variability in the timing of ovulation in relation to sexual
swelling patterns. *BMC Evolutionary Biology*. **16:140** (2016), doi:10.1186/s12862-016-
0691-3.

13. S. E. Street, C. P. Cross, G. R. Brown, Exaggerated sexual swellings in female nonhuman
primates are reliable signals of female fertility and body condition. *Animal Behaviour*. **112**,
203–212 (2016).

14. M. Surbeck, K. E. Langergraber, B. Fruth, L. Vigilant, G. Hohmann, Male reproductive
skew is higher in bonobos than chimpanzees. *Current Biology*. **27**, R640–R641 (2017).

15. S. Ishizuka, Y. Kawamoto, T. Sakamaki, N. Tokuyama, K. Toda, H. Okamura, T. Furuichi,
Paternity and kin structure among neighbouring groups in wild bonobos at Wamba. *Royal*
Society Open Science. **5**, 171006 (2018).

16. M. Surbeck, R. Mundry, G. Hohmann, Mothers matter! Maternal support, dominance status
and mating success in male bonobos (*Pan paniscus*). *Proceedings of the Royal Society B:*
Biological Sciences. **278**, 590–598 (2011).

17. M. Heistermann, U. Möhle, H. Vervaecke, L. Van Elsacker, J. K. Hodges, Application of
urinary and fecal steroid measurements for monitoring ovarian function and pregnancy in
the bonobo (*Pan paniscus*) and evaluation of perineal swelling patterns in relation to
endocrine events. *Biology of Reproduction*. **55**, 844–853 (1996).

1078 18. M. H. Jurke, L. R. Hagey, S. Jurke, N. M. Czekala, Monitoring hormones in urine and feces
1079    of captive bonobos (*Pan paniscus*). *Primates*. **41**, 311–319 (2000).

1080 19. H. Ryu, thesis, Kyoto University, Japan, Kyoto (2017).

1081 20. C. Hashimoto, H. Ryu, K. Mouri, K. Shimizu, T. Sakamaki, T. Furuichi, Physical,
1082    behavioral, and hormonal changes in the resumption of sexual receptivity during postpartum
1083    infertility in female bonobos at Wamba. *Primates* (2022), doi:10.1007/s10329-021-00968-
1084    w.

1085 21. A. J. Wilcox, D. Dunson, D. D. Baird, The timing of the "fertile window" in the menstrual
1086    cycle: day specific estimates from a prospective study. *BMJ□: British Medical Journal*.
1087    **321**, 1259–1262 (2000).

1088 22. J. Goodall, *The chimpanzees of Gombe: patterns of behavior* (The Belknap Press of Harvard
1089    University Press, Cambridge, Mass, 1986).

1090 23. T. Deschner, M. Heistermann, K. Hodges, C. Boesch, Timing and probability of ovulation
1091    in relation to sex skin swelling in wild West African chimpanzees, *Pan troglodytes verus*.
1092    *Animal Behaviour*. **66**, 551–560 (2003).

1093 24. T. Deschner, M. Heistermann, K. Hodges, C. Boesch, Female sexual swelling size, timing
1094    of ovulation, and male behavior in wild West African chimpanzees. *Hormones and
1095    Behavior*. **46**, 204–215 (2004).

1096 25. M. E. Thompson, R. W. Wrangham, Male mating interest varies with female fecundity in
1097    *Pan troglodytes schweinfurthii* of Kanyawara, Kibale national park. *International Journal of
1098    Primatology*. **29**, 885–905 (2008).

1099 26. C. L. Nunn, The evolution of exaggerated sexual swellings in primates and the graded-
1100    signal hypothesis. *Animal Behaviour*. **58**, 229–246 (1999).

1101 27. C. L. Fitzpatrick, J. Altmann, S. C. Alberts, Exaggerated sexual swellings and male mate
1102    choice in primates: testing the reliable indicator hypothesis in the Amboseli baboons.
1103    *Animal Behaviour*. **104**, 175–185 (2015).

1104 28. C. E. G. Tutin, Mating patterns and reproductive strategies in a community of wild
1105    chimpanzees (*Pan troglodytes schweinfurthii*). *Behavioral Ecology and Sociobiology*. **6**,
1106    29–38 (1979).

1107 29. B. B. Smuts, R. W. Smuts, in *Advances in the Study of Behavior*, J. S. R. Peter J.B. Slater,
1108    Ed. (Academic Press, 1993;
1109    http://www.sciencedirect.com/science/article/pii/S0065345408604040), vol. Volume 22, pp.
1110    1–63.

1111 30. M. N. Muller, S. M. Kahlenberg, M. E. Thompson, R. W. Wrangham, Male coercion and
1112    the costs of promiscuous mating for female chimpanzees. *Proceedings of the Royal Society
1113    B: Biological Sciences*. **274**, 1009–1014 (2007).

31. M. N. Muller, S. M. Kahlenberg, R. W. Wrangham, in *Sexual coercion in primates and*
humans□: an evolutionary perspective on male aggression against females, M. N. Muller,
R. W. Wrangham, Eds. (Harvard University Press, Cambridge Mass, 2009), pp. 184–217.

32. D. P. Watts, Coalitionary mate guarding by male chimpanzees at Ngogo, Kibale National
Park, Uganda. *Behavioral Ecology and Sociobiology*. **44**, 43–55 (1998).

33. J. Wallis, Synchrony of estrous swelling in captive group-living chimpanzees (*Pan*
troglodytes). *International Journal of Primatology*. **6**, 335–350 (1985).

34. M. E. Pereira, Asynchrony within estrous synchrony among ringtailed lemurs (Primates:
Lemuridae). *Physiology & Behavior*. **49**, 47–52 (1991).

35. T. Paoli, in *Sexual coercion in primates and humans□: an evolutionary perspective on male*
aggression against females, M. N. Muller, R. W. Wrangham, Eds. (Harvard University
Press, Cambridge Mass, 2009), pp. 410–423.

36. A. R. Parish, Sex and food control in the "uncommon chimpanzee": How bonobo females
overcome a phylogenetic legacy of male dominance. *Ethology and Sociobiology*. **15**, 157–
179 (1994).

37. J. Gillman, The effect of multiple injections of progesterone on the turgescent perineum of
the baboon (*Papio porcarius*). *Endocrinology*. **26**, 1072–1077 (1940).

38. J. Kato, T. Onouchi, K. Oshima, The presence of progesterone receptors in the sexual skin
of the monkey. *Steroids*. **36**, 743–749 (1980).

39. P. Martin, P. Bateson, *Measuring behaviour: an introductory guide* (Cambridge University
Press, Cambridge, ed. 2nd, 1993).

40. A. F. Zuur, E. N. Ieno, N. J. Walker, A. A. Saveliev, G. M. Smith, *Mixed effects models and*
extensions in ecology with R (Springer, New York, 2009;
http://www.springer.com/life+sciences/ecology/book/978-0-387-87457-9).

41. M. N. Muller, M. E. Thompson, R. W. Wrangham, Male chimpanzees prefer mating with
old females. *Current Biology*. **16**, 2234–2238 (2006).

42. F. B. Bercovitch, Coalitions, cooperation and reproductive tactics among adult male
baboons. *Animal behaviour*. **36**, 1198–1209 (1988).

43. L. R. Gesquiere, E. O. Wango, S. C. Alberts, J. Altmann, Mechanisms of sexual selection:
sexual swellings and estrogen concentrations as fertility indicators and cues for male consort
decisions in wild baboons. *Hormones and Behavior*. **51**, 114–125 (2007).

44. L. Rigaill, J. P. Higham, P. C. Lee, A. Blin, C. Garcia, Multimodal sexual signaling and
mating behavior in olive baboons (*Papio anubis*). *American Journal of Primatology*. **75**,
774–787 (2013).

45.   C. Young, B. Majolo, M. Heistermann, O. Schülke, J. Ostner, Male mating behaviour in
relation to female sexual swellings, socio-sexual behaviour and hormonal changes in wild
Barbary macaques. *Hormones and Behavior*. **63**, 32–39 (2013).

46.   M. Surbeck, T. Deschner, A. Weltring, G. Hohmann, Social correlates of variation in
urinary cortisol in wild male bonobos (*Pan paniscus*). *Hormones and Behavior*. **62**, 27–35
(2012).

47.   M. Surbeck, T. Deschner, G. Schubert, A. Weltring, G. Hohmann, Mate competition,
testosterone and intersexual relationships in bonobos, *Pan paniscus*. *Animal Behaviour*. **83**,
659–669 (2012).

48.   T. Furuichi, Agonistic interactions and matrifocal dominance rank of wild bonobos (*Pan
paniscus*) at Wamba. *International Journal of Primatology*. **18**, 855–875 (1997).

49.   N. Tokuyama, T. Furuichi, Do friends help each other? Patterns of female coalition
formation in wild bonobos at Wamba. *Animal Behaviour*. **119**, 27–35 (2016).

50.   M. Surbeck, C. Girard-Buttoz, L. Samuni, C. Boesch, B. Fruth, C. Crockford, R. M. Wittig,
G. Hohmann, Attractiveness of female sexual signaling predicts differences in female
grouping patterns between bonobos and chimpanzees. *Commun Biol*. **4**, 1–11 (2021).

51.   H. Ryu, D. A. Hill, T. Furuichi, Prolonged maximal sexual swelling in wild bonobos
facilitates affiliative interactions between females. *Behaviour*. **152**, 285–311 (2015).

52.   G. Hohmann, U. Gerloff, D. Tautz, B. Fruth, Social bonds and genetic ties: kinship,
association and affiliation in a community of bonobos (*Pan paniscus*). *Behaviour*. **136**,
1219–1235 (1999).

53.   C. Boesch, G. Kohou, H. Néné, L. Vigilant, Male competition and paternity in wild
chimpanzees of the Taï forest. *American Journal of Physical Anthropology*. **130**, 103–115
(2006).

54.   A. F. Dixson, *Primate sexuality: comparative studies of the prosimians, monkeys, apes and
human beings* (Oxford University Press, New York, 2012).

55.   U. Gerloff, B. Hartung, B. Fruth, G. Hohmann, D. Tautz, Intracommunity relationships,
dispersal pattern and paternity success in a wild living community of bonobos (*Pan
paniscus*) determined from DNA analysis of faecal samples. *Proceedings of the Royal
Society of London. Series B: Biological Sciences*. **266**, 1189–1195 (1999).

56.   A. Bateman, Intra-sexual selection in Drosophila. *Heredity*. **2**, 349–368 (1948).

57.   M. A. Emery, P. L. Whitten, Size of sexual swellings reflects ovarian function in
chimpanzees (*Pan troglodytes*). *Behavioral Ecology and Sociobiology*. **54**, 340–351 (2003).

58.   J. P. Higham, A. M. MacLarnon, C. Ross, M. Heistermann, S. Semple, Baboon sexual
swellings: Information content of size and color. *Hormones and Behavior*. **53**, 452–462
(2008).

59. D. W. Bullock, C. A. Paris, R. W. Goy, Sexual behaviour, swelling of the sex skin and
plasma progesterone in the pigtail macaque. *Reproduction*. **31**, 225–236 (1972).

60. K. S. Carlisle, R. M. Brenner, W. Montagna, Hormonal regulation of sex skin in *Macaca*
nemestrina. *Biology of Reproduction*. **25**, 1053–1063 (1981).

61. H. Ozasa, K. G. Gould, Demonstration and characterization of estrogen receptor in
chimpanzee sex skin: correlation between nuclear receptor levels and degree of swelling.
Endocrinology. **111**, 125–131 (1982).

62. A. J. Wilcox, C. R. Weinberg, J. F. O'Connor, D. D. Baird, J. P. Schlatterer, R. E. Canfield,
E. G. Armstrong, B. C. Nisula, Incidence of early loss of pregnancy. *New England Journal*
of Medicine. **319**, 189–194 (1988).

63. A. J. Wilcox, D. D. Baird, C. R. Weinberg, Time of implantation of the conceptus and loss
of pregnancy. *New England Journal of Medicine*. **340**, 1796–1799 (1999).

64. A. R. Parish, F. B. M. de Waal, D. Haig, The other "closest living relative": How bonobos
(*Pan paniscus*) challenge traditional assumptions about females, dominance, intra- and
intersexual interactions, and hominid evolution. *Annals of the New York Academy of*
Sciences. **907**, 97–113 (2000).

65. N. Burley, The evolution of concealed ovulation. *The American naturalist*, 835–858 (1979).

66. S. Kahlenberg, M. E. Thompson, R. Wrangham, Female competition over core areas in *Pan*
troglodytes schweinfurthii, Kibale National Park, Uganda. *International Journal of*
Primatology. **29**, 931–947 (2008).

67. T. Furuichi, G. Idani, H. Ihobe, C. Hashimoto, Y. Tashiro, T. Sakamaki, M. N. Mulavwa, K.
Yangozene, S. Kuroda, in *Long-term field studies of primates*, P. M. Kappeler, D. P. Watts,
Eds. (Springer, Berlin, 2012), pp. 413–433.

68. C. Hashimoto, Y. Tashiro, D. Kimura, T. Enomoto, E. J. Ingmanson, G. Idani, T. Furuichi,
Habitat use and ranging of wild bonobos (*Pan paniscus*) at Wamba. *International Journal of*
Primatology. **19**, 1045–1060 (1998).

69. M. N. Mulavwa, T. Furuichi, K. Yangozene, M. Yamba-Yamba, B. Motema-Salo, G. Idani,
H. Ihobe, C. Hashimoto, Y. Tashiro, N. Mwanza, in *The Bonobos*, T. Furuichi, J.
Thompson, Eds. (Springer New York, 2008; http://dx.doi.org/10.1007/978-0-387-74787-
3_7), *Developments in Primatology: Progress and Prospects*, pp. 121–134.

70. C. Hashimoto, Context and development of sexual behavior of wild bonobos (*Pan paniscus*)
at Wamba, Zaire. *International journal of primatology*. **18**, 1–21 (1997).

71. A. E. Pusey, Behavioural changes at adolescence in chimpanzees. *Behaviour*. **115**, 203–246
(1990).

72. T. Sakamaki, I. Behncke, M. Laporte, M. Mulavwa, H. Ryu, H. Takemoto, N. Tokuyama, S.
Yamamoto, T. Furuich, in *Dispersing primate females: life history and social strategies in*
male-philopatric species, T. Furuich, J. Yamagiwa, F. Aureli, Eds. (Springer Japan, Tokyo,

Japan, 2015; http://www.springer.com/in/book/9784431554790), *Primatology Monographs*,
pp. 127–164.

73. C. Hashimoto, T. Furuichi, Y. Tashiro, What factors affect the size of chimpanzee parties in
the Kalinzu Forest, Uganda? Examination of fruit abundance and number of estrous
females. *International Journal of Primatology*. **22**, 947–959 (2001).

74. S. Kuroda, Grouping of the pygmy chimpanzees. *Primates*. **20**, 161–183 (1979).

75. T. Kano, The social group of pygmy chimpanzees (*Pan paniscus*) of Wamba. *Primates*. **23**,
171–188 (1982).

76. K. E. Graham, T. Furuichi, R. W. Byrne, The gestural repertoire of the wild bonobo (*Pan
paniscus*): a mutually understood communication system. *Animal Cognition*, 1–7 (2016).

77. H. Vervaecke, H. de Vries, L. van Elsacker, Dominance and its behavioral measures in a
captive group of bonobos (*Pan paniscus*). *International Journal of Primatology*. **21**, 47–68
(2000).

78. A. J. Wilcox, C. R. Weinberg, D. D. Baird, Timing of sexual intercourse in relation to
ovulation - effects on the probability of conception, survival of the pregnancy, and sex of
the baby. *New England Journal of Medicine*. **333**, 1517–1521 (1995).

79. D. B. Dunson, D. D. Baird, A. J. Wilcox, C. R. Weinberg, Day-specific probabilities of
clinical pregnancy based on two studies with imperfect measures of ovulation. *Human
Reproduction*. **14**, 1835–1839 (1999).

80. D. Leiva, H. De Vries, *Steepness: testing steepness of dominance hierarchies* (2014;
https://CRAN.R-project.org/package=steepness).

81. H. de Vries, An improved test of linearity in dominance hierarchies containing unknown or
tied relationships. *Animal behaviour*. **50**, 1375–1389 (1995).

82. H. G. Landau, On dominance relations and the structure of animal societies: I. Effect of
inherent characteristics. *The Bulletin of Mathematical Biophysics*. **13**, 1–19 (1951).

83. G. Csardi, T. Nepusz, The igraph software package for complex network research.
InterJournal. **Complex Systems**, 1695 (2006).

84. R Core Team, *R: A language and environment for statistical computing* (R Foundation for
Statistical Computing, Vienna, Austria, 2020).

85. RStudio Team, *RStudio: integrated development environment for R* (RStudio, PBC, Boston,
MA, 2020; http://www.rstudio.com/).

86. A. F. Zuur, E. N. Ieno, C. S. Elphick, A protocol for data exploration to avoid common
statistical problems. *Methods in Ecology and Evolution*. **1**, 3–14 (2010).

1254

Acknowledgments

We thank our research assistants at Wamba for their great help and members of the Department of Ecology and Social Behavior and CICASP of the Primate Research Institute of Kyoto University for their support for fieldwork and valuable comments. We thank in particular K. Hosaka, T. Miyabe-Nishiwaki, F. Bercovitch, M. Nishikawa, A. MacIntosh, K. Graham, Y. Kawaguchi for valuable comments and reviews on the manuscript. We finally thank the Ministry of Scientific and Technological Research of the D.R. Congo for research permissions (MIN.ESURS/SG-RST/13/2013, MIN.ESURS/SG-RST/007/2014, MIN.ESURS/SG-RST/026/2014 to HR) and the Research Center of Ecology and Forestry for helping us and maintaining Luo Scientific Reserve.

Funding:
Environment Research and Technology Development Fund D-1007 (TF)
Japan Society for the Promotion of Science (JSPS) Grants-in-aid for Scientific Research 22255007 (TF)
JSPS Grants-in-aid for Scientific Research 25304019 (CH)
JSPS Grants-in-aid for Scientific Research 25257409 (HI),
JSPS Asia-Africa Science Platform Program 2012-2014 (TF)

Author contributions:
Conceptualization: HR
Methodology: HR, CH, KM, KS
Investigation: HR
Visualization: HR
Supervision: TF, DAH, CH
Project Administration: TF
Writing—original draft: HR
Writing—review & editing: HR, CH, DAH, KM, KS, TF

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text and supplementary materials. Raw and other related data may be requested from the authors.

Supplementary Materials

Supplementary materials for this article are available in a separate file.

 Figures and Tables

1293

1294 **Fig. 1. Morphological changes in sexual swelling of a female named Fk.** (A) non swelling;
1295   swelling score 1, (B) intermediate swelling; swelling score 2, (C) maximal swelling;
1296   swelling score 3.

1297

Fig. 2. The number of males in intensive following across all 250 observation days. (A)
the number of males in intensive following after females increased with the size of
sexual swelling. (B) the number of males in intensive following after females increased
with infant age of females. Sexual swelling 1 is non-swelling, 2 is intermediate
swelling, 3 is the maximal swelling. Infant age ranges -1 to 5 and the minus value
represents pregnant periods which included post conception swelling which resulted in
a successful delivery. Infant age 0 is from birth to within 1year, and infant age 1 is
from 1 to <2 years and so on. Due to many zero, medians are 0, and only a small box
that shows 75th percentile and an upper whisker is seen in figure 2B. The points are the
number of males in the intensive following of the day divided by daily observation
hours.

Fig. 3. Male intensive following after females with maximal swelling. (A) More males
intensively followed a female with an older infant and detumescence of the maximal
swelling of the female was close. The fitted line (red) was generated by the 'lowess'
function with default value in r which is based on locally weighted least squares
regression. (B) The probability of males' intensive following increased when a female
with an older infant and the onset of maximal swelling of the female started earlier.
The fitted line (red) was generated by logistic regression and each. Each dot is a single
data point for each day. Day 0 is the day that the maximal swelling of a given female
became detumescent (A) or one day before the onset of maximal swelling (B).

1318

Fig. 4. Male intensive following concerning the day of ovulation (14 cycles in 5 females).
The number of males each day who intensively followed females in relation to the days
from ovulation increased when ovulation was approaching. Detumescence of maximal
swelling started after 1.9 ± 2.4 days from ovulation day on average. The median
number of males in the intensive following after a given female was highest on the day
of ovulation. Due to many zero numbers of males, the median is mostly zero except the
periovulatory phase and one day after ovulation. Day 0 is the day that ovulation
occurred. The upper and lower edges of the box represent 75th and 25th percentile
respectively. A line within a box is median and whiskers are 1.5 times of interquartile
range, and the dots are all data points.

1329

Fig. 5. Probability of ovulation and male behaviors. (A) The probability of ovulation and
fertility days from the onset of maximal swelling phase, and (B) from the
detumescence of the maximal swelling phase. Black closed circles and lines are the
day-specific ovulation probability, and red open circles and lines are the day-specific
probability of fertility. (C) The number of males who intensively followed females
with the onset of maximal swelling is widely distributed as found in the probability of
ovulation in A. (D) The number of males who intensively followed females concerning
days from the detumescence of maximal swelling shows a similar distribution as found
in the probability of ovulation in B. Day 1 is the day of onset of maximal swelling
phase (A, C). Day 0 is the day of detumescence (B, D). The upper and lower edges of
the box represent 75th and 25th percentile respectively. A line within a box is median
and whiskers are 1.5 times of interquartile range, and the dots are all data points.

1342

Fig. 6. Prediction plots of males' copulation, intervention, solicitation of copulation. (A to
E) The predicted copulation rate within a dyad (GLMM-2A) in relation to the
ovulation day, infant age, male rank, male age, and female age. (F to J) The predicted
rate of intervention of copulation by the male within a female-male dyad in relation to
the ovulation day, infant age, male rank, male age, and female age. (K to O) The
predicted rate of solicitation of copulation by a dyadic male in relation to ovulation
day, infant age, male rank, male age, and female age. Asterisks (*) on each plot
indicate *P*-value (ns: not significant, *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$).

1351

Fig. 7. The number of male-male aggression concerning females with maximal swelling.
(A) When males' intensive following behavior was observed, there were more
aggressions per hour between males. (B) The number of male-male aggression also
increased when there were more females with maximal swelling in a daily ranging
party. Each dot represents each data point (total 250). The upper and lower edges of the
box represent 75th and 25th percentile respectively. A line within a box is median and
whiskers are 1.5 times of interquartile range.

1359

1360

 Table 1. Summary statistics of GLMM-1A to 1D.

GLMM results of the number of males who intensively followed a given female in
relation to infant age, sexual swelling, days from detumescence/onset of maximal
swelling, ovulation, female age, and the number of males and females with maximal
swelling. All predictor variables were standardized (mean of 0 and a standard deviation
of 1) and the maximal variance influence factor, VIF was 2.95 for days from onset of
MS in GLMM-1C. Asterisks indicates significance, *: $P<0.05$, **: $P<0.01$, ***:
$P<0.001$. FMS: female with maximal swelling, MS: maximal swelling.

Models	Predictors	Estimates	SE	Z	P
GLMM-1A (Figure 2)	Intercept	-7.98	0.85	-9.43	<0.001***
	Infant age	2.76	0.52	5.33	<0.001***
	Swelling score	2.48	0.33	7.44	<0.001***
	Female age	-0.02	0.59	-0.04	0.971
	Number of males	-0.17	0.17	-1.01	0.312
	Number of FMS	-0.03	0.15	-0.18	0.854
GLMM-1B (Figure 3A)	Intercept	-3.00	0.21	-14.42	<0.001***
	Infant age[1]	0.39	0.18	2.22	0.026*
	Days from detumescence[2]	1.25	0.21	6.06	<0.001***
	Female age	0.10	0.12	0.85	0.393
	Number of males	-0.10	0.12	-0.81	0.419
	Number of FMS	-0.07	0.12	-0.55	0.582
	Ovulation undetected vs detected	-1.13	0.28	-4.03	<0.001***
	Interaction between 1 & 2	1.36	0.25	5.44	<0.001***
GLMM-1C (Figure 3B)	Intercept	-3.00	0.38	-7.89	<0.001***
	Infant age[3]	3.44	0.55	6.21	<0.001***
	Days from onset of MS[4]	2.24	0.39	5.82	<0.001***
	Female age	0.84	0.38	2.20	0.028*
	Number of males	-0.02	0.22	-0.08	0.939
	Interaction between 3 & 4	1.92	0.47	4.12	<0.001***
GLMM-1D (Figure 4)	Intercept	-2.97	0.38	-7.87	<0.001***
	Infant age[5]	1.59	0.42	3.83	<0.001***
	Days from ovulation[6]	1.74	0.23	7.58	<0.001***
	Female age	-0.26	0.31	-0.82	0.413
	Number of males	0.17	0.14	1.23	0.217
	Number of FMS	0.10	0.13	0.78	0.434
	Interaction between 5 & 6	-0.04	0.27	-0.14	0.887

1369
1370

 Table 2. Summary statistics of GLMM-2A to GLM-2E

GLMM (2A to 2D) results of the number of copulations, interventions, and
solicitations of copulation between male-female dyads in relation to days from
ovulation (-14 to 0), infant age, female age, male rank (1: highest to 10: lowest in
GLMM-2A, 1 to 3: high and 4 to 10: low in GLMM-2B), and male age in 104 days.
GLM-2E shows results of the number of male-male aggression in relation to the
existence of male intensive following and the number of females with maximal
swelling. All predictor variables were standardized and the maximal VIF was 1.26 for
female age in GLMM-2C. Asterisks indicates significance, *: $P<0.05$, **: $P<0.01$, ***:
$P<0.001$. FMS: female with maximal swelling.

Models	Predictors	Estimates	SE	Z	P
GLMM-2A Copulation	Intercept	-4.89	0.27	-18.19	<0.001***
	Days from ovulation	0.24	0.10	2.49	0.013*
	Infant age	-0.48	0.18	-2.65	0.008**
	Male rank (ordinal 1 to 10)	-0.44	0.17	-2.65	0.008**
	Male age	-1.28	0.22	-5.79	<0.001***
	Female age	0.02	0.19	0.11	0.911
GLMM-2B Copulation	Intercept	-4.07	0.30	-13.61	<0.001***
	Days from ovulation[1]	0.30	0.10	3.01	0.003**
	Infant age[2]	-0.50	0.17	-2.86	0.004**
	Male rank low (base) vs high	-1.35	0.34	-3.99	<0.001***
	Male age	-1.23	0.22	-5.62	<0.001***
	Interaction 1, 2 & male rank (high)	-0.42	0.14	-3.01	0.003**
	Interaction 1, 2 & male rank (low)	0.24	0.14	1.70	0.090
GLMM-2C. Intervention	Intercept	-7.94	0.82	-9.70	<0.001***
	Days from ovulation	0.85	0.25	3.43	<0.001***
	Infant age	0.81	0.36	2.26	0.024*
	Male rank	-2.14	0.57	-3.77	<0.001***
	Male age	-0.58	0.41	-1.42	0.157
	Female age	-0.47	0.25	-1.93	0.053
GLMM-2D. Solicitation	Intercept	-4.53	0.23	-19.59	<0.001***
	Days from ovulation	0.29	0.10	2.84	0.005**
	Infant age	-0.10	0.15	-0.68	0.499
	Male rank	-0.34	0.14	-2.45	0.014*
	Male age	-1.25	0.19	-6.43	<0.001***
	Female age	0.15	0.15	1.00	0.318
GLM-2E. Male-male aggression	Intercept	-2.32	0.08	-27.41	<0.001***
	Intensive following (x: o)	-0.58	0.11	-5.07	<0.001***
	Number of FMS	0.20	0.06	3.42	<0.001***

1381