Modulation of glucose metabolism by 2-Deoxy-D-Glucose (2DG) promotes

IL-17 producing human T cell subsets

Xin Chen[1,2,3], Lina Kozhaya[1], Chuxi Wang[1,2], Lindsey Placek[1], Ece Karhan[1], Derya Unutmaz[1,2]

[1] Jackson Laboratory for Genomic Medicine, Farmington, Connecticut, USA.

[2] Department of Immunology, University of Connecticut School of Medicine, Farmington, CT USA.

[3] Institute for Immunity, Transplantation, and Infection, Stanford University, Stanford, CA USA.

* Corresponding author E-mail: derya@mac.com

Abstract

Activation and differentiation of T cells are closely linked to their cellular metabolic programs.

Glycolysis and mitochondrial metabolism are thought to be critical in modulating T cell function.

Here we asked to what extent inhibition of glycolysis, using 2-Deoxy-D-Glucose (2DG), regulate

activation, effector function, or differentiation of human T cell subsets. We found that glycolysis is

required for T cell receptor (TCR) -mediated activation and proliferation of human naive CD4+ T

cells but had less of an impact on memory subsets. CD4+ T cells cultured in the presence of 2DG

displayed higher level of IL-17-secreting cells (Th17) from memory or *in vitro* differentiated naive

regulatory T cell (Tregs) subsets. Moreover, the mucosal associated invariant T (MAIT) cell subset

survived or expanded better and secreted higher IL-17 in the presence of 2DG. Remarkably, we

found that the 2DG effect was reversed by mannose but not by glucose. Collectively, these

findings suggest that 2DG could enrich IL-17 secreting human effector T cell subsets and their

cellular functions. Our finding provides a framework to manipulate glycolytic pathways in human

T cells in infectious diseases such as COVID19 and in enhancing cancer immunotherapy.

Introduction

Metabolism has been recognized as an important regulator in T cell activation and lineage differentiation (Araki et al., 2010; Chapman et al., 2020; Jacobs et al., 2008; MacIver et al., 2013; Palmer et al., 2015; van der Windt & Pearce, 2012). Upon T cell activation, TCR signals and co-stimulation activate the phosphatidtyl-inositide-3 kinase (PI3K)/Akt/mTORC1 signaling pathway for the increased anabolic needs of effector T cells, which become more sensitive to metabolic regulation (Ho et al., 2015; Palmer et al., 2015; Sena et al., 2013). Importantly, an efficient immune response involves various T cell subsets, which have different metabolic requirements for development and effector functions (MacIver et al., 2013; Pearce et al., 2013). It is still unclear how metabolic pathways regulate the functions of T cell subsets specifically, but understanding the underlining mechanisms would allow modulation of immunity in chronic disease and cancer (Johnson et al., 2018).

Subsets of T cells that secrete IL-17 are the critical mediators in antimicrobial and pro-inflammatory responses, as well as in pathogenesis in autoimmune or chronic inflammatory diseases (Bettelli et al., 2006; Damasceno et al., 2020; Platt et al., 2020; Xu et al., 2020; Yasuda et al., 2019). Th17 cells and a subset of Tregs are the major sources of IL-17 secretion, which are also characterized by the expression of the transcription factor RORC and the chemokine receptor CCR6 (Singh et al., 2008; Valmori et al., 2010; Wan et al., 2011). Several mouse studies have revealed a role of glucose metabolism in Th17 differentiation, via mTORC1, Myc, and HIF1a signaling (Kastirr et al., 2015; Perl, 2016; Sasaki et al., 2016; Shen & Shi, 2019; Shi et al., 2011). Inhibition of glycolysis with 2-Deoxy-Glucose (2DG) or 3-bromopyruvate in mice for a short period of time impaired Th17 differentiation, phenocopying the effect of HIF1a deficiency (Okano et al., 2017; Shi et al., 2011). However, how glycolytic metabolism regulates the differentiation and functions of human Th17 and IL-17 secreting Tregs remains unclear.

Another source of IL-17 producing T cells are mucosal associated invariant T (MAIT) cells (Coulter

et al., 2017; Willing et al., 2018). Human MAIT cells can be identified by a semi-invariant T cell

receptor(TCR) alpha chain(Va7.2) and CD161 expression (Chiba et al., 2018). MAIT cells can be

activated by a broad range of bacteria and yeasts, and can discriminate and fine-tune their

functional responses to complex human microbiota (Constantinides et al., 2019; Corbett et al.,

2020; Hinks & Zhang, 2020; Ioannidis et al., 2020; Tastan et al., 2018). Thus, IL17-secreting MAIT

cells play a role in infections and autoimmune diseases such as Tuberculosis and Multiple

Sclerosis (Coulter et al., 2017; Willing et al., 2018). While the frequency and distribution of MAIT

cells are currently under intense investigation, there is still much to understand in the metabolic

regulation of the function of the human MAIT cell subset.

In this study, we utilized 2DG, an analog and potent glycolysis inhibitor, to investigate the role of

glucose metabolism in the activation, differentiation, and effector function of human T cell subsets.

We showed that 2DG suppressed human T cell activation, cytokine production, and proliferation

upon early T cell receptor (TCR) activation. However, 2DG treated naive T cells showed a greater

reduced proliferative capacity and glycolytic function compared to memory T cells. Remarkably,

2DG treatment greatly enriched IL-17 producing CD4+ T cells in long term culture and during *in*

vitro differentiation of naive or naive regulatory T cells (TNreg) or from the MAIT cell subset.

Together, our findings suggest a differential impact of glycolysis in different subsets and

differentiation stages of human T cells.

Results

Inhibitor of glucose uptake, 2DG, suppresses early activation of CD4+ T cells

It has been reported that 2DG inhibits glucose metabolism (Xi et al., 2014), which may in turn regulate T cell activation (Buck et al., 2015; Palmer et al., 2015). Therefore, we first examined the effect of 2DG on T cell activation by assessing the expression of the IL-2 receptor alpha (CD25) and the glucose transporter 1 (GLUT1), as both are upregulated upon T cell activation for IL-2 signaling and glucose uptake, respectively (Chapman et al., 2019). Human primary CD4+ T cells were activated with anti-CD3/CD28 beads in media alone or in media supplemented with 0.3mM, 1mM, or 3mM 2DG. The expression of CD25 and GLUT1 in T cells was significantly reduced by 3mM 2DG compared to the control (**Figure 1A-B**), but lower concentrations of 2DG (0.3mM or 1mM) did not reach statistical significance (data not shown). Because 2DG suppresses glycolysis (Xi et al., 2014), we also activated T cell subsets in the glucose-free media as a comparison. In contrast to the effect seen with 2DG, T cells activated in glucose-free-media showed higher GLUT1 expression and CD25 expression comparable to the control condition (**Figure 1A-B**). We also observed a reduced expression of program cell death 1 (PD1) and Lymphocyte Activating 3 (LAG3) on 2DG-treated T cells, which are known to be induced upon TCR activation (Lichtenegger et al., 2018) (**Figure 1C-D**). However, the expression of PD1 or LAG3 was not significantly different between T cells activated in glucose-free or regular media (**Figure 1C-D**).

TCR-activation of T cells leads to a rapid production of cytokines such as TNF, IFNγ, and IL-2. As such, to determine whether 2DG also suppresses cytokine production, we collected activated T cell supernatants 2 days post CD3/CD28 activation and analyzed the cytokine levels using a cytometric bead assay. We found that the levels of TNF and IFNγ secretion were significantly reduced but there was no difference in IL-2 secretion in the presence of 2DG culture compared to control media (**Figure 1E**). Together, these data suggest that 2DG suppresses early human T

101   cell activation and cytokine production.

2DG has different effects on activation or effector functions of naive and memory T cells.

It has been reported that naive and memory T cells may have different metabolic requirements

for activation and proliferation (Almeida et al., 2016; MacIver et al., 2013; van der Windt & Pearce,

2012). To determine whether 2DG modulates the activation of naive vs memory CD4+ T cells

differently, we sorted these subsets from human CD4+ T cells using CCR7+CD45RO- (naive)

and CCR7+/-CD45RO+ (memory) markers and activated with anti-CD3/CD28 beads in media

alone, in 3mM 2DG, or in glucose-free media. Both T cell subsets displayed a similar trend of

reduction in GLUT1 and CD25 expression on day 1 post activation **(Figure 2A)**; however, on day

4 post-activation, 2DG-treated memory T cells upregulated GLUT1 to levels comparable to control

cells whereas most naive T cells remained GLUT1 negative **(Figure 2B-C)**. Similar to unsorted

CD4+ T cells (**Figure 1A**), activation of naive or memory T cells in glucose-free media did not

significantly affect the expression of GLUT1 or CD25 (**Figure 2A-C**). Addition of IL-2 in these

activation experiments did not restore the 2DG-mediated down-regulation of GLUT1 or CD25

expression in either naive or memory T cells (**Supplemental Figure 1A-C**). Since we have

observed that 2DG suppresses T cell activation and glucose uptake, we hypothesized that 2DG

would also reduce T cell proliferative capacity. To address this, we labeled naive and memory T

cells with Celltrace violet (CTV) dye, and then activated with anti-CD3/CD28 coated beads.

Indeed, in 2DG-treated or in glucose-free media, naive or memory T cell subsets proliferated less

compared to control media (**Figure 2E**). Addition of IL-2 did not restore the low proliferation of

2DG-treated T cell subsets (S**upplemental Figure 1D**).

We next monitored the expansion of these activated T cells during 2-week culture in IL-2.

Expansion of 1 mM 2DG treated naive T cells were 30 fold after 14 days, compared to 150 fold

in the control, whereas memory T cells in the presence of 1mM 2DG expanded 85 fold compared

to 110 fold in the control media. We then asked whether 2DG has differential effects in the

glycolytic function of naive and memory subsets by using the Seahorse glucose stress test assay.

After T cell expansion in 2DG or control media for 14 days, 2DG was washed away and cells were

resuspended in Seahorse XF base media, and the metabolic functions were analyzed by the

Seahorse XFe96 Analyzer. The extracellular acidification rate (ECAR) was assessed after the

addition of glucose (gluc), oligomycin (oligo), and 2DG at indicated times and the glycolytic

capacity was determined. 2DG-treated memory T cells had comparable glycolytic function to the

control groups; however, 2DG-treated naive T cells showed significantly reduced glycolysis in

response to addition of glucose (**Figure 2F-G**).

2DG enhances IL-17 producing CD4+ memory subsets.

We next further explored which effector T cell subsets are modulated by suppressing glycolysis.

We first observed a significantly increased CD161+ and CCR6+ populations in 2DG-treated CD4+

T cells compared to control (**Figure 3A and 3B**). Since nearly all IL-17 secreting cells express

CCR6 and CCR6+ Th17 cells are reported enriched in the CD161+ subset (Acosta-Rodriguez et

al., 2007; Cosmi et al., 2008; Wan et al., 2011), we hypothesized that 2DG would preferentially

affect the function of IL-17 secreting T cell subsets. Indeed, after two weeks in culture with 2DG,

we observed that IL-17+ subsets were increased in both CCR6+ and CD161+ populations

compared to cells in control media, but interestingly, there was no difference in the low glucose

condition (**Figure 3C**). After 2DG was removed from the culture and cells reactivated with phorbol

myristate acetate (PMA) and calcium ionophore (ionomycin), we determined intracellular cytokine

secretion in the presence of brefeldin a (GolgiStop). We found that the frequency of cells that

expressed not only IL-17 but also IFNγ, IL-4, and IL-21 were significantly increased (**Figure 3C**

and 3D). However, no effect was seen in IL-2 and TNF production from CD4+ T cells

(**Supplemental Figure 3**). Taken together, these data demonstrate that 2DG-treatment can

modify the cytokine production of lineage-committed cells in long term *in vitro* cultures.

2DG enriches IL-17+ cells within Th17 lineage-committed CCR6+ memory T cell subset.

Since we observed a significant increase in IL-17 cytokine production from 2DG-treated T cells,

we hypothesized that 2DG could enhance the IL-17 production from Th17 cells compared with

other effector T cells. To test this hypothesis, we sorted CD4+ T cells into CCR6- and CCR6+

memory T cell subsets, activated them in the presence or absence of 2DG, and expanded for two

weeks in IL-2, as described above (**Figure 3A**). Consistent with our prior results, 2DG-treated

CCR6- and CCR6+ CD4+ T cells displayed higher CD161 expression than control cells (**Figure**

4A). IL-17 production from purified and expanded CCR6+ T cells were increased in the presence

of 2DG (**Figure 4B**). Notably, IFNγ+ and IL-21+ producing CCR6+ T cells were also increased by

2DG (**Figure 4B-E**). Interestingly, the percentage of IL-10+ and IL-17+ T cells was significantly

increased in 2DG-treated CCR6+ T cells (**Figure 4F and 4G**). Therefore, we concluded that 2DG

increased CD161+ within both CCR6+ and CCR6- T cells, and also enhanced IL-17, IFNγ, IL-21,

and IL-10 production from the sorted CCR6+ memory T cell subset.

2DG increases *in vitro* generation of IL-17-producing Tregs from human naive regulatory

T cell (TNreg) precursors

A subset of Tregs that secretes IL-17 can preferentially arise from human naive precursors

(CD25+CD45RO-CCR7+ cells, termed TNreg) in the presence of polarizing-cytokines IL-1β, IL-

23, and TGF-β (Mercer et al., 2014; Valmori et al., 2010). To explore how 2DG regulates Th17

cell differentiation from TNregs, we used this cytokine polarization protocol (**Figure 5A**) to

differentiate and expand IL-17-secreting cells from highly purified naive or precursors naive

regulatory T cells (TNregs) in media alone or in the presence of 1 mM 2DG. The FOXP3+ cells

during polarization of TNregs with 2DG were statistically similar in frequency (**Figure 5B**).

However, 2DG significantly increased generation of IL-17+ Tregs from TNreg precursors (**Figure**

5C). Since HELIOS expression defines a phenotypically distinct population of Tregs (Thornton et

al., 2019), and *in vitro*-generated IL-17+ cells from TNreg precursors did not express HELIOS

(Mercer et al., 2014), we then asked whether 2DG could induce IL-17, IFNγ, and IL-21 production

from HELIOS+ or HELIOS- populations of naive and TNregs. We also compared the 2DG effects

between HELIOS+ and HELIOS- population of naive cells since HELIOS plays a role in naive T

cell differentiation (Ng et al., 2019). The frequency of IL-17+ T cells witihn HELIOS- naive T cells

was increased by 2DG in polarizing condition, but not within HELIOS+ subset (**Supplemental**

Figure 4). 2DG also significantly increased IL-17 and IFNγ populations from HELIOS- population

from TNregs in the polarization condition (**Figure 5D-F**). Notably, when comparing the intracellular

IL-21 from HELIOS+ and HELIOS- subset of Tregs, the frequency of IL-21+ cells was significantly

higher in 2DG-treated HELIOS- subset (**Figure 5G and 5H**). But there was no significant

difference in IL-21 cytokines secretion in HELIOS+ or HELIOS- Treg subsets in 2DG-treated naive

T cells compared to controls (**Figure 5H**). Intracellular IL-10 was also increased in HELIOS-

population of TNregs in polarizing cytokines, and was significantly increased in HELIOS+ and

HELIOS- population from naive T cells (**Figure 5J**). In addition, increased cytokine production of

IL-17A, IL-17F, IL-10, and IFNγ was also confirmed by corresponding levels of cytokines detected

in the culture supernatants (**Figure 5K**). Taken together, 2DG enhanced the generation of IL-17-

producing cells from both naive and TNregs, which was restricted to HELIOS- subsets. 2DG also

increased the production of IFNγ, IL-21, and IL-10 from TNregs in the polarizing condition.

Mannose reverses or rescues 2DG effects on T cell activation and effector functions.

2DG has previously been shown to inhibit *N*-linked glycosylation (Andresen et al., 2012; Xi et al.,

2014), which can be modulated through a branch of the glucose metabolism pathway called the

hexosamine biosynthetic pathway (HBP) (Akella et al., 2019). Further, mannose, which is a major

component of sugar moeity in glycoproteins, can reverse the apoptotic effect of 2DG on cancer

cells (Ahadova et al., 2015; Gu et al., 2017; Kurtoglu et al., 2007). Thus, we sought to determine

whether mannose can also reverse the 2DG effects on short term activation of T cells (**Figure 2**).

Accordingly, we activated CD4+ naive T cells with anti-CD3/CD28 beads in media alone, with

3mM 2DG or mannose alone, or with 3mM 2DG supplemented with 3mM mannose. While we did

not observe any difference in CD25 and GLUT1 expression on day 1 post activation

(**Supplemental Figure 5**), mannose reversed the 2DG-mediated downregulation of CD25 and

GLUT1 expression on day 4 (**Figure 6A and 6B**). Additional equimolar glucose did not reverse

the 2DG effects (**Figure 6B**). Equal molar of mannose significantly reverses the proliferative

capacity from 1 fold expansion with 2DG alone to 50 fold expansion with 2DG and mannose

comparing to 157 fold cell expansion in control media (**Figure 6C**). However, after long-term

culture, mannose did not rescue the impaired glycolytic function of 2DG-treated naive T cells

(**Figure 6D and 6E**).

Next, we asked whether mannose could reverse 2DG effects on IL-17 enrichment. After 14 days

cell expansion, we observed a reduction of CCR6+ and CD161+ population in a mannose-does

dependent manner (**Figure 6F and 6G**). In addition, the 2DG mediated-reduction of IL-17, IFNγ,

and IL-21 were also reversed by mannose (**Figure 6G**). Addition of equimolar glucose into 2DG

culture did not reverse the 2DG-mediated enrichment of IL-17 production (**Figure 6F and 6G**).

Together these findings suggest that the inhibitory effects of T cells activation and IL-17

production on 2DG-treated human T cells are mannose-dependent.

Discussion

In this study, we demonstrated that the glucose analog 2DG have remarkably different effects on

activation and differentiation of human T cell subsets. 2DG suppresses early human T cell

activation through the TCR, as assessed by cytokine production, and proliferation. However, in

long term cultures 2DG also profoundly increases the frequency of IL-17 production from lineage-

committed memory CCR6+CD4+ (Th17) T cells, MAIT cells and Tregs. In addition, inhibition of

glycolysis by 2DG in these memory subsets (Th17, MAIT and Tregs) does not affect their

expansion upon activation. On the other hand, naive CD4+ T cells are more sensitive to the

inhibition of glycolysis by 2DG and therefore have reduced their survival and function. Importantly,

we also demonstrated that the 2DG-mediated IL-17 enrichment could be reversed by addition of

equimolar mannose.

A major question derived from our findings is that how inhibition of glycolysis can significantly

increase the IL-17 production from already lineage-committed/differentiated memory Th17 cells?

Several mice studies have revealed a role of glucose metabolism in Th17 differentiation (Kastirr

et al., 2015; Perl, 2016; Sasaki et al., 2016; Shen & Shi, 2019; Shi et al., 2011) but showed

discrepancies on the effect of inhibiting glycolysis in Th17 differentiation and functions. For

example, in one study Tregs treated with 0.2mM 2DG for 5 days reduced IL-17 production *in vitro*

(Li et al., 2019) and in another study reduced production of IL-17 by T cells after 5 day-2DG

treatment was observed (Shi et al., 2011). However, another study reported an enhancement of

Th17 cell polarization and IL-17 production after treatment of 2mM 2DG (Brucklacher-Waldert et

al., 2017). In our study, we observed reduced IL-17 production by 2DG treated T cells for during

early activation period. However, frequency of IL-17 producing subset was greately increased

after long-term T cell culture (12-14 days), during reactivation in the absence of 2DG. Th17 can

be characterized as "pathogenic" and "non-pathogenic" Th17 cells (Wu et al., 2018). Pathogenic

Th17 cells express more effector molecules such as CXCL3, CCL4, CCL5, IL-3, and IL-22,

whereas non-pathogenic Th17 cells exhibit upregulation of immune suppressive molecules and

cytokines such as IL-10 (Gaublomme et al., 2015; Lee et al., 2012). We hypothesized that the

increased subset of IL-17-secreting Th17 cells belongs to the "non-pathogenic" subset, as we

also found an increase in IL-10 production from the same cells. Since IL-17+ Th17 cells are crucial

for mediating mucosal immunity against fungi and bacteria infection in human subjects (Brembilla

et al., 2018; McDonald, 2012), our findings may suggest that the metabolic changes affect the

function of Th17 against commensal microbes in the gastrointestinal tract which would lead to

mucosal damage and mucosal-relative diseases such as inflammatory bowel disease (IBD)

(Galvez, 2014). Since the functions of Th17 cells in the small intestine can be closely regulated

by gut microbiome (Evans-Marin et al., 2018; Garidou et al., 2015), our findings also suggest that

a dysregulated metabolites from dysbiotic ileum microbiota (Visconti et al., 2019) could impact

Th17 homeostasis, and were sufficient to induce metabolic disease such as Type 2 Diabetes

(T2D) (Garidou et al., 2015).

We found that 2DG also increased the frequency of IL-17+ Tregs and IL-17 cytokine secretion

from these cells in the presence of polarizing cytokines *in vitro* differentiation. These IL-17-

producing Tregs were found to be accumulated in the inflamed joints of patients with rheumatoid

arthritis and were functionally suppressive (Afzali et al., 2013; Jung et al., 2017). However, we did

not observe a significant difference in suppressive capacity of 2DG-treated Tregs in proliferation

assay (data not shown), which is consistent with a human Tregs study *in vitro* (Tanimine et al.,

2019). Inhibition of glycolysis may also contribute to mucosal diseases, as IL-17+Tregs may

contribute to the development of colon cancer (Knochelmann et al., 2018; Marshall et al., 2016),

and possibly to IBD (Galvez, 2014).

As noted, an increased level of IL-10 from memory T cells (such as Tregs in the presence of

polarizing cytokines, CCR6- and CCR6+ memory T cells) was also observed throughout the

experiments. IL-10 is an anti-inflammatory cytokines, plays a central role in limiting the regulation

of immune responses (Iyer & Cheng, 2012). Elevated IL-10 signaling can inhibit antigen

presenting cells maturation, and chemokine secretion of the host during chronic viral infection

(Granelli-Piperno et al., 2004) and autoimmune disease (Braat et al., 2003). It can be highly

produced by a subset of Tregs, called Type 1 regulatory cells (Tr1) (Zeng et al., 2015), which is

associated with autoimmune diseases such as IBD, multiple sclerosis(MS), and type 1 diabetes

mellitus when their frequency was found reduced (Jia et al., 2019). Thus our data may suggest a

potential role of glycolysis in the function of IL-10 secreting subsets and related diseases.

Another subset of cells, where found differential and specific effect of 2DG were MAIT cells, which

are characterized with expression of CD161 and a semi-invariant Va7.2+TCR, are activated by a

ligand from riboflavin metabolism called 5-ARU in the context of MR-1 molecule (Godfrey et al.,

2019). MAIT cells are also unique in that they can be CD8+CD4- or CD8-CD4- and can also

produce IL-17 and mostly migrate to mucosal regions, thus potentially play an important role in

mucosal homeostasis and microbiome regulation (Oh & Unutmaz, 2019). In our experiments, we

had found that 2DG-treated cells compared to control induced higher frequency of CD161+ cells,

which is also highly expressed on MAIT cells (CD161hiVa7.2+) (Cosmi et al., 2008; Kleinschek

et al., 2009). We also identified a distinct subset exhibiting medium-expression of CD161 which

is also enhanced by 2DG, but is neither Va7.2+ (MAIT) nor CCR6+ (Th17) cells. The identity of

this subset is not fully clear, but Klenerman et al reported a distinct functional subset in human T

cells which had mid-level CD161 expression, produced IL-17/IFNγ, and were found to be highly

enriched in chronic inflammation (Billerbeck et al., 2010). Future studies will be needed to better

characterize this subset and its functions, as these have also shown different glucose tolerance

than other effector cells.

Our findings that 2DG enhances IL-17 production also in MAIT cells, in addition to Th17 cells,

suggest a shared metabolic pathways that regulate the effector functions of these mucosa-

associated subsets. Interestingly IL-17+ MAIT cells are found specifically enriched from the

peripheral blood of multiple sclerosis (MS) patient, implicating a proinflammatory role in

autoimmune diseases (Willing et al., 2018). IL-17+ MAIT are also associated with inflamed

mucosal tissue, and are found activated and produce more IL-17 in IBD patients (Serriari et al.,

2014). In addition to the gut, CD8+ MAIT cells have also been shown to be resident in normal

skin and are thought to play a role in skin-associated inflammations, such as psoriasis and

dermatitis herpetimorfis (Willing et al., 2018) and tissue repair (Constantinides et al., 2019).

Therefore, it is tempting to speculate that 2DG or similar analogs of glucose could be used to

modulate pathogenic responses of these T cell subsets.

An increase in IL-21 production in 2DG-treated memory T cells was also shown significant in our

study. IL-21 has been documented to regulate the differentiation and function of several CD4+ T

cells subsets, including Th17 cells (Leonard & Wan, 2016; Nurieva et al., 2007; Zhou et al., 2007),

Th2 (Frohlich et al., 2007; Lajoie et al., 2014), and T follicular helper cells (TFH) (Vogelzang et

al., 2008). Thus, the enhanced IL-17 and IL-4 cytokine production in 2DG treated T cells may be

mediated partially by IL-21 signaling. IL-21 was also reported to induce IL-10 production in Th17

polarizing condition in mice (Spolski et al., 2009), which is consistent with our observation that

differentiation of IL-17 secreting subsets from TNreg with 2DG increase IL-21 and IL-10 secretion.

Furthermore, the functional significance of IL-21 in regulating effector functions of CD8+ T cells

is highlighted by its critical role in sustaining anti-viral CD8+T cells during chronic LCMV infections

(Cui et al., 2011; Elsaesser et al., 2009) and its potent effects to induce and expand cytotoxic

CD8+ T cells for cancer immunotherapy (Davis et al., 2015; Santegoets et al., 2013). An increase

production of IL-21 and IL-4 cytokines may suggest that 2DG enhance TFH cells, which have high

expression level of CXCR5 and produce high level of IL-21 (Crotty, 2014). This hypothesis is also

supported by that exogenous Ag-specific TFH cells do not require glycolysis (Choi et al., 2018).

TFH is critical for the formation and maintenance of germinal centers and provide help for B cells

generating antibody responses after immunizations (Bentebibel et al., 2013; Duan et al., 2014;

Spolski & Leonard, 2010). Therefore, regulating TFH cells with glucose metabolism would

potentially enhance development of new or improved vaccines (Crotty, 2014).

One potential mechanism of an increase frequency of IL-17-secreting T cell subsets in 2DG-

treated cells is that 2DG alters T cells metabolism from glucose to alternative energy sources or

metabolic pathways such as fatty acid oxidation or pentose phosphate pathway. Previous studies

showed an inhibition of glucose metabolism with a glucose transporter inhibitor (CG-5) promoted

fatty acid oxidation and the pentose phosphate pathway in CD4+ T cells in a similar manner as

with 2DG (Li et al., 2019). Fatty acid metabolism has also been suggested to be involved in Th17

inflammation, as blockade of carnitine palmitoyltransferase 1 (CPT1), an enzyme responsible for

catalyzing the breakdown of long-chain fatty acids, inhibits Th17-associated cytokine production

from type 2 diabetes (T2D) patients (Nicholas et al., 2019).

One of our important findings towards mechanism of action of 2DG in our system is that 2DG-

mediated increases in IL-17-secreting cells can be reversed by addition of equimolar mannose

(Berthe et al., 2018; Kurtoglu et al., 2007). This suggests that another possible mechanism of

2DG is inhibiting the initial step of glycolysis affects the downstream metabolic pathway of

mannose. Since mannose is a dominant monosaccharide in N-linked glycosylation (Imperiali &

O'Connor, 1999), our observation may further suggest that part of the 2DG effects on T cell

function are through modifying sugar moieties generated in central carbon metabolism (e.g. N-

linked glycosylation). N-linked glycosylation plays a role in modulating activation and cytokine

signaling (Baum & Cobb, 2017; Dean et al., 1979; Hauser et al., 2016), and thereby may affect

differentiation and function of T cell subsets. An evidence to support this notion was our finding

that mannose reversed 2DG-mediated down-regulation of CD25 expression upon activation.

Similarly, another inhibitor of N-linked glycosylation, glucosamine, attenuated CD25 surface

retention on T cells (Chien et al., 2015), and down-regulation of N-linked glycosylation promotes

Th17 differentiation (Chien et al., 2015). However, surprisingly, addition of equimolar glucose

could not reverse 2DG-mediated downregulation of CD25. We reason that 2DG efficiently inhibits

glycolysis, therefore 2DG-treated cells could not utilize additional glucose. This speculation could

be supported by seahorse glycolytic analysis showing that neither additional mannose nor

glucose could reverse 2DG-mediated glycolytic capacity. Overall, it is conceivable that 2DG may

have dual mechanism in enhancing the survival and effector functions of IL-17 secreting subsets.

Recently, 2DG has also been considered as a potential therapy during viral infections such as

COVID-19 (Yang et al., 2021). Indeed, 2DG was approved by the Indian Council of Medical

Research (OCMR) to treat COVID-19 patients in India (Sahu & Kumar, 2021). 2DG shortens the

time of COVID patients recovered from the infection and there was a higher proportion of patients

with improved symptoms who were free from supplemental oxygen dependence (Sahu & Kumar,

2021) . The underling mechanism of the effect of 2DG in these COVID patients is still unclear, but

it was suggested that inhibiting glycolysis with 2DG could affected viral life cycle (Passalacqua et

al., 2019) and an increased glucose level could enhance the suppressive function of monocytes

and therefore enhanced SARS-CoV-2 infection (Codo et al., 2020). Given our findings, we also

speculate that 2DG effects may have been through ameloriation of excessive or uncontrolled

immune response during COVID19, which is observed during later severe disease stages. 2DG

has also been tested to treat advanced cancer and proven safe in phase I studies (Raez et al.,

2013; Stein et al., 2010). Our data shows long-term culture of human primary T cells with 2DG

enriched IL-17 secreting T cells, which have been reported to exhibit pathogenic features (Basdeo

et al., 2015). However, a significant increase in IFNγ production by 2DG may enhance therapeutic

anti-tumor activity (Medrano et al., 2017). Taking together, our study provides insights for future

clinical trials and strategies for development of 2DG-related cancer therapies.

Materials and Methods

T cell purification, activation, and culture

PBMCs from healthy individuals (New York Blood Center, New York, NY) were isolated using Ficoll-Paque plus (GE Healthcare). CD4+ T cells were isolated using Dynal CD4+ isolation kits (Invitrogen) and were 99% pure. Purified CD4+ cells were sorted in some experiments by flow cytometry (FACSAria; BD Biosciences) based on CD45RO, CCR7, CD25, and chemokine receptors expression into: 1) naive T cells (CD45RO-CCR7+CD25-), 2) memory T cells (CD45RO+CD25-), 3) naive regulatory T cells (Tregs; CD45RO-CD25+), 4) Th17 cells (CD45RO+CCR6+). Sorted subsets were 98% pure. All purified cells were kept at 37°C and 5% CO_2 in complete RPMI 1640 media (RPMI 1640 supplemented with 10% FBS; Atlanta Biologicals, Lawrenceville, GA), 8% GlutaMAX (Life Technologies), 8% sodium pyruvate, 8% MEM vitamins, 8% MEM nonessential amino acid, and 1% penicillin/streptomycin (all from Corning Cellgro). To activate cells for expansion *in vitro*, anti-CD3/CD28 Dynabeads (Invitrogen) were used at a bead: cell ratio of 1:2 and the activated cells were cultured in complete RPMI 1640 medium (Thermo Fisher Scientific) or complete 1640 RMPI no glucose medium supplemented with IL-2 (10 ng/ml). For MAIT experiments, MAIT cells were activated by adding riboflavin metabolite 5ARU (Toronto Research, Ontario, Canada) into PBMC culture.

Flow cytometry staining and analysis

Cells were stained with fluorochrome-conjugated Abs in FACS buffer(PBS+2% FCS and 0.1% sodium azide) for 30 min at 4°C (or room temperature for CCR6 staining). Abs used in surface staining are IL-2R (CD25), PD1, CD161, Va7.2, CCR6 (from BioLegend), LAG3, CD3, and CD8

(from Invitrogen). For intracellular cytokine staining, cells were stimulated for 4 h at 37°C with

PMA (10ng/ml for PBMC and CD4+ T cells for 40 ng/ml) and ionomycin (500 ng/ml) (both from

Sigma-Aldrich) together with GolgiStop (BD Biosciences). Cells were then stained with fixable

viability dye (eBioscience) in PBS and surface marker Abs in PBS or FACS for 30 minutes, then

fixed and permeabilized using eBioscience fixation/permeabilization buffers for 30 min at 4°C

according to the manufacturer's instructions, before staining for intracellular cytokines for 30 min

at 4°C. Abs used for intracellular cytokines are IFNγ, TNF, IL-17A, IL-10, and IL-21(BioLegend).

Ab used for Regulatory T cells transcription factor intracellular staining are Foxp3 and Helios Abs

(BioLegend). Flow cytometry analyses were performed using an LSRFortessa X-20 flow

cytometer (BD Biosciences) and SP6800 spectral cell analyzer (Sony Biotechnology).

Seahorse metabolism assay

Seahorse assays were performed according to the manufacturer's instructions to analyze

glycolysis metabolism using the Seahorse XF Glycolysis Stress Test Kit (Agilent Technologies).

Briefly, 14 day post activation, naive and memory T cells expanded in indicated conditions were

washed and resuspended in glucose-free media (Gibco). 300k cells per well were spun down

onto plates coated with Cell-Tak (Fisher scienctific). Four replicates were set up for each

condition. Glucose, oligomycin, and 2DG were serially injected to measure metabolic function.

Plates were analyzed using an XF^e 24 Extracellular Flux Analyzer (Agilent Technologies).

Glycolysis was calculated as average post-glucose ECAR values minus average basal ECAR

values.

In vitro **cytokine polarization assay**

Sorted TN and TNreg were activated with anti-CD3/CD28 beads and cultured in complete RPMI

media containing IL-2 10ng/ml (Chiron), together with IL-1β (10ng/ml), TGF-β (10ng/ml), and IL-

23 (100ng/ml) (R&D Systems). Cells were expanded for 2 weeks in media replenished for 2DG

and IL-2.

Statistical analysis

Data recorded by flow cytometry were analyzed using FlowJo (Tree Star, Ashland, OR). Statistical

analyses were performed using GraphPad Prism 6.0 software (GraphPad Software, La Jolla, CA).

Error bars represent SEM. Results were compared using two-tailed t tests. Bonferroni corrections

were applied for multiple comparisons. For all experiments, significance was defined as $*p < 0.05$,

$**p < 0.01$, $***p<0.001$

Conflict of Interest Statement

424   The authors declare that the research was conducted in the absence of any commercial or

425   financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

428   The research in this study was supported by National Institute of Health (NIH) grants

429   U54NS105539 and R01AI121920 to D.U.

Figures Legends:

Figure 1. 2DG suppresses early activation of CD4+ human T cells. T cell subsets were activated with anti-CD3/CD28 beads in media alone, 3mM 2DG or in glucose-free media. (A) Surface expression of CD25 and GLUT1 on resting and 24h-activated CD4+ T cells with indicated conditions. (B) Fold difference in GLUT1 expression and mean florescent intensity (MFI) of CD25 on 24h-activated CD4+ T cells in 3mM 2DG or in glucose-free media compared to control. (C) Flow cytometry plot of LAG3 and PD1 expression on resting and activated CD4+ T cell in media alone, 3mM 2DG, or in glucose-free media for 24h. (D) Statistical analysis of fold difference in PD1, LAG3, and double positive expression of CD4+ T cells activated for 24 h in 3mM 2DG and in glucose-free media. (E) CD4+ T cells were activated with anti-CD3/CD28 beads. Cell supernatant were collected from the indicated conditions after 2 days post beads activation. The levels of TNF, IFNg, and IL-2 cytokines were determined by FlowCytomix Multiplex bead assay. Data represent three independent experiments. **p <0.01, ***p <0.001

Figure 2. 2DG has differential effects on activation of naive and memory T cell subsets. (A) Naive and memory CD4+ T cells were activated with anti-CD3/CD28 beads in media alone, 3mM 2DG, or in glucose-free media. Media was not supplemented with IL-2. Representative flow cytometry plot of GLUT1 and CD25 surface expression on resting, and day 1(A)- or day 4(B)- post activated Naive (top) and memory T cells (bottom) in indicated conditions were shown. (C) Fold difference in frequency of GLUT1 expression or mean fluorescence intensity of CD25 expression on naive or memory T cells on day 1 (black)- and day 4 (grey)-post activation in indicated conditions. (D) Naive and memory CD4+ T cells were labeled with CellTrace violet (CTV) dye followed by anti-CD3/CD28 beads activation in indicated conditions. Representative histogram plot of CTV-labeled naive (up) and memory (bottom) CD4+ T cells in 3mM 2DG (red) or in glucose-

free media (blue) after day 4 and day 6 post TCR activation. (E) Representative fold expansion

of naive and memory CD4+ T cells in 1 or 3mM 2DG or in glucose-free media after 2 week T cells

expansion. (F and G) Naive and memory CD4+ T cells were activated and expanded in media

alone (control, black) or 1mM 2DG (red) for 14 days. Metabolic functions were analyzed using

Seahorse glucose stress test assay. The extracellular acidification rate (ECAR) was assessed

after the addition of glucose (gluc), oligomycin (oligo), and 2DG at indicated times and the

glycolytic capacity was determined. Each set of data is representative of three donors. $*p < 0.05$,

$**p < 0.01$, $***p<0.001$

Figure 3. 2DG enhances IL-17 production in primary CD4+ T cells. Purified CD4+ T cells

were activated with aCD3/CD28 beads in media alone, 3mM 2DG, and in media with 0.5mM

glucose. (A) The representative flow cytometry plot of CCR6 and CD161 expression on day14

post- activated T cells were shown. (B) Statistical analysis of fold difference in frequency of

CCR6+ and CD161+ T cells in indicated conditions. (C) After 14 days' cell culture, T cells activated

and expanded in indicated condition were re-stimulated with PMA/ionomycin for 4h, followed by

intracellular staining as described in the method section. The percentage of IL-17 production from

CD161+ and CCR6+ T cells were shown. (D) Statistical analysis of fold difference in frequency of

IL17+, IFNg, IL-4, and IL-21+ T cells from indicated conditions was shown. Each set of data is

representative of three donors. $*p < 0.05$, $**p < 0.01$, $***p<0.001$

Figure 4: 2DG enriches the frequency of IL-17 from sorted CCR6+ memory T cells. (A) sorted

CCR6+ and CCR6- memory cells were activated with aCD3/CD28 beads in media alone or with

3mM 2DG for 14 days. The frequency of CD161+ cells in day-14 post activated sorted CCR6+

and CCR6- subsets in 3mM 2DG compared to control were shown. (B) 14 days post activated T

cells were re-stimulated with PMA and ionomycin for 4h, followed by surface and intracellular

staining. The frequency of IFNg and IL-17 was determined from sorted CCR6- and CCR6+ T cells.

(C) Statistic analysis of the frequency of IL-17+, IFNg +, and IL17+ IFNg+ T cells from CCR6- and

CCR6+ T cells as in Figure 4B. The representative plot (D) and the statistical analysis (E) of the

frequency of IL-21, IL-10, and IL-17 was determined from sorted CCR6- and CCR6+ T cells after

14 days expansion and re-activation with PMA and ionomycin. The representative plot (F) and

the statistical analysis (G) of the frequency of IL-10 and IL-17 was determined from sorted CCR6-

and CCR6+ T cells after 14 days expansion as in Figure 4D and E. Each set of data is the

representative of three donors. *p < 0.05, ***p < 0.001

Figure 5. 2DG enhances *in vitro* generation of IL-17-producing cells from TNreg cells. (A)

Polarization protocol with Naive or TNreg cells. (B) Representative FOXP3/HELIOS expression

of TNreg in two-week polarization cultures with or without 1mM 2DG. (C) Fold difference of IL-17

production from 1mM 2DG treated TNreg cells with polarizing cytokines (P.C.) or without (-)

compared with control after 14-day expansion. (D) Representative flow cytometry plots of

intracellular IFNγ and IL-10 secretion within HELIOS+ and HELIOS- gated populations of TNreg

after 14-day expansion. (E) Statistical analysis of IL-17 production of indicated subsets from 1mM

treated naive T cell (left)/TNreg(right) with or without polarizing cytokines. (F) Statistical analysis

of IFNγ from 1mM treated TNreg in polarizing cytokines (G) Representative flow cytometry plots

of intracellular IL-21 and IL-17 secretion within HELIOS+ and HELIOS- gated populations of

TNreg after 14-day expansion. (H) Statistical analysis of IL-21 production of indicated subsets

from 1mM treated naive T cell (left)/TNreg(right) with or without polarizing cytokines. (I)

Representative flow cytometry plots of intracellular IL-10 and IL-17 secretion within HELIOS+ and

HELIOS- gated populations of TNreg after 14-day expansion. (J) Statistical analysis of fold

difference of IL-10 production of indicated subsets from 1mM treated naïve T cell

(left)/TNreg(right) with or without polarizing cytokines. (K) Day-14 expanded TNreg with or without

polarization condition were re-stimulated with anti-CD3/CD28 beads. Cell supernatant were

collected from the indicated conditions after 2 days post beads activation. IL-17A, IL-17F, IFNγ,

and IL-10 productions from supernatant were determined by FlowCytomix Multiplex bead assay.

Data represent three independent experiments. *p < 0.05, **p < 0.01, ***p<0.01

Figure 6. Mannose reverses or rescues 2DG effects on T cell subsets. (A) Naive CD4+ T

cells were activated with aCD3/CD28 beads in indicated conditions. Representative flow

cytometry plots of GLUT1 and CD25 surface expression. (B) Statistical analysis of fold difference

in GLUT1 and Mean Florescent Intensity (MFI) of CD25 on activated naive T cells as in Figure

6A. (C) Representative fold expansion of naive CD4+ T cells in indicated conditions after 2-week

T cells expansion. (D and E) Naive T cells were activated and expanded in the indicated

conditions for 14 days. 2DG was washed away and the metabolic functions were analyzed using

Seahorse glucose stress test assay. The extracellular acidification rate (ECAR) was assessed

after the addition of glucose (gluc), oligomycin (oligo), and 2DG at indicated times and the

glycolytic capacity was determined. (F) CD4+ T cells were activated with aCD3/CD28 beads in

media alone, 3mM 2DG alone, 3mM 2DG plus 1mM or 3mM mannose. 14-days post activated T

cells were re-stimulated with PMA and ionomycin for 4h, followed by surface and intracellular

staining. Representative flow cytometry plots and statistical analysis of T cells expressed CCR6,

CD161, IL-21, and IL-17. Data represent three independent experiments. *p < 0.05, **p < 0.01,

***p<0.01

Supplemental Figure Legends

Supplemental Figure 1. Addition of IL-2 did not restore 2DG effects on naive and memory

T cell subsets. (A) Naive and memory CD4+ T cells were activated with aCD3/CD28 beads in

media alone, 3mM 2DG, or in glucose-free media. Media was supplemented with IL-2.

Representative flow cytometry plots of GLUT1 and CD25 surface expression on resting, and day

1-(A) or day 4- post(B) activated Naïve (top) and memory T cells (bottom) in indicated conditions

were shown. (C) Fold difference in frequency of GLUT1 expression or mean fluorescence

intensity of CD25 expression on naive or memory T cells on day 1- and day 4-post activation in

media without IL-2. (D) Naive and memory CD4+ T cells were labeled with CTV dye followed by

aCD3/CD28 beads activation in media alone (control), 3mM 2DG, or in glucose-free media.

Representative histogram plot of CTV-labeled naïve (up) and memory (bottom) CD4+ T cells in

3mM 2DG (Red) or in glucose-free media(blue) after day 4 and day 6 post beads activation. Data

represent three independent experiments. ***p<0.01

Supplemental Figure 2. GLUT1 expression was down-regulated on naive T cells but not

memory T cells after long-term culture. Naive and memory CD4+ T cells were activated with

aCD3/CD28 beads in media alone (control in grey), or 1mM 2DG (in black). Media was not

supplemented with IL-2. Fold difference in frequency of GLUT1 expression on naïve or memory

T cells on day 14-post activation was shown. Data represent three independent experiments.

***p<0.01

Supplemental Figure 3. 2DG did not change IL-2 and TNF production by CD4+ T cells.

Purified CD4+ T cells were activated with aCD3/CD28 beads in media alone, with various doses

of 2DG, and in glucose-free media. After 14 days' cell culture, T cells activated and culture in

indicated condition were re-stimulated with PMA/ionomycin for 4h, followed by intracellular

staining as described in the method section. The percentage of IL-2 and TNF production from

CD4+ T cells were shown. Data represent three independent experiments.

Supplemental Figure 4. 2DG enhances *in vitro* generation of IL-17-producing cells from

naive T cells. Fold difference of IL-17 production from 1mM 2DG treated naive T cells with

polarizing cytokines (P.C.) or without (-) compared with control after 14-day expansion.

Supplemental Figure 5. Mannose did not reverse GLUT1 expression 2DG-treated naive T

cells. Naive CD4+ T cells were activated with aCD3/CD28 beads in media alone, 3mM 2DG

alone, 3mM 2DG plus 1mM or 3mM mannose. Representative flow cytometry plots of GLUT1 and

CD25 surface expression on naive T cells activated for 24 h in indicated conditions were shown.

Data represent three independent experiments.

References

Acosta-Rodriguez, E. V., Rivino, L., Geginat, J., Jarrossay, D., Gattorno, M., Lanzavecchia, A., Sallusto, F., & Napolitani, G. (2007). Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol*, *8*(6), 639-646. https://doi.org/10.1038/ni1467

Afzali, B., Mitchell, P. J., Edozie, F. C., Povoleri, G. A., Dowson, S. E., Demandt, L., Walter, G., Canavan, J. B., Scotta, C., Menon, B., Chana, P. S., Khamri, W., Kordasti, S. Y., Heck, S., Grimbacher, B., Tree, T., Cope, A. P., Taams, L. S., Lechler, R. I., . . . Lombardi, G. (2013). CD161 expression characterizes a subpopulation of human regulatory T cells that produces IL-17 in a STAT3-dependent manner. *Eur J Immunol*, *43*(8), 2043-2054. https://doi.org/10.1002/eji.201243296

Ahadova, A., Gebert, J., von Knebel Doeberitz, M., Kopitz, J., & Kloor, M. (2015). Dose-dependent effect of 2-deoxy-D-glucose on glycoprotein mannosylation in cancer cells. *IUBMB Life*, *67*(3), 218-226. https://doi.org/10.1002/iub.1364

Akella, N. M., Ciraku, L., & Reginato, M. J. (2019). Fueling the fire: emerging role of the hexosamine biosynthetic pathway in cancer. *BMC Biol*, *17*(1), 52. https://doi.org/10.1186/s12915-019-0671-3

Almeida, L., Lochner, M., Berod, L., & Sparwasser, T. (2016). Metabolic pathways in T cell activation and lineage differentiation. *Semin Immunol*, *28*(5), 514-524. https://doi.org/10.1016/j.smim.2016.10.009

Andresen, L., Skovbakke, S. L., Persson, G., Hagemann-Jensen, M., Hansen, K. A., Jensen, H.,
& Skov, S. (2012). 2-deoxy D-glucose prevents cell surface expression of NKG2D ligands
through inhibition of N-linked glycosylation. *J Immunol*, *188*(4), 1847-1855.
https://doi.org/10.4049/jimmunol.1004085

Araki, K., Youngblood, B., & Ahmed, R. (2010). The role of mTOR in memory CD8 T-cell
differentiation. *Immunol Rev*, *235*(1), 234-243. https://doi.org/10.1111/j.0105-
2896.2010.00898.x

Basdeo, S. A., Moran, B., Cluxton, D., Canavan, M., McCormick, J., Connolly, M., Orr, C., Mills,
K. H., Veale, D. J., Fearon, U., & Fletcher, J. M. (2015). Polyfunctional, Pathogenic
CD161+ Th17 Lineage Cells Are Resistant to Regulatory T Cell-Mediated Suppression in
the Context of Autoimmunity. *J Immunol*, *195*(2), 528-540.
https://doi.org/10.4049/jimmunol.1402990

Baum, L. G., & Cobb, B. A. (2017). The direct and indirect effects of glycans on immune function.
Glycobiology, *27*(7), 619-624. https://doi.org/10.1093/glycob/cwx036

Bentebibel, S. E., Lopez, S., Obermoser, G., Schmitt, N., Mueller, C., Harrod, C., Flano, E.,
Mejias, A., Albrecht, R. A., Blankenship, D., Xu, H., Pascual, V., Banchereau, J., Garcia-
Sastre, A., Palucka, A. K., Ramilo, O., & Ueno, H. (2013). Induction of
ICOS+CXCR3+CXCR5+ TH cells correlates with antibody responses to influenza
vaccination. *Sci Transl Med*, *5*(176), 176ra132.
https://doi.org/10.1126/scitranslmed.3005191

Berthe, A., Zaffino, M., Muller, C., Foulquier, F., Houdou, M., Schulz, C., Bost, F., De Fay, E.,
Mazerbourg, S., & Flament, S. (2018). Protein N-glycosylation alteration and glycolysis
inhibition both contribute to the antiproliferative action of 2-deoxyglucose in breast cancer
cells. *Breast Cancer Res Treat*, *171*(3), 581-591. https://doi.org/10.1007/s10549-018-
4874-z

Bettelli, E., Carrier, Y., Gao, W., Korn, T., Strom, T. B., Oukka, M., Weiner, H. L., & Kuchroo, V.
K. (2006). Reciprocal developmental pathways for the generation of pathogenic effector
TH17 and regulatory T cells. *Nature*, *441*(7090), 235-238.
https://doi.org/10.1038/nature04753

Billerbeck, E., Kang, Y. H., Walker, L., Lockstone, H., Grafmueller, S., Fleming, V., Flint, J.,
Willberg, C. B., Bengsch, B., Seigel, B., Ramamurthy, N., Zitzmann, N., Barnes, E. J.,
Thevanayagam, J., Bhagwanani, A., Leslie, A., Oo, Y. H., Kollnberger, S., Bowness, P., .
. . Klenerman, P. (2010). Analysis of CD161 expression on human CD8+ T cells defines
a distinct functional subset with tissue-homing properties. *Proc Natl Acad Sci U S A*,
107(7), 3006-3011. https://doi.org/10.1073/pnas.0914839107

Braat, H., Peppelenbosch, M. P., & Hommes, D. W. (2003). Interleukin-10-based therapy for
inflammatory bowel disease. *Expert Opin Biol Ther*, *3*(5), 725-731.
https://doi.org/10.1517/14712598.3.5.725

Brembilla, N. C., Senra, L., & Boehncke, W. H. (2018). The IL-17 Family of Cytokines in Psoriasis:
IL-17A and Beyond. *Front Immunol*, *9*, 1682. https://doi.org/10.3389/fimmu.2018.01682

Brucklacher-Waldert, V., Ferreira, C., Stebegg, M., Fesneau, O., Innocentin, S., Marie, J. C., &
Veldhoen, M. (2017). Cellular Stress in the Context of an Inflammatory Environment
Supports TGF-beta-Independent T Helper-17 Differentiation. *Cell Rep*, *19*(11), 2357-
2370. https://doi.org/10.1016/j.celrep.2017.05.052

Buck, M. D., O'Sullivan, D., & Pearce, E. L. (2015). T cell metabolism drives immunity. *J Exp Med*,
212(9), 1345-1360. https://doi.org/10.1084/jem.20151159

Chapman, N. M., Boothby, M. R., & Chi, H. (2019). Metabolic coordination of T cell quiescence
and activation. *Nat Rev Immunol*. https://doi.org/10.1038/s41577-019-0203-y

Chapman, N. M., Boothby, M. R., & Chi, H. (2020). Metabolic coordination of T cell quiescence
and activation. *Nat Rev Immunol*, *20*(1), 55-70. https://doi.org/10.1038/s41577-019-0203-
y

Chiba, A., Murayama, G., & Miyake, S. (2018). Mucosal-Associated Invariant T Cells in
Autoimmune Diseases. *Front Immunol*, *9*, 1333.
https://doi.org/10.3389/fimmu.2018.01333

Chien, M. W., Lin, M. H., Huang, S. H., Fu, S. H., Hsu, C. Y., Yen, B. L., Chen, J. T., Chang, D.
M., & Sytwu, H. K. (2015). Glucosamine Modulates T Cell Differentiation through Down-
regulating N-Linked Glycosylation of CD25. *J Biol Chem*, *290*(49), 29329-29344.
https://doi.org/10.1074/jbc.M115.674671

Choi, S. C., Titov, A. A., Abboud, G., Seay, H. R., Brusko, T. M., Roopenian, D. C., Salek-
Ardakani, S., & Morel, L. (2018). Inhibition of glucose metabolism selectively targets
autoreactive follicular helper T cells. *Nat Commun*, *9*(1), 4369.
https://doi.org/10.1038/s41467-018-06686-0

Codo, A. C., Davanzo, G. G., Monteiro, L. B., de Souza, G. F., Muraro, S. P., Virgilio-da-Silva, J.
V., Prodonoff, J. S., Carregari, V. C., de Biagi Junior, C. A. O., Crunfli, F., Jimenez
Restrepo, J. L., Vendramini, P. H., Reis-de-Oliveira, G., Bispo Dos Santos, K., Toledo-
Teixeira, D. A., Parise, P. L., Martini, M. C., Marques, R. E., Carmo, H. R., . . . Moraes-
Vieira, P. M. (2020). Elevated Glucose Levels Favor SARS-CoV-2 Infection and Monocyte
Response through a HIF-1alpha/Glycolysis-Dependent Axis. *Cell Metab*, *32*(3), 437-446
e435. https://doi.org/10.1016/j.cmet.2020.07.007

Constantinides, M. G., Link, V. M., Tamoutounour, S., Wong, A. C., Perez-Chaparro, P. J., Han,
S. J., Chen, Y. E., Li, K., Farhat, S., Weckel, A., Krishnamurthy, S. R., Vujkovic-Cvijin, I.,
Linehan, J. L., Bouladoux, N., Merrill, E. D., Roy, S., Cua, D. J., Adams, E. J., Bhandoola,
A., . . . Belkaid, Y. (2019). MAIT cells are imprinted by the microbiota in early life and
promote tissue repair. *Science*, *366*(6464). https://doi.org/10.1126/science.aax6624

Corbett, A. J., Awad, W., Wang, H., & Chen, Z. (2020). Antigen Recognition by MR1-Reactive T
Cells; MAIT Cells, Metabolites, and Remaining Mysteries. *Front Immunol*, *11*, 1961.
https://doi.org/10.3389/fimmu.2020.01961

Cosmi, L., De Palma, R., Santarlasci, V., Maggi, L., Capone, M., Frosali, F., Rodolico, G., Querci,
V., Abbate, G., Angeli, R., Berrino, L., Fambrini, M., Caproni, M., Tonelli, F., Lazzeri, E.,
Parronchi, P., Liotta, F., Maggi, E., Romagnani, S., & Annunziato, F. (2008). Human
interleukin 17-producing cells originate from a CD161+CD4+ T cell precursor. *J Exp Med*,
205(8), 1903-1916. https://doi.org/10.1084/jem.20080397

Coulter, E. H., Miller, L., McCorkell, S., McGuire, C., Algie, K., Freeman, J., Weller, B., Mattison,
P. G., McConnachie, A., Wu, O., & Paul, L. (2017). Validity of the activPAL3 activity
monitor in people moderately affected by Multiple Sclerosis. *Med Eng Phys*, *45*, 78-82.
https://doi.org/10.1016/j.medengphy.2017.03.008

Crotty, S. (2014). T follicular helper cell differentiation, function, and roles in disease. *Immunity*,
41(4), 529-542. https://doi.org/10.1016/j.immuni.2014.10.004

Cui, W., Liu, Y., Weinstein, J. S., Craft, J., & Kaech, S. M. (2011). An interleukin-21-interleukin-
10-STAT3 pathway is critical for functional maturation of memory CD8+ T cells. *Immunity*,
35(5), 792-805. https://doi.org/10.1016/j.immuni.2011.09.017

Damasceno, L. E. A., Prado, D. S., Veras, F. P., Fonseca, M. M., Toller-Kawahisa, J. E., Rosa,
M. H., Publio, G. A., Martins, T. V., Ramalho, F. S., Waisman, A., Cunha, F. Q., Cunha,
T. M., & Alves-Filho, J. C. (2020). PKM2 promotes Th17 cell differentiation and
autoimmune inflammation by fine-tuning STAT3 activation. *J Exp Med*, *217*(10).
https://doi.org/10.1084/jem.20190613

Davis, M. R., Zhu, Z., Hansen, D. M., Bai, Q., & Fang, Y. (2015). The role of IL-21 in immunity
and cancer. *Cancer Lett*, *358*(2), 107-114. https://doi.org/10.1016/j.canlet.2014.12.047

Dean, R. H., Polterauer, P., Hunter, C. E., Erath, H., Meng, H. C., & Scott, H. W., Jr. (1979). Effect
of intestinal venous diversion on lipid metabolism. *Surg Forum*, *30*, 85-86.
https://www.ncbi.nlm.nih.gov/pubmed/538706

Duan, Z., Chen, X., Liang, Z., Zeng, Y., Zhu, F., Long, L., McCrae, M. A., Zhuang, H., Shen, T.,
& Lu, F. (2014). Genetic polymorphisms of CXCR5 and CXCL13 are associated with non-
responsiveness to the hepatitis B vaccine. *Vaccine*, *32*(41), 5316-5322.
https://doi.org/10.1016/j.vaccine.2014.07.064

Elsaesser, H., Sauer, K., & Brooks, D. G. (2009). IL-21 is required to control chronic viral infection.
Science, *324*(5934), 1569-1572. https://doi.org/10.1126/science.1174182

Evans-Marin, H., Rogier, R., Koralov, S. B., Manasson, J., Roeleveld, D., van der Kraan, P. M.,
Scher, J. U., Koenders, M. I., & Abdollahi-Roodsaz, S. (2018). Microbiota-Dependent
Involvement of Th17 Cells in Murine Models of Inflammatory Arthritis. *Arthritis Rheumatol*,
70(12), 1971-1983. https://doi.org/10.1002/art.40657

Frohlich, A., Marsland, B. J., Sonderegger, I., Kurrer, M., Hodge, M. R., Harris, N. L., & Kopf, M.
(2007). IL-21 receptor signaling is integral to the development of Th2 effector responses
in vivo. *Blood*, *109*(5), 2023-2031. https://doi.org/10.1182/blood-2006-05-021600

Galvez, J. (2014). Role of Th17 Cells in the Pathogenesis of Human IBD. *ISRN Inflamm*, *2014*,
928461. https://doi.org/10.1155/2014/928461

Garidou, L., Pomie, C., Klopp, P., Waget, A., Charpentier, J., Aloulou, M., Giry, A., Serino, M.,
Stenman, L., Lahtinen, S., Dray, C., Iacovoni, J. S., Courtney, M., Collet, X., Amar, J.,
Servant, F., Lelouvier, B., Valet, P., Eberl, G., . . . Burcelin, R. (2015). The Gut Microbiota
Regulates Intestinal CD4 T Cells Expressing RORgammat and Controls Metabolic
Disease. *Cell Metab*, *22*(1), 100-112. https://doi.org/10.1016/j.cmet.2015.06.001

Gaublomme, J. T., Yosef, N., Lee, Y., Gertner, R. S., Yang, L. V., Wu, C., Pandolfi, P. P., Mak,
T., Satija, R., Shalek, A. K., Kuchroo, V. K., Park, H., & Regev, A. (2015). Single-Cell
Genomics Unveils Critical Regulators of Th17 Cell Pathogenicity. *Cell*, *163*(6), 1400-1412.
https://doi.org/10.1016/j.cell.2015.11.009

Godfrey, D. I., Koay, H. F., McCluskey, J., & Gherardin, N. A. (2019). The biology and functional
importance of MAIT cells. *Nat Immunol*, *20*(9), 1110-1128.
https://doi.org/10.1038/s41590-019-0444-8

Granelli-Piperno, A., Golebiowska, A., Trumpfheller, C., Siegal, F. P., & Steinman, R. M. (2004).
HIV-1-infected monocyte-derived dendritic cells do not undergo maturation but can elicit
IL-10 production and T cell regulation. *Proc Natl Acad Sci U S A*, *101*(20), 7669-7674.
https://doi.org/10.1073/pnas.0402431101

Gu, L., Yi, Z., Zhang, Y., Ma, Z., Zhu, Y., & Gao, J. (2017). Low dose of 2-deoxy-D-glucose kills
acute lymphoblastic leukemia cells and reverses glucocorticoid resistance via N-linked
glycosylation inhibition under normoxia. *Oncotarget*, *8*(19), 30978-30991.
https://doi.org/10.18632/oncotarget.16046

Hauser, M. A., Kindinger, I., Laufer, J. M., Spate, A. K., Bucher, D., Vanes, S. L., Krueger, W. A.,
Wittmann, V., & Legler, D. F. (2016). Distinct CCR7 glycosylation pattern shapes receptor
signaling and endocytosis to modulate chemotactic responses. *J Leukoc Biol*, *99*(6), 993-
1007. https://doi.org/10.1189/jlb.2VMA0915-432RR

Hinks, T. S. C., & Zhang, X. W. (2020). MAIT Cell Activation and Functions. *Front Immunol*, *11*,
1014. https://doi.org/10.3389/fimmu.2020.01014

Ho, P. C., Bihuniak, J. D., Macintyre, A. N., Staron, M., Liu, X., Amezquita, R., Tsui, Y. C., Cui,
G., Micevic, G., Perales, J. C., Kleinstein, S. H., Abel, E. D., Insogna, K. L., Feske, S.,
Locasale, J. W., Bosenberg, M. W., Rathmell, J. C., & Kaech, S. M. (2015).

Phosphoenolpyruvate Is a Metabolic Checkpoint of Anti-tumor T Cell Responses. *Cell*, *162*(6), 1217-1228. https://doi.org/10.1016/j.cell.2015.08.012

Imperiali, B., & O'Connor, S. E. (1999). Effect of N-linked glycosylation on glycopeptide and glycoprotein structure. *Curr Opin Chem Biol*, *3*(6), 643-649. https://www.ncbi.nlm.nih.gov/pubmed/10600722

Ioannidis, M., Cerundolo, V., & Salio, M. (2020). The Immune Modulating Properties of Mucosal-Associated Invariant T Cells. *Front Immunol*, *11*, 1556. https://doi.org/10.3389/fimmu.2020.01556

Iyer, S. S., & Cheng, G. (2012). Role of interleukin 10 transcriptional regulation in inflammation and autoimmune disease. *Crit Rev Immunol*, *32*(1), 23-63. https://doi.org/10.1615/critrevimmunol.v32.i1.30

Jacobs, S. R., Herman, C. E., Maciver, N. J., Wofford, J. A., Wieman, H. L., Hammen, J. J., & Rathmell, J. C. (2008). Glucose uptake is limiting in T cell activation and requires CD28-mediated Akt-dependent and independent pathways. *J Immunol*, *180*(7), 4476-4486. https://doi.org/10.4049/jimmunol.180.7.4476

Jia, X., Zhai, T., Wang, B., Yao, Q., Li, Q., Mu, K., & Zhang, J. A. (2019). Decreased number and impaired function of type 1 regulatory T cells in autoimmune diseases. *J Cell Physiol*, *234*(8), 12442-12450. https://doi.org/10.1002/jcp.28092

Johnson, M. O., Wolf, M. M., Madden, M. Z., Andrejeva, G., Sugiura, A., Contreras, D. C., Maseda, D., Liberti, M. V., Paz, K., Kishton, R. J., Johnson, M. E., de Cubas, A. A., Wu, P., Li, G., Zhang, Y., Newcomb, D. C., Wells, A. D., Restifo, N. P., Rathmell, W. K., . . . Rathmell, J. C. (2018). Distinct Regulation of Th17 and Th1 Cell Differentiation by Glutaminase-Dependent Metabolism. *Cell*, *175*(7), 1780-1795 e1719. https://doi.org/10.1016/j.cell.2018.10.001

Jung, M. K., Kwak, J. E., & Shin, E. C. (2017). IL-17A-Producing Foxp3(+) Regulatory T Cells and Human Diseases. *Immune Netw*, *17*(5), 276-286. https://doi.org/10.4110/in.2017.17.5.276

Kastirr, I., Crosti, M., Maglie, S., Paroni, M., Steckel, B., Moro, M., Pagani, M., Abrignani, S., & Geginat, J. (2015). Signal Strength and Metabolic Requirements Control Cytokine-

Induced Th17 Differentiation of Uncommitted Human T Cells. *J Immunol*, *195*(8), 3617-3627. https://doi.org/10.4049/jimmunol.1501016

Kleinschek, M. A., Boniface, K., Sadekova, S., Grein, J., Murphy, E. E., Turner, S. P., Raskin, L., Desai, B., Faubion, W. A., de Waal Malefyt, R., Pierce, R. H., McClanahan, T., & Kastelein, R. A. (2009). Circulating and gut-resident human Th17 cells express CD161 and promote intestinal inflammation. *J Exp Med*, *206*(3), 525-534. https://doi.org/10.1084/jem.20081712

Knochelmann, H. M., Dwyer, C. J., Bailey, S. R., Amaya, S. M., Elston, D. M., Mazza-McCrann, J. M., & Paulos, C. M. (2018). When worlds collide: Th17 and Treg cells in cancer and autoimmunity. *Cell Mol Immunol*, *15*(5), 458-469. https://doi.org/10.1038/s41423-018-0004-4

Kurtoglu, M., Gao, N., Shang, J., Maher, J. C., Lehrman, M. A., Wangpaichitr, M., Savaraj, N., Lane, A. N., & Lampidis, T. J. (2007). Under normoxia, 2-deoxy-D-glucose elicits cell death in select tumor types not by inhibition of glycolysis but by interfering with N-linked glycosylation. *Mol Cancer Ther*, *6*(11), 3049-3058. https://doi.org/10.1158/1535-7163.MCT-07-0310

Lajoie, S., Lewkowich, I., Herman, N. S., Sproles, A., Pesce, J. T., Wynn, T. A., Grusby, M. J., Hamid, Q., & Wills-Karp, M. (2014). IL-21 receptor signalling partially mediates Th2-mediated allergic airway responses. *Clin Exp Allergy*, *44*(7), 976-985. https://doi.org/10.1111/cea.12341

Lee, Y., Awasthi, A., Yosef, N., Quintana, F. J., Xiao, S., Peters, A., Wu, C., Kleinewietfeld, M., Kunder, S., Hafler, D. A., Sobel, R. A., Regev, A., & Kuchroo, V. K. (2012). Induction and molecular signature of pathogenic TH17 cells. *Nat Immunol*, *13*(10), 991-999. https://doi.org/10.1038/ni.2416

Leonard, W. J., & Wan, C. K. (2016). IL-21 Signaling in Immunity. *F1000Res*, *5*. https://doi.org/10.12688/f1000research.7634.1

Li, W., Qu, G., Choi, S. C., Cornaby, C., Titov, A., Kanda, N., Teng, X., Wang, H., & Morel, L. (2019). Targeting T Cell Activation and Lupus Autoimmune Phenotypes by Inhibiting Glucose Transporters. *Front Immunol*, *10*, 833. https://doi.org/10.3389/fimmu.2019.00833

790 Lichtenegger, F. S., Rothe, M., Schnorfeil, F. M., Deiser, K., Krupka, C., Augsberger, C., Schluter,
791   M., Neitz, J., & Subklewe, M. (2018). Targeting LAG-3 and PD-1 to Enhance T Cell
792   Activation by Antigen-Presenting Cells. *Front Immunol*, *9*, 385.
793   https://doi.org/10.3389/fimmu.2018.00385

794 MacIver, N. J., Michalek, R. D., & Rathmell, J. C. (2013). Metabolic regulation of T lymphocytes.
795   *Annu Rev Immunol*, *31*, 259-283. https://doi.org/10.1146/annurev-immunol-032712-
796   095956

797 Marshall, E. A., Ng, K. W., Kung, S. H., Conway, E. M., Martinez, V. D., Halvorsen, E. C.,
798   Rowbotham, D. A., Vucic, E. A., Plumb, A. W., Becker-Santos, D. D., Enfield, K. S.,
799   Kennett, J. Y., Bennewith, K. L., Lockwood, W. W., Lam, S., English, J. C., Abraham, N.,
800   & Lam, W. L. (2016). Emerging roles of T helper 17 and regulatory T cells in lung cancer
801   progression and metastasis. *Mol Cancer*, *15*(1), 67. https://doi.org/10.1186/s12943-016-
802   0551-1

803 McDonald, D. R. (2012). TH17 deficiency in human disease. *J Allergy Clin Immunol*, *129*(6), 1429-
804   1435; quiz 1436-1427. https://doi.org/10.1016/j.jaci.2012.03.034

805 Medrano, R. F. V., Hunger, A., Mendonca, S. A., Barbuto, J. A. M., & Strauss, B. E. (2017).
806   Immunomodulatory and antitumor effects of type I interferons and their application in
807   cancer therapy. *Oncotarget*, *8*(41), 71249-71284.
808   https://doi.org/10.18632/oncotarget.19531

809 Mercer, F., Khaitan, A., Kozhaya, L., Aberg, J. A., & Unutmaz, D. (2014). Differentiation of IL-17-
810   producing effector and regulatory human T cells from lineage-committed naive precursors.
811   *J Immunol*, *193*(3), 1047-1054. https://doi.org/10.4049/jimmunol.1302936

812 Ng, M. S. F., Roth, T. L., Mendoza, V. F., Marson, A., & Burt, T. D. (2019). Helios enhances the
813   preferential differentiation of human fetal CD4(+) naive T cells into regulatory T cells. *Sci*
814   *Immunol*, *4*(41). https://doi.org/10.1126/sciimmunol.aav5947

815 Nicholas, D. A., Proctor, E. A., Agrawal, M., Belkina, A. C., Van Nostrand, S. C., Panneerseelan-
816   Bharath, L., Jones, A. R. t., Raval, F., Ip, B. C., Zhu, M., Cacicedo, J. M., Habib, C., Sainz-
817   Rueda, N., Persky, L., Sullivan, P. G., Corkey, B. E., Apovian, C. M., Kern, P. A.,
818   Lauffenburger, D. A., & Nikolajczyk, B. S. (2019). Fatty Acid Metabolites Combine with

Reduced beta Oxidation to Activate Th17 Inflammation in Human Type 2 Diabetes. *Cell Metab*, *30*(3), 447-461 e445. https://doi.org/10.1016/j.cmet.2019.07.004

Nurieva, R., Yang, X. O., Martinez, G., Zhang, Y., Panopoulos, A. D., Ma, L., Schluns, K., Tian, Q., Watowich, S. S., Jetten, A. M., & Dong, C. (2007). Essential autocrine regulation by IL-21 in the generation of inflammatory T cells. *Nature*, *448*(7152), 480-483. https://doi.org/10.1038/nature05969

Oh, J., & Unutmaz, D. (2019). Immune cells for microbiota surveillance. *Science*, *366*(6464), 419-420. https://doi.org/10.1126/science.aaz4014

Okano, T., Saegusa, J., Nishimura, K., Takahashi, S., Sendo, S., Ueda, Y., & Morinobu, A. (2017). 3-bromopyruvate ameliorate autoimmune arthritis by modulating Th17/Treg cell differentiation and suppressing dendritic cell activation. *Sci Rep*, *7*, 42412. https://doi.org/10.1038/srep42412

Palmer, C. S., Ostrowski, M., Balderson, B., Christian, N., & Crowe, S. M. (2015). Glucose metabolism regulates T cell activation, differentiation, and functions. *Front Immunol*, *6*, 1. https://doi.org/10.3389/fimmu.2015.00001

Passalacqua, K. D., Lu, J., Goodfellow, I., Kolawole, A. O., Arche, J. R., Maddox, R. J., Carnahan, K. E., O'Riordan, M. X. D., & Wobus, C. E. (2019). Glycolysis Is an Intrinsic Factor for Optimal Replication of a Norovirus. *mBio*, *10*(2). https://doi.org/10.1128/mBio.02175-18

Pearce, E. L., Poffenberger, M. C., Chang, C. H., & Jones, R. G. (2013). Fueling immunity: insights into metabolism and lymphocyte function. *Science*, *342*(6155), 1242454. https://doi.org/10.1126/science.1242454

Perl, A. (2016). Activation of mTOR (mechanistic target of rapamycin) in rheumatic diseases. *Nat Rev Rheumatol*, *12*(3), 169-182. https://doi.org/10.1038/nrrheum.2015.172

Platt, M. P., Bolding, K. A., Wayne, C. R., Chaudhry, S., Cutforth, T., Franks, K. M., & Agalliu, D. (2020). Th17 lymphocytes drive vascular and neuronal deficits in a mouse model of postinfectious autoimmune encephalitis. *Proc Natl Acad Sci U S A*, *117*(12), 6708-6716. https://doi.org/10.1073/pnas.1911097117

Raez, L. E., Papadopoulos, K., Ricart, A. D., Chiorean, E. G., Dipaola, R. S., Stein, M. N., Rocha
Lima, C. M., Schlesselman, J. J., Tolba, K., Langmuir, V. K., Kroll, S., Jung, D. T., Kurtoglu,
M., Rosenblatt, J., & Lampidis, T. J. (2013). A phase I dose-escalation trial of 2-deoxy-D-
glucose alone or combined with docetaxel in patients with advanced solid tumors. *Cancer*
Chemother Pharmacol, *71*(2), 523-530. https://doi.org/10.1007/s00280-012-2045-1

Sahu, K. K., & Kumar, R. (2021). Role of 2-Deoxy-D-Glucose (2-DG) in COVID-19 disease: A
potential   game-changer.   *J   Family   Med   Prim   Care*,   *10*(10),   3548-3552.
https://doi.org/10.4103/jfmpc.jfmpc_1338_21

Santegoets, S. J., Turksma, A. W., Powell, D. J., Jr., Hooijberg, E., & de Gruijl, T. D. (2013). IL-
21 in cancer immunotherapy: At the right place at the right time. *Oncoimmunology*, *2*(6),
e24522. https://doi.org/10.4161/onci.24522

Sasaki, C. Y., Chen, G., Munk, R., Eitan, E., Martindale, J., Longo, D. L., & Ghosh, P. (2016).
p((7)(0)S(6)K(1)) in the TORC1 pathway is essential for the differentiation of Th17 Cells,
but not Th1, Th2, or Treg cells in mice. *Eur J Immunol*, *46*(1), 212-222.
https://doi.org/10.1002/eji.201445422

Sena, L. A., Li, S., Jairaman, A., Prakriya, M., Ezponda, T., Hildeman, D. A., Wang, C. R.,
Schumacker, P. T., Licht, J. D., Perlman, H., Bryce, P. J., & Chandel, N. S. (2013).
Mitochondria are required for antigen-specific T cell activation through reactive oxygen
species   signaling.   *Immunity*,   *38*(2),   225-236.
https://doi.org/10.1016/j.immuni.2012.10.020

Serriari, N. E., Eoche, M., Lamotte, L., Lion, J., Fumery, M., Marcelo, P., Chatelain, D., Barre, A.,
Nguyen-Khac, E., Lantz, O., Dupas, J. L., & Treiner, E. (2014). Innate mucosal-associated
invariant T (MAIT) cells are activated in inflammatory bowel diseases. *Clin Exp Immunol*,
176(2), 266-274. https://doi.org/10.1111/cei.12277

Shen, H., & Shi, L. Z. (2019). Metabolic regulation of TH17 cells. *Mol Immunol*, *109*, 81-87.
https://doi.org/10.1016/j.molimm.2019.03.005

Shi, L. Z., Wang, R., Huang, G., Vogel, P., Neale, G., Green, D. R., & Chi, H. (2011). HIF1alpha-
dependent glycolytic pathway orchestrates a metabolic checkpoint for the differentiation
of   TH17   and   Treg   cells.   *J   Exp   Med*,   *208*(7),   1367-1376.
https://doi.org/10.1084/jem.20110278

Singh, S. P., Zhang, H. H., Foley, J. F., Hedrick, M. N., & Farber, J. M. (2008). Human T cells that
877   are able to produce IL-17 express the chemokine receptor CCR6. *J Immunol*, *180*(1), 214-
878   221. https://doi.org/10.4049/jimmunol.180.1.214

879 Spolski, R., Kim, H. P., Zhu, W., Levy, D. E., & Leonard, W. J. (2009). IL-21 mediates suppressive
880   effects via its induction of IL-10. *J Immunol*, *182*(5), 2859-2867.
881   https://doi.org/10.4049/jimmunol.0802978

882 Spolski, R., & Leonard, W. J. (2010). IL-21 and T follicular helper cells. *Int Immunol*, *22*(1), 7-12.
883   https://doi.org/10.1093/intimm/dxp112

884 Stein, M., Lin, H., Jeyamohan, C., Dvorzhinski, D., Gounder, M., Bray, K., Eddy, S., Goodin, S.,
885   White, E., & Dipaola, R. S. (2010). Targeting tumor metabolism with 2-deoxyglucose in
886   patients with castrate-resistant prostate cancer and advanced malignancies. *Prostate*,
887   *70*(13), 1388-1394. https://doi.org/10.1002/pros.21172

888 Tanimine, N., Germana, S. K., Fan, M., Hippen, K., Blazar, B. R., Markmann, J. F., Turka, L. A.,
889   & Priyadharshini, B. (2019). Differential effects of 2-deoxy-D-glucose on in vitro expanded
890   human regulatory T cell subsets. *PLoS One*, *14*(6), e0217761.
891   https://doi.org/10.1371/journal.pone.0217761

892 Tastan, C., Karhan, E., Zhou, W., Fleming, E., Voigt, A. Y., Yao, X., Wang, L., Horne, M., Placek,
893   L., Kozhaya, L., Oh, J., & Unutmaz, D. (2018). Tuning of human MAIT cell activation by
894   commensal bacteria species and MR1-dependent T-cell presentation. *Mucosal Immunol*,
895   *11*(6), 1591-1605. https://doi.org/10.1038/s41385-018-0072-x

896 Thornton, A. M., Lu, J., Korty, P. E., Kim, Y. C., Martens, C., Sun, P. D., & Shevach, E. M. (2019).
897   Helios(+) and Helios(-) Treg subpopulations are phenotypically and functionally distinct
898   and express dissimilar TCR repertoires. *Eur J Immunol*, *49*(3), 398-412.
899   https://doi.org/10.1002/eji.201847935

900 Valmori, D., Raffin, C., Raimbaud, I., & Ayyoub, M. (2010). Human RORgammat+ TH17 cells
901   preferentially differentiate from naive FOXP3+Treg in the presence of lineage-specific
902   polarizing factors. *Proc Natl Acad Sci U S A*, *107*(45), 19402-19407.
903   https://doi.org/10.1073/pnas.1008247107

van der Windt, G. J., & Pearce, E. L. (2012). Metabolic switching and fuel choice during T-cell
differentiation and memory development. *Immunol Rev*, *249*(1), 27-42.
https://doi.org/10.1111/j.1600-065X.2012.01150.x

Visconti, A., Le Roy, C. I., Rosa, F., Rossi, N., Martin, T. C., Mohney, R. P., Li, W., de Rinaldis,
E., Bell, J. T., Venter, J. C., Nelson, K. E., Spector, T. D., & Falchi, M. (2019). Interplay
between the human gut microbiome and host metabolism. *Nat Commun*, *10*(1), 4505.
https://doi.org/10.1038/s41467-019-12476-z

Vogelzang, A., McGuire, H. M., Yu, D., Sprent, J., Mackay, C. R., & King, C. (2008). A
fundamental role for interleukin-21 in the generation of T follicular helper cells. *Immunity*,
29(1), 127-137. https://doi.org/10.1016/j.immuni.2008.06.001

Wan, Q., Kozhaya, L., ElHed, A., Ramesh, R., Carlson, T. J., Djuretic, I. M., Sundrud, M. S., &
Unutmaz, D. (2011). Cytokine signals through PI-3 kinase pathway modulate Th17
cytokine production by CCR6+ human memory T cells. *J Exp Med*, *208*(9), 1875-1887.
https://doi.org/10.1084/jem.20102516

Willing, A., Jager, J., Reinhardt, S., Kursawe, N., & Friese, M. A. (2018). Production of IL-17 by
MAIT Cells Is Increased in Multiple Sclerosis and Is Associated with IL-7 Receptor
Expression. *J Immunol*, *200*(3), 974-982. https://doi.org/10.4049/jimmunol.1701213

Wu, X., Tian, J., & Wang, S. (2018). Insight Into Non-Pathogenic Th17 Cells in Autoimmune
Diseases. *Front Immunol*, *9*, 1112. https://doi.org/10.3389/fimmu.2018.01112

Xi, H., Kurtoglu, M., & Lampidis, T. J. (2014). The wonders of 2-deoxy-D-glucose. *IUBMB Life*,
66(2), 110-121. https://doi.org/10.1002/iub.1251

Xu, H., Agalioti, T., Zhao, J., Steglich, B., Wahib, R., Vesely, M. C. A., Bielecki, P., Bailis, W.,
Jackson, R., Perez, D., Izbicki, J., Licona-Limon, P., Kaartinen, V., Geginat, J., Esplugues,
E., Tolosa, E., Huber, S., Flavell, R. A., & Gagliani, N. (2020). The induction and function
of the anti-inflammatory fate of TH17 cells. *Nat Commun*, *11*(1), 3334.
https://doi.org/10.1038/s41467-020-17097-5

Yang, C., Zuo, Q., Liu, X., Zhao, Q., Pu, H., Gao, L., Zhao, L., Guo, Z., Lin, Y., Liu, J., Bi, J., &
Yin, G. (2021). Small molecule screening identified cepharanthine as an inhibitor of
porcine reproductive and respiratory syndrome virus infection in vitro by suppressing

933   integrins/ILK/RACK1/PKCalpha/NF-kappaB signalling axis. *Vet Microbiol*, *255*, 109016.

934    https://doi.org/10.1016/j.vetmic.2021.109016

935 Yasuda, K., Kitagawa, Y., Kawakami, R., Isaka, Y., Watanabe, H., Kondoh, G., Kohwi-

936    Shigematsu, T., Sakaguchi, S., & Hirota, K. (2019). Satb1 regulates the effector program

937    of encephalitogenic tissue Th17 cells in chronic inflammation. *Nat Commun*, *10*(1), 549.

938    https://doi.org/10.1038/s41467-019-08404-w

939 Zeng, H., Zhang, R., Jin, B., & Chen, L. (2015). Type 1 regulatory T cells: a new mechanism of

940    peripheral immune tolerance. *Cell Mol Immunol*, *12*(5), 566-571.

941    https://doi.org/10.1038/cmi.2015.44

942 Zhou, L., Ivanov, II, Spolski, R., Min, R., Shenderov, K., Egawa, T., Levy, D. E., Leonard, W. J.,

943    & Littman, D. R. (2007). IL-6 programs T(H)-17 cell differentiation by promoting sequential

944    engagement of the IL-21 and IL-23 pathways. *Nat Immunol*, *8*(9), 967-974.

945    https://doi.org/10.1038/ni1488

Figure 1

Figure 2

Figure 3

A

B

C

D

Figure 4

Figure 5

Figure 6

A

	Control	3mM 2DG	+1mM mannose	+3mM mannose	3mM mannose

B

GLUT1 (Fold)

Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose
+3mM 2DG

CD25 (Fold)

Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose
+3mM 2DG

C Naive T Cell Expansion after 2 weeks culture

Control	3mM mannose	+ 3mM 2DG			
		None	1mM mannose	3mM mannose	3mM glucose
157±5	100±5	1	30±5	50±5	16±5

D

ECAR(mpH/min)

— Control
— 3mM mannose
— 3mM 2DG
— 3mM 2DG+3mM glucose
— 3mM 2DG+3mM mannose

E

ECAR(mpH/min)

Control, 3mM mannose, + 3mM 2DG (None, 3mM mannose, 3mM glucose)

F

	Control	3mM 2DG	+1mM Mannose	+3mM Mannose	3mM Mannose

CCR6 / CD161
IL-17 / CD161

G

CD161 Expression — Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose, +3mM 2DG

IFNγ — Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose, +3mM 2DG

IL-17 — Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose, +3mM 2DG

IL-21 — Control, 3mM mannose, 3mM 2DG, 1mM mannose, 3mM mannose, 3mM glucose, +3mM 2DG

Supplemental Figure 1

A — Day 1 post activation: Resting, None, 3mM 2DG, No glucose

B — Day 4 post activation: None, 3mM 2DG, No glucose

(Naive / Memory rows; axes GLUT1 vs CD25)

C — Bar graphs: GLUT1 (Fold) Naive; GLUT1 (old) Memory; CD25 (Fold) Naive; CD25 (Fold) Memory. Conditions: Control, 2DG, No glucose. Legend: Day 1 (black), Day 4 (gray). *** significance markers.

D — CTV histograms: Day 0, Day 4, Day 6. Rows: Naive, Memory. Legend: Control (gray), 2DG(3mM) (red), No glucose (blue). X-axis: CTV.

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5