Chromatin structure influences rate and spectrum of spontaneous mutations in *Neurospora crassa*

Mariana Villalba de la Peña[1], Pauliina A. M. Summanen[1],

Martta Liukkonen[1], Ilkka Kronholm[1]

[1]Department of Biological and Environmental Science, University of Jyväskylä, FI-40014 Jyväskylä, Finland

Running Head: Influence of chromatin on mutation rate

Keywords: fungi, mutation accumulation, genomics, epigenetics, H3K9 trimethylation, H3K27 trimethylation

Corresponding author:

Ilkka Kronholm

Department of Biological and Environmental Science,

University of Jyväskylä,

P.O. Box 35, FI-40014

Jyväskylä,

Finland

Fax +358 14 617 239

Email: ilkka.kronholm@jyu.fi

Abstract

Mutation rates in different species have been extensively studied. However, much less is known about the variation in mutation rate throughout the genome. Understanding patterns of mutation rate variation would reveal which evolutionary processes shape patterns of genetic variation. Chromatin modifications have been suggested to be an important factor in determining mutation rate variation in eukaryotic genomes. To study mutation rate variation we performed a mutation accumulation experiment in the filamentous fungus *Neurospora crassa*, and sequenced the genomes of the 40 MA lines that had been propagated asexually for approximately 1015 [1003, 1026] mitoses. We detected 1322 mutations in total. We observed that mutation rate was higher in regions of low GC, in domains of H3K9 trimethylation, in centromeric regions, and in domains of H3K27 trimethylation. The rate of single nucleotide mutations in euchromatin was $2.46\,[2.19, 2.77] \times 10^{-10}$. In contrast, mutation rate in H3K9me3 domains was tenfold higher: $2.43\,[2.25, 2.62] \times 10^{-9}$. We also observed that the spectrum of single nucleotide mutations was different in H3K9me3 and euchromatic domains. Mutation spectrum was also markedly different than the one previously reported for sexual reproduction. Our statistical model of mutation rate variation predicted a moderate amount of extant genetic variation. Suggesting that mutation rate is an important factor in determining levels of natural genetic variation. Furthermore, we characterized mutation rates of structural variants, complex mutations, and the effect of local sequence context on mutation rate. Our study highlights that chromatin modifications are associated with mutation rate and accurate evolutionary inferences should take variation in mutation rate across the genome into account.

Introduction

New mutations are the source of all genetic diversity, and evolutionary change ultimately depends on the input of new mutations into the population. However, organisms also pay a substantial cost for their ability to evolve, as deleterious mutations are more common than beneficial mutations (Eyre-Walker and Keightley, 2007), and mutations can lead to adverse outcomes, such as decline in fitness, hereditary diseases, and cancer. Therefore, knowledge of the rates and spectrum of spontaneous mutations are fundamental to our understanding of evolution and certain aspects of medicine.

Spontaneous mutations are rare events and used to be difficult to study. However, new sequencing technologies have made it possible to capture a large number of spontaneous mutations for analysis (Katju and Bergthorsson, 2018). Several studies have now estimated mutation rates by directly observing mutations by sequencing mutation accumulation lines or parent-offspring trios (Ossowski et al., 2010; Ness et al., 2012; Keightley et al., 2014; Zhu et al., 2014; Sung et al., 2015; Ness et al., 2015; Keightley et al., 2015; Keith et al., 2016; Wang et al., 2020). These studies have estimated the rate and spectrum of spontaneous mutations to high precision.

The process of mutation is stochastic, but not all mutations are equally likely. While this has been appreciated for a long time for certain classes of mutations, such as transitions and transversions, there is also variation in mutation rate that seems to depend on the structural features of the genome, such as the organization of chromatin (Makova and Hardison, 2015). In particular, nucleosome positioning (Tolstorukov et al., 2011; Chen et al., 2012; Li and Luscombe, 2020) and chromatin structure have a strong influence on mutation rates (Schuster-Böckler and Lehner, 2012; Polak et al., 2015; Weng et al., 2019; Monroe et al., 2022). Chromatin structure is associated with chemical modifications of histone H3. In particular, the methylation status of certain lysine residues, such as methylation of H3K9 and H3K27, is associated with closed and silenced chromatin, called heterochromatin, while methylation of H3K36 is associated with open actively transcribed chromatin, called euchromatin (Kouzarides, 2007). Heterochromatin appears to have higher mutation rates than euchromatin (Makova and Hardison, 2015). In addition, local sequence

context, such as GC-content, also has a strong effect on mutation rate (Makova and Hardison, 2015; Ness et al., 2015; Sung et al., 2015). Although we know that chromatin structure can shape mutation rates, most data come from humans and a few model species.

Furthermore, to what extent variation in mutation rate determines patterns of observed genetic diversity along with other evolutionary mechanisms is understood mostly from population genetic data, rather than from direct observations of mutation, with few exceptions e.g. Monroe et al. (2022). There are some tests for selection, such as dN/dS ratios, which are not affected by mutation rate. However, tests based on site frequency spectrum to infer mutational effects or demography can benefit if data about mutation rate is available (Keightley and Eyre-Walker, 2007), especially if the goal is the examine different categories of genes or regions of the genome. Furthermore, to understand how different evolutionary forces, such as background selection, gene conversion, demographic processes, and adaptive evolution jointly shape patterns of diversity across the genome, obtaining estimates of mutation rate will allow us to parameterize population genetic models (Campos et al., 2017; Castellano et al., 2020; Johri et al., 2020). Ultimately taking all possible effects into account will allow us to make better estimates of how natural selection shapes genetic diversity.

To examine patterns of mutation rate variation we performed a mutation accumulation (MA) experiment (Halligan and Keightley, 2009) in the filamentous fungus *Neurospora crassa*. In MA experiments an ancestor is split into multiple lines, and these lines are bottlenecked every generation which minimizes the efficiency of natural selection. (Figure 1A). This way even deleterious mutations can accumulate in these lines (Halligan and Keightley, 2009). We sequenced the genomes of these mutation accumulation lines using short read sequencing. *Neurospora crassa* is a filamentous fungus with a facultative sexual cycle, and it can produce both asexual and sexual spores. It has a small genome of 42 Mb, and the vegetative mycelium is haploid. Furthermore, *N. crassa* has a genome defence mechanism, called repeat-induced point mutation (RIP), that detects duplicated regions of the genome in premeiotic cells, and induces C $\rightarrow$ T transitions in the duplicated sequences (Selker, 1990). Because RIP does not induce the same specific mutations in both copies, large repeated arrays are seldom perfect in *N. crassa* as the duplicated sequences divergence from

each other due to the action of RIP. As the sequences diverge, efficiency of RIP decreases (Cambareri et al., 1991). Existence of imperfect repeats makes it possible to use short read sequencing to sequence repetitive regions, for example centromeric regions (Smith et al., 2011), where read mapping is normally difficult in plants and animals.

A previous study in *N. crassa* investigated mutation rate during sexual reproduction and revealed that mutation rate is especially high in regions of the genome targeted by RIP (Wang et al., 2020). However, only a small number of mutations was collected during asexual reproduction; not enough to infer variation in mutation rate across the genome. Our study complements Wang et al. (2020) by allowing us to characterize the determinants of mutation rate and spectrum during asexual reproduction, when RIP is not active. We used information on chromatin structure of *N. crassa* to model variation in mutation rate across the genome. We also resequenced strains obtained from natural populations, and compared the predictions of our mutation model to patterns of natural genetic variation to assess whether natural genetic variation reflects observed mutation rate variation. Furthermore, we examined the effect of local sequence context on mutation rate, and characterize patterns of structural variants and complex mutations.

Results

We initiated the MA experiment with two ancestors that were isogenic, except for the mating type locus. We split both ancestors into 20 lines, giving 40 MA lines in total. We then plated asexual spores (conidia) on plates, picked a single colony, and transferred this colony into a test tube and let the mycelium grow and make asexual spores. Then we plated these spores again to isolate a single spore, this process was repeated for 40 transfers (Figure 1B).

Number of mitoses in the experiment

Neurospora crassa is a filamentous organism, and it does not have a defined germline. All parts of the mycelium are capable of producing structures that produce asexual spores. Thus, the num-

Figure 1: A) In the mutation accumulation experiment ancestors were split into multiple lines, which were propagated via single spore descent. B) Lines in the MA experiment were transferred by always picking a colony originating from a single spore from a plate, moving this to a test tube with Vogel's Medium to allow sporulation, then diluting spores and spreading them to a sorbose plate. C) Micrograph of *N. crassa* mycelium, showing nuclei fluorescing green and cell walls in red. D) Distribution of mutations in the MA-lines.

ber of transfers during the MA experiment does not correspond to a generation in a natural way. Therefore, a reasonable unit for measuring mutation rate is the number of mutations per mitosis.

We estimated the number of mitoses the MA-lines went through based on counting nuclei in different phases of a transfer (Figure 1B, C). Based on our estimate, the MA lines went through 25 [25, 26] (median, [95% HPDI]) mitoses in a single transfer. For the whole experiment of 40 transfers, this means that the MA lines went through 1015 [1003, 1026] mitoses.

Mutations in the MA-lines

To detect mutations in the MA lines, we sequenced the genomes of the MA ancestors and the MA lines using short read sequencing with 150 bp paired end reads. Sequencing depth was more than 30x on average, and approximately 98% of reads were mapped to the reference genome (Table S1). Reference genome of *N. crassa* contains 41 108 926 bp and we called 98.7% of those bases on average. We used a pipeline based on the GATK best practises to call the mutations, followed by manual inspection of alignments.

After sequencing the MA lines, it became apparent that one of the lines had many of the same mutations as another line, likely due to a mislabeling or contamination at some part of the MA experiment. This line was excluded from the analysis, leaving 39 MA-lines in the data.

Accurate mutation calls are crucial for estimating mutation rates. Majority of our mutation calls had a maximum genotype quality score of 99 and were unambiguous (Figure S1), we also used simulations to check that were able to call mutations accurately in all regions of the genome (see supplementary results). We further verified some of the mutations by Sanger-sequencing. We focused verification efforts on mutation calls that were of lower quality based on visual inspection of alignments. Because if those calls were correct, then mutations with higher quality scores would be correct as well. For point mutations we selected 29 mutations to be confirmed. PCR or sequencing failed in 6 out 29 point mutations, and the remaining 23 point mutations were all confirmed. For the 37 small indels, PCR or sequencing failed from 10, while 20 were confirmed, and 7 were false positives. Of the 16 SVs tested, PCR or sequencing failed for 9, 5 were confirmed and 2

were false positives. Since these mutations were a sample of the mutations with the worst genotype qualities, our genotyping for point mutations was very accurate (see suplementary results), while there is some uncertainty for small indels and SVs. All mutations that were false positives were excluded from the data.

In total we observed 1322 mutations, with a median of 33 mutations per MA line. One of the MA lines had an excess amount of mutations (Figure 1D), and it is possible that a mutation happened in this line that increased mutation rate. There was a trend of increased C:G $\rightarrow$ A:T transversions in this line, but the rate was not statistically different from the rest of the MA lines (Figure S2). We did not observe any obvious candidate mutation that affected a DNA repair gene.

The breakdown of different mutation types among the MA lines was 1077 single nucleotide mutations, 134 insertions, 97 deletions, 9 complex mutations where a single mutational event created multiple adjacent nucleotide changes, and 5 translocations. Total mutation rate during asexual propagation was 0.03 [0.03, 0.04] mutations / genome / mitosis.

Mutation rate variation across the genome

To examine did chromatin structure influence mutation rates, we assembled publicly available data for H3K9 trimethylation (H3K9me3), H3K27 trimethylation (H3K27me3), H3K36 dimethylation, and location of *N. crassa* centromeres (Jamieson et al., 2013; Bicocca et al., 2018; Smith et al., 2011). We also examined regions of the genome containing ancestral duplicates defined by Wang et al. (2020), but the duplicated regions were almost perfectly correlated with H3K9me3 domains (Figure 2A), so we did not use them in further analysis. Furthermore, H3K9me3 and H3K36me2 domains were almost perfect mirror images of each other (Figure 2A), so H3K36me2 domains were excluded as they contain the same information as the absence of H3K9me3.

Next, we examined the distribution of mutations across the seven *N. crassa* chromosomes. We observed that mutations were not uniformly distributed along the chromosomes, but were concentrated in centromeric regions and regions of the genome marked by H3K9me3 (Figure 2A). Examining relative mutation rates confirmed that mutation rate was over six fold higher in centromeric

and H3K9me3 domains (Figure 2B), while the effect of H3K27me3 domains was much smaller: mutation rate in H3K27me3 domains relative to euchromatin was only 1.4 [1.1, 1.78] times higher.

We also observed that GC-content displayed a bimodal distribution and was lower in H3K9me3 and centromeric domains, with little overlap with distribution of GC-content in euchromatin (Figure 2C). Does the higher mutation rate in H3K9me3 domains arise from the lower GC-content itself, some other factor related to the chromatin modifications, or a combination of both? To exclude a case where a trend appears when comparing subgroups but disappears within subgroups, we examined the effect of GC-content on mutation rate within the different domains, and observed that lower GC-content is associated with higher mutation rates within each domain, at different ranges of GC-content (Figure S3). Although the pattern was unclear in H3K27me3 domains, as few mutations were observed in H3K27me3 regions that had low GC-content.

To investigate the joint effects of GC-content and chromatin modifications we fitted models with different predictors, including GC-content, H3K9me3, H3K27me3, and centromeric domains. Based on model comparisons the model with the best predictions included the effect of GC-content, the effects of H3K9me3, H3K27me3, and centromeric domains, and the interaction between GC-content and H3K9me3 domain (Table S2). Based on model weights, the next best model that included additional interaction between H3K27 domain and GC-content was also plausible (Table S2). However, the overall predictions for these two models were similar. There were only a few mutations in low GC areas of H3K27me domains, so there is uncertainty in estimating a different slope for H3K27me3 domains; therefore we prefer the first model with the highest weight. GC-content had a strong effect on mutation rate, with areas of low GC having higher mutation rates (Figure 2D). Within H3K9me3 domains, GC-content has a smaller effect on mutation rate, and centromeric regions have a statistically detectable increase in mutation rate in addition to the effect of the H3K9me domain (Figure 2D, Table S3), even if centromeric regions always have H3K9me3 as well. H3K27me3 also increased mutation rate on top the GC-effect (Table S3).

Figure 2: A) Distribution of mutations along the seven chromosomes. Black vertical lines indicate mutations. Centromeric regions, H3K27 trimethylation, H3K36 dimethylation, H3K9 trimethylation domains, and duplicated regions are shown. B) Relative mutation rates for different genomic domains.H3K9 ex. centromeric are H3K9me3 domains where overlaps with centromeric regions have been excluded. Posterior medians and 95% HPD intervals are shown. C) Violin plots overlaid with boxplots for GC-content in different domains. D) Model estimates (on a log-scale) for mutation rate from a model with GC-content, H3K9me3, H3K27me3, and centromeric domains as predictors. Lines are posterior medians and envelopes are 95% HPD intervals.

Genetic variation in natural populations and mutation rate

To cross-validate our mutation model results, and to investigate role of mutation in maintenance of genetic variation across the genome, we examined if natural genetic variation can be explained by our mutation model. We calculated nucleotide polymorphism, θ_W, which measures how many polymorphic bases are found in a given length of sequence corrected for sample size, across the genomes of natural strains in 200 bp windows. We observed that mean θ_W was higher in the other domains compared to euchromatin (Figure 3A). The median estimate of θ_W was 0.0150 [0.0149, 0.0151] in euchromatin, difference to centromeric regions was 0.0213 [0.0210, 0.0216] units, difference to H3K9me3 domains was 0.0159 [0.0156, 0.0161] units, and difference to H3K27me3 domains 0.0117 [0.0115, 0.0119] units.

Figure 3: Nucleotide polymorphism, θ_W, was calculated across the genome in 200 bp windows, $n = 202310$. A) Violin plots overlaid with boxplots for distribution of θ_W in different domains. B) Relationship with θ_W and predicted mutation rate. For plotting data was been binned into hexes, because of the high number of overlapping points, and number of windows falling into each hex is shown by the legend.

To examine how genetic variation segregating in natural populations depends on mutation rate variation, we used our mutation model (Table S3) to predict variation in θ_W across the genome. We calculated the predicted mutation rate for each 200 bp window across the genome and observed that a simple linear model predicted moderate amount of variation in θ_W (Figure 3B). The slope of the regression line was 0.0096 [0.0095, 0.0097], so a tenfold increase in the predicted mutation rate meant an increase of 0.0096 in θ_W. A measure of model fit, the Bayesian R^2 value was 0.22 [0.21,

0.22]. Although this may seem a low R^2, one should take into account that this is after our mutation
model has been challenged with completely new data, and other evolutionary mechanisms than
mutation also influence θ_W. Choice of window size was not important: we tested different window
sizes and found the same relationship between predicted mutation rate and θ_W (Figure S4). Larger
windows even improved the fit, as there were less windows where $\theta_W = 0$, thus our choice of 200
bp window was conservative. Therefore mutation rate has a substantial influence on the amount of
genetic variation that is present across the genome in *N. crassa*.

Rate and spectrum of single nucleotide mutations

Next, we examined the rate and spectrum of different types of mutations. The rate of single nu-
cleotide mutations (SNM) across the whole genome was 6.7 [6.32, 7.11] $\times 10^{-10}$ mutations / bp /
mitosis. SNM rate in euchromatic regions was 2.46 [2.19, 2.77] $\times 10^{-10}$ mutations / bp / mitosis,
and SNM rate in H3K9me3 domains was 2.43 [2.25, 2.62] $\times 10^{-9}$ mutations / bp / mitosis. The
ratio of transition to transversion rates over the whole genome was 1.08 [0.96, 1.21], which is on
the low end of reported Ts/Tv ratios. The Ts/Tv ratio of euchromatic regions was 1.49 [1.17, 1.91],
which was higher than the Ts/Tv ratio in H3K9me domains 0.93 [0.8, 1.08].

As can be seen from the different transition over transversion ratios, the spectra of single nu-
cleotide mutations was different for H3K9me3 domains versus rest of the genome (Figure 4A). We
calculated ratios of the relative mutation rates in different domains, by taking different nucleotide
or trinucleotide frequencies (see below) into account in H3K9me3 domains and euchromatin. C:G
$\rightarrow$ G:C transversions were more common in H3K9 domains (Figure 4B). There was also weak
evidence that A:T $\rightarrow$ C:G and A:T $\rightarrow$ G:C mutations could have different rates in the different
domains. Their ratios were barely different from one when nucleotide frequencies were taken into
account, but the ratios barely overlapped with one when mutation rates were corrected for trinu-
cleotide frequencies (Figure 4B). However, A:T $\rightarrow$ T:A transversions had a lower rate in H3K9me3
domains after correcting for trinucleotide frequencies. Over the whole genome the four different
transversions occur at similar rates (Figure 4A), and the 2 transitions at higher rates. The A:T $\rightarrow$

G:C transition was the most common SNM, and more common than the C:G → T:A transition. The ratio of A:T → G:C to C:G → T:A transitions was 1.23 [1.04, 1.44].

Figure 4: A) Spectrum of relative SNM rates, dashed line shows the expected rate if all mutations occurred at equal frequencies. Nucleotide frequencies were taken into account in calculating the relative rates. Error bars are 95 % HPD intervals. B) Ratios of relative mutation rates in H3K9me3 over euchromatin. Points show ratios corrected for nucleotide frequencies and diamonds show ratios corrected for trinucleotide frequenices. Estimates are medians and range shows 95% HPD interval of the ratio. If the interval estimate is different from one, mutation rate is different in H3K9me3 domains and euchromatin.

Effects of local base composition

In order to understand what factors are influencing SNM rates we looked at the effect of local base pair context. For each SNM we extracted the two adjacent base pairs to get the trinucleotide

context. We combined trinucleotides with respect to sequence complementary as we do not know in which strand the mutation originally occurred, this leaves 32 trinucleotide classes. We calculated trinucleotide frequencies and observed that over the whole genome the observed frequencies were approximately at expected frequencies based on GC-content (Figure S5). However, in regions marked by H3K9 trimethylation there were strong departures from expected trinucleotide frequencies (Figure S5). Prompted by this observation we investigated whether differences in trinucleotide mutation rate could explain the observed differences in mutation rates across the genome. We compared different models with the trinucleotide classes and the effects of epigenetic domains. The model that gave the best predictions included an effect of the trinucleotide classes, effect of H3K9me3, centromeric regions, H3K27me3 regions, but no interactions between trinucleotide class and any of the epigenetic domains (Table S4). We observed the same results regarding the epigenetic domains as before: mutation rate was 8.1 [6.8, 9.5] times higher in H3K9me3 domains, centromeric regions had an additional increase on top of H3K9me3, and there was a small, 1.5 [1.1, 2.0] fold, increase in mutation rate in H3K27me3 domains (Figure S6). Thus, differences in trinucleotide composition rates were not driving the mutation rate differences in the different domains.

After taking into account trinucleotide frequencies, and the effects of the epigenetic domains, mutations were not equally distributed across the different trinucleotide classes (Figure 5A). Trinucleotide class GAT:ATC had the lowest relative mutation rate, while TCT:AGA had the highest. Trinucleotides with adjacent C:G pairs tend to have high relative mutation rates.

To investigate the effects of C:G and A:T base pairs in either 5' or 3' flanking positions, we fit a linear model that included the effects of the flanking bases, the mutating base pair, and we also incorporated the uncertainty in the relative mutation rate for each trinucleotide class. The model predictions are shown in Figure 5B, there was a tendency for trinucleotides with C:G as the mutating base to have higher relative mutation rate compared to A:T trinucleotides. However, this effect was not significant (Figure 5C). Similarly, for A:T trinucleotides there was a tendency for 5' C:G to protect against mutation, but this effect was not significant as 95% HPD interval barely

includes zero (Figure 5C). When the mutating base pair was C:G, neither 5' or 3' base had any detectable effect (Figure 5C).

Deletions, insertions, and translocations

In addition to point mutations, we examined the structural variants that occurred in the MA lines. We observed 97 deletions and 134 insertions in the MA lines. Frequent length for deletions and insertions were changes of one base pair, 88 out of 96 1 bp indels occurred in homopolymer stretches. The second most common length was 3 bp, which were predominantly changes in microsatellite repeats. Some large deletions were observed: the largest deletion was 17.8 kb, there were 3 deletions around 8.8 kb, and one around 2.5 kb. Otherwise most deletions were under 100 bp (Figure 6A). The largest observed insertion was 130 bp, with most insertions under 20 bp (Figure 6B).

Table 1: Mutation rates (mutations / genome / mitosis) for deletions and insertions and their ratio.

	Deletions	Insertions	Insertion / deletion ratio
All indels	0.0024 [0.0020, 0.0029]	0.0034 [0.0028, 0.0039]	1.38 [1.06, 1.77]
Repeats excluded	0.0011 [0.0008, 0.0014]	0.0003 [0.0002, 0.0005]	0.3 [0.14, 0.51]
Repeats only	0.0014 [0.0010, 0.0017]	0.0030 [0.0025, 0.0036]	2.25 [1.6, 3.05]

Mutation rates for insertions and deletions are shown in table 1. When all deletions and insertions were included in the analysis, insertions had slightly higher rates than deletions (Table 1). However, when we excluded homopolymers, microsatellites, and other repeats from the analysis we observed that deletions had a three times higher mutation rate than insertions (Table 1). Given that mean length of deletions that do not occur in repeats was 1160 bp, which was much longer than the 27 bp mean length for insertions, there was mutation pressure to lose DNA. Some of the large deletions observed in our data are outliers, but deletions tended to be longer at all scales (Figure 6). For mutations that occurred in repeats, mutation rate of insertions was over two fold higher than deletions.

Homopolymer stretches had particularly high rates indel mutations, we observed 92 mutations in homopolymers, most mutations in homopolymers were indels of 1 bp. More mutations occurred in A:T than in C:G homopolymers (Figure 6C). A:T homopolymer loci are approximately 1.7

Figure 5: A) Relative mutation rates for the 32 different classes of trinucleotides. B) Model predictions for effects of flanking bases. C) Estimates of model coefficients for effects of flanking bases. β_b is the effect of focal C:G relative to A:T, β_5 is the effect of flanking 5' C:G relative to A:T, β_3 is the effect of flanking 3' C:G relative to A:T, B_{I5} is the interaction effect of 5' C:G when the focal base is C:G, and B_{I3} is the interaction effect of 3' C:G when the focal base is C:G. Range shows 95% HPD interval of the relative mutation rate.

Figure 6: A) Distribution of deletion lengths in the range of 0 to 75 bp, inset shows the full distribution that includes few large deletions of several kb. B) Distribution of insertion lengths. C) Distribution of homopolymer lengths for those mutations that occurred in either A:T or C:G homopolymer stretches. D) Model estimates for mutation rate in homopolymers, A:T homopolymers had an overall higher mutation rate, and longer homopolymers had higher mutation rates.

times more common in the genome, but even when adjusting for frequencies, mutation rate in A:T homopolymers was 1.79 [1.4, 2.22] $\times 10^{-8}$ mutations / locus / mitosis compared to rate in C:G homopolymers of 8.15 [4.89, 11.86] $\times 10^{-9}$ mutations / locus / mitosis. Thus, mutations in A:T homopolymers were 2.2 [1.27, 3.49] times more common than in C:G polymers. We also observed that longer homopolymers had higher mutation rates. In a model with polymer length and polymer type, length had the same effect in for both A:T and C:G homopolymers (Figure 6D). This suggest that replication slippage, the mechanism suggested to be involved in indel mutations in repeats, tends to occur more often in longer repeats as expected.

When we analyzed how indels occurred in the different regions of the genome, we observed that deletions had a higher mutation rate in centromeric regions and in regions marked by H3K9me3 (Figure S7) compared to euchromatin. This was also true when only deletions in repeats were considered. For deletions excluding repeats, H3K9me3 and H3K27me3 domains had a higher mutation rate than euchromatin (Figure S7). For insertions we did not observe any differences in mutation rate in different domains (Figure S7).

We observed 5 translocations in the MA lines, two of these were among the SVs confirmed by PCR and Sanger sequencing. Three translocations were from one chromosome to another, and two occurred among unmapped contigs. Mean translocation length was 316 bp. The translocation rate was 1.19 [0.33, 2.41] $\times 10^{-4}$ translocations / genome / mitosis. Translocations were much rarer than other insertions or deletions, thus we don't have enough mutations to do further analyses of their properties.

Complex mutations

We observed 9 cases were two SNMs or 1 bp indels occurred within few base pairs of each other in the same MA line. While it is possible that two independent mutations occurred next to each other, it is unlikely. These changes were more likely caused by a single mutational event. The observed complex mutational events are listed in table 2. To confirm whether these changes were caused by a single mutational event or two independent events, we used Sanger sequencing to check

the genotypes of MA lines from intermediate transfers stored during the experiment. We always observed that the two changes appeared together (Table 2). The most parsimonious explanation is that these changes appeared as a result of single mutational events, likely caused by DNA repair error by an error prone DNA-polymerase.

We treated complex mutations as single events in calculations were all mutations were used to calculate overall mutation rates. The rate of complex mutations was 5.42 [2.22, 9.3] $\times 10^{-12}$ mutations / bp / mitosis. The rate of SNMs over rate of complex mutations was 123.7 [57.48, 231.49], making point mutations over 100-fold more common than complex events.

Table 2: Complex mutations. Chr. is the chromosome or contig, position is the coordinate of the most 5' end change, dist. is the distance between nucleotide changes. Sequence column shows the ancestor sequence on first row and the MA line sequence on second row, mutated bases are in bold. Final column contains the MA experiment transfer numbers (mutations absent, mutations present) of those transfers when was the last transfer where none of the changes could be observed in the MA line, and the transfer when all changes were observed in the MA line. We always observed all changes in complex mutations to be present together. tv. = transversion, ts. = transition

MA-line	Chr.	Position	Dist. (bp)	Sequence	Changes	Mutation occurred between transfers
Line 27	1	1194971	2	`GG-TGC` `GG`**`TTT`**`C`	Insertion (T) and G → T tv.	(35, 38)
Line 31	1	4086493	1	`GTAA` `G`**`AG`**`A`	T → A tv. and A → G ts.	(10, 40)
Line 18	1	5852116	NA	`GATACTATCGTAATATATCACTCTTGAACTAC` `G`**`TAGTTCAAGAGTAT`**`----------------C`	Deletion and insertion	(20, 25)
Line 21	1	9717311	3	`CGAACT` `C-AA`**`A`**`T`	Deletion (G) and C → A tv.	(15, 20)
Line 24	3	742257	1	`AAAA` `A`**`TT`**`A`	Two A → T tv.	(10, 15)
Line 35	3	1051825	2	`ATTTG` `A`**`CTG`**`G`	T → C ts. and T → G tv.	(15, 20)
Line 12	6	2848187	2	`CTATA` `C-A`**`A`**`A`	Deletion (T) and T → A tv.	not checked
Line 23	contig 8	176939	4	`ATTTTTT` `A`**`C`**`TTT`**`C`**`T`	Two T → C ts.	(7, 8)
Line 20	contig 9	121389	3	`AATTTA` `A`**`T`**`TT`**`AA`	A → T tv. and T → A tv.	(30, 35)

Comparison of mutation rate and spectra during meiosis and mitosis

Wang et al. (2020) observed an extremely high mutation rate during meiosis, as a result of C:G → T:A transitions induced by RIP. We re-analyzed their data and combined it with data about chromatin modifications (see supplementary results for details). We observed heterogeneity in activa-

tion of RIP that was not taken into account in the original analysis (see Figure S8 and supplementary results). Based on our analysis, mutation rate during meiosis in euchromatin was 1.07 [0.60, 1.67] $\times 10^{-8}$ mutations / meiosis / bp, and 2.54 [0.11, 7.55] $\times 10^{-7}$ in H3K9me3 domains. Mutation rate was substantially higher during meiosis than mitosis, as observed by Wang et al. (2020), but not as high as their analysis suggested. Comparing mutation spectra in mitosis and meiosis shows that in euchromatic regions, A:T $\rightarrow$ C:G, and C:G $\rightarrow$ G:C transversions were more common in meiosis, while C:G $\rightarrow$ A:T transversions were less common (Figure S9). In heterochromatin C:G $\rightarrow$ T:A transitions overwhelm all other mutations in meiosis (Figure S8, S9).

Discussion

We have estimated spontaneous mutation rate during asexual growth to high precision in *Neurospora crassa*. Our estimate of point mutation rate across the whole genome of 6.7 [6.32, 7.11] $\times 10^{-10}$ mutations / bp / mitosis is higher, although close to an estimate of asexual mutation rate of 6.03×10^{-10} obtained by Wang et al. (2020) with only 64 observed mutations. Previous estimate from marker gene studies suggested that mutation rate is 4.10–4.66×10^{-9} (Lynch et al., 2016), but neither our nor the results of Wang et al. (2020) agree with this. Point mutation rate in euchromatic regions was 0.007 [0.006, 0.008] mutations / genome / mitosis, which is in line with results obtained by Drake (1991) who observed that mutation rate per genome for microbes seems to be around 0.003 mutations per genome per generation with approximately two fold variation around this mean. Thus, asexual mutation rate in *N. crassa* in euchromatic regions seems to be rather typical for a microbe.

N. crassa has a striking difference in rate and spectrum of mutation during sexual and asexual reproduction (Wang et al., 2020). During sexual reproduction a genome defence mechanism called RIP is activated which recognizes duplicated regions in premeiotic cells and induces C:G $\rightarrow$ T:A transitions in those regions (Selker, 1990). The biological function of RIP is presumably to protect the genome against transposons and other selfish genetic elements. Mutations induced during

sexual reproduction happen mainly in ancestral duplications (Wang et al., 2020), and these regions nearly completely overlap with regions of H3K9 trimethylation. In contrast, during asexual reproduction, while H3K9me3 domains have a higher mutation rate than euchromatin, the difference is much smaller and the spectrum of mutations is different than in meiosis. For euchromatic regions, our analysis supports a higher mutation rate during meiosis than in mitosis, although not as high as suggested by Wang et al. (2020). We also observed that the spectrum of mutations was different in meiosis, but notably there was no difference in C:G $\rightarrow$ T:A transitions in euchromatin. Moreover, since gene density is much higher in euchromatic regions, action of RIP likely does not result in as high genetic load as suggested by Wang et al. (2020).

While many studies have reported effects of chromatin structure on mutation rate, these have often been based on indirect inference from species divergence and polymorphism (Sasaki et al., 2009; Ying et al., 2010; Washietl et al., 2008). Our study adds to the growing body of direct observations that chromatin structure is an important determinant of spontaneous mutation rate. We observed extensive variation across the genome in mutation rate and mutation spectra due to chromatin modifications. Mutation rate for SNMs in domains marked by H3K9 trimethylation that determines heterochromatic regions in *N. crassa* was tenfold higher than in euchromatic regions. In *N. crassa* centromeric regions determined by the presence of centromeric histone variant CenH3 always overlap with H3K9 trimethylation (Smith et al., 2011). In centromeric regions there was an additional effect of increased mutation rate on top of the H3K9me3 effect. Increased mutation rate in heterochromatic and centromeric regions has also been found by Weng et al. (2019) in the plant *Arabidopsis thaliana*. These results are also in line with observations from human cancer cells that have higher rates of mutation in heterochromatic regions (Schuster-Böckler and Lehner, 2012; Polak et al., 2015). In *N. crassa* facultative heterochromatin is marked by H3K27 trimethylation (Jamieson et al., 2013), and we observed that mutation rate was slightly elevated in H3K27me3 domains, although this effect was much smaller than for H3K9 trimethylation.

In *N. crassa* methylation in H3K9 and H3K36 are almost completely mutually exclusive. However, H3K36me2 is not a straightforward mark of euchromatin as it can be deposited by two en-

zymes: SET-2 and ASH1. Genes marked with H3K36 methylation by SET-2 are actively tran-
scribed, while genes marked with H3K36 methylation by ASH1 are silenced and can be further
marked by H3K27 trimethylation (Bicocca et al., 2018). SET-2 is responsible for most of H3K36
methylation, and we considered all regions that lacked H3K9me3 and H3K27 trimethylation to
be euchromatin. Monroe et al. (2022) observed that in *Arabidopsis* mutation rate was affected by
several different epigenetic marks, and they suggested that mutation rate was lower in genes that
were actively transcribed, and perhaps even fine tuned for highly expressed genes, possibly due to
the presence of H3K36 methylation or other epigenetic marks of active transcription. While we
also observed that H3K36 domains (absence of H3K9me3) had a lower mutation rate compared to
heterochromatin, so in that sense genes also had a lower mutation rate. However, our dataset does
not contain enough mutations to address if gene expression levels quantitatively influence mutation
rate in *N. crassa*.

What is the mechanism causing mutation rate variation across the genome? In addition to
increased mutation rate in regions with lower GC-content we observed an effect of chromatin mod-
ifications. Chromatin structure is involved in DNA lesion and repair, and this may explain why
heterochromatic regions have a higher mutation rate. In yeast, actively transcribed regions contain
acetylation at H3K56 which suppresses spontaneous mutations (Kadyrova et al., 2013). Moreover,
analyses based on human tumours suggest that DNA mismatch repair works more efficiently in eu-
chromatin than in heterochromatin and it is DNA repair that is variable, not the supply of mutations
themselves (Supek and Lehner, 2015). Recently, Habig et al. (2021) investigated mutation rate in
relation to epigenetic modification in the fungus *Zymoseptoria tritici*. They observed that H3K27
trimethylation increased mutation rate as in this study. Furthermore, Habig et al. (2021) observed
that a mutant lacking H3K27me3 had reduced mutation rate in regions that had H3K27me3 in the
wild type. This suggested that H3K27me3 is causal, and DNA accessibility may be the reason.
Interestingly, Habig et al. (2021) did not detect an effect for H3K9me3 like we observed, which
may indicate that species specific differences in heterochromatin.

Further evidence supporting DNA accessibility hypothesis was gathered by a study performed

by Yazdi et al. (2015) in which the authors observed that, when using DNA repair machinery knock-out yeast strains, mutation rate did not increase in high nucleosome occupancy regions like in wild type strains. Differential exposure of heterochromatin to natural mutagens, such as oxidative damage, does not seem to explain our results. As mutations typically associated with oxidative damage: G $\rightarrow$ C transversions, C $\rightarrow$ T transitions, and G $\rightarrow$ T transversions (McBride et al., 1991; Cheng et al., 1992), were not systematically overrepresented in regions with H3K9 trimethylation, only the relative amount of G $\rightarrow$ C transversions differed between euchromatic and H3K9 trimethylated domains.

Among the mutations that were observed, sequence context had an effect on mutation rate. It is known that local sequence context can influence the probability of mutation and this phenomenon has been frequently observed (Ness et al., 2015). However, the mechanisms of why some sequence contexts are prone to mutation are less clear. For example, the formation rate of the thymine-thymine cyclobutane dimer, a form of DNA damage caused by sunlight, is dependent on flanking sequence (Law et al., 2013), but the biophysical basis is unclear (Law et al., 2013). We did observe a tendency of 5' C:G base pair to protect against mutation when the mutating base pair is A:T, and this phenomenon has been observed before in the context of TT-dimers (Law et al., 2013). However, to clarify this effect further, much more, or possibly strand specific data would need to be collected. We also observed a tendency of trinucleotides with with the mutation base pair being C:G having higher relative mutation rates, this could be because this class of trinucleotides includes many with GG dimers that have been found susceptible to oxidation (Hanrahan et al., 1997).

We did not find statistical support for a model that had unique trinucleotide mutation rates, in regions of H3K9me3. However, lack of overall effect may hide a few trinucleotides that differ. One hypothesis for different trinucleotide mutation rates in heterochromatin would be that the presence of DNA methylation increases mutation rate in certain contexts. In *N. crassa* DNA methylation occurs only at H3K9 trimethylated domains (Tamaru and Selker, 2001), and deamination of 5-methylcytosine is a known cause of mutations (Cooper et al., 2010). However, deamination should cause mainly C:G $\rightarrow$ T:A transitions, and we did not observe an excess of C:G $\rightarrow$ A:T transitions

in H3K9me3 domains. Instead we observed excess of C:G → G:C transversions in H3K9me3 domains. Therefore DNA methylation seems to contribute quite little to spontaneous genetic mutations in *N. crassa*, possibly because DNA methylation is so rare in this species (Hosseini et al., 2020).

For spontaneous insertions and deletions, we observed that deletions were more common than insertions when repeated sequences were excluded. Since deletions also tended to be longer, mutations have the tendency to reduce the size of the genome. Similar patterns in mutational bias have been observed for *Drosophila melanogaster* (Leushkin et al., 2013). For repeated sequences, we analyzed homopolymer sequences in more detail. We found that while A:T homopolymers are more common in *N. crassa*, they also have a higher mutation rate. This is in contrast to observations in nematodes, where C:G homopolymers had much higher mutation rates than A:T homopolymers (Denver et al., 2004) suggesting that there are species specific differences in mutation rates of these sequences.

We identified mutations with multiple base changes due to a single mutational event. Such complex mutations are thought to arise from the action of error-prone translesion DNA polymerases, such as Pol ζ (Stone et al., 2012). A recent large scale survey of human trios identified many multinucleotide mutations, and observed that mutations that were 2–10 bp apart showed an over-representation of A:T → T:A, and A:T → G:C mutations (Besenbacher et al., 2016). While we did not observe enough complex mutations to conduct a statistical analysis, our results are compatible with this pattern as most nucleotide changes in complex mutations were either A:T → T:A or A:T → G:C. However, higher mutation rate in AT-rich regions could also be important. We did not observe GC → AA or GA → TT tandem mutations, that are the most common tandem mutation in humans (Harris and Nielsen, 2014; Besenbacher et al., 2016). In *N. crassa* complex mutations represented only 0.7% of observed mutations, which is smaller than approximately 3% observed in humans, thus it may be that the impact of multinucleotide mutations on population genetic inference may be smaller in *N. crassa* than in humans.

Conclusions

We have observed variation in mutation rate and mutation spectrum across the genome of *N. crassa*. It appears that variation in mutation rate in eukaryotes due to chromatin modifications will be the norm rather than the exception. This raises an issue that there is no single mutation rate for a given species, as chromatin state has such a large effect. In the future it will be more informative to speak about mutation rates specific to different chromatin domains. The implication for evolutionary genetics is that variation in mutation rates must be taken into account in order to quantify which evolutionary forces act on the genome. Genes residing in different regions of the genome can differ substantially in their mutation loads, and it seems that variation in mutation rate does also have a large part in determining how much genetic variation is segregating in a given region of the genome. To accurately estimate imprints of selection on natural genetic variation, we must account for variation in mutation rates in evolutionary models. Therefore, detailed models of mutation rates are needed for different species. This study is the first step in order to do this in *N. crassa*, a model microbial eukaryote.

Methods

Mutation accumulation experiment

We used two ancestors for the MA experiment: B 26708 and B 26709, these strains have been generated by backcrossing mating type mat a from 4200 into 2489 background nine times (Kronholm et. al. 2020). There were 20 MA lines for both ancestors, giving 40 lines in total. Common protocols for culturing N. crassa were followed (see supplementary methods). We propagated the MA lines for 40 transfers.

To estimate the number of mitoses that that happened in the MA experiment during a transfer, we used to strain that had been tagged with fluorescent protein at histone H1 (Freitag et al. 2004). Using this strain we counted the numbers of nuclei that were present in each phase of one transfer of the MA experiment by confocal microscopy (Figure 1C), and used this data to estimate the number

of mitoses the lines went through (see supplementary methods).

Strains from natural populations

We obtained 33 strains from the Fungal Genetics Stock Center (McCluskey et al., 2010), and re-sequenced these strains. In addition we obtained genome sequencing data for additional 23 strains from Zhao et al. (2015). In total, including the laboratory strain 2489, we had 57 strains with sequencing data (Table S5).

DNA extraction

We extracted DNA from the strains using Phenol-Chloroform extraction, and further purified it using polyethyleneglycol precipitation. DNA quality was checked on aragose gels. For a detailed protocol see supplementary methods.

Genome sequencing

Sample DNA was sent for sequencing at Novogene (Cambridge, UK) using Illumina platform with paired-end 150 bp libraries. Libraries were prepared by fragmenting the DNA by sonication, adapter ligation, and PCR-amplification. Libraries were sequenced to 30x target coverage. Reads with adapter sequences, $> 10\%$ N's, or $> 50\%$ bases of low quality Q score ≤ 5 were removed.

Read mapping and genotyping

We mapped reads against the *N. crassa* reference genome (assembly version NC12). The strain used for the original genome project was 2489, so the MA experiment ancestors should be nearly identical to the reference. We mapped reads against the *N. crassa* reference genome. The strain used for the original genome project was 2489, so the MA experiment ancestors should be nearly identical to the reference. Reads were mapped using BWA-MEM (Li, 2013).

We used the GATK version 4.2.0.0 (McKenna et al., 2010) pipeline to call SNMs and indels. First we ran haplotypecaller for each sample and then jointly genotyped all samples together. *Neurospora crassa* is haploid, but we ran haplotype caller in diploid mode as mapping errors manifest as heterozygous sites in haploids (Kronholm et al., 2017), see also Li (2014) and Ness et al. (2012). We used wormtable (Kelleher et al., 2013) and custom python scripts to filter for high quality sites. To produce the final set of mutations, we checked all candidate mutations manually by inspecting alignment in IGV (Thorvaldsdóttir et al., 2013). SNPs were genotyped using the same pipeline, see supplementary methods for details.

For structural variants, we first compared the performance of different SV genotyping algorithms using simulations. We selected DELLY and Lumpy for genotyping. See supplementary materials for details.

Validation of mutations with Sanger sequencing

To verify a sample of the observed mutations, we performed Sanger sequencing using standard methods. We selected mutations that passed our threshold but had the lowest quality scores. We attempted to confirm 83 annotated low quality score mutations, 30 of them were point mutations, 37 small indels and 16 larger structural variants. List of primers and confirmed mutations are given in the supplementary file S1. Additionally, complex mutations, mutations where a suspected single event created multiple changes, were selected for verification and we sequenced time points from the middle of the MA-experiment to see if the multiple changed sites appeared together in the MA-line.

Chromatin modifications

We used publicly available data to determine where chromatin modifications occurred. ChIP-seq reads for H3K9 trimethylation and H3K27 trimethylation were obtained from Jamieson et al. (2013), accession numbers SRX248101 and SRX248097. Jamieson et al. (2013) used the same strain, 2489, that we used as the ancestor of the MA experiment. They also tested whether different

growth media had an effect on H3K27me3 regions, and observed that they were stable. Jamieson et al. (2013) extracted chromatin from mycelium, which gives rise to the asexual spores used for transfers in our experiment, so the chromatin states they observed shoud be relevant for our experiment. Taken together, this data provides a very good approximation of the chromatin states the ancestors of the MA lines had in our experiment. Data for H3K36 methylation was obtained from Bicocca et al. (2018), accession number SRX4549854. Data for centromeric regions were obtained from Smith et al. (2011). See supplementary methods for details.

Statistical analysis

We used Bayesian statistics in all statistical modeling. This approach is different from the traditional frequentist approach. We refer readers who are unfamiliar with this approach to McElreath (2015), which is an excellent introduction to the Bayesian statistics. As a rule of thumb for comparison to frequentist methods: if a 95% interval estimate of a parameter does not contain zero, the parameter is statistically different from zero, i. e. significant. We tested differences in mutation rates by computing ratios from posterior distributions. If the interval of the ratio did not include one, the mutation rates were statistically different. Estimates were reported as medians and 95% highest posterior density intervals in square brackets.

Mutation rate estimates

Mutation rates were estimated using Bayesian Poisson models implemented with the Stan language (Carpenter et al., 2017) interfaced from R 3.6.0 (R Core Team, 2019) with the "brms" package (Bürkner, 2017). The basic model for estimating mutation rate was

$$y_i \sim \text{Poisson}(\lambda_i) \tag{1}$$

$$\log(\lambda_i) = \alpha$$

$$\alpha \sim \text{N}(0, 10)$$

where y_i is the number of mutations in ith MA-line, λ_i is the poisson rate parameter, and α is the intercept. The linear model part was modified accordingly if other predictors were used. We can then calculate mutation rate, μ, from posterior distributions as

$$\mu = \frac{\exp(\alpha)}{Ntm} \tag{2}$$

where N is the number of called nucleotides, t is the number of transfers the MA-lines went through, and m is the number of mitoses per transfer. Note that we used the posterior distribution for m, so any uncertainty in number of mitoses is incorporated into our estimate of mutation rate. To get mutation rate per genome N is removed from the denominator. We used a weakly regularizing prior for α. Settings for MCMC estimation were: 1000 iterations of warm-up followed by 3000 iterations of sampling with four independent chains. MCMC convergence was monitored by traceplots and $\hat{R}$ values. No convergence problems were observed. For details of the other models see supplementary methods.

Mutation rate in repeats

Homopolymer sequences in the genome were detected using MISA (Beier et al., 2017), we used 5 bp as the minimum homopolymer length and extracted repeat counts for all homopolymer loci in the genome. Counts of homopolymers of different lengths were used as an offset term in a model used to estimate mutation rates in homopolymers.

Population genetics

We estimated nucleotide polymorphism, θ_W, from the SNP data for the population sample following Ferretti et al. (2012), which allows us to deal with missing data.

Data access

Sequencing data generated in this study has been deposited to the short read archive, project number PRJNA839531, with sequence accession numbers SRX15345434–SRX15345507. Other data and scripts are available at https://github.com/ikron/mutation_ms and as supplemental material.

Acknowledgments

This study was supported by a grant from the Academy of Finland (no. 321584) to IK. We thank Dr. Visa Ruokolainen for help with confocal microscopy, the Finnish CSC-IT Center for Science Ltd. for providing computational resources, and Dr. Matthieu Bruneaux for comments on the manuscript.

Author contributions

I.K. conceived the study. P.A.M.S., M. V., M.L., and I.K performed experiments. I.K., M.V., and P.A.M.S. analyzed the data. I.K. and M.V wrote the manuscript. All authors edited the final manuscript.

References

Beier, S., Thiel, T., Münch, T., Scholz, U., and Mascher, M., 2017. MISA-web: a web server for microsatellite prediction. *Bioinformatics*, **33**(16):2583–2585.

Besenbacher, S., Sulem, P., Helgason, A., Helgason, H., Kristjansson, H., Jonasdottir, A., Jonasdottir, A., Magnusson, O. T., Thorsteinsdottir, U., Masson, G., *et al.*, 2016. Multi-nucleotide *de novo* mutations in humans. *PLOS Genetics*, **12**(11):1–15.

Bicocca, V. T., Ormsby, T., Adhvaryu, K. K., Honda, S., and Selker, E. U., 2018. ASH1-catalyzed H3K36 methylation drives gene repression and marks H3K27me2/3-competent chromatin. *eLife*, **7**:e41497.

Bürkner, P.-C., 2017. brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, **80**(1):1–28.

Cambareri, E. B., Singer, M. J., and Selker, E. U., 1991. Recurrence of repeat-induced point mutation (RIP) in *Neurospora crassa*. *Genetics*, **127**(4):699–710.

Campos, J. L., Zhao, L., and Charlesworth, B., 2017. Estimating the parameters of background selection and selective sweeps in *Drosophila* in the presence of gene conversion. *Proceedings of the National Academy of Sciences*, **114**(24):E4762–E4771.

Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P., and Riddell, A., *et al.*, 2017. Stan: A probabilistic programming language. *Journal of Statistical Software*, **76**(1):1–32.

Castellano, D., Eyre-Walker, A., and Munch, K., 2020. Impact of Mutation Rate and Selection at Linked Sites on DNA Variation across the Genomes of Humans and Other Homininae. *Genome Biology and Evolution*, **12**(1):3550–3561.

Chen, X., Chen, Z., Chen, H., Su, Z., Yang, J., Lin, F., Shi, S., and He, X., 2012. Nucleosomes suppress spontaneous mutations base-specifically in eukaryotes. *Science*, **335**(6073):1235–1238.

Cheng, K. C., Cahill, D. S., Kasai, H., Nishimura, S., and Loeb, L. A., 1992. 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G-T and A-C substitutions. *Journal of Biological Chemistry*, **267**(1):166–172.

Cooper, D. N., Mort, M., Stenson, P. D., Ball, E. V., and Chuzhanova, N. A., 2010. Methylation-mediated deamination of 5-methylcytosine appears to give rise to mutations causing human inherited disease in CpNpG trinucleotides, as well as in CpG dinucleotides. *Human Genomics*, **4**(6):406.

Denver, D. R., Morris, K., Kewalramani, A., Harris, K. E., Chow, A., Estes, S., Lynch, M., and Thomas, W. K., 2004. Abundance, distribution, and mutation rates of homopolymeric nucleotide runs in the genome of *Caenorhabditis elegans*. *Journal of Molecular Evolution*, **58**(5):584–595.

Drake, J. W., 1991. A constant rate of spontaneous mutation in DNA-based microbes. *Proceedings of the National Academy of Sciences*, **88**(16):7160–7164.

Eyre-Walker, A. and Keightley, P. D., 2007. The distribution of fitness effects of new mutations. *Nature Reviews Genetics*, **8**(8):610–618.

Ferretti, L., Raineri, E., and Ramos-Onsins, S., 2012. Neutrality tests for sequences with missing data. *Genetics*, **191**(4):1397–1401.

Habig, M., Lorrain, C., Feurtey, A., Komluski, J., and Stukenbrock, E. H., 2021. Epigenetic modifications affect the rate of spontaneous mutations in a pathogenic fungus. *Nature Communications*, **12**(1):5869.

Halligan, D. L. and Keightley, P. D., 2009. Spontaneous mutation accumulation studies in evolutionary genetics. *Annual Review of Ecology, Evolution, and Systematics*, **40**(1):151–172.

Hanrahan, C. J., Bacolod, M. D., Vyas, R. R., Liu, T., Geacintov, N. E., Loechler, E. L., and Basu, A. K., 1997. Sequence specific mutagenesis of the major (+)-anti-benzo[a]pyrene diol epoxide–DNA adduct at a mutational hot spot *in vitro* and in *Escherichia coli* cells. *Chem. Res. Toxicol.*, **10**(4):369–377.

Harris, K. and Nielsen, R., 2014. Error-prone polymerase activity causes multinucleotide mutations in humans. *Genome Research*, **24**(9):1445–1454.

Hosseini, S., Meunier, C., Nguyen, D., Reimegård, J., and Johannesson, H., 2020. Comparative analysis of genome-wide DNA methylation in *Neurospora*. *Epigenetics*, **15**(9):972–987. PMID: 32228351.

Jamieson, K., Rountree, M. R., Lewis, Z. A., Stajich, J. E., and Selker, E. U., 2013. Regional control of histone H3 lysine 27 methylation in *Neurospora*. *Proceedings of the National Academy of Sciences*, **110**(15):6027–6032.

Johri, P., Charlesworth, B., and Jensen, J. D., 2020. Toward an evolutionarily appropriate null model: Jointly inferring demography and purifying selection. *Genetics*, **215**(1):173–192.

Kadyrova, L. Y., Mertz, T. M., Zhang, Y., Northam, M. R., Sheng, Z., Lobachev, K. S., Shcherbakova, P. V., and Kadyrov, F. A., 2013. A reversible histone H3 acetylation cooperates with mismatch repair and replicative polymerases in maintaining genome stability. *PLOS Genetics*, **9**(10):1–16.

Katju, V. and Bergthorsson, U., 2018. Old trade, new tricks: Insights into the spontaneous mutation process from the partnering of classical mutation accumulation experiments with high-throughput genomic approaches. *Genome Biology and Evolution*, **11**(1):136–165.

Keightley, P. D. and Eyre-Walker, A., 2007. Joint inference of the distribution of fitness effects of deleterious mutations and population demography based on nucleotide polymorphism frequencies. *Genetics*, **177**(4):2251–2261.

Keightley, P. D., Ness, R. W., Halligan, D. L., and Haddrill, P. R., 2014. Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics*, **196**(1):313–320.

Keightley, P. D., Pinharanda, A., Ness, R. W., Simpson, F., Dasmahapatra, K. K., Mallet, J., Davey, J. W., and Jiggins, C. D., 2015. Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Molecular Biology and Evolution*, **32**(1):239–243.

Keith, N., Tucker, A. E., Jackson, C. E., Sung, W., Lucas Lledó, J. I., Schrider, D. R., Schaack, S., Dudycha, J. L., Ackerman, M., Younge, A. J., *et al.*, 2016. High mutational rates of large-scale duplication and deletion in *Daphnia pulex*. *Genome Research*, **26**(1):60–69.

Kelleher, J., Ness, R. W., and Halligan, D. L., 2013. Processing genome scale tabular data with wormtable. *BMC Bioinformatics*, **14**(1):1–5.

Kouzarides, T., 2007. Chromatin modifications and their function. *Cell*, **128**(4):693 – 705.

Kronholm, I., Bassett, A., Baulcombe, D., and Collins, S., 2017. Epigenetic and genetic contributions to adaptation in *Chlamydomonas*. *Molecular Biology and Evolution*, **34**(9):2285–2306.

Law, Y. K., Forties, R. A., Liu, X., Poirier, M. G., and Kohler, B., 2013. Sequence-dependent thymine dimer formation and photoreversal rates in double-stranded DNA. *Photochem. Photobiol. Sci.*, **12**:1431–1439.

Leushkin, E. V., Bazykin, G. A., and Kondrashov, A. S., 2013. Strong mutational bias toward deletions in the *Drosophila melanogaster* genome is compensated by selection. *Genome Biology and Evolution*, **5**(3):514–524.

Li, C. and Luscombe, N. M., 2020. Nucleosome positioning stability is a modulator of germline mutation rate variation across the human genome. *Nature Communications*, **11**(1):1363.

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:1303.3997 [q-bio.GN]*, .

Li, H., 2014. Toward better understanding of artifacts in variant calling from high-coverage samples. *Bioinformatics*, **30**(20):2843–2851.

Lynch, M., Ackerman, M. S., Gout, J.-F., Long, H., Sung, W., Thomas, W. K., and Foster, P. L., 2016. Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet*, **17**(11):704–714.

Makova, K. D. and Hardison, R. C., 2015. The effects of chromatin organization on variation in mutation rates in the genome. *Nature Reviews Genetics*, **16**(4):213–223.

McBride, T. J., Preston, B. D., and Loeb, L. A., 1991. Mutagenic spectrum resulting from DNA damage by oxygen radicals. *Biochemistry*, **30**(1):207–213.

McCluskey, K., Wiest, A., and Plamann, M., 2010. The fungal genetics stock center: a repository for 50 years of fungal genetics research. *J Biosci*, **35**(1):119–126.

McElreath, R., 2015. *Statistical Rethinking - A Bayesian course with examples in R and Stan*. CRC Press, New York.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., *et al.*, 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, **20**(9):1297–1303.

Monroe, J. G., Srikant, T., Carbonell-Bejerano, P., Becker, C., Lensink, M., Exposito-Alonso, M., Klein, M., Hildebrandt, J., Neumann, M., Kliebenstein, D., *et al.*, 2022. Mutation bias reflects natural selection in *Arabidopsis thaliana*. *Nature*, **602**:101–105.

Ness, R. W., Morgan, A. D., Colegrave, N., and Keightley, P. D., 2012. Estimate of the spontaneous mutation rate in *Chlamydomonas reinhardtii*. *Genetics*, **192**:1447–1454.

Ness, R. W., Morgan, A. D., Vasanthakrishnan, R. B., Colegrave, N., and Keightley, P. D., 2015. Extensive de novo mutation rate variation between individuals and across the genome of *Chlamydomonas reinhardtii*. *Genome Research*, **25**:1739–1749.

Ossowski, S., Schneeberger, K., Lucas-Lledó, J. I., Warthmann, N., Clark, R. M., Shaw, R. G., Weigel, D., and Lynch, M., 2010. The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science*, **327**(5961):92–94.

Polak, P., Karlić, R., Koren, A., Thurman, R., Sandstrom, R., Lawrence, M. S., Reynolds, A., Rynes, E., Vlahoviček, K., Stamatoyannopoulos, J. A., *et al.*, 2015. Cell-of-origin chromatin organization shapes the mutational landscape of cancer. *Nature*, **518**(7539):360–364.

R Core Team, 2019. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Sasaki, S., Mello, C. C., Shimada, A., Nakatani, Y., Hashimoto, S.-i., Ogawa, M., Matsushima, K., Gu, S. G., Kasahara, M., Ahsan, B., *et al.*, 2009. Chromatin-associated periodicity in genetic variation downstream of transcriptional start sites. *Science*, **323**(5912):401–404.

Schuster-Böckler, B. and Lehner, B., 2012. Chromatin organization is a major influence on regional mutation rates in human cancer cells. *Nature*, **488**(7412):504–507.

Selker, E. U., 1990. Premeiotic instability of repeated sequences in *Neurospora crassa*. *Annual Review of Genetics*, **24**(1):579–613. PMID: 2150906.

Smith, K. M., Phatale, P. A., Sullivan, C. M., Pomraning, K. R., and Freitag, M., 2011. Heterochromatin is required for normal distribution of *Neurospora crassa* CenH3. *Molecular and Cellular Biology*, **31**(12):2528–2542.

Stone, J. E., Lujan, S. A., and Kunkel, T. A., 2012. DNA polymerase zeta generates clustered mutations during bypass of endogenous DNA lesions in *Saccharomyces cerevisiae*. *Environmental and Molecular Mutagenesis*, **53**(9):777–786.

Sung, W., Ackerman, M. S., Gout, J.-F., Miller, S. F., Williams, E., Foster, P. L., and Lynch, M., 2015. Asymmetric context-dependent mutation patterns revealed through mutation–accumulation experiments. *Molecular Biology and Evolution*, **32**(7):1672–1683.

Supek, F. and Lehner, B., 2015. Differential DNA mismatch repair underlies mutation rate variation across the human genome. *Nature*, **521**(7550):81–84.

Tamaru, H. and Selker, E. U., 2001. A histone H3 methyltransferase controls DNA methylation in *Neurospora crassa*. *Nature*, **414**(6861):277–283.

Thorvaldsdóttir, H., Robinson, J. T., and Mesirov, J. P., 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Briefings in Bioinformatics*, **14**(2):178–192.

Tolstorukov, M. Y., Volfovsky, N., Stephens, R. M., and Park, P. J., 2011. Impact of chromatin structure on sequence variability in the human genome. *Nature Structural & Molecular Biology*, **18**(4):510–515.

Wang, L., Sun, Y., Sun, X., Yu, L., Xue, L., He, Z., Huang, J., Tian, D., Hurst, L. D., and Yang, S., *et al.*, 2020. Repeat-induced point mutation in *Neurospora crassa* causes the highest known mutation rate and mutational burden of any cellular life. *Genome Biology*, **21**(1):142.

Washietl, S., Machné, R., and Goldman, N., 2008. Evolutionary footprints of nucleosome positions in yeast. *Trends in Genetics*, **24**(12):583–587.

Weng, M.-L., Becker, C., Hildebrandt, J., Neumann, M., Rutter, M. T., Shaw, R. G., Weigel, D., and Fenster, C. B., 2019. Fine-grained analysis of spontaneous mutation spectrum and frequency in *Arabidopsis thaliana*. *Genetics*, **211**(2):703–714.

Yazdi, P. G., Pedersen, B. A., Taylor, J. F., Khattab, O. S., Chen, Y.-H., Chen, Y., Jacobsen, S. E., and Wang, P. H., 2015. Increasing nucleosome occupancy is correlated with an increasing mutation rate so long as DNA repair machinery is intact. *PLoS One*, **10**(8):e0136574.

Ying, H., Epps, J., Williams, R., and Huttley, G., 2010. Evidence that localized variation in primate sequence divergence arises from an influence of nucleosome placement on DNA repair. *Molecular biology and evolution*, **27**(3):637–649.

Zhao, J., Gladieux, P., Hutchison, E., Bueche, J., Hall, C., Perraudeau, F., and Glass, N. L., 2015. Identification of allorecognition loci in *Neurospora crassa* by genomics and evolutionary approaches. *Molecular Biology and Evolution*, **32**(9):2417–2432.

Zhu, Y. O., Siegal, M. L., Hall, D. W., and Petrov, D. A., 2014. Precise estimates of mutation rate and spectrum in yeast. *Proceedings of the National Academy of Sciences*, **111**(22):E2310–E2318.

Supplementary Information

Supplemental information for article: Chromatin structure influences rate and spectrum of spontaneous mutations in *Neurospora crassa*. By Mariana Villalba de la Peña, Pauliina A. M. Summanen, Martta Liukkonen, Ilkka Kronholm.

Supplementary methods

Mutation accumulation experiment

We started the MA-experiment with two different strains: 2489 *mat A* and 2489 *mat a*. We have previously generated these strains by backcrossing mating type *mat a* from strain 4200 into 2489 nine times (Kronholm et al., 2020). The strains have different mating types, but theoretically the strains should share over 98% of rest of their genetic background. Their Fungal Genetics Stock Center ID's are: B 26708 and B 26709. We used 20 lines for both these strains, giving 40 MA-lines in total. We used two different mating types to later have the possibility to perform crosses between the MA lines. However, for this study the mating types of the lines do not matter as all propagation was asexual.

Common protocols for culturing *N. crassa* were followed, and sorbose plates were used to induce colonial morphology on plates (Davis and de Serres, 1970). The experiment was started by picking a single colony from a sorbose plate for both ancestors and transferring that colony into a 75×12 mm test tube with flat surface of 1 mL of Vogel's Medium (VM) with 1.5% agar and 1.5% sucrose (Metzenberg, 2003). Tubes were incubated at 25 °C for 3 days to allow conidia (asexual spores) to develop. Then we picked small amount of conidia with a loop into a tube with 1.4 mL of 0.01% Tween-80, we then pipetted 1 µL of this conidial suspension into a 50 µL water droplet on a sorbose plate and spread it. We incubated the plates at room temperature for 2 days and picked single colonies to establish the MA-lines. The MA-lines were transferred the same way, so that a single colony was always picked randomly from a sorbose plate to propagate the MA-line (Figure 1B). We tested that 2 days of incubation was enough time for all colonies to appear on plates.

Combining the time of 2 days on plates and 3 days in a tube, a single transfer took 5 days. We propagated the MA lines for 40 transfers, the ancestors and the MA lines were stored frozen in suspended animation until sequencing.

Estimating the number of mitoses in the MA-experiment

To estimate mutation rate per mitosis, we needed to estimate how many mitoses happened in the MA lines during the experiment. To estimate the number of mitoses that happened during one transfer, we needed to obtain data about the number of nuclei present in each phase of a transfer: in a colony on a sorbose plate, in the mycelium in a test tube, and the conidia produced in the test tube. To estimate the density of nuclei per μm^2 of hyphae we used the strain *mat A his-3$^+$::Pccg-1-hH1$^+$-sgfp$^+$* (FGSC# 9518) which expressed a green fluorescent protein that had been fused into histone H1 (Freitag et al., 2004). We grew the strain on plates with either normal VM medium or sorbose medium, cut out a piece of the agar, and mounted it on a glass coverslip using the inverted agar block method (Lichius and Zeilinger, 2019). We used Congo Red to stain cell walls: a 20 µL droplet with 2 µM Congo Red was pipetted to a glass coverslip and an agar block with the side carrying the mycelium was placed face down in the droplet.

Samples were imaged with a Nikon A1R confocal microscope, GFP was excited with a 488 nm laser and detected with a 515/30 emission filter, Congo Red was excited with a 561 nm laser and detected with a 595/50 emission filter. Plan apochromat air objectives 20x (numerical aperture 0.75) and 40x (numerical aperture 0.95) were used. Laser power was set as low as possible to avoid saturated pixels. We imaged vertical stacks of the mycelium, and used imageJ2 (Rueden et al., 2017) to measure the area covered by hyphae in sections of the image, and counted the number of nuclei in these areas (Figure 1C).

We then estimated the number of nuclei in the different phases of a transfer, and calculated the number of mitoses that the MA-lines went through. Number of nuclei can only increase when the old nuclei divide. If we know the number of initial nuclei and the number of nuclei at time t, we

can calculate the number of mitoses, m, that separate these time points from the equation

$$m = \log_2 \left(\frac{n_t}{n_i} \right) \tag{S1}$$

where n_i is the initial number of nuclei, and n_t number of nuclei at time t. Thus, in order to estimate number of mitoses that happened in the MA lines during one transfer, we need to count how many nuclei were present in the colonies on sorbose plates that were picked and transferred to slants, and how many nuclei were in the mycelium that formed in the test tube, and finally how many nuclei were in the conidia that formed in the test tube (Figure 1B). This allows us to calculate how many mitoses happened during one transfer of the MA-experiment, from one spore to a spore. There are multiple sources of uncertainty in these calculations, so we used a Bayesian framework to do the calculations using posterior distributions of the estimates to incorporate all sources of uncertainty in the final estimate.

Nuclei were counted from the microscope images using Fiji2 version 2.0.0-rc-54/1.51g (Schindelin et al., 2012). Short sections of mycelium were surrounded with the rectangular selection tool and the area inside was measured. All nuclei with more than 50% of their diameter inside the selection were counted manually. Multiple sections were counted from each image, with no overlap. In some fainter images, the contrast was enhanced with the enhance contrast tool, with the default value 0.3% saturated pixels and no histogram equalization. To estimate the number of nuclei in a given area of hyphae, we used the counts of nuclei and hyphal areas measured from the microscope images to obtain number of nuclei per μm^2. We had images for both VM and sorbose plates, in total we collected 519 measurements. To estimate average density of nuclei for VM and sorbose

811   we used a model where we allowed standard deviations to differ for VM and sorbose media:

$$y_i \sim \mathrm{N}(\mu_i, \sigma_i) \qquad (\mathrm{S2})$$

$$\mu_i = \rho + \beta_s x_i$$

$$\log(\sigma_i) = \alpha_\sigma + \beta_\sigma x_i$$

$$\rho, \beta_s \sim \mathrm{N}(0, 0.1)$$

$$\alpha_\sigma, \beta_\sigma \sim \mathrm{hT}(3, 0, 10)$$

where y_i is the ith density measurement, ρ is the intercept, β_s is the effect of sorbose medium, x_i

an indicator variable for sorbose, α_σ is the intercept for standard deviation, and β_σ is the effect of

sorbose medium on standard deviation. Average density of nuclei in VM medium is ρ and density

of nuclei in sorbose is obtained as $\rho_s = \rho + \beta_s$.

To estimate the average size of colonies on sorbose plates, we plated conidia on sorbose plates

as in the MA-experiment and photographed the plates. Millimeter paper was used as a scale.

Colony area was measured from these images with ImageJ2 version 2.0.0-rc-43/1.50e. The pixels

per millimeter calibration value was set by measuring the number of pixels per 1 mm of millimeter

paper. The images were enhanced with the sharpen tool to make the colony outlines more distinct.

The colony area was measured using the elliptical selection tool. We used 10 different genotypes

from different MA-lines and timepoints in this experiment, including the 2 ancestors. We collected

a dataset with 482 area measurements. To estimate average colony size, we fitted a multilevel

model

$$y_i \sim \mathrm{N}(\mu_i, \sigma) \tag{S3}$$

$$\mu_i = \alpha_{g[i]}$$

$$\alpha_g \sim \mathrm{N}(\bar{\alpha}, \sigma_g)$$

$$\bar{\alpha} \sim \mathrm{N}(0, 3)$$

$$\sigma, \sigma_g \sim \mathrm{hT}(3, 0, 10)$$

where y_i is the ith area measurement, $\bar{\alpha}$ is the overall mean, α_j is the mean for jth genotype, σ_g is the genotype standard deviation, and σ is the error standard deviation. Standard deviations had a weakly informative prior, which was the half-location scale version of Student's t-distribution, where 3 is the degrees of freedom, 0 is the location, and 10 is the scale parameter. We estimated the number of nuclei in a sorbose plate colony, n_s, as

$$n_s = \bar{\alpha} \times 10^6 \times \rho_s \tag{S4}$$

the average colony size is multiplied by 10^6 to transform the unit from mm^2 to $\mathrm{\mu m}^2$.

Once the sorbose colony is transferred to the test tube, the mycelium will cover the surface of the growth media. We estimated the number of nuclei present in the mycelium, n_v by multiplying the surface of the media in the test tube with the density of nuclei in the hyphae in VM medium:

$$n_v = \pi(d/2)^2 \times 10^6 \times \rho \tag{S5}$$

where d is the diameter of the test tubes (in mm) used in the experiment, area is multiplied by 10^6 to transform the unit to $\mathrm{\mu m}^2$.

To estimate the number of conidia produced by the mycelium in the test tube, we counted conidia by suspending them in $1~\mathrm{mL}$ of 0.01% Tween-80, making a 10000-fold dilution of the

suspension, and plating 10 µL of the dilution on sorbose plates. We counted the colonies that were formed, and estimated the original number of conidia produced. We used 10 different genotypes, including the ancestors from the MA-experiment to estimate produced conidia. We collected 71 measurements, the model was

$$y_i \sim \mathrm{N}(\mu_i, \sigma) \tag{S6}$$

$$\mu_i = \nu_{g[i]}$$

$$\nu_g \sim \mathrm{N}(\bar{\nu}, \sigma_g)$$

$$\bar{\nu} \sim \mathrm{hT}(3, 40, 21)$$

$$\sigma, \sigma_g \sim \mathrm{hT}(3, 0, 21)$$

where y_i is the ith conidial number measurement, $\bar{\nu}$ is the overall mean, α_j is the mean for jth genotype, σ_g is the genotype standard deviation, and σ is the error standard deviation. Priors followed Student's t-distribution. The number of nuclei contained by the conidia, n_c was estimated as

$$n_c = 2\bar{\nu} \tag{S7}$$

since the mode of nuclei in conidia of *N. crassa* is two.

Number of mitotic divisions separating two time points can be calculated from equation S1. First, we need to calculate the number of divisions that happened when a single spore grows to a colony on sorbose plate, then the number of divisions when the colony grows to a lawn of mycelium in the test tube, and finally the number of divisions it takes to form the final number of conidia. Thus, using the posterior distributions of numbers of nuclei in the different phases of the transfer and equation S1, we can calculate the number of mitoses that happen during a transfer, m, as:

$$m = \log_2\left(\frac{n_s}{2}\right) + \log_2\left(\frac{n_s + n_v}{n_s}\right) + \log_2\left(\frac{n_s + n_v + n_c}{n_s + n_v}\right)$$

which simplifies to

$$m = \log_2\left(n_s + n_v + n_c\right) - 1 \qquad \text{(S8)}$$

this estimate of number of mitoses incorporates all sources of measurement error since posterior distributions are used in every step of the calculations.

DNA extraction

To get high quality DNA for sequencing, the natural strains, MA-lines, and the ancestors were grown in 5 mL of liquid VM for two days at 25 °C with shaking. We harvested the mycelium and freeze dried it over night in a lyophilizer. Dried mycelium was then ground with a glass bead in Qiagen Tissue Lyzer for two times 20 s with frequency of 25 s^{-1}. Then 500 μL of extraction buffer was added (10 mM Tris pH 8, 0.1 M EDTA, 150 mM NaCl, and 2% SDS), and the powdered tissue dissolved by shaking. Then samples were extracted with 750 μL of 25:24:1 Phenol:Chloroform:Isoamylalcohol and keeping the aqueous phase. We added 2 μL of RNAse A (10 mg/mL) and 50 U of RNAse I to each sample and incubated them for 1 h at 37 °C. Samples were then extracted with 750 μL of chloroform, 1 mL of 100% ethanol was added, and DNA precipitated for 1 h at −20 °C. Then DNA was pelleted with centrifugation at 4 °C, ethanol aspirated, pellet washed with 70% ethanol, and air dried. We then added 77.5 μL of TE-buffer to elute the samples and incubated at 37 °C to help dissolve the pellets. We observed that occasional small DNA fragments would remain in the samples and to remove these we did a polyethyleneglycol precipitation: we added 12.5 μL of 4 M NaCl, mixed and added 12 μL 50% PEG (P3350), mixed and precipiated DNA over night at 4 °C. DNA was then pelleted with centrifugation and supernatant aspirated, pellet was washed twice with 70% ethanol, and aspirated. Pellets were eluted to 55 μL of TE-buffer as above. DNA concentrations were measured with Qubit Broad Range Kit, and DNA quality was checked by running 2 μL of sample on an 0.8% agarose gel.

Read mapping and genotyping

To be able to map reads to the mating type locus in *mat a* strains, we included the mating type *a* region, as well as the mitochondrial genome, as additional contigs. Reads were mapped using BWA-MEM version 0.7.12-r1039 with default parameters (Li, 2013). Alignment files were sorted and indexed with samtools and read groups were added with picardtools. See table S1 for alignment metrics.

We used the GATK version 4.2.0.0 (McKenna et al., 2010) pipeline to call single nucleotide mutations (SNMs) and small indels. First, we ran Haplotypecaller for each sample individually to make a g.vcf file. Haplotypecaller was run with otherwise default parameters, emitting all sites, and in diploid mode. We then consolidated all of the samples together into a database using the GenomicsBDImport function in GATK. Samples were then jointly genotyped with the Genotype-GVCFs function to produce a vcf file with all samples.

We used wormtable version 0.1.5 (Kelleher et al., 2013) to convert the vcf file into an indexed database and then a custom python script to filter for high quality sites. For a site to be included as a candidate mutation, first we required the genotypes of the ancestor and the MA-line to differ for that site. Second, the site had to have five or more reads from both the ancestor and the sample. Third, the site had to have genotype quality greater or equal to 30 for both the ancestor and the sample, and finally sites that were called heterozygous in either the ancestor or the sample were excluded. There was also a filter that a site could not be called as a mutation if all of the MA lines had the same genotype. Sites were considered as invariant if their reference genotype quality was greater or equal to 30.

To produce the final dataset of curated mutations, we checked all candidate mutations manually by inspecting the alignments from BWA and or Haplotypecaller in IGV (Thorvaldsdóttir et al., 2013). Based on our manual inspection our filtering criteria were stringent enough, for our high coverage haploid genomes, to remove mapping errors and leave only real mutations, as only very few candidate mutations had to be rejected based on manual inspection and most mutations were unambiguous.

For genotyping SNPs in the strains sampled from natural populations, the above pipeline was used to call the genotypes. Other variants than SNPs were excluded. For a site to be included, it had to be polymorphic in the sample, with a mean read depth five or greater, genotype quality 30 or greater, and mapping quality 40 or greater across all samples. Then these same criteria were applied for each individual sample, and if a sample failed to meet the quality filters, its genotype was recorded as missing data. Heterozygous sites were excluded. Sites were also excluded if > 90% of samples had missing data. Sites were called as monomorphic if mean reference genotype quality was 30 or greater and read depth 5 of greater across all samples. Then these same criteria were applied to individual samples, genotypes were recorded as missing data if a sample did not pass the filters.

Genotyping structural variants

There are several algorithms available to detect structural variants (SVs) from short-read sequencing data. However, because this kind of data is prone to base calling and alignment errors, none of the available computational algorithms can accurately and sensitively detect all types and sizes of SVs (Kosugi et al., 2019). To overcome this limitation it is common to use several algorithms and merge their outputs to increase sensitivity and precision. First, we assessed the performance of four different SVs algorithms (DELLY, Lumpy, PINDEL and SVaba) using simulated data.

We evaluated the performance of different SV callers on simulated data created using SUR-VIVOR version 1.0.7 (Jeffares et al., 2017). SURVIVOR simulates SVs by first modifying a fasta reference file by randomly altering locations according to given parameters of length and number of different SVs types (insertions, deletions, duplications, inversions and translocations). Reads are simulated based on the modified fasta and SVs are detected using the preferred SV caller. Finally SURVIVOR compares the SVs detected against the known simulated SVs, based on this FDR and sensitivity can be calculated.

We simulated four sets with 18, 40, 50 and 120 structural variants with mutation rate of 0.001 on the reference genome (assembly NC12). The number of each type of SV simulated in each set is

presented in the table S6. In set number four we simulated 20 complex SVs in which inversions and deletions occur in the same location. For duplications the min and max length parameter was set to 100-1000 bp, for INDELs 20-500 bp, for translocations 1000-3000 bp and for inversions 600-800 bp.

The SV length distribution across our four simulated sets are very similar between each other (Figure S10), and coincide with the literature which indicates that short SVs are more common than large ones (Jeffares et al., 2017). The number of reads, the error rate and the coverage of the simulated data represent our sequenced reads. The inflated number of SV per genome is for testing purposes.

Based on the modified fasta we created 150 bp pair end reads with an error rate of 0.003% and a mean coverage of 30X using DWGSIM version 0.1.11 (Homer, 2021). Simulated reads were then aligned to the reference genome using BWA-MEM (Li, 2013) with default parameters, and SVs were called using DELLY version 0.8.7, LUMPY version 0.2.13, PINDEL version 0.2.5b9 and SVaba version 1.1.0 (Rausch et al., 2012; Layer et al., 2014; Ye et al., 2009; Wala et al., 2018). Finally we used SURVIVOR to evaluate the performance of each SV caller. The SV calls were considered correct if the simulated and detected SVs were 1) of the same type 2) on same chromosome and 3) both start and stop locations were within 50 bp. The callers that performed the best were DELLY and LUMPY as they showed high sensitivity score and low false discovery rate (FDR) score (Table S6), and they were selected to call SVs on the MA lines.

For calling SVs in the MA lines we first aligned the reads to the reference genome using BWA-MEM, excluded duplicated reads with SAMBLASTER version 0.1.26 (Faust and Hall, 2014), and extracted the discordant paired-end and split-read alignments using SAMTOOLS version 1.9 (Danecek et al., 2021). DELLY was used as indicated in the recommended workflow (Rausch et al., 2012). For LUMPY the read and insert lengths were extracted from alignment files using SAM-TOOLS and the SVs were genotyped using SVTyper version 0.7.1 (Chiang et al., 2015). To filter out SVs that were present in the ancestor we used SnpSift version 5.0e (Cingolani et al., 2012). We removed those calls with a genotype quality score lower than 30 and read depth below 10. The anal-

ysis with both callers were carried out in somatic-germline mode, considering MA line as somatic and the ancestor as the germline. The signature of a translocation are reads with discordant mate pairs, where both mates are consistantly mapped to other chromosomes for example. Translocation length was determined from the break points of these discordant reads. All of the SVs detected by each caller were manually verified by inspecting the alignment files in IGV.

Genotyping copy number variants

To evaluate the performance of copy number variant (CNV) detection algorithms, we simulated 32 CNVs using SECNVs version 2.7.1 (Xing et al., 2020), then simulated 150 bp paired end reads with an error rate of 0.03% and a mean coverage of 30X using DWGSIM. We scanned for copy number variants (CNVs) using two detection programs, CNVnator version 0.4.1 (Abyzov et al., 2011) and CNV-seq version 0.2-7 (Xie and Tammi, 2009). CNV-seq was used with default parameters while CNVnator was used with two different bin sizes, 75 and 1670. Bins of 75 bp allowed the detection of small events, while bins of 1670 bp, which is the average gene length of *N. crassa* (Galagan et al., 2003), allowed the detection larger-scale events. The merged data set performed better that any of the callers individually by showing the lowest FDR rate score of 0.482, and good sensitivity score of 0.906 (Table S7).

For genotyping CNVs in the MA lines we excluded MA line sites if the start or stop location of these where within 500 bp of any site detected in the ancestor. Also, we only retained the sites that were detected by both callers CNVnator and CNVseq (if 1000 bp or less overlapped at the start or end location). The remaining sites were manually verified by inspecting the alignment file in IGV. However, we did not find any evidence of copy number changes in the MA lines.

Chromatin modifications

To determine regions of the genome where chromatin modifications occur, chip-seq reads for H3K9me3, H3K27me3, and H3K36me2 were aligned to the reference genome using BWA-MEM, and duplicate reads were removed by Picard tools. Domains of chromatin modifications were

identified using RSEG 0.4.9 (Song and Smith, 2011). Data for centromeric regions were obtained from Smith et al. (2011) and coordinate corrections for NC12 from Wang et al. (2020). The centromeric regions are defined based on the presence of centromeric histone 3 variant: CENPA. Smith et al. (2011) collected chip-seq data against CENPA and other centromeric proteins. Centromeric sequences in *N. crassa* are composed of AT-rich sequences of degraded transposable elements. However, the repeat arrays are heterogenous due to action of RIP, making almost all sequence sufficiently unique to be able to map short reads to the genome (Smith et al., 2011).

Furthermore, we used the data of duplicated regions that were defined by Wang et al. (2020). Wang et al. (2020) identified duplicated regions using BLAST, with the criteria of at least 100 bp alignment length and at least 65% sequence identity.

Analysis of relative mutation rate for different classes

For cases where relative mutation rates were computed for different classes of mutations the model was:

$$y_i \sim \text{Poisson}(\lambda_i) \tag{S9}$$

$$\log(\lambda_i) = \log \tau_j + \alpha_{[j]}$$

$$\alpha_{[j]} \sim \text{N}(0, 10)$$

where τ_j is an offset term for class j that allows taking into account differences in the abundance of certain classes (McElreath, 2015), such as higher frequency of A's and T's than G's and C's in the genome. Priors for different predictors remained the same as in equation 1. Furthermore, if we calculate the expected number of mutations for different classes under the assumptions that all mutations in all classes are equally likely, as $\tau_j = f_j n$, where f_j is the frequency of class j and n is the total number of observed mutations, and use τ_j, the expected number of mutations, as the offset

 parameter, then $\exp(\alpha_{[j]})$ yields the relative mutation rate of class j.

Mutation rate variation across the genome

To model the effects of epigenetic domains and GC-content on mutation rate we used the following model:

$$y_i \sim \text{Poisson}(\lambda_i) \tag{S10}$$

$$\log(\lambda_i) = \log \tau_i + \alpha + \beta_{GC} x_i + \beta_{K9} d_i + \beta_{K27} g_i + \beta_C c_i + \beta_I x_i d_i$$

$$\alpha, \beta \sim \text{N}(0, 10)$$

where y_i is the number of mutations a class of i intervals contained, τ_i is the number of class i intervals in total, x_i the GC-content of those intervals, d_i indicates presence or absence of H3K9me3, g_i indicates presence or absence of H3K27me3, and c_i indicates presence or absence of centromeric region. β coefficients are the corresponding effects and α is the intercept.

We tested different linear models (Table S2) and model comparison were done using the widely applicable information criterion (WAIC) (McElreath, 2015; Vehtari et al., 2017). When we assessed how well did the mutation model predict the natural genetic variation we used the predicted mutation rates S10 as a response and θ_W calculated from a population sample of strains as a predictor in a simple regression model. Bayesian version of R^2 (Gelman et al., 2019) was used to assess the model fit.

Effects of local sequence context

To analyze effects of local base composition on mutation rate, we estimated the effects of the trinucleotides from a model that included the effects of the epigenetic domains. First, we extracted the adjacent basepairs for every point mutation. There are 64 different trinucleotides, but as we

cannot know in which strand the mutation originally occurred we grouped the trinucleotides into 32 different classes based on sequence complementarity. For example, trinucleotides ATA and TAT are complementary and were grouped. Then we calculated how many times a given trinucleotide occurs in the genome in all three reading frames. Relative mutation rate was analyzed using the following model:

$$y_i \sim \text{Poisson}(\lambda_i) \tag{S11}$$
$$\log(\lambda_i) = \log \tau_t + \beta_t x_{[}t] + \beta_{K9} d_t + \beta_{K27} g_t + \beta_C c_t$$
$$\beta \sim \text{N}(0, 10)$$

We compared different linear models (Table S4) as above. We did not include intercept in this model as we wanted estimates for all trinucleotide classes and not set one class as the intercept against which the other are compared. This does not alter any biological conclusions.

We further investigated whether how the flanking base pairs influenced the relative mutation rates of the trinucleotides. We extracted estimates of relative mutation rate for the trinucleotides from model S11 and used these as a response and predicted relative mutation rate with with identities of flanking base pairs and the mutating base. Since our estimates of relative mutation rates contain uncertainty, we included the estimated error of relative mutation rate in the model. The model was:

$$y_{obs,i} \sim \mathrm{N}(y_{est,i}, y_{sd,i}) \tag{S12}$$

$$y_{est,i} \sim \mathrm{N}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_b x_i + \beta_5 z_i + \beta_3 g_i + \beta_{I5} x_i z_i + \beta_{I3} x_i g_i$$

$$\alpha, \beta \sim \mathrm{N}(0, 10)$$

$$\sigma \sim \mathrm{hT}(3, 0, 10)$$

where $y_{obs,i}$ is the median of ith observed relative mutation rate, $y_{sd,i}$ is the observed standard deviation of ith relative mutation rate, $y_{est,i}$ is the ith estimated relative mutation rate, α is the intercept, β_b is the effect of C:G relative to A:T for the mutating base, β_5 is the effect of C:G relative to A:T for the 5' flanking base pair, β_3 is the effect of C:G relative to A:T for the 3' flanking base pair, β_{I5} is the interaction effect of 5' CG when the mutating base is C:G, and β_{I3} is the interaction effect of 3' C:G when the mutating base pair is C:G. x_i, z_i, and g_i are indicators whether the basepair is C:G. We used the half location-scale version of Student's t-distribution as a prior for standard deviation with 3 degree's of freedom, location 0, and scale 10.

Supplementary results

Accuracy of mutation calling

Estimating mutation rates and particularly estimating differences in mutation rate in different parts of the genome requires accurate mutation calls. As some regions of the genome, such as centromeric regions, may contain repetitive sequences it is important to verify that mutations are called accurately in all regions of the genome, and that no region has an excess of false positive mutations. First we examined sequencing coverage throughout the genome, GC-content does have an effect on sequencing coverage as regions of low GC can be preferentially amplified during library construction, and we observed slight elevation on normalized coverage around 35% GC (Figure

S11). However, overall we observed that sequencing coverage was rather uniform across regions of different GC-content (Figure S11). Centromeric regions and regions marked by H3K9me3, have low GC-content, and while we did observed that coverage went down in regions of $< 15\%$ GC those regions constitute a very small fraction of the genome. Next, we explored the accuracy of our mutation calls after the mutations had been called by our pipeline and manually inspected in IGV. We observed that overwhelming majority of mutations had the highest possible genotype quality score determined by the GATK pipeline (Figure S1). Median genotype quality for mutations is the highest possible value of 99, only 8.6% of mutations have genotype quality less than 80 and only 1.9% less than 50. Distribution of quality scores is similar in different regions of the genome (Figure S1). While there is slightly more mutations that have lower quality scores than 99 regions marked by H3K9me3 and centromeric regions than in euchromatic regions (Figure S1), overwhelming majority of mutations in those regions have the highest genotype quality score of 99.

If most of the mutations have genotype quality scores of 99, then what kind of confidence we have in those mutation calls? We illustrate genotype quality scores with alignments viewed in IGV that show mutations in different regions of the genome and different genotype quality scores (Figure S12, S13, S14, S15, S16, S17, S18, S19, S20). When mutations had genotype quality score of 99 they were unambiguous (Figure S12, S15, S18). When genotype qualities were around 70 mutations could still be distinguished from unambiguously, even if few reads did not support the mutation or the mutations were in repetitive regions. When mutation genotype qualities were around 45 this was usually a sign that the region had lower mapping quality due to repeats or duplications (Figure S14, S17, S20). Despite of this, even in these regions, real mutations could be distinguished from mapping errors by looking at which reads supported the mutation and which did not (Figure S17, S20).

The reason we chose to do Sanger-verification for the mutations with the lowest genotype qualities was because for mutations with genotype quality of 99, there was no doubt that these were real mutations. We reasoned that since most mutations were unambiguous, verifying a random sample

of mutations by Sanger sequencing was not efficient use of resources. While there certainly are some reasons to go for a random sample we chose to verify mutations with the lowest genotype qualities, furthermore we also wanted to focus verification efforts on complex mutations to make sure that we accurately called these rare events. We verified 23 base pair changes, of which 12 were in complex mutations and 11 as single nucleotide mutations. Of the 11 SNMs 5 were in regions marked by H3K9 (excluding centromes), 3 in centromeric regions, and 3 in euchromatin. Of the 12 base pair changes in complex mutations, 3 mutations were in H3K9 regions (5 base changes in total), 1 mutation in centromeric region (2 base pair changes), and 3 mutations in euchromatin (5 base pair changes). Therefore we confirmed point mutations by Sanger sequencing in centromeric, H3K9me3, and euchromatic regions. We confirmed all point mutations where PCR-amplification and Sanger sequencing were successful, so we never detected a false positive point mutation.

Why were the genotype qualities of the mutations so good in our experiment? There are several factors in this study that contributed excellent genotype calls. First, the ancestors for the MA-lines were derived from line 2489 (synonym OR74a), which was the strain used for the original genome project (Galagan et al., 2003). Therefore, the reference genome used for read mapping corresponds to the genome of the MA-line ancestors. This is seen in alignment metrics as 98% reads are mapped to the genome in the ancestors and MA-lines (Table S1). As such, there are likely not many reads that would erroneously map to an incorrect location because their true source of origin was missing from the reference genome. Second, as explained in the introduction, repetitive sequences tend to diverge from each other in *N. crassa* due to the action of RIP. RIP does not induce the exact same mutations to the duplications, so over time duplicated arrays, such as those often found in centromeric regions, tend to diverge from one another, to the extent that short reads can be mapped to the genome in regions where it is not often possible to the same extent in other species (Smith et al., 2011). Third, the small genome of *N. crassa* made it possible to sequence the samples to a high depth, on average over 50x in many samples (Table S1). This allowed us to discriminate between true mutations and mapping errors. With this kind of sequencing depth, sequencing errors are simply not an issue anymore and they have no impact on calling the mutations, e.g. Figure S16

shows a mutation in repetitive region that as a consequence has higher frequency of sequencing errors, but with so many reads identifying the real mutation is not a problem. Finally, *N. crassa* is haploid. Combined with high sequencing depth, this makes identifying mutations easy. The only important errors are read mapping errors that may cause some sites to appear as heterozygous. But as heterozygous sites are not expected to occur in our experiment we can filter out sites called as heterozygous. We did inspect heterozygous sites manually, as it is possible that some mutations could have been present in a heterokaryotic state (nuclei with different genotypes in the same mycelium). However, we did not find any evidence of true mutations in heterokaryotic state. Whenever sites appeared as heterozygous, multiple sites were found close together (Figure S17), indicating that read mapping errors were the more likely explanation. Because of these factors, our study differs substantially from studies that need to call heterozygous sites from data with low sequencing depth and the problem of calling genotypes correctly is of different nature.

In summary, overwhelming majority of mutations that our pipeline detected had the highest possible genotype quality of 99, and this was true in regions of the genome with potentially more repetitive and duplicated regions like in centromeric regions and regions marked by H3K9 methylation. Those mutations that had genotype quality of 99 were unambiguously real mutations. Thus, even if we would filter out every mutation with genotype quality less than 99, we would still detect the observed pattern that mutation rate was higher in regions marked by H3K9 trimethylation and in centromeric regions. Differential mutation calling in different regions of the genome cannot explain the observed results.

Simulating variation in mutation rate

Despite our very high genotype qualities, we attempted to further understand could repetitive sequences or other sequence features of heterochromatin in the *N. crassa* genome hinder our ability to correctly estimate differences in mutation rates in different regions of the genome. We simulated data under two different scenarios. First, we simulated a scenario where mutation rate was set to be higher in H3K9me3 domains, with a rate of 2×10^{-5} mutations per site, compared to the rest

of the genome, with a rate of 3×10^{-6} mutations per site. In the second scenario, we simulated a uniform mutation rate across the genome, with a rate of 2×10^{-6} mutations per site. We simulated mutations to the *N. crassa* genome using the program Mutation-Simulator (Kühl et al., 2021). We simulated 40 different MA lines for each scenario with a transition / transversion rate of 1.08. We then generated simulated reads from these simulated genomes, using DWGSM (Homer, 2021), with 30X sequencing depth and read length of 150 bp. We tried to imitate the conditions of our real sequenced data, so we set the standard deviation of the base quality scores to two and the per base sequencing error rate to 0.003. The ancestor of the MA lines was simulated by generating reads from the reference genome of *N. crassa*. To call the simulated mutations from the simulated reads, we ran the same pipeline as we used for the experimental data. Thus, we had two simulated scenarios, and for each scenario we had information about the true number of mutations that happened in the simulation, and number of mutations we called with our pipeline from the simulated read data.

In the scenario with higher mutation rate in H3K9me3 regions, we ended up with a total of 1759 mutations, of which 719 were in H3K9me3 domains, 990 in euchromatin, and 50 in unspecific domains. With our pipeline we detected a total of 1705 mutations, of which 692 were in H3K9me3 domains, 964 in euchromatin and 49 in unspecific domains. All of the called mutations were true positives. However, we failed to call 54 true mutations, that is these were false negatives. In a similar manner, in the scenario with uniform mutation rate, we ended up with a total of 3078 mutations, of which 562 were in H3K9me3 domains, 2245 in euchromatin, and 271 in unspecific domains. Our pipeline detected 2978 mutations in total, of which 535 were in H3K9 domains, 2177 in euchromatin, and 266 in unspecific domains. Again, there were no false positive calls. We failed to detect 100 mutations in this set. In general, the number of false negatives was higher in H3K9me3 regions, with proportion of false negatives 3.75% and 4.80% in H3K9me3 regions, and 2.62% and 3.02% in euchromatin in for the different and uniform mutation rate scenarios respectively.

We found that the estimated mutation rate was higher in H3K9me3 regions in the scenario where the true mutation rate was higher in H3K9me3 (Figure S21), the mutation rate ratio of H3K9me3 / euchromatin was 3.39 [3.06, 3.72]. This mutation rate ratio was not statistically different from the

one calculated from the true simulated mutations: difference was 0.28 [-0.15, 0.74], which includes zero in the interval estimate. Furthermore, when we simulated a uniform mutation rate across the genome, we found no difference among called and true datasets (Figure S21). The mutation rate ratio of H3K9me3 / euchromatin was 1.15 [1.05, 1.27], there was no statistical difference in the rate ratios between called and true simulated mutations: difference was 0.08 [-0.05, 0.23], which includes zero in the interval estimate.

With this simulation data we show that our pipeline can confidently detect a difference in mutation rates in different regions of the genome. This shows that sequence features of the H3K9me3 regions, such as repetitive sequences, do not interfere with mutation calling in a manner that would lead to gross biases in mutation rate estimates in the different domains. While simulated read data cannot capture all of the properties of real data, because of sequences missing from the reference or assembly errors, it does give us confidence that we will be able to detect a real difference in mutation rates. Moreover, since we did not observe any false positive mutations, we are confident that mutation calling cannot generate spurious results in our case. We did observed slightly higher proportions of false negative mutations in H3K9me3 regions. However, if this bias is true for real data, this would make our estimates of elevated mutation rates in H3K9me3 regions more conservative.

Re-analysis of data from Wang et al. 2020

Wang et al. (2020) estimated rate of spontaneous mutation during meiosis in *N. crassa*. During meiosis a genome defence mechanism called repeat-induced point mutation (RIP) induces mainly $C \rightarrow T$ transitions in duplicated regions of the genome resulting in a very high overall mutation rate (Wang et al., 2020). While not made explicit by Wang et al. (2020), the duplicated regions correspond almost completely to H3K9 methylated domains. In order to better compare our results for asexual mutation rate in different domains to the sexual mutation rate estimated in their study, we re-analyzed the data from Wang et al. (2020) provided in their supplementary material, and included the information about chromatin domains. Their data are comprised of mutations in sequenced tetrads, which correspond to the products of a single meiosis. We included only those

tetrads originating from crosses between non-mutant strains. This leaves 67 tetrads in the data that originate from five different crosses.

First we split the mutations to those that occurred in euchromatin and to those that occurred in H3K9 trimethylated domains. We observed that the numbers of mutations occurring in euchromatin and H3K9me3 domains for a given tetrad had very different distributions (Figure S8A), number of mutations occurring per tetrad in H3K9 domains had a very long tail. When we examined mutations per tetrad by cross, we observed that for mutations that occurred in euchromatic regions there were a median of 22 mutations per tetrad with some differences among the five crosses, but variation among tetrads from different crosses was similar (Figure S8B). However, for mutations that occurred in H3K9me3 domains, there were a median of 38 mutations per tetrad but huge variation among tetrads, even within tetrads from a single cross (Figure S8B). For example, some tetrads from the same cross had 20 to 40 mutations, while others could have hundreds. In cross E the range of mutations was from 27 in one tetrad to 1187 in another. Variation among mutations in H3K9 domains per tetrad suggest that while there probably were some genetic influences on mutation rate in the different crosses, there was substantial heterogeneity in the activation of RIP that was independent of genetic effects.

We calculated mutation rate per meiosis for the euchromatic regions of the genome using a multilevel model with cross as a random factor. The model was

$$y_i \sim \text{Poisson}(\lambda_i) \tag{S13}$$

$$\log(\lambda_i) = \bar{\alpha} + \alpha_{c[i]}$$

$$\bar{\alpha} \sim \text{N}(0, 10)$$

$$\alpha_c \sim \text{N}(0, \sigma_c)$$

$$\sigma_c \sim \text{hT}(3, 0, 10)$$

where y_i is the number of mutations in euchromatic regions in the ith tetrad, $\bar{\alpha}$ is the average inter-

cept, α_c is deviation from average intercept for each cross, and σ_c is the cross standard deviation. Prior for σ_c was the half-location scale version of Student's t-distribution, with 3 degrees of freedom, location 0, and scale 10. Based on posterior predictive checks, this model fitted the data. Mutation rate was calculated from posterior distribution of $\bar{\alpha}$ as

$$\mu = \frac{\exp(\bar{\alpha})}{N n_t} \tag{S14}$$

where N is the number of called nucleotides, and n_t is the number of tetrads. The mutation rate in euchromatic regions during sexual reproduction was 1.07 [0.6, 1.67] $\times 10^{-8}$ mutations / meiosis / bp.

The data for mutations that occurred in H3K9me3 domains are clearly overdispersed. To calculate mutation rate per meiosis for H3K9 trimethylated domains we also modelled the heterogeneity among tetrads. We fitted a gamma-poisson model, also called a negative binomial model, to the data. A gamma-poisson model allows each observation, a tetrad in our case, to have a different poisson rate allowing us to model this heterogeneity in observed rates (McElreath, 2015). We fitted a model

$$y_i \sim \text{Gamma-Poisson}(\lambda_i, \phi) \tag{S15}$$
$$\log(\lambda_i) = \bar{\alpha} + \alpha_{c[i]}$$
$$\bar{\alpha} \sim \text{N}(0, 10)$$
$$\alpha_c \sim \text{N}(0, \sigma_c)$$
$$\sigma_c \sim \text{hT}(3, 0, 10)$$
$$\phi \sim \Gamma(0.01, 0.01)$$

where y_i is the number of mutations in H3K9me3 domains in the ith tetrad, ϕ is the dispersion parameter, and other parameters were same as above. The prior for ϕ was a gamma distribution with

shape of 0.01 and scale 0.01. Posterior predictive check indicated that the model fit the data reasonably well. Then mutation rate was calculated from average intercept as above. The mutation rate in H3K9 trimethylated regions during sexual reproduction was 2.54 [0.11, 7.55] $\times 10^{-7}$ mutations / meiosis / bp. As a result of rate heterogeneity there is quite a bit of uncertainty in the estimate. The ratio of mutation rates in H3K9me3 regions over euchromatic regions was 23.7 [0.99, 76.38]. While the 95% interval of the ratio slightly overlaps one due to large uncertainly in mutation rate in H3K9me3 regions, mutation rate those regions seems higher.

We examined the spectrum of mutations that occurred for euchromatic and H3K9 methylated regions separately in the same way we did for asexual mutations. We observed that in H3K9 regions there was a substantial overrepresentation of C:G $\rightarrow$ T:A transitions due to the action of RIP (Figure S8C). However, the mutation spectra that occurred in euchromatic regions was much more similar to the one we observed during asexual reproduction in euchromatic regions. There was no difference in the relative mutation rate of C:G $\rightarrow$ T:A transitions during sexual and asexual reproduction in euchromatic regions. Some of the transversions did have different relative rates: A:T $\rightarrow$ C:G, and C:G $\rightarrow$ G:C transversions had higher rate during sexual reproduction, while C:G $\rightarrow$ A:T transversions had a lower relative mutation rate during sexual reproduction (Figure S9).

Our analysis gives somewhat different results compared to those of Wang et al. (2020), who only calculated mutation rates across the whole genome, and did not take variation among tetrads or crosses into account. We do find higher mutation rates during sexual reproduction than during asexual reproduction, suggesting that in *N. crassa* meiosis is mutagenic in addition to the RIP effect in H3K9me3 domains. However, the mutation rate per meiosis was much smaller than that estimated by Wang et al. (2020). H3K9 trimethylated regions contain mainly degraded transposable elements, and are quite gene poor. If we compare non-synonymous mutations in euchromatic and H3K9 regions, of those mutations that occurred in euchromatic regions 22.16% were non-synonymous, while only 0.17% of mutations were non-synonymous in H3K9 methylated regions. Thus, the very high mutation rate observed in H3K9 regions due to action of RIP, does not necessarily translate into a high genetic load. We suggest that mutation load during sexual reproduction

in *N. crassa* may not be as high as it has been suggested by Wang et al. (2020).

Supplementary References

Abyzov, A., Urban, A. E., Snyder, M., and Gerstein, M., 2011. CNVnator: An approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Research*, **21**(6):974–984.

Chiang, C., Layer, R. M., Faust, G. G., Lindberg, M. R., Rose, D. B., Garrison, E. P., Marth, G. T., Quinlan, A. R., and Hall, I. M., 2015. Speedseq: ultra-fast personal genome analysis and interpretation. *Nature Methods*, **12**(10):966–968.

Cingolani, P., Platts, A., Coon, M., Nguyen, T., Wang, L., Land, S., Lu, X., and Ruden, D., 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, **6**(2):80–92.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., *et al.*, 2021. Twelve years of SAMtools and BCFtools. *GigaScience*, **10**(2). giab008.

Davis, R. H. and de Serres, F. J., 1970. Genetic and microbiological research techniques for *Neurospora crassa*. *Methods in Enzymology*, **17**:79–143.

Faust, G. G. and Hall, I. M., 2014. SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics*, **30**(17):2503–2505.

Freitag, M., Hickey, P. C., Raju, N. B., Selker, E. U., and Read, N. D., 2004. GFP as a tool to analyze the organization, dynamics and function of nuclei and microtubules in *Neurospora crassa*. *Fungal Genetics and Biology*, **41**(10):897–910.

Galagan, J. E., Calvo, S. E., Borkovich, K. A., Selker, E. U., Read, N. D., Jaffe, D., FitzHugh, W.,

Ma, L.-J., Smirnov, S., Purcell, S., *et al.*, 2003. The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature*, **422**(6934):859–868.

Gelman, A., Goodrich, B., Gabry, J., and Vehtari, A., 2019. R-squared for Bayesian regression models. *The American Statistician*, **73**(3):307–309.

Homer, N., 2021. DWGSIM: Whole genome simulator for next-generation sequencing. github repository.

Jeffares, D. C., Jolly, C., Hoti, M., Speed, D., Shaw, L., Rallis, C., Balloux, F., Dessimoz, C., Bähler, J., and Sedlazeck, F. J., *et al.*, 2017. Transient structural variations have strong effects on quantitative traits and reproductive isolation in fission yeast. *Nature Communications*, **8**:14061.

Kelleher, J., Ness, R. W., and Halligan, D. L., 2013. Processing genome scale tabular data with wormtable. *BMC Bioinformatics*, **14**(1):1–5.

Kosugi, S., Momozawa, Y., Liu, X., Terao, C., Kubo, M., and Kamatani, Y., 2019. Comprehensive evaluation of structural variation detection algorithms for whole genome sequencing. *Genome Biology*, **20**(1):117.

Kronholm, I., Ormsby, T., McNaught, K. J., Selker, E. U., and Ketola, T., 2020. Marked *Neurospora crassa* strains for competition experiments and Bayesian methods for fitness estimates. *G3: Genes|Genomes|Genetics*, **10**:1261–1270.

Kühl, M., Stich, B., and Ries, D., 2021. Mutation-Simulator: fine-grained simulation of random mutations in any genome. *Bioinformatics*, **37**(4):568–569.

Layer, R. M., Chiang, C., Quinlan, A. R., and Hall, I. M., 2014. LUMPY: a probabilistic framework for structural variant discovery. *Genome Biology*, **15**(6):R84.

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:1303.3997 [q-bio.GN]*, .

Lichius, A. and Zeilinger, S., 2019. Application of membrane and cell wall selective fluorescent dyes for live-cell imaging of filamentous fungi. *JoVE*, (153):e60613.

McElreath, R., 2015. *Statistical Rethinking - A Bayesian course with examples in R and Stan*. CRC Press, New York.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., *et al.*, 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, **20**(9):1297–1303.

Metzenberg, R. L., 2003. Vogel's medium N salts: Avoiding the need for ammonium nitrate. *Fungal Genetics Newsletter*, **50**:14.

Rausch, T., Zichner, T., Schlattl, A., Stütz, A. M., Benes, V., and Korbel, J. O., 2012. DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics*, **28**(18):i333–i339.

Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T., and Eliceiri, K. W., 2017. ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*, **18**(1):529.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., *et al.*, 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods*, **9**(7):676–682.

Smith, K. M., Phatale, P. A., Sullivan, C. M., Pomraning, K. R., and Freitag, M., 2011. Heterochro-matin is required for normal distribution of *Neurospora crassa* CenH3. *Molecular and Cellular Biology*, **31**(12):2528–2542.

Song, Q. and Smith, A. D., 2011. Identifying dispersed epigenomic domains from ChIP-Seq data. *Bioinformatics*, **27**(6):870–871.

Thorvaldsdóttir, H., Robinson, J. T., and Mesirov, J. P., 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Briefings in Bioinformatics*, **14**(2):178–192.

Vehtari, A., Gelman, A., and Gabry, J., 2017. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, **27**(5):1413–1432.

Wala, J. A., Bandopadhayay, P., Greenwald, N. F., O'Rourke, R., Sharpe, T., Stewart, C., Schumacher, S., Li, Y., Weischenfeldt, J., Yao, X., *et al.*, 2018. SvABa: genome-wide detection of structural variants and indels by local assembly. *Genome Research*, **28**(4):581–591.

Wang, L., Sun, Y., Sun, X., Yu, L., Xue, L., He, Z., Huang, J., Tian, D., Hurst, L. D., and Yang, S., *et al.*, 2020. Repeat-induced point mutation in *Neurospora crassa* causes the highest known mutation rate and mutational burden of any cellular life. *Genome Biology*, **21**(1):142.

Xie, C. and Tammi, M. T., 2009. CNV-seq, a new method to detect copy number variation using high-throughput sequencing. *BMC Bioinformatics*, **10**(1):80.

Xing, Y., Dabney, A. R., Li, X., Wang, G., Gill, C. A., and Casola, C., 2020. SECNVs: a simulator of copy number variants and whole-exome sequences from reference genomes. *Frontiers in Genetics*, **11**:82.

Ye, K., Schulz, M. H., Long, Q., Apweiler, R., and Ning, Z., 2009. Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics*, **25**(21):2865–2871.

 Supplementary Figures

Figure S1: Distribution of genotype qualities of observed mutations given by GATK. Distributions are shown for the whole genome, euchromatin, H3K9me3 domains excluding centromeric regions, and centromeric regions.

Figure S2: A) Mutation spectra for the MA line 21, this line had an excess number of mutations. B) Ratios of relative mutation rates for line 21 / rest of the MA lines. Intervals for C:G → A:T transversions and C:G → T:A transitions barely overlap one.

Figure S3: Relative mutation rates for windows of 200 bp binned for GC-content at 2.5 percentage point intervals. Ticks on the horizontal axis are at the end points of intervals. Note that y-axis is on $\log_{10}$ scale, dashed line indicates relative mutation rate of one. Some bins did not contain any mutations, so estimates are missing for those bins. A) Euchromatic regions B) H3K9 domains C) H3K27 domains D) Centromeric regions.

Figure S4: Regression between predicted mutation rate and observed nucleotide polymorphism in a sample of strains from natural populations. Results have been calculated for different window sizes. n is the number of windows, β is the slope of the regression line, and R^2 is the Baysian R^2 value, a measure of model fit.

Figure S5: Observerved deviations of trinucleotide frequencies from expectations for different parts of the genome. Observed trinucleotide frequencies were divided by their expected frequencies based on GC-content. Dashed line shows the expected ratio of one.

Figure S6: A) Relative mutation rates for the 32 different trinucleotide classes. B) Model estimates for relative mutation rates for centromeric, H3K9 and H3K27 domains from the trinucleotide model. Estimates are medians and range shows 95% HPD intervals.

Figure S7: Mutation rate for indels in different domains relative to euchromatin. Dashed line shows relative mutation rate of one. Facets show deletions and insertions for all indels, for indels that did not occur in repeats, and for indels that occurred in repeated sequences. Estimates are medians and ranges show 95% HPD intervals. Intervals that do not overlap with one, that is, those where mutation rate is higher than in euchromatin are colored red.

Figure S8: Mutations that occurred during sexual reproduction. Data is from Wang et al. (2020). Note that y-axis scales are different in different panels. A) Distribution of number of mutations in the tetrads in euchromatin and H3K9 methylated domains. B) Number of mutations per tetrad for the different crosses. C) Spectrum of mutations for different regions of the genome. Error bars are 95% HPD intervals.

Figure S9: Ratios of relative mutation rates during meiosis over mitosis. Points are medians and range shows 95% HPD interval of the ratio. If the interval estimate is higher than one, mutation rate in meiosis is higher, if the interval estimate is lower than one, mutation rate in mitosis is higher.

Figure S10: Densities for the length of distributions of SVs simulated using survivor. The characteristics of each simulated set is specified in the supplementary table S6.

Figure S11: Sequencing coverage plotted against GC-content of the genome for the *mat A* ancestor. Other samples had similar profiles.

Figure S12: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 11. Mutation is in chromosome 2, position 4 299 675, in euchromatin. Genotype quality of the mutation is 99.

Figure S13: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 1. Mutation is in chromosome 4, position 5 520 332, in euchromatin. Genotype quality of the mutation is 75.

Figure S14: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 25. Mutation is in chromosome 1, position 7 379 443, in euchromatin. Genotype quality of the mutation is 45 as the mutation is located in a repetitive region. This mutation was confirmed by Sanger sequencing.

Figure S15: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 10. Mutation is in chromosome 4, position 903 059, in centromeric region. Genotype quality of the mutation is 99.

Figure S16: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 13. Mutation is in chromosome 1, position 3 924 538, in centromeric region. Genotype quality of the mutation is 67 as the mutation is located in a repetitive region.

Figure S17: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 25. Mutation is in chromosome 5, position 1 046 639, in centromeric region. Genotype quality of the mutation is 45 as the mutation is located in region with reduced mapping quality. Some reads that do not support the mutation map to this location. However, those reads also have other changes that are not supported by other reads. This suggest that reads not supporting the mutation are mapping errors. This mutation was confirmed by Sanger sequencing.

Figure S18: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 10. Mutation is in chromosome 5, position 597 516, in region marked by H3K9 methylation. Genotype quality is 99.

Figure S19: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 18. Mutation is in chromosome 4, position 5 657 442, in region marked by H3K9 methylation. Genotype quality is 72.

Figure S20: Screenshot of a mutation viewed in IGV. Upper track is the ancestor and lower track is MA-line 21. Mutation is in chromosome 7, position 141 198, in region marked by H3K9 methylation. Genotype quality is 45. Some reads do not support the mutation. However, those reads have other changes that suggest the read maps to the wrong place. This mutation was confirmed by Sanger sequencing.

Figure S21: H3K9me3 / euchromatin mutation rate ratio in simulated data. Estimates calculated from called mutations from the simulated reads are in red, and estimates calculated using the true simulated mutations are in blue. The two different scenarios are: mutation rate was higher in regions of the genome marked by H3K9me3, and mutation rate was uniform across the genome. Points are means and range shows 95% HPD interval of the ratio. Interval estimates overlap in both scenarios.

 Supplementary Tables

Table S1: Summary of alignment metrics for genomes used in this study. The ancestors used to start the MA-experiment were: B 26708, which is 2489 *mat A*, and B 26709, which is 2489 *mat a*. Lines L1–L20 are *mat A* and L21–L40 are *mat a*.

Natural strains				MA-lines			
Line	Number of reads	Depth	Mapped reads (%)	Line	Number of reads	Depth	Mapped reads (%)
10882	30683221	112	87.2	B 26708	15183018	55	98.5
10883	20920000	76	69.7	B 26709	18679696	68	98.3
10884	32294020	118	72.8	L1	13215012	48	98.2
10886	15265173	56	92.7	L2	15658340	57	98.2
10892	32124218	117	62.3	L3	14699884	54	98.3
10904	14718008	54	91.1	L4	16424902	60	98.2
10906	14935778	55	89.7	L5	16328002	60	97.6
10907	32213115	118	68.1	L6	15189850	55	98.3
10908	13924689	51	91.7	L7	14006386	51	98.4
10912	14829204	54	92	L8	13888092	51	98
10914	30158241	110	61.1	L9	14386986	53	98.3
10915	24410441	89	61.1	L10	15458492	56	98.6
10918	40163515	147	65.4	L11	13884490	51	98.1
10923	16231892	59	92	L12	15913482	58	98.2
10925	37031022	135	85.2	L13	20192750	74	97.8
10926	32682749	119	66	L14	17445964	64	98.1
10927	17741573	65	73.1	L15	14776588	54	98.5
10928	15048163	55	92	L16	14562124	53	98
10932	13402421	49	91.1	L17	16043040	59	97.9
10935	22304835	81	81.1	L18	14755826	54	97.7
10937	29757556	109	85.4	L19	13712542	50	98.1
10943	32838278	120	72.8	L20	18685746	68	96.9
10946	17261401	63	90.1	L21	14814794	54	98.5
10948	13835447	50	91	L22	15849832	58	98
10950	14364201	52	98	L23	17528602	64	98.3
10951	13354998	49	89.5	L24	14415798	53	97.8
10983	38349194	140	72.8	L25	14773870	54	98.1
1131	16974127	62	88.7	L26	17436754	64	98.2
1133	14540257	53	88.3	L27	14047860	51	98.2
1165	15095250	55	90	L28	16295790	59	98.3
3210	13681263	50	90.4	L29	16244750	59	98.1
3211	13908534	51	89.4	L30	19847786	72	98.1
3223	14095397	51	92.1	L31	19613222	72	97.9
3943	143993668	525	82.6	L33	33430658	122	97.9
3975	13671543	50	88.8	L34	15456254	56	97.9
4708	14310459	52	87.7	L35	14405808	53	98
4712	12602174	46	86	L36	15567642	57	97.8
4716	18336337	67	86.7	L37	15232240	56	98
4730	15588817	57	87.4	L38	14265830	52	98
4824	12921275	47	88.3	L39	16540462	60	98.1
5910	12494976	46	83.4	L40	18856392	69	98.3
6203	13266522	48	89.3				
851	12956372	47	89.7				
8783	15381794	56	88				
8790	13205097	48	90.1				
8816	15452652	56	89.6				
8819	13736103	50	87.5				
8845	15868843	58	86.9				
8850	12695261	46	89.3				
P4452	172179164	628	86.9				
P4463	21708426	79	69.7				
P4468	21129155	77	68.4				
P4471	24828717	91	72.4				
P4476	92452134	337	86.8				
P4479	22220857	81	72.5				
P4489	34624044	126	72.5				

Table S2: Model comparisons among different models of mutation rate predicted by GC-content and chromatin modifications. Model terms are different linear model parts, α is the intercept, β_{GC} is the slope effect of GC-content, β_{K9} is the effect of H3K9 domain, β_{K27} is the effect of H3K27 domain, β_C is the effect of centromeric domain, β_I is the interaction effect between GC-content and H3K9 domain, β_{I2} is the interaction effect between GC-content and centromeric domain, β_{I3} is the interaction effect between GC-content and H3K27 domain. d_i, g_i, and c_i are indicator variables, and x_i is GC-content in percentage points. WAIC = widely applicable information criterion, SE = standard error.

Model terms	WAIC	diff ($\pm$ SE)	weight
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i + \beta_{K27}g_i + \beta_Cc_i + \beta_Ix_id_i$	454.47	0 (0)	0.63
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i + \beta_{K27}g_i + \beta_Cc_i + \beta_Ix_id_i + \beta_{I3}x_ig_i$	456.86	2.39 (2.19)	0.19
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i + \beta_Cc_i + \beta_Ix_id_i$	458.67	4.2 (6.94)	0
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i$	458.84	4.37 (10.59)	0
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i + \beta_Cc_i + \beta_Ix_id_i + \beta_{I2}x_ic_i$	460.35	5.88 (7.17)	0
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i + \beta_Ix_id_i$	495.86	41.39 (19.86)	0
$\alpha + \beta_{GC}x_i + \beta_{K9}d_i$	496.65	42.18 (22.43)	0
$\alpha + \beta_{GC}x_i$	546.62	92.15 (33.84)	0
$\alpha + \beta_{K9}d_i + \beta_Cc_i$	614.83	160.36 (42.53)	0
$\alpha + \beta_{K9}d_i$	645.82	191.35 (49.19)	0
$\alpha + \beta_Cc_i$	1290.02	835.55 (255.52)	0
α	1689.56	1235.09 (264.15)	0

Table S3: Model estimates for a model predicting mutation rate by GC-content, centromeric, H3K9, and H3K27 domains, α is the intercept, β_{GC} is the slope effect of GC-content, β_{K9} is the effect of H3K9me domain, β_{K27} is the effect of the H3K27me3 domain, β_C is the effect of centromeric domain, and β_I is the interaction effect between GC-content and H3K9me domain.

Parameter	Estimate [95% HPDI]
α	-2.61 $[-3.34, -1.88]$
β_C	0.51 $[0.35, 0.67]$
β_{K9}	-0.14 $[-0.93, 0.63]$
β_{K27}	0.32 $[0.08, 0.55]$
β_{GC}	-0.06 $[-0.08, -0.05]$
β_I	0.02 $[0.00, 0.04]$

Table S4: Model comparison among different models mutation rate predicted by trinucleotide class and chromatin modifications. Model terms are different linear model parts. α is the intercept, β_t is a vector of effects for the 32 trinucleotide classes, β_{K9} is the effect of H3K9 domain, β_{K27} is the effect of H3K27 domain, β_C is the effect of centromeric domain, β_I is the interaction effect between trinucleotide class and H3K9 domain, β_{I2} is the interaction effect between trinucleotide class and centromeric domain. d_i, g_i, and c_i are indicator variables, and $x_{[t]}$ is the trinucleotide class. WAIC = widely applicable information criterion, SE = standard error.

Model terms	WAIC	diff ($\pm$ SE)	weight
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_{K27} g_i + \beta_C c_i$	621.31	0 (0)	0.96
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_C c_i$	627.79	6.47 (6.95)	0.04
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_{K27} g_i + \beta_C c_i + \beta_I x_{[t]} d_i$	640.35	19.04 (14.69)	0
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_C c_i + \beta_I x_{[t]} d_i$	647.17	25.85 (16.26)	0
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_C c_i + \beta_I x_{[t]} d_i + \beta_{I2} x_{[t]} c_i$	652.37	31.06 (18.55)	0
$\beta_t x_{[t]} + \beta_{K9} d_i$	693.47	72.15 (21.8)	0
$\beta_t x_{[t]} + \beta_{K9} d_i + \beta_I x_{[t]} d_i$	717.79	96.48 (26.11)	0
$\beta_{K9} d_i + \beta_C c_i$	996.07	374.76 (67.31)	0
$\beta_{K9} d_i$	1043.35	422.04 (67.64)	0
$\beta_C c_i$	1370.5	749.19 (114.3)	0
α	1776.5	1155.19 (135.53)	0
$\beta_t x_{[t]}$	1916.35	1295.03 (134.38)	0

Table S5: Natural strains with sequencing data included in this study. Strains were obtained from FGSC. 33 strains were sequenced in this study and data for 23 strains were obtained from Zhao et al. (2015).

Strain	Source	Strain	Source
10948	This study	P4452	(Zhao et al., 2015)
10886	This study	P4463	(Zhao et al., 2015)
10932	This study	P4468	(Zhao et al., 2015)
1165	This study	P4471	(Zhao et al., 2015)
8816	This study	P4476	(Zhao et al., 2015)
3223	This study	P4479	(Zhao et al., 2015)
8845	This study	10882	(Zhao et al., 2015)
10908	This study	10883	(Zhao et al., 2015)
10904	This study	10884	(Zhao et al., 2015)
851	This study	10892	(Zhao et al., 2015)
1131	This study	10907	(Zhao et al., 2015)
8850	This study	10914	(Zhao et al., 2015)
8819	This study	10915	(Zhao et al., 2015)
4708	This study	10918	(Zhao et al., 2015)
4712	This study	10925	(Zhao et al., 2015)
6203	This study	10926	(Zhao et al., 2015)
4824	This study	10927	(Zhao et al., 2015)
8783	This study	10935	(Zhao et al., 2015)
8790	This study	10937	(Zhao et al., 2015)
3975	This study	10943	(Zhao et al., 2015)
10928	This study	10983	(Zhao et al., 2015)
10912	This study	3943	(Zhao et al., 2015)
3210	This study	P4489	(Zhao et al., 2015)
10923	This study		
10950	This study		
10951	This study		
10946	This study		
3211	This study		
10906	This study		
5910	This study		
4730	This study		
1133	This study		
4716	This study		

Table S6: Detecting structural variants with different callers from simulated data.

set	Deletion	Duplication	Inversion	Translocation	Insertion	Inv-del	Total number of SV	Delly sensitivity	Delly FDR	Lumpy sensitivity	Lumpy FDR	SVaba sensitivity	SVaba FDR	Pindel sensitivity	Pindel FDR
1	1	4	4	4	5	0	18	0.90	0	0.90	0.55	0.60	0.60	0.55	0.65
2	7	15	5	8	5	0	40	0.84	0.02	0.84	0.54	0.51	0.55	0.48	0.60
3	8	10	10	15	7	0	50	0.88	0	0.88	0.54	0.64	0.52	0.38	0.61
4	6	20	20	20	14	0	100	0.54	0.33	0.54	0.60	0.36	0.77	0.28	0.94

1331

Table S7: Calling CNVs on simulated data using different approaches.

	Sensitivity score	FDR score
CNVnator (1670 bin size)	0.375	0.556
CNVnator (75 bin size)	0.968	0.797
CNV-seq	0.937	0.999
Merged data	0.906	0.482