Life history diversity and signals of trade-offs in a large group of chloroviruses

Eva J.P. Lievens*, Irina V. Agarkova, David D. Dunigan, James L. Van Etten, and Lutz Becks

*Aquatic Ecology and Evolution Group, Limnological Institute, University of Konstanz, Mainaustraße

252, 78464 Konstanz, Germany; eva.j.lievens@gmail.com

Abstract

Life cycle traits shape viral evolution through their effects on epidemiology, their diversity, and their

trade-offs. Despite their importance, characterizations of viral life history diversity are hampered by

the difficulty of measuring viral phenotypes. In this study, we overcame these challenges to describe

the life history diversity of the genus *Chlorovirus*, which contains lytic dsDNA viruses that infect

unicellular green algae. We developed efficient methods to measure adsorption rate, depolarization

probability, lysis completion probability, lysis time, burst size, specific infectivity, and mortality rate.

We used these methods to quantify the life history traits of 35 chloroviruses from three host range

groups (NC64A, Osy, and SAG viruses). The trait estimates varied between 5- and 75-fold across all

the virus strains, with most traits varying substantially both within and across groups. We then used

the largest virus group in our study, the NC64A viruses (*n* = 20), to look for correlations that could

signal trade-offs between traits. This revealed that viruses with high depolarization probabilities also

had higher burst sizes, but lower probabilities of completing lysis. We also found a strong negative

correlation between growth rate and survival. Finally, we discovered biphasic or 'tailing' decay in all

the chlorovirus groups. The nature and diversity of these traits have major implications for the

ecology and evolutionary potential of the chloroviruses, and reinforce the importance of studying

viral life histories.

Keywords

life history, phenotypic diversity, trade-offs, phycodnavirus, chlorovirus, *Chlorella*, flow virometry,

one-step growth assay, persistence assay, non-linear models

Introduction

Life histories play an important role in the ecology and evolution of microbial viruses. Viral life histories encompass 'performance' traits such as adsorption rate, lysogeny probability, progeny number, and mortality rate (performance sensu DeLong et al. 2022). The values of these traits determine a virus' population dynamics and fitness (Weitz 2016), and shape its response to selection (Abedon et al. 2003, Dessau et al. 2012, Heineman and Brown 2012, Berngruber et al. 2013). Virus populations with diverse life cycle traits have more material upon which selection can act. Conversely, trade-offs between life history traits are likely to act as evolutionary constraints (Goldhill and Turner 2014, Keen 2014). Functional constraints have been found between e.g. lysis time and burst size (Wang 2006) and growth and extracellular survival (De Paepe and Taddei 2006), leading viruses to different ecological strategies (Suttle 2007). Moreover, through their effect on host mortality, the diversity and trade-offs in viral life cycles can affect entire microbial ecosystems (Suttle 2007, Record et al. 2016, Horas et al. 2018).

Despite the importance of life history traits for microbial viruses, there are not many studies that characterize their natural diversity. Measuring viral life cycle traits requires either microscopic resolution (e.g. Timm and Yin 2012, Vincent et al. 2021, DeLong et al. 2022), which is challenging, or tracking virus concentrations over time (Hiatt 1964, Hyman and Abedon 2009), which is labor-intensive when applied to multiple strains. Thus, most studies only phenotype a few viruses at a time. Many of the studies that do analyze multiple (> 3) microbial viruses investigate the phenotypic diversity within a given host species (e.g. Baudoux and Brussaard 2005, De Paepe and Taddei 2006, Stenholm et al. 2008, Frois-Moniz 2014, Kahn Mirzaei and Nilsson 2015, Zago et al. 2015, Litt and Jaroni 2017) or viral ecotype (e.g. Kukkaro and Bamford 2009, Mojica and Brussaard 2014, Jurczak-Kurek et al. 2016, Zaburlin et al. 2017, Edwards and Steward 2018), and not within a virus clade (though see Ceyssens et al. 2011). Most of these studies also focus on a subset of the life cycle traits (e.g. Baudoux and Brussaard 2005, Stenholm et al. 2008, Kukkaro and Bamford 2009, Frois-Moniz 2014, Zago et al. 2015, Jurczak-Kurek et al. 2016, Zaburlin et al. 2017, Edwards and Steward 2018). The missing information on phenotypic diversity makes it difficult to parameterize epidemiological models and predict evolutionary responses, and is particularly striking when contrasted with our growing understanding of viral genetic diversity (DeLong et al. 2022).

In this study, we quantified the life history of a large group of chloroviruses. Chloroviruses (family *Phycodnaviridae*, genus *Chlorovirus*) are dsDNA viruses that infect unicellular 'chlorella-like' green algae (Van Etten et al. 2020). They have a lytic life cycle (Fig. 1). Upon adsorption, the virus digests a hole in the host cell wall through which the host' plasma membrane and virion internal membrane

fuse (Thiel et al. 2010, Milrot et al. 2017, Agarkova et al. 2021). Virion-associated proteins then

depolarize the host's plasma membrane, allowing viral DNA and proteins to enter (Thiel et al. 2010,

Milrot et al. 2017) and preventing infection by other chloroviruses (mutual exclusion, Greiner et al.

2009). Viral DNA is replicated in or near the nucleus, packaged into capsids in the cytoplasm, and

eventually mature virions escape by lysis (Meints et al. 1986). Virions are large (~190 nm, Yan et al.

2000, Quispe et al. 2017) and have environmentally dependent decay (Van Etten et al. 1983,

Agarkova et al. 2014). The chloroviruses are known to vary in genome content and plaque size (Van

Etten et al. 1988, Rodrigues et al. 2022), but the life cycle traits have almost exclusively been

measured in the type strain PBCV-1 (exceptions: Van Etten et al. 1988, Greiner et al. 2009, Quispe et

al. 2017, Schiano di Visconte et al. 2022).

To better understand the diversity of chlorovirus life history traits, we developed efficient methods

that combine flow cytometry, classic kinetic assays, and statistical modeling. We used these

methods to quantify the full life cycle of 35 chlorovirus strains (adsorption, depolarization,

completion with lysis time and burst, and mortality; Fig. 1), including some newly identified traits

(e.g. the robust fraction; Fig. 1). We then compared the phenotypic diversity within and across three

host range groups, and tested for signals of trade-offs between traits. This is the most thorough

exploration of chlorovirus life histories to date, and provides rare insights into the diversity of

phytoplankton viruses.

Methods

We measured the life history traits of 35 chlorovirus strains using new versions of the classic one-

step growth and persistence assays. The modified assays are higher throughput and allow

quantification of the entire life cycle (Fig. 1); they rely on the distinction between virions and

infectious virions. In what follows, viruses are described as 'virions', 'depolarizing-capable virions', or

'infectious virions', and we use the terms 'multiplicity of particles (MOP)', 'multiplicity of adsorption

(MOA)', and 'multiplicity of infection (MOI)'. These terms are defined for our purposes in Table 1.

Viruses, algae, and experimental conditions

The genus *Chlorovirus* is divided into four virus groups, which are defined by their host range and

share a high degree of protein sequence similarity (≥ 92.6 %, Jeanniard et al. 2013, see also

Rodrigues et al. 2022). Throughout this study, we treat the group level as corresponding to the

species level, within which evolution can occur (though the NC64A viruses are paraphyletic, see

below). We used viruses from three groups: NC64A viruses, which infect the ciliate endosymbionts

Chlorella variabilis NC64A and *C. variabilis* Syngen 2-3; Only Syngen viruses (Osy viruses), which

replicate in *C. variabilis* Syngen 2-3 but not *C. variabilis* NC64A; and SAG viruses, which infect the

heliozoon endosymbiont *Chlorella heliozoae* SAG 3.83. All of the described procedures took place in

the type hosts (e.g. the SAG viruses were amplified and assayed in SAG 3.83).

The algal strains NC64A, Syngen 2-3, and SAG 3.83 were stored on agar slants at 4°C and inoculated

into liquid medium before use. We used a modified version of BBM (Nichols and Bold 1965), with

ammonium chloride substituted for sodium nitrate as a nitrogen source and twice the concentration

of trace element solution 4 (first used by Frickel et al. 2016). We keep the abbreviation BBM in order

to distinguish our medium from the enriched "MBBM" typically used in this model system (e.g. Van

Etten et al. 1983). Growth was on a shaker under the standard conditions (see below). The algae

used for the described assays were in late exponential phase (~2×10^6 algae/ml in this medium).

We phenotyped 20 NC64A viruses, 7 Osy viruses, and 8 SAG viruses (Table 2). These strains were

isolated from natural ponds or streams around the world between 1981 and 2017, and have been

maintained as part of a stock collection since then. Lysates were stored at 4°C. Before starting the

assays, we refreshed and amplified the virus stocks: 0.5 ml of the original lysate was inoculated into

10 ml of 2×10^6 algae/ml solution, left on a shaker under the standard conditions (see below) until

they lysed (24 h or 48 h), and filtered through 0.2 µm. The resulting filtrates were stored at 4°C and

their virion concentration was measured by flow cytometry (see below). The filtrates with

insufficient virion concentrations were given one more round of this amplification treatment.

Algal growth and virus replication took place under the 'standard conditions', which were 20°C and

constant light. Unless otherwise specified, 'shaker' refers to an orbital shaker with diameter 10 mm

and frequency 120 rpm. 'Culture plates' always refers to 96-well flat-bottomed tissue culture plates

(Techno Plastic Products, CH); 'deep well plates' always refers to 96-well 2.2 ml PP deep well plates

(VWR International, USA). Any 4°C storage was also dark storage.

Virion quantification with flow cytometry

To quantify virions, we used a flow cytometry protocol based on Brussaard (2004) and Wen et al.

(Wen et al. 2004). Our version does not require fixation and can handle up to 96 samples at a time

with 97-99 % repeatability (data not shown). Briefly, virus samples were mixed with SYBR Green I

stain (Merck, Germany) and Tris-EDTA buffer in a PCR plate, to a final volume of 150 µl and a final

SYBR Green concentration of 0.5 X. Depending on the expected virion concentration, 7.5 or 30 µl of

virus sample was used. The samples were heated to 80°C for 10 min, allowed to cool for 15 min, and

intact virions were counted by a flow cytometer. We use a BD FACSVerse (BD Biosciences, USA),

which has an automated loader and records the analyzed volume. Each sample was run for 30 s (40-

50 µl), and the virion population was identified based on its side scatter and green fluorescence. A

detailed protocol is available on protocols.io (Lievens 2022a).

Modified one-step growth (mOSG) assay

In a classic one-step growth assay, viruses and hosts are mixed at a given MOI and allowed to adsorb

for a short time. The solutions are then diluted to synchronize the infections, and the increase in

plaque-forming units is used to calculate the lysis time and burst size (Hyman and Abedon 2009).

Our modified one-step growth (mOSG) differed in two ways. First, we worked with virions, not

infectious virions: we inoculated based on MOP, and measured the increase in virion concentration

(cf. Brown and Bidle 2014). Second, we varied the MOP across replicates (cf. Brown and Bidle 2014,

Taylor et al. 2018). These differences allowed us to calculate the depolarization probability (d) in

addition to the adsorption constant (k), lysis time ($\mu_l \pm \sigma_l$), and burst size (b_d) (Fig. 1).

The mOSG assay is represented schematically for one virus strain in Fig. 2; a detailed protocol is

available on protocols.io (Lievens 2022b). We combined algae and viruses in culture plates. Each well

contained 0.1 ml solution with 1.5×10^7 algae/ml and either MOP 0.5, 1, 2, 5, or 10 of a given virus

filtrate (MOP 20 was included for a few strains where there was room on the plates). The virus

volumes were pipetted separately, so that the different MOPs were statistically independent.

Immediately after combining algae and viruses, the solutions were mixed for 15 min to induce

adsorption (1200 rpm on a microplate shaker with orbital diameter 1 mm). The infections were then

synchronized by 1,000-fold dilution into nine sets of deep well plates (2 µl solution + 1998 µl BBM).

The deep well plates were sealed with transparent foil and left under the standard conditions; they

were inverted every hour to prevent sedimentation. At time points 0, 2, 4, …, and 16 h after dilution,

we collected a set of deep well plates for sampling. These independent collections maximized the

statistical independence of the data. We also prepared one deep well plate containing a 1 : 10,000

dilution, whose purpose was to check for signals of secondary infection in the growth curves (not

shown in Fig. 2; see Supp. Fig. S1M). This plate was sampled after 16 h. Sampling was done as

follows: deep well plates were centrifuged for 15 min at 2000 g to separate virions from algae;

supernatants (containing virions) were transferred into culture plates, sealed, and stored at 4°C. The

virion concentration in the supernatants was quantified by flow cytometry within one week after

sampling.

We measured the concentration of free virions at each time point. For each virus strain, we modeled

this concentration as the sum of leftover unadsorbed virions and newly released progeny virions:

$V(t) = \frac{1}{\delta} * \left(A_a * M * e^{-k*A_a*t_a} + A_a * \left(1 - e^{-d*(1-e^{-k*A_a*t_a})*M} \right) * F\left(t; \mu_l, \sigma_l^2, \alpha\right) * b_d \right)$ (Eq. 1).

154
unadsorbed virions   *release of progeny virions*

$V(t)$ is the free virion concentration at t hours after dilution, M is the MOP, and δ is the dilution

factor. To model the concentration of unadsorbed virions, we assumed that there were enough

binding sites available for all virions (Meints et al. 1988), that all virions were capable of adsorbing,

and that virions adsorbed at a constant rate. In this case, the concentration of free virions at the end

of the adsorption period is described by the adsorption constant k, the algal concentration during

the adsorption period A_a, the duration of the adsorption period in minutes t_a, and the initial virion

concentration $A_a * M$ (Hyman and Abedon 2009, Eq. 18.2). The release of progeny virions over time

depends on the concentration of depolarized cells, the burst size per depolarized cell, and the

distribution of lysis time. To model the concentration of depolarized cells at the end of the

adsorption period, we assumed that adsorbed virions were randomly distributed across host cells,

and that depolarizing-capable and depolarizing-incapable virions adsorbed at the same rate. The

concentration is then derived from the Poisson distribution with expected number of events $d *$

$\left(1 - e^{-k*A_a*t_a}\right) * M$, which is the depolarization probability d (i.e. the probability that a given virion

is depolarizing-capable) multiplied by the MOA. Next, we considered that every depolarized cell

produced b_d progeny virions, with b_d the average burst size per depolarized cell (the true burst size

probably varied across cells (Delbrück 1945, Timm and Yin 2012, Lievens et al. accepted), but

without single-cell information we cannot take this into account). Crucially, we assumed that b_d was

unaffected by MOP. This assumption was based on the finding that depolarization leads to mutual

exclusion in chloroviruses (all depolarized cells are infected by max. one virion, Greiner et al. 2009),

and was supported by single-cell burst size data for a related chlorovirus (Lievens et al. accepted).

Finally, under the simplification that all depolarized host cells produced b_d, the distribution of lysis

time could be approximated by a truncated normal distribution. The proportion of lysed cells over

time $F(t)$ is the distribution's cumulative distribution function, with mean lysis time μ_l, standard

deviation σ_l, and truncation value α. We set the truncation value α to 0 for simplicity (for the

majority of the virus strains, including α as a separate parameter did not improve the fit nor affect

the other parameter estimates, Supp. Fig. S2). The model is derived with supporting literature in

Supp. Method S1, the model parameters are summarized in Table 3, and the effects of the

parameters are demonstrated graphically in Supp. Fig. S3.

For each viral strain, we fit Eq. 1 to our data using nonlinear least squares fitting ('nls' in base R

version 3.6.1, R Core Team 2014). Function $F(t)$ was calculated with the 'EnvStats' package (Millard

2013). The data was *ln*-transformed and curated before fitting: obvious outliers were removed

(virion concentration implausibly high or low compared to the data for other MOPs at the same time

point, or compared to other time points of that MOP), as were time points with secondary infections (aberrant 1 : 1,000 vs. 1 : 10,000 dilutions data at time point 16 h, or secondary increases at later time points for low MOPs). For the fitting, all parameters were given appropriate lower limits (0 for k and d; 0.5 for σ_l; 1 for μ_l and b_d). k, μ_l, σ_l, and b_d were given unrealistically high upper limits; d was given a meaningful upper limit of 1. To ensure the parameter estimates were robust, we used a variety of initial parameters for each model (example in Supp. Fig. S1, panels A-D). The quality of the model fit was assessed by visually judging the fit and distribution of residuals (example in Supp. Fig. S1, panels I-O). Non-parametric bootstrapping was used to obtain 95 % CIs for each parameter (package 'nlstools', Baty et al. 2015 p. 20) (example in Supp. Fig. S1, panels E-H).

We tested the repeatability of the mOSG assay in separate experiments with four MOPs (0.5, 1, 5, 10) and time points up to 14 h (results shown in Supp. Fig. S4). The repeatabilities of the point estimates were 0.86 for k, 0.82 for d, 0.51 for μ_l, 0.44 for σ_l, and 0.65 for b_d (n = 10, 17, 8, 8, and 16 strains, respectively, each strain tested twice; calculated following Nakagawa and Schielzeth 2010). When an effect of the testing day was included to account for environmental variation, repeatability increased to 0.76 for μ_l, 0.57 for σ_l, and 0.83 for b_d. Note that we used more MOPs, more time points, and a single testing day in this mOSG assay, so our power was certainly higher.

Modified persistence (mP) assay

A classic persistence assay (also called survival, inactivation, or decay assay) exposes a virus solution to an environment and measures the decline in infectious virions over time. At each time point, the concentration of infectious virions is estimated via serial dilution methods, e.g. plaque or most probable number assays (e.g. Suttle and Chan 1994, De Paepe and Taddei 2006). In contrast, our modified persistence (mP) assay started with solutions with known virion concentrations, allowing us to estimate the initial frequency of infectious virions (*s*, the specific infectivity) as well as their mortality rate (*m*) (Fig. 1). The data also supported the existence of an additional trait, the robust fraction (*r*, see below) (Fig. 1).

We quantified the persistence of infectious virions under our standard conditions. The assay is represented schematically for one virus strain in Fig. 2; a detailed protocol is available on protocols.io (Lievens 2022c). First, we produced "fresh" viruses by amplifying viruses in 10 ml of 1×10^6 algae/ml solution. The amplification was started with a modest number of virions (~5×10^5 virions/ml) and incubated for 24 h on a shaker under the standard conditions. After 24 h, solutions were filtered through 0.45 µm, and the virion concentration in the filtrates was immediately quantified by flow cytometry. We then prepared virion solutions with different concentrations: in deep well plates, we combined filtrates and BBM to produce 1.5 ml solutions containing 5×10^4,

5×10^3, 5×10^2, and 5×10^1 virions/ml of a given virus strain. The filtrate volumes were pipetted

separately, so that the different dilutions were statistically independent. The solutions were mixed

and aliquoted into five sets of culture plates, which were sealed with transparent foil and left under

the standard conditions. After 0, 7, 14, 21, and 28 days, we used a set of culture plates to measure

the concentration of infectious virions in each solution: 10 μl from each solution was added to 16

culture wells containing 190 μl of 1.05×10^6 algae/ml solution (also in culture plates). These were

kept on a shaker under the standard conditions for 4 days, after which we measured the OD_{680} to

determine which culture wells contained viable algae (virus-negative) and which did not (virus-

positive). Using aliquots instead of longitudinal sampling maximized the statistical independence of

the data.

We measured the number of virus-positive vs. virus-negative wells produced at each time point. For

each virus strain, we modeled the proportion of virus-positive wells as

$P(t) = 1 - e^{-(V*s*r + V*s*(1-r)*e^{-m*t})}$ (Eq. 2).

This is a biphasic decay model, which assumes that infectious virions can be either "fragile", i.e.

susceptible to decay, or "robust", i.e. immortal. We used this model because the exponential decay

model was a poor fit for many of our virus strains (Supp. Fig. S5, Supp. Method S1). This biphasic

parametrization was the simplest extension of the exponential model, and fit our data very well. In

the model, $P(t)$ is the proportion of virus-positive wells after t days under the standard conditions,

and $V*s*r + V*s*(1-r)*e^{-m*t}$ is the expected number of infectious virions added to each

culture well. V is the initial number of virions (e.g. 10 μl of a 5×10^4 virions/ml solution = 500 virions),

s is the frequency of infectious virions when the solution is fresh, and r is the fraction of those

virions that is robust. The fragile infectious virions (fraction $1-r$) decay exponentially with

mortality rate m. The probability that at least one infectious virion was added to a well is derived

from the Poisson distribution. The model is derived with supporting literature in Supp. Method S1,

the model parameters are summarized in Table 3, and the effects of the parameters are

demonstrated graphically in Supp. Fig. S6.

For each viral strain, we fit Eq. 2 to the data using generalized nonlinear regression models, with the

number of virus-positive and -negative wells as a binomial response variable (in R version 3.6.1, R

Core Team 2014, package 'gnlm', Swihart and Lindsey 2019). The parameters s, r, and m were

inverse logit-transformed to keep them in the interval [0,1] (m can be larger than 1, but in practice

the mortality rates were much lower). To ensure the parameter estimates were robust, we used a

variety of initial parameters for each model (example in Supp. Fig. S7, panels A-C). The quality of the

model fit was assessed by visually judging the fit and distribution of residuals (example in Supp. Fig.

S7, panels G-L). Parametric bootstrapping was used to obtain 95 % CIs for each parameter (example

in Supp. Fig. S7, panels D-F).

Biphasic decay was confirmed in a separate experiment, as was the consistency of the fits for s, r,

and m (results shown in Supp. Fig. S8).

Comparison of mOSG and mP assays

By comparing the results of the mOSG and mP assays, we could derive the final traits in the life cycle:

the probability that a depolarized cell completes the life cycle (c) and the burst size per completed

cell (b_c) (Fig. 1). We define "completion" as lysis with release of virions. The completion probability c

could be derived from the specific infectivity s and the depolarization probability d (see Fig. 1):

$c = s/d$ (Eq. 3).

We could then derive the average burst size per completed cell b_c as

$b_c = b_d/c$ (Eq. 4),

because the burst size of depolarized cells that don't complete the life cycle is 0 (see Fig. 1). The

bootstraps from the mOSG and mP assays were used to calculate confidence intervals for c and b_c.

Eq. 3 combines estimates from two experiments, and should thus be approached with caution. One

difference was that the filtrates used for the mOSG assay were stored at 4°C for 2-8 days before the

assay, and the mP assay filtrates were not. However, previous experiments showed that s is

unaffected by 4°C storage for at least 9 days (J. Clot & E.J.P. Lievens, unpublished data for strain

PBCV-1), so this difference was unlikely to affect our calculations. A second difference was that the

algae were centrifuged before the mOSG experiment to concentrate them. This may have affected

the algae's susceptibility to depolarization (i.e. d) or stability when infected (i.e. c), although

previous experiments did not support an effect of centrifugation on s (i.e. on $d*c$; E.J.P. Lievens &

M. Spagnuolo, unpublished data for strain PBCV-1). If the algae were changed by centrifugation, we

would expect the strains in each virus group to be affected equally. Finally, due to Covid-19

restrictions there was a 3-month delay between the two experiments, and we had to use separate

batches of the host algae. The batches were grown from the same agar slants, however, so at most

we could expect an environmental effect on s that affected all strains within a group.

Growth assay

Finally, we added a test of relative growth to the mP assay (not shown in Fig. 2), using the "fresh"

filtrates prepared at the start of that assay. For each viral strain, fresh filtrate was combined with the

appropriate alga in a culture plate (2 wells per strain), to a final volume of 0.2 ml containing

1×10^6 algae/ml and 2,500 or 25,000 virions/ml. The plate was put on a shaker under the standard

conditions for 24 h, and then centrifuged for 15 min at 2000 g to separate algae and viruses. The

virion concentration of the supernatants was measured by flow cytometry; the relative increase in

virions/ml was used as the measure of viral growth.

Statistical analysis of the trait results

Our statistical analyses explored the diversity of the traits and looked for potential trade-offs. The

analyses excluded trait estimates from incomplete assays (d, μ_l, σ_l, b_d, c, and b_c for strains that did

not finish growing in the mOSG assay; r and m for strains that did not decay in the mP assay),

estimates that were poorly fitted by the statistical model, and estimates with uninformatively broad

CIs (criteria: CI > 0.67 for d, s, and c; CI spanning more than an order of magnitude for b_d, r, and b_c; CI

overlapping with the upper fitting constraint for k, μ_l, σ_l, and m). The retained trait estimates were

analyzed in R version 3.6.1 (R Core Team 2014).

To examine the trait diversity within and among virus groups, we fit ANOVAs to each trait measured

in the mOSG and mP assays. First, we partitioned variance among the virus groups using the full

dataset (n = 20 NC64A viruses, 8 SAG viruses, 7 Osy viruses; Table 2). Second, we used the strains

with a known phylogenetic position to compare the variance partitioning across virus groups vs.

clades (n = 14 NC64A viruses, 1 Osy virus, 6 SAG viruses; Table 2). The per-clade ANOVA accounted

for the paraphyletic nature of the NC64A viruses, which are divided into a PBCV-1 clade and an NY-

2A clade by the Osy strain OSyNE-5 (Quispe et al. 2017, Rodrigues et al. 2022). For each series of

ANOVAs, we corrected for multiple testing with Holm's method (Holm 1979). Trait k was

transformed by $\times 10^9$ for computational reasons, and traits b_d, b_c, and r were *ln*-transformed to

normalize their distributions.

We also used correlational analyses to look for signals of trade-offs between the traits. This analysis

focused on the NC64A viruses, as they were assayed in the same algal strain. First, we looked for

associations between the independent life history traits (traits marked in blue in Fig. 1 plus genome

size, see Supp. Table 1). We tested for pairwise correlations using Spearman's rank correlations,

weighted by the precision of each estimate ($1/\sqrt{}$(product of the CI breadth of trait 1 and trait 2,

package 'weights', Pasek 2021). To ensure that weights were comparable in scale, trait k was

transformed by $\times 10^8$ and trait b_c was *ln*-transformed. The correlations were corrected for multiple

testing using Holm's method (Holm 1979). We also applied a Principal Component Analysis (PCA) to

the independent traits. To deal with the missing values in our dataset, we used the 'Ipca' and

'PairCor' methods described in Dray & Josse (2015) (Ipca from package 'missMDA', Josse and Husson

2016, PairCor code and biplot extension following Podani et al. 2021). Ipca uses an iterative

algorithm to impute missing values, while PairCor uses a pairwise correlation approach to avoid

missing values. If the two missing value methods produced inconsistent results, we considered that

the PCA was not robust. Trait k was transformed by $\times 10^8$ for computational reasons, and trait b_c

was *ln*-transformed to linearize it. Second, we tested for specific constraints that have been found or

suggested in the literature (listed in Table 4). As above, we used weighted Spearman's rank

correlations and corrected for multiple testing using Holm's method.

As trade-offs can be masked by quality effects in correlational analyses (Stearns 1989), we repeated

the analyses in the previous paragraph with the 'highest quality' NC64A viruses. Our proxy for

quality was the relative growth of the viruses in the growth assay. The only methodological

differences were the linearizing transformations for the PCA: trait b_c was not *ln*-transformed, and

trait r was logit-transformed.

Results

The trait estimates and their 95 % CIs are shown in the upper panels of Fig. 3. Fig. 3 does not include

estimates from failed or incomplete assays; estimates that were excluded as a result of poor model

fitting are colored red (see Supp. Table S1 for more details). Hereafter we consider only the

'retained' trait estimates (colored black in Fig. 3, listed in Supp. Table S1).

Evaluation of model fitting

Overall, Eqs. 1 and 2 fit the data well and produced reliable trait estimates. Fewer than 10 % of the

estimates were excluded for having uninformatively broad 95 % CIs (red in Fig. 3). The estimates'

precision did depend on the parameter, with the depolarization probability d and completion

probability c tending to have large CIs compared to the plausible phenotypic range ([0,1]). In some

cases, precision was probably decreased by compensating effects between parameters: the

bootstrapped fits revealed positive correlations between parameters μ_l & σ_l, μ_l & b_d, σ_l & b_l, and s &

m, and negative correlations between parameters d & b_d and s & r (Supp. Fig. S9). These effects were

small compared to the scale of phenotypic variation, however, so they did not prevent the models

from making useful distinctions between parameters (Supp. Fig. S10). An exception occurred in the

subset of viral strains with $s > 0.05$: in this subset the fitting variability of s & m and s & r was on the

same scale as the phenotypic variability, so their correlated fits are likely to confound further

analyses (Supp. Fig. S10C&F).

Trait diversity within- and among chlorovirus groups

The virus traits were remarkably diverse. There was a 5- to 75-fold variation in retained estimates for

every trait, and the trait ranges often overlapped between chlorovirus groups (Fig. 3, lower panels).

To investigate this further, we partitioned the trait variance within and among groups using ANOVAs.

First, we partitioned variance among the virus groups (NC64A viruses, SAG viruses, and Osy viruses)

using the full dataset (Supp. Table S2). Group explained a significant proportion of the variance for

the adsorption constant k, the mean lysis time μ_l, the SD of lysis time σ_l, and the burst size per

depolarized cell b_d (proportions respectively 0.33, 0.86, 0.69, 0.57; p respectively 0.01, < 0.0001,

< 0.0001, < 0.01).

Among the viruses with a known phylogenetic position, the per-group (NC64A viruses, SAG viruses,

and Osy viruses) and per-clade (PBCV-1 clade, NY-2A clade, SAG clade, and OSy clade) ANOVAs

yielded similar variance partitioning results (Supp. Table S2). There were two notable exceptions:

clade explained four times more of the variance in specific infectivity s (0.45 vs. 0.11 per group) and

twelve times more of the variance in mortality rate m (0.37 vs. 0.03). These effects were due to the

higher s and m values in the PBCV-1 clade compared to the NY-2A clade. Clade also explained more

of the variance in burst size per completed cell b_c (0.09 vs. 0.001), but this difference is unreliable

due to the many missing values in the NY-2A clade. Only the mean lysis time μ_l and the SD of lysis

time σ_l varied significantly by group or clade (p respectively < 0.0001 and < 0.01).

It should be noted that these analyses could not distinguish between an effect of the group (i.e. the

type host) and an effect of relatedness (phylogenetic dependence, Garland et al. 1993). Indeed,

when the "known phylogenetic position" ANOVAs were rerun as phylogenetic ANOVAs, none of the

group effects were statistically significant (data not shown).

Correlations between traits in the NC64A viruses

Correlations between life cycle traits can suggest the existence of constraints or life history

strategies. We used our largest virus group, the NC64A viruses, to look for such associations.

First, we analyzed the associations among independent traits (traits marked in blue in Fig. 1 plus

genome size). Only three pairwise correlations were statistically significant (Fig. 4A and Supp. Fig.

S11, Supp. Table S3): the burst size per completed cell b_c and depolarization probability d were

negatively correlated with the completion probability c (resp. ρ = -0.89 and -0.84, p = 0.002 and 0.01)

and positively correlated with each other (ρ = 0.87, p = 0.002). When this dataset was used for PCA,

the PairCor and Ipca methods were inconsistent (data not shown), so these PCA results are not

interpreted further.

Second, we looked for additional correlations that we specifically hypothesized might occur (Fig. 4B

and Supp. Fig. S11, Supp. Table S3). Short-term virion quality (represented by the specific infectivity

s) was significantly correlated with long-term virion quality in the form of mortality rate m ($\rho = 0.83$,

$p = 0.0003$), but not in the form of the robust fraction r. Survival (represented by the mortality rate

m) was significantly correlated with reproduction in the form of relative growth ($\rho = 0.76$, $p = 0.002$),

but not with the 'multiplication rate' as in De Paepe & Taddei (2006). Finally, offspring quantity

(burst size per infected cell b_d) and quality (represented by the specific infectivity s) were not

significantly correlated.

Correlations between traits in the fastest-growing NC64A viruses

Because overall quality can confound the detection of constraints (Stearns 1989), we repeated the

analyses from the previous section on the 12 fastest-growing NC64A viruses. These were: AN69C,

CA-4A, CV-K1, CvsA1, IL-3A, IL-5-2s1, MA-1E, NE-JV-4, NYs-1, PBCV-1, WNE-10B-L2, and XZ-4A (see

Fig. 4C, Supp. Table S1).

For most pairwise comparisons of independent traits, the correlation coefficients were similar for

the restricted and full datasets (Fig. 5A, Supp. Table S3). However, only the positive correlation

between the burst size per completed cell b_c and depolarization probability d remained significant in

the restricted dataset ($\rho = 0.86$, $p = 0.02$). The PCAs of the independent traits were robust to the

missing-value method. They supported the association between burst size per completed cell b_c,

depolarization probability d, and completion probability c, and associated this axis with the

adsorption constant k (Fig. 5B). The mean and SD of lysis time μ_l and σ_l were positively associated,

and mortality rate m was negatively associated with the robust fraction r.

None of the additional correlations were significant in the fastest-growing NC64A viruses, and the

positive correlation between relative growth and mortality rate m was lost (Fig. 5C, Supp. Table S3).

As all the fastest-growing strains had specific infectivity $s > 0.05$, we did not consider the correlations

between s and mortality rate m or robust fraction r (see above & Supp. Fig. S10C&F).

Discussion

Viral life histories shape their epidemiology, ecosystem effects, and evolutionary potential. In this

study, we modified classic methods to quantify the life cycle traits of 35 chloroviruses from the

NC64A, Osy, and SAG virus groups. We found high trait diversity among these viruses, with evidence

for trade-offs between some traits (summarized in Fig. 6). We discuss these findings below, and also

highlight new insights into the chlorovirus life cycle.

The traits we measured included genome, virion, and virocell performance traits (sensu DeLong et al.

2022); we focus on their performance aspects here. The virion and virocell phenotypes were

probably mediated by interactions between the viral genotype, the host genotype, and the

environment (G_V x G_H x E interactions), which should be kept in mind when comparing traits

between groups and with other studies.

High trait diversity within and across chlorovirus groups

The first goal of our study was to explore the trait diversity of the chloroviruses, which influences

their evolvability and the range of their ecological effects. The strains we used were known to vary in

genome size and plaque size (Supp. Table S1). We can now add that they vary substantially in every

life cycle trait, both within and between virus groups (Fig. 3).

We can assume that the within-group diversity in chlorovirus traits is a consequence of genetic

variation, as the strains were tested in the same host and environment (G_H and E kept constant). This

genetic association would allow an evolutionary response to occur if selection favors a particular

phenotype, which may happen during the progression of an epidemic (Frank 1996, Bolker et al.

2010, Berngruber et al. 2013), as host densities change (Abedon et al. 2003, Heineman and Bull

2007), or in response to abiotic changes (Alizon et al. 2009, Dessau et al. 2012, Heineman and Brown

2012). The occurrence of chlorovirus evolution in nature is supported by work on NC64A, Osy, and

SAG viruses in natural ponds (Quispe et al. 2016), which showed that the frequencies of plaque size

phenotypes varied over the course of a year. Combining trait data with genomic data (e.g. Jeanniard

et al. 2013, Rodrigues et al. 2022) could allow the creation of a genotype-to-phenotype map to

further explore chlorovirus evolution (DeLong et al. 2022).

Attributing the between-group variation to genetic differences is more difficult, as the viruses were

not tested on a common host (only E kept constant). It is intriguing, however, that the SAG viruses

had shorter lysis times and slower adsorption. The SAG viruses infect *C. heliozoae*, which forms

symbioses with the heliozoan *Acanthocystis turfacea*, while the NC64A and Osy viruses infect *C.*

variabilis, which symbioses with the ciliate *Paramecium bursaria* (Van Etten et al. 2020). Predation of

the ciliate host is known to affect Osy-NE-ZA1 growth (DeLong et al. 2016, 2018, Al-Ameeli et al.

2021), and it is possible that chlorovirus traits respond to the broader ecological context of the

heliozoan or ciliate hosts.

The natural phenotypic diversity is likely to be even higher than the 5- to 75-fold range we

measured. Previous work has shown that the expression of chlorovirus traits depends on the host

genotype (G_H effects Quispe et al. 2017) and on environmental (E) effects such as light conditions

(Van Etten et al. 1983, I.V. Agarkova unpublished data), temperature (Horas et al. 2018, I.V.

Agarkova unpublished data), nutrient ratios (Clasen and Elser 2007, Cheng et al. 2015), and the

host's growth phase (Van Etten et al. 1983, E.J.P. Lievens unpublished data). It is likely that these

effects interact with virus strain (G_V x G_H and G_V x E effects) and with each other (G_H x E and

G_V x G_H x E effects), further enriching the overall phenotypic diversity. Interestingly, the relative lysis

times we found corresponded well to previous studies of multiple chloroviruses, but the relative

burst sizes did not (Supp. Table S1, Van Etten et al. 1988, Schiano di Visconte et al. 2022). This

suggests that different traits have different G_V x E effects, as found in other host-parasite systems

(Hall et al. 2017). Future research could examine the extent and functional basis of this (lack of)

plasticity.

Evidence for constraints and life history strategies in the NC64A viruses

We used our largest virus group, the NC64A viruses, to look for trait correlations that could suggest

the existence of trade-offs among life cycle traits (summarized in Fig. 6). While correlations cannot

prove or disprove that a functional constraint exists (Stearns 1989, Zera and Harshman 2001,

Agrawal et al. 2010), they are an important explorative step and allow comparisons with the existing

literature.

Among the independent life cycle traits, we found that virus strains with higher depolarization

probabilities (*d*) also had higher burst sizes per completed cell (b_c) and lower completion

probabilities (*c*). This three-trait association was detectable across the NC64A viruses and in the

high-quality subset (Fig. 4A, non-significant in Fig. 5A&B). It arises primarily from the low variation in

specific infectivity *s* and the positive correlation between *d* and b_d (Supp. Fig. S12). The association is

not counteracted by the broad confidence intervals around *d* and *c*, and is further supported by the

SAG viruses (Supp. Figs. S11 & S12). It is unclear why depolarization and the number of progeny

should come at the expense of completion. If this is a trade-off, the production of many

depolarizing-capable virions could be more error-prone or overly demanding on host resources. The

relationship with the adsorption constant *k* in the fastest-growing NC64A viruses (Fig. 5B) could fit

both possible explanations, assuming that well-adsorbing virions are more complicated or resource-

intensive to produce.

There were no other significant correlations between the independent traits. In particular, there was

no relationship between the mean lysis time μ_l and burst size per completed cell b_c, which has been

suggested as a general trade-off for lytic viruses (Goldhill and Turner 2014). Replacing b_c with the

infectious burst size ($s*b_c$) or nucleotide production per completed cell (genome size*b_c) did not

produce correlations with μ_l either (data not shown). The one-dimensional trade-off between lysis

time and burst size may be obscured by the comparative nature of the analysis, or it may not exist

for the NC64A viruses.

A clear pattern emerged, however, from the additional correlations: viruses with a lower long-term

success (higher mortality rate m) had a higher short-term success (higher specific infectivity s and

relative growth, Fig. 4B). These correlations were driven by the slowest-growing strains, which all

belonged to the NY-2A clade (AR158, MA-1D, NY-2A, NY-2B, Supp. Table S1). This finding could

reflect a trade-off between survival and growth, as found in phages (Table 4; correlations: De Paepe

and Taddei 2006, Dessau et al. 2012, confirmed trade-offs: Heineman and Brown 2012, García-

Villada and Drake 2013), with an extreme survival strategy evolving in the NY-2A clade. Observations

of the short- and long-term success of chloroviruses under more natural conditions would be

necessary to support this possibility. The occurrence of very slow-growing strains in the NY-2A clade

could also reflect an ecological difference, for example that these strains are best adapted to a host

genotype whose ecology favors very slow decay.

New insights into chlorovirus life cycles: failure to depolarize vs. failure to complete

Not all depolarized cells lead to completed infections (Quispe et al. 2017). Defective virions are a

well-known aspect of virus biology (Klasse 2015), and in chloroviruses they include empty capsids

and capsids with different membrane conformations (strain PBCV-1, Agarkova et al. 2014). The

chlorovirus' failure rate was generally high: depending on the strain, only 0.4-24 % of freshly-

produced virions were infectious (specific infectivity s, Fig. 3). Our estimates of s were lower than

those found previously, e.g. 10 % compared to 25-30 % for PBCV-1 (Van Etten et al. 1983). This

difference might be explained by the lower quality of the medium (Bratbak et al. 1998, cf. Tseng and

Myers 2014, Maat and Brussaard 2016, Bachy et al. 2018) or by a difference in sensitivity of liquid

culture vs. plaque assays. It is also possible that s was underestimated in our mP assay because wells

inoculated with very small numbers of infectious virions did not lyse within four days (especially

likely for slow-growing strains). Interestingly, we were able to separate viral failure into performance

traits: an inability to depolarize the host cell and an inability to complete the life cycle.

The depolarization probability varied widely between strains (Fig. 3). The 'depolarization' step in our

model actually includes three events: first, the host cell wall is digested by a conserved virion

protein, A561L (Agarkova et al. 2021); next, the viral and host membranes fuse, most likely due to

turgor pressure in the host cell (Thiel et al. 2010); finally, virion-associated K^+ channel proteins

depolarize the cell (Frohns et al. 2006, Thiel et al. 2010). The ability to successfully depolarize a host

is therefore dependent on specific, pre-packaged virion proteins (it is a virion trait, DeLong et al.

2022). We hypothesize that the interstrain variation in depolarization probability is caused by

variation in the efficiency or copy number of the A561L or K^+ channel proteins (K+ channel proteins

vary across strains, Frohns et al. 2006, PBCV-1 has ~60 copies of A651L, Dunigan et al. 2012).

Transcriptomic or proteomic studies on multiple virus strains should shed light on the underlying

mechanism.

Surprisingly, we find that an inability to complete the life cycle is the biggest hurdle for

chloroviruses: in all but one strain, fewer than half of the depolarized host cells produced progeny

virions (completion probability c = 0.01-0.42, Fig. 3). These low values may have been affected by an

underestimate in our measurement of the specific infectivity s (see above), which would have led to

an underestimate of c (and an overestimate of b_c). Alternately, they may have been biased by our

assumption that all virions were capable of adsorbing: if there was a "residual fraction" of non- or

slow-adsorbing virions (Gallet et al. 2012), the specific infectivity would actually be $s = P$(capable of

adsorbing)*d*c, and we would also have underestimated c (and overestimated b_c). Yet even if the

specific infectivity was twice as high, or half of the virions were incapable of adsorbing, the corrected

completion probability would fall below 50 % for most strains. Completion probability is a virocell

trait that depends on both host and virus (DeLong et al. 2022). If the virus effect is the most

important, then we expect the completion probability to reflect the proportion of virions with

inherent faults (Agarkova et al. 2014). Alternately, the interaction between virus and host could be

most important: algal cells probably vary in competence, e.g. through variation in their physiological

states (cf. Clasen and Elser 2007, Cheng et al. 2015) or expression of defense pathways (Rowe et al.

2013), and some virus strains could be more susceptible to this variation. More susceptible virus

strains could be evolutionarily mismatched with the host or susceptible to trade-offs with some

other virus trait (Goldhill and Turner 2014). As discussed above, the cost of completion probability

could be paid in depolarization quality or progeny number; if not, this does not invalidate the

possibility of a trade-off constraining completion probability.

New insights into chlorovirus life cycles: biphasic persistence

We discovered that many chloroviruses have biphasic or "tailing" persistence, with 0-10 % of the

initial infectious virions being resistant to decay in our model (robust fraction r, Fig. 1 & 3). In reality

these virions almost certainly would lose the ability to infect the host, but this was not detected in

our experiment. Biphasic decay is found in many virus species, and can be caused by several factors.

1) Changes in medium properties through time (Hiatt 1964). In this case we would have expected

the same biphasic pattern to occur across all strains, so this is not likely. 2) Adhesion to organic

particles (Agarkova et al. 2014, Pinon and Vialette 2018). In this case, we would expect to see a

larger robust fraction in the fast-growing strains, because they caused more lysis in the first step of

the mP assay, and thus produced more organic matter. We do not see this pattern (e.g. r between

0.1 and 5 % for the fast-growing NC64A viruses, Fig. 3). 3) Viral aggregation (Pinon and Vialette

2018). Chloroviruses are likely to aggregate, and different strains could vary in their tendency to do

so. 4) Two concurrent mortality processes causing differing degrees of damage (Koch 1960, Hiatt

1964, Roy et al. 1982, cf. Beck et al. 2014). If the first process (e.g. damage to the capsid) occurs

rapidly but leaves some probability of successful infection, and the second process (e.g. DNA

denaturation) occurs slowly and leads to complete inactivation, biphasic kinetics emerge. The

variation between strains could be due to variation in the stability of the two targets of inactivation;

the strains where $r = 0$ would then lack one target or have both decay at the same rate. 5) The

presence of a more resistant subpopulation with slower decay. These might be virions with fewer

production errors (Beauchemin et al. 2019), or virions with a specifically different phenotype (Hiatt

1964, Brouwer et al. 2017).

For possibilities 3-5, we could expect selection to optimize decay in response to each strain's native

environment. Selection could act synergistically on the mortality rate m and robust fraction r to

produce high vs. low-persistence types (though we don't see this pattern in the NC64A viruses, Fig.

4B). Alternatively, the robust fraction could reflect an adaptation to uncertainty in the environment.

If long-term persistence comes at a cost of short-term success (see above), then producing two

virion types could allow chloroviruses to profit from patches of concentrated algae (Frank 1996,

Berngruber et al. 2013) and to survive their seasonal environments (Long and Short 2016). In this

case, biphasic persistence could be a bet-hedging strategy (constant r) or a plastic strategy (r varying

in response to environmental cues); both types have been demonstrated for various virus traits (bet-

hedging: Gallet et al. 2012, Leggett et al. 2013, Erez et al. 2017, Chaturvedi et al. 2020, plasticity:

Bruce et al. 2021).

It is striking that we find biphasic decay in all the chlorovirus groups, because the existing work on

phytoplankton viruses typically assumes exponential decay (i.e. mortality at a constant rate; Suttle

and Chan 1994, Cottrell and Suttle 1995, Garza and Suttle 1998, Šulčius et al. 2015, Long and Short

2016, Demory et al. 2017, 2021a, 2021b). Though assuming exponential decay may be appropriate in

many cases, it seems unlikely that the chloroviruses are the only algal viruses with a robust fraction

(e.g. possible non-linearity in Tomaru et al. 2005). Finding biphasic decay in other phytoplankton

viruses could resolve some questions about their ecology, such as their ability to form 'seed banks'

that persist outside blooms (Long and Short 2016). It would also affect our understanding of aquatic

microbial communities and nutrient cycling, in which viral infection and decay play important roles

(Mojica and Brussaard 2014).

Applicability of the methods and future work

We hope that these methods will be useful to other virologists interested in life history traits. By

leveraging the distinction between MOP and MOI, the mOSG and mP assays measure more traits

than assays based on one or the other (e.g. Van Etten et al. 1983, Tomaru et al. 2005, Taylor et al.

2018). The high-throughput flow cytometry also makes them more efficient than e.g. quantification

through plaque assays. The assays should be applicable to any culturable lytic virus that can be

counted on a flow cytometer, which now includes many aquatic (Brussaard 2004) and human viruses

(Lippé 2018). Precision could be increased by increasing the sampling resolution, including more

MOPs, or replicating MOP treatments. The statistical models also can be edited to reflect life cycle

differences. For example, our models assume that burst size is constant because multiple infections

do not occur (discussed below). This is equivalent to a system where multiple infections occur, but

do not affect burst size. If multiple infections are possible and do affect burst size, the equations

could be extended to include a facilitation or competition factor. Finally, in systems where no

information is available, some inferences can be made from the relative amplitudes of the mOSG

curves. These should be related to MOP via a Poisson probability function if burst size is independent

of MOP (Supp. Fig. S13).

Further work on the chlorovirus life cycles could support or refute some aspects of our models. As

discussed above, we assumed there was no "residual fraction" during adsorption, and that robust

infectious virions never decay (Fig. 1). These are probably unrealistic assumptions, which could be

tested by longer studies of adsorption or persistence. We also assumed that all mOSG parameters

were independent of MOP. Mutual exclusion prevents coinfection in chloroviruses (Greiner et al.

2009), and we assumed that established infections were unaffected by other virions attaching to or

digesting the cell wall of the host. Our data was consistent with this assumption, but it had limited

power to detect deviations for d and b_d (Supp. Fig. S13). A recent study confirmed that burst size was

unaffected by MOP in a chlorovirus related to PBCV-1 (MOP 1 vs. 10, Lievens et al. accepted), but

previous work did find an effect of MOI on burst size in PBCV-1 (MOI 0.1-50, Van Etten et al. 1983). It

may be that these effects are restricted to very high MOPs/MOIs, when 'lysis from without' and dual

infection can occur (Tessman 1985, Chase et al. 1989). Future work may need to incorporate MOP-

dependencies, especially at higher MOPs (cf. Brown and Bidle 2014, Taylor et al. 2018).

Conclusion

A proper understanding of the ecological impact and evolutionary potential of viruses rests on a knowledge of their life cycle traits. We showed that the chloroviruses have high phenotypic diversity across their entire life history, both within and across host range groups. This suggests that they are evolutionarily flexible, although they may be constrained by e.g. a survival-reproduction trade-off or depolarization-completion trade-off. The chloroviruses also had large variation in relative growth, mortality rate, and long-term persistence capabilities, which are likely to have strong effects on their ecology. These results confirm the importance of studying viral life histories, and will hopefully prompt similar research into other virus clades.

Acknowledgements

E.J.P.L. thanks J. Pantel for feedback on the statistical modeling, J. Clot for help with the viruses, R. Hermann for transferring algae during lockdown, and the University of Konstanz Young Scholar Fund for material support. We acknowledge the flow cytometry centre (FlowKon) at the University of Konstanz for the use of flow cytometers and the expert support in instrument usage and data analysis. We also thank the editor and two anonymous reviewers for constructive feedback.

Data availability

Data and analyses are available at doi:10.5281/zenodo.6573770.

References

Abedon, S. T., P. Hyman, and C. Thomas. 2003. Experimental Examination of Bacteriophage Latent-Period Evolution as a Response to Bacterial Availability. Applied and Environmental Microbiology 69:7499–7506.

Agarkova, I., B. Hertel, X. Zhang, L. Lane, A. Tchourbanov, D. D. Dunigan, G. Thiel, M. G. Rossmann, and J. L. Van Etten. 2014. Dynamic attachment of Chlorovirus PBCV-1 to *Chlorella variabilis*. Virology 466–467:95–102.

Agarkova, I. V., L. C. Lane, D. D. Dunigan, C. F. Quispe, G. A. Duncan, E. Milrot, A. Minsky, A. Esmael, J. S. Ghosh, and J. L. Van Etten. 2021. Identification of a chlorovirus PBCV-1 protein involved in degrading the host cell wall during virus infection. Viruses 13:1–19.

Agrawal, A. A., J. K. Conner, and S. Rasmann. 2010. Tradeoffs and Negative Correlations in Evolutionary Ecology. Pages 243–268 *in* M. A. Bell, W. F. Eanes, D. J. Futuyma, and J. S. Levinton, editors. Evolution After Darwin: the First 150 Years. Sinauer Associates.

Al-Ameeli, Z. T., M. A. Al-Sammak, J. P. DeLong, S. Thomas, D. D. Dunigan, and J. L. Van Etten. 2021. Catalysis of chlorovirus production by the foraging of *Bursaria truncatella* on *Paramecia bursaria* containing endosymbiotic algae. Microbial Ecology 9:2170.

Alizon, S., A. Hurford, N. Mideo, and M. Van Baalen. 2009. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. Journal of Evolutionary Biology 22:245–59.

Bachy, C., C. J. Charlesworth, A. M. Chan, J. F. Finke, C. H. Wong, C. L. Wei, S. Sudek, M. L. Coleman, C. A. Suttle, and A. Z. Worden. 2018. Transcriptional responses of the marine green alga

Micromonas pusilla and an infecting prasinovirus under different phosphate conditions.
Environmental Microbiology 20:2898–2912.

Baty, F., C. Ritz, S. Charles, M. Brutsche, J. Flandrois, and D.-M. M. 2015. A Toolbox for Nonlinear
Regression in R: The Package nlstools. Journal of Statistical Software 66:1–21.

Baudoux, A.-C., and C. P. D. Brussaard. 2005. Characterization of different viruses infecting the
marine harmful algal bloom species *Phaeocystis globosa*. Virology 341:80–90.

Beauchemin, C. A. A., Y.-I. Kim, Q. Yu, G. Ciaramella, and J. P. DeVincenzo. 2019. Uncovering critical
properties of the human respiratory syncytial virus by combining in vitro assays and in silico
analyses. PLoS ONE 14:e0214708.

Beck, S. E., R. A. Rodriguez, K. G. Linden, T. M. Hargy, T. C. Larason, and H. B. Wright. 2014.
Wavelength dependent UV inactivation and DNA damage of adenovirus as measured by cell
culture infectivity and long range quantitative PCR. Environmental Science and Technology
48:591–598.

Belshaw, R., A. Gardner, A. Rambaut, and O. G. Pybus. 2008. Pacing a small cage: mutation and RNA
viruses. Trends in Ecology and Evolution 23:188–193.

Berngruber, T. W., R. Froissart, M. Choisy, and S. Gandon. 2013. Evolution of virulence in emerging
epidemics. PLOS Pathogens 9:e1003209.

Bolker, B. M., A. Nanda, and D. Shah. 2010. Transient virulence of emerging pathogens. Journal of
the Royal Society Interface 7:811–822.

Bratbak, G., A. Jacobsen, M. Heldal, K. Nagasaki, and F. Thingstad. 1998. Virus production in
Phaeocystis pouchetii and its relation to host cell growth and nutrition. Aquatic Microbial
Ecology 16:1–9.

Brouwer, A. F., M. C. Eisenberg, J. V. Remais, P. A. Collender, R. Meza, and J. N. S. Eisenberg. 2017.
Modeling Biphasic Environmental Decay of Pathogens and Implications for Risk Analysis.
Environmental Science and Technology 51:2186–2196.

Brown, C. M., and K. D. Bidle. 2014. Attenuation of virus production at high multiplicities of infection
in *Aureococcus anophagefferens*. Virology 466–467:71–81.

Bruce, J. B., S. Lion, A. Buckling, E. R. Westra, and S. Gandon. 2021. Regulation of prophage induction
and lysogenization by phage communication systems. Current Biology 31:5046–5051.

Brussaard, C. P. D. 2004. Optimization of procedures for counting viruses by flow cytometry. Applied
and Environmental Microbiology 70:1506–1513.

Ceyssens, P.-J., T. Glonti, A. M. Kropinski, R. Lavigne, N. Chanishvili, L. Kulakov, N. Lashki, M.
Tediashvili, and M. Merabishvili. 2011. Phenotypic and genotypic variations within a single
bacteriophage species. Virology Journal 8:134.

Chase, T. E., J. A. Nelson, D. E. Burbank, and J. L. Van Etten. 1989. Mutual Exclusion Occurs in a
Chlorella-like Green Alga Inoculated with Two Viruses. Journal of General Virology 70:1829–
1836.

Chaturvedi, S., J. Klein, N. Vardi, C. Bolovan-Fritts, M. Wolf, K. Du, L. Mlera, M. Calvert, N. J.
Moorman, F. Goodrum, B. Huang, and L. S. Weinberger. 2020. A molecular mechanism for
probabilistic bet hedging and its role in viral latency. Proceedings of the National Academy of
Sciences of the United States of America 117:17240–17248.

Cheng, Y. S., J. Labavitch, and J. S. VanderGheynst. 2015. Organic and Inorganic Nitrogen Impact
Chlorella variabilis Productivity and Host Quality for Viral Production and Cell Lysis. Applied
Biochemistry and Biotechnology 176:467–479.

Clasen, J. L., and J. J. Elser. 2007. The effect of host *Chlorella* NC64A carbon : phosphorus ratio on the

686   production of *Paramecium bursaria Chlorella* Virus-1. Freshwater Biology 52:112–122.

687 Cottrell, M. T., and C. A. Suttle. 1995. Dynamics of a lytic virus infecting the photosynthetic marine
688   picoflagellate *Micromonas pusilla*. Limnology and Oceanography 40:730–739.

689 De Paepe, M., and F. Taddei. 2006. Viruses' life history: towards a mechanistic basis of a trade-off
690   between survival and reproduction among phages. PLOS Biology 4:e193.

691 Delbrück, M. 1945. The burst size distribution in the growth of bacterial viruses (bacteriophages).
692   Journal of Bacteriology 50:131–135.

693 DeLong, J. P., Z. Al-Ameeli, G. Duncan, J. L. Van Etten, and D. D. Dunigan. 2016. Predators catalyze an
694   increase in chloroviruses by foraging on the symbiotic hosts of zoochlorellae. Proceedings of
695   the National Academy of Sciences of the United States of America 113:13780–13784.

696 DeLong, J. P., Z. Al-Ameeli, S. Lyon, J. L. Van Etten, and D. D. Dunigan. 2018. Size-dependent catalysis
697   of chlorovirus population growth by a messy feeding predator. Microbial Ecology 75:847–
698   853.

699 DeLong, J. P., M. A. Al-Sammak, Z. T. Al-Ameeli, D. D. Dunigan, K. F. Edwards, J. J. Fuhrmann, J. P.
700   Gleghorn, H. Li, K. Haramoto, A. O. Harrison, M. F. Marston, R. M. Moore, S. W. Polson, B. D.
701   Ferrell, M. E. Salsbery, C. R. Schvarcz, J. Shirazi, G. F. Steward, J. L. Van Etten, and K. E.
702   Wommack. 2022. Towards an integrative view of virus phenotypes. Nature Reviews
703   Microbiology 20:83–94.

704 Demory, D., L. Arsenieff, N. Simon, C. Six, F. Rigaut-Jalabert, D. Marie, P. Ge, E. Bigeard, S. Jacquet, A.
705   Sciandra, O. Bernard, S. Rabouille, and A.-C. Baudoux. 2017. Temperature is a key factor in
706   *Micromonas*-virus interactions. ISME Journal 11:601–612.

707 Demory, D., R. Liu, Y. Chen, F. Zhao, A. R. Coenen, Q. Zeng, and J. S. Weitz. 2021a. Linking Light-
708   Dependent Life History Traits with Population Dynamics for *Prochlorococcus* and
709   Cyanophage. mSystems 5:e00586-19.

710 Demory, D., J. S. Weitz, A.-C. Baudoux, S. Touzeau, N. Simon, S. Rabouille, A. Sciandra, and O.
711   Bernard. 2021b. A thermal trade-off between viral production and degradation drives virus-
712   phytoplankton population dynamics. Ecology Letters 24:1133–1144.

713 Dessau, M., D. Goldhill, R. L. McBride, P. E. Turner, and Y. Modis. 2012. Selective Pressure Causes an
714   RNA Virus to Trade Reproductive Fitness for Increased Structural and Thermal Stability of a
715   Viral Enzyme. PLoS Genetics 8.

716 Dray, S., and J. Josse. 2015. Principal component analysis with missing values: a comparative survey
717   of methods. Plant Ecology 216:657–667.

718 Dunigan, D. D., R. L. Cerny, A. T. Bauman, J. C. Roach, L. C. Lane, I. V. Agarkova, K. Wulser, G. M.
719   Yanai-Balser, J. R. Gurnon, J. C. Vitek, B. J. Kronschnabel, A. Jeanniard, G. Blanc, C. Upton, G.
720   A. Duncan, O. W. McClung, F. Ma, and J. L. Van Etten. 2012. *Paramecium bursaria Chlorella*
721   Virus 1 Proteome Reveals Novel Architectural and Regulatory Features of a Giant Virus.
722   Journal of Virology 86:8821–8834.

723 Edwards, K. F., and G. F. Steward. 2018. Host Traits Drive Viral Life Histories across Phytoplankton
724   Viruses. American Naturalist 191:566–581.

725 Erez, Z., I. Steinberger-Levy, M. Shamir, S. Doron, A. Stokar-Avihail, Y. Peleg, S. Melamed, A. Leavitt,
726   A. Savidor, S. Albeck, G. Amitai, and R. Sorek. 2017. Communication between viruses guides
727   lysis-lysogeny decisions. Nature 541:488–493.

728 Frank, S. A. 1996. Models of parasite virulence. Quarterly Review of Biology 71:37–78.

729 Frickel, J., M. Sieber, and L. Becks. 2016. Eco-evolutionary dynamics in a co-evolving host-virus
730   system. Ecology Letters 19:450–459.

Frohns, F., A. Käsmann, D. Kramer, B. Schäfer, M. Mehmel, M. Kang, J. L. Van Etten, S. Gazzarrini, A.
Moroni, and G. Thiel. 2006. Potassium Ion Channels of *Chlorella* Viruses Cause Rapid
Depolarization of Host Cells during Infection. Journal of Virology 80:2437–2444.
Frois-Moniz, K. 2014. Host/virus interactions in the marine cyanobacterium *Prochlorococcus*. Thesis,
Massachusetts Institute of Technology.
Furió, V., A. Moya, and R. Sanjuán. 2005. The cost of replication fidelity in an RNA virus. Proceedings
of the National Academy of Sciences of the United States of America 102:10233–10237.
Gallet, R., T. Lenormand, and I. N. Wang. 2012. Phenotypic Stochasticity Protects Lytic Bacteriophage
Populations From Extinction During The Bacterial Stationary Phase. Evolution 66:3485–3494.
García-Villada, L., and J. W. Drake. 2013. Experimental selection reveals a trade-off between
fecundity and lifespan in the coliphage Qß. Open Biology 3:130043.
Garland, T. Jr., A. W. Dickerman, C. M. Janis, and J. A. Jones. 1993. Phylogenetic analysis of
covariance by computer simulation. Systematic Biology 42:265–292.
Garza, D. R., and C. A. Suttle. 1998. The Effect of Cyanophages on the Mortality of *Synechococcus*
spp. and Selection for UV Resistant Viral Communities. Microbial Ecology 36:281–292.
Goldhill, D. H., and P. E. Turner. 2014. The evolution of life history trade-offs in viruses. Current
Opinion in Virology 8:79–84.
Greiner, T., F. Frohns, M. Kang, J. L. Van Etten, A. Käsmann, A. Moroni, B. Hertel, and G. Thiel. 2009.
Chlorella viruses prevent multiple infections by depolarizing the host membrane. Journal of
General Virology 90:2033–2039.
Hall, M. D., G. Bento, and D. Ebert. 2017. The evolutionary consequences of stepwise infection
processes. Trends in Ecology & Evolution 32:612–623.
Heineman, R. H., and S. P. Brown. 2012. Experimental Evolution of a Bacteriophage Virus Reveals the
Trajectory of Adaptation across a Fecundity/Longevity Trade-Off. PLoS ONE 7.
Heineman, R. H., and J. J. Bull. 2007. Testing optimality with experimental evolution: Lysis time in a
bacteriophage. Evolution 61:1695–1709.
Hiatt, C. W. 1964. Kinetics of the inactivation of viruses. Bacteriological Reviews 28:150–163.
Holm, S. 1979. A simple sequentially rejective multiple test procedure. Scandinavian Journal of
Statistics 6:65–70.
Horas, E. L., L. Theodosiou, and L. Becks. 2018. Why Are Algal Viruses Not Always Successful? Viruses
10:474.
Hyman, P., and S. T. Abedon. 2009. Practical Methods for Determining Phage Growth Parameters.
Pages 175–202 *in* M. R. J. Clokie and A. M. Kropinski, editors. Bacteriophages: Methods and
Protocols, Volume 1: Isolation, Characterization, and Interactions, vol. 401. Humana Press.
Jeanniard, A., D. D. Dunigan, J. R. Gurnon, I. V. Agarkova, M. Kang, J. Vitek, G. Duncan, O. W.
McClung, M. Larsen, J. M. Claverie, J. L. Van Etten, and G. Blanc. 2013. Towards defining the
chloroviruses: A genomic journey through a genus of large DNA viruses. BMC Genomics 14.
Josse, J., and F. Husson. 2016. missMDA: A Package for Handling Missing Values in Multivariate Data
Analysis. Journal of Statistical Software 70:1–31.
Jurczak-Kurek, A., T. Gąsior, B. Nejman-Faleńczyk, S. Bloch, A. Dydecka, G. Topka, A. Necel, M.
Jakubowska-Deredas, M. Narajczyk, M. Richert, A. Mieszkowska, B. Wróbel, G. Węgrzyn, and
A. Węgrzyn. 2016. Biodiversity of bacteriophages: Morphological and biological properties of
a large group of phages isolated from urban sewage. Scientific Reports 6:1–17.
Kahn Mirzaei, M., and A. S. Nilsson. 2015. Isolation of Phages for Phage Therapy: A Comparison of
Spot Tests and Efficiency of Plating Analyses for Determination of Host Range and Efficacy.

PloS ONE 10:e0118557.

Keen, E. C. 2014. Tradeoffs in bacteriophage life histories. Bacteriophage 4:e28365.

Klasse, P. J. 2015. Molecular Determinants of the Ratio of Inert to Infectious Virus Particles. Progress
in Molecular Biology and Translational Science 129:285–326.

Koch, G. 1960. Influence of assay conditions on infectivity of heated poliovirus. Virology 12:601–603.

Kukkaro, P., and D. H. Bamford. 2009. Virus–host interactions in environments with a wide range of
ionic strengths. Environmental Microbiology Reports 1:71–77.

Leggett, H. C., R. Benmayor, D. J. Hodgson, and A. Buckling. 2013. Experimental evolution of adaptive
phenotypic plasticity in a parasite. Current Biology 23:139–142.

Lievens, E. J. P. 2022a. Virion quantification by flow cytometry, without fixation or freezing.
dx.doi.org/10.17504/protocols.io.6qpvr6q93vmk/v1.

Lievens, E. J. P. 2022b. Modified one-step growth (mOSG) assay.
dx.doi.org/10.17504/protocols.io.4r3l2op9pv1y/v2.

Lievens, E. J. P. 2022c. Modified persistence (mP) assay.
dx.doi.org/10.17504/protocols.io.81wgb6knylpk/v2.

Lievens, E. J. P., M. Spagnuolo, T. Réveillon, and L. Becks. accepted. Delayed lysis time at high
multiplicities of particles in a chlorovirus-*Chlorella* interaction.

Lippé, R. 2018. Flow Virometry: A Powerful Tool To Functionally Characterize Viruses. Journal of
Virology 92:e01765-17.

Litt, P. K., and D. Jaroni. 2017. Isolation and Physiomorphological Characterization of *Escherichia coli*
O157:H7-Infecting Bacteriophages Recovered from Beef Cattle Operations. International
Journal of Microbiology 2017:e7013236.

Long, A. M., and S. M. Short. 2016. Seasonal determinations of algal virus decay rates reveal
overwintering in a temperate freshwater pond. ISME Journal 10:1602–1612.

Maat, D. S., and C. P. D. Brussaard. 2016. Both phosphorus- and nitrogen limitation constrain viral
proliferation in marine phytoplankton. Aquatic Microbial Ecology 77:87–97.

Meints, R. H., D. E. Burbank, J. L. Van Etten, and D. T. A. Lamport. 1988. Properties of the chlorella
receptor for the virus PBCV-1. Virology 164:15–21.

Meints, R. H., K. Lee, and J. L. Van Etten. 1986. Assembly site of the virus PBCV-1 in a *Chlorella*-like
green alga: ultrastructural studies. Virology 154:240–245.

Millard, S. P. 2013. EnvStats: An R Package for Environmental Statistics. Springer, New York.

Milrot, E., E. Shimoni, T. Dadosh, K. Rechav, T. Unger, J. L. Van Etten, and A. Minsky. 2017. Structural
studies demonstrating a bacteriophage-like replication cycle of the eukaryote-infecting
Paramecium bursaria chlorella virus-1. PLOS Pathogens 13:e1006562.

Mojica, K. D. A., and C. P. D. Brussaard. 2014. Factors affecting virus dynamics and microbial host–
virus interactions in marine environments. FEMS Microbiology Ecology 89:495–515.

Nakagawa, S., and H. Schielzeth. 2010. Repeatability for Gaussian and non-Gaussian data: a practical
guide for biologists. Biological Reviews 85:935–956.

Nichols, H. W., and H. C. Bold. 1965. *Trichosarcina polymorpha* Gen. et Sp. Nov. Journal of Phycology
1:34–38.

Pasek, J. 2021. weights: Weighting and Weighted Statistics, version 1.0.4.

Pinon, A., and M. Vialette. 2018. Survival of viruses in water. Intervirology 61:214–222.

Podani, J., T. Kalapos, B. Barta, and D. Schmera. 2021. Principal component analysis of incomplete
data – A simple solution to an old problem. Ecological Informatics 61:101235.

Quispe, C. F., A. Esmael, O. Sonderman, M. McQuinn, I. Agarkova, M. Battah, G. A. Duncan, D. D.

Dunigan, P. L. Timothy, C. DeCastro, I. Speciale, F. Ma, and J. L. Van Etten. 2017.
Characterization of a New Chlorovirus Type with Permissive and Non-permissive Features on
Phylogenetically Related Algal Strains. Virology 500:103–113.

Quispe, C. F., O. Sonderman, A. Seng, B. Rasmussen, G. Weber, C. Mueller, D. D. Dunigan, and J. L.
Van Etten. 2016. Three-year survey of abundance, prevalence and genetic diversity of
chlorovirus populations in a small urban lake. Archives of Virology 161:1839–1847.

R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Record, N. R., D. Talmy, and S. Våge. 2016. Quantifying tradeoffs for marine viruses. Frontiers in
Marine Science 3:251.

Rodrigues, R. A. L., V. F. Queiroz, J. Ghosh, D. D. Dunigan, and J. L. Van Etten. 2022. Functional
Genomic Analyses Reveal an Open Pan-genome for the Chloroviruses and a Potential for
Genetic Innovation in New Isolates. Journal of Virology 96:e01367-21.

Rowe, J. M., D. D. Dunigan, G. Blanc, J. R. Gurnon, Y. Xia, and J. L. Van Etten. 2013. Evaluation of
higher plant virus resistance genes in the green alga, *Chlorella variabilis* NC64A, during the
early phase of infection with *Paramecium bursaria* chlorella virus-1. Virology 442:101–113.

Roy, D., E. S. K. Chian, and R. S. Engelbrecht. 1982. Mathematical model for enterovirus inactivation
by ozone. Water Research 16:667–673.

Schiano di Visconte, G., M. J. Allen, and A. Spicer. 2022. New Insights from the High-Resolution
Monitoring of Microalgae–Virus Infection Dynamics. Viruses 14:466.

Stearns, S. C. 1989. Trade-Offs in Life-History Evolution. Functional Ecology 3:259–268.

Stenholm, A. R. nnest, I. Dalsgaard, and M. Middelboe. 2008. Isolation and Characterization of
Bacteriophages Infecting the Fish Pathogen *Flavobacterium psychrophilum*. Applied and
Environmental Microbiology 74:4070–4078.

Šulčius, S., E. Šimoliūnas, J. Staniulis, J. Koreivienė, P. Baltrušis, R. Meškys, and R. Paškauskas. 2015.
Characterization of a lytic cyanophage that infects the bloom-forming cyanobacterium
Aphanizomenon flos-aquae. FEMS Microbiology Ecology 91:1–7.

Suttle, C. A. 2007. Marine viruses - Major players in the global ecosystem. Nature Reviews
Microbiology 5:801–812.

Suttle, C. A., and A. M. Chan. 1994. Dynamics and distribution of cyanophages and their effect on
marine *Synechococcus* spp. Applied and Environmental Microbiology 60:3167–3174.

Swihart, B., and J. Lindsey. 2019. gnlm: Generalized Nonlinear Regression Models, version 1.1.1.

Taylor, B. P., J. S. Weitz, C. P. D. Brussaard, and M. G. Fischer. 2018. Quantitative Infection Dynamics
of *Cafeteria Roenbergensis* Virus. Viruses 10:468.

Tessman, I. 1985. Genetic recombination of the DNA plant virus PBCV1 in a *Chlorella*-like alga.
Virology 145:319–322.

Thiel, G., A. Moroni, D. D. Dunigan, and J. L. Van Etten. 2010. Initial Events Associated with Virus
PBCV-1 Infection of *Chlorella* NC64A. Progress in Botany 71:169–183.

Timm, A., and J. Yin. 2012. Kinetics of virus production from single cells. Virology 424:11–17.

Tomaru, Y., H. Tanabe, S. Yamanaka, and K. Nagasaki. 2005. Effects of temperature and light on
stability of microalgal viruses, HaV, HcV and HcRNAV. Plankton Biology and Ecology 52:1–6.

Tseng, M., and J. H. Myers. 2014. The relationship between parasite fitness and host condition in an
insect -virus system. PLoS ONE 9:e106401.

Van Etten, J. L., I. V. Agarkova, and D. D. Dunigan. 2020. Chloroviruses. Viruses 12:20.

Van Etten, J. L., D. E. Burbank, Y. Xia, and R. H. Meints. 1983. Growth cycle of a virus, PBCV-1, that

866     infects *Chlorella*-like algae. Virology 126:117–125.

867 Van Etten, J. L., A. M. Schuster, and R. H. Meints. 1988. Viruses of eukaryotic *Chlorella*-like algae.

868     Pages 411–428 *in* Y. Koltin and M. J. Leibowitz, editors. Viruses of Fungi and Simple

869     Eukaryotes. Marcel Dek. New York and Basel.

870 Vincent, F., U. Sheyn, Z. Porat, D. Schatz, and A. Vardi. 2021. Visualizing active viral infection reveals

871     diverse cell fates in synchronized algal bloom demise. Proceedings of the National Academy

872     of Sciences of the United States of America 118:e2021586118.

873 Wang, I.-N. 2006. Lysis Timing and Bacteriophage Fitness. Genetics 172:17–26.

874 Weitz, J. 2016. Quantitative Viral Ecology: Dynamics of Viruses and Their Microbial Hosts. Princeton

875     University Press.

876 Wen, K., A. C. Ortmann, and C. A. Suttle. 2004. Accurate estimation of viral abundance by

877     epifluorescence microscopy. Applied and Environmental Microbiology 70:3862–3867.

878 Yan, X., N. H. Olson, J. L. Van Etten, M. Bergoin, M. G. Rossmann, and T. S. Baker. 2000. Structure and

879     assembly of large lipid-containing dsDNA viruses. Nature Structural Biology 7:101–103.

880 Zaburlin, D., A. Quiberoni, and D. Mercanti. 2017. Changes in Environmental Conditions Modify

881     Infection Kinetics of Dairy Phages. Food and Environmental Virology 9:270–276.

882 Zago, M., B. Bonvini, L. Rossetti, A. Meucci, G. Giraffa, and D. Carminati. 2015. Biodiversity of

883     *Lactobacillus helveticus* bacteriophages isolated from cheese whey starters. Journal of Dairy

884     Research 82:242–247.

885 Zera, A. J., and L. G. Harshman. 2001. The Physiology of Life History Trade-Offs in Animals. Annual

886     Review of Ecological Systems 32:95–126.

 Tables

890 **Table 1. Definitions used in this paper.**

Term	Our definition	Ratio to host cells
Virions	Virus particles that can be detected by flow cytometry (excludes e.g.DNA-free capsids[1]).	Multiplicity of particles (MOP)
Adsorbed virions	Virions that are adsorbed to a host cell. Virions that have not adsorbed may be incapable or may have failed to adsorb by chance.	Multiplicity of adsorption (MOA)[2]
Depolarizing-capable virions	Virions that can depolarize a host cell if they adsorb to one. We assume that chance plays no role in depolarization success: failure to depolarize after adsorption is always due to incapability or mutual exclusion (the cell has already been depolarized by another depolarizing-capable virion[3]).	-
Infectious virions	Virions that can catalyze the death of an algal population, i.e. finish a life cycle and spread to other cells. Non-infectious virions can be incapable of adsorbing, depolarizing the host[1,4], or completing the infection[5].	Multiplicity of infection (MOI)

891 [1](Agarkova et al. 2014), [2](Hyman and Abedon 2009), [3](Greiner et al. 2009), [4](Frohns et al. 2006), [5](Quispe et al.
892 2017)

893 **Table 2. List of viral strains.** We used two lysates of reference strain AR158; lysate AR158[1] was seven

894 years older than AR158[2]. Phylogenetic positions based on Quispe et al. (2017) and Rodrigues et al.

895 (2022).

Group	Clade	Strains
NC64A viruses	PBCV-1 clade	AN69C, CV-K1 (= CviKI), CvsA1, IL-3A, KS-1B, MA-1E, NE-JV-4, PBCV-1
	NY-2A clade	AR158[1], AR158[2], IL-5-2s1, MA-1D, NY-2A, NY-2B, NYs-1
	unknown	CA-4A, SH-6A, WNE-10B-L2, WNE-11A-L2 (= WNE-11A-L1), XZ-4A
Osy viruses	Osy clade	OSyNE-5
	unknown	OSyNE-4B-L2, OSyNE-4B-M2, OSyNE-4B-S2, OSyNE-5B-M2, OSyNE-5B-S1, OSyNE-ZA-1
SAG viruses	SAG clade	Canal-1, GM0701.1, MN0810.1, MO0605SPH, NTS-1, TN603.4.2
	unknown	NES-4A-S1, NES-5A-L1

897 **Table 3. List of life history traits.**

Model	Parameter	Trait
Eq. 1	k	adsorption constant[1]
Eq. 1	d	depolarization probability (probability that a virion is depolarizing-capable)
Eq. 3	c	completion probability (probability that a depolarized cell will lyse and produce progeny virions)
Eq. 1	α (set to 0)	truncation value of lysis time (earliest possible lysis time)
Eq. 1	μ_l	mean lysis time
Eq. 1	σ_l	standard deviation of lysis time
Eq. 1	b_d	average burst size per depolarized cell
Eq. 4	b_c	average burst size per completed cell
Eq. 2	s	specific infectivity (proportion of virions that are infectious when the solution is fresh)
Eq. 2	r	robust fraction (fraction of initially infectious virions that do not decay)
Eq. 2	m	mortality rate of the fragile infectious virions

898 [1](Hyman and Abedon 2009)

899 **Table 4. Additional correlations that we hypothesized could occur between traits, based on trade-**

900 **offs found in or suggested by the literature.**

Trade-off	Expected correlations	Basis
Short-term vs. long-term virion quality	- specific infectivity s positively correlated with mortality rate m - specific infectivity s negatively correlated with the robust fraction r	Reflection of the general expectation that trade-offs occur between adaptations to different environments[1], in this case short-term and long-term outside-host conditions.
Survival vs. reproduction	- mortality rate m positively correlated with multiplication rate, calculated as b_c*s/μ_l following De Paepe & Taddei[2] - mortality rate m positively correlated with relative growth in the growth assay	Classic trade-off[3] that is also expected to apply to viruses[4,5]. Seminal example in De Paepe & Taddei[2].
Offspring quantity & quality	- burst size per infected cell b_d negatively correlated with specific infectivity s	Version of the classic quantity/quality trade-off[3,5], but at $t = 0$. Related to the speed-fidelity trade-off[6,7], though other processes than mutation may cause defective virions.

901 [1](Agrawal et al. 2010), [2](De Paepe and Taddei 2006), [3](Stearns 1989), [4](Goldhill and Turner 2014), [5](Keen 2014)
902 [6](Furió et al. 2005), [7](Belshaw et al. 2008)
903

Figures

Fig. 1. Life cycle of the chloroviruses, as assumed in our statistical models. See the text and Table 3

for definitions of the traits and model assumptions. Traits marked in blue are assumed to be

independent of each other; traits marked in red arise from combinations of two or more other traits

(see gray box). Traits in bold are estimated from the modified one-step growth (mOSG) or modified

persistence (mP) assay results; traits not in bold are derived from comparing the two assays. "*TrN*":

truncated normal distribution (truncated at 0 h).

Fig. 2. Schematic representation of the mOSG and mP assays for one virus strain. Both assays started from the same filtrate, represented by a 15 ml tube. Culture plates are depicted with round wells; deep well plates with square wells. The solid gray boxes describe the sampling procedure that was repeated at every time point. On the right, the gray arrows lead to examples of the data (points) and the model fits (gray lines). The data shown here is for virus strain SH-6A.

Fig. 3. Within- and across-group trait diversity revealed by the mOSG and mP assays. Upper panels: Trait estimates and 95 % CIs. Estimates and CIs

colored in black were retained for further analysis; estimates and traits colored red were excluded from the analyses due to poor or imprecise model fit.

Missing traits are due to failed or incomplete assays. See Supp. Table S1 for details and exact values. **Lower panels: Distributions of the retained estimates.**

Shown as violin plots with internal box plots. Only the estimates used in the analyses are included in these distributions (i.e. those shown in black in the

upper panels).

Fig. 4. Correlations between traits in the NC64A viruses. Shade represents high (dark green) vs. low (light green) relative growth in the growth assay (see panel C). **A.** Pairwise correlations between independent traits. **B.** Additional correlations that we hypothesized might occur ("mult. rate" multiplication rate; "rel. growth" relative growth). **A&B.** Each point represents one virus strain. Panels with significant Spearman correlations are marked with stars and a gray background. Note that the analyses were weighted by the precision of the retained estimates, and this weighting is not represented in the figure. A version of these panels with 95 % CIs can be found in Supp. Fig. S11. **C.** Results of the growth assay, expressed as an x-fold increase in virion concentration in 24 h. The 12 fastest-growing strains are marked.

Fig. 5. Correlations between traits in the fastest-growing NC64A viruses. A&C. Comparison of
Spearman's ρ correlation coefficients between independent traits (**A**) or for additional trait
combinations (**C**) when all strains vs. the fastest-growing strains are used (see Fig. 4C). Each point
represents one trait combination. The gray area marks the zone where -0.5 ≤ Δρ ≤ 0.5; points outside
this zone are labeled with the trait combination they represent. **B.** PCA of the independent traits in
the fastest-growing strains. The biplot shown here was produced by the Ipca method; the PairCor
result was extremely similar. The variance explained by each principal component (PC) is given on
the axis. Each point represents one virus strain.

Fig. 6. Summary figure, showing a simplified life cycle and correlation results. The main life cycle
steps (adsorption, depolarization, completion with burst, and decay) are shown, as well as relevant
compound traits (specific infectivity, relative growth). In the NC64A viruses, there were two main
correlational patterns that suggested life history trade-offs. The traits associated with these are
marked in respectively blue and red.

<u>Supplementary Material</u>

Detailed legends provided in each file.

Supplementary Method S1. Derivation of Eqs. 1 & 2.

Supplementary Table S1. Trait table for the chloroviruses, including selected traits from the
literature and the traits estimated in this manuscript.

Supplementary Table S2. ANOVA tables partitioning the trait variance within and among
chlorovirus groups.

Supplementary Table S3. Correlations between traits in the NC64A virus group.

Supp. Fig. S1. Example of statistical fitting for the mOSG model.

Supp. Fig. S2. Effect of imposing ⬚ = 0 in the mOSG model.

Supp. Fig. S3. Sensitivity of the mOSG model to its parameters.

Supp. Fig. S4. Repeatability of the mOSG assay.

Supp. Fig. S5. Effect of imposing r = 0 in the mP model.

Supp. Fig. S6. Sensitivity of the mP model to its parameters.

Supp. Fig. S7. Example of statistical fitting for the mP model.

Supp. Fig. S8. Repeatability of the mP assay.

Supp. Fig. S9. Correlations between model parameters arising from the fitting process.

Supp. Fig. S10. Effect of correlations between model parameters arising from the fitting process.

Supp. Fig. S11. Correlations between traits in the NC64A viruses, with 95 % CIs represented.

Supp. Fig. S12. Derivation of the independent traits c and b_c.

Supp. Fig. S13. Exploration of the assumption that mOSG parameters are independent of MOP.