Title: Hypoxia-inducible factor 1 signaling drives placental aging and can provoke preterm labor

Authors: Erin J. Ciampa[1], Padraich Flahardy[1], Harini Srinivasan[2], Christopher Jacobs[2], Linus Tsai[2],

S. Ananth Karumanchi[3], Samir M. Parikh[4]

Affiliations:

[1] Department of Anesthesia, Critical Care, and Pain Medicine, Beth Israel Deaconess Medical

Center, Harvard Medical School, Boston MA, USA

[2] Division of Endocrinology, Beth Israel Deaconess Medical Center, Harvard Medical School,

Boston MA, USA

[3] Department of Medicine, Cedars-Sinai Medical Center, Los Angeles CA, USA

[4] Division of Nephrology, Departments of Medicine and Pharmacology, University of Texas

Southwestern Medical School, Dallas TX, USA

Address correspondence to: Samir Parikh, MD. University of Texas Southwestern Medical

Center, Department of Internal Medicine. 5323 Harry Hines Blvd. Dallas, TX 75390-9030. Tel:

214-648-2754. Email: samir.parikh@utsouthwestern.edu.

Abstract:

Most cases of pre-term labor have unknown cause, and the burden of preterm birth is

immense. Placental aging has been proposed to promote labor onset, but specific mechanisms

remain elusive. We report findings stemming from unbiased transcriptomic analysis of mouse

placenta, which revealed that hypoxia-inducible factor 1 (HIF-1) stabilization is a hallmark of

advanced gestational timepoints, accompanied by mitochondrial dysregulation and cellular

senescence; we detected similar effects in aging human placenta. In parallel in primary mouse

trophoblasts and human choriocarcinoma cells, we modeled HIF-1 induction and demonstrated

resultant mitochondrial dysfunction and cellular senescence. Transcriptomic analysis revealed

that HIF-1 stabilization recapitulated gene signatures observed in aged placenta. Further,

conditioned media from trophoblasts following HIF-1 induction promoted contractility in

immortalized uterine myocytes, suggesting a mechanism by which the aging placenta may drive

the transition from uterine quiescence to contractility at the onset of labor. Finally,

pharmacological induction of HIF-1 via intraperitoneal administration of dimethyloxalyl glycine

(DMOG) to pregnant mice caused preterm labor. These results provide clear evidence for

placental aging in normal pregnancy, and demonstrate how HIF-1 signaling in late gestation

may be a causal determinant of the mitochondrial dysfunction and senescence observed within

the trophoblast as well as a trigger for uterine contraction.

Introduction:

Preterm birth (birth prior to 37 completed weeks of gestation) is a massive global health burden: it leads all causes of death in neonates and children to the age of 5 worldwide[1], and survivors face a broad array of short- and long-term health challenges[2]. Most preterm births in the United States are due to spontaneous onset of preterm labor, with unknown underlying cause[3]. Preterm labor has proven difficult to treat; there currently exist no highly effective interventions that prevent spontaneous preterm birth (SPTB)[4,5]. The dearth of effective treatments stems from our lack of understanding about the pathways regulating spontaneous onset of labor, both preterm and at term.

The placenta defines the maternal-fetal interface; it is capable of profoundly influencing both maternal and fetal physiology[6-8], and it is subject to remodeling in response to its local environment[9]. Placental disease is known to contribute to other disorders of pregnancy including preeclampsia[10-12] and intrauterine growth restriction[13,14] and there is increasing interest in its potential role as a driver of preterm labor[15-17].

Recent studies have found that placentas from pregnancies that resulted in SPTB have unique metabolomic and transcriptomic signatures from term counterparts[18-20]. The reported differences broadly implicate the stress response, inflammation, and various metabolic pathways. It has been challenging to narrow these findings or translate them for mechanistic relevance, given the lack of suitable culture systems that integrate experimental manipulation of placental cells with the uterine myocyte response. Furthermore, interpretation studies

profiling human placentas from SPTB is sharply limited by the lack of gestational age-matched

controls. Without these, the effects of gestational age cannot be distinguished from factors

driving premature labor. The placenta encounters a dynamic local environment across its

lifetime and faces evolving needs of the growing fetus, so the context of normal gestational

age-related changes is vital for the correct interpretation of molecular characteristics

distinguishing a SPTB placenta. Defining the molecular-level changes in the *healthy* placenta as

it approaches the end of gestation, and their potential effects on the timing of labor onset

would therefore address important knowledge gaps.

Here we report findings from unbiased transcriptomic analysis of healthy mouse placenta,

which highlighted hypoxia-inducible factor 1 (HIF-1) signaling as a hallmark of advanced

gestational timepoints, accompanied by mitochondrial dysfunction and cellular senescence. We

detected some similar effects in human placentas, then modeled HIF-1 induction with two

different stimuli and in two trophoblast cell models, demonstrating that mitochondrial

dysfunction and cellular senescence arise secondary to HIF-1 stabilization. Whole transcriptome

analysis revealed that upon HIF-1 stabilization, a trophoblast cell line acquires signatures that

recapitulate the aged placenta. Finally, we show that conditioned media from these cells is

sufficient to potentiate a contractile phenotype in uterine myocytes, implicating a mechanism

by which the aging placenta may help drive the transition from uterine quiescence to

contractility at the onset of labor. This mechanism is further reinforced by an *in vivo* model in

which administration of the prolyl hydroxylase inhibitor dimethyloxalyl glycine (DMOG) to

pregnant mice induces HIF-1 signaling in the placenta and causes SPTB.

Results

To illuminate gestational age-dependent transcriptional changes in across a healthy pregnancy,

we collected whole mouse placentas at 48h intervals spanning embryonic day 13.5 to 17.5

(e13.5-e17.5) and quantified mRNA and protein targets that emerged from network analysis of

an independently published microarray dataset from healthy mouse pregnancies. Upon

searching Gene Expression Omnibus (GEO)[21] datasets for "placenta AND transcriptome", we

were surprised that among 326 results, only nine included data from normal placenta across a

series of gestational timepoints extending to late pregnancy[22-28] (**Figure 1**). We applied

weighted gene correlation network analysis (WGCNA) to a microarray study of mouse

placenta[22] (GEO accession GSE11224) spanning e8.5 to postnatal day 0 (p0) to assess the mRNA

signature of the aging mouse placenta, using the dataset with the best temporal resolution

across gestation. Distinct clusters emerged, each reflecting a group of genes whose expression

changes across timepoints in a unified way (**Figure 2A;** accompanying statistics available at

Mendeley Data, doi: 10.17632/g6vrw9jjn4.1). Genes in the 'blue' cluster showed increasingly

positive correlation with advancing gestational timepoints; KEGG functional pathways

significantly overrepresented among these genes include HIF1 signaling, AMPK signaling, and

cellular senescence. Genes in the 'turquoise' cluster showed increasingly negative correlation to

advancing timepoints; KEGG functional pathways significantly overrepresented among these

genes include the citric acid cycle, mitochondrial complex I biogenesis, and the mitotic cell

cycle.

Figure 1. Systematic search flow for placental transcriptomics datasets. 326 GEO datasets were identified by the search terms "placenta" and "transcriptome"; nine met criteria for containing placental transcriptomic data representing normal physiology at a range of gestational timepoints spanning through the final 1/3 of pregnancy. Dataset selected for further analysis in yellow.

Figure 2. Mouse placental aging is characterized by cellular senescence, HIF-1 signaling, and mitochondrial dysregulation. WGCNA yielded 20 gene clusters. Functional pathways overrepresented in clusters found to increase (blue) and decrease (turquoise) across gestation highlight enhanced cellular senescence, increased HIF-1 signaling, and decreased mitochondrial synthesis and respiration late in pregnancy (A). Expression of senescence marker *Glb1* peaks at e17.5 (B; one-way ANOVA p=0.0048). HIF-1 protein abundance is higher at e17.5 versus e13.5 and e15.5 (C; one-way ANOVA p=0.019), as is expression of HIF-1 targets *Hk2* and *Glut1* (D; two-way ANOVA p<0.0001 for gestational age factor). (See Figure 2 - Supplement 1 for analysis of gene expression changes across timepoints by placental sex.) Mitochondrial abundance, reflected by COX IV protein, decreases with gestational age (E, one-way ANOVA p=0.0064), and mitochondrial DNA lesion rate peaks at e17.5 in the regions of the D-loop (one-way ANOVA p=0.0001), COII/ATPase6 (p=0.0027), and ND5 (p=0.036) (F). B-F, Each data point represents a biological replicate (eg. RNA, protein, or DNA extracted from an individual placenta, in turn collected from one of 2-4 pregnant dams per group). See Figure 2 - Source Data 1 for uncropped, unedited blots.

To validate the dynamic pathway activity suggested by the WGCNA, we quantified mRNA

expression of the senescence marker *Glb1*[29] in our mouse placentas, which mirrored the

WGCNA finding that cellular senescence in the placenta peaks in the final days of gestation

(**Figure 2B**). HIF-1 protein abundance was found to peak at e17.5 (**Figure 2C**) and HIF-1 targets

Hk2 and *Glut1* likewise confirmed increasing HIF-1 activation with advancing gestational age

through e17.5 (**Figure 2D**). Of note, a modest but significant fetal sex-dependent difference in

HIF-1 target expression (but not *Glb1*) was also observed across timepoints (**Figure 2 -**

Supplement 1). To assess changes in mitochondrial abundance across gestation, we measured

COX IV protein expression and found it declined progressively across mouse gestation (**Figure**

2E) as predicted by the WGCNA.

Having observed a pattern of declining mitochondrial abundance in the placenta with advancing

gestational age, we next investigated the mitochondrial DNA (mtDNA) lesion rate. The

mitochondrial genome is particularly vulnerable to reactive oxygen species (ROS) insults, and

mtDNA damage participates in a vicious cycle with mitochondrial dysfunction and further ROS

production; these effects are observed in a number of age-related diseases in various tissues[30]

and may drive age-associated loss of function[31]. We employed a semi-long run real-time qPCR

approach[32] to quantify relative mtDNA lesion rates in mouse placentas, normalized to e13.5.

There was a measurable increase in the mtDNA lesion rate in the D-loop and COII/ATPase 6

regions of the mitochondrial genome at e17.5 (**Figure 2F**). Together, these results reflect a

series of coordinated changes as gestation progresses —namely HIF-1 signaling induction,

A

two-way ANOVA: p = 0.83 (sex), p = 0.01 (gestational age)

B

two-way ANOVA: p = 0.25 (sex), p < 0.0001 (gestational age)

C

two-way ANOVA: p < 0.0001 (sex), p < 0.0001 (gestational age),
p = 0.026 (interaction)

Figure 2 - Supplement 1. Gestational age-dependent variability in expression of Hif-1 target *Glut1*, but not *Hk2*, is affected by placental sex. Subgroup analysis did not reveal sex-dependent expression changes in *Glb1* (A) or *Hk2* (B), but did demonstrate consistently higher *Glut1* expression in male placentas at all gestational ages and an interaction between the effects of sex and gestational age on *Glut1* expression (C; two-way ANOVA p<0.0001 for sex, p<0.0001 for gestational age, and p=0.028 for interaction). Each data point represents a biological replicate (RNA isolated from an individual placenta, collected from one of 2-4 dams per group).

decreasing mitochondrial abundance, accumulating ROS stress, and escalating cellular

senescence — confirming the patterns we discovered through reanalysis of published

transcriptomic data.

We next probed human placentas for the same gestational age-dependent changes. Studying

normal human placenta at a range of gestational ages beyond the second trimester is

challenging, as placental sampling is usually only possible after delivery, and most deliveries

associated with healthy pregnancies occur at term. We therefore sought to capitalize on rare

exceptions such as cases of placenta previa, vasa previa, or uterine dehiscence, where

iatrogenic preterm delivery is indicated for reasons unrelated to placental health. With an

objective to compare term versus preterm placentas, yet minimize confounding factors that

accompany labor and disease states that could be expected to affect placental health, we

designed a case-control study using the following exclusion criteria: onset of labor prior to

delivery (spontaneous or induced); maternal history of hypertension, asthma, diabetes, or

autoimmune disease; pregnancy complicated by gestational hypertension, gestational diabetes,

preeclampsia, multiples, fetal anomalies, placenta accreta spectrum disorder, or smoking

during pregnancy.

Placentas from 9 cesarean deliveries occurring before 35 weeks gestation and 11 cesarean

deliveries occurring after 39 weeks gestation were studied (**Table 1**). Maternal characteristics

including race, nulliparity, and obesity were not statistically different across the early versus

late groups; there was a small but statistically significant difference in maternal age at the time

	< 35 weeks	> 39 weeks	p value
	n = 9	n = 11	
Gestational age at delivery (weeks)	34.0 ± 0.3	39.5 ± 0.1	< 0.0001
Maternal age (years)	31.9 ± 0.9	36.2 ± 1.0	0.008
Maternal BMI > 40	1 (11)	0 (0)	0.26
Maternal race			
White	7 (78)	9 (82)	0.13
Black	2 (22)	0 (0)	
Asian	0 (0)	2 (18)	
Primiparous	4 (44)	4 (36)	0.71
Female neonate	5 (55)	8 (73)	0.42
Indication for delivery	Placenta previa (3)	Scheduled repeat (7)	
	Vasa previa (4)	Breech presentation (3)	
	Thinned lower uterine segment (2)	Elective (1)	

Table 1. Maternal and fetal characteristics. Data summarized by mean ± SEM or n (%). p-values calculated via t-test (continuous variables) or Chi-square contingency table (categorical variables).

of delivery (31.9 ± 0.9 years versus 36.2 ± 1.0 for early gestation versus late, p = 0.008). The

distribution of fetal sex was not different among groups.

We found a trend toward increased expression of *GLB1* and HIF-1 targets *HK-2* and *GLUT1* in

the > 39-week cohort (**Figure 3A**), consistent with the gestational age-dependent effect seen in

mouse placenta. We also examined mitochondrial abundance in the two groups and found that

mitochondrial RNA transcripts *ATP6* and *COX2* were significantly decreased (**Figure 3B**) and COX

IV protein abundance was lower at the later timepoint (**Figure 3C**), mirroring the mouse

findings. Of note, power calculation to reject the null hypothesis for difference between means

for some of these measurements (with α=0.05 and β=0.2) suggests a sample size of greater

than 35 per group is required, assuming a similar effect size as seen in mouse data (eg.

expecting *GLB1* fold change difference of 40% across groups), and greater variability than seen

for mouse data (eg. expected standard deviation of *GLB1* fold change equal to 0.6, versus 0.3 in

mice). Practical constraints, especially given our strict exclusion criteria, make a study of this

size unfeasible; nonetheless, we have included analysis of 20 human placentas here in

recognition of the vital importance of translating mouse findings to human biology, even

preliminarily. The data should be interpreted in the context of these statistical realities.

The co-appearance of cellular senescence, HIF-1 signaling, and mitochondrial dysregulation in

the placenta as it approaches the end of gestation led us to hypothesize that in aging placental

cells, HIF-1 induction could be upstream of mitochondrial dysregulation and cellular

senescence, as is seen in other systems in emerging aging research[33-35]. To test this hypothesis,

Figure 3. Senescence, HIF-1 signaling, and decreased mitochondrial abundance characterize late-gestation human placenta. Expression of senescence marker *GLB1* and HIF-1 targets *HK-2* and *GLUT1* trends higher in placentas from > 39-week cohort vs < 35-week cohort (A; 2-way ANOVA gestational age factor p=0.057). Mitochondrial abundance, reflected by mitochondrial genes *ATP6* and *COX2* (B; two-way ANOVA gestational age factor p=0.042) and COX IV protein (C; p=0.0036) decreases with advancing gestational age. Each data point represents a biological replicate (RNA or protein isolated from an individual placenta). See Figure 3 - Source Data 1 for uncropped, unedited blots.

we established a pharmacological model of HIF-1 induction in primary mouse trophoblasts

using cobalt chloride, a prolyl hydroxylase inhibitor that stabilizes HIF-1α[36,37] and has been

widely used to model hypoxia. After six hours of CoCl$_2$ exposure, we confirmed HIF-1 protein

accumulation in cultured trophoblasts (**Figure 4A**). After 48 hours of CoCl$_2$ exposure, mouse

trophoblasts exhibit decreased mitochondrial abundance, by *Cox2* expression (**Figure 4B**) and

Cox IV protein abundance (**Figure 4C**), and an increase in senescence-associated beta

galactosidase (SA-βGal), encoded by *Glb1* (**Figure 4D**) and detected as a blue stain in an X-gal

assay for senescence (**Figure 4E**). These findings suggest that HIF-1 stabilization induces

subsequent mitochondrial dysfunction and senescence in trophoblasts.

Primary trophoblasts undergo spontaneous syncytialization in culture, a phenomenon that

limits the duration of study and may also confound the interpretation of experimental changes

in key metabolic factors[38]. We therefore also modeled HIF-1 activation in JAR cells, a

trophoblast cell line that does not undergo syncytialization[39]. Consistent with our results in

primary cells, we found that HIF-1 is stabilized in CoCl$_2$-treated JAR cells (**Figure 5A**), and after 6

days of exposure, mitochondrial DNA and protein abundance declined (**Figure 5B-C**). We

further defined the time course of mitochondrial downregulation: by 72 hours the effect began

to appear (**Figure 5 – Supplement 1**). Furthermore, at the 6-day timepoint we found that CoCl$_2$

exposure impairs mitochondrial polarization (**Figure 5D**), and leads to accumulation of

mitochondrial reactive oxygen species (**Figure 5E**). Additionally, we observed pronounced signs

of cellular senescence: morphological hallmarks (cellular swelling) and β-galactosidase

overexpression (**Figure 5F**); growth arrest which persists for days after removal of the HIF-1

Figure 4. Short-term HIF-1 stabilization in primary mouse trophoblasts leads to decreased mitochondrial abundance and cellular senescence. HIF-1 is detected in cultured trophoblasts exposed to $CoCl_2$ (A). After 48 hours of $CoCl_2$ exposure, trophoblasts exhibit decreased mitochondrial abundance reflected by *Cox2* (B; p=0.014) and COX IV (C; p=0.0047) expression levels. Senescence marker *Glb1* is increased (D; p=0.038) and SA-βGal accumulation is noted by X-gal assay (E; p=0.012). Each data point represents a technical replicate (eg. protein, RNA, or β-Gal measured from an individual well of cells grown in treated or control condition). See Figure 4 - Source Data 1 for uncropped, unedited blots.

Figure 5. Long-term HIF-1 stabilization in JAR cells leads to mitochondrial dysfunction, cellular senescence, and metabolic reprogramming. JAR cell exposure to CoCl$_2$ stabilizes HIF1 (A). After 6 days, mitochondrial abundance is decreased as reflected by a drop in the mitochondrial:nuclear DNA copy number (B) and a decrease in COX IV protein (C). (See Figure 5 - Supplement 1 for timecourse of COX IV downregulation.) Cells also exhibit augmented signs of mitochondrial dysfunction, including MtSox (D) and TMRE staining (E; two-way ANOVA CoCl$_2$ factor p<0.0001). SA-βGal staining reflects a high proportion of senescent cells (F) and growth arrest is confirmed by cell counting following a 6-day pre-treatment with CoCl$_2$ (G; two-way ANOVA p<0.0001 for interaction of CoCl$_2$ factor with time). (See Figure 5 - Supplement 2 for assessment of cell death by propidium iodide staining.) SASP candidates VEGF, TNFα, IL-1α, and IL10 are dysregulated after CoCl$_2$ exposure (H; *, adjusted p<0.01). RNA Seq revealed upregulation of 2188 and downregulation of 1389 genes (I; genes with | log$_2$(FC) | > 1 and -log(FDR)>2 indicated in red) after CoCl$_2$ treatment, with geneset enrichment analysis revealing several pathways significantly dysregulated after CoCl$_2$ treatment recapitulating changes seen in transcriptomic analysis of late versus early gestation mouse placenta. Scale marker = 200 μm. See Figure 5 - Supplement 3 for assessment of effects of HIF-1 stabilization in JAR cells using DMOG. Each data point represents a technical replicate (measurement from an independent well of cells grown in treatment vs control condition). See Figure 5 - Source Data 1 for uncropped, unedited blots.

stabilizing compound (**Figure 5G**); and a senescence-associated secretory phenotype (SASP,

Figure 5H) reflected by increases in expression of VEGF, TNFα and IL-1α and a decrease in the

anti-inflammatory cytokine IL-10.

We conducted additional studies to confirm that cell death was not a primary contributor to

the lack of proliferation. Propidium iodide staining with quantitative fluorescence cytometry

indicated that CoCl$_2$ treatment only increased cell death by 0-3% (**Figure 5 - Supplement 2**).

Importantly, cells remained adherent and continued to acidify culture medium beyond 14 days

of CoCl$_2$ exposure, providing confidence that HIF-1 stabilization induces a phenotype

characterized by predominantly viable cells that are no longer proliferating, namely cellular

senescence. Finally, to assess whether the effects we observed were attributable to HIF-1

stabilization and not an off-target effect of CoCl$_2$, we also evaluated an alternative prolyl

hydroxylase inhibitor, dimethyloxalylglycine (DMOG)[40], and found similar effects on HIF-1,

mitochondrial abundance and cellular senescence (**Figure 5 – Supplement 3**).

To determine if other features of the aged placenta phenotype were recapitulated in this

model, we performed whole transcriptome analysis via RNA-Seq. Differential expression (log$_2$

fold change > 1; FDR < 0.05) was found for 2188 upregulated and 1389 downregulated genes

(**Figure 5I**). Gene set enrichment analysis revealed that chemical hypoxia led to upregulation of

inflammatory signaling, and downregulation of the TCA cycle, mitochondrial biogenesis, and

respiratory electron transport. These data therefore recapitulated metabolic patterns that

emerged from time course transcriptomic analysis of intact placentas (Figure 2A).

Figure 5 - Supplement 1. The mitochondrial effects of HIF-1 stabilization in JAR cells begin to appear on day 3 following CoCl$_2$ exposure. Each data point represents a technical replicate (independent well of cells grown in control vs treatment condition).

Figure 5 - Supplement 2. Increased number of JAR cells stain with propidium iodide, but the absolute number remains low following 6 days of CoCl$_2$ treatment. Each data point represents a technical replicate (independent well of cells grown in control vs treatment condition).

Figure 5 - Supplement 3. DMOG stabilizes HIF-1 in JAR cells (A) and induces similar effects as CoCl$_2$ on COX IV protein (B), SA-βGal expression (C), and cell growth (D) after 4 days. Each data point represents a technical replicate (measurement from an independent well of cells grown in treatment vs control condition). See Figure 5 - Source Data 1 for uncropped, unedited blots.

The onset of labor is accompanied by a transformation of the uterus from quiescence into a

distinct physiologic state in which it generates powerful, coordinated contractions. This

phenotypic change is characterized by upregulation of a cadre of contraction associated

proteins (CAPs), notably: cyclooxygenase-2 (COX-2, encoded by *PTGS2*), prostaglandin F2α

receptor (encoded by *FP*), interleukin-6 (IL-6), and connexin 43 (Cx43, encoded by *GJA1*)

Phenotypic switching can be modeled in primary and immortalized uterine myocytes upon

stimulation with inflammatory mediators such as IL1β, PGF2α, and thrombin[41-43].

To test whether primary placental metabolic disruptions could crosstalk with uterine myocytes

to trigger the contractile phenotype, we collected conditioned media from our JAR cell model

of HIF-1-driven senescence and measured the effect on expression of CAPs in uterine myocytes

(hTERT-HM cells). We observed a robust and specific effect: co-stimulation of uterine myocytes

with IL-1β plus conditioned media from JAR cells treated with $CoCl_2$ (but not from untreated

JAR cells) potentiated the induction of *PTGS2*, *GJA1*, *FP*, and *IL-6* expression (**Figure 6A-D**). We

next employed a collagen lattice assay previously described for assessing contractility of

myometrial cells *in vitro*[41,44], and demonstrated that myocyte contraction is augmented upon

exposure to conditioned media from JAR cells treated with $CoCl_2$ but not from untreated JAR

cells (**Figure 6E**). These transcriptional and functional results collectively demonstrate that

uterine myocytes are responsive to the secretome of JAR cells driven to senescence via HIF-1

induction.

Figure 6. Conditioned media (CM) from JAR cells following HIF-1 stabilization induces expression of contractile-associated proteins and augments contraction in immortalized human uterine myocytes. hTERT-HM expression of *PTGS2* (A), *GJA1* (B), *FP* (C), and *IL6* (D) was induced by CM from JAR cells following $CoCl_2$ treatment (but not in control conditions), similar to (in some cases potentiating) the effect of stimulation of myocytes by exogenous IL-1β. Percent well area occupied by hTERT-HM cells embedded in collagen matrix is significantly smaller after stimulation with IL-1β plus JAR CM from $CoCl_2$ condition, reflecting greater degree of hTERT-HM cellular contraction (E). Each data point represents a technical replicate (measurement from an independent well of cells grown in treatment vs control condition).

To test whether HIF-1 induction drives labor onset *in vivo*, we administered DMOG

intraperitoneally to pregnant mice on gestational day e16.5. Analysis of placentas recovered 12

hours following injection indicated that HIF-1 protein is stabilized in placenta following

maternal DMOG injection (**Figure 7A**), and transcription of HIF-1 target genes *Hk-2* and *Glut1*

was significantly increased (**Figure 7B**). Following injection of DMOG, gestational length was

significantly shortened compared to injection of vehicle alone (**Figure 7C-D**).

Discussion

The subject of whether the healthy placenta undergoes aging (with accompanying dysfunction)

within its 40-week lifespan has been of great interest and debate for many years. Age-related

functional decline is a near-universal phenomenon affecting many tissues, but the underlying

biochemical driving factors vary widely depending on the context of cell type and local

stressors. Mechanisms well described in other tissues are rooted in genomic instability,

mitochondrial dysfunction and oxidative stress, nutrient deprivation and metabolic

insufficiency, or loss of proteostasis[45]. While characterization of the placenta as an aging organ

appeared in the literature beginning nearly fifty years ago[46,47], a counter-argument has been

offered that there is no logical reason for the placenta to undergo accelerated aging relative to

the fetus, since both share the same genes and environment[48]. A third viewpoint proposes that

aging in the placenta is indicative of a disease state—not normal progression of healthy

gestation—one usually reflective of maladaptive responses to oxidative stress[49-53]. Finally,

recent reports have noted prominent signs of aging in another fetally derived tissue, the

chorioamniotic membranes, in healthy pregnancies[54], and have linked aging of the fetal

Figure 7. Maternal DMOG injection on e16.5 stabilizes placental HIF-1 and induces preterm labor. HIF-1α protein is detected in placental lysates 12 hours following DMOG injection but not vehicle (A). HIF-1 targets *Hk-2* and *Glut1* are upregulated following DMOG injection (p = 0.002 for DMOG vs vehicle, two-way ANOVA) (B). Gestational length is significantly shortened following DMOG injection versus vehicle (C-D). Each data point represents a biological replicate (in A and B, each measurement from an individual placenta collected from one of 2 pregnant dams). See Figure 7 - Source Data 1 for uncropped, unedited blots.

membranes to the onset of labor[55,56]. The results presented here span mouse and human

placentas to provide evidence for placental aging in normal pregnancy. Mechanistic dissection

of this phenomenon further demonstrates how hypoxia in late gestation may be both a causal

determinant of mitochondrial dysfunction and senescence observed within the trophoblast as

well as a trigger to induce uterine contraction.

To our knowledge, this is the first report to propose that placental aging in healthy pregnancy is

characterized by induction of HIF-1 signaling, accompanied by downstream effects including

mitochondrial dysfunction and cellular senescence. These findings have implications for the role

of the placenta in signaling that promotes the onset of labor, particularly given our

demonstration that upon HIF-1 stabilization, trophoblasts can induce inflammatory changes in

uterine myocytes and potentiate their contractility, and our *in vivo* demonstration that

administration of DMOG stabilizes placental HIF-1 and leads to preterm labor in mice.

Important next steps prompted by this work include determining the specific components of

the secretome of HIF-1-activated trophoblasts which are responsible for inducing myocyte

transformation. Additionally, having established that HIF-1 signaling is upstream of

mitochondrial dysfunction and cellular senescence in late-gestation trophoblasts in normal

pregnancies, future studies can investigate whether factors that modulate this pathway could

increase or decrease vulnerability to preterm labor. In studies targeted to uterine decidua, a

link between cellular senescence and labor onset has previously been established, with

evidence implicating phospho-Akt and mTORC1 signaling upstream of prostaglandin synthesis

as a key mechanism in this model[57,58]. It is possible that senescence signaling from the placenta

and decidua converge, and labor is provoked upon these convergent endpoint signals

surpassing a threshold, regardless of their origin within the gestational compartment.

Our results suggest that placental HIF-1 becomes stabilized late in gestation, and this could be

secondary to the onset of hypoxia sensing as oxygen demand outstrips supply in the

fetoplacental unit. However, placental hypoxia has been difficult to examine *in vivo*. Despite

evolving approaches for measurement of placental oxygenation, findings have varied widely

and adequate spatiotemporal resolution has so far been challenging[59]. However, it is apparent

that due to arteriovenous shunting and high metabolic extraction rates, the placenta

experiences hypoxia throughout pregnancy with a reported partial pressure of oxygen 30-50

mm Hg in the intervillous space, so perhaps a relevant question is what *prevents* HIF-1

stabilization and its downstream effects early in gestation. It is possible that the gestational-

age-dependent HIF-1 induction we observed reflects not a change in oxygen availability, but

rather a change in other HIF-regulatory factors such as nicotinamide adenine dinucleotide

(NAD^+) depletion, as has been demonstrated to occur in muscle and is accompanied by

metabolic reprogramming and mitochondrial decline[60]. Importantly, replenishment of NAD^+

was shown to restore markers of mitochondrial function in these aged cells, suggesting that

mitochondrial effects of HIF-1-driven aging are reversible in some systems. Separately, a recent

study which compared the miR-nomes of 1st- versus 3rd-trimester human placenta highlighted

differential expression of miRNAs with functional links to silencing of hypoxia response and

cellular senescence pathways in the first trimester[61]. Future investigations into the impact of

placental HIF-1 regulation will require a more precise understanding of placental oxygenation *in*

vivo and a sophisticated approach to model these factors in placental explants or other cellular

systems.

The shift in metabolic phenotype away from oxidative respiration in late pregnancy as

demonstrated by our transcriptomics analysis is consistent with earlier findings from

respirometry studies in mouse placenta[62], where rates of oxidative phosphorylation were noted

to decline from e14 to e19, particularly in the labyrinthine zone. It is still unclear if this shift

reflects changing substrate availability, or oxygen content, or perhaps the metabolic needs of

the developing fetus. A better understanding of the dynamic metabolism of the placenta across

gestation will be crucial for developing strategies for optimizing placental health, with major

implications for both maternal and fetal outcomes. For example, placental metabolomics

studies with a gestational time series could help corroborate the metabolic implications

interpreted from our transcriptomic data, while single-cell transcriptomics across advancing

gestational age in healthy placentas would offer critical cell-type localization of specific

metabolic events noted to occur late in pregnancy. Future studies aimed at targeted

manipulation of placental metabolism *in vivo*, whether pharmacologically or genetically, are

needed to define the physiologic relevance of these findings.

Our data have implications for other adverse pregnancy outcomes beyond preterm birth,

including intrauterine growth restriction and preeclampsia. An important study previously

examined mice with placental expression of a constitutively active HIF-1 mutant and found that

when induced from the beginning of gestation, placental HIF-1 signaling drives abnormal

placentogenesis and impaired spiral artery remodeling, leading to placental hypoperfusion,

fetal growth restriction, and a preeclampsia-like clinical syndrome[63]. In the context of these

earlier results, our study highlights the critical gestational age-dependence of HIF-1 effects,

having demonstrated that in a normally developed mouse placenta, induction of HIF-1 signaling

during the final days of gestation leads to labor onset. Timing may be key to understanding how

two distinct pregnancy complications – preterm labor and preeclampsia – appear to emerge

separately from overlapping pathophysiologic mechanisms[64-68].

In summary, we report a molecular characterization of placental aging phenomena occurring in

normal pregnancies, stemming from induction of HIF-1 signaling and downstream

mitochondrial dysfunction and cellular senescence. These findings establish aging in the

placenta as a feature of normal gestation; identify HIF-1 signaling as an upstream trigger

leading to mitochondrial dysfunction and senescence in the placenta; demonstrate that the

secretome of senescent trophoblasts is sufficient to potentiate uterine myocyte transformation

and contractility; and establish that HIF-1 induction *in vivo* can induce preterm labor. These

findings may have important implications for illuminating the factors that determine gestational

length both in health and disease.

Data Availability:

Statistics accompanying WGCNA (Figure 2) are accessible through Mendeley Data, doi:

10.17632/g6vrw9jjn4.1. (https://data.mendeley.com/datasets/g6vrw9jjn4/1)

The RNA Seq dataset has been deposited in NCBI's Gene Expression Omnibus[21] and is accessible

through GEO Series accession number GSE199278.

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE199278)

Additional source data (original images for all blots) available in attached files.

Methods:

Mouse placenta collection: Wild-type C57BL/6 mice (Jackson Laboratory #000664) were fed a

chow diet and housed at 20°C in a 12-hour light/12-hour dark cycle. Nulliparous females (< 6

months of age) were housed for a single dark cycle (midnight = gestational day 0) with a stud

male. On gestational day e13.5-17.5, placentas were isolated via laparotomy from pregnant

females anesthetized via surgical-plane isoflurane. Placentas were quartered, immersion-rinsed

in dH$_2$O and blotted dry, then snap frozen in liquid nitrogen and stored at -80°C prior to use.

Mouse trophoblast isolation: Protocol described in full in Pennington *et al*[69]. On gestational day

e15.5, placentas were collected via laparotomy from pregnant wild-type C57BL/6 female mice

anesthetized with surgical-plane isoflurane. After dissection, placentas were quartered,

immersion-rinsed in dH2O, then placed in ice-cold digestion buffer: DMEM plus HEPES 20 mM,

collagenase 1mg/mL, and DNase I 4000U/mL. Placentas were digested for 20-40 minutes at

37°C with trituration and examined under a microscope until optimal digestion achieved.

Digested cells were passed through a 70 micron strainer, washed in DMEM, then separated

using a Percoll gradient. Trophoblast fraction was washed once more then plated in complete

medium: DMEM with HEPES 20 mM, FBS 10%, and supplemented with sodium pyruvate, pen-

strep-glutamine, non-essential amino acids, and gentamicin.

Mouse DMOG injection and measurement of gestational length: On gestational day e16.5, dams

received an intraperitoneal injection of DMOG (Selleck Chemicals) 7.5 mg in 0.3 mL sterile

saline (approximately 250 mg/kg) versus vehicle alone. Video recordings were used to measure

gestational length, defined as interval from midnight of timed mating period until birth of the

first pup.

Human placental specimens. Placenta samples were collected at the time of cesarean delivery,

within 20 minutes of delivery of the placenta. Placental tissue was sampled from approximately

1cm deep to the maternal surface after dissecting away membranes. Cotyledons from all 4

quadrants were collected and minced together prior to immersion rinse in dH_2O and storage of

separate aliquots by snap freezing versus immersion in RNA Later. All samples were stored at -

80°C prior to use.

Cell culture.

JAR choriocarcinoma cells (ATCC HTB-144) were grown in RPMI1640 media (4.5 g/L glucose)

with 10% FBS. Adherent JAR cells were treated with HIF1-stabilizing agents dissolved in 1X PBS

(pH 7.4): cobalt chloride hexahydrate (100 µM $CoCl_2$, Millipore Sigma) and

dimethyloxalylglycine, N-(methoxyoxoacetyl)-glycine methyl ester (1 mM DMOG, Millipore

Sigma). JAR cells were treated with 100 μM CoCl$_2$ or 1 mM DMOG in culture media as indicated

prior to endpoint assays.

Preparation of conditioned media: On day 5 of JAR cell treatment with CoCl$_2$ (versus control

condition), media exchange was performed to apply serum-free media for both conditions. 24

hours later, conditioned media samples were collected and applied to 10 kDa molecular weight

cutoff filters (Amicon Ultra), with centrifugation for 20 minutes at 4000 x g, 4°C. Filters were

washed with one volume of hTERT-HM base medium (see below), with repeat centrifugation.

Concentrated proteins retained in suspension above the filter were collected and stored at -

80°C.

hTERT-HM cells (generously supplied by Dr. Jennifer Condon, Wayne State) were grown in

DMEM/F12 medium (Gibco) with 10% FBS and Antibiotic-Antimycotic (1X, Gibco). Cells were

stimulated by addition of recombinant human IL-1 (R&D Systems) and/or filter-concentrated

JAR cell conditioned media as indicated, 6 hours prior to collection of cellular RNA.

RT – qPCR. Total RNA was reverse transcribed to cDNA with the Superscript III reverse

transcriptase (Invitrogen) system with random hexamer primers (Invitrogen) according to

manufacturer's instructions. cDNA was amplified via real-time quantitative PCR with TaqMan

Fast Advanced Master Mix (Applied Biosystems) or SYBR Green PCR Master mix (Qiagen) in a

QuantStudio 6 Flex Real-Time PCR System (Applied Biosystems). In mouse samples, target gene

expression was normalized to endogenous levels of housekeeping gene, β-actin. In human

samples, target gene expression was normalized to endogenous levels of housekeeping gene,

YWHAZ (TaqMan) or β-actin (SYBR). Mouse and human primers are listed in Table 2.

Collagen lattice contraction assay: As previously described[41,44], hTERT-HM cells were suspended

in a collagen gel matrix (Cellmatrix collagen type I-A, Fisher Scientific, prepared with MEM,

NaOH and HEPES) then plated in 12-well dishes and grown in hTERT-HM medium. When cells

reached approximately 60% confluence, gel matrix was detached from plate and experimental

treatments were applied in fresh cell culture medium. After 1 hour incubation, a photograph

was taken of each well for measurement of percent well area occupied by collagen disc,

reflecting cellular contraction.

mtDNA damage assay. Mitochondrial DNA damage was assayed from total DNA via a semi-long

run real-time PCR approach, as described elsewhere[32]. Briefly, separate qPCR reactions were

assayed to compare a long versus short amplicon representing each genomic region. A lesion

rate per 10 kb of mtDNA was calculated as:

$\text{lesion rate} = \dfrac{10000\ (bp)}{size\ of\ long\ fragment\ (bp)} * (1 - FC_{long-short}),$

with $FC_{long-short} = 2^{-\Delta\Delta Ct}$ calculated in the usual method.

Relative mtDNA copy number. qPCR assays for nuclear and mitochondrial genes were

performed using total DNA as a template. Relative mitochondrial DNA was calculated as:

follows:

$\text{mt:nuc DNA copy number} = 2 * 2^{\Delta Ct}$, where $\Delta Ct = Ct_{nuclear\ gene} - Ct_{mitochondrial\ gene}.$

Immunoblots. Antibodies are listed in Table 3.

SA-β-Galactosidase Assay. Adherent cells were fixed and stained for SA-βGal via a Senescence

Detection kit (Abcam, ab65351) per manufacturer's instructions. Representative images were

used to calculate the fraction of SA-β-Gal-positive cells, scored by an observer blinded to

treatment condition.

Live cell staining. Adherent cells were stained using the TMRE-Mitochondrial Membrane

Potential Assay Kit (Abcam, ab113852) per manufacturer's instructions. Select wells were

pretreated with 20 µM FCCP, a decoupling agent, as a positive control for 20 minutes before

TMRE and Hoechst counterstain was applied. Live JAR cells were analyzed by the Celigo Image

Cytometer (Nexcelom BioScience) for fluorescence quantification and gating. Adherent live JAR

cells were stained with MitoSOX™ Red Mitochondrial Superoxide Indicator kit (Invitrogen) and

analyzed using the cytometer in a similar manner.

Cell counting. JAR cells were cultured for 6 days ± cobalt chloride exposure prior to cell

counting. On day 6, cells from each condition were replated in a 96-well plate in $CoCl_2$-free

media at a density of 5 x 10^3 cells per well. Celigo Image Cytometer was used for automated cell

counting (total cells per well) at 24-hour intervals.

Table 2. Primers

Mouse

Target	Sequence	Platform
GLB1	Mm00515342_m1	TaqMan
HK-2	Mm00443385_m1	TaqMan
GLUT1	Mm00441473_m1	TaqMan
D loop (long, 801 bp amplicon)	F: CGTACATTAAACTATTTTCCCCAAG R: GAGTTTTGGTTCACGGAACAT	SYBR
COII/ATPase 6 (long, 855 bp amplicon)	F: TTGGTCTACAAGACGCCACA R: ATTTTGGTGAAGGTGCCAGT	SYBR
ND5 (long, 930 bp amplicon)	F: CGCCTACTCCTCAGTTAGCC R: ATGGTGACTCAGTGCCAGGT	SYBR
ND2/ND1 (long, 832 bp amplicon)	F: GGATGAGCCTCAAACTCCAA R: ATGATGGCAAGGGTGATAGG	SYBR
D loop (short)	F: TGACTATCCCCTTCCCCATT R: TTGTTGGTTTCACGGAGGAT	SYBR
COII/ATPase 6 (short)	F: TCTCCCCTCTCTACGCATTC R: CGGTTAATACGGGGTTGTTG	SYBR
ND5 (short)	F: GGCCTCACATCATCACTCCT R: GCTGTGGATCCGTTCGTAGT	SYBR
ND2/ND1 (short)	F: GGATGAGCCTCAAACTCCAA R: GGCTCGTAAAGCTCCGAATA	SYBR
ACTINB	Mm00607939_s1	TaqMan

Human

Target	Sequence	Platform
HK-2	Hs00606086_m1	TaqMan
GLUT1	Hs00892681_m1	TaqMan
IL-1α	Hs00174092_m1	TaqMan
IL-1β	Hs01555410_m1	TaqMan
TNFα	Hs00174128_m1	TaqMan
IL-6	Hs00174131_m1	TaqMan
IL-10	Hs00961622_m1	TaqMan
MT-ATP-6	Hs02596862_g1	TaqMan
MT-CO2	Hs02596865_g1	TaqMan
ND1	F: CCATAAAACCCGCCACACT R: GAGCGATGGTGAGAGCTAAGGT	SYBR
18S	F: CGCAGCTAGGAATAATGGAATAGG R: CATGGCCTCAGTTCCGAAA	SYBR
CNX43	Hs.PT.58.38338544	SYBR
COX2	Hs.PT.58.77266	SYBR
ACTB	Hs.PT.39a.22214847	SYBR
YWHAZ	Hs01122445_g1	TaqMan

Table 3. Antibodies

Target	Species	Working Concentration	Source
Cox IV	Mouse IgG mAb	1:1,000	Cell Signaling Technology, #11967S
β-Actin-HRP	Rabbit IgG mAb	1:2,000	Cell Signaling Technology, #5125S
HIF1	Rabbit	1:1000	Cell Signaling Technology, #14179S

RNA-Seq. Following 6 days of treatment with CoCl2 100 ⵂM versus control media, total RNA was

isolated from JAR cells as described above. Extracted RNA (300ng) was treated with NEBNext

rRNA Depletion Kit v2 (E7400X) and cDNA was generated using random hexamer priming and

Maxima H Minus Reverse Transcriptase. cDNA was converted into double stranded cDNA using

NEBNext mRNA Second Strand Synthesis Module (E6111L) and sequencing libraries were

generated by tagmentation using Nextera XT DNA Library Preparation Kit (Illumina FC-131) with

12 cycles of PCR amplification. Sequencing libraries were analyzed by Qubit and Agilent

Bioanalyzer, pooled at a final concentration of 1.2pM, and sequenced on an Illumina

Nextseq500 instrument 36x8x36 read structure.

Transcriptomics Analysis: Sequencing reads were demultiplexed and trimmed for adapters

using bcl2fastq (v2.20.0). Secondary adapter trimming, NextSeq/Poly(G) tail trimming, and read

filtering was performed using fastp (v0.20.0)[70]; low quality reads and reads shorter than 18nt

after trimming were removed from the read pool. Salmon (v1.1.0)[71] was used to simultaneously

map and quantify reads to transcripts in the GENCODE 33 genome annotation of the

GRCh38/hg38 human assembly. Salmon was run using full selective alignment, with sequence-

specific and fragment GC-bias correction turned on (--seqBias and --gcBias options,

respectively). Transcript abundances were collated and summarized to gene abundances using

the tximport package for R[72]. Normalization and differential expression analysis were

performed using edgeR[73,74]. For differential gene expression analysis, genes were considered

significant if they passed a fold change (FC) cutoff of $\log_2 FC > 1$ and a false discovery rate (FDR)

cutoff of FDR < 0.05. Functional enrichment analyses were performed using g:Profiler[75].

WGCNA[76] was performed after removing genes with low expression as previously described[77].

Statistics. Student's two-tailed unpaired t-test, ordinary one-way, and two-way ANOVA

statistical tests were applied as indicated to compare biological replicates in each experiment.

Data were excluded as outliers using the interquartile range method (lower limit = 1^{st} quartile –

1.5 x IQR; upper limit = 3^{rd} quartile + 1.5 x IQR).

Study Approval. All animal experiments were approved by the Beth Israel Deaconess Medical

Center Institutional Animal Care and Use Committee (protocol #008-2022). Human placental

specimens and data were biobanked and accessed under protocols approved by the Beth Israel

Deaconess Medical Center Institutional Review Board, and written informed consent was

obtained before subject participation (protocols #2008P000061, 2020P000997, 2021P000897).

Author Contributions:

EJC, LT, SAK, and SMP designed the studies. HS, CJ, and LT performed WCGNA and JAR cell

transcriptomic assay and analyses. EJC and PF conducted *in vitro*, human placenta, and mouse

experiments and analyzed their results. EJC, SAK, and SMP wrote the manuscript with input

and approval from all authors.

Acknowledgements:

We thank Dr. Jennifer Condon of Wayne State University School of Medicine for generously

providing the hTERT-HM cell line.

We thank Saira Salahuddin for maintaining the placenta biorepository and helping access the

study samples and accompanying clinical data.

Figures were created using Biorender.com.

Research reported in this publication was supported by the Institute of General Medical

Sciences of the National Institutes of Health under award GM007592-41 and the Foundation for

Anesthesia Education and Research (FAER) (EJC).

The authors declare no competing interests.

References:

1. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J.E., Cousens, S., Mathers, C., and

Black, R.E. (2015). Global, regional, and national causes of child mortality in 2000-13,

with projections to inform post-2015 priorities: an updated systematic analysis. Lancet

385, 430-440. 10.1016/S0140-6736(14)61698-6.

2. Blencowe, H., Cousens, S., Chou, D., Oestergaard, M., Say, L., Moller, A.B., Kinney, M.,

Lawn, J., and Born Too Soon Preterm Birth Action, G. (2013). Born too soon: the global

epidemiology of 15 million preterm births. Reprod Health *10 Suppl 1*, S2. 10.1186/1742-

4755-10-S1-S2.

3. Goldenberg, R.L., Culhane, J.F., Iams, J.D., and Romero, R. (2008). Epidemiology and

causes of preterm birth. Lancet *371*, 75-84. 10.1016/S0140-6736(08)60074-4.

4. Smith, V., Devane, D., Begley, C.M., Clarke, M., and Higgins, S. (2009). A systematic

review and quality assessment of systematic reviews of randomised trials of

interventions for preventing and treating preterm birth. Eur J Obstet Gynecol Reprod

Biol *142*, 3-11. 10.1016/j.ejogrb.2008.09.008.

5. Romero, R., Dey, S.K., and Fisher, S.J. (2014). Preterm labor: one syndrome, many

causes. Science *345*, 760-765. 10.1126/science.1251816.

6. Kiserud, T., Ebbing, C., Kessler, J., and Rasmussen, S. (2006). Fetal cardiac output,

distribution to the placenta and impact of placental compromise. Ultrasound Obstet

Gynecol *28*, 126-136. 10.1002/uog.2832.

7.   Shaut, C.A., Keene, D.R., Sorensen, L.K., Li, D.Y., and Stadler, H.S. (2008). HOXA13 Is

essential for placental vascular patterning and labyrinth endothelial specification. PLoS

Genet *4*, e1000073. 10.1371/journal.pgen.1000073.

8.   Lykke, J.A., Langhoff-Roos, J., Sibai, B.M., Funai, E.F., Triche, E.W., and Paidas, M.J.

(2009). Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and

type 2 diabetes mellitus in the mother. Hypertension *53*, 944-951.

10.1161/HYPERTENSIONAHA.109.130765.

9.   Genbacev, O., Joslin, R., Damsky, C.H., Polliotti, B.M., and Fisher, S.J. (1996). Hypoxia

alters early gestation human cytotrophoblast differentiation/invasion in vitro and

models the placental defects that occur in preeclampsia. J Clin Invest *97*, 540-550.

10.1172/JCI118447.

10.   Plaks, V., Rinkenberger, J., Dai, J., Flannery, M., Sund, M., Kanasaki, K., Ni, W., Kalluri, R.,

and Werb, Z. (2013). Matrix metalloproteinase-9 deficiency phenocopies features of

preeclampsia and intrauterine growth restriction. Proc Natl Acad Sci U S A *110*, 11109-

11114. 10.1073/pnas.1309561110.

11.   Li, F., Fushima, T., Oyanagi, G., Townley-Tilson, H.W., Sato, E., Nakada, H., Oe, Y.,

Hagaman, J.R., Wilder, J., Li, M., et al. (2016). Nicotinamide benefits both mothers and

pups in two contrasting mouse models of preeclampsia. Proc Natl Acad Sci U S A *113*,

13450-13455. 10.1073/pnas.1614947113.

12.   Romero, R., and Chaiworapongsa, T. (2013). Preeclampsia: a link between trophoblast

dysregulation and an antiangiogenic state. J Clin Invest *123*, 2775-2777.

10.1172/JCI70431.

13.   McIntyre, K.R., Vincent, K.M.M., Hayward, C.E., Li, X., Sibley, C.P., Desforges, M.,

Greenwood, S.L., and Dilworth, M.R. (2020). Human placental uptake of glutamine and

glutamate is reduced in fetal growth restriction. Sci Rep *10*, 16197. 10.1038/s41598-

020-72930-7.

14.   Xu, Y.Y., Liu, Y., Cui, L., Wu, W.B., Quinn, M.J., Menon, R., and Zhang, H.J. (2021).

Hypoxic effects on the mitochondrial content and functions of the placenta in fetal

growth restriction. Placenta *114*, 100-107. 10.1016/j.placenta.2021.09.003.

15.   Koga, K., Cardenas, I., Aldo, P., Abrahams, V.M., Peng, B., Fill, S., Romero, R., and Mor, G.

(2009). Activation of TLR3 in the trophoblast is associated with preterm delivery. Am J

Reprod Immunol *61*, 196-212. 10.1111/j.1600-0897.2008.00682.x.

16.   Pique-Regi, R., Romero, R., Tarca, A.L., Sendler, E.D., Xu, Y., Garcia-Flores, V., Leng, Y.,

Luca, F., Hassan, S.S., and Gomez-Lopez, N. (2019). Single cell transcriptional signatures

of the human placenta in term and preterm parturition. Elife *8*. 10.7554/eLife.52004.

17.   Beharier, O., Tyurin, V.A., Goff, J.P., Guerrero-Santoro, J., Kajiwara, K., Chu, T., Tyurina,

Y.Y., St Croix, C.M., Wallace, C.T., Parry, S., et al. (2020). PLA2G6 guards placental

trophoblasts against ferroptotic injury. Proc Natl Acad Sci U S A *117*, 27319-27328.

10.1073/pnas.2009201117.

18.   Elshenawy, S., Pinney, S.E., Stuart, T., Doulias, P.T., Zura, G., Parry, S., Elovitz, M.A.,

Bennett, M.J., Bansal, A., Strauss, J.F., 3rd, et al. (2020). The Metabolomic Signature of

the Placenta in Spontaneous Preterm Birth. Int J Mol Sci *21*. 10.3390/ijms21031043.

19.   Lien, Y.C., Zhang, Z., Cheng, Y., Polyak, E., Sillers, L., Falk, M.J., Ischiropoulos, H., Parry, S.,

and Simmons, R.A. (2021). Human Placental Transcriptome Reveals Critical Alterations in

Inflammation and Energy Metabolism with Fetal Sex Differences in Spontaneous

Preterm Birth. Int J Mol Sci *22*. 10.3390/ijms22157899.

20.   Paquette, A.G., Brockway, H.M., Price, N.D., and Muglia, L.J. (2018). Comparative

transcriptomic analysis of human placentae at term and preterm delivery. Biol Reprod

98, 89-101. 10.1093/biolre/iox163.

21.   Edgar, R., Domrachev, M., and Lash, A.E. (2002). Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. Nucleic Acids Res *30*, 207-210.

10.1093/nar/30.1.207.

22.   Knox, K., and Baker, J.C. (2008). Genomic evolution of the placenta using co-option and

duplication and divergence. Genome Res *18*, 695-705. 10.1101/gr.071407.107.

23.   Zhou, Q.Y., Fang, M.D., Huang, T.H., Li, C.C., Yu, M., and Zhao, S.H. (2009). Detection of

differentially expressed genes between Erhualian and Large White placentas on day 75

and 90 of gestation. BMC Genomics *10*, 337. 10.1186/1471-2164-10-337.

24.   Loux, S.C., Dini, P., El-Sheikh Ali, H., Kalbfleisch, T., and Ball, B.A. (2019). Characterization

of the placental transcriptome through mid to late gestation in the mare. PLoS One *14*,

e0224497. 10.1371/journal.pone.0224497.

25.   Soncin, F., Khater, M., To, C., Pizzo, D., Farah, O., Wakeland, A., Arul Nambi Rajan, K.,

Nelson, K.K., Chang, C.W., Moretto-Zita, M., et al. (2018). Comparative analysis of

mouse and human placentae across gestation reveals species-specific regulators of

placental development. Development *145*. 10.1242/dev.156273.

26.   Maeda, K.J., Showmaker, K.C., Johnson, A.C., Garrett, M.R., and Sasser, J.M. (2019).

Spontaneous superimposed preeclampsia: chronology and expression unveiled by

temporal transcriptomic analysis. Physiol Genomics *51*, 342-355.

10.1152/physiolgenomics.00020.2019.

27. Morey, R., Farah, O., Kallol, S., Requena, D.F., Meads, M., Moretto-Zita, M., Soncin, F.,

Laurent, L.C., and Parast, M.M. (2021). Transcriptomic Drivers of Differentiation,

Maturation, and Polyploidy in Human Extravillous Trophoblast. Front Cell Dev Biol *9*,

702046. 10.3389/fcell.2021.702046.

28. Steinhauser, C.B., Lambo, C.A., Askelson, K., Burns, G.W., Behura, S.K., Spencer, T.E.,

Bazer, F.W., and Satterfield, M.C. (2021). Placental Transcriptome Adaptations to

Maternal Nutrient Restriction in Sheep. Int J Mol Sci *22*. 10.3390/ijms22147654.

29. Lee, B.Y., Han, J.A., Im, J.S., Morrone, A., Johung, K., Goodwin, E.C., Kleijer, W.J., DiMaio,

D., and Hwang, E.S. (2006). Senescence-associated beta-galactosidase is lysosomal beta-

galactosidase. Aging Cell *5*, 187-195. 10.1111/j.1474-9726.2006.00199.x.

30. Jang, J.Y., Blum, A., Liu, J., and Finkel, T. (2018). The role of mitochondria in aging. J Clin

Invest *128*, 3662-3670. 10.1172/JCI120842.

31. Trifunovic, A., Wredenberg, A., Falkenberg, M., Spelbrink, J.N., Rovio, A.T., Bruder, C.E.,

Bohlooly, Y.M., Gidlof, S., Oldfors, A., Wibom, R., et al. (2004). Premature ageing in mice

expressing defective mitochondrial DNA polymerase. Nature *429*, 417-423.

10.1038/nature02517.

32. Rothfuss, O., Gasser, T., and Patenge, N. (2010). Analysis of differential DNA damage in

the mitochondrial genome employing a semi-long run real-time PCR approach. Nucleic

Acids Res *38*, e24. 10.1093/nar/gkp1082.

33.   Bratic, A., and Larsson, N.G. (2013). The role of mitochondria in aging. J Clin Invest *123*,

619   951-957. 10.1172/JCI64125.

620   34.   Wiley, C.D., and Campisi, J. (2016). From Ancient Pathways to Aging Cells-Connecting

621   Metabolism and Cellular Senescence. Cell Metab *23*, 1013-1021.

622   10.1016/j.cmet.2016.05.010.

623   35.   Wiley, C.D., and Campisi, J. (2021). The metabolic roots of senescence: mechanisms and

624   opportunities for intervention. Nat Metab *3*, 1290-1301. 10.1038/s42255-021-00483-8.

625   36.   Maxwell, P.H., Wiesener, M.S., Chang, G.W., Clifford, S.C., Vaux, E.C., Cockman, M.E.,

626   Wykoff, C.C., Pugh, C.W., Maher, E.R., and Ratcliffe, P.J. (1999). The tumour suppressor

627   protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature

628   *399*, 271-275. 10.1038/20459.

629   37.   Jaakkola, P., Mole, D.R., Tian, Y.M., Wilson, M.I., Gielbert, J., Gaskell, S.J., von

630   Kriegsheim, A., Hebestreit, H.F., Mukherji, M., Schofield, C.J., et al. (2001). Targeting of

631   HIF-alpha to the von Hippel-Lindau ubiquitylation complex by O2-regulated prolyl

632   hydroxylation. Science *292*, 468-472. 10.1126/science.1059796.

633   38.   Nursalim, Y.N.S., Blenkiron, C., Groom, K.M., and Chamley, L.W. (2020). Growing human

634   trophoblasts in vitro: a review of the media commonly used in trophoblast cell culture.

635   Reproduction *160*, R119-R128. 10.1530/REP-19-0605.

636   39.   Rothbauer, M., Patel, N., Gondola, H., Siwetz, M., Huppertz, B., and Ertl, P. (2017). A

637   comparative study of five physiological key parameters between four different human

638   trophoblast-derived cell lines. Sci Rep *7*, 5892. 10.1038/s41598-017-06364-z.

40. Epstein, A.C., Gleadle, J.M., McNeill, L.A., Hewitson, K.S., O'Rourke, J., Mole, D.R.,

Mukherji, M., Metzen, E., Wilson, M.I., Dhanda, A., et al. (2001). C. elegans EGL-9 and

mammalian homologs define a family of dioxygenases that regulate HIF by prolyl

hydroxylation. Cell *107*, 43-54. 10.1016/s0092-8674(01)00507-4.

41. Nishimura, F., Mogami, H., Moriuchi, K., Chigusa, Y., Mandai, M., and Kondoh, E. (2020).

Mechanisms of thrombin-Induced myometrial contractions: Potential targets of

progesterone. PLoS One *15*, e0231944. 10.1371/journal.pone.0231944.

42. Rauk, P.N., and Chiao, J.P. (2000). Interleukin-1 stimulates human uterine prostaglandin

production through induction of cyclooxygenase-2 expression. Am J Reprod Immunol

43, 152-159. 10.1111/j.8755-8920.2000.430304.x.

43. Leimert, K.B., Verstraeten, B.S.E., Messer, A., Nemati, R., Blackadar, K., Fang, X.,

Robertson, S.A., Chemtob, S., and Olson, D.M. (2019). Cooperative effects of sequential

PGF2alpha and IL-1beta on IL-6 and COX-2 expression in human myometrial cellsdagger.

Biol Reprod *100*, 1370-1385. 10.1093/biolre/ioz029.

44. Devost, D., and Zingg, H.H. (2007). Novel in vitro system for functional assessment of

oxytocin action. Am J Physiol Endocrinol Metab *292*, E1-6.

10.1152/ajpendo.00529.2005.

45. Campisi, J., Kapahi, P., Lithgow, G.J., Melov, S., Newman, J.C., and Verdin, E. (2019).

From discoveries in ageing research to therapeutics for healthy ageing. Nature *571*, 183-

192. 10.1038/s41586-019-1365-2.

46. Rosso, P. (1976). Placenta as an aging organ. Curr Concepts Nutr *4*, 23-41.

47.   Martin, B.J., and Spicer, S.S. (1973). Ultrastructural features of cellular maturation and

aging in human trophoblast. J Ultrastruct Res *43*, 133-149. 10.1016/s0022-

5320(73)90074-9.

48.   Fox, H. (1997). Aging of the placenta. Arch Dis Child Fetal Neonatal Ed *77*, F171-175.

10.1136/fn.77.3.f171.

49.   Sultana, Z., Maiti, K., Aitken, J., Morris, J., Dedman, L., and Smith, R. (2017). Oxidative

stress, placental ageing-related pathologies and adverse pregnancy outcomes. Am J

Reprod Immunol *77*. 10.1111/aji.12653.

50.   Cindrova-Davies, T., Fogarty, N.M.E., Jones, C.J.P., Kingdom, J., and Burton, G.J. (2018).

Evidence of oxidative stress-induced senescence in mature, post-mature and

pathological human placentas. Placenta *68*, 15-22. 10.1016/j.placenta.2018.06.307.

51.   Biron-Shental, T., Sukenik-Halevy, R., Sharon, Y., Goldberg-Bittman, L., Kidron, D., Fejgin,

M.D., and Amiel, A. (2010). Short telomeres may play a role in placental dysfunction in

preeclampsia and intrauterine growth restriction. Am J Obstet Gynecol *202*, 381 e381-

387. 10.1016/j.ajog.2010.01.036.

52.   Chen, K.H., Chen, L.R., and Lee, Y.H. (2011). Exploring the relationship between preterm

placental calcification and adverse maternal and fetal outcome. Ultrasound Obstet

Gynecol *37*, 328-334. 10.1002/uog.7733.

53.   Maiti, K., Sultana, Z., Aitken, R.J., Morris, J., Park, F., Andrew, B., Riley, S.C., and Smith, R.

(2017). Evidence that fetal death is associated with placental aging. Am J Obstet Gynecol

217, 441 e441-441 e414. 10.1016/j.ajog.2017.06.015.

54. Bonney, E.A., Krebs, K., Saade, G., Kechichian, T., Trivedi, J., Huaizhi, Y., and Menon, R.

(2016). Differential senescence in feto-maternal tissues during mouse pregnancy.

Placenta *43*, 26-34. 10.1016/j.placenta.2016.04.018.

55. Menon, R., Behnia, F., Polettini, J., Saade, G.R., Campisi, J., and Velarde, M. (2016).

Placental membrane aging and HMGB1 signaling associated with human parturition.

Aging (Albany NY) *8*, 216-230. 10.18632/aging.100891.

56. Menon, R., Richardson, L.S., and Lappas, M. (2019). Fetal membrane architecture, aging

and inflammation in pregnancy and parturition. Placenta *79*, 40-45.

10.1016/j.placenta.2018.11.003.

57. Hirota, Y., Daikoku, T., Tranguch, S., Xie, H., Bradshaw, H.B., and Dey, S.K. (2010).

Uterine-specific p53 deficiency confers premature uterine senescence and promotes

preterm birth in mice. J Clin Invest *120*, 803-815. 10.1172/JCI40051.

58. Hirota, Y., Cha, J., Yoshie, M., Daikoku, T., and Dey, S.K. (2011). Heightened uterine

mammalian target of rapamycin complex 1 (mTORC1) signaling provokes preterm birth

in mice. Proc Natl Acad Sci U S A *108*, 18073-18078. 10.1073/pnas.1108180108.

59. Nye, G.A., Ingram, E., Johnstone, E.D., Jensen, O.E., Schneider, H., Lewis, R.M.,

Chernyavsky, I.L., and Brownbill, P. (2018). Human placental oxygenation in late

gestation: experimental and theoretical approaches. J Physiol *596*, 5523-5534.

10.1113/JP275633.

60. Gomes, A.P., Price, N.L., Ling, A.J., Moslehi, J.J., Montgomery, M.K., Rajman, L., White,

J.P., Teodoro, J.S., Wrann, C.D., Hubbard, B.P., et al. (2013). Declining NAD(+) induces a

pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. Cell

155, 1624-1638. 10.1016/j.cell.2013.11.037.

61.   Gonzalez, T.L., Eisman, L.E., Joshi, N.V., Flowers, A.E., Wu, D., Wang, Y., Santiskulvong,

C., Tang, J., Buttle, R.A., Sauro, E., et al. (2021). High-throughput miRNA sequencing of

the human placenta: expression throughout gestation. Epigenomics *13*, 995-1012.

10.2217/epi-2021-0055.

62.   Sferruzzi-Perri, A.N., Higgins, J.S., Vaughan, O.R., Murray, A.J., and Fowden, A.L. (2019).

Placental mitochondria adapt developmentally and in response to hypoxia to support

fetal growth. Proc Natl Acad Sci U S A *116*, 1621-1626. 10.1073/pnas.1816056116.

63.   Albers, R.E., Kaufman, M.R., Natale, B.V., Keoni, C., Kulkarni-Datar, K., Min, S., Williams,

C.R., Natale, D.R.C., and Brown, T.L. (2019). Trophoblast-Specific Expression of Hif-

1alpha Results in Preeclampsia-Like Symptoms and Fetal Growth Restriction. Sci Rep *9*,

2742. 10.1038/s41598-019-39426-5.

64.   Rasmussen, S., Ebbing, C., and Irgens, L.M. (2017). Predicting preeclampsia from a

history of preterm birth. PLoS One *12*, e0181016. 10.1371/journal.pone.0181016.

65.   Mando, C., De Palma, C., Stampalija, T., Anelli, G.M., Figus, M., Novielli, C., Parisi, F.,

Clementi, E., Ferrazzi, E., and Cetin, I. (2014). Placental mitochondrial content and

function in intrauterine growth restriction and preeclampsia. Am J Physiol Endocrinol

Metab *306*, E404-413. 10.1152/ajpendo.00426.2013.

66.   Fujimaki, A., Watanabe, K., Mori, T., Kimura, C., Shinohara, K., and Wakatsuki, A. (2011).

Placental oxidative DNA damage and its repair in preeclamptic women with fetal growth

restriction. Placenta *32*, 367-372. 10.1016/j.placenta.2011.02.004.

67. Davy, P., Nagata, M., Bullard, P., Fogelson, N.S., and Allsopp, R. (2009). Fetal growth

restriction is associated with accelerated telomere shortening and increased expression

of cell senescence markers in the placenta. Placenta *30*, 539-542.

10.1016/j.placenta.2009.03.005.

68. Burton, G.J., Yung, H.W., Cindrova-Davies, T., and Charnock-Jones, D.S. (2009). Placental

endoplasmic reticulum stress and oxidative stress in the pathophysiology of unexplained

intrauterine growth restriction and early onset preeclampsia. Placenta *30 Suppl A*, S43-

48. 10.1016/j.placenta.2008.11.003.

69. Pennington, K.A., Schlitt, J.M., and Schulz, L.C. (2012). Isolation of primary mouse

trophoblast cells and trophoblast invasion assay. J Vis Exp, e3202. 10.3791/3202.

70. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics *34*, i884-i890. 10.1093/bioinformatics/bty560.

71. Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides

fast and bias-aware quantification of transcript expression. Nat Methods *14*, 417-419.

10.1038/nmeth.4197.

72. Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. F1000Res *4*, 1521.

10.12688/f1000research.7563.2.

73. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. Bioinformatics *26*,

139-140. 10.1093/bioinformatics/btp616.

74. Chen, Y., Lun, A.T., and Smyth, G.K. (2016). From reads to genes to pathways:

differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR

quasi-likelihood pipeline. F1000Res *5*, 1438. 10.12688/f1000research.8987.2.

75. Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019).

g:Profiler: a web server for functional enrichment analysis and conversions of gene lists

(2019 update). Nucleic Acids Res *47*, W191-W198. 10.1093/nar/gkz369.

76. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation

network analysis. BMC Bioinformatics *9*, 559. 10.1186/1471-2105-9-559.

77. Bentsen, M.A., Rausch, D.M., Mirzadeh, Z., Muta, K., Scarlett, J.M., Brown, J.M.,

Herranz-Perez, V., Baquero, A.F., Thompson, J., Alonge, K.M., et al. (2020).

Transcriptomic analysis links diverse hypothalamic cell types to fibroblast growth factor

1-induced sustained diabetes remission. Nat Commun *11*, 4458. 10.1038/s41467-020-

17720-5.

758