Structural insights into the regulation of acetylation-dependent genome condensation

Ziad Ibrahim[1], Tao Wang[2], Olivier Destaing[2], Nicola Salvi[3], Naghmeh Hoghoughi[2], Alexandra Rusu[1], Jinjun Gao[4], Leonardo Feletto[1], Nicolas Reynoird[2], Thomas Schalch[1], Yingming Zhao[4], Martin Blackledge[3], Saadi Khochbin[2] & Daniel Panne[1,*]

[1]Leicester Institute of Structural and Chemical Biology, Department of Molecular and Cell Biology, University of Leicester, Leicester, UK

[2]CNRS UMR 5309, INSERM U1209, Université Grenoble Alpes, Institute for Advanced Biosciences, Grenoble, France.

[3]Institut de Biologie Structurale, CNRS, CEA, UGA, Grenoble, France

[4]Ben May Department of Cancer Research, The University of Chicago, Chicago, IL 60637, USA.

*Correspondence: daniel.panne@leicester.ac.uk

Summary
Histone modifications are deposited by chromatin modifying enzymes and read out by proteins that recognize the modified state. BRD4-NUT is an oncogenic fusion protein of the acetyl lysine reader BRD4 that binds to the acetylase p300 and enables formation of large hyperacetylated chromatin 'megadomains'. We here examine how 'reading' and 'writing' contribute to larger-scale chromatin architecture. We show that NUT contains an acidic transcriptional activation domain that binds to the TAZ2 domain of p300. We use NMR to investigate the structure of the complex and found that the TAZ2 domain has an autoinhibitory role for p300. NUT-TAZ2 interaction or mutants that interfere with autoinhibition by TAZ2 allosterically activate p300. p300 activation results in a self-organizing, acetylation-dependent feed-forward reaction that enables large-scale intra- and interchromosomal interactions by bromodomain multivalent acetyl-lysine binding. We discuss how the acetylation-dependent 'read-write' reaction drives condensation and the implications for chromatin organisation, gene regulation and dysregulation in disease.

Keywords: Chromatin modification, Acetylation, bromodomain, BRD4, acetyltransferase, p300, transcription factor, activation domain, gene regulation, NMR

Introduction

Chromosome capture experiments have led to the identification of two major mechanisms that are behind large-scale chromosome organisation (Lieberman-Aiden et al., 2009; Schwarzer et al., 2017): (1) at shorter genomic distances, genomes are organized into topologically associating domains (Dixon et al., 2012; Nora et al., 2012) which arise through the process of loop extrusion by cohesin; (2) at longer-range genomic distances, genomes are spatially compartmentalised into active, euchromatic and inactive, heterochromatic regions (Misteli, 2020; Nuebler et al., 2018).

One of the main mechanisms driving genome compartmentalisation is thought to be selective self-association of active or inactive regions and physical constraints including association with architectural elements (Solovei et al., 2016; van de Werken et al., 2017). One hypothesis is that 'readers' of histone modifications seed 'condensation' reactions as a function of the histone modification state (Misteli, 2020; Wang et al., 2019). It has been proposed that compartmentalisation arises specifically due to attractions between heterochromatic regions, possibly driven by polymer-polymer interactions and liquid-liquid phase separation (LLPS) (Erdel and Rippe, 2018; Falk et al., 2019). Whether heterochromatic compartments form through LLPS or other mechanisms is controversial (Erdel et al., 2020; Larson et al., 2017; Strom et al., 2021; Strom et al., 2017).

Histone acetylation counteracts heterochromatinization and has long been associated with chromatin decondensation and euchromatic chromatin regions (Ho et al., 2014). Targeting of heterochromatin with acidic transcription factor (TF) activation domains (ADs) or treatment with deacetylase inhibitors results in hyperacetylation, decondensation and relocation to the nuclear interior even in the absence of active transcription (Carpenter et al., 2005; Chambeyron and Bickmore, 2004; Mueller et al., 2001; Taddei et al., 2001; Therizols et al., 2014; Toth et al., 2004; Tumbar et al., 1999b).

Numerous TFs associate through their ADs with a small set of co-activators including Mediator and the acetylases CBP and its paralogue p300 (also known as KAT3A and KAT3B respectively; hereafter 'p300' for short) (Allen and Taatjes, 2015; Avantaggiati et al., 1996; Green, 2005; Malik and Roeder, 2010; Merika et al., 1998; Reiter et al., 2017). p300 has long been known to be important in integrating multiple signal transduction pathways in the nucleus and enabling enhancer-mediated transcription (Goodman and Smolik, 2000; Kamei et al., 1996; Narita et al., 2021). Intriguingly, it has been proposed that spatial sequestration of heterochromatin at the nuclear lamina requires TF-mediated retention of p300 in euchromatin (Cabianca et al., 2019). Quantitative Mass Spectrometry experiments show that p300 is present at limiting concentrations in the nucleus while HDAC co-repressors are in large excess (Gillespie et al., 2020). As p300 is responsible for up to a third of nuclear acetylation (Weinert et al., 2018), TF-mediated competition for and retention of these essential and limiting co-activators in euchromatic regions could be a key driver event that not only shapes gene specific transcription but also eu- and heterochromatic genome compartmentalisation both locally and globally.

It has been proposed that transcriptional control may be driven by the formation of phase-separated condensates (Hnisz et al., 2017). Accordingly, condensates of TF ADs (Boija et al.,

2018), co-activators and the transcriptional machinery have all been proposed to be relevant for transcriptional regulation (Cho et al., 2018; Chong et al., 2018; Guo et al., 2019; Sabari et al., 2018; Shrinivas et al., 2019; Zamudio et al., 2019). p300 contains an ordered catalytic 'core' domain including a HAT- and bromodomain (BD) and several domains that form promiscuous interactions with the disordered ADs of hundreds of cellular TFs (Fig. 1a) (Delvecchio et al., 2013; Dyson and Wright, 2016). These interactions domains are separated by long, intrinsically disordered regions (IDRs). p300 is activated by autoacetylation (Dancy and Cole, 2015), a reaction that can be controlled by cellular signalling through TF activation and dimerisation (Ortega et al., 2018). The p300 core contains a RING and TAZ2 domain that have an autoinhibitory function (Delvecchio et al., 2013; Ortega et al., 2018). Perturbation of the autoinhibitory function of the RING and TAZ2 domains by mutation leads to constitutive p300 activation and formation of nuclear droplets that co-localise with p53. These droplets are dispersed upon to HAT or BD inhibition, suggesting that a form of phase separation (we here use the more general term 'condensation') occurs through an acetylation-dependent 'read-write' mechanism (Ortega et al., 2018). Other data indicate that p300 forms nuclear condensates through the 'core' domain (Zhang et al., 2021). Co-condensation of p300 with TFs on enhancer loci is thought to enable activation of p300's catalytic activity thus allowing the recruitment of acetylation-dependent coactivators, including the BD - and extraterminal domain-containing (BET) protein BRD4 (Ma et al., 2021).

BET proteins have long been associated with the control of large-scale nuclear structure and chromatin organization. BRDT, a testes-specific BET, mediates histone acetylation-dependent chromatin condensation (Pivot-Pajot et al., 2003) and is required for stage-specific gene expression and acetylation-dependent histone removal during spermatogenesis (Gaucher et al., 2012; Goudarzi et al., 2016; Shiota et al., 2018). Similarly, BRD4 has been linked both to gene-specific (Zhao et al., 2011) and higher-order chromatin organisation (Devaiah et al., 2016; Wang et al., 2012b). In yeast, the BET protein BDF1 maintains physical genome barriers by preventing spreading of heterochromatin (Ladurner et al., 2003). *In vitro* data show that under specific buffer conditions, chromatin can undergo LLPS which can be modulated by acetylation and BRD4 binding with BD multivalency being a key determinant factor (Gibson et al., 2019)

BRD4 also forms condensates in cells (Sabari et al., 2018). *In vitro* and *in vivo* experiments with systems inducing high local concentration suggest that a C-terminal IDR drives the condensation reaction (Sabari et al., 2018; Shin et al., 2018). However, a BRD4 isoform lacking the C-terminal IDR also forms condensates in cells (Han et al., 2020). Other data indicate that BRD4 clustering does not depend on IDR-mediated condensation but on DNA and BD-acetyl-lysine recognition which possibly enables long-range enhancer-promoter communication (Li et al., 2020). At a larger scale, such acetylation-dependent long-range interactions could enable selective self-association of euchromatic regions and provide a link between genome topology and gene regulation. It is therefore important to understand the underlying 'read-write' mechanisms that drive such long-range interactions.

BRD4-NUT is an oncogenic fusion protein that drives NUT carcinoma, one of the most aggressive human solid malignancies known (Eagen and French, 2021). NUT is normally expressed in post-meiotic spermatogenic cells, where it interacts with p300 and triggers genome-wide histone hyperacetylation, which enables recruitment of BRDT, histone-to-protamine exchange and chromatin compaction during spermatogenesis (Shiota et al., 2018).

In the oncogenic fusion protein, NUT binds to and activates p300 which drives an acetylation- and BRD4 BD-dependent 'feed-forward' loop that leads to formation of hyperacetylated nuclear 'foci' visible by light microscopy (Eagen and French, 2021; Reynoird et al., 2010; Yan et al., 2011), that resemble foci seen in cells expressing constitutively activated p300 (Ortega et al., 2018). These foci correspond to large chromosomal 'megadomains' (100 kb to 2 Mb) that are enriched for BRD4-NUT, p300 and acetylated histones (Alekseyenko et al., 2015; Alekseyenko et al., 2017; Wang and You, 2015). Megadomains form in the absence of transcription and engage in very long-range acetylation-dependent inter- and intrachromosomal interactions (Rosencrance et al., 2020).

The BRD4-NUT model thus exemplifies how a modification of BRD4 enables recruitment and activation of p300 and establishment of a self-organizing feed-forward reaction that results in chromatin hyperacetylation and long-range contact. The underlying mechanisms could have broad implications for our understanding of how 'reading' and 'writing' of chromatin acetylation control gene expression and how dysregulation of this process leads to aggressive forms of disease.

Here we use the BDR4-NUT/p300 system to dissect the mechanism that drives large-scale chromatin organization. We determined the element of BRD4-NUT required for interaction with p300. We found that the NUT portion of the fusion protein contains an acidic AD that interacts with the p300 TAZ2 domain, a common binding site for numerous TF ADs. We investigate the solution behavior of the complex and found that interaction with the TAZ2 domain stimulates p300 HAT activity. Our data indicate that the TAZ2 domain has an auto-inhibitory function that is relieved upon TAZ2 ligand-binding. We systematically probed which elements of BRD4-NUT and p300 drive large scale chromatin condensation in cells. We found that condensation depends on a feed-forward mechanism that involves allosteric activation of p300 by binding of NUT to the p300 TAZ2 domain, nucleation of the condensation reaction through BRD4 and p300 BD-acetyl-lysine binding and that BD multivalency is required. We found that other TAZ2 ligands including p53 and E1A or mutations that interfere with TAZ2 autoinhibition also activate p300 and induce similar condensation reactions. Constitutive activation of p300 is associated with cancer. Our data reveal how activation of p300 establishes a self-organising positive feedback mechanism and how an acetylation-dependent read-write mechanism drives larger-scale chromatin architecture.

Results

Identification of an acidic activation domain in NUT
As a first step in characterization of the p300-NUT complex, we examined regions of the two proteins required for interaction (Fig. 1a,b). Previous data showed that a region spanning BRD4-NUT amino acids 347-588 (NUT F1c; NUT numbering is used; 1060-1307 in BRD4-NUT) directly interacts with p300 (Reynoird et al., 2010). To map the interaction further, we produced a series of deletions within NUT and assayed their interaction with p300 using co-sedimentation and GST pulldown assays. NUT fragments that contained amino acids 427-527 generally co-sedimented with the p300 core-CH3, the isolated p300 CH3 or TAZ2 domains (Fig. S1a,b; Fig. S2a-e).

The TAZ2 domain forms dynamic interactions with short ~30 amino acid transcription factor activation domains (ADs) that are unrelated in sequence but generally share net negative charge, hydrophobic, and aromatic residues that are typically intrinsically disordered (IDR) in

Figure 1. Identification of an acidic activation domain in NUT. **a**, Domain structure of p300 and **b**, BRD4-NUT constructs. **c**, IUPred, prediction of intrinsic disorder; FOLD, folding prediction using the foldindex (blue), PLAAC (**P**rion-**L**ike **A**mino **A**cid **C**omposition; orange) and PAPA (**P**rion **A**ggregation **P**rediction **A**lgorithm; gray) analysis, NCPR (**N**et **C**harge **P**er **R**esidue with a sliding window of 10 residues of the BRD4-NUT protein. **d**, SEC-MALLS analysis of the TAZ2 domain of p300 alone (black), NUT4 alone (blue) and the complex between TAZ2 and NUT4 (red). SDS-PAGE analysis of the peak fractions of the TAZ2-NUT4 complex is shown. **e,** Competition analysis: Cos7 cells were transfected with constructs expressing GFP-BRD4-NUT (green) alone or co-transfected with RFP-NUT-F1c and RFP-NUT4 (red). Immunofluorescence with anti-p300 antibody was used to visualize endogenous p300. Scale bars, 10 µm.

isolation (Dyson and Wright, 2016). Accordingly, sequence analysis revealed that the region of NUT is predicted to be disordered, and negatively charged with modest propensity to self-aggregate (Fig. 1c). A luciferase assay confirmed that the isolated NUT region can stimulate transcription (Fig. S1c), in agreement with previous data showing that tethering of NUT on chromatin is sufficient to induce transcriptional activation (Wang and You, 2015). We therefore conclude that NUT contains an acidic AD that directly interacts with the TAZ2 domain of p300 (Fig. 1b).

The high percentage of negatively charged amino acid residues in ADs is thought to prevent hydrophobic collapse while the distribution of charges maintains the conformational ensemble in a more expanded state leaving the charged and hydrophobic side chains exposed to the solvent and binding partners (Staller et al., 2018). Accordingly, analysis by size-exclusion chromatography–multi-angle laser-light scattering (SEC–MALS) showed that isolated NUT4 is monomeric and monodisperse in solution with a large hydrodynamic radius, consistent with an extended conformation (Fig. 1d). The TAZ2:NUT4 complex formed a 1:1 complex with a reduced hydrodynamic radius, indicative of conformational compaction upon binding (Fig. 1d).

Co-expression of NUT F1c can prevent condensate formation in the nucleus by acting as a dominant negative inhibitor for the p300-NUT interaction (Reynoird et al., 2010; Wang and You, 2015). Accordingly, overexpression of RFP-NUT F1c, -NUT4, or -NUT2 but not -NUT1 or -NUT3 reduced p300 foci in COS cells (Fig. 1e, Fig. S2g). As AD function is usually confined to short sequences, we further mapped the interaction by limited proteolysis followed by in-gel acid hydrolysis and mass spectroscopy (Fig. S1d). This led to an additional series of constructs that all bound the TAZ2 domain (Table 1). Fragments spanning amino acids 396-482 (NUT9) or a peptide spanning 458-482 (NUT10) all bound to the TAZ2 domain (Table 1). We conclude that the acidic AD of NUT interacts with the TAZ2 domain of p300 (Fig. 1b).

TAZ2-NUT interaction.
NMR spectra of NUT7, spanning 448-492, feature little dispersion of ^{1}H resonances (Fig. 2a), indicating that the protein is largely disordered in solution. We used a series of 3D experiments (see Methods) to assign backbone resonances of NUT7. Secondary structure propensities (SSP) calculated using ^{13}Cα, ^{13}C' and ^{15}N chemical shifts show that amino acids 451-472 have a significant tendency to sample helical conformations in solution (Fig. 2b). We conclude that the NUT7 AD is an IDR in isolation with a pre-organised N-terminal α helix.

We monitored binding to the TAZ2 domain by collecting a series of ^{1}H, ^{15}N-HSQC spectra of ^{15}N-labeled NUT7 mixed with increasing concentrations of TAZ2 (Fig. 2a). Chemical shift perturbations (CSP) showed that several NUT7 backbone amide resonances, including A450, L451, G460-A464, V467 and E475 were perturbed upon addition of an equimolar amount of TAZ2 while resonances from the C-terminus of NUT7 do not appear to shift significantly upon addition of TAZ2 (Fig. 2c). Thus, the N-terminal α-helical segment of NUT7 is involved in TAZ2 binding. Addition of equimolar amounts of unlabelled NUT7 to ^{15}N-labeled TAZ2 showed significant CSPs of resonances corresponding to residues R1737, I1739, M1761, R1763, V1765, Q1766, T1768, K1783, Q1784, I1809 (Fig. 2a, c), that are mostly located on the surface of TAZ2.

Figure 2. TAZ2:NUT interaction. **a**, (Left) Overlay of ^{1}H-^{15}N HSQC spectra of free TAZ2 (black) and after addition of one molar equivalent of NUT7 (448-492) (blue). Arrows indicate chemical shift changes for individual cross-peaks. (Right) Overlay of ^{1}H-^{15}N HSQC spectra of free NUT7 (448-492) and after adding half (orange) and one (brown) molar equivalent of TAZ2. Arrows indicate chemical shift changes for individual cross-peaks. **b**, Secondary structure propensities (SSP) plot of the NUT (448-492) fragment in its free state. positive values on the Y-axis indicate a-helical propensity, the X-axis is the amino acid sequence. **c**, (Left) Chemical shift perturbations on TAZ2 upon adding one molar equivalent of unlabeled NUT7 (448-492). (Right) Chemical shift perturbations on NUT7 (448-492) upon adding one molar equivalent of unlabeled TAZ2. **d**, (Left) Ribbon representation of lowest energy NMR model of the TAZ2 (blue)-NUT (orange) complex. Hydrophobic residues on the surface of NUT are shown. (middle) Zoom-in view of the interface between NUT (orange) and TAZ2 (blue). Residues on both sides implicated in the interaction as well as the hydrogen bonding network are shown. (Right) acidic residues on the surface of NUT (top) as well as hydrophobic and basic residues on the surface of TAZ2 (bottom) are shown. **e,** TAZ2 and NUT7 (448-492) are surface rendered and colored according to electrostatic potential (left) and sequence conservation (right).

To understand the binding mode, we measured both ^{13}C- and ^{15}N-HSQC-NOESY spectra on a sample of ^{15}N,^{13}C-labelled TAZ2 to which one equivalent of unlabelled NUT7 was added. We also measured ^{15}N-HSQC-NOESY spectra on ^{15}N-labelled NUT7 in admixture with one equivalent of unlabelled TAZ2. The NOESY spectra contain very few cross-peaks that can be unambiguously assigned to intermolecular contacts most likely due to extensive dynamic fluctuations of NUT7 even in the bound state. One contact is made between the backbone amide hydrogen of S1726 in TAZ2 and the backbone amide hydrogen of residue L463 and the Hβ of A464 in NUT7. Because of relative sparsity and ambiguity of structural restraints, we used HADDOCK (Dominguez et al., 2003) to derive a structural model of the interaction consistent with the observed spectroscopic changes (Fig. 2d and e). NMR restraints and structural statistics are summarized in Table 2. On the basis of this model, it appears that the N-terminus of NUT7 folds into an amphipathic helix that binds a hydrophobic groove in the interface between helices α1/α2/α3 on the surface of TAZ2 (Fig. 2d). Only the NUT backbone residues P448 to A464 are well ordered while residues at the C-terminus (Q465-D492) do not interact with TAZ2 and remain disordered. A set of hydrophobic side chains of NUT L449, A450, L451, L455, L461 are in contact with TAZ2 residues I1735, M1761, V1764, I1781 (Fig. 2d). The interaction appears to be further stabilised by electrostatic interactions involving NUT residues E453, E454, E456, E458, E459 and TAZ2 residues R1737, S1754, S1757 and K1760 (Fig. 2d). In order to test our model, we mutated amino acids that, according to the model, are not interacting with TAZ2 (E459A, L466A and L474A), These mutations have very limited impact on the HSQC spectrum of TAZ2 in complex with NUT7, indicating that the mutant has the same binding mode as the wild-type construct (Fig. S3d). In summary, similar to other TAZ2:AD interactions, a positively charged TAZ2 domain interacts with a negatively charged NUT AD (Fig. 2e). The key interfacial binding surfaces are conserved in p300 and NUT (Fig. 2e).

Analysis of the TAZ2:NUT interaction
As ADs are frequently bi-partite and the NUT7 fragment in our structure only partially engages the TAZ2 binding surface, we further explored the region N-terminal of NUT7. We found using HSQC competition, that the NUT7 fragment is unable to displace p53 from the surface of TAZ2 when added to the complex (Fig. S3e). Biolayer Interferometry (BLI) experiments showed that NUT8 (396-470) bound to TAZ2 with a dissociation constant of K_d = 0.96 +/- 0.29 μM (Fig. 3a, b). Alanine substitutions of conserved residues in the region spanning 447-462 (mut3, Fig. 3c), the main binding epitope in NUT7, did not abolish TAZ2 binding in GST-pulldowns, but reduced TAZ2 binding affinity (Fig. 3b, Fig. S3a,b). Additional Alanine substitution of L436, L440, L441 (mut4, Fig. 3c) further reduced TAZ2 affinity (Fig. 3b). In contrast, binding was largely abolished by additional mutation to Alanine of L410, L411, Y413, L427, C428, F423, V424, V427, V430 and I431 (mut1, Fig. 3b-d, Fig. S3a). As alanine repeat expansion can alter transcription factor condensation properties (Basu et al., 2020), we also generated a NUT variant containing mostly Glycine, Serine and charge reversal mutations (mut2, Fig. 3c). Mut2 also failed to bind TAZ2 in BLI and GST-pulldown experiments (Fig. 3b-d, Fig. S3a).

As mutation of isolated binding epitopes reduced but did not abolish TAZ2 binding, we conclude that the NUT AD is made up of subdomains that contain short amphipathic interaction motifs that synergise to enable high affinity binding. TF ADs frequently bind p300 through a minimal ΦXXΦΦ motif (X is any residue and Φ is hydrophobic), frequently LXXLL (Plevin et al., 2005). The NUT AD contains several such conserved motifs most of which are predicted to engage TAZ2 (Fig. 3c,e).

Figure 3. Analysis of the TAZ2-NUT interaction. a, Representative Biolayer Interferometry (BLI) profiles of p300 TAZ2-NUT binding from which the steady-state data were derived. **b**, Steady state analysis of p300 TAZ2 interaction with wild-type and NUT mutants using BLI. Error bars represent the standard deviations from three technical replicate experiments. **c**, Sequences of the NUT AD and mutants (in bold red) used. Top: Skylign plot showing NUT AD sequence conservation; Bottom: predicted secondary structure. **d**, GST pulldown analysis of TAZ2 and NUT8 (396-470) wild type and mutants 1 and 2. **e,** AlphaFold prediction of NUT (396-482) structure. **f**, (left) AlphaFold prediction of the TAZ2-NUT complex structure. Right: mutated residues in NUT mutant 1 (blue) and 2 (green) are shown.

In our structure, the isolated AD sub-domain in NUT7 is positioned at the interface of the TAZ2 α1, α2 and α3 helices, which frequently functions as a docking site for TF ADs that typically adopt helical structure upon binding (Fig. S3f). AlphaFold (AF) computational models predict 5 helices for NUT (Fig. 3e) that engage the protein interaction surfaces of TAZ2 (Fig. 3f, Fig. S3c). In the model, the NUT7 AD is re-positioned while the TAZ2 α1, α2, α3 surface is occupied by α3 of the NUT AD. Structure determination of the TAZ2 domain in complex with the entire NUT AD was not possible as NMR spectra suffered from severe broadening (data not shown). As (1) isolated AD sub-domains can interact with alternate TAZ2 surfaces (Krois et al., 2016) and (2) as the residues that impair binding are located in the predicted TAZ2-NUT binding interface, we believe this model provides a reasonable starting point for further structural characterization.

The role of the TAZ2 domain in p300 regulation and substrate acetylation
Previous data show that NUT stimulates p300 auto- and histone acetylation in cells and *in vitro* (Reynoird et al., 2010; Shiota et al., 2018). The AF model of p300 suggests that the TAZ2 domain is positioned over the p300 HAT active site (Fig. S4f, top). The TAZ2 C-terminal helix (amino acids 1806-1836) is sandwiched between the RING and HAT domains proximal to the lysine substrate-binding tunnel in a conformation that is incompatible with substrate access to the HAT active site. In contrast, a model in the presence of NUT AD shows that the TAZ2 domain is displaced from the HAT active site (Fig. S4f, bottom). One possibility is that the TAZ2 domain auto-inhibits the HAT domain and that this auto-inhibition is relieved upon ligand-binding to the TAZ2 domain. As the TAZ2 domain binds numerous transcription factor ADs, such a TAZ2-mediated control mechanism may be broadly relevant and a key aspect of how TFs control chromatin acetylation and gene regulation.

To test this model, we incubated increasing amounts of GST-NUT8wt with p300s (amino acids 324-2094) in the presence of [^{14}C] acetyl coenzyme A. Addition of NUT WT stimulated p300 auto- and NUT acetylation in a dose-dependent manner (Fig. 4a). Other TAZ2 ligands, Adenoviral E1A or the transcription factor p53 also showed a dose-depended stimulation of p300s auto-acetylation (Fig. 4c,d). In contrast, the TAZ2-binding mutant NUT8mut1 did not show stimulation of p300 auto-acetylation (Fig. 4a). Despite the identical number of lysine amino acids, NUT8mut1 was also not acetylated by p300 (Fig. 4a). These data suggest that the TAZ2 domain autoinhibits p300 activity, possibly through HAT active site restriction (Fig. S4f). Accordingly, acetylation reactions with the core domain of p300 (Fig. 1a; p300 core amino acids 1048-1664), lacking TAZ2, show that p300 autoacetylation is not stimulated by the presence of NUT8wt and both NUT8wt and NUT8mut1 substrates are acetylated at equal rates (Fig. 4b). We conclude that the interaction of TF ADs with the TAZ2 domain allosterically regulates p300 HAT activation by relief of autoinhibition.

Recent data indicate that p300 directly engages nucleosome core particles (NCPs) through its ZZ (Zhang et al., 2018) and TAZ2 domains (Sheahan et al., 2020). Would then TAZ2-engagement also orient NCPs with respect to the HAT active site and specify nucleosome acetylation patterns? In accordance with this possibility, it has been shown that the TAZ2 domain is required for efficient acetylation of H3K27, while curbing activity towards other lysine residues within nucleosomes (Sheahan et al., 2020). To test this model, we generated homogeneous 'structure-grade' NCPs as shown by single particle cryoEM (Fig. S4g, Table 3). p300 variants containing the TAZ2 domain, as well as the isolated TAZ2 domain directly bound to NCPs in EMSA experiments (Fig. 4g, Fig. S4d).

Figure 4. Role of the TAZ2 domain in p300 regulation and substrate acetylation. a, p300s or (**b**) p300 core were incubated for with [^{14}C]Acetyl-CoA in the presence or absence (C: p300 alone) of increasing concentrations of NUT8 (396-470) wild type or mutant1, (**c**) GST-E1a or (**d**) NCPs. Top two images: Analysis by SDS–PAGE followed by Coomassie staining and (bottom two images) ^{14}C phosphor imaging. Representative data of three independent experiments are shown. **e**, EMSA analysis of p300 core-CH3 binding to NCPs. Binding reactions were analyzed by native-PAGE and stained with Syber Safe (left) and Coomassie stain (right). **f,** Cross-link (XL-MS) analysis of the p300 core-CH3:NCP complex. Right: Graphical representation of the p300:histone and histones intermolecular cross-links. Left: p300 residues that cross-link to the NCP histones are shown in red on the p300 core (4BHW) and TAZ2 (3IO2) structures. **h**, Quantitative mass spectrometry of NCP acetylation. Reconstituted NCPs were incubated with Acetyl-CoA and p300s or the p300 core. Quantification of the site-specific histone H2B, H3 and H4 acetylation by label-free HPLC/MS/MS. Error bars represent standard deviations from two independent technical replicate experiments.

Cross-linking mass spectrometry experiments showed that surface residues from the p300 Bromo-, RING, PHD and TAZ2 domains directly engage the N-terminal regions of Histone H3 and H2A (Fig. 4f, Fig. S4h). This suggests that NCP engagement by the TAZ2 domain orients the enzyme on NCP substrate. To assess if this interaction contributes to substrate specificity, we assayed NCP acetylation using label free quantitative mass spectrometry. While p300 core, lacking TAZ2, acetylated most lysine residues indiscriminately in Histones H2B, H3 and H4, p300s showed a more selective acetylation pattern thus suggesting that direct NCP binding contributes modestly to substrate specificity (Fig. 4g).

Mechanism of condensation of BRD4-NUT and p300
BRD4-NUT forms hyperacetylated chromatin megadomains in patient tumors (Alekseyenko et al., 2015; Reynoird et al., 2010; Rosencrance et al., 2020). Formation of such megadomains can be reproduced by expression of BRD4-NUT in heterologous cell lines (Alekseyenko et al., 2015; Reynoird et al., 2010; Rosencrance et al., 2020). As there is strong evidence that this system reproduces the phenoptype seen in patient cells, we consider it as physiologically relevant, at least with respect to the biophysical properties of the condensation reaction. We generated a series of GFP-BRD4-NUT variants and assessed their propensity to interact with p300 and form co-condensates in the cell nucleus. Co-immunoprecipitation experiments showed that wild-type GFP-BRD4-NUT but not the AD mutants bound to endogenous p300 (Fig. 5a). In agreement with our *in vitro* data, the wild-type but not mutant variants showed a signal at the level of GFP-BRD4-NUT when probed with an anti-acetyl lysine antibody (Fig. 5a). GST-pulldowns in nuclear extracts using a purified GST-NUT8wt fragment showed signal for p300 only for the wild-type but not the AD mutant variants (Fig. 5b). Thus, mutation of the NUT AD interferes with p300 interaction and BRD4-NUT acetylation in cells, as also seen with the corresponding fragments *in vitro* (Fig. 4a).

Co-transfection experiments showed that GFP-BRD4-NUTwt and RFP-p300 formed nuclear co-condensates (Fig. 5c; row 1; for quantification see Table 4) both in co-transfected and RFP-p300 negative cells. Thus, condensation of GFP-BRD4-NUTwt not only concentrates but also depletes p300 from other areas of the nucleus. These condensates were independent of the NUT AD as mutant variants still formed condensates that co-localized with p300 (Fig. 5c; row 2&3). Immunofluorescence experiments showed that the NUT mutant condensates still co-localised with acetylated chromatin (Fig. S5a). Mutation of the acetyl-lysine binding pocket in BD1 (N140A) and BD2 (N433A) completely abolished GFP-BRD4-NUT and RFP-p300 co-condensation (Fig. 5c; row 4). To assess the model that BRD4 multivalency drives the condensation reaction (Gibson et al., 2019), we mutated BD1 and BD2 individually. We found that mutation of BD1 but not BD2 largely abolished condensation (Fig. 5c; row 5, 6). Thus, the acetyl-lysine binding ability of BD1 is required to nucleate BRD4-NUT and p300 co-condensation.

We reported previously that BRDT interacts with nucleosomes through its first (BD1), but not second (BD2) bromodomain, and that acetylated Histone recognition by BD1 is complemented by a direct BD1–DNA interaction (Miller et al., 2016). Another report indicates that DNA binding promotes condensation of a short BRD4S 'isoform' *in vitro* (Han et al., 2020). We therefore introduced charge inversion mutations in a conserved, basic surface patch in BD1 (R68E, K72E, K76E) that is involved in DNA binding in BRDT (Miller et al., 2016) and in the equivalent BD2 amino acids (S358E, K362E, A366E). Mutation of the basic patch abolished condensation (Fig. 5c; row 7). Thus, we propose that BRD4 BD-DNA

interaction facilitates acetylation-dependent NCP binding and chromatin condensation in cells.

We also probed the BRD4 N-terminal cluster of phosphorylation sites (NPS) and the basic residue-enriched interaction domain (BID), elements that have been proposed to enable BRD4 regulation by phosphorylation (Wu et al., 2013). Deletion of the NPS did not interfere with condensation while removal of the BID reduced condensation (Fig. 5c; row 8, 9). A minimal BD1-BD2 variant fused to the AD but not an AD Mut1 variant (Fig. 1b) still formed p300 co-condensates (Fig. 5c; row 10, 11, Movie S1). The BD1-BD2-AD fusion protein but not the AD Mut1 variant was still able to co-IP p300 from cells indicating that the ability to interact with p300 is maintained in the fusion protein (Fig. S5c). We therefore conclude that, at a minimal level, both BRD4 BD acetyl-lysine binding and p300 recruitment are required for condensation.

Addition of 1,6-hexanediol, the p300/CBP HAT inhibitor A-485 or the BET inhibitor JQ1 all abolished condensation of GFP-BRD4-NUTwt, mut1, mut2 and ΔNPS confirming that both acetylation by p300/CBP and BRD4 BD substrate engagement are required (Fig. 5d, Table 6). HDAC inhibition results in expansion of Histone acetylation and retargets lysine-acetyl readers including BRD4 thus partially mimicking the effect of BD inhibition (Slaughter et al., 2021). In agreement with this model, addition of the HDAC inhibitor Trichostatin A abolished condensation of all variants (Fig. 5d). Immunofluorescence experiments further showed that the BRD4-NUT condensates colocalise with HDAC1 thus confirming that these sites correspond to high acetylation turnover (Fig. S5b).

As mutation of the NUT AD does not abolish p300 recruitment into the condensate we further assessed the contribution of the CBP/p300 BD. We used the CBP30 inhibitor that has~34-fold higher selectivity for CBP/p300 BD as compared to BRD4 BD1 (Hammitzsch et al., 2015). Addition of CBP30 did not impact the condensation of GFP-BRD4-NUTwt and ΔNPS. However, condensation of the NUT AD mutants GFP-BRD4-NUTmut1 and mut2 was sensitive to CBP30 treatment (Fig. 5d, Table 7). Thus, the NUT AD mutants reveal that co-condensation can also occur through a p300 BD-acetyl-lysine-dependent recruitment mechanism. Thus, two interactions drive BRD4-NUT and p300 co-condensation: (1) interaction of p300 with the AD domain in NUT and (2) p300 BD-acetyl-lysine engagement. Disruption of both interactions, but not each interaction individually, abolishes the co-condensation reaction.

Condensation requires BD oligomerization
Our observation that mutation of BD1 but not BD2 largely abolished BRD4-NUT condensation in cells (Fig. 5c; row 5, 6) suggests that condensation in cells is not simply driven by BRD4 BD divalency. We therefore tested the requirements of acetylation-dependent chromatin condensation *in vitro*. Previous data show that addition of BRD4 or an engineered multivalent protein containing five copies (BD_5) of BRD4 BD1 induces condensation of p300-acetylated nucleosome arrays *in vitro* (Gibson et al., 2019). Condensation in this *in vitro* system depends on BD acetyl-lysine binding and multivalency (Gibson et al., 2019).

We generated dodecameric nucleosome arrays (Fig. S5d). In agreement with previous data (Gibson et al., 2019), non-acetylated nucleosome arrays formed round droplets that were dissolved by p300 acetylation (Fig. S5e), in agreement with the model that acetylation disrupts internucleosomal interactions (Shogren-Knaak et al., 2006). As expected, addition of

Figure 5. Mechanism of condensation of BRD4-NUT and p300. a, Cos7 cells were transfected with GFP-BRD4-NUT variants or GFP alone as the control. Co-immunoprecipitation (IP) experiments were performed using anti-GFP antibody and immuno-blotted (IB) for p300, GFP and Acetylated-Lysine. Input samples were loaded as a control. Experiments were independently repeated at least three times. **b,** Purified GST-NUT8 wild type and mutants 1 & 2 were incubated with Cos7 cells extract for 4 hours. The complexes were pull-down by Glutathione Sepharose beads and analysed by SDS-PAGE followed by Coomassie staining (bottom) or by IB for p300 (top). **c**, Cos7 cells were transfected with the indicated GFP-BRD4-NUT and RFP-p300 constructs and analysed by fluorescence microscopy, BRD4-NUT (green) and p300 (red). Scale bars, 10 μm. Quantification of foci formation of different BRD4-NUT variants and their co-localization with p300 are shown in Tables 4 and 5. **d**, Cos7 cells were transfected with the indicated GFP-BRD4-NUT constructs and treated with DMSO (control), 1,6-hexanediol, p300 HAT inhibitor A485, pan-BET inhibitor JQ1, HDAC inhibitor TSA or p300 bromodomain inhibitor CBP30. Scale bars, 10 μm. Quantification of foci formation in presence of CBP30 is shown in Table 6. **e**, Reconstituted chromatin containing 12 Nucleosome Core Particles (12x NCP). Fluorescence microscopy images of 12x NCP chromatin templates without and with acetylation by p300 and with GFP-labelled bromodomain-containing proteins. Scale bars, 10 μm.

GFP-bromo5 allowed droplet formation of acetylated but not of non-acetylated nucleosome arrays (Fig. 5e). This condensation reaction was driven by BD1-acetyl-lysine binding as a N140A mutation in the acetyl-lysine binding pocket abolished droplet formation (Fig. 5e, middle).

As a minimal BRD4 construct containing BD1-BD2 (amino acids 1-459) induced condensation in cells when fused to the NUT AD, we tested if BD1-BD2 divalency is sufficient to drive droplet formation *in vitro*. However, we found that BD1-BD2 did not enable droplet formation of acetylated or non-acetylated nucleosome arrays (Fig. 5e, right). As a consequence, mutation of the BD acetyl-lysine binding pockets (BD1 N140A-BD2; BD1-BD2 N433A or BD1 N140A -BD2 N433A) also did not have impact droplet formation under these conditions (Fig. S5f). We conclude that the BD divalency of BRD4 is not sufficient to drive condensation *in vitro*. As large-scale condensation in cells is sensitive to BD1 but not BD2 mutation (Fig. 5c), our model is that multivalency is established by BRD4 oligomerisation.

Dynamics of condensation of BRD4-NUT and p300
Time-lapse imaging showed that the condensates remained stable during the experiment (Fig. 6a, Movie S2). Addition of 1,6-hexanediol resulted in rapid and transient dissolution of the condensate (Fig. 6b). After reformation, a second addition of 1,6-hexanediol dissolved the condensates again illustrating their dynamic and transient nature. To study the consequences of the BRD4-NUT mutations on turnover of the condensate we performed Fluorescence Recovery After Photobleaching (FRAP) experiments. Wild-type GFP-BRD4-NUT and RFP-p300 recovered ~50% of signal after ~20 sec (Fig. 6c). Thus, while droplets appear stable at the mesoscale, the molecules are highly mobile at the nanoscale and in constant exchange with the nucleoplasm. Mutation of BRD4-NUT mostly interfered with droplet retention as shown by increased FRAP recovery rates (Fig. 6d). p300 dynamics was directly coupled to that of the BRD4-NUT mutants except for the GFP-BRD4-NUT BD2 and the BD1-BD2-AD constructs which showed slower p300 recovery rates indicating some degree of functional uncoupling.

To assay internal droplet dynamics, we performed half-bleach FRAP experiments (Fig. 6e). We analysed the change in fluorescence intensity over time along a defined A-B axis in the bleached (red square) and unbleached area. Kymograph analysis showed that the fluorescence signal in the 'unbleached' region (orange line) and the droplet 'center' (blue line) recovered much quicker and to completion as compared to the 'periphery' of the droplet (grey line), which only partially recovered signal (Fig. 6f). Our interpretation is that recovery at the center occurs more quickly because high local concentration in the droplet center maintains a feed-forward acetylation reaction that traps quickly diffusing molecules. At the droplet periphery, lower concentrations and competition with HDACs destabilizes droplet assembly leading to reduced recovery rates. As there is (1) constant exchange with the nucleoplasm and (2) no preferential equilibration between the bleached and unbleached area we argue that these condensates are 'liquid'-like at the nanoscale but not at the mesoscale where the viscoelastic properties of the 'solid' chromatin scaffold constrain dynamic exchange (Strickfaden et al., 2020). As a result, we do not see droplet fission or fusion at least during the time-course of our experiments.

Discussion
Chromatin acetylation has long been linked to transcriptional activation, by altering chromatin structure and by enabling binding of BD containing proteins (Dhalluin et al., 1999;

Figure 6. Dynamics of condensation of BRD4-NUT and p300. a, Example image of nuclear droplets. Red square: Time laps imaging of BRD4-NUT and p300 droplets; Scale bar, 4 μm. **b,** Time laps imaging after addition of 3% [w/v] 1,6 Hexanediol. Scale bar, 12 μm. **c,** Top: Example image of droplets (red square) after photobleaching. Scale bar, 2 μm. Bottom: Quantification of the FRAP experiments. 3 independent technical replicates were done. Averages and standard error of mean (SEM; grey bars) are shown. **d,** Quantification of the characteristic time of recovery rates for different GFP-BRD4-NUT mutants (top) and RFP-p300 (bottom). Three to four technical replicate FRAP experiments were done for each mutant (40 to 65 foci measured). Data shown are mean +/-standard error of mean (SEM). Statistical differences in   recovery rates were determined by Student's t-test. *p≤0.05; **p≤0.01; ***p≤0.001; n.s. non-significant difference as compared to WT. **e,** Half-bleach FRAP experiment. Top: The area before (left) and after (right) laser bleach are shown. The red box indicates the bleached area. Scale bar, 0.5 μm. Bottom: The change in fluorescence intensity along the A-B axis was analysed in the Kymograph. **f,** Top: Plot of fluorescence recovery from the Kymograph for the unbleached area, center and periphery of the droplet. Bottom: cartoon of the internal and external exchange.

Eberharter and Becker, 2002; Strahl and Allis, 2000). How acetylation promotes transcription has remained elusive. p300 is responsible for up to a third of nuclear acetylation (Weinert et al., 2018). The BRD4-NUT fusion reveals how constitutive activation of p300 establishes a self-organizing, acetylation-dependent feed-forward reaction that results in large-scale intra- and interchromosomal interactions with an oncogenic outcome. In addition to its well-known role in enhancer function and control of gene expression (Ma et al., 2021; Narita et al., 2021), p300 is also involved in compartmentalisation of the genome into eu- and heterochromatin (Cabianca et al., 2019). We argue that key aspects of these essential reactions involve modulation of an acetylation-dependent condensation reaction we describe here.

Regulation of p300 is central for establishment of such acetylation-dependent condensation reactions (Fig. 7a). We propose that in the deacetylated state p300 is auto-inhibited by the RING, TAZ2 domains and the AIL. p300 is activated by trans-autoacetylation (Karanam et al., 2006; Ortega et al., 2018; Thompson et al., 2004). Due to low p300 concentration(Gillespie et al., 2020), self-collision and activation do not occur spontaneously. Instead, as HDACs are estimated to be ~100-fold more abundant (Gillespie et al., 2020), p300 most frequently collides with HDAC complexes (Black et al., 2008; Hogg et al., 2021) which maintain p300 in the inactive conformation. In this inactive state p300 is evenly dispersed in the nucleus (Ortega et al., 2018).

Induced proximity, by TF binding, enables p300 self-collision, trans-autoacetylation and activation (Ortega et al., 2018). The BRD4-NUT system reveals an autoinhibitory role for the TAZ2 domain which is relieved upon NUT AD binding (Fig. 4). Accordingly, other well-characterized TAZ2 ligands including the ADs of E1A and p53 (Hsu et al., 2018; Krois et al., 2016; Pelka et al., 2009) also stimulate p300 auto- and substrate acetylation (Fig. 4), as also reported for other TAZ2 ligands (Hansson et al., 2009; Kraus et al., 1999). Thus, we propose that the TAZ2 domain plays a role in allosteric HAT regulation: TF AD binding results in displacement of the TAZ2 domain from its auto-inhibitory position resulting in HAT activation thus directly coupling TF AD-mediated p300 binding to HAT activation (Fig. S4f).

Perturbation of the autoinhibitory function of the RING and TAZ2 domains by mutation all lead to constitutive p300 activation and condensation reactions similar to what is seen with BRD4-NUT (Delvecchio et al., 2013; Ortega et al., 2018). Importantly, comprehensive studies of cancer cell lines show that truncation of the C-terminal helix of TAZ2, which we predict is required to stabilise the autoinhibitory conformation (Fig. S4f), also lead to p300 HAT gain-of-function in disease (Ghandi et al., 2019; Kumar et al., 2021). An acetylation-dependent condensation reaction may also explain how Adenoviral E1A causes genome-wide relocalisation of p300 and activation of genes that promote cell growth and division (Ferrari et al., 2014; Ferrari et al., 2008; Horwitz et al., 2008). Not unlike BRD4-NUT, this condensation reaction is also sensitive to p300 HAT, BD and HDAC inhibition (Ferrari et al., 2014). Therefore, while p300 is thought of as a classic tumor suppressor, with loss of function mutations occurring in a number of different cancers (Attar and Kurdistani, 2017), constitutive activation by mutation or by interaction with activating ligands such as BRD4-NUT and E1A can lead to acetylation-dependent condensation with an oncogenic outcome.

As numerous TFs interact with p300 trough their ADs (Dyson and Wright, 2016; Goodman and Smolik, 2000), we predict that the condensation reaction establishes intra- and interchromosomal interactions that are normally involved in enhancer signaling (Fig. 7b). Many TF ADs, including NUT (Fig. 1), are IDRs that form dynamic interactions with the different protein interaction domains of p300 (Dames et al., 2002; Feng et al., 2009; Ferreon

BRD4-NUT megadomain

Figure 7. Molecular model for p300 activation in physiological setting and disease. **a**, p300 is maintained in the inactive state by HDACs. Activation requires TF binding which results autoacetylation and removal of autoinhibition. **b,** In enhancer signalling, TFs interact through their ADs with the different protein interaction domains of p300 which results in scaffolding of dynamic TF assemblies. Local TF and histone acetylation allows binding of BD containing proteins including BRD4 thus establishing positive feed-back. HDACs are located in these condensates and enable dynamic turnover thus resulting in bistability and response to changes in TF signaling. **c,** In BRD4-NUT NMC, the acetylation-dependent reaction drives formation of intra- and interchromosomal interactions resulting in the formation of a Megadomain. **d,** TF binding and acetylation establish open Euchromatin. TFs compete for limiting p300 co-activators but TF-DNA binding outside enhancer clusters remains transient and, due to high HDAC activity, does not effectively establish an acetylation-dependent condensate. **e,** Depletion and partitioning of p300 into euchromatic compartments results in hypoacetylation due to high HDAC activity. Hypoacetylated compartments become a substrate for heterochromatinization.

et al., 2009; He et al., 2011; Krois et al., 2016; Lee et al., 2010; Marceau et al., 2019; Miller Jenkins et al., 2015; Mukherjee et al., 2013; Wang et al., 2012a; Wojciak et al., 2009). Due to such multivalent AD interactions, p300 is able to scaffold TF assembly on complex enhancer elements thereby facilitating combinatorial control of enhancer-mediated transcription (Chan and La Thangue, 2001; Panne, 2008; Panne et al., 2007).

TF-mediated recruitment and activation of p300 result in local histone and TF acetylation and establishment of positive feed-back through binding by BD containing proteins. High rates of self-collision of p300 maintain this positive feedback and establish a self-organizing feed-forward reaction resulting in co-condensation of p300, BET proteins BRD2, BRD3, and BRD4 and numerous transcriptional and chromatin regulators, which are all present in BRD4-NUT condensates (Alekseyenko et al., 2017). The condensation reaction is limited by HDACs which are usually present at active enhancers and promoters (Wang et al., 2009; Whyte et al., 2012), and prevent hyperacetylation and transcriptional inhibition (Gryder et al., 2019a; Gryder et al., 2019b). HDAC1 is localised in the BRD4-NUT condensate (Fig. S5b). Our FRAP experiments show that the condensates are 'liquid'-like at the nanoscale and propose that HDACs enable continuous and dynamic turnover of such condensates. If the feed-forward condensation reaction is sufficiently strong, it will overcome this inhibitory effect by HDACs provided that local high concentrations are maintained. As a result, the BRD4-NUT condensate is essentially stable at the mesoscale despite continuous internal dynamic exchange. In enhancer signaling, this dynamic bistability may then render the condensate more immediately responsive to dynamic and transient changes in TF signaling and explain the short transcriptional burst kinetics seen in cells (Fukaya et al., 2016; Li et al., 2019).

The reaction redistributes p300 (and BD containing proteins) into the condensate thus leading to depletion from other genomic regions (Fig. 5) (Reynoird et al., 2010). The condensation reaction thus leads to high local acetylation (in the condensate) but global histone hypoacetylation elsewhere (Schwartz et al., 2011). Counterintuitively, treatment with HDAC inhibitors does not lead to BRD4-NUT 'hypercondensation'. Instead, HDACi restores global histone acetylation (Schwartz et al., 2011) and rapidly dissolves the BRD4-NUT/p300 condensate (Fig. 5). In agreement with recent data (Slaughter et al., 2021), our model is that HDAC inhibition and resultant increased global histone acetylation redistributes BD containing proteins thereby decreasing their effective local concentration resulting in droplet dissolution. Due to the diffusible nature of the reactants, maintenance of global histone hypoacetylation is therefore a key aspect of how such condensates arise. We therefore argue that HDACs contribute to transcriptional regulation by maintaining (1) dynamic bistability of enhancer condensates and (2) global hypoacetylation which in turn 'enables' focal enhancer condensation. Thus, the dual nature of HDAC function with roles in gene activation and repression (Kidder and Palmer, 2012; Rundlett et al., 1996; Vogelauer et al., 2000; Wang et al., 2009) would then be directly linked to modulation of these condensation reactions.

While histone H3K27 acetylation has long been seen as a signature mark for active enhancers (Creyghton et al., 2010; Rada-Iglesias et al., 2010), p300 also acetylates numerous non-histone substrates including many TFs (Fig. 4) (Weinert et al., 2018). Therefore, we do not see a role for a defined 'code' in how such modifications contribute to the signalling reaction (Zhang et al., 2020). Instead, promiscuous lysine acetylation of TFs and histones drive emergent multivalency that kinetically traps diffusing BD-domain containing factors such as p300 and BRD4. While individual BDs typically have weak binding affinity and specificity (Filippakopoulos and Knapp, 2012), local proximity facilitates rebinding after dissociation

thus enabling an acetylation-dependent feed-forward co-condensation reaction. We propose that this biophysical mechanism enables formation of an 'exploratory' network of interactions that generates bistability by positive feedback and enables focal accumulation of regulatory factors on enhancers and establishment of long-range interactions that control promoter activity (Li et al., 2019). Successful establishment of this 'exploratory' network could then explain the long-standing observation that multiple TF DNA binding sites synergistically activate transcription without the requirement of physical interaction between TFs (Carey et al., 1990; Panne et al., 2007).

It has long-been known that TFs activate some genes while at the same time repressing others through competitive coactivator 'squelching' (De Bosscher et al., 2003; Gill and Ptashne, 1988; Guertin et al., 2014; Hottiger et al., 1998; Kamei et al., 1996; Meyer et al., 1989; Portuguez et al., 2019; Triezenberg et al., 1988). Accordingly, in BRD4-NUT, the sequestration of p300 inhibits p53 and the apoptotic cell response while knock-down of BRD4-NUT releases p300, and restores p53-dependent cell differentiation and apoptosis (Reynoird et al., 2010). Thus, we predict that competition for diffusible co-activators and successful establishment of a continual acetylation-dependent condensation reaction is a key regulatory control mechanism: only enhancers containing the appropriate set of clustered TF binding sites will effectively compete in establishing such an acetylation-dependent condensate. Due to high concentrations of HDACs (Gillespie et al., 2020) and rapid deacetylation kinetics across the genome (Hogg et al., 2021; Weinert et al., 2018), the acetylation-dependent condensation reaction is transient. It is such transient and dynamic acetylation that establishes euchromatic compartments without necessarily leading to transcriptional activation (Fig. 7d).

There is increasing evidence suggesting a role for LLPS of intrinsically disordered proteins in different nuclear processes (Han et al., 2020; Sabari et al., 2018; Shin et al., 2018). IDR-driven condensation reactions of chromatin-binding proteins including BRD4 and p300 have been implicated in gene regulation (Boija et al., 2018; Chong et al., 2018; Guo et al., 2019; Han et al., 2020; Ma et al., 2021; Sabari et al., 2018; Shin et al., 2018; Zamudio et al., 2019) and in disease (Ahn et al., 2021; Boija et al., 2021). As BRD4-NUT and p300 are largely disordered (Fig. 1; 72% and 42% predicted IDR, respectively) our assumption was that IDR-mediated interactions would contribute to the condensation reaction.

While the NUT AD region can indeed form spherical condensates *in vitro* as analysed by SAXS (Fig. S6b,c), we do not think that this is relevant for the condensation reaction in cells. Even a minimal BRD4 BD1-BD2 construct, removing a large fraction of IDR sequence from BRD4-NUT, enables condensation in cells provided it contains a functional AD (Fig. 5c). As an equivalent NUT AD mutant (that fails to bind to the TAZ2 domain) does not enable the condensation we exclude the possibility that residual structural disorder, e.g. the 200 amino acid linker between the BD1 and BD2 domains of BRD4 disorder drives the condensation reaction. We and others have shown that the condensation reaction is inhibited by (1) the BET BD inhibitor JQ1 (Reynoird et al., 2010; Wang and You, 2015), (2) mutation of the acetyl-lysine binding pocket of BRD4 BD1 or, (3) the DNA binding surfaces of BRD4 BD1/BD2 or, (4) p300 HAT inhibition (Fig. 5). In addition, several p300 HAT gain-of-function mutations, that interfere with the autoinhibitory function of the RING and TAZ2 domains, enable a similar acetylation- and BD-dependent condensation reaction (Ortega et al., 2018). We therefore conclude that the reaction is driven by p300 activation and engagement of acetylated chromatin by the BDs. In accordance, single-gene imaging data support the notion that promoter-enhancer communication and transcription control depend

on TF and BRD4 clustering driven by specific DNA and BD binding rather than IDR-mediated interactions (Li et al., 2020).

Further, as (1) divalent BD1-BD2 did not enable condensation of acetylated nucleosome arrays *in vitro* (Fig. 5), and (2) as of mutation of BRD4 BD1 is sufficient to abolish condensation in cells, our model is that the condensate does not arise due to BRD4 BD1-BD2 divalency. Instead, as BET proteins form higher-order oligomers on acetylated chromatin (Pivot-Pajot et al., 2003; Wang et al., 2012b), our model is that BD-mediated oligomerisation and emergent multivalency drive condensation.

TFs locate their target DNA sites using 'facilitated diffusion' (Halford, 2009; Mirny et al., 2009). We propose that following TF-DNA binding, the combined action of p300 scaffolding, TF acetylation and formation of an acetylation-dependent condensate slows down the dynamic search behaviour thus increasing local concentration, residency time and transcription activation. Accordingly, acetylation-mediated increase of TF residence time has been described for p53 (Loffreda et al., 2017; Luo et al., 2004) and could thus control TF turnover and transcriptional regulation (Gurdon et al., 2020; Lickwar et al., 2012). Binding events that occur outside TF enhancer clusters are then unable to compete for limited co-activators and fail to establish such an acetylation-condensation reaction and therefore remain largely dynamic and transcriptionally inactive (Fisher et al., 2012; Kribelbauer et al., 2019; Li et al., 2008).

The co-activator depletion reaction may also be one of the key reaction mechanisms that enables the physical segregation of transcriptionally active euchromatin from repressed heterochromatin (Fig. 7e), and explain the general link between the transcriptional state and 3D chromatin compartmentalisation (Rowley and Corces, 2018). Targeting of heterochromatin with acidic TF ADs or treatment with HDAC inhibitors generally counteracts heterochromatinization even in the absence of active transcription (Ahmad and Henikoff, 2001; Aparicio and Gottschling, 1994; Carpenter et al., 2005; Chambeyron and Bickmore, 2004; Mueller et al., 2001; Taddei et al., 2001; Therizols et al., 2014; Toth et al., 2004; Tumbar et al., 1999a; Tumbar et al., 1999b). TF-mediated retention of p300 in euchromatin enables establishment of heterochromatin in worms (Cabianca et al., 2019), while depletion of p300 and chromatin hypoacetylation facilitates methylation by PRC2 (Hogg et al., 2021). Thus, competition for and partitioning of limiting key acetylases by TFs into euchromatin and depletion from heterochromatin, together with BD-mediated higher order chromatin folding (Gaucher et al., 2012; Pivot-Pajot et al., 2003; Wang et al., 2012b), may drive key aspects of 'epigenetic' genome compartmentalization. Phenomena such as stochastic and exclusive choice of monoallelic gene expression or X-chromosome inactivation in mammals may then also depend on depletion of limiting coactivators and establishment of positive feedback so that only the 'winning' allele drives gene expression while 'competing' alleles are silenced (Apostolou and Thanos, 2008; Jeppesen and Turner, 1993; Monahan et al., 2019; Spilianakis et al., 2005; Zylicz et al., 2019).

Methods

Constructs. cDNA encoding residues for NUT-F1c (347-588), NUT1 (347-446), NUT2 (427-527), NUT3 (508-588), NUT4 (347-480), NUT5 (347-514), NUT6 (351-470), NUT7 (448-492) (wild type and mutants), NUT8 (424-492) (wild type and mutants), NUT9 (395-482) and p53 (1-61) were cloned into the *Nco*I/*Xho*I restriction sites of vectors pETM-30 or pETM11 (EMBL) with a tobacco etch virus (TEV)-cleavable N-terminal His$_6$ glutathione S-transferase (His$_6$-GST) tag for pETM30 or an N-

terminal His$_6$-tag for pETM11. The p300 TAZ2 domain constructs (TAZ2 1723-1812, TAZ2 1723-1822, TAZ2 1723-1836) were codon optimized and cloned into the *NcoI*/*BamH*I restriction sites of pETM-11 (EMBL) with a TEV-cleavable N-terminal His$_6$-tag. Cysteines which do not participate in zinc coordination were mutated to Alanine or Serine (C1738, C1746, C1769, C1770). The CH3 domain of p300 (1660-1815) was cloned using *EcoR*I/*Xho*I restriction sites in pGEX-5X1 vector with a TEV-cleavable N-terminal glutathione S-transferase (GST) tag. p300s (324-2094) and p300 core-CH3 (1043-1828 and 1043-1817) were produced as described previously (Delvecchio et al., 2013). The BRD4 constructs: GFP-BD1-BD2, GFP-BD1-BD2 N140A/N433A, GFP-BD1-BD2 N140A and GFP-BD1-BD2 N433A were cloned into the *NcoI*/*Xho*I restriction sites of pETM11. The vectors expressing the GFP-(Bromo)5 and the GFP-(Bromo)5 N140A were gifts from Michael Rosen lab (University of Texas Southwestern). The co-expression vector encoding the *S. cerevisiae* histones was a gift from Stephen Harrison lab (Harvard Medical school). The vector expressing the E1A protein was a gift from Andy Turnell lab (University of Birmingham). For cell-based assays, all the GFP-tagged BRD4-NUT variants (Table 2), RFP-tagged NUT-F1c, NUT1, NUT2, NUT3 and NUT4 and RFP tagged p300 were cloned into pEGFP-C1 vector (Clontech). All constructs were confirmed by DNA sequencing.

Expression and purification. Expression and purification of Flag-tagged p300s and GST-tagged p300 core-CH3 constructs was done as reported previously (Delvecchio et al., 2013; Ortega et al., 2018). All p300 TAZ2 constructs were produced as inclusion bodies in *Escherichia coli* (*E. coli*) BL21 (DE3). Cells were grown to mid-log phase (OD600 ~ 0.6) at 37°C in Luria Broth containing 50 μg/ml kamamycin and 200 μM ZnSO$_4$. Protein expression was induced by addition of 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) at 18°C for 16h. Uniformly ^{15}N- and ^{15}N/^{13}C-labelled samples were produced from cells grown in M9 minimal medium containing 50 μg/ml Kanamycin, 200 μM ZnSO$_4$ and 1 g/l of ^{15}NH$_4$Cl and 2g/l ^{13}C glucose. Cells were harvested by centrifugation for 20 mins at 5000 g in a SLC-6000 rotor (Sorvall) and resuspended in 50 ml/L culture lysis buffer 1 (20 mM Tris-HCl pH 7.5, 300 mM NaCl, 5 μM ZnSO$_4$, 1 mM Dithiothreitol (DTT)), supplemented with EDTA-free protease inhibitors (Roche), and 10 μl of benzonase nuclease (Sigma-Aldrich). Cells were sonicated and centrifuged in a SS-34 rotor (Sorvall) for 1h at 12000 rpm. The pellet was then washed three times in the lysis buffer and the final pellet resuspended in buffer 2 (20 mM Tris pH 7.5, 300 mM NaCl, 1 mM Dithiothreitol (DTT), 6 M urea) for one hour at 4 °C on a rotating device. The protein solution was centrifuged as above and the soluble fraction applied to Ni^{2+}-charged IMAC Sepharose 6 FF (Cytiva) beads preequilibrated with buffer 2 containing 20 mM Imidazole. The column was washed with 20 column volumes (CVs) of buffer 2 containing 20 mM Imidazole and the bound protein eluted with the buffer 2 containing 300 mM Imidazole. The protein was refolded by dialysis overnight against buffer 3 (20 mM Tris pH 7.0, 100 mM NaCl, 200 μM ZnSO$_4$ and 2 mM DTT).

The refolded TAZ2 was incubated with His-tagged TEV protease (1:100 w/w) for 14-16 h at 4 °C. Subtractive Ni-NTA chromatography was used to remove the residual His-tag and TEV protease. The cleaved TAZ2 was further purified on a 5ml HiTrap SP HP cation exchange column (Cytiva) preequilibrated in buffer 4 (20 mM Tris pH 7.0, 100 mM NaCl, 5 μM ZnSO$_4$ and 1 mM Dithiothreitol (DTT)) and eluted using a 20 CV NaCl gradient. The peak containing TAZ2 was further purified using a HiLoad 16/60 Superdex 75 column (Cytiva) preequilibrated with buffer 5 (20 mM Tris pH 7.0, 300 mM NaCl, 5 μM ZnSO$_4$ and 1 mM DTT). The purified protein was concentrated using a prewashed Amicon Ultra-10 concentrator (molecular weight cut off 10 kDa; EMD Millipore), flash frozen in liquid N2 and stored in -80°C.

Unlabeled, ^{15}N- and ^{15}N/^{13}C-labelled NUT fragments were produced as insoluble inclusion bodies in *E. coli* and purified as described above. After Ni-NTA purification, the protein was dialyzed and refolded overnight at 4°C against 1L buffer 5 (20 mM Tris pH 7.0, 300 mM NaCl, 1 mM DTT and 10% glycerol). Refolded His-GST-NUT fusion proteins were incubated with His-tagged TEV protease (1:100 w/w) for 14-16 h at 4 °C. Subtractive Ni-NTA chromatography was used to remove the residual His-GST-tag and TEV protease. The cleaved NUT proteins were applied to a 5ml HiTrap Q HP anion exchange column (Cytiva) in buffer 4 without ZnSO$_4$ and eluted in a 20 CV NaCl

gradient. The peak containing NUT was further purified by gel filtration on a Superdex 75 Increase 10/300 GL column (Cytiva) equilibrated in buffer 5 without $ZnSO_4$. The final proteins were concentrated in a prewashed Amicon Ultra-15 Centrifugal filter (molecular weight cut off 3 kDa; EMD Millipore), flash-frozen in liquid nitrogen and stored at -80 °C.

The GST-E1A amino acids 1-289 and GST-p53 AD amino acids 1-61 proteins were expressed in *E. coli* BL21 (DE3) cells. Cell pellets were resuspended in lysis buffer resuspended and sonicated in lysis buffer (20 mM Tris-HCl pH 7.5, 300 mM NaCl, 0.5 mM TCEP), supplemented with EDTA-free protease inhibitors (Roche), and 10 µl of benzonase nuclease (Sigma-Aldrich) and lysed by sonication. The lysate was clarified by centrifugation for 1h in a SS-34 rotor (Sorvall) at 12000 rpm and applied to a Glutathione Sepharose 4 Fast Flow resin according to instructions from the manufacturer (Cytiva). The resin was washed with 20 CV lysis buffer and the protein was eluted in lysis buffer containing 20 mM reduced L-Glutathione. The protein solution was then dialyzed overnight against lysis buffer to remove L-Glutathione and then applied to a 5 ml HiTrap Q HP anion exchange column (Cytiva) and further purified by gel filtration on a Superdex 75 Increase 10/300 GL column (Cytiva) as previously described for the NUT fragments.

The BRD4 GFP-BD1-BD2, GFP-BD1-BD2 N140A/N433A, GFP-BD1-BD2 N140A and GFP-BD1-BD2 N433A were all expressed in *E. coli* BL21 (DE3) cells. The cells were sonicated in lysis buffer (20 mM Tris-HCl pH 7.5, 300 mM NaCl, 0.5 mM TCEP) and the lysate was clarified as previously described for the other proteins. The clarified lysate was applied to Ni^{2+} charged IMAC Sepharose 6 FF (Cytiva) beads preequilibrated with lysis buffer containing 20 mM Imidazole. The column was washed with 20 column volumes (CVs) of lysis buffer and the bound proteins eluted with the lysis buffer containing 300 mM Imidazole. Imidazole was removed by dialysis overnight against 1L lysis buffer and the proteins were loaded on a 5 ml HiTrap SP HP cation exchange column (Cytiva) as described for TAZ2 and further purified using a HiLoad 16/600 Superdex 200 column (Cytiva) preequilibrated with 20 mM Tris pH 7.0, 300 mM NaCl and 1 mM DTT. The purified proteins were concentrated using a prewashed Amicon Ultra-10 concentrator (molecular weight cut off 30 kDa; EMD Millipore), flash frozen in liquid N2 and stored in -80°C.

The GFP-(Bromo)5 and GFP-(Bromo)5 N140A were purified as described previously (Gibson et al., 2019). The co-expressed *S. cerevisiae* histones octamer was purified as described in Migl et al. (Migl et al., 2020). NUT peptides were ordered from Proteogenix.

HAT assays. The p300 autoacetylation and substrate acetylation assays were performed using [^{14}C] acetyl-CoA (Perkin-Elmer). Autoacetylation of p300 was quantified by autoradiography after SDS-PAGE gel analysis. The acetylation reaction was performed for 30 minutes at 30°C in 1x HAT buffer (20 mM Tris-HCl pH 7.5, 100 mM NaCl, 1 mM DTT, 10 % glycerol and 1x Complete EDTA-free protease inhibitor (Roche)) with 2.5 µM p300s or p300 core and 50 µM [^{14}C] acetyl-CoA with or without increasing concentrations of the different substrates (NUT fragments, NCP, E1A). 5 µl of the reaction was quenched by addition of 2.5 µl of 3x SDS gel loading buffer followed by analysis on a 4-20% SDS-PAGE gel. The gels were stained with Quick Coomassie Stain (Generon), fixed for 30 min in enhancer solution (KODAK ENLIGHTNING Rapid Autoradiography Enhancer, Perkin Elmer) and dried for 2 hours using a Bio-Rad Gel Dryer. The ^{14}C signal was quantified using phosphor imaging (Typhoon, Cytiva).

***In vitro* nucleosome acetyltransferase assays.** Nucleosome acetylation reactions were performed in 20 µl volume in acetylation buffer (20 mM Tris pH 7.5, 100 mM NaCl, 1 mM DTT, 10% glycerol and complete Protease Inhibitor EDTA-free (Roche)) with 50 µM Acetyl-CoA, 2.5 µM of p300s or p300 core and 2 µM of nucleosome core particles (NCPs). Control experiments contained only NCPs and Acetyl-CoA without p300. The reactions were incubated for 30 minutes at 30°C and stopped by addition of 1x SDS loading buffer. Each condition was performed in duplicate. The samples were then analyzed by SDS-PAGE and mass spectrometry.

Multi angle laser light scattering-size exclusion chromatography. SEC-MALLS was performed at a flow rate of 0.5 ml/min on a Superdex 75 Increase 10/300 GL column equilibrated in buffer (20 mM HEPES pH7.0, 300 mM NaCl, 5 µM ZnSO$_4$, 0.5 mM TCEP). 50 µl of TAZ2, NUT4 or the TAZ2:NUT4 complex at 2 mg/ml were injected onto the column and multi angle laser light scattering was recorded with a laser emitting at 690 nm using a DAWN-EOS detector (Wyatt Technology Corp). The refractive index was measured using a RI2000 detector (Schambeck SFD). The molecular weight was calculated from differential refractive index measurements across the center of the elution peaks using the Debye model for protein using ASTRA software version 6.0.5.3.

Biolayer interferometry assay. BLI measurements were performed with a ForteBio Octet RED384 instrument and ForteBio biosensors. Data analyses was done using ForteBio Data Analysis 9.0 software. Kinetics assays were carried out at 25 °C using settings of Standard Kinetics Acquisition rate (5.0 Hz, averaging by 20) at a sample plate shake speed of 1000 rpm. The GST-tagged NUT8 wild type and mutants were loaded onto Anti-GST biosensors and the TAZ2 protein was used as an analyte. The GST-NUT-loaded Anti-GST biosensors were dipped in kinetic buffer (1x PBS, 0.1% BSA, 0.02 % Tween 20) to establish a baseline time course and then dipped into wells containing TAZ2 (1723-1836) at various concentrations diluted in the same kinetic buffer to monitor NUT-TAZ2 binding. The dissociation step was monitored by dipping the biosensors back into the wells used to collect the baseline time course. To monitor binding due to nonspecific interactions of TAZ2 with the biosensors, Anti-GST biosensors were used as controls for every analyte concentration and a double referencing method was used for data subtraction and analysis. The subtracted binding curves were analysed and plotted as a steady-state response to obtain the dissociation constant K_d value.

Limited proteolysis. To identify stable variants of NUT we performed limited proteolysis by Trypsin and Chymotrypsin. 100µl of the purified CH3-NUT-F1c complex (250 µg) and TAZ2-NUT4 complex (250 µg) were incubated at 4 °C with Trypsin or Chymotrypsin with a ratio of 1:100 (w/w). At the indicated time points (30 min, 2 hours, overnight), 30 µl of the reaction was quenched by addition of 15 µl of 3x SDS gel loading buffer followed by analysis on 4-20 % SDS-PAGE gel. The stable fragments on gel were analysed by acid hydrolysis and mass spectrometry at the Proteomics Core Facility (EMBL, Heidelberg).

SAXS analysis. X-ray scattering data were collected on the NUT-F1c sample at 100 µM in phase separation buffer: 20 mM Tris pH 7.2 mM DTT, 125 mM NaCl and 10% (w/v) PEG 6000 at the B21 BioSAXS beamline of the Diamond Light Source (Chilton, Oxfordshire, UK). The data were analysed with the ATSAS package (Franke et al., 2017). Data collected from the sample were substracted from buffer scattering. R_g values were obtained from the Guinier approximation $sR_g < 1.3$ using Primus (Petoukhov et al., 2012). Distance distribution functions p(r) were computed from the entire scattering curve using GNOM (Petoukhov et al., 2012).

***In vitro* NUT phase separation assay.** Phase separation of purified GST-NUT-F1c *in vitro* was done phase separation buffer. The concentrations of GST-NUT-F1c used were 0-200 µM. A GST control was used in the same concentration range under identical conditions. The droplets were visually observed and turbidity of the sample (OD$_{600}$) quantified using NanoDrop (Thermo scientific).

GST pulldowns. For GST pulldowns, 50 µg of GST-tagged NUT constructs were mixed at an equimolar concentration with binding partners in 50 µl pulldown buffer (20 mM HEPES pH 8.0, 300 mM NaCl, 0.5 mM TCEP, 5µM ZnSO$_4$, 0.1% Triton X-100) containing 25 µl of a 50 % slurry of GST Sepharose 4 Fast Flow beads (Cytiva) per reaction. Reactions were incubated for 30 min at 4 °C. Twenty-five microliters of the reaction were withdrawn as the reaction input and the remainder was washed 5 times with 500 µl of pulldown buffer. Samples were boiled in 1x sample loading buffer for 5 min and analysed by SDS-PAGE.

Luciferase reporter assays. Cos7 cells were co-transfected for 36 hours with β-Galactosidase reporter plasmid and UAS-promoter luciferase reporter plasmid either alone or with GAL4 DNA

binding domain (DBD) plasmid or increasing amount of GAL4-DBD-NUT-F1c plasmid. Cells were then washed and incubated 15 min at RT with Luciferase lysis buffer (1% Triton X-100, 10% Glycerol, 2 mM EDTA, 25 mM Tris-HCl pH 7.8). Luciferase activity was measured directly after mixing 10µl of cellular extract with 50 µl of Luciferase substrate (Luciferase assay kit, Stratagene). Normalization was made by measuring the β-Gal activity after 1-hour incubation of 7.5 µl of cellular extract with 50 µl of β-Gal substrate (β-Gal detection kit II, Clontech).

Proteomics of histone modifications. For the identification and quantification of histone acetylation following *in vitro* HAT assays, we followed the previously published procedure (Goudarzi et al., 2016; Shiota et al., 2018). In brief, reconstituted NCPs were incubated with 100µM acetyl-CoA, as well as purified p300s (340-2,094) or the p300 core domain (1,043-1,666) for 30 min at 30°C. HPLC-MS/MS was used to quantify site specific H2A, H2B, H3 and H4 acetylation by label-free quantification. The peak area of each histone peptide was then normalized. Two technical replicate experiments were done.

Isolation and Purification of 1x601 Widom DNA. Carrier plasmids of 1x Widom 601 repeats were purified using QIAGEN Plasmid Giga kit from 6L LB culture according to the manufacturer's instructions. 1x arrays DNA were cut from carrier plasmid DNA by incubation overnight with 5,000 units of *Eco*RV-HF restriction endonuclease (New England Biolabs) in 1x CutSmart Buffer (20 mM Tris,OAc, pH 7.9, 50 mM KOAc, 10 mM Mg[OAc]$_2$, 0.1 mg/mL BSA). The excised DNA fragments were separated from the linearized plasmid by PEG precipitation. 0.192 volume of NaCl and 8% PEG 6000 were added to the DNA sample and incubated for 1h on ice, the sample was then centrifuged for 20 min at 4 °C, 27000 g. the *Eco*RV fragments contained in the supernatant were precipitated by the addition of 2.5 volumes of 100% cold ethanol. The DNA fragments were air dried for 10 min and dissolved in buffer TE (10 mM Tris/HCl, pH 8.0, 1 mM EDTA). Purity of the extracted 12x601 and 1x601 fragments were verified on 0.7% agarose gels and the concentration measured by 260 nm absorbance on a NanoDrop spectrophotometer (Thermo Scientific).

Isolation and Purification of 12x Widom 601 Array DNA. 12x601 DNA template was prepared as described previously (Dorigo et al., 2003). In short, 12-mer DNA with177 bp repeats of the 601 sequence separated from the rest of the plasmid by *Eco*RV sites were expressed in *E.coli* strain DH5a. The DNA was then extracted by alkaline lysis and purified on a Sepharose-6 column. The DNA template was separated from the plasmid by *Eco*RV-HF digestion and PEG6000 precipitation.

Histones octamer preparation. Wild type human histones (H2A, H2B, H3 and H4) and the cysteine mutant Xenopus H4 S47C were individually expressed in *E. coli* strain Rosetta 2 DE3 pLysS and purified following the protocol described (Luger et al., 1999), but using a RESOURCE S 6 ml cation exchange chromatography column. For octamer reconstitution, equimolar ratios of the four histones were refolded together in the presence of 5 mM β-mercaptoethanol. The octamer was then separated from dimers by gel filtration using a Superdex 200 column (Cytiva). Two batches of octamers have been generated: one fully wild-type human octamer (unlabelled octamer) and an octamer formed by wild-type human H2A, H2B, H3 and Alexa Fluor™ 594 C5 Maleimide labelled *Xenopus* H4 S47C (labelled octamer).

Fluorescent labelling of histones. The *Xenopus* H4 S47C mutant histone was labelled using Alexa Fluor™ 594 C5 Maleimide following the protocol from (Gibson et al., 2019). In short, lyophilised H4 S47C was resuspended in 7M urea, 500mM Tris-HCl pH 8 and incubated with Tris-neutralized TCEP to a final concentration of 2 mM for 1 hr and 30 min at RT. Alexa Fluor™ 594 C5 Maleimide stock solution was added dropwise to a 1.5 molar excess and the mix was incubated for 4 hr at RT in the dark. The reaction was quenched by the addition of β-mercaptoethanol. The histone was then purified from the free dye by ion exchange chromatography using a HiTrap SP HP cation exchange column followed by dialysis into ultra-pure water with 5 mM β-mercaptoethanol.

Mononucleosome assembly. DNA and *Saccharomyces cerevisiae* histones octamer were mixed in a 1:0.5 molar ratio. Nucleosomes were reconstituted by salt gradient dialysis method and dialysed against buffer 20 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM DTT with continuous decrease of NaCl concentrations (1.5, 1, 0.6 and 0.25 M) at 4 °C, 3h each dialysis step with the third dialysis step proceeding overnight using mini dialysis units (10,000 Da MWCO, Pierce Slide-a-lyzer). The reconstituted nucleosomes were quality controlled by electrophoretic mobility shift assays on 5% acrylamide native gels with 0.25x TBE running buffer.

12x Nucleosome array assembly, acetylation and phase separation. Chromatin was reconstituted *in vitro* by assembling the 12-mer DNA with histone octamers by step-wise salt dialysis as described previously (Dorigo et al., 2003), but using KCl instead of NaCl. The optimal octamer to DNA ratio was determined empirically. A mix of labelled and unlabelled histone octamer was used. buffer. The ratio of labelled to unlabelled histone octamer was also determined empirically. The reconstituted 12-mer nucleosome arrays were quality controlled by *Sca*I restriction enzyme digestion to isolate single nucleosome particles and by electrophoretic mobility shift assays on 1% agarose gel in 0.5x TAE 12x601 nucleosome arrays at 2 μM mono-nucleosome concentration were incubated with 1 μM of the p300 core domain in Acetylation buffer (20 mM Tris-HCl pH 7.0, 100 mM NaCl, 10% glycerol, 1 mM DTT) for 30 min at 30 °C. The non-acetylated chromatin reactions were incubated under identical conditions but without p300 core enzyme. After acetylation, 5 μM BRD4 GFP-Bromo$_5$, GFP-Bromo$_5$ N140A, 10 μM GFP-BD1-BD2, 10 μM GFP-BD1-BD2 N140A, 10 μM GFP-BD1-BD2 N433A or 10 μM GFP-BD1-BD2 N140A/N433A were added to nucleosomal arrays for a final concentration of 1 μM mono-nucleosome in phase separation buffer (25 mM Tris, pH7.5, 150 mM KOAc, 1 mM MgOAc, 5% glycerol, 5 mM DTT, 0.1 mM EDTA, 0.1 mg/mL BSA) (Gibson et al., 2019).

Microscopy of chromatin droplets. For confocal microscopy image acquisition, a Zeiss LSM 980 Airyscan 2 microscope equipped with a Plan-Apochromat 63x/1.40 Oil DIC f/ELYRA and ZEN Blue 3 acquisition and analysis software was used. Samples were imaged in 384 well glass bottom plates for microscopy which were prepared as described in (Gibson et al., 2019). In short, microscopy plates were washed with 5% Hellmanex, etched with 1 M NaOH and treated with 25 mg/mL 5K mPEG-silane. Prior to use, individual wells were passivated by incubation with freshly prepared 100 mg/mL BSA.

Cross-linking mass spectrometry. Purified solutions of both NCPs and p300 core-CH3 (1043-1817) were diluted to a concentration between 0.5 and 1 mg/ml in 20 mM HEPES pH 7.0, 300 mM NaCl, 1 mM DTT and 5 μM ZnCl$_2$ and cross linked using a homobifunctional, isotopically-coded crosslinker DiSuccinimidyl Suberate DSS-H12/D12 (Creative Molecules Inc.) at a concentration of 2 mM for 30 min at 35°C 600 rpm. The cross-linking reaction was then quenched with 0.1 volumes of 1M Ammonium bicarbonate for 10 min at 35 °C 600 rpm. The sample was then treated with 0.8 volumes of 10 M urea and 250 mM Ammonium bicarbonate as well as 0.05 volumes RapiGest SF Surfactant (Waters, cat. No. 186008090) and sonicated for 1 min in an ultrasound bath. DTT was later added with a final concentration of 10 mM, incubated for 30 min at 37 °C and freshly prepared Iodoacetamide was added to a final concentration of 15 mM and incubated for 30 min at room temperature in the dark. The protein was sequentially digested with Endoproteinase Lys-C and Trypsin. The enzymes were added at an enzyme to substrate ratio of 1:100 for LysC and 1:50 for Trypsin and the reaction incubated for 3-4 h at 37°C. The reaction was started by LysC followed by Trypsin addition after 4h. After digestion, the sample was acidified with Trifluoroacetic acid (1% v/v final concentration) at 37°C for 30 min, spun down at 17,000 g for 5 min and the peptides were fractionated by peptide size-exclusion chromatography. The mass spectrometry experiment and cross-links identification were performed by the EMBL Proteomics core facility.

Immunofluorescence and drug treatment. Cos7 cells were seeded in glass coverslips (Lab-Tek), and GFP and RFP bearing constructs were transfected by Lipofectamine 2000. Twenty-four hours later, cells were washed with PBS and fixed with 4% paraformaldehyde for 15 min, followed by

permeabilization with 0.2% Triton-X100 and 1 × PBS for 10 min at room temperature. After being blocked with 5% dry milk, 0.2% Tween 20, and 1 × PBS at RT for 30 min, slides were incubated with anti-acH4 antibody (1:100 dilution, Millipore) or anti-HDAC1 antibody (1:100 dilution, Cell Signaling Technology) in 1% dry milk, 0.1% Tween 20, and 1 × PBS in a humidified chamber at 4℃ overnight and then washed three times for 5 min each in the antibody dilution buffer. The secondary antibodies were diluted at 1:500 in the same buffer and incubated in a humidified chamber for 30 min at 37℃, and then washed as for the primary antibodies. The DNA was counterstained by DAPI, and the slides were mounted in Dako fluorescent mounting medium. For the drug treatment assays, twenty-four hours after being transfected with GFP and RFP bearing constructs, Cos7 cells were treated with DMSO (Sigma), 1,6-hexanediol (10%, 1 hour, Sigma), A485 (3 μM, 1 hour, Selleck), JQ1 (1 μM, 1 hour, Selleck), TSA (0.3 μM, 12 hours, Selleck) or CBP30 (1 μM, 1 hour, Selleck), then the cells were washed with PBS and fixed with 4% paraformaldehyde for 15 min.

Co-immunoprecipitation. Cell pellets were lysed in 300 mM NaCl buffer (20 mM HEPES-KOH pH 7.9, 300 mM NaCl, 3 mM $MgCl_2$, 0.1% NP-40, 1X protease inhibitor cocktail (Roche), 1X phosphatase inhibitor cocktail-1 (Sigma) and 10 mM sodium butyrate) for 30 min at 4 ℃ with rotation. After a 15 min centrifugation at 12,000g at 4℃, the supernatants were collected and separated into input and immunoprecipitate (IP) lysates. IP lysates were mixed with GFP-selector (pre-washed with 300 mM NaCl buffer three times, NanoTag) and rotated overnight at 4 ℃. After washing with 300 mM NaCl buffer, IPs were released by adding Laemmli sample buffer to beads and boiling for 5 min at 100℃.

Fluorescence recovery after photobleaching (FRAP). Live imaging and fluorescence recovery after photobleaching experiments (FRAP) were performed with the LSM710 Confocor3 confocal microscope (Carl Zeiss), equipped with the Plan-Apochromat C 63X /1.4 oil-immersion objective. FRAP area, time and time interval (1 image every 783ms) were maintained identical between the different BRD4-NUT mutants and repeat measurements (N=3-4, 45 to 65 foci measured per conditions) in order to photobleach 3-6 foci at the same time. To compare recovery kinetics against BRD4-NUT wild-type, FRAP measurements were fitted to a single exponential curve, $I_{(t)} = I_{(0)} + k_1 e^{-k2t}$ (performed with ZEN software; Carl Zeiss Microscopy) to determine the characteristic time of recovery. Values of τ (1/ k2) were compiled and statistical significance was tested using the Student's t-test (Excel).

NMR experiments. Backbone resonances of both proteins free in solution were assigned by measuring HNCO, HNCA, and HN(CO)CA spectra on ^{15}N, ^{13}C-labelled samples at 600 (NUT7) and 700 (TAZ2) MHz. We were able to assign 87% (95%) of the NH pairs, 72% (82%) of the CO resonances and 88% (96%) of the CA resonances in the backbone of NUT7 (TAZ2).

Side-chain resonances of both proteins free in solution were assigned by measuring HBHA(CO)NH, H(CCO)NH, and CC(CO)NH spectra on ^{15}N, ^{13}C-labelled samples at 600 (NUT7) and 700 (TAZ2) MHz. We were able to assign 70% (55%) of the HA resonances, 57% (45%) of the HB resonances, and 45% (53%) of the CB resonances in NUT7 (TAZ2), as well as part of the γ-protons.

To determine the binding site of NUT7 on TAZ2, we measured ^{1}H-^{15}N HSQC spectra on ^{15}N-labelled TAZ2 before and after adding unlabelled NUT7. For each of the assigned peaks in the spectra we calculate the chemical shift perturbations (CSP) defined as

$$CSP = \sqrt{\frac{1}{2}\left(\Delta\delta_H^2 + (0.14 \cdot \Delta\delta_N)^2\right)},$$

in which $\Delta\delta_H$ and $\Delta\delta_N$ are the chemical shift differences upon adding NUT7 in the ^{1}H and ^{15}N dimensions, respectively. We used the procedure described by Schumann et al. (Schumann et al., 2007) to determine whether measured CSPs are significant or not. We used the same approach to map the binding site on NUT7 monitoring a titration of a solution of ^{15}N-labelled NUT7 with unlabelled TAZ2 using ^{1}H-^{15}N HSQC spectra.

Both ^{13}C- and ^{15}N-HSQC-NOESY spectra on a sample of ^{15}N,^{13}C-labelled TAZ2 to which one equivalent of unlabelled NUT7 was added and ^{15}N-HSQC-NOESY spectra on ^{15}N-labelled NUT7 in admixture with one equivalent of unlabelled TAZ2 were measured using standard pulse sequences. All NOESY spectra were measured at 950 MHz with a mixing time of 100 ms.

HADDOCK docking. We used the HADDOCK protocol (Dominguez et al., 2003) with experimental information from titrations and NOESY spectra to derive a structural model of the interaction. We used the NMRbox platform (Maciejewski et al., 2017) to carry out the calculations. In a first step, we used the results of the titration experiments to derive ambiguous distance restraints (AIRs) describing the binding interface, and we used these AIRs as input for HADDOCK to dock extended conformations of NUT7 on the TAZ2 structure derived from PDB 2mzd. The best 200 structures resulting from the calculations were clustered based on the fraction of common contacts (FCC). The resulting clusters were sorted according to their agreement with the AIRs. Structures in the best cluster display the interaction of residues P448-E454 in NUT7 with TAZ2. Thus, we used the Flexible-Meccano algorithm (Ozenne et al., 2012) to generate an ensemble of 100 conformers of NUT7, in which the conformation of residues P448-E454 is the same as in the structure resulted from the HADDOCK calculation, whereas the rest of the chain behaves like a statistical coil. Then, we used HADDOCK again to dock this ensemble of structures on the same TAZ2 conformation as above. In this run, in addition to the same AIRs as in the preliminary run, we used distance restraints derived from NOESY spectra between S1726 in TAZ2 and residues L463 and A464 in NUT7. We also used TALOS-N (Shen and Bax, 2013) to derive restraints on the backbone dihedral angles of residues P448-E454 from chemical shift values. We repeated this calculation ten times with different initial seeds and velocities. Finally, the best 200 structures from each of the ten runs were clustered based on the FCC. The resulting clusters were sorted according to their agreement with the AIRs and NOE-based restraints. Then, from the best-ranked cluster in each of the runs, we selected the structure with the best HADDOCK score (which is a linear combination of various energy terms and buried surface area), thereby obtaining a final set of 10 structures.

Cryo-electron microscopy grid preparation, data collection and image processing
Quantifoil R1.2/1.3 Au 300 grids were glow discharged for 5 min at 40 mA on a Quorum Gloqube glow-discharge unit prior to application of the sample. 3 µl of 1 µM final concentration of the mononucleosome particles in buffer 20 mM Tris-HCl, pH8.0, 250 mM NaCl, 1 mM EDTA, 1 mM DTT were applied to the grid, blotted for 3 s at blot force 10 and plunge frozen into liquid ethane using a Vitrobot Mark IV (FEI Thermo Fisher), which was set at 4°C and 100% humidity. Cryo-EM data was collected on a Thermo Fisher Scientific Titan Krios G3 transmission electron microscope at the LISCB at the University of Leicester. Electron micrographs were recorded using a K3 direct electron detector (Gatan Inc.) at a dose rate of 15 e$^-$/pix/sec and a calibrated pixel size 1.086 Å. Focusing was performed over a range between -2.3 and -1.1 µm in 0.3 µm intervals. Image processing of the NCPs was performed as follows: movie stacks were corrected for beam-induced motion and then integrated using MotionCor2 (Zheng et al., 2017). All frames were retained and a patch alignment of 4 x 4 was used. Contrast transfer function (CTF) parameters for each micrograph were estimated by Gctf (Zhang, 2016). Integrated movies were inspected with Relion-3.1 for further image processing (243 movies). Initial particle sets were picked manually, extracted, and subjected to 2D classification in Relion 3.1 (Zivanov et al., 2018). Resulting classes corresponding to protein complexes were subsequently employed as references for auto-picking in Relion 3.1. Particle extraction was carried out from micrographs using a box size of 128 pixels (pixel size: 1.08 Å/pixel). An initial dataset of 80764 particles was cleaned by several iterations of 2D classification, classes exhibiting clear secondary structure features were selected for 3D classification. After iterative rounds of 3D classifications, well defined classes were selected and global refinement was performed.

References

Ahmad, K., and Henikoff, S. (2001). Modulation of a transcription factor counteracts heterochromatic gene silencing in Drosophila. Cell *104*, 839-847.

Ahn, J.H., Davis, E.S., Daugird, T.A., Zhao, S., Quiroga, I.Y., Uryu, H., Li, J., Storey, A.J., Tsai, Y.H., Keeley, D.P.*, et al.* (2021). Phase separation drives aberrant chromatin looping and cancer development. Nature *595*, 591-595.

Alekseyenko, A.A., Walsh, E.M., Wang, X., Grayson, A.R., Hsi, P.T., Kharchenko, P.V., Kuroda, M.I., and French, C.A. (2015). The oncogenic BRD4-NUT chromatin regulator drives aberrant transcription within large topological domains. Genes Dev *29*, 1507-1523.

Alekseyenko, A.A., Walsh, E.M., Zee, B.M., Pakozdi, T., Hsi, P., Lemieux, M.E., Dal Cin, P., Ince, T.A., Kharchenko, P.V., Kuroda, M.I.*, et al.* (2017). Ectopic protein interactions within BRD4-chromatin complexes drive oncogenic megadomain formation in NUT midline carcinoma. Proc Natl Acad Sci U S A *114*, E4184-E4192.

Allen, B.L., and Taatjes, D.J. (2015). The Mediator complex: a central integrator of transcription. Nat Rev Mol Cell Biol *16*, 155-166.

Aparicio, O.M., and Gottschling, D.E. (1994). Overcoming telomeric silencing: a trans-activator competes to establish gene expression in a cell cycle-dependent way. Genes Dev *8*, 1133-1146.

Apostolou, E., and Thanos, D. (2008). Virus Infection Induces NF-kappaB-dependent interchromosomal associations mediating monoallelic IFN-beta gene expression. Cell *134*, 85-96.

Attar, N., and Kurdistani, S.K. (2017). Exploitation of EP300 and CREBBP Lysine Acetyltransferases by Cancer. Cold Spring Harb Perspect Med *7*.

Avantaggiati, M.L., Carbone, M., Graessmann, A., Nakatani, Y., Howard, B., and Levine, A.S. (1996). The SV40 large T antigen and adenovirus E1a oncoproteins interact with distinct isoforms of the transcriptional co-activator, p300. EMBO Journal *15*, 2236-2248.

Basu, S., Mackowiak, S.D., Niskanen, H., Knezevic, D., Asimi, V., Grosswendt, S., Geertsema, H., Ali, S., Jerkovic, I., Ewers, H.*, et al.* (2020). Unblending of Transcriptional Condensates in Human Repeat Expansion Disease. Cell *181*, 1062-1079 e1030.

Black, J.C., Mosley, A., Kitada, T., Washburn, M., and Carey, M. (2008). The SIRT2 deacetylase regulates autoacetylation of p300. Mol Cell *32*, 449-455.

Boija, A., Klein, I.A., Sabari, B.R., Dall'Agnese, A., Coffey, E.L., Zamudio, A.V., Li, C.H., Shrinivas, K., Manteiga, J.C., Hannett, N.M.*, et al.* (2018). Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. Cell *175*, 1842-1855 e1816.

Boija, A., Klein, I.A., and Young, R.A. (2021). Biomolecular Condensates and Cancer. Cancer Cell *39*, 174-192.

Cabianca, D.S., Munoz-Jimenez, C., Kalck, V., Gaidatzis, D., Padeken, J., Seeber, A., Askjaer, P., and Gasser, S.M. (2019). Active chromatin marks drive spatial sequestration of heterochromatin in C. elegans nuclei. Nature *569*, 734-739.

Carey, M., Lin, Y.S., Green, M.R., and Ptashne, M. (1990). A mechanism for synergistic activation of a mammalian gene by GAL4 derivatives. Nature *345*, 361-364.

Carpenter, A.E., Memedula, S., Plutz, M.J., and Belmont, A.S. (2005). Common effects of acidic activators on large-scale chromatin structure and transcription. Mol Cell Biol *25*, 958-968.

Chambeyron, S., and Bickmore, W.A. (2004). Chromatin decondensation and nuclear reorganization of the HoxB locus upon induction of transcription. Gene Dev *18*, 1119-1130.

Chan, H.M., and La Thangue, N.B. (2001). p300/CBP proteins: HATs for transcriptional bridges and scaffolds. J Cell Sci *114*, 2363-2373.

Cho, W.K., Spille, J.H., Hecht, M., Lee, C., Li, C., Grube, V., and Cisse, II (2018). Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. Science *361*, 412-415.

Chong, S., Dugast-Darzacq, C., Liu, Z., Dong, P., Dailey, G.M., Cattoglio, C., Heckert, A., Banala, S., Lavis, L., Darzacq, X.*, et al.* (2018). Imaging dynamic and selective low-complexity domain interactions that control gene transcription. Science *361*.

Creyghton, M.P., Cheng, A.W., Welstead, G.G., Kooistra, T., Carey, B.W., Steine, E.J., Hanna, J., Lodato, M.A., Frampton, G.M., Sharp, P.A.*, et al.* (2010). Histone H3K27ac separates

active from poised enhancers and predicts developmental state. Proc Natl Acad Sci U S A *107*, 21931-21936.

Dames, S.A., Martinez-Yamout, M., De Guzman, R.N., Dyson, H.J., and Wright, P.E. (2002). Structural basis for Hif-1 alpha /CBP recognition in the cellular hypoxic response. Proc Natl Acad Sci U S A *99*, 5271-5276.

Dancy, B.M., and Cole, P.A. (2015). Protein lysine acetylation by p300/CBP. Chem Rev *115*, 2419-2452.

De Bosscher, K., Vanden Berghe, W., and Haegeman, G. (2003). The interplay between the glucocorticoid receptor and nuclear factor-kappaB or activator protein-1: molecular mechanisms for gene repression. Endocr Rev *24*, 488-522.

Delvecchio, M., Gaucher, J., Aguilar-Guerreri, C., Ortega, E., and Panne, D. (2013). Structure of the p300 catalytic core and implications for chromatin targeting and HAT regulation. Nature Structural and Molecular Biology *21*.

Devaiah, B.N., Case-Borden, C., Gegonne, A., Hsu, C.H., Chen, Q., Meerzaman, D., Dey, A., Ozato, K., and Singer, D.S. (2016). BRD4 is a histone acetyltransferase that evicts nucleosomes from chromatin. Nat Struct Mol Biol *23*, 540-548.

Dhalluin, C., Carlson, J.E., Zeng, L., He, C., Aggarwal, A.K., and Zhou, M.M. (1999). Structure and ligand of a histone acetyltransferase bromodomain. Nature *399*, 491-496.

Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012). Topological domains in mammalian genomes identified by analysis of chromatin interactions. Nature *485*, 376-380.

Dominguez, C., Boelens, R., and Bonvin, A.M. (2003). HADDOCK: a protein-protein docking approach based on biochemical or biophysical information. Journal of the American Chemical Society *125*, 1731-1737.

Dorigo, B., Schalch, T., Bystricky, K., and Richmond, T.J. (2003). Chromatin fiber folding: requirement for the histone H4 N-terminal tail. J Mol Biol *327*, 85-96.

Dyson, H.J., and Wright, P.E. (2016). Role of Intrinsic Protein Disorder in the Function and Interactions of the Transcriptional Coactivators CREB-binding Protein (CBP) and p300. J Biol Chem *291*, 6714-6722.

Eagen, K.P., and French, C.A. (2021). Supercharging BRD4 with NUT in carcinoma. Oncogene *40*, 1396-1408.

Eberharter, A., and Becker, P.B. (2002). Histone acetylation: a switch between repressive and permissive chromatin. Second in review series on chromatin dynamics. EMBO Rep *3*, 224-229.

Erdel, F., Rademacher, A., Vlijm, R., Tunnermann, J., Frank, L., Weinmann, R., Schweigert, E., Yserentant, K., Hummert, J., Bauer, C.*, et al.* (2020). Mouse Heterochromatin Adopts Digital Compaction States without Showing Hallmarks of HP1-Driven Liquid-Liquid Phase Separation. Mol Cell *78*, 236-249 e237.

Erdel, F., and Rippe, K. (2018). Formation of Chromatin Subcompartments by Phase Separation. Biophys J *114*, 2262-2270.

Falk, M., Feodorova, Y., Naumova, N., Imakaev, M., Lajoie, B.R., Leonhardt, H., Joffe, B., Dekker, J., Fudenberg, G., Solovei, I.*, et al.* (2019). Heterochromatin drives compartmentalization of inverted and conventional nuclei. Nature *570*, 395-399.

Feng, H., Jenkins, L.M., Durell, S.R., Hayashi, R., Mazur, S.J., Cherry, S., Tropea, J.E., Miller, M., Wlodawer, A., Appella, E.*, et al.* (2009). Structural basis for p300 Taz2-p53 TAD1 binding and modulation by phosphorylation. Structure *17*, 202-210.

Ferrari, R., Gou, D., Jawdekar, G., Johnson, S.A., Nava, M., Su, T., Yousef, A.F., Zemke, N.R., Pellegrini, M., Kurdistani, S.K.*, et al.* (2014). Adenovirus small E1A employs the lysine acetylases p300/CBP and tumor suppressor Rb to repress select host genes and promote productive virus infection. Cell Host Microbe *16*, 663-676.

Ferrari, R., Pellegrini, M., Horwitz, G.A., Xie, W., Berk, A.J., and Kurdistani, S.K. (2008). Epigenetic reprogramming by adenovirus e1a. Science *321*, 1086-1088.

Ferreon, J.C., Martinez-Yamout, M.A., Dyson, H.J., and Wright, P.E. (2009). Structural basis for subversion of cellular control mechanisms by the adenoviral E1A oncoprotein. Proceedings of the National Academy of Sciences of the United States of America *106*, 13260-13265.

Filippakopoulos, P., and Knapp, S. (2012). The bromodomain interaction module. FEBS Lett *586*, 2692-2704.

Fisher, W.W., Li, J.J., Hammonds, A.S., Brown, J.B., Pfeiffer, B.D., Weiszmann, R., MacArthur, S., Thomas, S., Stamatoyannopoulos, J.A., Eisen, M.B.*, et al.* (2012). DNA regions bound at low occupancy by transcription factors do not drive patterned reporter gene expression in Drosophila. Proc Natl Acad Sci U S A *109*, 21330-21335.

Franke, D., Petoukhov, M.V., Konarev, P.V., Panjkovich, A., Tuukkanen, A., Mertens, H.D.T., Kikhney, A.G., Hajizadeh, N.R., Franklin, J.M., Jeffries, C.M.*, et al.* (2017). ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering from macromolecular solutions. J Appl Crystallogr *50*, 1212-1225.

Fukaya, T., Lim, B., and Levine, M. (2016). Enhancer Control of Transcriptional Bursting. Cell *166*, 358-368.

Gaucher, J., Boussouar, F., Montellier, E., Curtet, S., Buchou, T., Bertrand, S., Hery, P., Jounier, S., Depaux, A., Vitte, A.L.*, et al.* (2012). Bromodomain-dependent stage-specific male genome programming by Brdt. EMBO J *31*, 3809-3820.

Ghandi, M., Huang, F.W., Jane-Valbuena, J., Kryukov, G.V., Lo, C.C., McDonald, E.R., 3rd, Barretina, J., Gelfand, E.T., Bielski, C.M., Li, H.*, et al.* (2019). Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature *569*, 503-508.

Gibson, B.A., Doolittle, L.K., Schneider, M.W.G., Jensen, L.E., Gamarra, N., Henry, L., Gerlich, D.W., Redding, S., and Rosen, M.K. (2019). Organization of Chromatin by Intrinsic and Regulated Phase Separation. Cell *179*, 470-484 e421.

Gill, G., and Ptashne, M. (1988). Negative effect of the transcriptional activator GAL4. Nature *334*, 721-724.

Gillespie, M.A., Palii, C.G., Sanchez-Taltavull, D., Shannon, P., Longabaugh, W.J.R., Downes, D.J., Sivaraman, K., Espinoza, H.M., Hughes, J.R., Price, N.D.*, et al.* (2020). Absolute Quantification of Transcription Factors Reveals Principles of Gene Regulation in Erythropoiesis. Mol Cell *78*, 960-974 e911.

Goodman, R.H., and Smolik, S. (2000). CBP/p300 in cell growth, transformation, and development. Genes Dev *14*, 1553-1577.

Goudarzi, A., Zhang, D., Huang, H., Barral, S., Kwon, O.K., Qi, S., Tang, Z., Buchou, T., Vitte, A.L., He, T.*, et al.* (2016). Dynamic Competing Histone H4 K5K8 Acetylation and Butyrylation Are Hallmarks of Highly Active Gene Promoters. Mol Cell *62*, 169-180.

Green, M.R. (2005). Eukaryotic transcription activation: right on target. Mol Cell *18*, 399-402.

Gryder, B.E., Pomella, S., Sayers, C., Wu, X.S., Song, Y., Chiarella, A.M., Bagchi, S., Chou, H.C., Sinniah, R.S., Walton, A.*, et al.* (2019a). Histone hyperacetylation disrupts core gene regulatory architecture in rhabdomyosarcoma. Nat Genet *51*, 1714-1722.

Gryder, B.E., Wu, L., Woldemichael, G.M., Pomella, S., Quinn, T.R., Park, P.M.C., Cleveland, A., Stanton, B.Z., Song, Y., Rota, R.*, et al.* (2019b). Chemical genomics reveals histone deacetylases are required for core regulatory transcription. Nat Commun *10*, 3004.

Guertin, M.J., Zhang, X., Coonrod, S.A., and Hager, G.L. (2014). Transient estrogen receptor binding and p300 redistribution support a squelching mechanism for estradiol-repressed genes. Mol Endocrinol *28*, 1522-1533.

Guo, Y.E., Manteiga, J.C., Henninger, J.E., Sabari, B.R., Dall'Agnese, A., Hannett, N.M., Spille, J.H., Afeyan, L.K., Zamudio, A.V., Shrinivas, K.*, et al.* (2019). Pol II phosphorylation regulates a switch between transcriptional and splicing condensates. Nature *572*, 543-+.

Gurdon, J.B., Javed, K., Vodnala, M., and Garrett, N. (2020). Long-term association of a transcription factor with its chromatin binding site can stabilize gene expression and cell fate commitment. Proc Natl Acad Sci U S A *117*, 15075-15084.

Halford, S.E. (2009). An end to 40 years of mistakes in DNA-protein association kinetics? Biochem Soc Trans *37*, 343-348.

Hammitzsch, A., Tallant, C., Fedorov, O., O'Mahony, A., Brennan, P.E., Hay, D.A., Martinez, F.O., Al-Mossawi, M.H., de Wit, J., Vecellio, M.*, et al.* (2015). CBP30, a selective CBP/p300 bromodomain inhibitor, suppresses human Th17 responses. Proc Natl Acad Sci U S A *112*, 10768-10773.

Han, X., Yu, D., Gu, R., Jia, Y., Wang, Q., Jaganathan, A., Yang, X., Yu, M., Babault, N., Zhao, C., *et al.* (2020). Roles of the BRD4 short isoform in phase separation and active gene transcription. Nat Struct Mol Biol *27*, 333-341.

Hansson, M.L., Popko-Scibor, A.E., Saint Just Ribeiro, M., Dancy, B.M., Lindberg, M.J., Cole, P.A., and Wallberg, A.E. (2009). The transcriptional coactivator MAML1 regulates p300 autoacetylation and HAT activity. Nucleic Acids Res *37*, 2996-3006.

He, J., Ye, J., Cai, Y., Riquelme, C., Liu, J.O., Liu, X., Han, A., and Chen, L. (2011). Structure of p300 bound to MEF2 on DNA reveals a mechanism of enhanceosome assembly. Nucleic Acids Res *39*, 4464-4474.

Hnisz, D., Shrinivas, K., Young, R.A., Chakraborty, A.K., and Sharp, P.A. (2017). A Phase Separation Model for Transcriptional Control. Cell *169*, 13-23.

Ho, J.W., Jung, Y.L., Liu, T., Alver, B.H., Lee, S., Ikegami, K., Sohn, K.A., Minoda, A., Tolstorukov, M.Y., Appert, A., *et al.* (2014). Comparative analysis of metazoan chromatin organization. Nature *512*, 449-452.

Hogg, S.J., Motorna, O., Cluse, L.A., Johanson, T.M., Coughlan, H.D., Raviram, R., Myers, R.M., Costacurta, M., Todorovski, I., Pijpers, L., *et al.* (2021). Targeting histone acetylation dynamics and oncogenic transcription by catalytic P300/CBP inhibition. Mol Cell *81*, 2183-2200 e2113.

Horwitz, G.A., Zhang, K., McBrian, M.A., Grunstein, M., Kurdistani, S.K., and Berk, A.J. (2008). Adenovirus small e1a alters global patterns of histone modification. Science *321*, 1084-1085.

Hottiger, M.O., Felzien, L.K., and Nabel, G.J. (1998). Modulation of cytokine-induced HIV gene expression by competitive binding of transcription factors to the coactivator p300. EMBO J *17*, 3124-3134.

Hsu, E., Pennella, M.A., Zemke, N.R., Eng, C., and Berk, A.J. (2018). Adenovirus E1A Activation Domain Regulates H3 Acetylation Affecting Varied Steps in Transcription at Different Viral Promoters. J Virol *92*.

Jeppesen, P., and Turner, B.M. (1993). The inactive X chromosome in female mammals is distinguished by a lack of histone H4 acetylation, a cytogenetic marker for gene expression. Cell *74*, 281-289.

Kamei, Y., Xu, L., Heinzel, T., Torchia, J., Kurokawa, R., Gloss, B., Lin, S.C., Heyman, R.A., Rose, D.W., Glass, C.K., *et al.* (1996). A CBP integrator complex mediates transcriptional activation and AP-1 inhibition by nuclear receptors. Cell *85*, 403-414.

Karanam, B., Jiang, L., Wang, L., Kelleher, N.L., and Cole, P.A. (2006). Kinetic and mass spectrometric analysis of p300 histone acetyltransferase domain autoacetylation. J Biol Chem *281*, 40292-40301.

Kidder, B.L., and Palmer, S. (2012). HDAC1 regulates pluripotency and lineage specific transcriptional networks in embryonic and trophoblast stem cells. Nucleic Acids Res *40*, 2925-2939.

Kraus, W.L., Manning, E.T., and Kadonaga, J.T. (1999). Biochemical analysis of distinct activation functions in p300 that enhance transcription initiation with chromatin templates. Molecular and Cellular Biology *19*, 8123-8135.

Kribelbauer, J.F., Rastogi, C., Bussemaker, H.J., and Mann, R.S. (2019). Low-Affinity Binding Sites and the Transcription Factor Specificity Paradox in Eukaryotes. Annu Rev Cell Dev Biol *35*, 357-379.

Krois, A.S., Ferreon, J.C., Martinez-Yamout, M.A., Dyson, H.J., and Wright, P.E. (2016). Recognition of the disordered p53 transactivation domain by the transcriptional adapter zinc finger domains of CREB-binding protein. Proc Natl Acad Sci U S A *113*, E1853-1862.

Kumar, M., Molkentine, D., Molkentine, J., Bridges, K., Xie, T., Yang, L., Hefner, A., Gao, M., Bahri, R., Dhawan, A., *et al.* (2021). Inhibition of histone acetyltransferase function radiosensitizes CREBBP/EP300 mutants via repression of homologous recombination, potentially targeting a gain of function. Nat Commun *12*, 6340.

Ladurner, A.G., Inouye, C., Jain, R., and Tjian, R. (2003). Bromodomains mediate an acetyl-histone encoded antisilencing function at heterochromatin boundaries. Molecular Cell *11*, 365-376.

Larson, A.G., Elnatan, D., Keenen, M.M., Trnka, M.J., Johnston, J.B., Burlingame, A.L., Agard, D.A., Redding, S., and Narlikar, G.J. (2017). Liquid droplet formation by HP1alpha suggests a role for phase separation in heterochromatin. Nature *547*, 236-240.

Lee, C.W., Martinez-Yamout, M.A., Dyson, H.J., and Wright, P.E. (2010). Structure of the p53 transactivation domain in complex with the nuclear receptor coactivator binding domain of CREB binding protein. Biochemistry *49*, 9964-9971.

Li, J., Dong, A., Saydaminova, K., Chang, H., Wang, G., Ochiai, H., Yamamoto, T., and Pertsinidis, A. (2019). Single-Molecule Nanoscopy Elucidates RNA Polymerase II Transcription at Single Genes in Live Cells. Cell *178*, 491-506 e428.

Li, J., Hsu, A., Hua, Y., Wang, G., Cheng, L., Ochiai, H., Yamamoto, T., and Pertsinidis, A. (2020). Single-gene imaging links genome topology, promoter-enhancer communication and transcription control. Nat Struct Mol Biol *27*, 1032-1040.

Li, X.Y., MacArthur, S., Bourgon, R., Nix, D., Pollard, D.A., Iyer, V.N., Hechmer, A., Simirenko, L., Stapleton, M., Luengo Hendriks, C.L.*, et al.* (2008). Transcription factors bind thousands of active and inactive regions in the Drosophila blastoderm. PLoS Biol *6*, e27.

Lickwar, C.R., Mueller, F., Hanlon, S.E., McNally, J.G., and Lieb, J.D. (2012). Genome-wide protein-DNA binding dynamics suggest a molecular clutch for transcription factor function. Nature *484*, 251-255.

Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O.*, et al.* (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science *326*, 289-293.

Loffreda, A., Jacchetti, E., Antunes, S., Rainone, P., Daniele, T., Morisaki, T., Bianchi, M.E., Tacchetti, C., and Mazza, D. (2017). Live-cell p53 single-molecule binding is modulated by C-terminal acetylation and correlates with transcriptional activity. Nat Commun *8*, 313.

Luger, K., Rechsteiner, T.J., and Richmond, T.J. (1999). Preparation of nucleosome core particle from recombinant histones. Methods Enzymol *304*, 3-19.

Luo, J., Li, M., Tang, Y., Laszkowska, M., Roeder, R.G., and Gu, W. (2004). Acetylation of p53 augments its site-specific DNA binding both *in vitro* and *in vivo*. Proc Natl Acad Sci U S A *101*, 2259-2264.

Ma, L., Gao, Z., Wu, J., Zhong, B., Xie, Y., Huang, W., and Lin, Y. (2021). Co-condensation between transcription factor and coactivator p300 modulates transcriptional bursting kinetics. Mol Cell.

Maciejewski, M.W., Schuyler, A.D., Gryk, M.R., Moraru, II, Romero, P.R., Ulrich, E.L., Eghbalnia, H.R., Livny, M., Delaglio, F., and Hoch, J.C. (2017). NMRbox: A Resource for Biomolecular NMR Computation. Biophys J *112*, 1529-1534.

Malik, S., and Roeder, R.G. (2010). The metazoan Mediator co-activator complex as an integrative hub for transcriptional regulation. Nat Rev Genet *11*, 761-772.

Marceau, A.H., Brison, C.M., Nerli, S., Arsenault, H.E., McShan, A.C., Chen, E., Lee, H.W., Benanti, J.A., Sgourakis, N.G., and Rubin, S.M. (2019). An order-to-disorder structural switch activates the FoxM1 transcription factor. Elife *8*.

Merika, M., Williams, A.J., Chen, G., Collins, T., and Thanos, D. (1998). Recruitment of CBP/p300 by the IFN beta enhanceosome is required for synergistic activation of transcription. Mol Cell *1*, 277-287.

Meyer, M.E., Gronemeyer, H., Turcotte, B., Bocquel, M.T., Tasset, D., and Chambon, P. (1989). Steroid hormone receptors compete for factors that mediate their enhancer function. Cell *57*, 433-442.

Migl, D., Kschonsak, M., Arthur, C.P., Khin, Y., Harrison, S.C., Ciferri, C., and Dimitrova, Y.N. (2020). Cryoelectron Microscopy Structure of a Yeast Centromeric Nucleosome at 2.7 A Resolution. Structure *28*, 363-370 e363.

Miller Jenkins, L.M., Feng, H., Durell, S.R., Tagad, H.D., Mazur, S.J., Tropea, J.E., Bai, Y., and Appella, E. (2015). Characterization of the p300 Taz2-p53 TAD2 complex and comparison with the p300 Taz2-p53 TAD1 complex. Biochemistry *54*, 2001-2010.

Miller, T.C., Simon, B., Rybin, V., Grotsch, H., Curtet, S., Khochbin, S., Carlomagno, T., and Muller, C.W. (2016). A bromodomain-DNA interaction facilitates acetylation-dependent bivalent nucleosome recognition by the BET protein BRDT. Nat Commun *7*, 13855.

Mirny, L., Slutsky, M., Wunderlich, Z., Tafvizi, A., Leith, J., and Kosmrlj, A. (2009). How a protein searches for its site on DNA: the mechanism of facilitated diffusion. Journal of Physics A: Mathematical and Theoretical *42*, 434013.

Misteli, T. (2020). The Self-Organizing Genome: Principles of Genome Architecture and Function. Cell *183*, 28-45.

Monahan, K., Horta, A., and Lomvardas, S. (2019). LHX2- and LDB1-mediated trans interactions regulate olfactory receptor choice. Nature *565*, 448-453.

Mueller, W.G., Walker, D., Hager, G.L., and McNally, J.G. (2001). Large-scale chromatin decondensation and recondensation regulated by transcription from a natural promoter. Mol Biol Cell *12*, 357a-357a.

Mukherjee, S.P., Behar, M., Birnbaum, H.A., Hoffmann, A., Wright, P.E., and Ghosh, G. (2013). Analysis of the RelA:CBP/p300 interaction reveals its involvement in NF-kappaB-driven transcription. PLoS Biol *11*, e1001647.

Narita, T., Ito, S., Higashijima, Y., Chu, W.K., Neumann, K., Walter, J., Satpathy, S., Liebner, T., Hamilton, W.B., Maskey, E.*, et al.* (2021). Enhancers are activated by p300/CBP activity-dependent PIC assembly, RNAPII recruitment, and pause release. Mol Cell *81*, 2166-2182 e2166.

Nora, E.P., Lajoie, B.R., Schulz, E.G., Giorgetti, L., Okamoto, I., Servant, N., Piolot, T., van Berkum, N.L., Meisig, J., Sedat, J.*, et al.* (2012). Spatial partitioning of the regulatory landscape of the X-inactivation centre. Nature *485*, 381-385.

Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N., and Mirny, L.A. (2018). Chromatin organization by an interplay of loop extrusion and compartmental segregation. Proc Natl Acad Sci U S A *115*, E6697-E6706.

Ortega, E., Rengachari, S., Ibrahim, Z., Hoghoughi, N., Gaucher, J., Holehouse, A.S., Khochbin, S., and Panne, D. (2018). Transcription factor dimerization activates the p300 acetyltransferase. Nature *562*, 538-544.

Ozenne, V., Bauer, F., Salmon, L., Huang, J.R., Jensen, M.R., Segard, S., Bernado, P., Charavay, C., and Blackledge, M. (2012). Flexible-meccano: a tool for the generation of explicit ensemble descriptions of intrinsically disordered proteins and their associated experimental observables. Bioinformatics *28*, 1463-1470.

Panne, D. (2008). The enhanceosome. Curr Opin Struct Biol *18*, 236-242.

Panne, D., Maniatis, T., and Harrison, S.C. (2007). An atomic model of the interferon-beta enhanceosome. Cell *129*, 1111-1123.

Pelka, P., Ablack, J.N., Torchia, J., Turnell, A.S., Grand, R.J., and Mymryk, J.S. (2009). Transcriptional control by adenovirus E1A conserved region 3 via p300/CBP. Nucleic Acids Res *37*, 1095-1106.

Petoukhov, M.V., Franke, D., Shkumatov, A.V., Tria, G., Kikhney, A.G., Gajda, M., Gorba, C., Mertens, H.D., Konarev, P.V., and Svergun, D.I. (2012). New developments in the program package for small-angle scattering data analysis. J Appl Crystallogr *45*, 342-350.

Pivot-Pajot, C., Caron, C., Govin, J., Vion, A., Rousseaux, S., and Khochbin, S. (2003). Acetylation-dependent chromatin reorganization by BRDT, a testis-specific bromodomain-containing protein. Mol Cell Biol *23*, 5354-5365.

Plevin, M.J., Mills, M.M., and Ikura, M. (2005). The LxxLL motif: a multifunctional binding sequence in transcriptional regulation. Trends Biochem Sci *30*, 66-69.

Portuguez, A.S., Grbesa, I., Tal, M., Deitch, R., Raz, D., Weismann, R., Schwartz, M., Loza, O., Sung, M.-H., Kaplan, T.*, et al.* (2019). Gene activation and repression by the glucocorticoid receptor are mediated by sequestering Ep300 and two modes of chromatin binding. bioRxiv.

Rada-Iglesias, A., Bajpai, R., Swigut, T., Brugmann, S.A., Flynn, R.A., and Wysocka, J. (2010). A unique chromatin signature uncovers early developmental enhancers in humans. Nature.

Reiter, F., Wienerroither, S., and Stark, A. (2017). Combinatorial function of transcription factors and cofactors. Curr Opin Genet Dev *43*, 73-81.

Reynoird, N., Schwartz, B.E., Delvecchio, M., Sadoul, K., Meyers, D., Mukherjee, C., Caron, C., Kimura, H., Rousseaux, S., Cole, P.A.*, et al.* (2010). Oncogenesis by sequestration of CBP/p300 in transcriptionally inactive hyperacetylated chromatin domains. EMBO J *29*, 2943-2952.

Rosencrance, C.D., Ammouri, H.N., Yu, Q., Ge, T., Rendleman, E.J., Marshall, S.A., and Eagen, K.P. (2020). Chromatin Hyperacetylation Impacts Chromosome Folding by Forming a Nuclear Subcompartment. Mol Cell *78*, 112-126 e112.

Rowley, M.J., and Corces, V.G. (2018). Organizational principles of 3D genome architecture. Nat Rev Genet *19*, 789-800.

Rundlett, S.E., Carmen, A.A., Kobayashi, R., Bavykin, S., Turner, B.M., and Grunstein, M. (1996). HDA1 and RPD3 are members of distinct yeast histone deacetylase complexes that regulate silencing and transcription. Proc Natl Acad Sci U S A *93*, 14503-14508.

Sabari, B.R., Dall'Agnese, A., Boija, A., Klein, I.A., Coffey, E.L., Shrinivas, K., Abraham, B.J., Hannett, N.M., Zamudio, A.V., Manteiga, J.C.*, et al.* (2018). Coactivator condensation at super-enhancers links phase separation and gene control. Science *361*.

Schumann, F.H., Riepl, H., Maurer, T., Gronwald, W., Neidig, K.P., and Kalbitzer, H.R. (2007). Combined chemical shift changes and amino acid specific chemical shift mapping of protein-protein interactions. J Biomol NMR *39*, 275-289.

Schwartz, B.E., Hofer, M.D., Lemieux, M.E., Bauer, D.E., Cameron, M.J., West, N.H., Agoston, E.S., Reynoird, N., Khochbin, S., Ince, T.A.*, et al.* (2011). Differentiation of NUT midline carcinoma by epigenomic reprogramming. Cancer Research *71*, 2686-2696.

Schwarzer, W., Abdennur, N., Goloborodko, A., Pekowska, A., Fudenberg, G., Loe-Mie, Y., Fonseca, N.A., Huber, W., C, H.H., Mirny, L.*, et al.* (2017). Two independent modes of chromatin organization revealed by cohesin removal. Nature *551*, 51-56.

Sheahan, T.W., Major, V., Webb, K.M., Bryan, E., and Voigt, P. (2020). The TAZ2 domain of CBP/p300 directs acetylation towards H3K27 within chromatin. bioRxiv.

Shen, Y., and Bax, A. (2013). Protein backbone and sidechain torsion angles predicted from NMR chemical shifts using artificial neural networks. J Biomol NMR *56*, 227-241.

Shin, Y., Chang, Y.C., Lee, D.S.W., Berry, J., Sanders, D.W., Ronceray, P., Wingreen, N.S., Haataja, M., and Brangwynne, C.P. (2018). Liquid Nuclear Condensates Mechanically Sense and Restructure the Genome. Cell *175*, 1481-1491 e1413.

Shiota, H., Barral, S., Buchou, T., Tan, M., Coute, Y., Charbonnier, G., Reynoird, N., Boussouar, F., Gerard, M., Zhu, M.*, et al.* (2018). Nut Directs p300-Dependent, Genome-Wide H4 Hyperacetylation in Male Germ Cells. Cell Rep *24*, 3477-3487 e3476.

Shogren-Knaak, M., Ishii, H., Sun, J.M., Pazin, M.J., Davie, J.R., and Peterson, C.L. (2006). Histone H4-K16 acetylation controls chromatin structure and protein interactions. Science *311*, 844-847.

Shrinivas, K., Sabari, B.R., Coffey, E.L., Klein, I.A., Boija, A., Zamudio, A.V., Schuijers, J., Hannett, N.M., Sharp, P.A., Young, R.A.*, et al.* (2019). Enhancer Features that Drive Formation of Transcriptional Condensates. Mol Cell *75*, 549-561 e547.

Slaughter, M.J., Shanle, E.K., Khan, A., Chua, K.F., Hong, T., Boxer, L.D., Allis, C.D., Josefowicz, S.Z., Garcia, B.A., Rothbart, S.B.*, et al.* (2021). HDAC inhibition results in widespread alteration of the histone acetylation landscape and BRD4 targeting to gene bodies. Cell Rep *34*, 108638.

Solovei, I., Thanisch, K., and Feodorova, Y. (2016). How to rule the nucleus: divide et impera. Curr Opin Cell Biol *40*, 47-59.

Spilianakis, C.G., Lalioti, M.D., Town, T., Lee, G.R., and Flavell, R.A. (2005). Interchromosomal associations between alternatively expressed loci. Nature *435*, 637-645.

Staller, M.V., Holehouse, A.S., Swain-Lenz, D., Das, R.K., Pappu, R.V., and Cohen, B.A. (2018). A High-Throughput Mutational Scan of an Intrinsically Disordered Acidic Transcriptional Activation Domain. Cell Syst *6*, 444-455 e446.

Strahl, B.D., and Allis, C.D. (2000). The language of covalent histone modifications. Nature *403*, 41-45.

Strickfaden, H., Tolsma, T.O., Sharma, A., Underhill, D.A., Hansen, J.C., and Hendzel, M.J. (2020). Condensed Chromatin Behaves like a Solid on the Mesoscale *In vitro* and in Living Cells. Cell *183*, 1772-1784 e1713.

Strom, A.R., Biggs, R.J., Banigan, E.J., Wang, X., Chiu, K., Herman, C., Collado, J., Yue, F., Ritland Politz, J.C., Tait, L.J.*, et al.* (2021). HP1alpha is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. Elife *10*.

Strom, A.R., Emelyanov, A.V., Mir, M., Fyodorov, D.V., Darzacq, X., and Karpen, G.H. (2017). Phase separation drives heterochromatin domain formation. Nature *547*, 241-245.

Taddei, A., Maison, C., Roche, D., and Almouzni, G. (2001). Reversible disruption of pericentric heterochromatin and centromere function by inhibiting deacetylases. Nat Cell Biol *3*, 114-120.

Therizols, P., Illingworth, R.S., Courilleau, C., Boyle, S., Wood, A.J., and Bickmore, W.A. (2014). Chromatin decondensation is sufficient to alter nuclear organization in embryonic stem cells. Science *346*, 1238-1242.

Thompson, P.R., Wang, D., Wang, L., Fulco, M., Pediconi, N., Zhang, D., An, W., Ge, Q., Roeder, R.G., Wong, J.*, et al.* (2004). Regulation of the p300 HAT domain via a novel activation loop. Nat Struct Mol Biol *11*, 308-315.

Toth, K.F., Knoch, T.A., Wachsmuth, M., Frank-Stohr, M., Stohr, M., Bacher, C.P., Muller, G., and Rippe, K. (2004). Trichostatin A-induced histone acetylation causes decondensation of interphase chromatin. J Cell Sci *117*, 4277-4287.

Triezenberg, S.J., Kingsbury, R.C., and McKnight, S.L. (1988). Functional dissection of VP16, the trans-activator of herpes simplex virus immediate early gene expression. Genes Dev *2*, 718-729.

Tumbar, T., Belmont, A.S., and Sudlow, G.P. (1999a). Targeting the VP16 acidic activation domain induces chances in large-scale chromatin structure and nuclear positioning. Mol Biol Cell *10*, 281a-281a.

Tumbar, T., Sudlow, G., and Belmont, A.S. (1999b). Large-scale chromatin unfolding and remodeling induced by VP16 acidic activation domain. J Cell Biol *145*, 1341-1354.

van de Werken, H.J.G., Haan, J.C., Feodorova, Y., Bijos, D., Weuts, A., Theunis, K., Holwerda, S.J.B., Meuleman, W., Pagie, L., Thanisch, K.*, et al.* (2017). Small chromosomal regions position themselves autonomously according to their chromatin class. Genome Res *27*, 922-933.

Vogelauer, M., Wu, J., Suka, N., and Grunstein, M. (2000). Global histone acetylation and deacetylation in yeast. Nature *408*, 495-498.

Wang, F., Marshall, C.B., Yamamoto, K., Li, G.Y., Gasmi-Seabrook, G.M., Okada, H., Mak, T.W., and Ikura, M. (2012a). Structures of KIX domain of CBP in complex with two FOXO3a transactivation domains reveal promiscuity and plasticity in coactivator recruitment. Proc Natl Acad Sci U S A *109*, 6078-6083.

Wang, L., Gao, Y., Zheng, X., Liu, C., Dong, S., Li, R., Zhang, G., Wei, Y., Qu, H., Li, Y.*, et al.* (2019). Histone Modifications Regulate Chromatin Compartmentalization by Contributing to a Phase Separation Mechanism. Mol Cell *76*, 646-659 e646.

Wang, R., Li, Q., Helfer, C.M., Jiao, J., and You, J. (2012b). Bromodomain protein Brd4 associated with acetylated chromatin is important for maintenance of higher-order chromatin structure. J Biol Chem *287*, 10738-10752.

Wang, R., and You, J. (2015). Mechanistic analysis of the role of bromodomain-containing protein 4 (BRD4) in BRD4-NUT oncoprotein-induced transcriptional activation. J Biol Chem *290*, 2744-2758.

Wang, Z., Zang, C., Cui, K., Schones, D.E., Barski, A., Peng, W., and Zhao, K. (2009). Genome-wide mapping of HATs and HDACs reveals distinct functions in active and inactive genes. Cell *138*, 1019-1031.

Weinert, B.T., Narita, T., Satpathy, S., Srinivasan, B., Hansen, B.K., Scholz, C., Hamilton, W.B., Zucconi, B.E., Wang, W.W., Liu, W.R.*, et al.* (2018). Time-Resolved Analysis Reveals Rapid Dynamics and Broad Scope of the CBP/p300 Acetylome. Cell *174*, 231-244 e212.

Whyte, W.A., Bilodeau, S., Orlando, D.A., Hoke, H.A., Frampton, G.M., Foster, C.T., Cowley, S.M., and Young, R.A. (2012). Enhancer decommissioning by LSD1 during embryonic stem cell differentiation. Nature *482*, 221-225.

Wojciak, J.M., Martinez-Yamout, M.A., Dyson, H.J., and Wright, P.E. (2009). Structural basis for recruitment of CBP/p300 coactivators by STAT1 and STAT2 transactivation domains. EMBO J *28*, 948-958.

Wu, S.Y., Lee, A.Y., Lai, H.T., Zhang, H., and Chiang, C.M. (2013). Phospho switch triggers Brd4 chromatin binding and activator recruitment for gene-specific targeting. Mol Cell *49*, 843-857.

Yan, J., Diaz, J., Jiao, J., Wang, R., and You, J. (2011). Perturbation of BRD4 protein function by BRD4-NUT protein abrogates cellular differentiation in NUT midline carcinoma. J Biol Chem *286*, 27663-27675.

Zamudio, A.V., Dall'Agnese, A., Henninger, J.E., Manteiga, J.C., Afeyan, L.K., Hannett, N.M., Coffey, E.L., Li, C.H., Oksuz, O., Sabari, B.R.*, et al.* (2019). Mediator Condensates Localize Signaling Factors to Key Cell Identity Genes. Mol Cell *76*, 753-766 e756.

Zhang, K. (2016). Gctf: Real-time CTF determination and correction. Journal of Structural Biology *193*, 1-12.

Zhang, T., Zhang, Z., Dong, Q., Xiong, J., and Zhu, B. (2020). Histone H3K27 acetylation is dispensable for enhancer activity in mouse embryonic stem cells. Genome Biol *21*, 45.

Zhang, Y., Brown, K., Yu, Y., Ibrahim, Z., Zandian, M., Xuan, H., Ingersoll, S., Lee, T., Ebmeier, C.C., Liu, J.*, et al.* (2021). Nuclear condensates of p300 formed though the structured catalytic core can act as a storage pool of p300 with reduced HAT activity. Nat Commun *12*, 4618.

Zhang, Y., Xue, Y., Shi, J., Ahn, J., Mi, W., Ali, M., Wang, X., Klein, B.J., Wen, H., Li, W.*, et al.* (2018). The ZZ domain of p300 mediates specificity of the adjacent HAT domain for histone H3. Nat Struct Mol Biol *25*, 841-849.

Zhao, R., Nakamura, T., Fu, Y., Lazar, Z., and Spector, D.L. (2011). Gene bookmarking accelerates the kinetics of post-mitotic transcriptional re-activation. Nature Cell Biology *13*, 1295-U1239.

Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017). MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. Nat Methods *14*, 331-332.

Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in RELION-3. Elife *7*.

Zylicz, J.J., Bousard, A., Zumer, K., Dossin, F., Mohammad, E., da Rocha, S.T., Schwalb, B., Syx, L., Dingli, F., Loew, D.*, et al.* (2019). The Implication of Early Chromatin Changes in X Chromosome Inactivation. Cell *176*, 182-197 e123.

Acknowledgements We thank Christos Savva and TJ Ragan for CryoEM support. We thank Nathan Cowieson from the B21 BioSAXS beamline of the Diamond Light Source; Yingming Zhao and Jinjun Gao for mass spectroscopy analysis; Luke Bailey, Sarah Northall and Dipti Vashi for help with histone preparations. Michael K. Rosen for plasmids encoding GFP-(Bomo)$_5$ and GFP-(Bromo)$_5$ N140A; Stephen M. Hinshaw and Stephen C. Harrison for plasmids encoding *S. cerevisiae* histones; Timothy J. Richmond for histone H4 mutant plasmids; Andy Turnell for E1A plasmids. Work in the S.K. laboratory was supported by ANR Episperm4 program, by Plan Cancer Pitcher, by MSD Avenir ERICAN programs, by the Cancer ITMO (Multi-Organisation Thematic Institute) of the French Alliance for Life Sciences and Health (AVIESAN) MIC program, by the "Université Grenoble Alpes" ANR-15-IDEX-02 SYMER and LIFE programs and by Fondation ARC program RF20190208471. Work in the D.P. laboratory was supported by grants from the Worldwide Cancer Research [16-0280], the Wellcome Trust [221881/Z/20/Z], Instruct-ERIC [PID: 18571] and the Medical Research Council [MR/W001667/1] to D.P. and the BBSRC (BB/S019510/1; BB/R016275/1).

Author contributions Z.I. designed and performed most experiments, analysed and validated the data with support by L.F. T.W., O.D. N.H and N.R performed cell biology experiments. N.S. performed NMR experiments. A.R performed chromatin analyses *in vitro*. Y. Z. and J. G. performed mass spectroscopy analysis. T.S., M.B., and S.K. provided supervision, funding acquisition and commented on the draft. D.P. was involved in conceptualization, supervision, project administration, funding acquisition. Z.I and D.P. wrote the original and revised drafts with input from all authors.

Competing interests: Y. Z. is a founder, board member, advisor to, and inventor on patents licensed to PTM Biolabs Inc (Hangzhou, China and Chicago, IL) and Maponos Therapeutics Inc. (Chicago, IL). The other authors declare no competing interests.

Table 1 – List of NUT constructs and binding experiments

Sample	Amino acid residues	Binding*
NUT F1c	347-588	+
NUT1	347-446	+/–
NUT2	427-527	+
NUT3	508-588	–
NUT4	347-480	+
NUT5	347-514	+
NUT6	351-470	+
NUT7	448-492	+
NUT7 mut1	448-492	+
NUT8	396-470	+
NUT8 mut1	396-470	–
NUT8 mut2	396-470	–
NUT8 mut3	396-470	+
NUT8 mut4	396-470	+
NUT9	395-482	+
NUT10	468-482	+

*GST-pulldown, co-sedimentation or BLI

```
NUT7 wt:    PLALIEELEQEEGLTLAQLVQKRLMALEEEEDAEAPPSFSGAQLD
NUT7 mut1:  PLALIEELEQEAGLTLAQAVQKRLMAAEEEEDAEAPPSFSGAQLD
NUT8 wt:    GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFLADLLSPEKQRDPLALIEELEQEEGLTLAQLVQKRLMALEEEEDAE
NUT8 Mut1:  GQQQEEEGMYPDPGAASAINEAASQKVAASKAEAAAHPQAAADAASPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
NUT8 Mut2:  GQQQEEEGMYPKPGALSKINELGSQKVAGSKVSAGIHGQSLADGSSPEKQRKPLGSIEKSEQKSGGSLAQLVQKRLMALEEEEDAE
NUT8 Mut3:  GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFLADLLSPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
NUT8 Mut4:  GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFAADAASPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
```

Table 2 - HADDOCK docking statistics

Distance constraints	
Ambiguous interaction restraints (AIRs)	14
Active residues on NUT7	5
Active residues on TAZ2	9
Intermolecular NOEs	2
Dihedral angle restraints*	
Phi/Psi	5/5
Structural statistics	
AIR Violations (mean ± s.d.)	2±1
Ramachandran analysis	
Residues in the favoured region (%)	87%
Residues in the additional allowed regions (%)	10%
Outliers	3%
Intermolecular energies after water refinement	
E_{vdw} (kcal mol^{-1})	-45 ± 7
E_{elec} (kcal mol^{-1})	-303 ± 32
Buried surface area (Å^2)	1498 ± 88

*Dihedral angle restraints derived from chemical shift values using TALOS-N are applied to NUT7 only. For the TAZ2 domain, the structure in PDB 2mzd is used as input to HADDOCK.

Table 3 - Cryo-EM data collection and map reconstruction

	Nucleosome Core Particle
Data collection	FEI Titan Krios G3
Detector	Gatan K3 direct electron detector
Voltage	300 kV
Pixel size (Å)	1.086
Nominal Magnification	81,000
Dose rate on Detector (e^- sec^{-1} px^{-1})	15
Dose rate on specimen (e^- sec^{-1} px^{-1})	17
Defocus range (μm)	-2.3 to -1.1
Reconstruction	
Symmetry imposed	C1
Initial particle images (no.)	80,764
Final particle images (no.)	33,202
Resolution, FSC=0.143 (Å)	6.61

Table 4 – Quantification of condensation of BRD4-NUT variants*

GFP-BRD4-NUT	RFP-p300 positive	RFP-p300 negative
WT	100/100	100/100
mut1	99/100	89/100
mut2	100/100	92/100
BD1-BD2	16/100	0/100
BD1	18/100	0/100
BD2	99/100	99/100
K/E	21/100	3/100
ΔNPS	100/100	100/100
ΔBID	100/100	9/100
BD1-BD2-AD	99/100	1/100
BD1-BD2-ADmut1	0/100	0/100

* Foci forming cells/Total cell count

Table 5 – Quantification of co-localization of BRD4-NUT and p300*

GFP-BRD4-NUT	Pearson's R value	Co-localization ratio
WT	0.81±0.01	0.90±0.02
ADmut1	0.84±0.01	0.87±0.02
ADmut2	0.76±0.02	0.90±0.01
ΔNPS	0.76±0.02	0.86±0.03

* At least 10 cells were analysed (n=10)

Table 6 – Impact of Drug treatment on condensation

GFP-BRD4-NUT	JQ1	A485	TSA	CBP30	1,6-Hexandiol
WT	No	No	No	Yes	No
ADmut1	No	No	No	No	No
ADmut2	No	No	No	No	No
ΔNPS	No	No	No	Yes	No

Table 7 – Quantification of foci formation of after CBP30 treatment*

GFP-BRD4-NUT	DMSO	CBP30
WT	100/100	54/100
ADmut1	92/100	3/100
ADmut2	91/100	2/100
ΔNPS	100/100	58/100

*Foci-forming cells/Total cell counted