# Structural insights into p300 regulation and acetylation-dependent genome organisation

Ziad Ibrahim[1], Tao Wang[2], Olivier Destaing[2], Nicola Salvi[3], Naghmeh Hoghoughi[2], Clovis Chabert[2], Alexandra Rusu[1], Jinjun Gao[4], Leonardo Feletto[1], Nicolas Reynoird[2], Thomas Schalch[1], Yingming Zhao[4], Martin Blackledge[3], Saadi Khochbin[2] & Daniel Panne[1,*]

[1]Leicester Institute of Structural and Chemical Biology, Department of Molecular and Cell Biology, University of Leicester, Leicester, UK

[2]CNRS UMR 5309, INSERM U1209, Université Grenoble Alpes, Institute for Advanced Biosciences, Grenoble, France.

[3]Institut de Biologie Structurale, CNRS, CEA, UGA, Grenoble, France

[4]Ben May Department of Cancer Research, The University of Chicago, Chicago, IL 60637, USA.

*Correspondence: daniel.panne@leicester.ac.uk

## Summary

**Histone modifications are deposited by chromatin modifying enzymes and read out by proteins that recognize the modified state. BRD4-NUT is an oncogenic fusion protein of the acetyl lysine reader BRD4 that binds to the acetylase p300 and enables formation of long-range intra- and interchromosomal interactions. We here examine how acetylation reading and writing enable formation of such interactions. We show that NUT contains an acidic transcriptional activation domain that binds to the TAZ2 domain of p300. We use NMR to investigate the structure of the complex and found that the TAZ2 domain has an autoinhibitory role for p300. NUT-TAZ2 interaction or mutations found in cancer that interfere with autoinhibition by TAZ2 allosterically activate p300. p300 activation results in a self-organizing, acetylation-dependent feed-forward reaction that enables long-range interactions by bromodomain multivalent acetyl-lysine binding. We discuss the implications for chromatin organisation, gene regulation and dysregulation in disease.**

**Keywords:** Chromatin modification, Acetylation, bromodomain, BRD4, acetyltransferase, p300, transcription factor, activation domain, gene regulation, NMR

## Introduction

Chromosome capture experiments have led to the identification of two major mechanisms that are behind large-scale chromosome organisation[1,2]: (1) at shorter genomic distances, genomes are organized into topologically associating domains[3,4] which arise through the process of loop extrusion by cohesin; (2) at longer-range genomic distances, genomes are spatially compartmentalised into active, euchromatic and inactive, heterochromatic regions[5,6].

One of the main mechanisms driving genome compartmentalisation is thought to be selective self-association of active or inactive regions and physical constraints including association with architectural elements[7,8]. One hypothesis is that 'readers' of histone modifications seed 'condensation' reactions as a function of the histone modification state[6,9]. It has been proposed that compartmentalisation arises specifically due to attractions between heterochromatic regions, possibly driven by polymer-polymer interactions and liquid-liquid phase separation (LLPS)[10,11]. Whether heterochromatic compartments form through LLPS or other mechanisms is controversial[12-15].

Histone acetylation counteracts heterochromatinization and has long been associated with chromatin decondensation and euchromatic chromatin regions[16]. Targeting of heterochromatin with acidic transcription factor (TF) activation domains (ADs) or treatment with deacetylase inhibitors results in hyperacetylation, decondensation and relocation to the nuclear interior even in the absence of active transcription[17-23].

Numerous TFs associate through their ADs with a small set of co-activators including Mediator and the acetylases CBP and its paralogue p300 (also known as KAT3A and KAT3B respectively; hereafter 'p300' for short)[24-29]. p300 has long been known to be important in integrating multiple signal transduction pathways in the nucleus and enabling enhancer-mediated transcription[30-32]. Intriguingly, it has been proposed that spatial sequestration of heterochromatin at the nuclear lamina requires TF-mediated retention of p300 in euchromatin[33]. Quantitative Mass Spectrometry experiments show that p300 is present at limiting concentrations in the nucleus while HDAC co-repressors are in large excess[34]. As p300 is responsible for up to a third of nuclear acetylation[35], TF-mediated competition for and retention of these essential and limiting co-activators in euchromatic regions could be a key driver event that not only shapes gene specific transcription but also eu- and heterochromatic genome compartmentalisation.

It has been proposed that transcriptional control may be driven by the formation of phase-separated condensates[36]. Accordingly, condensates of TF ADs[37], co-activators and the transcriptional machinery have all been proposed to be relevant for transcriptional regulation[38-43]. p300 contains an ordered catalytic 'core' domain including a HAT- and bromodomain (BD) and several domains that form promiscuous interactions with the disordered ADs of hundreds of cellular TFs (Fig. 1a)[44,45]. These interactions domains are separated by long, intrinsically disordered regions (IDRs). p300 is activated by autoacetylation[46], a reaction that can be controlled by cellular signalling through TF activation and dimerisation[47]. The p300 core contains a RING and TAZ2 domain that have an autoinhibitory function[44,47]. Perturbation of the autoinhibitory function of the RING and TAZ2 domains by mutation leads to constitutive p300

activation and formation of nuclear droplets that co-localise with p53. These droplets are
dispersed upon HAT or BD inhibition, suggesting that a form of phase separation (we here use
the more general term 'condensation') occurs through an acetylation-dependent 'read-write'
mechanism[47]. Other data indicate that p300 forms nuclear condensates through the 'core'
domain[48]. Co-condensation of p300 with TFs on enhancer loci is thought to enable activation of
p300's catalytic activity thus allowing the recruitment of acetylation-dependent coactivators,
including the BD - and extraterminal domain-containing (BET) protein BRD4[49].
94
BET proteins have long been associated with the control of large-scale nuclear structure and
chromatin organization. BRDT, a testes-specific BET, mediates histone acetylation-dependent
chromatin condensation[50] and is required for stage-specific gene expression and acetylation-
dependent histone removal during spermatogenesis[51-53]. Similarly, BRD4 has been linked both to
gene-specific[54] and higher-order chromatin organisation[55,56]. In yeast, the BET protein BDF1
maintains physical genome barriers by preventing spreading of heterochromatin[57]. *In vitro* data
show that under specific buffer conditions, chromatin can undergo LLPS which can be
modulated by acetylation and BRD4 binding[58]
103
BRD4 also forms condensates in cells[39]. *In vitro* and *in vivo* experiments with systems inducing
high local concentration suggest that a C-terminal IDR drives the condensation reaction[39,59].
However, a BRD4 isoform lacking the C-terminal IDR also forms condensates in cells[60]. Other
data indicate that BRD4 clustering does not depend on IDR-mediated condensation but on DNA
and BD-acetyl-lysine recognition which possibly enables long-range enhancer-promoter
communication[61]. At a larger scale, such acetylation-dependent long-range interactions could
enable selective self-association of euchromatic regions and provide a link between genome
topology and gene regulation. It is therefore important to understand the underlying 'read-write'
mechanisms that drive such long-range interactions.
113
BRD4-NUT is an oncogenic fusion protein that drives NUT carcinoma, one of the most
aggressive human solid malignancies known[62]. NUT is normally expressed in post-meiotic
spermatogenic cells, where it interacts with p300 and triggers genome-wide histone
hyperacetylation, which enables recruitment of BRDT, histone-to-protamine exchange and
chromatin compaction during spermatogenesis[53]. In the oncogenic fusion protein, NUT binds to
and activates p300 which drives an acetylation- and BRD4 BD-dependent 'feed-forward' loop
that leads to formation of hyperacetylated nuclear 'foci' visible by light microscopy[62-64], that
resemble foci seen in cells expressing constitutively activated p300[47]. These foci correspond to
large chromosomal 'megadomains' (100 kb to 2 Mb) that are enriched for BRD4-NUT, p300 and
acetylated histones[65-67]. Megadomains form in the absence of transcription and engage in very
long-range acetylation-dependent inter- and intrachromosomal interactions[68].
125
The BRD4-NUT model thus exemplifies how a modification of BRD4 enables recruitment and
activation of p300 and establishment of a self-organizing feed-forward reaction that results in
chromatin hyperacetylation and long-range contact. The underlying mechanisms could have
broad implications for our understanding of how 'reading' and 'writing' of chromatin acetylation
control gene expression and how dysregulation of this process leads to aggressive forms of
disease.

132
Here we use the BDR4-NUT/p300 system to dissect the mechanism that drives large-scale
chromatin organization. We determined the element of BRD4-NUT required for interaction with
p300. We found that the NUT portion of the fusion protein contains an acidic AD that interacts
with the p300 TAZ2 domain, a common binding site for numerous TF ADs. We investigate the
solution behavior of the complex and found that interaction with the TAZ2 domain stimulates
p300 HAT activity. Our data indicate that the TAZ2 domain has an auto-inhibitory function that
is relieved upon TAZ2 ligand-binding. We systematically probed which elements of BRD4-NUT
and p300 drive large scale chromatin condensation in cells. We found that condensation depends
on a feed-forward mechanism that involves allosteric activation of p300 by binding of NUT to
the p300 TAZ2 domain, nucleation of the condensation reaction through BRD4 and p300 BD-
acetyl-lysine binding and that BD multivalency is required. We found that Adenoviral E1A,
another TAZ2 ligand, or mutations associated with cancer that interfere with TAZ2
autoinhibition also activate p300. Our data reveal how activation of p300 establishes a self-
organising positive feedback mechanism and how an acetylation-dependent read-write
mechanism drives larger-scale chromatin architecture.
148
**Results**
**Identification of an acidic activation domain in NUT**
Previous data showed that a region spanning BRD4-NUT amino acids 347-588 (NUT F1c; NUT
numbering is used; 1060-1307 in BRD4-NUT) directly interacts with p300 (Fig. 1a,b)[63]. To map
the interaction further, we produced a series of deletions within NUT and assayed their
interaction with p300 using co-sedimentation and GST pulldown assays. NUT fragments that
contained amino acids 396-482 generally co-sedimented with the p300 core-CH3, the isolated
p300 CH3 or TAZ2 domains (Fig. S1a,b; Fig. S2a-f).
157
The TAZ2 domain forms dynamic interactions with short ~30 amino acid transcription factor
activation domains (ADs) that are unrelated in sequence but generally share net negative charge,
hydrophobic, and aromatic residues that are typically intrinsically disordered (IDR) in
isolation[45]. Accordingly, sequence analysis revealed that the region of NUT is predicted to be
disordered, and negatively charged with modest propensity to self-aggregate (Fig. 1c). A
luciferase assay confirmed that the isolated NUT region can stimulate transcription (Fig. S1c), in
agreement with previous data showing that tethering of NUT on chromatin is sufficient to induce
transcriptional activation[67]. We therefore conclude that NUT contains an acidic AD that directly
interacts with the TAZ2 domain of p300 (Fig. 1b).
167
The high percentage of negatively charged amino acid residues in ADs is thought to prevent
hydrophobic collapse while the distribution of charges maintains the conformational ensemble in
a more expanded state leaving the charged and hydrophobic side chains exposed to the solvent
and binding partners[69]. Accordingly, analysis by size-exclusion chromatography–multi-angle
laser-light scattering (SEC–MALS) showed that a NUT fragment spanning amino acids 347-480
(NUT4) is monomeric and monodisperse in solution with a large hydrodynamic radius,
consistent with an extended conformation (Fig. 1d). The TAZ2:NUT4 complex formed a 1:1
complex with a reduced hydrodynamic radius, indicative of conformational compaction upon
binding (Fig. 1d).
177

**Figure 1. Identification of an acidic activation domain in NUT**. **a**, Domain structure of p300 and **b**, BRD4-NUT constructs. **c,** IUPred, prediction of intrinsic disorder; FOLD, folding prediction using the foldindex (blue), PLAAC (**P**rion-**L**ike **A**mino **A**cid **C**omposition; orange) and PAPA (**P**rion **A**ggregation **P**rediction **A**lgorithm; gray) analysis, NCPR (**N**et **C**harge **P**er **R**esidue with a sliding window of 10 residues of the BRD4-NUT protein. **d**, SEC-MALLS analysis of the TAZ2 domain of p300 alone (black), NUT4 alone (blue) and the complex between TAZ2 and NUT4 (red). SDS-PAGE analysis of the peak fractions of the TAZ2-NUT4 complex is shown. **e,** Competition analysis: Cos7 cells were transfected with constructs expressing GFP-BRD4-NUT (green) alone or co-transfected with RFP-NUT-F1c and RFP-NUT4 (red). Immunofluorescence with anti-p300 antibody was used to visualize endogenous p300. Scale bars, 10 μm.

Co-expression of NUT F1c can prevent condensate formation in the nucleus by acting as a
dominant negative inhibitor for the p300-NUT interaction[63,67]. Accordingly, overexpression of
RFP-NUT F1c, -NUT4, or -NUT2 but not -NUT1 or -NUT3 reduced p300 foci in COS cells
(Fig. 1e, Fig. S2g). As AD function is usually confined to short sequences, we further mapped
the interaction by limited proteolysis followed by in-gel acid hydrolysis and mass spectrometry
(Fig. S1d). Fragments spanning amino acids 396-482 (NUT9) or a peptide spanning 458-482
(NUT10) all bound to the TAZ2 domain (Table S1).
185
**TAZ2-NUT interaction.**
NMR spectra of NUT7, spanning 448-492, feature little dispersion of $^1$H resonances (Fig. 2a),
indicating that the protein is largely disordered in solution. We used a series of 3D experiments
(see Methods) to assign backbone resonances of NUT7. Secondary structure propensities (SSP)
calculated using $^{13}$Cα, $^{13}$C' and $^{15}$N chemical shifts show that amino acids 451-472 have a
significant tendency to sample helical conformations in solution (Fig. 2b). We conclude that the
NUT7 AD is an IDR in isolation with a pre-organised N-terminal α helix.
193
We monitored binding to the TAZ2 domain by collecting a series of $^1$H, $^{15}$N-HSQC spectra of
$^{15}$N-labeled NUT7 mixed with increasing concentrations of TAZ2 (Fig. 2a). Chemical shift
perturbations (CSP) showed that several NUT7 backbone amide resonances, including A450,
L451, G460-A464, V467 and E475 were perturbed upon addition of an equimolar amount of
TAZ2 while resonances from the C-terminus of NUT7 do not appear to shift significantly upon
addition of TAZ2 (Fig. 2c). Thus, the N-terminal α-helical segment of NUT7 is involved in
TAZ2 binding. Addition of equimolar amounts of unlabelled NUT7 to $^{15}$N-labeled TAZ2
showed significant CSPs of resonances corresponding to residues R1737, I1739, M1761, R1763,
V1765, Q1766, T1768, K1783, Q1784, I1809 (Fig. 2a, c), that are mostly located on the surface
of TAZ2.
204
To understand the binding mode, we measured both $^{13}$C- and $^{15}$N-HSQC-NOESY spectra on a
sample of $^{15}$N,$^{13}$C-labelled TAZ2 to which one equivalent of unlabelled NUT7 was added. We
also measured $^{15}$N-HSQC-NOESY spectra on $^{15}$N-labelled NUT7 in admixture with one
equivalent of unlabelled TAZ2. The NOESY spectra contain very few cross-peaks that can be
unambiguously assigned to intermolecular contacts most likely due to extensive dynamic
fluctuations of NUT7 even in the bound state. One contact is made between the backbone amide
hydrogen of S1726 in TAZ2 and the backbone amide hydrogen of residue L463 and the Hβ of
A464 in NUT7. Because of relative sparsity and ambiguity of structural restraints, we used
HADDOCK[70] to derive a structural model of the interaction consistent with the observed
spectroscopic changes (Fig. 2d and e). NMR restraints and structural statistics are summarized in
Table S2. On the basis of this model, it appears that the N-terminus of NUT7 folds into an
amphipathic helix that binds a hydrophobic groove in the interface between helices α1/α2/α3 on
the surface of TAZ2 (Fig. 2d). Only the NUT backbone residues P448 to A464 are well ordered
while residues at the C-terminus (Q465-D492) do not interact with TAZ2 and remain disordered.
A set of hydrophobic side chains of NUT L449, A450, L451, L455, L461 are in contact with
TAZ2 residues I1735, M1761, V1764, I1781 (Fig. 2d). The interaction is further stabilised by
electrostatic interactions involving NUT residues E453, E454, E456, E458, E459 and TAZ2
residues R1737, S1754, S1757 and K1760 (Fig. 2d). In summary, similar to other TAZ2:AD

**Figure 2. TAZ2:NUT interaction**. **a**, (Left) Overlay of $^1$H-$^{15}$N HSQC spectra of free TAZ2 (black) and after addition of one molar equivalent of NUT7 (448-492) (blue). Arrows indicate chemical shift changes for individual cross-peaks. (Right) Overlay of $^1$H-$^{15}$N HSQC spectra of free NUT7 (448-492) and after adding half (orange) and one (brown) molar equivalent of TAZ2. Arrows indicate chemical shift changes for individual cross-peaks. **b**, Secondary structure propensities (SSP) plot of the NUT (448-492) fragment in its free state. positive values on the Y-axis indicate a-helical propensity, the X-axis is the amino acid sequence. **c**, (Left) Chemical shift perturbations on TAZ2 upon adding one molar equivalent of unlabeled NUT7 (448-492). (Right) Chemical shift perturbations on NUT7 (448-492) upon adding one molar equivalent of unlabeled TAZ2. **d**, (Left) Ribbon representation of lowest energy NMR model of the TAZ2 (blue)-NUT (orange) complex. Hydrophobic residues on the surface of NUT are shown. (middle) Zoom-in view of the interface between NUT (orange) and TAZ2 (blue). Residues on both sides implicated in the interaction as well as the hydrogen bonding network are shown. (Right) acidic residues on the surface of NUT (top) as well as hydrophobic and basic residues on the surface of TAZ2 (bottom) are shown. **e,** TAZ2 and NUT7 (448-492) are surface rendered and colored according to electrostatic potential (left) and sequence conservation (right).

223  interactions, a positively charged TAZ2 domain interacts with a negatively charged NUT AD
224  (Fig. 2e). The key interfacial binding surfaces are conserved in p300 and NUT (Fig. 2e).
225
**Analysis of the TAZ2:NUT interaction**
As ADs are frequently bi-partite and the NUT7 fragment in our structure only partially engages
the TAZ2 binding surface, we further explored the region N-terminal of NUT7. We found using
HSQC competition, that the NUT7 fragment is unable to displace p53 from the surface of TAZ2
when added to the complex (Fig. S3d). Biolayer Interferometry (BLI) experiments showed that
NUT8 (396-470) bound to TAZ2 with a dissociation constant of $K_d = 0.96$ +/- 0.29 µM (Fig. 3a,
b). Alanine substitutions of conserved residues in the region spanning 447-462 (mut3, Fig. 3c),
the main binding epitope in NUT7, did not abolish TAZ2 binding in GST-pulldowns, but
reduced TAZ2 binding affinity (Fig. 3b, Fig. S3a,b). Additional Alanine substitution of L436,
L440, L441 (mut4, Fig. 3c) further reduced TAZ2 affinity (Fig. 3b). In contrast, binding was
largely abolished by additional mutation to Alanine of L410, L411, Y413, L427, C428, F423,
V424, V427, V430 and I431 (mut1, Fig. 3b-d, Fig. S3a). As alanine repeat expansion can alter
transcription factor condensation properties[71], we also generated a NUT variant containing
mostly Glycine, Serine and charge reversal mutations (mut2, Fig. 3c). Mut2 also failed to bind
TAZ2 in BLI and GST-pulldown experiments (Fig. 3b-d, Fig. S3a).
241
242  As mutation of isolated binding epitopes reduced but did not abolish TAZ2 binding, we conclude
243  that the NUT AD is made up of subdomains that contain short amphipathic interaction motifs
244  that synergise to enable high affinity binding. TF ADs frequently bind p300 through a minimal
245  $\Phi XX\Phi\Phi$ motif (X is any residue and $\Phi$ is hydrophobic), frequently LXXLL[72]. The NUT AD
246  contains several such conserved motifs most of which are predicted to engage TAZ2 (Fig. 3c,e).
247
In our structure, the isolated AD sub-domain in NUT7 is positioned at the interface of the TAZ2
$\alpha 1$, $\alpha 2$ and $\alpha 3$ helices, which frequently functions as a docking site for TF ADs that typically
adopt helical structure upon binding (Fig. S3f). AlphaFold (AF) computational models predict 5
helices for NUT (Fig. 3e) that engage the protein interaction surfaces of TAZ2 (Fig. 3f, Fig.
S3c,g)[73,74]. In the model, the NUT7 AD is re-positioned while the TAZ2 $\alpha 1$, $\alpha 2$, $\alpha 3$ surface is
occupied by $\alpha 3$ of the NUT AD. Structure determination of the TAZ2 domain in complex with
the entire NUT AD was not possible as NMR spectra suffered from severe broadening (data not
shown). As (1) isolated AD sub-domains can interact with alternate TAZ2 surfaces[75] and (2) as
the residues that impair binding are located in the predicted TAZ2-NUT binding interface, these
data provide a reasonable model for NUT-p300 interaction.
258
**The role of the TAZ2 domain in p300 regulation and substrate acetylation**
Previous data show that NUT stimulates p300 auto- and histone acetylation in cells and *in*
*vitro*[53,63]. The AF model of p300 suggests that the TAZ2 domain is positioned over the p300
HAT active site (Fig. 4a). The TAZ2 C-terminal helix (amino acids 1806-1836) is sandwiched
between the RING and HAT domains proximal to the lysine substrate-binding tunnel in a
conformation that is incompatible with substrate access. In contrast, a model in the presence of
NUT AD shows that the TAZ2 domain is displaced from the HAT active site (Fig. 4b). One
possibility is that the TAZ2 domain auto-inhibits the HAT domain and that this auto-inhibition is
relieved upon ligand-binding to the TAZ2 domain. As the TAZ2 domain binds numerous

**Figure 3. Analysis of the TAZ2-NUT interaction. a**, Representative Biolayer Interferometry (BLI) profiles of p300 TAZ2-NUT binding from which the steady-state data were derived. **b**, Steady state analysis of p300 TAZ2 interaction with wild-type and NUT mutants using BLI. Error bars represent the standard deviations from three technical replicate experiments. **c**, Sequences of the NUT AD and mutants (in bold red) used. Top: Skylign plot showing NUT AD sequence conservation; Bottom: predicted secondary structure. **d**, GST pulldown analysis of TAZ2 and NUT8 (396-470) wild type and mutants 1 and 2. **e,** AlphaFold prediction of NUT (396-482) structure. **f**, Left: AlphaFold prediction of the TAZ2-NUT complex structure. Right: mutated residues in NUT mutant 1 (blue) and 2 (green) are shown.

268  transcription factor ADs, such a TAZ2-mediated control mechanism may be broadly relevant and
269  a key aspect of how TFs control chromatin acetylation and gene regulation.
270
To test this model, we incubated increasing amounts of GST-NUT8wt with p300s (amino acids
324-2094) in the presence of [$^{14}$C] acetyl coenzyme A. Addition of NUT WT but not mut1
stimulated p300 autoacetylation in a dose-dependent manner (Fig. 4c). Despite the identical
number of lysine amino acids, only NUT WT but not mut1 was acetylated by p300. Other NUT
fragments that contain the AD also activated p300 (Fig. S4a,b). These data suggest that the
TAZ2 domain autoinhibits p300 and that AD-TAZ2 interaction relieves autoinhibition.
Accordingly, acetylation reactions with the core domain of p300, lacking TAZ2 (Fig. 1a; p300
amino acids 1048-1664), showed that p300 autoacetylation is not stimulated by the presence of
NUT WT and both WT and mutant substrates are acetylated at equal rates (Fig. 4d). Another
classic TAZ2 ligand, Adenoviral E1A also showed a dose-depended stimulation of p300s auto-
and substrate acetylation (Fig. 4e).
282
Recent data indicate that p300 directly engages nucleosome core particles (NCPs) through its
ZZ[76] and TAZ2 domains[77]. Would then NCP engagement also activate p300? In accordance with
this possibility, it has been shown that the TAZ2 domain is required for efficient acetylation of
H3K27, while curbing activity towards other lysine residues within nucleosomes[77]. To test this
model, we generated homogeneous 'structure-grade' NCPs as shown by single particle cryoEM
(Fig. S4f, Table S3). p300 variants containing the TAZ2 domain, as well as the isolated TAZ2
domain directly bound to NCPs in EMSA experiments (Fig. S4c,d,e). Addition of increasing
amounts of NCPs stimulated auto-acetylation of p300s but not p300 core (Fig. 4f).
291
Cross-linking mass spectrometry experiments showed that surface residues from the p300
Bromo-, RING, PHD and TAZ2 domains directly engage the N-terminal regions of Histone H3
and H2A (Fig. S4i,j). Label free quantitative mass spectrometry experiments showed that p300
core acetylated most lysine residues indiscriminately in Histones H2B, H3 and H4 while p300s
showed a more selective acetylation pattern (Fig. S4g,h). We therefore conclude that direct NCP
binding through the TAZ2 domain can stimulate p300 activation but contributes modestly to
substrate specificity.
299
To further test the autoinhibition model, we truncated amino acid residues Δ1813-1838 of the
TAZ2 domain (Fig. 4a). Deletion of the C-terminal helix resulted in the formation of p300
nuclear condensates (Fig. 4g) that are typically seen upon p300 activation[47]. p300 mutations
found in cancer cell lines that truncate the TAZ2 domain (Fig. S4k) also result in p300
activation[78,79]. Together, we conclude that the TAZ2 domain autoinhibits p300 HAT activity.
Interaction of TF ADs with the TAZ2 domain allosterically regulates p300 HAT activation by
relief of autoinhibition. Mutations that interfere with TAZ2 autoinhibition are found in cancer.
307
**Mechanism of condensation of BRD4-NUT and p300**
BRD4-NUT forms hyperacetylated chromatin megadomains in patient tumors[63,65,68]. Formation
of such megadomains can be reproduced by expression of BRD4-NUT in heterologous cell
lines[63,65,68]. As there is strong evidence that this system reproduces the phenoptype seen in
patient cells, we consider it as physiologically relevant, at least with respect to the biophysical
properties of the condensation reaction. We generated a series of GFP-BRD4-NUT variants and

**Figure 4. Role of the TAZ2 domain in p300 regulation and substrate acetylation. a**, AlphaFold prediction of p300 core-CH3 (1048-1838) structure. The structure is coloured according to domain structure. The TAZ2 domain is positioned in the HAT active site thus autoinhibiting substrate access. The TAZ2 C-terminal helix deletion (Δ1813-1838) is coloured in magenta **b,** NUT AD binding displaces the TAZ2 domain from the HAT active site thus enabling allosteric HAT activation. **c,** p300s or **d**, p300 core were incubated with [14C]Acetyl-CoA in the presence or absence (C: p300 alone) of increasing concentrations of wild type or mut1 NUT8 (396-470), **e**, Increasing concentrations of GST-E1a we incubated with p300s, **f**, Increasing concentrations of NCPs were incubated with p300s or p300 core. Panels c-f: Top two images: Analysis by SDS–PAGE followed by Coomassie staining. Bottom two images: 14C phosphor imaging. Representative data of three independent experiments are shown. **g**, Cos7 cells were transfected with the HA-tagged p300 WT (top) or p300 Δ1813-1838 (bottom) and analysed by anti HA immunofluorescence. Four representative nuclei are shown. The percentage of cells showing condensates (n=100 cells) is indicated. Scale bars, 5 μm.

assessed their propensity to interact with p300 and form co-condensates in the cell nucleus. Co-
immunoprecipitation experiments showed that wild-type GFP-BRD4-NUT but not the AD
mutants bound to endogenous p300 (Fig. 5a). In agreement with our *in vitro* data, the wild-type
but not mutant variants showed a signal at the level of GFP-BRD4-NUT when probed with an
anti-acetyl lysine antibody (Fig. 5a). GST-pulldowns in nuclear extracts using a purified GST-
NUT8wt fragment showed signal for p300 only for the wild-type but not the AD mutant variants
(Fig. 5b). Thus, mutation of the NUT AD interferes with p300 interaction and BRD4-NUT
acetylation in cells, as also seen with the corresponding fragments *in vitro* (Fig. 4c).
322
Co-transfection experiments showed that GFP-BRD4-NUTwt and RFP-p300 formed nuclear co-
condensates (Fig. 5c; row 1; for quantification see Tables S4 and S5) both in co-transfected and
RFP-p300 negative cells. Thus, condensation of GFP-BRD4-NUTwt not only concentrates but
also depletes p300 from other areas of the nucleus. These condensates were independent of the
NUT AD as mutant variants still formed condensates that co-localized with p300 (Fig. 5c; row
2&3) and acetylated chromatin (Fig. S5a). Mutation of the acetyl-lysine binding pocket in BD1
(N140A) and BD2 (N433A) completely abolished GFP-BRD4-NUT and RFP-p300 co-
condensation (Fig. 5c; row 4). To assess the model that BRD4 multivalency drives the
condensation reaction[58], we mutated BD1 and BD2 individually. We found that mutation of BD1
but not BD2 largely abolished condensation (Fig. 5c; row 5, 6). Thus, the acetyl-lysine binding
ability of BD1 is required to nucleate BRD4-NUT and p300 co-condensation.
334
We reported previously that BRDT interacts with nucleosomes through its first (BD1), but not
second (BD2) bromodomain, and that acetylated Histone recognition by BD1 is complemented
by a direct BD1–DNA interaction[80]. Another report indicates that DNA binding promotes
condensation of a short BRD4S 'isoform' *in vitro*[60]. We therefore introduced charge inversion
mutations in a conserved, basic surface patch in BD1 (R68E, K72E, K76E) that is involved in
DNA binding in BRDT[80] and in the equivalent BD2 amino acids (S358E, K362E, A366E).
Mutation of the basic patch abolished condensation (Fig. 5c; row 7). Thus, we propose that
BRD4 BD-DNA interaction facilitates acetylation-dependent NCP binding and chromatin
condensation in cells.
344
We also probed the BRD4 N-terminal cluster of phosphorylation sites (NPS) and the basic
residue-enriched interaction domain (BID), elements that have been proposed to enable BRD4
regulation by phosphorylation[81]. Deletion of the NPS did not interfere with condensation while
removal of the BID reduced condensation (Fig. 5c; row 8, 9). A minimal BD1-BD2 variant fused
to the AD but not an AD Mut1 variant (Fig. 1b) still formed p300 co-condensates (Fig. 5c; row
10, 11, Movie S1). The BD1-BD2-AD fusion protein but not the AD Mut1 variant was still able
to co-IP p300 from cells indicating that the ability to interact with p300 is maintained in the
fusion protein (Fig. S5c). Fusion of the minimal BD1-BD2 variant to the AD of the Adenoviral
E1A (Fig. 1b) also resulted in co-condensation with p300 (Fig. 5c; row 12). We therefore
conclude that, at a minimal level, both BRD4 BD acetyl-lysine binding and TF AD-mediated
p300 recruitment and activation are required for co-condensation.
356
357  Addition of 1,6-hexanediol, the p300/CBP HAT inhibitor A-485 or the BET inhibitor JQ1 all
358  abolished condensation of GFP-BRD4-NUTwt, mut1, mut2 and ΔNPS confirming that both
359  acetylation by p300/CBP and BRD4 BD substrate engagement are required (Fig. 5d, Table S6

**Figure 5. Mechanism of condensation of BRD4-NUT and p300. a,** Cos7 cells were transfected with GFP-BRD4-NUT variants or GFP alone as the control. Co-immunoprecipitation (IP) experiments were performed using anti-GFP antibody and immuno-blotted (IB) for p300, GFP and Acetylated-Lysine. Input samples were loaded as a control. Experiments were independently repeated at least three times. **b,** Purified GST-NUT8 wild type and mutants 1 & 2 were incubated with Cos7 cells extract for 4 hours. The complexes were pull-down by Glutathione Sepharose beads and analysed by SDS-PAGE followed by Coomassie staining (bottom) or by IB for p300 (top). **c**, Cos7 cells were transfected with the indicated GFP-BRD4-NUT and RFP-p300 constructs and analysed by fluorescence microscopy, BRD4-NUT (green) and p300 (red). Scale bars, 10 μm. Quantification of foci formation of different BRD4-NUT variants and their co-localization with p300 are shown in Tables S4 and S5. **d**, Cos7 cells were transfected with the indicated GFP-BRD4-NUT constructs and treated with DMSO (control), 1,6-hexanediol, p300 HAT inhibitor A485, pan-BET inhibitor JQ1, HDAC inhibitor TSA or p300 bromodomain inhibitor CBP30. Scale bars, 10 μm. Quantification of foci formation in presence of CBP30 is shown in Table S6. **e**, Shematic representation of GFP-labelled BRD4 mutants and Alexa Fluor 594 labelled reconstituted chromatin containing non-acetylated (12x NCP) or acetylated (Ac 12X NCP) 12 Nucleosome Core Particles. Fluorescence microscopy images of non-acetylated or acetylated 12x NCP chromatin incubated with GFP-BD1-BD2. Scale bars, 10 μm.

and S7). HDAC inhibition results in expansion of Histone acetylation and retargets lysine-acetyl
readers including BRD4 thus partially mimicking the effect of BD inhibition[82]. In agreement
with this model, addition of the HDAC inhibitor Trichostatin A abolished condensation of all
variants (Fig. 5d). Immunofluorescence experiments further showed that the BRD4-NUT
condensates colocalise with HDAC1 thus confirming that these sites correspond to high
acetylation turnover (Fig. S5b).
366
As mutation of the NUT AD did not abolish p300 recruitment into the condensate we further
assessed the contribution of the CBP/p300 BD. We used the CBP30 inhibitor that has~34-fold
higher selectivity for CBP/p300 BD as compared to BRD4 BD1[83]. Addition of CBP30 did not
impact the condensation of GFP-BRD4-NUTwt and ΔNPS. However, condensation of the NUT
AD mutants GFP-BRD4-NUTmut1 and mut2 was sensitive to CBP30 treatment (Fig. 5d, Table
S7). Thus, the NUT AD mutants reveal that co-condensation can also occur through a p300 BD-
acetyl-lysine-dependent recruitment mechanism. Thus, two interactions drive BRD4-NUT and
p300 co-condensation: (1) interaction of p300 with the AD domain in NUT and (2) p300 BD-
acetyl-lysine engagement. Disruption of both interactions, but not each interaction individually,
abolishes the co-condensation reaction.
377
**Condensation requires BD multivalency**
Our observation that mutation of BD1 but not BD2 largely abolished BRD4-NUT condensation
in cells (Fig. 5c; row 5, 6) suggests that condensation in cells is not simply driven by BRD4 BD
divalency. We therefore tested the requirements of acetylation-dependent chromatin
condensation *in vitro*. Previous data show that addition of BRD4 or an engineered multivalent
protein containing five copies (BD1)$_5$ of BRD4 BD1 induces condensation of p300-acetylated
nucleosome arrays *in vitro*[58].
385
We generated dodecameric nucleosome arrays (Fig. 5e, Fig. S5d). In agreement with previous
data[58], non-acetylated nucleosome arrays formed round droplets that were dissolved by p300
acetylation (Fig. S5e), in agreement with the model that acetylation disrupts internucleosomal
interactions[84]. As expected, addition of GFP-(BD1)$_5$ allowed droplet formation of acetylated but
not of non-acetylated nucleosome arrays (Fig. S5f). This condensation reaction was driven by
acetyl-lysine binding as a N140A mutation in the BD1 acetyl-lysine binding pocket abolished
droplet formation (Fig. S5g).
393
As a minimal BRD4 construct containing BD1-BD2 (amino acids 1-459) induced condensation
in cells when fused to the NUT AD, we tested if BD1-BD2 divalency is sufficient to drive
droplet formation *in vitro*. BRD4 BD1-BD2 enabled droplet formation of acetylated but not of
non-acetylated nucleosome arrays (Fig. 5e, right) in the same way as the GFP-(BD1)$_5$. Mutation
of the BD acetyl-lysine binding pockets individually or in combination abolished droplet
formation (Fig. S5g). We conclude that the BD divalency of BRD4 is sufficient to drive
condensation *in vitro*. As large-scale condensation in cells is sensitive to BD1 but not BD2
mutation (Fig. 5c), our model is that BRD4 oligomerisation contributes to condensation.
402
403    **Dynamics of condensation of BRD4-NUT and p300**
404    Time-lapse imaging showed that the condensates remained stable during the experiment (Fig. 6a,
405    Movie S2). Addition of 1,6-hexanediol resulted in rapid and transient dissolution of the

**Figure 6. Dynamics of condensation of BRD4-NUT and p300. a,** Example image of nuclear droplets. Red square: Time laps imaging of BRD4-NUT and p300 droplets; Scale bar, 4 µm. **b,** Time laps imaging after addition of 3% [w/v] 1,6 Hexanediol. Scale bar, 12 µm. **c**, Top: Example image of droplets (red square) after photobleaching. Scale bar, 2 µm. Bottom: Quantification of the FRAP experiments. 3 independent technical replicates were done. Averages and standard error of mean (SEM; grey bars) are shown. **d**, Quantification of the characteristic time of recovery rates for different GFP-BRD4-NUT mutants (top) and RFP-p300 (bottom). Three to four technical replicate FRAP experiments were done for each mutant (40 to 65 foci measured). Data shown are mean +/- standard error of mean (SEM). Statistical differences in recovery rates were determined by Student's t-test. *p≤0.05; **p≤0.01; ***p≤0.001; n.s. non-significant difference as compared to WT. **e**, Half-bleach FRAP experiment. Top: The area before (left) and after (right) laser bleach are shown. The red box indicates the bleached area. Scale bar, 0.5 µm. Bottom: The change in fluorescence intensity along the A-B axis was analysed in the Kymograph. **f**, Top: Plot of fluorescence recovery from the Kymograph for the unbleached area, center and periphery of the droplet. Bottom: cartoon of the internal and external exchange.

condensate (Fig. 6b). After reformation, a second addition of 1,6-hexanediol dissolved the
condensates again illustrating their dynamic and transient nature. To study the consequences of
the BRD4-NUT mutations on turnover of the condensate we performed Fluorescence Recovery
After Photobleaching (FRAP) experiments. Wild-type GFP-BRD4-NUT and RFP-p300
recovered ~50% of signal after ~20 sec (Fig. 6c). Thus, while droplets appear stable at the
mesoscale, the molecules are highly mobile at the nanoscale and in constant exchange with the
nucleoplasm. Mutation of BRD4-NUT mostly interfered with droplet retention as shown by
increased FRAP recovery rates (Fig. 6d). p300 dynamics was directly coupled to that of the
BRD4-NUT mutants except for the GFP-BRD4-NUT BD2 and the BD1-BD2-AD constructs
which showed slower p300 recovery rates indicating some degree of functional uncoupling.
416
To assay internal droplet dynamics, we performed half-bleach FRAP experiments (Fig. 6e). We
analysed the change in fluorescence intensity over time along a defined A-B axis in the bleached
(red square) and unbleached area. Kymograph analysis showed that the fluorescence signal in the
'unbleached' region (orange line) and the droplet 'center' (blue line) recovered much quicker
and to completion as compared to the 'periphery' of the droplet (grey line), which only partially
recovered signal (Fig. 6f). Our interpretation is that recovery at the center occurs more quickly
because high local concentration in the droplet centre maintains a feed-forward acetylation
reaction that traps quickly diffusing molecules. At the droplet periphery, lower concentrations
and competition with HDACs destabilizes droplet assembly leading to reduced recovery rates.
As there is (1) constant exchange with the nucleoplasm and (2) no preferential equilibration
between the bleached and unbleached area we argue that these condensates are 'liquid'-like at
the nanoscale but not at the mesoscale where the viscoelastic properties of the 'solid' chromatin
scaffold constrain dynamic exchange[85]. As a result, we do not see droplet fission or fusion at
least during the time-course of our experiments.
431
**Discussion**
Chromatin acetylation has long been linked to transcriptional activation, by altering chromatin
structure and by enabling binding of BD containing proteins[86-88]. How acetylation promotes
transcription has remained elusive. p300 is responsible for up to a third of nuclear acetylation[35].
The BRD4-NUT fusion reveals how constitutive activation of p300 establishes a self-organizing,
acetylation-dependent feed-forward reaction that results in large-scale intra- and
interchromosomal interactions with an oncogenic outcome[62]. In addition to its well-known role
in enhancer function and control of gene expression[32,49], p300 is also involved in
compartmentalisation of the genome into eu- and heterochromatin[33]. We argue that key aspects
of these reactions involve modulation of an acetylation-dependent condensation reaction we
describe here.
443
Regulation of p300 is central for establishment of an acetylation-dependent condensation
reaction (Fig. 7a). We propose that in the deacetylated state p300 is auto-inhibited by the RING,
TAZ2 domains and the AIL. p300 is activated by trans-autoacetylation[47,89,90]. Due to low p300
concentration[34], self-collision and activation do not occur spontaneously. Instead, as HDACs are
estimated to be ~100-fold more abundant[34], p300 most frequently collides with HDAC
complexes[91,92] which maintain p300 in the inactive conformation. In this inactive state p300 is
evenly dispersed in the nucleus[47].
451

**Figure 7. Molecular model for p300 activation in physiological setting and disease**. **a**, p300 is maintained in the inactive state by HDACs. Activation requires TF binding which results autoacetylation and removal of autoinhibition. **b**, In enhancer signalling, TFs interact through their ADs with the different protein interaction domains of p300 which results in scaffolding of dynamic TF assemblies. Local TF and histone acetylation allows binding of BD containing proteins including BRD4 thus establishing positive feed-back. HDACs are located in these condensates and enable dynamic turnover thus resulting in bistability and response to changes in TF signaling. **c**, In BRD4-NUT NMC, the acetylation-dependent reaction drives formation of intra- and interchromosomal interactions resulting in the formation of a Megadomain. **d**, TF binding and acetylation establish open Euchromatin. TFs compete for limiting p300 co-activators but TF-DNA binding outside enhancer clusters remains transient and, due to high HDAC activity, does not effectively establish an acetylation-dependent condensate. **e**, Depletion and partitioning of p300 into euchromatic compartments results in hypoacetylation due to high HDAC activity. Hypoacetylated compartments become a substrate for heterochromatinization.

Induced proximity, by TF binding, enables p300 self-collision, trans-autoacetylation and activation[47]. The BRD4-NUT system reveals an autoinhibitory role for the TAZ2 domain which is relieved upon NUT AD binding (Fig. 4). Accordingly, the well-characterized TAZ2 ligand E1A[93,94] also stimulates p300 auto- and substrate acetylation (Fig. 4), as also reported for other TAZ2 ligands[95,96]. Thus, we propose that the TAZ2 domain plays a role in allosteric HAT regulation: TF AD binding results in displacement of the TAZ2 domain from its auto-inhibitory position resulting in HAT activation thus directly coupling TF AD-mediated p300 binding to HAT activation.

Perturbation of the autoinhibitory function of the RING and TAZ2 domains by mutation all lead to constitutive p300 activation and condensation reactions similar to what is seen with BRD4-NUT[44,47]. Importantly, comprehensive studies of cancer cell lines show that truncation of the TAZ2 domain (Fig. S4k) leads to p300 HAT gain-of-function in disease[78,79]. An acetylation-dependent condensation reaction may also explain how Adenoviral E1A causes genome-wide relocalisation of p300 and activation of genes that promote cell growth and division[97-99]. Not unlike BRD4-NUT, this condensation reaction is also sensitive to p300 HAT, BD and HDAC inhibition[99]. Therefore, while p300 is thought of as a classic tumor suppressor, with loss of function mutations occurring in a number of different cancers[100], constitutive activation by mutation or by interaction with activating ligands such as BRD4-NUT and E1A can lead to acetylation-dependent condensation with an oncogenic outcome.

As numerous TFs interact with p300 trough their ADs[31,45], we predict that the condensation reaction establishes intra- and interchromosomal interactions that are normally involved in enhancer signaling (Fig. 7b). Many TF ADs, including NUT (Fig. 1), are IDRs that form dynamic interactions with the different protein interaction domains of p300[75,101-110]. Due to such multivalent AD interactions, p300 is able to scaffold TF assembly on complex enhancer elements thereby facilitating combinatorial control of enhancer-mediated transcription[111-113].

TF-mediated recruitment and activation of p300 result in local histone and TF acetylation and establishment of positive feed-back through binding by BD containing proteins. High rates of self-collision of p300 maintain this positive feedback and establish a self-organizing feed-forward reaction resulting in co-condensation of p300, BET proteins BRD2, BRD3, and BRD4 and numerous transcriptional and chromatin regulators[66]. The condensation reaction is limited by HDACs which are usually present at active enhancers and promoters[114,115], and prevent hyperacetylation and transcriptional inhibition[116,117]. HDAC1 is localised in the BRD4-NUT condensate (Fig. S5b). Our FRAP experiments show that the condensates are 'liquid'-like at the nanoscale and we propose that HDACs enable continuous and dynamic turnover. If the feed-forward condensation reaction is sufficiently strong, it will overcome this inhibitory effect by HDACs provided that local high concentrations are maintained. As a result, the BRD4-NUT condensate is essentially stable at the mesoscale despite continuous internal dynamic exchange. In enhancer signaling, this dynamic bistability may then render the condensate more immediately responsive to dynamic and transient changes in TF signaling and explain the short transcriptional burst kinetics seen in cells[118,119].

The reaction redistributes p300 (and BD containing proteins) into the condensate thus leading to depletion from other genomic regions (Fig. 5)[63]. The condensation reaction thus leads to high

local acetylation (in the condensate) but global histone hypoacetylation elsewhere[120].
Counterintuitively, treatment with HDAC inhibitors does not lead to BRD4-NUT
'hypercondensation'. Instead, HDACi restores global histone acetylation[120] and rapidly dissolves
the BRD4-NUT/p300 condensate (Fig. 5). In agreement with recent data[82], our model is that
HDAC inhibition and resultant increased global histone acetylation redistributes BD containing
proteins thereby decreasing their effective local concentration resulting in droplet dissolution.
Due to the diffusible nature of the reactants, maintenance of global histone hypoacetylation is
therefore a key aspect of how such condensates arise. We therefore argue that HDACs contribute
to transcriptional regulation by maintaining (1) dynamic bistability of enhancer condensates and
(2) global hypoacetylation which in turn 'enables' focal enhancer condensation. Thus, the dual
nature of HDAC function with roles in gene activation and repression[114,121-123] would then be
directly linked to modulation of these condensation reactions.
510
While histone H3K27 acetylation has long been seen as a signature mark for active
enhancers[124,125], p300 also acetylates numerous non-histone substrates including many TFs[35].
Therefore, we do not see a role for a defined 'code' in how such modifications contribute to the
signalling reaction[126]. Instead, promiscuous lysine acetylation of TFs and histones drive
emergent multivalency that kinetically traps diffusing BD-domain containing factors such as
p300 and BRD4. While individual BDs typically have weak binding affinity and specificity[127],
local proximity facilitates rebinding after dissociation thus enabling an acetylation-dependent
feed-forward co-condensation reaction (Fig. 6). We propose that this biophysical mechanism
enables formation of an 'exploratory' network of interactions that generates bistability by
positive feedback and enables focal accumulation of regulatory factors on enhancers and
establishment of long-range interactions that control promoter activity[119]. Successful
establishment of this 'exploratory' network could then explain the long-standing observation that
multiple TF DNA binding sites synergistically activate transcription without the requirement of
physical interaction between TFs[112,128].
525
It has long-been known that TFs activate some genes while at the same time repressing others
through competitive coactivator 'squelching'[30,129-135]. Accordingly, in BRD4-NUT, the
sequestration of p300 inhibits p53 and the apoptotic cell response while knock-down of BRD4-
NUT releases p300, and restores p53-dependent cell differentiation and apoptosis[63]. Thus, we
predict that competition for diffusible co-activators and successful establishment of a continual
acetylation-dependent condensation reaction is a key regulatory control mechanism: only
enhancers containing the appropriate set of clustered TF binding sites will effectively compete
and establish such an acetylation-dependent condensate. As TFs locate their target DNA sites
using 'facilitated diffusion'[136,137], we propose that following TF-DNA binding, the combined
action of p300 scaffolding, TF acetylation and formation of an acetylation-dependent condensate
slows down the dynamic search behaviour thus increasing local concentration, residency time
and transcription activation. Accordingly, acetylation-mediated increase of TF residence time has
been described for p53[138,139] and could thus control TF turnover and transcriptional
regulation[140,141].
540
Binding events that occur outside TF enhancer clusters are then unable to compete for limited
co-activators, fail to establish such an acetylation-condensation reaction and therefore remain
transcriptionally inactive[142-144]. Due to high concentrations of HDACs[34] and rapid deacetylation

kinetics across the genome[35,91], the acetylation-dependent condensation reaction is transient. It is
such transient and dynamic acetylation that establishes euchromatic compartments without
necessarily leading to transcriptional activation (Fig. 7d).

There is increasing evidence suggesting a role for LLPS of intrinsically disordered proteins in
different nuclear processes[39,59,60]. IDR-driven condensation reactions of chromatin-binding
proteins including BRD4 and p300 have been implicated in gene regulation[37-39,41,42,49,59,60] and in
disease[145,146]. As BRD4-NUT and p300 are largely disordered (Fig. 1; 72% and 42% predicted
IDR, respectively) our assumption was that IDR-mediated interactions would contribute to the
condensation reaction.

While the NUT AD region can indeed form spherical condensates *in vitro* as analysed by SAXS
(Fig. S6), we do not think that this is relevant for the condensation reaction in cells. Even a
minimal BRD4 BD1-BD2 construct, removing a large fraction of IDR sequence from BRD4-
NUT, enables condensation in cells provided it contains a functional AD (Fig. 5c). As an
equivalent NUT AD mutant (that fails to bind to the TAZ2 domain) does not enable the
condensation we exclude the possibility that residual structural disorder, e.g. the 200 amino acid
linker between the BD1 and BD2 domains of BRD4, drives the condensation reaction. We and
others have shown that the condensation reaction is inhibited by (1) the BET BD inhibitor
JQ1[63,67], (2) mutation of the acetyl-lysine binding pocket of BRD4 BD1 or, (3) the DNA binding
surfaces of BRD4 BD1/BD2 or, (4) p300 HAT inhibition (Fig. 5). *In vitro*, divalency of BD1-
BD2 was essential for condensation of acetylated nucleosome arrays (Fig. 5e). BET proteins are
thought to oligomerize[50,56] and BRD4 phosphorylation enables dimerization[147]. We propose that
the combination of BD divalency and BET oligomerization provide multivalency that enables
condensation of acetylated chromatin in cells. Several p300 HAT gain-of-function mutations,
that interfere with the autoinhibitory function of the RING and TAZ2 domains, enable a similar
acetylation- and BD-dependent condensation reactions[47]. We therefore conclude that the reaction
is driven by p300 activation and engagement of acetylated chromatin by the BDs. In accordance,
single-gene imaging data support the notion that promoter-enhancer communication and
transcription control depend on TF and BRD4 clustering driven by specific DNA and BD
binding rather than IDR-mediated interactions[61].

The co-activator depletion reaction may also be one of the key reaction mechanisms that enables
the physical segregation of transcriptionally active euchromatin from repressed heterochromatin
(Fig. 7e), and explain the general link between the transcriptional state and 3D chromatin
compartmentalisation[148]. Targeting of heterochromatin with acidic TF ADs or treatment with
HDAC inhibitors generally counteracts heterochromatinization even in the absence of active
transcription[17-23,149-151]. TF-mediated retention of p300 in euchromatin enables establishment of
heterochromatin in worms[33], while depletion of p300 and chromatin hypoacetylation facilitates
methylation by PRC2[91]. Thus, competition for and partitioning of limiting key acetylases by TFs
into euchromatin and depletion from heterochromatin, together with BD-mediated higher order
chromatin folding[50,51,56], may drive key aspects of 'epigenetic' genome compartmentalization.
Phenomena such as stochastic and exclusive choice of monoallelic gene expression or X-
chromosome inactivation in mammals may then also depend on depletion of limiting
coactivators and establishment of positive feedback so that only the 'winning' allele drives gene
expression while 'competing' alleles are silenced[152-156].

## References

1. Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* **326**, 289-293, doi:10.1126/science.1181369 (2009).

2. Schwarzer, W. *et al.* Two independent modes of chromatin organization revealed by cohesin removal. *Nature* **551**, 51-56, doi:10.1038/nature24281 (2017).

3. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376-380, doi:10.1038/nature11082 (2012).

4. Nora, E. P. *et al.* Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**, 381-385, doi:10.1038/nature11049 (2012).

5. Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N. & Mirny, L. A. Chromatin organization by an interplay of loop extrusion and compartmental segregation. *Proc Natl Acad Sci U S A* **115**, E6697-E6706, doi:10.1073/pnas.1717730115 (2018).

6. Misteli, T. The Self-Organizing Genome: Principles of Genome Architecture and Function. *Cell* **183**, 28-45, doi:10.1016/j.cell.2020.09.014 (2020).

7. van de Werken, H. J. G. *et al.* Small chromosomal regions position themselves autonomously according to their chromatin class. *Genome Res* **27**, 922-933, doi:10.1101/gr.213751.116 (2017).

8. Solovei, I., Thanisch, K. & Feodorova, Y. How to rule the nucleus: divide et impera. *Curr Opin Cell Biol* **40**, 47-59, doi:10.1016/j.ceb.2016.02.014 (2016).

9. Wang, L. *et al.* Histone Modifications Regulate Chromatin Compartmentalization by Contributing to a Phase Separation Mechanism. *Mol Cell* **76**, 646-659 e646, doi:10.1016/j.molcel.2019.08.019 (2019).

10. Erdel, F. & Rippe, K. Formation of Chromatin Subcompartments by Phase Separation. *Biophys J* **114**, 2262-2270, doi:10.1016/j.bpj.2018.03.011 (2018).

11. Falk, M. *et al.* Heterochromatin drives compartmentalization of inverted and conventional nuclei. *Nature* **570**, 395-399, doi:10.1038/s41586-019-1275-3 (2019).

12. Larson, A. G. *et al.* Liquid droplet formation by HP1alpha suggests a role for phase separation in heterochromatin. *Nature* **547**, 236-240, doi:10.1038/nature22822 (2017).

13. Strom, A. R. *et al.* Phase separation drives heterochromatin domain formation. *Nature* **547**, 241-245, doi:10.1038/nature22989 (2017).

14. Erdel, F. *et al.* Mouse Heterochromatin Adopts Digital Compaction States without Showing Hallmarks of HP1-Driven Liquid-Liquid Phase Separation. *Mol Cell* **78**, 236-249 e237, doi:10.1016/j.molcel.2020.02.005 (2020).

15. Strom, A. R. *et al.* HP1alpha is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. *Elife* **10**, doi:10.7554/eLife.63972 (2021).

16. Ho, J. W. *et al.* Comparative analysis of metazoan chromatin organization. *Nature* **512**, 449-452, doi:10.1038/nature13415 (2014).

17. Tumbar, T., Sudlow, G. & Belmont, A. S. Large-scale chromatin unfolding and remodeling induced by VP16 acidic activation domain. *J Cell Biol* **145**, 1341-1354, doi:DOI 10.1083/jcb.145.7.1341 (1999).

18. Carpenter, A. E., Memedula, S., Plutz, M. J. & Belmont, A. S. Common effects of acidic activators on large-scale chromatin structure and transcription. *Mol Cell Biol* **25**, 958-968, doi:10.1128/MCB.25.3.958-968.2005 (2005).

19. Therizols, P. *et al.* Chromatin decondensation is sufficient to alter nuclear organization in embryonic stem cells. *Science* **346**, 1238-1242, doi:10.1126/science.1259587 (2014).

20. Taddei, A., Maison, C., Roche, D. & Almouzni, G. Reversible disruption of pericentric heterochromatin and centromere function by inhibiting deacetylases. *Nat Cell Biol* **3**, 114-120, doi:10.1038/35055010 (2001).

21. Mueller, W. G., Walker, D., Hager, G. L. & McNally, J. G. Large-scale chromatin decondensation and recondensation regulated by transcription from a natural promoter. *Mol Biol Cell* **12**, 357a-357a (2001).

| 640 | 22 | Toth, K. F. *et al.* Trichostatin A-induced histone acetylation causes decondensation of interphase |
| 641 | | chromatin. *J Cell Sci* **117**, 4277-4287, doi:10.1242/jcs.01293 (2004). |

22 Toth, K. F. *et al.* Trichostatin A-induced histone acetylation causes decondensation of interphase
chromatin. *J Cell Sci* **117**, 4277-4287, doi:10.1242/jcs.01293 (2004).

23 Chambeyron, S. & Bickmore, W. A. Chromatin decondensation and nuclear reorganization of the
HoxB locus upon induction of transcription. *Gene Dev* **18**, 1119-1130, doi:10.1101/gad.292104
(2004).

24 Avantaggiati, M. L. *et al.* The SV40 large T antigen and adenovirus E1a oncoproteins interact
with distinct isoforms of the transcriptional co-activator, p300. *EMBO Journal* **15**, 2236-2248
(1996).

25 Green, M. R. Eukaryotic transcription activation: right on target. *Mol Cell* **18**, 399-402,
doi:10.1016/j.molcel.2005.04.017 (2005).

26 Malik, S. & Roeder, R. G. The metazoan Mediator co-activator complex as an integrative hub for
transcriptional regulation. *Nat Rev Genet* **11**, 761-772, doi:10.1038/nrg2901 (2010).

27 Allen, B. L. & Taatjes, D. J. The Mediator complex: a central integrator of transcription. *Nat Rev*
*Mol Cell Biol* **16**, 155-166, doi:10.1038/nrm3951 (2015).

28 Merika, M., Williams, A. J., Chen, G., Collins, T. & Thanos, D. Recruitment of CBP/p300 by the
IFN beta enhanceosome is required for synergistic activation of transcription. *Mol Cell* **1**, 277-
287 (1998).

29 Reiter, F., Wienerroither, S. & Stark, A. Combinatorial function of transcription factors and
cofactors. *Curr Opin Genet Dev* **43**, 73-81, doi:10.1016/j.gde.2016.12.007 (2017).

30 Kamei, Y. *et al.* A CBP integrator complex mediates transcriptional activation and AP-1
inhibition by nuclear receptors. *Cell* **85**, 403-414 (1996).

31 Goodman, R. H. & Smolik, S. CBP/p300 in cell growth, transformation, and development. *Genes*
*Dev* **14**, 1553-1577 (2000).

32 Narita, T. *et al.* Enhancers are activated by p300/CBP activity-dependent PIC assembly, RNAPII
recruitment, and pause release. *Mol Cell* **81**, 2166-2182 e2166, doi:10.1016/j.molcel.2021.03.008
(2021).

33 Cabianca, D. S. *et al.* Active chromatin marks drive spatial sequestration of heterochromatin in C.
elegans nuclei. *Nature* **569**, 734-739, doi:10.1038/s41586-019-1243-y (2019).

34 Gillespie, M. A. *et al.* Absolute Quantification of Transcription Factors Reveals Principles of
Gene Regulation in Erythropoiesis. *Mol Cell* **78**, 960-974 e911, doi:10.1016/j.molcel.2020.03.031
(2020).

35 Weinert, B. T. *et al.* Time-Resolved Analysis Reveals Rapid Dynamics and Broad Scope of the
CBP/p300 Acetylome. *Cell* **174**, 231-244 e212, doi:10.1016/j.cell.2018.04.033 (2018).

36 Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase Separation
Model for Transcriptional Control. *Cell* **169**, 13-23, doi:10.1016/j.cell.2017.02.007 (2017).

37 Boija, A. *et al.* Transcription Factors Activate Genes through the Phase-Separation Capacity of
Their Activation Domains. *Cell* **175**, 1842-1855 e1816, doi:10.1016/j.cell.2018.10.042 (2018).

38 Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions that control
gene transcription. *Science* **361**, doi:10.1126/science.aar2555 (2018).

39 Sabari, B. R. *et al.* Coactivator condensation at super-enhancers links phase separation and gene
control. *Science* **361**, doi:10.1126/science.aar3958 (2018).

40 Cho, W. K. *et al.* Mediator and RNA polymerase II clusters associate in transcription-dependent
condensates. *Science* **361**, 412-415, doi:10.1126/science.aar4199 (2018).

41 Zamudio, A. V. *et al.* Mediator Condensates Localize Signaling Factors to Key Cell Identity
Genes. *Mol Cell* **76**, 753-766 e756, doi:10.1016/j.molcel.2019.08.016 (2019).

42 Guo, Y. E. *et al.* Pol II phosphorylation regulates a switch between transcriptional and splicing
condensates. *Nature* **572**, 543-+, doi:10.1038/s41586-019-1464-0 (2019).

43 Shrinivas, K. *et al.* Enhancer Features that Drive Formation of Transcriptional Condensates. *Mol*
*Cell* **75**, 549-561 e547, doi:10.1016/j.molcel.2019.07.009 (2019).

44  Delvecchio, M., Gaucher, J., Aguilar-Guerreri, C., Ortega, E. & Panne, D. Structure of the p300
catalytic core and implications for chromatin targeting and HAT regulation. *Nature Structural*
*and Molecular Biology* **21**, doi:10.1038/nsmb.2642 (2013).
45  Dyson, H. J. & Wright, P. E. Role of Intrinsic Protein Disorder in the Function and Interactions of
the Transcriptional Coactivators CREB-binding Protein (CBP) and p300. *J Biol Chem* **291**, 6714-
6722, doi:10.1074/jbc.R115.692020 (2016).
46  Dancy, B. M. & Cole, P. A. Protein lysine acetylation by p300/CBP. *Chem Rev* **115**, 2419-2452,
doi:10.1021/cr500452k (2015).
47  Ortega, E. *et al.* Transcription factor dimerization activates the p300 acetyltransferase. *Nature*
**562**, 538-544, doi:10.1038/s41586-018-0621-1 (2018).
48  Zhang, Y. *et al.* Nuclear condensates of p300 formed though the structured catalytic core can act
as a storage pool of p300 with reduced HAT activity. *Nat Commun* **12**, 4618,
doi:10.1038/s41467-021-24950-8 (2021).
49  Ma, L. *et al.* Co-condensation between transcription factor and coactivator p300 modulates
transcriptional bursting kinetics. *Mol Cell*, doi:10.1016/j.molcel.2021.01.031 (2021).
50  Pivot-Pajot, C. *et al.* Acetylation-dependent chromatin reorganization by BRDT, a testis-specific
bromodomain-containing protein. *Mol Cell Biol* **23**, 5354-5365, doi:10.1128/MCB.23.15.5354-
5365.2003 (2003).
51  Gaucher, J. *et al.* Bromodomain-dependent stage-specific male genome programming by Brdt.
*EMBO J* **31**, 3809-3820, doi:10.1038/emboj.2012.233 (2012).
52  Goudarzi, A. *et al.* Dynamic Competing Histone H4 K5K8 Acetylation and Butyrylation Are
Hallmarks of Highly Active Gene Promoters. *Mol Cell* **62**, 169-180,
doi:10.1016/j.molcel.2016.03.014 (2016).
53  Shiota, H. *et al.* Nut Directs p300-Dependent, Genome-Wide H4 Hyperacetylation in Male Germ
Cells. *Cell Rep* **24**, 3477-3487 e3476, doi:10.1016/j.celrep.2018.08.069 (2018).
54  Zhao, R., Nakamura, T., Fu, Y., Lazar, Z. & Spector, D. L. Gene bookmarking accelerates the
kinetics of post-mitotic transcriptional re-activation. *Nature Cell Biology* **13**, 1295-U1239,
doi:10.1038/ncb2341 (2011).
55  Devaiah, B. N. *et al.* BRD4 is a histone acetyltransferase that evicts nucleosomes from chromatin.
*Nat Struct Mol Biol* **23**, 540-548, doi:10.1038/nsmb.3228 (2016).
56  Wang, R., Li, Q., Helfer, C. M., Jiao, J. & You, J. Bromodomain protein Brd4 associated with
acetylated chromatin is important for maintenance of higher-order chromatin structure. *J Biol*
*Chem* **287**, 10738-10752, doi:10.1074/jbc.M111.323493 (2012).
57  Ladurner, A. G., Inouye, C., Jain, R. & Tjian, R. Bromodomains mediate an acetyl-histone
encoded antisilencing function at heterochromatin boundaries. *Molecular Cell* **11**, 365-376,
doi:Doi 10.1016/S1097-2765(03)00035-2 (2003).
58  Gibson, B. A. *et al.* Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell*
**179**, 470-484 e421, doi:10.1016/j.cell.2019.08.037 (2019).
59  Shin, Y. *et al.* Liquid Nuclear Condensates Mechanically Sense and Restructure the Genome.
*Cell* **175**, 1481-1491 e1413, doi:10.1016/j.cell.2018.10.057 (2018).
60  Han, X. *et al.* Roles of the BRD4 short isoform in phase separation and active gene transcription.
*Nat Struct Mol Biol* **27**, 333-341, doi:10.1038/s41594-020-0394-8 (2020).
61  Li, J. *et al.* Single-gene imaging links genome topology, promoter-enhancer communication and
transcription control. *Nat Struct Mol Biol* **27**, 1032-1040, doi:10.1038/s41594-020-0493-6 (2020).
62  Eagen, K. P. & French, C. A. Supercharging BRD4 with NUT in carcinoma. *Oncogene* **40**, 1396-
1408, doi:10.1038/s41388-020-01625-0 (2021).
63  Reynoird, N. *et al.* Oncogenesis by sequestration of CBP/p300 in transcriptionally inactive
hyperacetylated chromatin domains. *EMBO J* **29**, 2943-2952, doi:emboj2010176 [pii]
10.1038/emboj.2010.176 (2010).

64    Yan, J., Diaz, J., Jiao, J., Wang, R. & You, J. Perturbation of BRD4 protein function by BRD4-
NUT protein abrogates cellular differentiation in NUT midline carcinoma. *J Biol Chem* **286**,
27663-27675, doi:10.1074/jbc.M111.246975 (2011).
65    Alekseyenko, A. A. *et al.* The oncogenic BRD4-NUT chromatin regulator drives aberrant
transcription within large topological domains. *Genes Dev* **29**, 1507-1523,
doi:10.1101/gad.267583.115 (2015).
66    Alekseyenko, A. A. *et al.* Ectopic protein interactions within BRD4-chromatin complexes drive
oncogenic megadomain formation in NUT midline carcinoma. *Proc Natl Acad Sci U S A* **114**,
E4184-E4192, doi:10.1073/pnas.1702086114 (2017).
67    Wang, R. & You, J. Mechanistic analysis of the role of bromodomain-containing protein 4
(BRD4) in BRD4-NUT oncoprotein-induced transcriptional activation. *J Biol Chem* **290**, 2744-
2758, doi:10.1074/jbc.M114.600759 (2015).
68    Rosencrance, C. D. *et al.* Chromatin Hyperacetylation Impacts Chromosome Folding by Forming
a Nuclear Subcompartment. *Mol Cell* **78**, 112-126 e112, doi:10.1016/j.molcel.2020.03.018
(2020).
69    Staller, M. V. *et al.* A High-Throughput Mutational Scan of an Intrinsically Disordered Acidic
Transcriptional Activation Domain. *Cell Syst* **6**, 444-455 e446, doi:10.1016/j.cels.2018.01.015
(2018).
70    Dominguez, C., Boelens, R. & Bonvin, A. M. HADDOCK: a protein-protein docking approach
based on biochemical or biophysical information. *Journal of the American Chemical Society* **125**,
1731-1737, doi:10.1021/ja026939x (2003).
71    Basu, S. *et al.* Unblending of Transcriptional Condensates in Human Repeat Expansion Disease.
*Cell* **181**, 1062-1079 e1030, doi:10.1016/j.cell.2020.04.018 (2020).
72    Plevin, M. J., Mills, M. M. & Ikura, M. The LxxLL motif: a multifunctional binding sequence in
transcriptional regulation. *Trends Biochem Sci* **30**, 66-69, doi:10.1016/j.tibs.2004.12.001 (2005).
73    Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*,
doi:10.1101/2021.10.04.463034 (2021).
74    Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-
589, doi:10.1038/s41586-021-03819-2 (2021).
75    Krois, A. S., Ferreon, J. C., Martinez-Yamout, M. A., Dyson, H. J. & Wright, P. E. Recognition
of the disordered p53 transactivation domain by the transcriptional adapter zinc finger domains of
CREB-binding protein. *Proc Natl Acad Sci U S A* **113**, E1853-1862,
doi:10.1073/pnas.1602487113 (2016).
76    Zhang, Y. *et al.* The ZZ domain of p300 mediates specificity of the adjacent HAT domain for
histone H3. *Nat Struct Mol Biol* **25**, 841-849, doi:10.1038/s41594-018-0114-9 (2018).
77    Sheahan, T. W., Major, V., Webb, K. M., Bryan, E. & Voigt, P. The TAZ2 domain of CBP/p300
directs acetylation towards H3K27 within chromatin. *bioRxiv* (2020).
78    Ghandi, M. *et al.* Next-generation characterization of the Cancer Cell Line Encyclopedia. *Nature*
**569**, 503-508, doi:10.1038/s41586-019-1186-3 (2019).
79    Kumar, M. *et al.* Inhibition of histone acetyltransferase function radiosensitizes CREBBP/EP300
mutants via repression of homologous recombination, potentially targeting a gain of function. *Nat*
*Commun* **12**, 6340, doi:10.1038/s41467-021-26570-8 (2021).
80    Miller, T. C. *et al.* A bromodomain-DNA interaction facilitates acetylation-dependent bivalent
nucleosome recognition by the BET protein BRDT. *Nat Commun* **7**, 13855,
doi:10.1038/ncomms13855 (2016).
81    Wu, S. Y., Lee, A. Y., Lai, H. T., Zhang, H. & Chiang, C. M. Phospho switch triggers Brd4
chromatin binding and activator recruitment for gene-specific targeting. *Mol Cell* **49**, 843-857,
doi:10.1016/j.molcel.2012.12.006 (2013).
82    Slaughter, M. J. *et al.* HDAC inhibition results in widespread alteration of the histone acetylation
landscape and BRD4 targeting to gene bodies. *Cell Rep* **34**, 108638,
doi:10.1016/j.celrep.2020.108638 (2021).

83    Hammitzsch, A. *et al.* CBP30, a selective CBP/p300 bromodomain inhibitor, suppresses human
Th17 responses. *Proc Natl Acad Sci U S A* **112**, 10768-10773, doi:10.1073/pnas.1501956112
(2015).
84    Shogren-Knaak, M. *et al.* Histone H4-K16 acetylation controls chromatin structure and protein
interactions. *Science* **311**, 844-847, doi:10.1126/science.1124000 (2006).
85    Strickfaden, H. *et al.* Condensed Chromatin Behaves like a Solid on the Mesoscale In Vitro and
in Living Cells. *Cell* **183**, 1772-1784 e1713, doi:10.1016/j.cell.2020.11.027 (2020).
86    Dhalluin, C. *et al.* Structure and ligand of a histone acetyltransferase bromodomain. *Nature* **399**,
491-496, doi:10.1038/20974 (1999).
87    Strahl, B. D. & Allis, C. D. The language of covalent histone modifications. *Nature* **403**, 41-45,
doi:10.1038/47412 (2000).
88    Eberharter, A. & Becker, P. B. Histone acetylation: a switch between repressive and permissive
chromatin. Second in review series on chromatin dynamics. *EMBO Rep* **3**, 224-229,
doi:10.1093/embo-reports/kvf053 (2002).
89    Thompson, P. R. *et al.* Regulation of the p300 HAT domain via a novel activation loop. *Nat*
*Struct Mol Biol* **11**, 308-315, doi:10.1038/nsmb740
nsmb740 [pii] (2004).
90    Karanam, B., Jiang, L., Wang, L., Kelleher, N. L. & Cole, P. A. Kinetic and mass spectrometric
analysis of p300 histone acetyltransferase domain autoacetylation. *J Biol Chem* **281**, 40292-
40301, doi:10.1074/jbc.M608813200 (2006).
91    Hogg, S. J. *et al.* Targeting histone acetylation dynamics and oncogenic transcription by catalytic
P300/CBP inhibition. *Mol Cell* **81**, 2183-2200 e2113, doi:10.1016/j.molcel.2021.04.015 (2021).
92    Black, J. C., Mosley, A., Kitada, T., Washburn, M. & Carey, M. The SIRT2 deacetylase regulates
autoacetylation of p300. *Mol Cell* **32**, 449-455, doi:S1097-2765(08)00696-5 [pii]
10.1016/j.molcel.2008.09.018 (2008).
93    Pelka, P. *et al.* Transcriptional control by adenovirus E1A conserved region 3 via p300/CBP.
*Nucleic Acids Res* **37**, 1095-1106, doi:10.1093/nar/gkn1057 (2009).
94    Hsu, E., Pennella, M. A., Zemke, N. R., Eng, C. & Berk, A. J. Adenovirus E1A Activation
Domain Regulates H3 Acetylation Affecting Varied Steps in Transcription at Different Viral
Promoters. *J Virol* **92**, doi:10.1128/JVI.00805-18 (2018).
95    Kraus, W. L., Manning, E. T. & Kadonaga, J. T. Biochemical analysis of distinct activation
functions in p300 that enhance transcription initiation with chromatin templates. *Molecular and*
*Cellular Biology* **19**, 8123-8135 (1999).
96    Hansson, M. L. *et al.* The transcriptional coactivator MAML1 regulates p300 autoacetylation and
HAT activity. *Nucleic Acids Res* **37**, 2996-3006, doi:gkp163 [pii]
10.1093/nar/gkp163 (2009).
97    Ferrari, R. *et al.* Epigenetic reprogramming by adenovirus e1a. *Science* **321**, 1086-1088,
doi:10.1126/science.1155546 (2008).
98    Horwitz, G. A. *et al.* Adenovirus small e1a alters global patterns of histone modification. *Science*
**321**, 1084-1085, doi:10.1126/science.1155544 (2008).
99    Ferrari, R. *et al.* Adenovirus small E1A employs the lysine acetylases p300/CBP and tumor
suppressor Rb to repress select host genes and promote productive virus infection. *Cell Host*
*Microbe* **16**, 663-676, doi:10.1016/j.chom.2014.10.004 (2014).
100    Attar, N. & Kurdistani, S. K. Exploitation of EP300 and CREBBP Lysine Acetyltransferases by
Cancer. *Cold Spring Harb Perspect Med* **7**, doi:10.1101/cshperspect.a026534 (2017).
101    Wojciak, J. M., Martinez-Yamout, M. A., Dyson, H. J. & Wright, P. E. Structural basis for
recruitment of CBP/p300 coactivators by STAT1 and STAT2 transactivation domains. *EMBO J*
**28**, 948-958, doi:emboj200930 [pii]
10.1038/emboj.2009.30 (2009).
102    Ferreon, J. C., Martinez-Yamout, M. A., Dyson, H. J. & Wright, P. E. Structural basis for
subversion of cellular control mechanisms by the adenoviral E1A oncoprotein. *Proceedings of the*

840            *National Academy of Sciences of the United States of America* **106**, 13260-13265,
841            doi:10.1073/pnas.0906770106 (2009).
842  103  Lee, C. W., Martinez-Yamout, M. A., Dyson, H. J. & Wright, P. E. Structure of the p53
843            transactivation domain in complex with the nuclear receptor coactivator binding domain of CREB
844            binding protein. *Biochemistry* **49**, 9964-9971, doi:10.1021/bi1012996 (2010).
845  104  Mukherjee, S. P. *et al.* Analysis of the RelA:CBP/p300 interaction reveals its involvement in NF-
846            kappaB-driven transcription. *PLoS Biol* **11**, e1001647, doi:10.1371/journal.pbio.1001647 (2013).
847  105  Miller Jenkins, L. M. *et al.* Characterization of the p300 Taz2-p53 TAD2 complex and
848            comparison with the p300 Taz2-p53 TAD1 complex. *Biochemistry* **54**, 2001-2010,
849            doi:10.1021/acs.biochem.5b00044 (2015).
850  106  He, J. *et al.* Structure of p300 bound to MEF2 on DNA reveals a mechanism of enhanceosome
851            assembly. *Nucleic Acids Res* **39**, 4464-4474, doi:10.1093/nar/gkr030 (2011).
852  107  Feng, H. *et al.* Structural basis for p300 Taz2-p53 TAD1 binding and modulation by
853            phosphorylation. *Structure* **17**, 202-210, doi:10.1016/j.str.2008.12.009 (2009).
854  108  Marceau, A. H. *et al.* An order-to-disorder structural switch activates the FoxM1 transcription
855            factor. *Elife* **8**, doi:10.7554/eLife.46131 (2019).
856  109  Dames, S. A., Martinez-Yamout, M., De Guzman, R. N., Dyson, H. J. & Wright, P. E. Structural
857            basis for Hif-1 alpha /CBP recognition in the cellular hypoxic response. *Proc Natl Acad Sci U S A*
858            **99**, 5271-5276, doi:10.1073/pnas.082121399 (2002).
859  110  Wang, F. *et al.* Structures of KIX domain of CBP in complex with two FOXO3a transactivation
860            domains reveal promiscuity and plasticity in coactivator recruitment. *Proc Natl Acad Sci U S A*
861            **109**, 6078-6083, doi:10.1073/pnas.1119073109 (2012).
862  111  Chan, H. M. & La Thangue, N. B. p300/CBP proteins: HATs for transcriptional bridges and
863            scaffolds. *J Cell Sci* **114**, 2363-2373 (2001).
864  112  Panne, D., Maniatis, T. & Harrison, S. C. An atomic model of the interferon-beta enhanceosome.
865            *Cell* **129**, 1111-1123, doi:S0092-8674(07)00656-3 [pii]
866            10.1016/j.cell.2007.05.019 (2007).
867  113  Panne, D. The enhanceosome. *Curr Opin Struct Biol* **18**, 236-242, doi:S0959-440X(07)00202-3
868            [pii]
869            10.1016/j.sbi.2007.12.002 (2008).
870  114  Wang, Z. *et al.* Genome-wide mapping of HATs and HDACs reveals distinct functions in active
871            and inactive genes. *Cell* **138**, 1019-1031, doi:S0092-8674(09)00841-1 [pii]
872            10.1016/j.cell.2009.06.049 (2009).
873  115  Whyte, W. A. *et al.* Enhancer decommissioning by LSD1 during embryonic stem cell
874            differentiation. *Nature* **482**, 221-225, doi:10.1038/nature10805 (2012).
875  116  Gryder, B. E. *et al.* Histone hyperacetylation disrupts core gene regulatory architecture in
876            rhabdomyosarcoma. *Nat Genet* **51**, 1714-1722, doi:10.1038/s41588-019-0534-4 (2019).
877  117  Gryder, B. E. *et al.* Chemical genomics reveals histone deacetylases are required for core
878            regulatory transcription. *Nat Commun* **10**, 3004, doi:10.1038/s41467-019-11046-7 (2019).
879  118  Fukaya, T., Lim, B. & Levine, M. Enhancer Control of Transcriptional Bursting. *Cell* **166**, 358-
880            368, doi:10.1016/j.cell.2016.05.025 (2016).
881  119  Li, J. *et al.* Single-Molecule Nanoscopy Elucidates RNA Polymerase II Transcription at Single
882            Genes in Live Cells. *Cell* **178**, 491-506 e428, doi:10.1016/j.cell.2019.05.029 (2019).
883  120  Schwartz, B. E. *et al.* Differentiation of NUT midline carcinoma by epigenomic reprogramming.
884            *Cancer Research* **71**, 2686-2696, doi:10.1158/0008-5472.CAN-10-3513 (2011).
885  121  Rundlett, S. E. *et al.* HDA1 and RPD3 are members of distinct yeast histone deacetylase
886            complexes that regulate silencing and transcription. *Proc Natl Acad Sci U S A* **93**, 14503-14508,
887            doi:10.1073/pnas.93.25.14503 (1996).
888  122  Kidder, B. L. & Palmer, S. HDAC1 regulates pluripotency and lineage specific transcriptional
889            networks in embryonic and trophoblast stem cells. *Nucleic Acids Res* **40**, 2925-2939,
890            doi:10.1093/nar/gkr1151 (2012).

123  Vogelauer, M., Wu, J., Suka, N. & Grunstein, M. Global histone acetylation and deacetylation in
yeast. *Nature* **408**, 495-498, doi:10.1038/35044127 (2000).
124  Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and predicts
developmental state. *Proc Natl Acad Sci U S A* **107**, 21931-21936, doi:10.1073/pnas.1016071107
(2010).
125  Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental enhancers in
humans. *Nature*, doi:nature09692 [pii]
10.1038/nature09692 (2010).
126  Zhang, T., Zhang, Z., Dong, Q., Xiong, J. & Zhu, B. Histone H3K27 acetylation is dispensable
for enhancer activity in mouse embryonic stem cells. *Genome Biol* **21**, 45, doi:10.1186/s13059-
020-01957-w (2020).
127  Filippakopoulos, P. & Knapp, S. The bromodomain interaction module. *FEBS Lett* **586**, 2692-
2704, doi:10.1016/j.febslet.2012.04.045 (2012).
128  Carey, M., Lin, Y. S., Green, M. R. & Ptashne, M. A mechanism for synergistic activation of a
mammalian gene by GAL4 derivatives. *Nature* **345**, 361-364 (1990).
129  Gill, G. & Ptashne, M. Negative effect of the transcriptional activator GAL4. *Nature* **334**, 721-
724, doi:10.1038/334721a0 (1988).
130  Meyer, M. E. *et al.* Steroid hormone receptors compete for factors that mediate their enhancer
function. *Cell* **57**, 433-442, doi:10.1016/0092-8674(89)90918-5 (1989).
131  Triezenberg, S. J., Kingsbury, R. C. & McKnight, S. L. Functional dissection of VP16, the trans-
activator of herpes simplex virus immediate early gene expression. *Genes Dev* **2**, 718-729,
doi:10.1101/gad.2.6.718 (1988).
132  Hottiger, M. O., Felzien, L. K. & Nabel, G. J. Modulation of cytokine-induced HIV gene
expression by competitive binding of transcription factors to the coactivator p300. *EMBO J* **17**,
3124-3134, doi:10.1093/emboj/17.11.3124 (1998).
133  De Bosscher, K., Vanden Berghe, W. & Haegeman, G. The interplay between the glucocorticoid
receptor and nuclear factor-kappaB or activator protein-1: molecular mechanisms for gene
repression. *Endocr Rev* **24**, 488-522, doi:10.1210/er.2002-0006 (2003).
134  Portuguez, A. S. *et al.* Gene activation and repression by the glucocorticoid receptor are mediated
by sequestering Ep300 and two modes of chromatin binding. *bioRxiv* (2019).
135  Guertin, M. J., Zhang, X., Coonrod, S. A. & Hager, G. L. Transient estrogen receptor binding and
p300 redistribution support a squelching mechanism for estradiol-repressed genes. *Mol
Endocrinol* **28**, 1522-1533, doi:10.1210/me.2014-1130 (2014).
136  Halford, S. E. An end to 40 years of mistakes in DNA-protein association kinetics? *Biochem Soc
Trans* **37**, 343-348, doi:10.1042/BST0370343 (2009).
137  Mirny, L. *et al.* How a protein searches for its site on DNA: the mechanism of facilitated
diffusion. *Journal of Physics A: Mathematical and Theoretical* **42**, 434013, doi:10.1088/1751-
8113/42/43/434013 (2009).
138  Loffreda, A. *et al.* Live-cell p53 single-molecule binding is modulated by C-terminal acetylation
and correlates with transcriptional activity. *Nat Commun* **8**, 313, doi:10.1038/s41467-017-00398-
7 (2017).
139  Luo, J. *et al.* Acetylation of p53 augments its site-specific DNA binding both in vitro and in vivo.
*Proc Natl Acad Sci U S A* **101**, 2259-2264, doi:10.1073/pnas.0308762101 (2004).
140  Lickwar, C. R., Mueller, F., Hanlon, S. E., McNally, J. G. & Lieb, J. D. Genome-wide protein-
DNA binding dynamics suggest a molecular clutch for transcription factor function. *Nature* **484**,
251-255, doi:10.1038/nature10985 (2012).
141  Gurdon, J. B., Javed, K., Vodnala, M. & Garrett, N. Long-term association of a transcription
factor with its chromatin binding site can stabilize gene expression and cell fate commitment.
*Proc Natl Acad Sci U S A* **117**, 15075-15084, doi:10.1073/pnas.2000467117 (2020).

142 Kribelbauer, J. F., Rastogi, C., Bussemaker, H. J. & Mann, R. S. Low-Affinity Binding Sites and
the Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol* **35**, 357-379,
doi:10.1146/annurev-cellbio-100617-062719 (2019).
143 Li, X. Y. *et al.* Transcription factors bind thousands of active and inactive regions in the
Drosophila blastoderm. *PLoS Biol* **6**, e27, doi:10.1371/journal.pbio.0060027 (2008).
144 Fisher, W. W. *et al.* DNA regions bound at low occupancy by transcription factors do not drive
patterned reporter gene expression in Drosophila. *Proc Natl Acad Sci U S A* **109**, 21330-21335,
doi:10.1073/pnas.1209589110 (2012).
145 Boija, A., Klein, I. A. & Young, R. A. Biomolecular Condensates and Cancer. *Cancer Cell* **39**,
174-192, doi:10.1016/j.ccell.2020.12.003 (2021).
146 Ahn, J. H. *et al.* Phase separation drives aberrant chromatin looping and cancer development.
*Nature* **595**, 591-595, doi:10.1038/s41586-021-03662-5 (2021).
147 Malvezzi, F. *et al.* Phosphorylation-dependent BRD4 dimerization and implications for
therapeutic inhibition of BET family proteins. *Commun Biol* **4**, 1273, doi:10.1038/s42003-021-
02750-6 (2021).
148 Rowley, M. J. & Corces, V. G. Organizational principles of 3D genome architecture. *Nat Rev*
*Genet* **19**, 789-800, doi:10.1038/s41576-018-0060-8 (2018).
149 Aparicio, O. M. & Gottschling, D. E. Overcoming telomeric silencing: a trans-activator competes
to establish gene expression in a cell cycle-dependent way. *Genes Dev* **8**, 1133-1146,
doi:10.1101/gad.8.10.1133 (1994).
150 Tumbar, T., Belmont, A. S. & Sudlow, G. P. Targeting the VP16 acidic activation domain
induces chances in large-scale chromatin structure and nuclear positioning. *Mol Biol Cell* **10**,
281a-281a (1999).
151 Ahmad, K. & Henikoff, S. Modulation of a transcription factor counteracts heterochromatic gene
silencing in Drosophila. *Cell* **104**, 839-847, doi:10.1016/s0092-8674(01)00281-1 (2001).
152 Jeppesen, P. & Turner, B. M. The inactive X chromosome in female mammals is distinguished by
a lack of histone H4 acetylation, a cytogenetic marker for gene expression. *Cell* **74**, 281-289,
doi:10.1016/0092-8674(93)90419-q (1993).
153 Spilianakis, C. G., Lalioti, M. D., Town, T., Lee, G. R. & Flavell, R. A. Interchromosomal
associations between alternatively expressed loci. *Nature* **435**, 637-645, doi:10.1038/nature03574
(2005).
154 Apostolou, E. & Thanos, D. Virus Infection Induces NF-kappaB-dependent interchromosomal
associations mediating monoallelic IFN-beta gene expression. *Cell* **134**, 85-96,
doi:10.1016/j.cell.2008.05.052 (2008).
155 Monahan, K., Horta, A. & Lomvardas, S. LHX2- and LDB1-mediated trans interactions regulate
olfactory receptor choice. *Nature* **565**, 448-453, doi:10.1038/s41586-018-0845-0 (2019).
156 Zylicz, J. J. *et al.* The Implication of Early Chromatin Changes in X Chromosome Inactivation.
*Cell* **176**, 182-197 e123, doi:10.1016/j.cell.2018.11.041 (2019).
157 Migl, D. *et al.* Cryoelectron Microscopy Structure of a Yeast Centromeric Nucleosome at 2.7 A
Resolution. *Structure* **28**, 363-370 e363, doi:10.1016/j.str.2019.12.002 (2020).
158 Franke, D. *et al.* ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering from
macromolecular solutions. *J Appl Crystallogr* **50**, 1212-1225, doi:10.1107/S1600576717007786
(2017).
159 Petoukhov, M. V. *et al.* New developments in the program package for small-angle scattering
data analysis. *J Appl Crystallogr* **45**, 342-350, doi:10.1107/S0021889812007662 (2012).
160 Dorigo, B., Schalch, T., Bystricky, K. & Richmond, T. J. Chromatin fiber folding: requirement
for the histone H4 N-terminal tail. *J Mol Biol* **327**, 85-96, doi:10.1016/s0022-2836(03)00025-1
(2003).
161 Luger, K., Rechsteiner, T. J. & Richmond, T. J. Preparation of nucleosome core particle from
recombinant histones. *Methods Enzymol* **304**, 3-19, doi:S0076-6879(99)04003-3 [pii] (1999).

162  Schumann, F. H. *et al.* Combined chemical shift changes and amino acid specific chemical shift
mapping of protein-protein interactions. *J Biomol NMR* **39**, 275-289, doi:10.1007/s10858-007-
9197-z (2007).
163  Maciejewski, M. W. *et al.* NMRbox: A Resource for Biomolecular NMR Computation. *Biophys J*
**112**, 1529-1534, doi:10.1016/j.bpj.2017.03.011 (2017).
164  Ozenne, V. *et al.* Flexible-meccano: a tool for the generation of explicit ensemble descriptions of
intrinsically disordered proteins and their associated experimental observables. *Bioinformatics* **28**,
1463-1470, doi:10.1093/bioinformatics/bts172 (2012).
165  Shen, Y. & Bax, A. Protein backbone and sidechain torsion angles predicted from NMR chemical
shifts using artificial neural networks. *J Biomol NMR* **56**, 227-241, doi:10.1007/s10858-013-
9741-y (2013).
166  Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193 (2017).
167  Zhang, K. Gctf: Real-time CTF determination and correction. *Journal of Structural Biology* **193**,
1-12, doi:10.1016/j.jsb.2015.11.003 (2016).
168  Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination in
RELION-3. *Elife* **7**, doi:10.7554/eLife.42166 (2018).

## Methods

**Constructs.** cDNA encoding residues for NUT-F1c (347-588), NUT1 (347-446), NUT2 (427-527), NUT3 (508-588), NUT4 (347-480), NUT5 (347-514), NUT6 (351-470), NUT7 (448-492) (wild type and mutants), NUT8 (424-492) (wild type and mutants), NUT9 (395-482) and p53 (1-61) were cloned into the *Nco*I/*Xho*I restriction sites of vectors pETM-30 or pETM11 (EMBL) with a tobacco etch virus (TEV)-cleavable N-terminal His$_6$ glutathione S-transferase (His$_6$-GST) tag for pETM30 or an N-terminal His$_6$-tag for pETM11. The p300 TAZ2 domain constructs (TAZ2 1723-1812, TAZ2 1723-1822, TAZ2 1723-1836) were codon optimized and cloned into the *Nco*I/*Bam*HI restriction sites of pETM-11 (EMBL) with a TEV-cleavable N-terminal His$_6$-tag. Cysteines which do not participate in zinc coordination were mutated to Alanine or Serine (C1738, C1746, C1769, C1770). The CH3 domain of p300 (1660-1815) was cloned using *Eco*RI/*Xho*I restriction sites in pGEX-5X1 vector with a TEV-cleavable N-terminal glutathione S-transferase (GST) tag. p300s (324-2094) and p300 core-CH3 (1043-1828 and 1043-1817) were produced as described previously[44]. The BRD4 constructs: GFP-BD1-BD2, GFP-BD1-BD2 N140A/N433A, GFP-BD1-BD2 N140A and GFP-BD1-BD2 N433A were cloned into the *Nco*I/*Xho*I restriction sites of pETM11. The vectors expressing the GFP-(Bromo)5 and the GFP-(Bromo)5 N140A were gifts from Michael Rosen lab (University of Texas Southwestern). The co-expression vector encoding the *S. cerevisiae* histones was a gift from Stephen Harrison lab (Harvard Medical school). The vector expressing the E1A protein was a gift from Andy Turnell lab (University of Birmingham). For cell-based assays, all the GFP-tagged BRD4-NUT variants, RFP-tagged NUT-F1c, NUT1, NUT2, NUT3 and NUT4 and RFP tagged p300 were cloned into pEGFP-C1 vector (Clontech), HA-tagged p300 constructs (WT, Δ1813-1838) were cloned into pcDNA3.1. All constructs were confirmed by DNA sequencing.

**Expression and purification.** Expression and purification of Flag-tagged p300s and GST-tagged p300 core-CH3 constructs was done as reported previously[44,47]. All p300 TAZ2 constructs were produced as inclusion bodies in *Escherichia coli* (*E. coli)* BL21 (DE3). Cells were grown to mid-log phase (OD600 ~ 0.6) at 37°C in Luria Broth containing 50 µg/ml kamamycin and 200 µM ZnSO$_4$. Protein expression was induced by addition of 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) at 18°C for 16h. Uniformly $^{15}$N- and $^{15}$N/$^{13}$C-labelled samples were produced from cells grown in M9 minimal medium containing 50 µg/ml Kanamycin, 200 µM ZnSO$_4$ and 1 g/l of $^{15}$NH$_4$Cl and 2g/l $^{13}$C glucose. Cells were harvested by centrifugation for 20 mins at 5000 g in a SLC-6000 rotor (Sorvall) and resuspended in 50 ml/L culture

lysis buffer 1 (20 mM Tris-HCl pH 7.5, 300 mM NaCl, 5 µM ZnSO$_4$, 1 mM Dithiothreitol (DTT)),
supplemented with EDTA-free protease inhibitors (Roche), and 10 µl of benzonase nuclease (Sigma-
Aldrich). Cells were sonicated and centrifuged in a SS-34 rotor (Sorvall) for 1h at 12000 rpm. The pellet
was then washed three times in the lysis buffer and the final pellet resuspended in buffer 2 (20 mM Tris
pH 7.5, 300 mM NaCl, 1 mM Dithiothreitol (DTT), 6 M urea) for one hour at 4 °C on a rotating device.
The protein solution was centrifuged as above and the soluble fraction applied to Ni$^{2+}$-charged IMAC
Sepharose 6 FF (Cytiva) beads preequilibrated with buffer 2 containing 20 mM Imidazole. The column
was washed with 20 column volumes (CVs) of buffer 2 containing 20 mM Imidazole and the bound
protein eluted with the buffer 2 containing 300 mM Imidazole. The protein was refolded by dialysis
overnight against buffer 3 (20 mM Tris pH 7.0, 100 mM NaCl, 200 µM ZnSO$_4$ and 2 mM DTT).
1050
The refolded TAZ2 was incubated with His-tagged TEV protease (1:100 w/w) for 14-16 h at 4 °C.
Subtractive Ni-NTA chromatography was used to remove the residual His-tag and TEV protease. The
cleaved TAZ2 was further purified on a 5ml HiTrap SP HP cation exchange column (Cytiva)
preequilibrated in buffer 4 (20 mM Tris pH 7.0, 100 mM NaCl, 5 µM ZnSO$_4$ and 1 mM Dithiothreitol
(DTT)) and eluted using a 20 CV NaCl gradient. The peak containing TAZ2 was further purified using a
HiLoad 16/60 Superdex 75 column (Cytiva) preequilibrated with buffer 5 (20 mM Tris pH 7.0, 300 mM
NaCl, 5 µM ZnSO$_4$ and 1 mM DTT). The purified protein was concentrated using a prewashed Amicon
Ultra-10 concentrator (molecular weight cut off 10 kDa; EMD Millipore), flash frozen in liquid N2 and
stored in -80°C.
1060
Unlabeled, $^{15}$N- and $^{15}$N/$^{13}$C-labelled NUT fragments were produced as insoluble inclusion bodies in *E.*
*coli* and purified as described above. After Ni-NTA purification, the protein was dialyzed and refolded
overnight at 4°C against 1L buffer 5 (20 mM Tris pH 7.0, 300 mM NaCl, 1 mM DTT and 10% glycerol).
Refolded His-GST-NUT fusion proteins were incubated with His-tagged TEV protease (1:100 w/w) for
14-16 h at 4 °C. Subtractive Ni-NTA chromatography was used to remove the residual His-GST-tag and
TEV protease. The cleaved NUT proteins were applied to a 5ml HiTrap Q HP anion exchange column
(Cytiva) in buffer 4 without ZnSO$_4$ and eluted in a 20 CV NaCl gradient. The peak containing NUT was
further purified by gel filtration on a Superdex 75 Increase 10/300 GL column (Cytiva) equilibrated in
buffer 5 without ZnSO$_4$. The final proteins were concentrated in a prewashed Amicon Ultra-15
Centrifugal filter (molecular weight cut off 3 kDa; EMD Millipore), flash-frozen in liquid nitrogen and
stored at -80 °C.
1072
The GST-E1A amino acids 1-289 and GST-p53 AD amino acids 1-61 proteins were expressed in *E. coli*
BL21 (DE3) cells. Cell pellets were resuspended and sonicated in lysis buffer (20 mM Tris-HCl pH 7.5,
300 mM NaCl, 0.5 mM TCEP), supplemented with EDTA-free protease inhibitors (Roche), and 10 µl of
benzonase nuclease (Sigma-Aldrich) and lysed by sonication. The lysate was clarified by centrifugation
for 1h in a SS-34 rotor (Sorvall) at 12000 rpm and applied to a Glutathione Sepharose 4 Fast Flow resin
according to instructions from the manufacturer (Cytiva). The resin was washed with 20 CV lysis buffer
and the protein was eluted in lysis buffer containing 20 mM reduced L-Glutathione. The protein solution
was then dialyzed overnight against lysis buffer to remove L-Glutathione and then applied to a 5 ml
HiTrap Q HP anion exchange column (Cytiva) and further purified by gel filtration on a Superdex 75
Increase 10/300 GL column (Cytiva) as previously described for the NUT fragments.
1083
The BRD4 GFP-BD1-BD2, GFP-BD1-BD2 N140A/N433A, GFP-BD1-BD2 N140A and GFP-BD1-BD2
N433A were all expressed in *E. coli* BL21 (DE3) cells. The cells were sonicated in lysis buffer (20 mM
Tris-HCl pH 7.5, 300 mM NaCl, 0.5 mM TCEP) and the lysate was clarified as previously described for
the other proteins. The clarified lysate was applied to Ni$^{2+}$ charged IMAC Sepharose 6 FF (Cytiva) beads
preequilibrated with lysis buffer containing 20 mM Imidazole. The column was washed with 20 column
volumes (CVs) of lysis buffer and the bound proteins eluted with the lysis buffer containing 300 mM
Imidazole. Imidazole was removed by dialysis overnight against 1L lysis buffer and the proteins were

loaded on a 5 ml HiTrap SP HP cation exchange column (Cytiva) as described for TAZ2 and further
purified using a HiLoad 16/600 Superdex 200 column (Cytiva) preequilibrated with 20 mM Tris pH 7.0,
300 mM NaCl and 1 mM DTT. The purified proteins were concentrated using a prewashed Amicon
Ultra-10 concentrator (molecular weight cut off 30 kDa; EMD Millipore), flash frozen in liquid N2 and
stored in -80°C.
1096
1097     The GFP-(Bromo)5 and GFP-(Bromo)5 N140A were purified as described previously[58]. The co-
1098     expressed *S. cerevisiae* histones octamer was purified as described previously[157]. NUT peptides were
1099     ordered from Proteogenix.
1100
**HAT assays.** The p300 autoacetylation and substrate acetylation assays were performed using [$^{14}$C]
acetyl-CoA (Perkin-Elmer). Autoacetylation of p300 was quantified by autoradiography after SDS-PAGE
gel analysis. The acetylation reaction was performed for 30 minutes at 30°C in 1x HAT buffer (20 mM
Tris-HCl pH 7.5, 100 mM NaCl, 1 mM DTT, 10 % glycerol and 1x Complete EDTA-free protease
inhibitor (Roche)) with 2.5 µM p300s or p300 core and 50 µM [$^{14}$C] acetyl-CoA with or without
increasing concentrations of the different substrates (NUT fragments, NCP, E1A). 5 µl of the reaction
was quenched by addition of 2.5 µl of 3x SDS gel loading buffer followed by analysis on a 4-20% SDS-
PAGE gel. The gels were stained with Quick Coomassie Stain (Generon), fixed for 30 min in enhancer
solution (KODAK ENLIGHTNING Rapid Autoradiography Enhancer, Perkin Elmer) and dried for 2
hours using a Bio-Rad Gel Dryer. The $^{14}$C signal was quantified using phosphor imaging (Typhoon,
Cytiva).
1112
***In vitro* nucleosome acetyltransferase assays.** Nucleosome acetylation reactions were performed in 20
µl volume in acetylation buffer (20 mM Tris pH 7.5, 100 mM NaCl, 1 mM DTT, 10% glycerol and
complete Protease Inhibitor EDTA-free (Roche)) with 50 µM Acetyl-CoA, 2.5 µM of p300s or p300 core
and 2 µM of nucleosome core particles (NCPs). Control experiments contained only NCPs and Acetyl-
CoA without p300. The reactions were incubated for 30 minutes at 30°C and stopped by addition of 1x
SDS loading buffer. Each condition was performed in duplicate. The samples were then analyzed by
SDS-PAGE and mass spectrometry.
1120
**Multi angle laser light scattering-size exclusion chromatography.** SEC-MALLS was performed at a
flow rate of 0.5 ml/min on a Superdex 75 Increase 10/300 GL column equilibrated in buffer (20 mM
HEPES pH7.0, 300 mM NaCl, 5 µM ZnSO$_4$, 0.5 mM TCEP). 50 µl of TAZ2, NUT4 or the TAZ2:NUT4
complex at 2 mg/ml were injected onto the column and multi angle laser light scattering was recorded
with a laser emitting at 690 nm using a DAWN-EOS detector (Wyatt Technology Corp). The refractive
index was measured using a RI2000 detector (Schambeck SFD). The molecular weight was calculated
from differential refractive index measurements across the center of the elution peaks using the Debye
model for protein using ASTRA software version 6.0.5.3.
1129
**Biolayer interferometry assay.** BLI measurements were performed with a ForteBio Octet RED384
instrument and ForteBio biosensors. Data analyses was done using ForteBio Data Analysis 9.0 software.
Kinetics assays were carried out at 25 °C using settings of Standard Kinetics Acquisition rate (5.0 Hz,
averaging by 20) at a sample plate shake speed of 1000 rpm. The GST-tagged NUT8 wild type and
mutants were loaded onto Anti-GST biosensors and the TAZ2 protein was used as an analyte. The GST-
NUT-loaded Anti-GST biosensors were dipped in kinetic buffer (1x PBS, 0.1% BSA, 0.02 % Tween 20)
to establish a baseline time course and then dipped into wells containing TAZ2 (1723-1836) at various
concentrations diluted in the same kinetic buffer to monitor NUT-TAZ2 binding. The dissociation step
was monitored by dipping the biosensors back into the wells used to collect the baseline time course. To
monitor binding due to nonspecific interactions of TAZ2 with the biosensors, Anti-GST biosensors were
used as controls for every analyte concentration and a double referencing method was used for data

subtraction and analysis. The subtracted binding curves were analysed and plotted as a steady-state
response to obtain the dissociation constant $K_d$ value.

1143

**Limited proteolysis.** To identify stable variants of NUT we performed limited proteolysis by Trypsin and
Chymotrypsin. 100µl of the purified CH3-NUT-F1c complex (250 µg) and TAZ2-NUT4 complex (250
µg) were incubated at 4 °C with Trypsin or Chymotrypsin with a ratio of 1:100 (w/w). At the indicated
time points (30 min, 2 hours, overnight), 30 µl of the reaction was quenched by addition of 15 µl of 3x
SDS gel loading buffer followed by analysis on 4-20 % SDS-PAGE gel. The stable fragments on gel were
analysed by acid hydrolysis and mass spectrometry at the Proteomics Core Facility (EMBL, Heidelberg).

1150

**SAXS analysis.** X-ray scattering data were collected on the NUT-F1c sample at 100 µM in phase
separation buffer: 20 mM Tris pH 7.0, 2 mM DTT, 125 mM NaCl and 10% (w/v) PEG 6000 at the B21
BioSAXS beamline of the Diamond Light Source (Chilton, Oxfordshire, UK). The data were analysed
with the ATSAS package[158]. Data collected from the sample were substracted from buffer scattering. $R_g$
values were obtained from the Guinier approximation $sR_g < 1.3$ using Primus[159]. Distance distribution
functions p(r) were computed from the entire scattering curve using GNOM[159].

1157

1158 *In vitro* **NUT phase separation assay.** Phase separation of purified GST-NUT-F1c *in vitro* was done
1159 phase separation buffer. The concentrations of GST-NUT-F1c used were 0-200 µM. A GST control was
1160 used in the same concentration range under identical conditions. The droplets were visually observed and
1161 turbidity of the sample ($OD_{600}$) quantified using NanoDrop (Thermo scientific).

1162

**GST pulldowns.** For GST pulldowns, 50 µg of GST-tagged NUT constructs were mixed at an equimolar
concentration with binding partners in 50 µl pulldown buffer (20 mM HEPES pH 8.0, 300 mM NaCl, 0.5
mM TCEP, 5µM ZnSO₄, 0.1% Triton X-100) containing 25 µl of a 50 % slurry of GST Sepharose 4 Fast
Flow beads (Cytiva) per reaction. Reactions were incubated for 30 min at 4 °C. Twenty-five microliters
of the reaction were withdrawn as the reaction input and the remainder was washed 5 times with 500 µl of
pulldown buffer. Samples were boiled in 1x sample loading buffer for 5 min and analysed by SDS-
PAGE.

1170

**Luciferase reporter assays.** Cos7 cells were co-transfected for 36 hours with β-Galactosidase reporter
plasmid and UAS-promoter luciferase reporter plasmid either alone or with GAL4 DNA binding domain
(DBD) plasmid or increasing amount of GAL4-DBD-NUT-F1c plasmid. Cells were then washed and
incubated 15 min at RT with Luciferase lysis buffer (1% Triton X-100, 10% Glycerol, 2 mM EDTA, 25
mM Tris-HCl pH 7.8). Luciferase activity was measured directly after mixing 10µl of cellular extract
with 50 µl of Luciferase substrate (Luciferase assay kit, Stratagene). Normalization was made by
measuring the β-Gal activity after 1-hour incubation of 7.5 µl of cellular extract with 50 µl of β-Gal
substrate (β-Gal detection kit II, Clontech).

1179

**Proteomics of histone modifications.** For the identification and quantification of histone acetylation
following *in vitro* HAT assays, we followed the previously published procedure[52,53]. In brief, reconstituted
NCPs were incubated with 100µM acetyl-CoA, as well as purified p300s (340-2,094) or the p300 core
domain (1,043-1,666) for 30 min at 30°C. HPLC-MS/MS was used to quantify site specific H2A, H2B,
H3 and H4 acetylation by label-free quantification. The peak area of each histone peptide was then
normalized. Two technical replicate experiments were done.

1186

**Isolation and Purification of 1x601 Widom DNA.** Carrier plasmids of 1x Widom 601 repeats were
purified using QIAGEN Plasmid Giga kit from 6L LB culture according to the manufacturer's
instructions. 1x arrays DNA were cut from carrier plasmid DNA by incubation overnight with 5,000 units
of *EcoR*V-HF restriction endonuclease (New England Biolabs) in 1x CutSmart Buffer (20 mM Tris,OAc,

pH 7.9, 50 mM KOAc, 10 mM Mg[OAc]$_2$, 0.1 mg/mL BSA). The excised DNA fragments were
separated from the linearized plasmid by PEG precipitation. 0.192 volume of NaCl and 8% PEG 6000
were added to the DNA sample and incubated for 1h on ice, the sample was then centrifuged for 20 min
at 4 °C, 27000 g. the *Eco*RV fragments contained in the supernatant were precipitated by the addition of
2.5 volumes of 100% cold ethanol. The DNA fragments were air dried for 10 min and dissolved in buffer
TE (10 mM Tris/HCl, pH 8.0, 1 mM EDTA). Purity of the extracted 12x601 and 1x601 fragments were
verified on 0.7% agarose gels and the concentration measured by 260 nm absorbance on a NanoDrop
spectrophotometer (Thermo Scientific).
1199
1200  **Isolation and Purification of 12x Widom 601 Array DNA.** 12x601 DNA template was prepared as
1201  described previously[160]. In short, 12-mer DNA with177 bp repeats of the 601 sequence separated
1202  from the rest of the plasmid by *Eco*RV sites were expressed in *E.coli* strain DH5a. The DNA was
1203  then extracted by alkaline lysis and purified on a Sepharose-6 column. The DNA template was
1204  separated from the plasmid by *Eco*RV-HF digestion and PEG6000 precipitation.
1205
**Histones octamer preparation.** Wild type human histones (H2A, H2B, H3 and H4) and the cysteine
mutant Xenopus H4 S47C were individually expressed in *E. coli* strain Rosetta 2 DE3 pLysS and purified
following the protocol described[161], but using a RESOURCE S 6 ml cation exchange chromatography
column. For octamer reconstitution, equimolar ratios of the four histones were refolded together in the
presence of 5 mM β-mercaptoethanol. The octamer was then separated from dimers by gel filtration using
a Superdex 200 column (Cytiva). Two batches of octamers have been generated: one fully wild-type
human octamer (unlabelled octamer) and an octamer formed by wild-type human H2A, H2B, H3 and
Alexa Fluor™ 594 C5 Maleimide labelled *Xenopus* H4 S47C (labelled octamer).
1214
**Fluorescent labelling of histones.** The *Xenopus* H4 S47C mutant histone was labelled using Alexa
Fluor™ 594 C5 Maleimide following the protocol from[58]. In short, lyophilised H4 S47C was resuspended
in 7M urea, 500mM Tris-HCl pH 8 and incubated with Tris-neutralized TCEP to a final concentration of
2 mM for 1 hr and 30 min at RT. Alexa Fluor™ 594 C5 Maleimide stock solution was added dropwise to
a 1.5 molar excess and the mix was incubated for 4 hr at RT in the dark. The reaction was quenched by
the addition of β-mercaptoethanol. The histone was then purified from the free dye by ion exchange
chromatography using a HiTrap SP HP cation exchange column followed by dialysis into ultra-pure water
with 5 mM β-mercaptoethanol.
1223
**Mononucleosome assembly.** DNA and *Saccharomyces cerevisiae* histones octamer were mixed in a
1:0.5 molar ratio. Nucleosomes were reconstituted by salt gradient dialysis method and dialysed against
buffer 20 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM DTT with continuous decrease of NaCl
concentrations (1.5, 1, 0.6 and 0.25 M) at 4 °C, 3h each dialysis step with the third dialysis step
proceeding overnight using mini dialysis units (10,000 Da MWCO, Pierce Slide-a-lyzer). The
reconstituted nucleosomes were quality controlled by electrophoretic mobility shift assays on 5%
acrylamide native gels with 0.25x TBE running buffer.
1231
**12x Nucleosome array assembly, acetylation and phase separation.** Chromatin was reconstituted *in*
*vitro* by assembling the 12-mer DNA with histone octamers by step-wise salt dialysis as described
previously[160], but using KCl instead of NaCl. The optimal octamer to DNA ratio was determined
empirically. A mix of labelled and unlabelled histone octamer was used. The ratio of labelled to
unlabelled histone octamer was also determined empirically. The reconstituted 12-mer nucleosome arrays
were quality controlled by *Sca*I restriction enzyme digestion to isolate single nucleosome particles and by
electrophoretic mobility shift assays on 2% agarose gel in 0.5x TBE
12x601 nucleosome arrays at 2 μM mono-nucleosome concentration were incubated with 1 μM of the
p300 core domain in Acetylation buffer (20 mM Tris-HCl pH 7.0, 100 mM NaCl, 10% glycerol, 1 mM

DTT) for 30 min at 30 °C. The non-acetylated chromatin reactions were incubated under identical
conditions but without p300 core enzyme. After acetylation, 5 μM BRD4 GFP-Bromo$_5$, GFP-Bromo$_5$
N140A, 10 μM GFP-BD1-BD2, 10 μM GFP-BD1-BD2 N140A, 10 μM GFP-BD1-BD2 N433A or 10
μM GFP-BD1-BD2 N140A/N433A were added to nucleosomal arrays for a final concentration of 1 μM
mono-nucleosome in phase separation buffer (25 mM Tris, pH7.5, 150 mM KOAc, 1 mM MgOAc, 5%
glycerol, 5 mM DTT, 0.1 mM EDTA, 0.1 mg/mL BSA)[58].

1247
**Microscopy of chromatin droplets.** For confocal microscopy image acquisition, a Zeiss LSM 980
Airyscan 2 microscope equipped with a Plan-Apochromat 63x/1.40 Oil DIC f/ELYRA and ZEN Blue 3
acquisition and analysis software was used. Samples were imaged in 384 well glass bottom plates for
microscopy which were prepared as described in[58]. In short, microscopy plates were washed with 5%
Hellmanex, etched with 1 M NaOH and treated with 25 mg/mL 5K mPEG-silane. Prior to use, individual
wells were passivated by incubation with freshly prepared 100 mg/mL BSA.

1254
**Cross-linking mass spectrometry.** Purified solutions of both NCPs and p300 core-CH3 (1043-1828)
were diluted to a concentration between 0.5 and 1 mg/ml in 20 mM HEPES pH 7.0, 300 mM NaCl, 1 mM
DTT and 5 μM ZnCl$_2$ and cross linked using a homobifunctional, isotopically-coded crosslinker
DiSuccinimidyl Suberate DSS-H12/D12 (Creative Molecules Inc.) at a concentration of 2 mM for 30 min
at 35°C 600 rpm. The cross-linking reaction was then quenched with 0.1 volumes of 1M Ammonium
bicarbonate for 10 min at 35 °C 600 rpm. The sample was then treated with 0.8 volumes of 10 M urea and
250 mM Ammonium bicarbonate as well as 0.05 volumes RapiGest SF Surfactant (Waters, cat. No.
186008090) and sonicated for 1 min in an ultrasound bath. DTT was later added with a final
concentration of 10 mM, incubated for 30 min at 37 °C and freshly prepared Iodoacetamide was added to
a final concentration of 15 mM and incubated for 30 min at room temperature in the dark. The protein
was sequentially digested with Endoproteinase Lys-C and Trypsin. The enzymes were added at an
enzyme to substrate ratio of 1:100 for LysC and 1:50 for Trypsin and the reaction incubated for 3-4 h at
37°C. The reaction was started by LysC followed by Trypsin addition after 4h. After digestion, the
sample was acidified with Trifluoroacetic acid (1% v/v final concentration) at 37°C for 30 min, spun
down at 17,000 g for 5 min and the peptides were fractionated by peptide size-exclusion chromatography.
The mass spectrometry experiment and cross-links identification were performed by the EMBL
Proteomics core facility.

1272
**Immunofluorescence and drug treatment.** Cos7 cells were seeded in glass coverslips (Lab-Tek), and
GFP and RFP bearing constructs were transfected by Lipofectamine 2000. Twenty-four hours later, cells
were washed with PBS and fixed with 4% paraformaldehyde for 15 min, followed by permeabilization
with 0.2% Triton-X100 and 1 × PBS for 10 min at room temperature. After being blocked with 5% dry
milk, 0.2% Tween 20, and 1 × PBS at RT for 30 min, slides were incubated with anti-acH4 antibody
(1:100 dilution, Millipore) or anti-HDAC1 antibody (1:100 dilution, Cell Signaling Technology) in 1%
dry milk, 0.1% Tween 20, and 1 × PBS in a humidified chamber at 4°C overnight and then washed three
times for 5 min each in the antibody dilution buffer. The secondary antibodies were diluted at 1:500 in the
same buffer and incubated in a humidified chamber for 30 min at 37°C, and then washed as for the
primary antibodies. The DNA was counterstained by DAPI, and the slides were mounted in Dako
fluorescent mounting medium. For the drug treatment assays, twenty-four hours after being transfected
with GFP and RFP bearing constructs, Cos7 cells were treated with DMSO (Sigma), 1,6-hexanediol
(10%, 1 hour, Sigma), A485 (3 μM, 1 hour, Selleck), JQ1 (1 μM, 1 hour, Selleck), TSA (0.3 μM, 12
hours, Selleck) or CBP30 (1 μM, 1 hour, Selleck), then the cells were washed with PBS and fixed with
4% paraformaldehyde for 15 min.

1288
**Co-immunoprecipitation.** Cell pellets were lysed in 300 mM NaCl buffer (20 mM HEPES-KOH pH 7.9,
300 mM NaCl, 3 mM MgCl$_2$, 0.1% NP-40, 1X protease inhibitor cocktail (Roche), 1X phosphatase

inhibitor cocktail-1 (Sigma) and 10 mM sodium butyrate) for 30 min at 4 °C with rotation. After a 15 min
centrifugation at 12,000g at 4°C, the supernatants were collected and separated into input and
immunoprecipitate (IP) lysates. IP lysates were mixed with GFP-selector (pre-washed with 300 mM NaCl
buffer three times, NanoTag) and rotated overnight at 4 °C. After washing with 300 mM NaCl buffer, IPs
were released by adding Laemmli sample buffer to beads and boiling for 5 min at 100°C.

1296

**Fluorescence recovery after photobleaching (FRAP).** Live imaging and fluorescence recovery after
photobleaching experiments (FRAP) were performed with the LSM710 Confocor3 confocal microscope
(Carl Zeiss), equipped with the Plan-Apochromat C 63X /1.4 oil-immersion objective. FRAP area, time
and time interval (1 image every 783ms) were maintained identical between the different BRD4-NUT
mutants and repeat measurements (N=3-4, 45 to 65 foci measured per conditions) in order to photobleach
3-6 foci at the same time. To compare recovery kinetics against BRD4-NUT wild-type, FRAP
measurements were fitted to a single exponential curve, $I_{(t)}=I_{(0)} + k_1 e^{-k2t}$ (performed with ZEN software;
Carl Zeiss Microscopy) to determine the characteristic time of recovery. Values of $\tau$ (1/ k2) were
compiled and statistical significance was tested using the Student's t-test (Excel).

**NMR experiments and AlphaFold calculations.** Backbone resonances of both proteins free in solution
were assigned by measuring HNCO, HNCA, and HN(CO)CA spectra on $^{15}$N,$^{13}$C-labelled samples at 600
(NUT7) and 700 (TAZ2) MHz. We were able to assign 87% (95%) of the NH pairs, 72% (82%) of the
CO resonances and 88% (96%) of the CA resonances in the backbone of NUT7 (TAZ2).

1310

1311     Side-chain resonances of both proteins free in solution were assigned by measuring HBHA(CO)NH,
1312     H(CCO)NH, and CC(CO)NH spectra on $^{15}$N,$^{13}$C-labelled samples at 600 (NUT7) and 700 (TAZ2) MHz.
1313     We were able to assign 70% (55%) of the HA resonances, 57% (45%) of the HB resonances, and 45%
1314     (53%) of the CB resonances in NUT7 (TAZ2), as well as part of the γ-protons.

1315

1316     To determine the binding site of NUT7 on TAZ2, we measured $^1$H-$^{15}$N HSQC spectra on $^{15}$N-labelled
1317     TAZ2 before and after adding unlabelled NUT7. For each of the assigned peaks in the spectra we
1318     calculate the chemical shift perturbations (CSP) defined as

1319
$$CSP = \sqrt{\frac{1}{2}(\Delta\delta_H^2 + (0.14 \cdot \Delta\delta_N^2)^2)},$$

1320     in which $\Delta\delta_H^2$ and $\Delta\delta_N^2$ are the chemical shift differences upon adding NUT7 in the $^1$H and $^{15}$N dimensions,
1321     respectively. We used the procedure described by Schumann et al.[162] to determine whether measured
1322     CSPs are significant or not. We used the same approach to map the binding site on NUT7 monitoring a
1323     titration of a solution of $^{15}$N-labelled NUT7 with unlabelled TAZ2 using $^1$H-$^{15}$N HSQC spectra.

1324

Both $^{13}$C- and $^{15}$N-HSQC-NOESY spectra on a sample of $^{15}$N,$^{13}$C-labelled TAZ2 to which one equivalent
of unlabelled NUT7 was added and $^{15}$N-HSQC-NOESY spectra on $^{15}$N-labelled NUT7 in admixture with
one equivalent of unlabelled TAZ2 were measured using standard pulse sequences. All NOESY spectra
were measured at 950 MHz with a mixing time of 100 ms. The AlphaFold computational models
shown in Fig. 3e and Fig. 4a were obtained from alphafold.ebi.ac.uk. Models shown in Fig. 3f
and Fig. 4b were calculated using AlphaFold v2.1.1 with multimer model v1 weights[73].

1331

**HADDOCK docking.** We used the HADDOCK protocol[70] with experimental information from titrations
and NOESY spectra to derive a structural model of the interaction. We used the NMRbox platform[163] to
carry out the calculations. In a first step, we used the results of the titration experiments to derive
ambiguous distance restraints (AIRs) describing the binding interface, and we used these AIRs as input
for HADDOCK to dock extended conformations of NUT7 on the TAZ2 structure derived from PDB
2mzd. The best 200 structures resulting from the calculations were clustered based on the fraction of

common contacts (FCC). The resulting clusters were sorted according to their agreement with the AIRs.
Structures in the best cluster display the interaction of residues P448-E454 in NUT7 with TAZ2. Thus,
we used the Flexible-Meccano algorithm[164] to generate an ensemble of 100 conformers of NUT7, in
which the conformation of residues P448-E454 is the same as in the structure resulted from the
HADDOCK calculation, whereas the rest of the chain behaves like a statistical coil. Then, we used
HADDOCK again to dock this ensemble of structures on the same TAZ2 conformation as above. In this
run, in addition to the same AIRs as in the preliminary run, we used distance restraints derived from
NOESY spectra between S1726 in TAZ2 and residues L463 and A464 in NUT7. We also used TALOS-
N[165] to derive restraints on the backbone dihedral angles of residues P448-E454 from chemical shift
values. We repeated this calculation ten times with different initial seeds and velocities. Finally, the best
200 structures from each of the ten runs were clustered based on the FCC. The resulting clusters were
sorted according to their agreement with the AIRs and NOE-based restraints. Then, from the best-ranked
cluster in each of the runs, we selected the structure with the best HADDOCK score (which is a linear
combination of various energy terms and buried surface area), thereby obtaining a final set of 10
structures.
1353
**Cryo-electron microscopy grid preparation, data collection and image processing**
Quantifoil R1.2/1.3 Au 300 grids were glow discharged for 5 min at 40 mA on a Quorum Gloqube glow-
discharge unit prior to application of the sample. 3 μl of 1 μM final concentration of the mononucleosome
particles in buffer 20 mM Tris-HCl, pH8.0, 250 mM NaCl, 1 mM EDTA, 1 mM DTT were applied to the
grid, blotted for 3 s at blot force 10 and plunge frozen into liquid ethane using a Vitrobot Mark IV (FEI
Thermo Fisher), which was set at 4°C and 100% humidity. Cryo-EM data was collected on a Thermo
Fisher Scientific Titan Krios G3 transmission electron microscope at the LISCB at the University of
Leicester. Electron micrographs were recorded using a K3 direct electron detector (Gatan Inc.) at a dose
rate of 15 e⁻/pix/sec and a calibrated pixel size 1.086 Å. Focusing was performed over a range between -
2.3 and -1.1 μm in 0.3 μm intervals. Image processing of the NCPs was performed as follows: movie
stacks were corrected for beam-induced motion and then integrated using MotionCor2[166]. All frames were
retained and a patch alignment of 4 x 4 was used. Contrast transfer function (CTF) parameters for each
micrograph were estimated by Gctf[167]. Integrated movies were inspected with Relion-3.1 for further
image processing (243 movies). Initial particle sets were picked manually, extracted, and subjected to 2D
classification in Relion 3.1[168]. Resulting classes corresponding to protein complexes were subsequently
employed as references for auto-picking in Relion 3.1. Particle extraction was carried out from
micrographs using a box size of 128 pixels (pixel size: 1.08 Å/pixel). An initial dataset of 80764 particles
was cleaned by several iterations of 2D classification, classes exhibiting clear secondary structure features
were selected for 3D classification. After iterative rounds of 3D classifications, well defined classes were
selected and global refinement was performed.
1374

**Acknowledgements** We thank Christos Savva and TJ Ragan for CryoEM support. We thank Nathan Cowieson from the B21 BioSAXS beamline of the Diamond Light Source; Yingming Zhao and Jinjun Gao for mass spectroscopy analysis; Luke Bailey, Sarah Northall and Dipti Vashi for help with histone preparations. Michael K. Rosen for plasmids encoding GFP-(Bomo)$_5$ and GFP-(Bromo)$_5$ N140A; Stephen M. Hinshaw and Stephen C. Harrison for plasmids encoding *S. cerevisiae* histones; Andrew Flaus for *H. sapiens* histones; Timothy J. Richmond for histone H4 mutant plasmids; Andy Turnell for E1A plasmids. Work in the S.K. laboratory was supported by ANR Episperm4 program, by Plan Cancer Pitcher, by MSD Avenir ERICAN programs, by the Cancer ITMO (Multi-Organisation Thematic Institute) of the French Alliance for Life Sciences and Health (AVIESAN) MIC program, by the "Université Grenoble Alpes" ANR-15-IDEX-02 SYMER and LIFE programs and by Fondation ARC program RF20190208471. Work in the T.S. laboratory was supported by the BBSRC (BB/S019510/1; BB/R016275/1). Work in the D.P. laboratory was supported by grants from the Worldwide Cancer Research [16-0280], the Wellcome Trust [221881/Z/20/Z], Instruct-ERIC [PID: 18571] and the Medical Research Council [MR/W001667/1].

**Author contributions** Z.I. designed and performed most experiments, analysed and validated the data with support by L.F. T.W., O.D., N.H., C.C. and N.R performed cell biology experiments. N.S. performed NMR experiments. A.R performed chromatin analyses *in vitro*. Y. Z. and J. G. performed mass spectroscopy analysis. T.S., M.B., and S.K. provided supervision, funding acquisition and commented on the draft. D.P. was involved in conceptualization, supervision, project administration, funding acquisition. Z.I and D.P. wrote the original and revised drafts with input from all authors.

**Competing interests**: Y. Z. is a founder, board member, advisor to, and inventor on patents licensed to PTM Biolabs Inc (Hangzhou, China and Chicago, IL) and Maponos Therapeutics Inc. (Chicago, IL). The other authors declare no competing interests.

# Supplemental Figures

**Figure S1. Identification of an acidic activation domain in NUT. a,** Schematic of the different NUT constructs used. Sequence coordinates correspond to NUTM1 (UNIPROT Q86Y26). **b**, GST pulldown analysis between the p300 core-CH3 or the isolated TAZ2 domain and different NUT fragments. **c**, Luciferase reporter assay: Cos7 cells were co-transfected with a luciferase reporter plasmid and 'empty' plasmid, or containing the Gal4 DNA binding domain (DBD) or increasing amounts of Gal4-NUT-F1c plasmid or NUT-F1c lacking the Gal4 DBD. Luciferase expression was measured by luminescence (Luciferase assay kit, Stratagene) (right). Error bars represent standard deviations of 3 independent technical replicate experiments. **d**, Limited proteolysis of the p300 CH3-NUT-F1c complex (left) or the p300 TAZ2-NUT4 complex (right). After overnight incubation with Trypsin or Chymotrypsin samples were analysed by SDS-PAGE followed by mass spectrometry. The identified NUT fragments are highlighted on the gels and their corresponding sequences are shown on the bottom. Position of the NUT AD is indicated **e**, SEC analysis of the p300 TAZ2 domain (black), NUT9 (396-482) (blue) or the TAZ2:NUT9 complex (red) on a Superdex 75 Increase 10/300 GL column. Bottom: SDS-PAGE analysis of the peak fractions of the TAZ2-NUT9 complex.

**Figure S2. Identification of an acidic activation domain in NUT. a,** Size-exclusion chromatography (SEC) analysis of p300 CH3-NUT-F1c complex on HiLoad Superdex 200 column. **b**, SEC analysis of p300 core-CH3-NUT4 complex on Superdex 200 Increase 10/300. **c**, p300 core-CH3-NUT1 complex on Superdex 200 increase 10/300 GL. **d**, p300 TAZ2-NUT4 complex on Superdex 75 increase 10/300 GL. **e**, p300 TAZ2-NUT6 complex on Superdex 75 increase 10/300 GL. Bottom of each panel: SDS-PAGE analysis of the peak fractions. **f**, SEC analysis of the TAZ2-NUT10 complex on Superdex 75 increase 10/300 GL column (top). Bottom: Reverse-Phase HPLC analysis of TAZ2-NUT10 peptide complex fractions followed by electrospray ionisation time of flight mass spectrometry (LC-ESI-TOF-MS). Table: Summary of the mass spectrometry analysis data showing the presence of TAZ2 (peak 1) and the NUT peptide (peak 4). **g**, Competition analysis: Cos7 cells were transfected with constructs expressing GFP-BRD4-NUT (green) alone or co-transfected with RFP-NUT1, 2 or 3 (red). Immunofluorescence with anti-p300 antibody was used to visualize condensation of endogenous p300.

**Figure S3. Analysis of the TAZ2-NUT interaction. a,** Representative Biolayer Interferometry (BLI) profiles of p300 TAZ2 interaction with NUT8 (396-470) mutants 1, 2, 3 and 4 from which the steady-state data shown in Fig. 3b were derived. The GST-NUT8 construct was immobilized on the biosensors and TAZ2 was used as analyte and titrated at the indicated concentrations. **b,** GST pulldown analysis of TAZ2 and NUT8 (396-470) wild type and mutants 3 and 4. **c,** AlphaFold model of the TAZ2-NUT complex surface rendered and coloured according to electrostatic potential. **d,** (Left) GST pulldown analysis of TAZ2 and NUT7 (448-492) mutant. Sequences of the NUT (448-492) wild type and mutant are shown below. (Middle) [1]H-[15]N HSQC spectrum of free TAZ2 (gray) compared to a 1:1 molar ratio admixture with NUT7 (blue) or 1:1 (orange) and 2:1 (red) admixtures with the NUT7 mutant. (Right) Chemical shift perturbations (CSPs) upon adding 1 equimolar amount of NUT7 wild type (gray bars) to TAZ2 or after adding 1 (blue) and 2 (red) molar equivalents of the NUT7 mutant. **e,** [1]H-[15]N HSQC spectra of TAZ2 when 1 equivalent of NUT7 wild type (blue), p53 TAD (1-61) (red) and both NUT7 and p53 (black) are added. **f,** Overview of different TF ADs in complex with the TAZ2 domain. The N – and C-termini of the TF ADs are indicated. **g**, Predicted Aligned Error of the TAZ2-NUT AlphaFold model. The colour at position (x,y) indicates AlphaFold's expected position error at residue x, when the predicted and true structures are aligned on residue y.

**Figure S4. Role of the TAZ2 domain in p300 regulation and substrate acetylation. a,** p300s was incubated with [14C]Acetyl-CoA in the presence or absence (C: p300s alone) of (**a**) NUT-F1c or NUT4, **b**, NUT1, NUT2, NUT3. Samples were analysed by SDS-PAGE followed by Coomassie staining and 14C phosphor imaging. Representative data of three independent experiments are shown. **c,** EMSA analysis of NCP binding of the p300 TAZ2 domain. **d**, NCP binding of p300 core. **e**, NCP binding of p300 core-CH3. Native PAGE gels were stained with SybrSafe DNA stain or Coomassie Blue. **f,** CryoEM density for the NCP used in biochemical studies. An atomic model of the NCP (PDBs: 3LZ0) is docked into the density. **g**, SDS-PAGE gel showing input material used for quantitative mass spectrometry experiments. **h,** Quantitative mass spectrometry of NCP acetylation. Reconstituted NCPs were incubated with Acetyl-CoA and p300s or the p300 core. Quantification of the site-specific histone H2B, H3 and H4 acetylation by label-free HPLC/MS/MS. Error bars represent standard deviations from two independent technical replicate experiments. **i,** Cross-link (XL-MS) analysis of the p300 core-CH3:NCP complex. p300 residues that cross-link to the NCP histones are shown in red on p300 core (4BHW) and TAZ2 (3IO2) structures. **j**, Graphical representation of the p300:histone and histones intermolecular cross-links and intra-p300 core-CH3 cross-links. **k**, p300 (top) and CBP (bottom) gain--of-function mutations in the TAZ2 domain.

**Figure S5. Mechanism of condensation of BRD4-NUT and p300**. **a,** Colocalization of BRD4-NUT condensates with acetylated H4 (H4Ac) or **b**, with HDAC1. Cos7 cells were transfected with the indicated GFP-BRD4-NUT constructs and analysed by immunofluorescence using anti-H4Ac antibody and anti-HDAC1 antibody. Scale bars, 10 µm. **c**, Cos7 cells were co-transfected with HA-p300 and the indicated GFP-BRD4-NUT variants or GFP alone as the control. Co-immunoprecipitation (IP) experiments were performed using anti-GFP antibody and immuno-blotted (IB) for GFP, p300 and Acetylated-Lysine. (Left panel) Input samples were loaded as a control. Experiments were independently repeated at least three times. **d**, EMSA analysis of 12x nucleosome templates assembly (left) on 2% agarose (12xNCP) native gels. The 12x nucleosome template was digested with *Sca*I restriction enzyme to isolate single NCPs. (Right) EMSA analysis of mononucleosome (1x NCP) assembly on 5% native PAGE stained with Sybr safe or Coomassie Blue. **e**, Fluorescence microscopy images of AF594-labeled dodecameric nucleosome arrays. Non-acetylated 12x NCP chromatin formed droplets (bottom panel) that are dissolved after p300 acetylation (top panel). Scale bars, 10 µm. **f**, Acetylated and non-acetylated 12x NCP chromatin templates incubated with GFP-labelled BRD4 (BD1)₅. Scale bars, 10 µm. **g**, Acetylated 12x NCP chromatin templates incubated with GFP-labelled BRD4 (BD1)₅ N140A, BD1-BD2 N140A/N433A, BD1-BD2 N140A or BD1-BD2 N433A.

**Figure S6. Condensation is driven by bromodomain mulivalency**.
**a,** Concentration-dependent droplet formation of GST-NUT-F1c in droplet formation buffer with 125 mM NaCl and 10% PEG-8000 measured by turbidity. **b**, Experimental SAXS profile (log intensities calculated as a function of momentum transfer) for a 100 $\mu$M solution of GST-NUT-F1c in droplet formation buffer (blue). The Guinier region is inset. **c**, Distance distribution function.

**Movie S1. Dynamics of condensation of BD1-BD2-AD and p300.**
Time laps imaging. Left panel: GFP- BD1-BD2-AD. Middle: RFP-p300. Right: Merge of GFP- BD1-BD2-AD and RFP-p300.

**Movie S2. Dynamics of condensation of BRD4-NUT and p300.**
Time laps imaging. Left panel: GFP-BRD4-NUT. Middle: RFP-p300. Right: Merge of GFP-BRD4-NUT and RFP-p300.

**Table S1 – List of NUT constructs and binding experiments**

| Sample | Amino acid residues | Binding* |
|---|---|---|
| **NUT F1c** | 347-588 | + |
| **NUT1** | 347-446 | +/– |
| **NUT2** | 427-527 | + |
| **NUT3** | 508-588 | – |
| **NUT4** | 347-480 | + |
| **NUT5** | 347-514 | + |
| **NUT6** | 351-470 | + |
| **NUT7** | 448-492 | + |
| **NUT7 mut1** | 448-492 | + |
| **NUT8** | 396-470 | + |
| **NUT8 mut1** | 396-470 | – |
| **NUT8 mut2** | 396-470 | – |
| **NUT8 mut3** | 396-470 | + |
| **NUT8 mut4** | 396-470 | + |
| **NUT9** | 395-482 | + |
| **NUT10** | 468-482 | + |

*GST-pulldown, co-sedimentation or BLI

```
NUT7 wt:    PLALIEELEQEEGLTLAQLVQKRLMALEEEEDAEAPPSFSGAQLD
NUT7 mut1:  PLALIEELEQEAGLTLAQAVQKRLMAAEEEEDAEAPPSFSGAQLD
NUT8 wt:    GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFLADLLSPEKQRDPLALIEELEQEEGLTLAQLVQKRLMALEEEEDAE
NUT8 Mut1:  GQQQEEEGMYPDPGAASAINEAASQKVAASKAEAAAHPQAAADAASPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
NUT8 Mut2:  GQQQEEEGMYPKPGALSKINELGSQKVAGSKVSAGIHGQSLADGSSPEKQRKPLGSIEKSEQKSGGSLAQLVQKRLMALEEEEDAE
NUT8 Mut3:  GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFLADLLSPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
NUT8 Mut4:  GQQQEEEGMYPDPGLLSYINELCSQKVFVSKVEAVIHPQFAADAASPEKQRAPAGAIAAAAQAAGAALAQLVQKRLMALEEEEDAE
```

**Table S2 - HADDOCK docking statistics**

| | |
|---|---|
| **Distance constraints** | |
| Ambiguous interaction restraints (AIRs) | 14 |
| Active residues on NUT7 | 5 |
| Active residues on TAZ2 | 9 |
| Intermolecular NOEs | 2 |
| **Dihedral angle restraints*** | |
| Phi/Psi | 5/5 |
| **Structural statistics** | |
| AIR Violations (mean ± s.d.) | 2±1 |
| Ramachandran analysis | |
|    Residues in the favoured region (%) | 87% |
|    Residues in the additional allowed regions (%) | 10% |
|    Outliers | 3% |
| **Intermolecular energies after water refinement** | |
| $E_{vdw}$ (kcal mol$^{-1}$) | -45 ± 7 |
| $E_{elec}$ (kcal mol$^{-1}$) | -303 ± 32 |
| Buried surface area (Å$^2$) | 1498 ± 88 |

*Dihedral angle restraints derived from chemical shift values using TALOS-N are applied to NUT7 only. For the TAZ2 domain, the structure in PDB 2mzd is used as input to HADDOCK.

**Table S3 - Cryo-EM data collection and map reconstruction**

| | Nucleosome Core Particle |
|---|---|
| **Data collection** | FEI Titan Krios G3 |
| Detector | Gatan K3 direct electron detector |
| Voltage | 300 kV |
| Pixel size (Å) | 1.086 |
| Nominal Magnification | 81,000 |
| Dose rate on Detector (e$^-$ sec$^{-1}$ px$^{-1}$) | 15 |
| Dose rate on specimen (e$^-$ sec$^{-1}$ px$^{-1}$) | 17 |
| Defocus range (μm) | -2.3 to -1.1 |
| **Reconstruction** | |
| Symmetry imposed | C1 |
| Initial particle images (no.) | 80,764 |
| Final particle images (no.) | 33,202 |
| Resolution, FSC=0.143 (Å) | 6.61 |

**Table S4 – Quantification of condensation of BRD4-NUT variants***

| GFP-BRD4-NUT | RFP-p300 positive | RFP-p300 negative |
|---|---|---|
| **WT** | 100/100 | 100/100 |
| **mut1** | 99/100 | 89/100 |
| **mut2** | 100/100 | 92/100 |
| **BD1-BD2** | 16/100 | 0/100 |
| **BD1** | 18/100 | 0/100 |
| **BD2** | 99/100 | 99/100 |
| **K/E** | 21/100 | 3/100 |
| **ΔNPS** | 100/100 | 100/100 |
| **ΔBID** | 100/100 | 9/100 |
| **BD1-BD2-AD** | 99/100 | 1/100 |
| **BD1-BD2-ADmut1** | 0/100 | 0/100 |
| **BD1-BD2-E1A** | 94/100 | 91/100 |

* Foci forming cells/Total cell count

**Table S5 – Quantification of co-localization of BRD4-NUT and p300***

| GFP-BRD4-NUT | Pearson's R value | Co-localization ratio |
|---|---|---|
| **WT** | 0.81±0.01 | 0.90±0.02 |
| **ADmut1** | 0.84±0.01 | 0.87±0.02 |
| **ADmut2** | 0.76±0.02 | 0.90±0.01 |
| **ΔNPS** | 0.76±0.02 | 0.86±0.03 |

* At least 10 cells were analysed (n=10)

**Table S6 – Impact of Drug treatment on condensation**

| GFP-BRD4-NUT | JQ1 | A485 | TSA | CBP30 | 1,6-Hexandiol |
|---|---|---|---|---|---|
| **WT** | No | No | No | Yes | No |
| **ADmut1** | No | No | No | No | No |
| **ADmut2** | No | No | No | No | No |
| **ΔNPS** | No | No | No | Yes | No |

**Table S7 – Quantification of foci formation of after CBP30 treatment***

| GFP-BRD4-NUT | DMSO | CBP30 |
|---|---|---|
| **WT** | 100/100 | 54/100 |
| **ADmut1** | 92/100 | 3/100 |
| **ADmut2** | 91/100 | 2/100 |
| **ΔNPS** | 100/100 | 58/100 |

*Foci-forming cells/Total cell counted