Does ankle push-off correct for errors in anterior-posterior foot placement relative to center-of-mass states?

Jian Jin[1], Jaap H. van Dieën[1], Dinant Kistemaker[1], Andreas Daffertshofer[1,2], Sjoerd M. Bruijn[1,2]

[1] *Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Amsterdam Movement Sciences*, Amsterdam, The Netherlands

[2] *Institute of Brain and Behavior Amsterdam*, Amsterdam, The Netherlands

I.   ABSTRACT

Understanding the mechanisms humans use to stabilize walking is vital for predicting falls in elderly. Modeling studies identified two potential mechanisms to stabilize gait in the anterior-posterior direction: foot placement control and ankle push-off control: Foot placement depends on position and velocity of the center-of-mass (CoM) and push-off modulates with deviations between actual and predicted CoM trajectories. While both control mechanisms have been reported in humans, it is unknown whether especially the latter one is employed in unperturbed steady-state walking. On the one hand, the covariance between CoM states and anterior-posterior foot placement served as a measure of foot placement control. On the other hand, we determined variations in ankle push-off as a function of differences between the actual foot placement and the one predicted from CoM states. We estimated the corresponding correlations and consider them indictors for push-off control based on foot placement errors. We found ankle push-off torque to be correlated to the foot placement error in 30 participants when walking at normal and slow speed, with mean correlation values of up to 0.45. Our study suggests that humans use a push-off strategy for correcting foot placement errors in steady-state walking.

Keywords: foot placement, push-off, gait stability.

List of symbols and abbreviations:

AP anterior-posterior

CoM center-of-mass

CoP center-of-pressure

GRF ground reaction force

II. INTRODUCTION

Walking is a seemingly simple task for most of us. Yet, it involves intricate feedback control and an ongoing integration of sensory inputs and generation of motor outputs (Rossignol et al., 2006; Warren et al., 2001; Zehr & Stein, 1999). A solid understanding of the mechanisms that underlie the stabilization of walking is crucial for identifying causes of falls in elderly (McGibbon, 2003). In-silico studies singled out basic principles of locomotion. Two examples are feedback control of foot placement and of ankle push-off. Both of these mechanisms have been shown to stabilize comparably simple walking models in the presence of perturbations. For instance, Hobbelen & Wisse, (2008) illustrated that modulating ankle push-off torque via feedback from the leading leg's angle substantially improved robustness against perturbations in a sagittal plane flat-feet walker. According to Byl & Tedrake, (2008), feedback-controlled foot placement with a constant push-off magnitude appears more robust than merely modulating the push-off magnitude in a point-feet walker model. In addition, in inverted pendulum models, a combined foot placement and push-off control based on feedback from mid-stance speed demonstrated good robustness (Bhounsule, 2015; Kelly & Ruina, 2015; Zaytsev et al., 2018). Also in human walking, one can find ample evidence that foot placement is based on CoM position and/or velocity. This applies to both steady-state (Redfern & Schumann, 1994; Wang & Srinivasan, 2014) and perturbed walking (Joshi & Srinivasan, 2019; Vlutters et al., 2018b). Moreover, after a perturbation, ankle push-off seems to modulate with deviations of CoM trajectories from the desired trajectories (Afschrift et al., 2021; Fettrow et al., 2019; van Mierlo et al., 2021; Vlutters et al., 2016). Thus far this evidence is restricted to perturbed walking, but we here subsume that humans also use a feedback-controlled push-off strategy to stabilize steady-state walking.

In human walking, foot placement along the anterior-posterior (AP) direction is correlated to CoM states, i.e., CoM position and velocity. Wang & Srinivasan, (2014) showed that CoM states at mid-stance and heel strike predict about 33% of the variance in AP foot placement. Recently, Liu, (2021) replicated this finding in steady-state walking and additionally in perturbed walking. This study by Liu revealed that a different foot placement model is needed to describe responses to slip-like and trip-like treadmill perturbations, suggesting that the control of foot placement may be different between steady-state and perturbed walking. By the same token, ankle push-off during the double stance phase seems to modulate with AP pelvis perturbations: van Mierlo et al., (2021) found adjustments of the double stance duration and the AP center-of-pressure (CoP) trajectory to correct perturbed CoM states towards desired states at the end of the double stance phase; and, Afschrift et al., (2021) demonstrated that for AP pelvis perturbations and for treadmill belt speed perturbations, the deviations in CoM states from the steady-state trajectory accurately predict the ankle moment after a neural

delay of about 100 ms. In steady-state walking, the percentage of explained variance of AP foot placement from the CoM states increases from 33% to 70% during the double stance phase (Wang & Srinivasan, 2014). That is, there may be modulations of the CoM states through ankle push-off control.

Since CoM states covary with AP foot placement, neither of them can be isolated as input to push-off control. A more appropriate feedback signal would be the *foot placement error*, first introduced by van Leeuwen and co-workers (2021). It is defined as the difference between actual foot placement and the ideal foot placement based on the CoM states at heel strike. This can be estimated by linearly regressing CoM states with foot placement (Wang & Srinivasan, 2014). As such, the covariance between CoM states and foot placement has been suggested to subserve gait stability both in simple walking models (Hof, 2008; Joshi & Srinivasan, 2019; Maki & McIlroy, 1999; Townsend, 1985) and in humans (Joshi & Srinivasan, 2019; Redfern & Schumann, 1994; Verrel et al., 2010; Wang & Srinivasan, 2014). Assuming the presence of foot placement control, deviations from this covariance can be viewed as foot placement *errors* that need to be corrected.

Fig. 1. Illustration of positive and negative foot placement errors. For certain CoM states (CoM AP position and velocity at heel strike), a too large step yields a positive foot placement error, while a too small step leads to a negative foot placement error. A vanishing foot placement error implies linear predictions of foot placement from CoM states.

As shown in Figure 1, a positive foot placement error in the AP direction indicates that foot is placed more forward with respect to a linear prediction from the CoM states. In this case, there is a higher collision loss due to larger step length (Adamczyk & Kuo, 2009), and the walker's post-collision total energy will be lower, requiring a stronger push-off to compensate for this larger collision loss. Here, we assumed the collision loss to be balanced by the push-off work on the CoM during the double stance phase, as observed in human walking (Kim & Park, 2012). We hence expected the push-off force/torque magnitude to be proportional to the AP foot placement error, i.e., a stronger push-off for larger positive foot placement errors, and a weaker push-off for more negative foot placement errors.

We examined push-off modulations by investigating how kinetic time series are modulated during the double stance phase as a function of the foot placement error. Next, we evaluated the correlations between foot placement errors and three kinetic time series: the combined AP ground reaction forces (GRFs), the trailing leg's AP GRFs, and the ankle moments. For every participant, we also compared the mean kinetics of ten most positive foot placement errors with the mean kinetics of ten most negative foot placement errors to study how the push-off may differ when correcting positive and negative foot placement errors. We selected the combined AP GRF because it represents the summed effect of ankle push-off by the trailing leg and braking by the leading leg. The use of the trailing leg's AP GRF was motivated by the idea that the CoM is mainly accelerated via the trailing leg's AP GRF (Hernández et al., 2009). The use of ankle moment was motivated by modeling studies that indicated ankle push-off to stabilize gait (Hobbelen & Wisse, 2008; Kerimoglu et al., 2021; Kim & Collins, 2017) and to provide the bulk of the trailing leg's propulsive force (Franz & Kram, 2014). This is due to a higher gain from ankle moments to the trailing leg's AP GRF compared to hip or knee moments (Biewener et al., 2004; Toney & Chang, 2016).

III. METHODS

We used existing data from steady-state normal and slow treadmill walking experiments (van Leeuwen et al., 2020a & 2020b). Below, we briefly sketch data collection and processing but we refer to van Leeuwen et al., (2020a) for more details. The data can be found at https://doi.org/10.5281/zenodo.4229851 and the code can be found at https://drive.google.com/drive/folders/1AnL9ajMk_Q3Gwzv_OrOp7A8DNwS8lniC?usp=sharing.

Participants. 30 participants were included (19 female, 11 male, age 30 ± 7 years, weight 70 ± 13 kg, height 1.73 ± 0.08 m; mean $\pm$ sd). None of them reported injuries or balance issues that could affect their gait pattern. All participants signed informed consent before the experiment. Ethical approval (VCWE-2018-159) had been granted by the ethics board of the Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, prior to conducting the experiment.

Protocol. Every participant walked on a treadmill at a constant belt speed of $1.25 \cdot \sqrt{\text{leg length}}$ m/s (*normal* walking condition) and $0.63 \cdot \sqrt{\text{leg length}}$ m/s (*slow* walking condition). A metronome served to impose stride frequency to minimize stride frequency variations. Participants were asked to match their right heel

strikes to the beat. The imposed frequency was customized as the average preferred stride frequency during the last hundred steps of the familiarization trial at each speed. The normal walking trials lasted five and the slow walking trials ten minutes each to ensure that data of at least 200 consecutive strides were collected. Normal and slow speed trials were randomized in order and separated by sufficient breaks to prevent fatigue.

Data collection and processing. Participants walked on an instrumented dual-belt treadmill (Motek-Forcelink, Amsterdam, Netherlands). Full-body kinematics were recorded using an active 3D motion analysis system and cluster markers on all segments (Optotrak, Northern Digital Inc, Waterloo ON, Canada). For every participant and condition, we analyzed the last 200 consecutive strides without data quality issues (e.g., limited marker visibility, large noise, etc.). GRF data were collected using a force plate integrated in the treadmill. Gait events (heel strike and toe-off) were detected based on the so-called "butterfly pattern" of the combined CoP trajectory. This method, validated by Roerdink et al., (2008), has the advantage that it does not require the setting of an ad-hoc threshold. Ankle moments in the sagittal plane were computed following the approach outlined in Hof, (1992).

A left (right) stride was defined from left (right) heel strike to left (right) heel strike and a left (right) step from left (right) toe-off to right (left) toe-off. Ankle joint moments were calculated using the GRFs and the lower body kinematics via a bottom-up inverse dynamics approach (Kingma et al., 1996). For subsequent analysis, the kinematic and kinetic time series of a stride and a step were time-normalized. We accounted for the variations in timing of heel strike and subsequent push-off within a step[1] by segmenting the time window of a step to single-stance and double stance sub-windows with fixed time length per walking condition.

Linear foot placement model. Similar to previous studies (van Leeuwen et al., 2020a; Wang & Srinivasan, 2014), we fitted a linear model between the CoM states and AP foot placement. This full body CoM state was derived from a weighted sum of the body segments' CoM, which in turn was estimated from the percent longitudinal distances of the body segments' CoM to neighboring bony landmarks, relative to the lengths of respective segments (De Leva, 1996). The predictor CoM states included the AP position (the horizontal distance from stance foot to CoM) and the AP velocity, i.e., COM_{pos} and COM_{vel}, respectively. The predictor and dependent variables were de-meaned prior to regression to ensure a zero intercept. The model reads as follows:

---

[1] The total samples of time window for the single stance phase, double stance phase in normal and slow walking were different on average (over all steps from all participants).

$$FP = \beta_{\text{pos}}(i) \cdot \text{CoM}_{\text{pos}}(i) + \beta_{\text{vel}}(i) \cdot \text{CoM}_{\text{vel}}(i) + \varepsilon_{\text{FP}}(i), \qquad (1)$$

where FP represents the (de-meaned) AP distance between the trailing and leading leg, $\beta_{\text{pos}}(i)$ and $\beta_{\text{vel}}(i)$ are the phase-specific regression coefficients obtained from the least square fit, and $\varepsilon_{\text{FP}}(i)$ denotes the residual from the linear regression, or, in short, the foot placement error. The index "i" indicates the time instant in a step (0-100% from heel strike to contralateral heel strike). We used the time instant at heel strike (100%) to compute the foot placement FP and the foot placement error $\varepsilon_{\text{FP}}(i = \text{heel strike})$, which we assume to be corrected by the subsequent push-off.

Correction of foot placement error. Using regression analysis, we evaluated how well the foot placement error at heel strike correlated to the combined AP GRF, the trailing leg's AP GRF, and the trailing leg's ankle moment during the double stance. The general model relating foot placement errors and the subsequent kinetic time series (indicated by F) was set to:

$$F(i + 1) = \beta_{\varepsilon_{\text{FP}}}(i + 1) \cdot \varepsilon_{\text{FP}}(i = \text{heel strike}) + \varepsilon(i + 1), \qquad (2)$$

where $\beta_{\varepsilon_{\text{FP}}}(i + 1)$ is the phase-specific linear regression coefficient at phase i in the next step, and $\varepsilon(i + 1)$ is the residual. We assume the resulting correlations to suggest feedback control. Yet, we have to admit that correlations may also arise from passive dynamics (Patil et al., 2019) or from intrinsic muscle properties (e.g., preflexes stabilizing movements through nonlinear viscoelastic properties of muscles when in contact; cf. Loeb, 1995). We return to this in the Discussion section.

In addition to the regression analysis, we compared the kinetic time series of the "most positive" with that of the "most negative" foot placement errors. Specifically, the foot placement errors of each participant (200 strides) were sorted from low to high, and the ten most positive and ten most negative foot placement error strides were analyzed, by extracting the mean kinetic time series from these sets of ten strides.

Statistics. All statistical tests and analysis were performed in Matlab (v2017b, The Mathworks Inc., Natick, MA), including the multi-linear regression of the foot placement model. Correlation values are presented as group level means and individual data. The mean correlations were obtained by 1) first applying Fisher's r-to-z transform to guarantee the normality of the correlation coefficients; 2) then averaging the z-values over all participants per condition; and, 3), inverting the Fisher transform to obtain the group mean correlation

coefficients per condition. We tested the correlation coefficients for significance using statistical parametric mapping (Friston et al., 2007) as implemented in the SPM1D toolbox (https://spm1d.org/). In brief, SPM1D allows for statistical testing of time series data, considering the interdependence (smoothness) between time samples. To test for differences between the kinetic time series of the "most positive" and "most negative" foot placement errors, we also performed an SPM1D paired t-test. The significance level of every test was set to α=0.05.

IV. RESULTS

The variance in AP foot placement explained by the CoM states (R^2) increased over the single stance phase (Fig. 2, left of the dashed vertical line) and reached values between 50% and 80% during the double stance phase in both normal and slow walking (Fig. 2, right of the dashed vertical line).

Fig. 2. Proportion of foot placement variance explained by CoM states, for normal (a) and slow walking (b). Every curve represents the explained variance for a participant. The dashed vertical lines correspond to foot placement, and the phases after heel strike correspond to the double stance phase.

Although a large proportion of foot placement variance could be explained by the CoM states at heel strike, a fair amount of unexplained variance remained, i.e., there were substantial foot placement errors. The distributions of foot placement errors across all participants for normal and slow walking were close to Gaussian with zero mean (Fig. 3a). The variance of foot placement errors in slow walking was larger than in normal walking (Fig. 3a). The group means of the 10 most positive and 10 most negative foot placement errors in slow walking had larger magnitudes compared to those in normal walking (Fig. 3b).

Fig. 3. (a) Distributions of foot placement errors for normal and slow walking for all participants. Gaussian distributions are fitted to the data in two conditions. (b) The group mean values of the averaged 10 most positive and 10 most negative foot placement errors per participant. Error bars indicate the standard deviation across participants.

Across participants the foot placement errors were significantly correlated with the combined AP GRF during the double stance phase in normal and slow walking (mean correlations up to 0.34 and 0.28, respectively; see Fig. 4a). High positive correlations occurred in the first half of the double stance phase and decreased to negative correlations over the latter half of the double stance phase. Correlations were close to zero prior to foot placement, which implies that foot placement can hardly be predicted based on the combined AP GRF prior to foot placement. Correlations were also close to zero at foot placement. This suggests that preemptive push-off indicated by the combined AP GRF did not cause foot placement errors. Correlations were significant and negative at the end of the stride cycle during normal walking (before the subsequent heel strike, mean values reached -0.43, Fig. 4a). See the *Supplementary Material* S2 for further details on the regression.

Likewise, the trailing leg's AP GRF displayed significant correlations with foot placement error at heel strike in normal and in slow walking (mean correlations up to 0.45 and 0.41, respectively; see Fig. 4b and *Supplementary Material* S3 for the corresponding statistical tests). Correlations were highest in the middle of the double stance phase, i.e., push-off modulations occurred mostly around this phase. Correlations were close to zero at the beginning and at the end of the double stance phase, suggesting that preemptive, early and late push-off did not influence foot placement errors.

Fig. 4. Correlations between foot placement errors at heel strike (left dash-dotted and dashed lines) and different kinetic time series in normal and slow walking. (a) combined AP GRF; (b) trailing leg's AP GRF; (c) ankle moment. The thin curves are correlations per participant and per foot, the thick ones are means over participants. The phases between the dash-dotted lines or the dashed lines correspond to the double stance phase in normal and slow walking, respectively. Note that the trailing leg's AP GRF and ankle moment were zero during the swing phases at the end of the stride cycle, thus the correlations were not shown.

The ankle moments during the double stance were significantly correlated with foot placement errors at heel strike in normal and slow walking (mean correlations up to 0.39 and 0.37, respectively; see Fig. 4c and *Supplementary Material* S4). Correlations with ankle moments were also highest in the middle of the double stance phase and were positive in the beginning of the double stance phase.

Fig. 5. Averaged kinetic time series corresponding to the "most positive" and "most negative" foot placement errors in normal and slow walking. (a) combined AP GRF; (b) trailing leg's AP GRF; (c) ankle moment. The phases between the dash-dotted lines or the dashed lines correspond to the double stance phase in normal and slow walking, respectively.

In line with the correlations, all the kinetic time series during the double stance phase were significantly larger for "most positive" foot placement errors than for "most negative" foot placement errors in both walking conditions (Figs. 5a-c and *Supplementary Material* S5). As shown in Figs. 4a-c, the combined AP GRF was higher in the first half and lower in the end of the double stance phase for "most positive" foot placement

errors; the trailing leg's AP GRF and ankle moment were higher in the middle of the double stance phase for "most positive" foot placement errors.

V. DISCUSSION

We sought to test the hypothesis that humans employ ankle push-off control to compensate for errors in AP foot placement during steady-state walking. Our correlations between combined and trailing leg's AP GRF and ankle moment with foot placement errors at heel strike appear to confirm this. This does not come as a surprise, as several modeling studies demonstrated the capacity of ankle push-off modulations in improving robustness against perturbations (Hobbelen & Wisse, 2008; Kim & Collins, 2013, 2017; Stephens & Atkeson, 2009). Moreover, correlations between push-off modulations and deviations in CoM trajectories from AP perturbations have already been found in perturbed human walking (Afschrift et al., 2021; Rafiee & Kiemel, 2020; van Mierlo et al., 2021; Vlutters et al., 2016). Such correlations arguably arise from feedback mechanisms. Recently, van Leeuwen et al., (2021) reported that errors in medio-lateral foot placement are corrected by ankle moments during the subsequent stance phase. We here supplemented these findings by showing evidence for a push-off mechanism used to correct for errors in foot placement during steady-state walking in the AP direction.

In steady-state normal walking, we found maximum correlations between foot placement error and trailing leg's AP GRF (up to 0.45), followed by correlations with ankle moment (up to 0.39), and with combined AP GRF (up to 0.34). Feedback control based on foot placement error may primarily serve for control of the push-off force from the trailing leg, which is strongly determined by the ankle moment. Push-off can be generated by ankle, knee, and hip muscles (Kuo, 2002). At the instant of maximum trailing leg power, the gain from the ankle moment to the trailing leg's AP GRF is ten times larger than the gain from the hip joint moment (Toney & Chang, 2016). It is therefore not unexpected that correlations for the ankle moment are close to the correlations for the trailing leg's AP GRF.

Peaks in the correlations between foot placement errors and kinetic variables occurred around the middle of the double stance phase (Figs. 4a-c), while the peak combined/trailing leg's AP GRF and the peak ankle moment all occurred already in the beginning of the double stance phase (Fig. 5). Both observations imply that modulations of the CoM states are less prevalent during the peak combined/trailing leg's AP GRF or peak ankle moment, but mostly around the middle of the double stance phase, at which time the CoM AP velocity

and ankle power is highest (Hernández et al., 2009; Kuhman & Hurt, 2019; Lipfert et al., 2014). Modulations of the CoM states may be achieved either by adjusting the trailing leg's AP GRF and ankle moment magnitude (e.g., Fig. 5c), or by modulating their timings (duration of the double stance phase and relative duration of particular force/torque in the double stance phase) (Kuhman & Hurt, 2019; Toney & Chang, 2016; van Mierlo et al., 2021; Williams & Martin, 2019). For instance, in slow walking the peak in combined AP GRF occurs later in too large foot placement error steps than in too small foot placement error steps (Fig. 5b), which may imply a prolonged activation of the muscles contributing to this force. Yet, the prolonged activation mechanism in slow walking cannot be directly verified or falsified from Figs. 4 and 5 due to the normalization of the double stance phase to a fixed window. The time delay between heel strike and the peak in the correlations between foot placement errors and ankle moment in normal walking was about 80 ms. Feedback delay in control of human ankle moments is arguably longer ($\approx$ 100 ms) due to signal transmission, sensory integration in the nervous system and the electromechanical delay (Afschrift et al., 2021; Welch & Ting, 2008). Rather than directly using foot placement error at heel strike, humans might predict their foot placement error shortly before heel strike as input for the (ankle) push-off modulation. The correlations between the CoM states and the AP foot placement shortly before heel strike were close to the ones at heel strike (Fig. 2). This implies a largely similar foot placement accuracy when predicted prior to or at foot placement. Whether and how humans estimate foot placement error around foot placement, however, requires further investigation.

The positive correlations between foot placement errors and ankle moment at initial double stance phase in normal speed (Fig. 4c) suggest limited ankle push-off modulations before contralateral heel strike. The correlations for ankle moment were positive throughout the double stance phase except at the end, and peak correlations did not appear until the middle of the double stance phase (Fig. 4c). Our findings are consistent with the study by Kuhman & Hurt, (2019) who found that at human preferred steady-state walking speed (around 1.4 m/s), the peak trailing leg's AP GRF, the onset and peak of trailing leg power and the peak ankle power all occurred after contralateral heel strike. Humans seemingly rely less on the more energy efficient preemptive push-off strategy to stabilize gait than suggested by modeling studies (Kuo, 2002; Ruina et al., 2005). Potentially, they opt for producing push-off somewhat later to incorporate sensory input information about foot placement error.

We showed that steady-state walking relies on more or less the same push-off strategy as perturbed walking, where a forward push led to more forward foot placement (Vlutters et al., 2018b), followed by CoP modulation (shorter double stance duration and longer CoP distance travelled, see van Mierlo et al., 2021) from either

weight shift of the leading leg (Hof, 2007) or plantarflexion ankle moment modulations (Gruben & Boehm, 2014; Vlutters et al., 2018b), contributing to a reduction of the CoM AP velocity over the double stance phase. van Mierlo et al., (2021) found CoP modulation to be absent for backward perturbations, presumably because adjustments in foot placement sufficiently reduced the foot placement error (see Fig. 5 from Vlutters et al., 2018b). According to Vlutters et al., (2018a), changing magnitude and direction of AP perturbations on the pelvis near foot contact hardly affected the AP foot placement in both the first and second recovery step. This suggests that a limited response time for foot placement modulation primarily calls for push-off modulation. Apparently, humans cope with perturbations by complementing foot placement with push-off control to correct the remaining foot placement errors.

Older adults have poorer foot placement accuracy, at least in the medio-lateral direction (Arvin et al., 2018). They have a reduced ankle push-off power generation (Franz, 2016; Hernández et al., 2009). One may therefore expect the older adults have poorer gait robustness due to larger foot placement errors and a reduced capacity to correct for these. Future work should compare age-related differences in foot placement error at the beginning and the end of the double stance phase and relate them to differences in push-off modulation. This can further clarify gait adaptations in the elderly, such as the distal-to-proximal redistribution of push-off power production (Franz, 2016).

We interpreted the modulation of the push-off force by the trailing leg in terms of gait stability. Admittedly, the modulation of push-off force might also be explained by speed regulation. On a treadmill, speed is controlled from step to step and may results in an over-correction of speed errors with respect to the treadmill speed (Dingwell & Cusumano, 2015). When walking at different steady-state speeds, humans can adjust the relative duration of the double stance phase (cf. the percentage for normal and slow walking in Fig. 4), which may lead to changes in magnitudes of AP GRF (Williams & Martin, 2019). Here, we also observed over-correction of foot placement errors by the combined AP GRF at the subsequent step (see the end of the stride cycle in Fig. 4a and Fig. 5a). The foot placement error is determined by two variables: CoM states and foot placement at heel strike. For an inverted pendulum model walking at a nominal speed, perturbations leading to positive foot placement errors reduced walking speed and perturbations leading to negative foot placement errors increased walking speed (Hof, 2008). Corrections of foot placement errors probably also correct speed errors with respect to the treadmill. Our current data do not allow for directly distinguishing whether maintaining gait stability or constant speed were the *primary* control goal. Either way, it is likely to be an improper dichotomy given their interdependence (Hof, 2008).

Our study suggests that push-off control corrects foot placement errors in steady-state walking. Push-off, however, also contributes to the leg swing (Zelik & Adamczyk, 2016) and to the foot trajectory in the early swing phase. This is evident from humans' responses to stepping target perturbations (Barton et al., 2019). Contrary to push-off control, swing leg control for foot placement is largely passive except at the beginning and end of the swing phase, as indicated by quick bursts of leg swing and retraction impulses (Doke et al., 2005). This passive ("predictable") pendular dynamics in the swing phase may hence be exploited by the feedforward/anticipative control of push-off for the desired foot placement. It is currently unknown whether in steady-state walking feedforward control of push-off is also used. As such is remains opaque whether push-off corrects foot placement error or push-off plans for the next foot placement or both. To further unravel the stabilizing mechanisms of push-off control, modeling approaches can elucidate when implementing feedforward and/or feedback push-off controllers based on foot placement errors for simple bipedal walkers under noisy conditions. Ryu & Kuo, (2021) showed pure feedback and pure feedforward control to be susceptible to sensor noise and process noise (e.g., uncertainly in the environment), respectively, and the combined feedforward and feedback controller was best for stable and robust walking.

Limitations.

Our study comes with several limitations. Before listing them, however, we want to repeat our statement-of-warning, namely, that the correlations we observed should be interpreted with care. Next to feedback control there might be other sources causing them, such as passive dynamics (Patil et al., 2019). Future studies may employ (perceptual or mechanical) perturbations (Fettrow et al., 2019; Roden-Reynolds et al., 2015) and/or EMG responses (Rankin et al., 2014; van Leeuwen et al., 2021; Vlutters et al., 2019) to disentangle contributions from feedback control and from passive dynamics.

We did not apply any normalization, to render our variables of interest easier to interpret. Admittedly, normalization can reduce inter-participant variability. For this, however, we customized each participant's treadmill speed, i.e., we maintained constant dimensionless speeds (Hof, 1996). Although this is a valid approach to account for participants' leg differences, controlling speeds and additionally stride frequency by metronome may cause our results to not directly be generalizable to natural walking. Since people seem to prefer walking at a fixed speed and stride length to minimize the metabolic cost of transport (Bertram, 2005; Bertram & Ruina, 2001; Donelan et al., 2001; Selinger et al., 2015), however, we expect the same push-off mechanisms to also hold for natural walking.

We observed an outlier with large negative correlations for the ankle moments only in slow walking in Figure 4c. This participant's right ankle moments were negatively correlated to (left) foot placement errors, and the foot placement of the right leg was consistently further forward than of the left leg. We suspect this participant used a different and uncommon walking strategy in slow walking. And there was another participant that might be considered an outlier when considering the correlations between the left trailing leg's AP GRF and right foot placement errors in slow walking in Figure 4b. However, this participant showed different behavior only prior to heel strike.

VI. CONCLUSION

We correlated foot placement errors with kinetic time series of ankle push-off to test the idea that foot placement errors are corrected for by push-off in steady-state walking. We found ankle push-off torque to be correlated to AP foot placement errors mainly around the middle of the double stance phase both in steady-state normal and slow walking, confirming our hypothesis that humans employ ankle push-off control to compensate for errors in AP foot placement during steady-state walking.

VII. ACKNOWLEDGEMENTS

S.M.B. and J.J. are funded by a VIDI grant no. (016.Vidi.178.014) from the Dutch Organization for Scientific Research (NWO).

VIII. REFERENCES

Adamczyk, P. G., & Kuo, A. D. (2009). Redirection of center-of-mass velocity during the step-to-step transition of human walking. *Journal of Experimental Biology*, *212*(16). https://doi.org/10.1242/jeb.027581

Afschrift, M., de Groote, F., & Jonkers, I. (2021). Similar sensorimotor transformations control balance during standing and walking. *PLoS Computational Biology*, *17*(6). https://doi.org/10.1371/journal.pcbi.1008369

Arvin, M., Hoozemans, M. J. M., Pijnappels, M., Duysens, J., Verschueren, S. M., & van Dieën, J. H. (2018). Where to step? Contributions of stance leg muscle spindle afference to planning of mediolateral foot placement for balance control in young and old adults. *Frontiers in Physiology*, *9*(1134). https://doi.org/10.3389/fphys.2018.01134

Barton, S. L., Matthis, J. S., & Fajen, B. R. (2019). Control strategies for rapid, visually guided adjustments of the foot during continuous walking. *Experimental Brain Research*, *237*(7). https://doi.org/10.1007/s00221-019-05538-7

Bertram, J. E. A. (2005). Constrained optimization in human walking: Cost minimization and gait plasticity. *Journal of Experimental Biology*, *208*(6). https://doi.org/10.1242/jeb.01498

Bertram, J. E. A., & Ruina, A. (2001). Multiple walking speed-frequency relations are predicted by constrained optimization. *Journal of Theoretical Biology*, *209*(4). https://doi.org/10.1006/jtbi.2001.2279

Bhounsule, P. A. (2015). Control of a compass gait walker based on energy regulation using ankle push-off and foot placement. *Robotica*, *33*(6). https://doi.org/10.1017/S0263574714000745

Biewener, A. A., Farley, C. T., Roberts, T. J., & Temaner, M. (2004). Muscle mechanical advantage of human walking and running:

Implications for energy cost. *Journal of Applied Physiology*, *97*(6). https://doi.org/10.1152/japplphysiol.00003.2004

Byl, K., & Tedrake, R. (2008). Approximate optimal control of the compass gait on rough terrain. In *Proceedings - IEEE International Conference on Robotics and Automation* (pp. 1258–1263). https://doi.org/10.1109/ROBOT.2008.4543376

De Leva, P. (1996). Adjustments to zatsiorsky-seluyanov's segment inertia parameters. *Journal of Biomechanics*, *29*(9). https://doi.org/10.1016/0021-9290(95)00178-6

Dingwell, J. B., & Cusumano, J. P. (2015). Identifying stride-to-stride control strategies in human treadmill walking. *PLoS ONE*, *10*(4). https://doi.org/10.1371/journal.pone.0124879

Doke, J., Donelan, J. M., & Kuo, A. D. (2005). Mechanics and energetics of swinging the human leg. *Journal of Experimental Biology*, *208*(3). https://doi.org/10.1242/jeb.01408

Donelan, J. M., Kram, R., & Kuo, A. D. (2001). Mechanical and metabolic determinants of the preferred step width in human walking. *Proceedings of the Royal Society B: Biological Sciences*, *268*(1480). https://doi.org/10.1098/rspb.2001.1761

Fettrow, T., Reimann, H., Grenet, D., Thompson, E., Crenshaw, J., Higginson, J., & Jeka, J. (2019). Interdependence of balance mechanisms during bipedal locomotion. *PLoS ONE*, *14*(12). https://doi.org/10.1371/journal.pone.0225902

Franz, J. R. (2016). The Age-Associated Reduction in Propulsive Power Generation in Walking. *Exercise and Sport Sciences Reviews*, *44*(4). https://doi.org/10.1249/JES.0000000000000086

Franz, J. R., & Kram, R. (2014). Advanced age and the mechanics of uphill walking: A joint-level, inverse dynamic analysis. *Gait and Posture*, *39*(1). https://doi.org/10.1016/j.gaitpost.2013.06.012

Friston, K. J., Ashburner, J. T., Kiebel, S. J., Nichols, T. E., & Penny, W. D. (2007). *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. *Statistical Parametric Mapping The Analysis of Functional Brain Images* (Vol. 8).

Gruben, K. G., & Boehm, W. L. (2014). Ankle torque control that shifts the center of pressure from heel to toe contributes non-zero sagittal plane angular momentum during human walking. *Journal of Biomechanics*, *47*(6). https://doi.org/10.1016/j.jbiomech.2014.01.034

Hernández, A., Silder, A., Heiderscheit, B. C., & Thelen, D. G. (2009). Effect of age on center of mass motion during human walking. *Gait and Posture*, *30*(2). https://doi.org/10.1016/j.gaitpost.2009.05.006

Hobbelen, D. G. E., & Wisse, M. (2008). Ankle actuation for limit cycle walkers. *International Journal of Robotics Research*, *27*(6). https://doi.org/10.1177/0278364908091365

Hof, A. L. (1992). An explicit expression for the moment in multibody systems. *Journal of Biomechanics*, *25*(10). https://doi.org/10.1016/0021-9290(92)90076-D

Hof, A. L. (1996). Scaling gait data to body size. *Gait and Posture*. https://doi.org/10.1016/0966-6362(95)01057-2

Hof, A. L. (2007). The equations of motion for a standing human reveal three mechanisms for balance. *Journal of Biomechanics*, *40*(2). https://doi.org/10.1016/j.jbiomech.2005.12.016

Hof, A. L. (2008). The "extrapolated center of mass" concept suggests a simple control of balance in walking. *Human Movement Science*, *27*(1). https://doi.org/10.1016/j.humov.2007.08.003

Joshi, V., & Srinivasan, M. (2019). A controller for walking derived from how humans recover from perturbations. *Journal of the Royal Society Interface*, *16*(157). https://doi.org/10.1098/rsif.2019.0027

Kelly, M., & Ruina, A. (2015). Non-linear robust control for inverted-pendulum 2D walking. In *Proceedings - IEEE International Conference on Robotics and Automation* (Vol. 2015-June). https://doi.org/10.1109/ICRA.2015.7139800

Kerimoglu, D., Karkoub, M., Ismail, U., Morgul, O., & Saranli, U. (2021). Efficient bipedal locomotion on rough terrain via compliant ankle actuation with energy regulation. *Bioinspiration & Biomimetics*, *16*(5). https://doi.org/10.1088/1748-3190/ac13b1

Kim, M., & Collins, S. H. (2013). Stabilization of a three-dimensional limit cycle walking model through step-to-step ankle control. In *IEEE International Conference on Rehabilitation Robotics* (pp. 1–6). https://doi.org/10.1109/ICORR.2013.6650437

Kim, M., & Collins, S. H. (2017). Once-Per-Step Control of Ankle Push-Off Work Improves Balance in a Three-Dimensional Simulation of Bipedal Walking. *IEEE Transactions on Robotics*, *33*(2). https://doi.org/10.1109/TRO.2016.2636297

Kim, S., & Park, S. (2012). The oscillatory behavior of the CoM facilitates mechanical energy balance between push-off and heel strike. *Journal of Biomechanics*, *45*(2). https://doi.org/10.1016/j.jbiomech.2011.10.009

Kingma, I., De Looze, M. P., Toussaint, H. M., Klijnsma, H. G., & Bruijnen, T. B. M. (1996). Validation of a full body 3-D dynamic linked segment model. *Human Movement Science*, *15*(6). https://doi.org/10.1016/S0167-9457(96)00034-6

Kuhman, D., & Hurt, C. P. (2019). The timing of locomotor propulsion in healthy adults walking at multiple speeds. *Human Movement Science*, *68*. https://doi.org/10.1016/j.humov.2019.102524

Kuo, A. D. (2002). Energetics of actively powered locomotion using the simplest walking model. *Journal of Biomechanical Engineering*, *124*(1). https://doi.org/10.1115/1.1427703

Lipfert, S. W., Günther, M., Renjewski, D., & Seyfarth, A. (2014). Impulsive ankle push-off powers leg swing in human walking. *Journal of Experimental Biology*, *217*(8). https://doi.org/10.1242/jeb.097345

Liu, C. (2021). *Understanding reactive balance control strategies in non-disabled and post-stroke gait*. University of Southern California.

Loeb, G. E. (1995). Control implications of musculoskeletal mechanics. In *Annual International Conference of the IEEE*

Engineering in Medicine and Biology - Proceedings (Vol. 17). https://doi.org/10.1109/iembs.1995.579743

Maki, B. E., & McIlroy, W. E. (1999). The control of foot placement during compensatory stepping reactions: Does speed of response take precedence over stability? *IEEE Transactions on Rehabilitation Engineering*, *7*(1). https://doi.org/10.1109/86.750556

McGibbon, C. A. (2003). Toward a better understanding of gait changes with age and disablement: Neuromuscular adaptation. *Exercise and Sport Sciences Reviews*, *31*(2). https://doi.org/10.1097/00003677-200304000-00009

Patil, N. S., Dingwell, J. B., & Cusumano, J. P. (2019). Correlations of pelvis state to foot placement do not imply within-step active control. *Journal of Biomechanics*, *97*. https://doi.org/10.1016/j.jbiomech.2019.109375

Rafiee, S., & Kiemel, T. (2020). Multiple strategies to correct errors in foot placement and control speed in human walking. *Experimental Brain Research*, *238*(12). https://doi.org/10.1007/s00221-020-05949-x

Rankin, B. L., Buffo, S. K., & Dean, J. C. (2014). A neuromechanical strategy for mediolateral foot placement in walking humans. *Journal of Neurophysiology*, *112*(2). https://doi.org/10.1152/jn.00138.2014

Redfern, M. S., & Schumann, T. (1994). A model of foot placement during gait. *Journal of Biomechanics*, *27*(11). https://doi.org/10.1016/0021-9290(94)90043-4

Roden-Reynolds, D. C., Walker, M. H., Wasserman, C. R., & Dean, J. C. (2015). Hip proprioceptive feedback influences the control of mediolateral stability during human walking. *Journal of Neurophysiology*, *114*(4). https://doi.org/10.1152/jn.00551.2015

Roerdink, M., Coolen, B. (H)., Clairbois, B. (H)E, Lamoth, C. J. C., & Beek, P. J. (2008). Online gait event detection using a large force platform embedded in a treadmill. *Journal of Biomechanics*, *41*(12). https://doi.org/10.1016/j.jbiomech.2008.06.023

Rossignol, S., Dubuc, R., & Gossard, J. P. (2006). Dynamic sensorimotor interactions in locomotion. *Physiological Reviews*, *86*(1). https://doi.org/10.1152/physrev.00028.2005

Ruina, A., Bertram, J. E. A., & Srinivasan, M. (2005). A collisional model of the energetic cost of support work qualitatively explains leg sequencing in walking and galloping, pseudo-elastic leg behavior in running and the walk-to-run transition. *Journal of Theoretical Biology*, *237*(2). https://doi.org/10.1016/j.jtbi.2005.04.004

Ryu, H. X., & Kuo, A. D. (2021). An optimality principle for locomotor central pattern generators. *Scientific Reports*, *11*(1). https://doi.org/10.1038/s41598-021-91714-1

Selinger, J. C., O'Connor, S. M., Wong, J. D., & Donelan, J. M. (2015). Humans Can Continuously Optimize Energetic Cost during Walking. *Current Biology*, *25*(18). https://doi.org/10.1016/j.cub.2015.08.016

Stephens, B., & Atkeson, C. (2009). Modeling and control of periodic humanoid balance using the linear biped model. In *9th IEEE-RAS International Conference on Humanoid Robots, HUMANOIDS09*. https://doi.org/10.1109/ICHR.2009.5379605

Toney, M. E., & Chang, Y. H. (2016). The motor and the brake of the trailing leg in human walking: leg force control through ankle modulation and knee covariance. *Experimental Brain Research*, *234*(10). https://doi.org/10.1007/s00221-016-4703-8

Townsend, M. A. (1985). Biped gait stabilization via foot placement. *Journal of Biomechanics*, *18*(1). https://doi.org/10.1016/0021-9290(85)90042-9

van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A., & Bruijn, S. M. (2020a). Active foot placement control ensures stable gait: Effect of constraints on foot placement and ankle moments. *Plos One*, *15*(12). https://doi.org/10.1371/journal.pone.0242215

van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A., & Bruijn, S. M. (2021). Ankle muscles drive mediolateral center of pressure control to ensure stable steady state gait. *Scientific Reports*, *11*(1). https://doi.org/10.1038/s41598-021-00463-8

van Leeuwen, M., Bruijn, S. M., van Dieën, J. H., & Daffertshofer, A. (2020b). Dataset and analyses for: Active foot placement control ensures stable gait: Effect of constraints on foot placement and ankle moments. Zenodo. https://doi.org/https://doi.org/10.5281/zenodo.4229851

van Mierlo, M., Vlutters, M., van Asseldonk, E. H. F., & van der Kooij, H. (2021). Centre of pressure modulations in double support effectively counteract anteroposterior perturbations during gait. *Journal of Biomechanics*, *126*. https://doi.org/10.1016/j.jbiomech.2021.110637

Verrel, J., Lövdén, M., & Lindenberger, U. (2010). Motor-equivalent covariation stabilizes step parameters and center of mass position during treadmill walking. *Experimental Brain Research*, *207*(1–2). https://doi.org/10.1007/s00221-010-2424-y

Vlutters, M., van Asseldonk, E. H. F., & van der Kooij, H. (2016). Center of mass velocity-based predictions in balance recovery following pelvis perturbations during human walking. *Journal of Experimental Biology*, *219*(10). https://doi.org/10.1242/jeb.129338

Vlutters, M., van Asseldonk, E. H. F., & van der Kooij, H. (2018a). Foot placement modulation diminishes for perturbations near foot contact. *Frontiers in Bioengineering and Biotechnology*, *6*(48). https://doi.org/10.3389/fbioe.2018.00048

Vlutters, M., van Asseldonk, E. H. F., & van der Kooij, H. (2018b). Lower extremity joint-level responses to pelvis perturbation during human walking. *Scientific Reports*, *8*(1). https://doi.org/10.1038/s41598-018-32839-8

Vlutters, Mark, van Asseldonk, E. H. F., & van der Kooij, H. (2019). Ankle muscle responses during perturbed walking with blocked ankle joints. *Journal of Neurophysiology*, *121*(5). https://doi.org/10.1152/jn.00752.2018

Wang, Y., & Srinivasan, M. (2014). Stepping in the direction of the fall: The next foot placement can be predicted from current upper body state in steady-state walking. *Biology Letters*, *10*(9). https://doi.org/10.1098/rsbl.2014.0405

Warren, J., Kay, B. A., Zosh, W. D., Duchon, A. P., & Sahuc, S. (2001). Optic flow is used to control human walking. *Nature Neuroscience*, *4*(2). https://doi.org/10.1038/84054

Welch, T. D. J., & Ting, L. H. (2008). A feedback model reproduces muscle activity during human postural responses to support-surface translations. *Journal of Neurophysiology*, *99*(2). https://doi.org/10.1152/jn.01110.2007

Williams, D. S., & Martin, A. E. (2019). Gait modification when decreasing double support percentage. *Journal of Biomechanics*, *92*. https://doi.org/10.1016/j.jbiomech.2019.05.028

Zaytsev, P., Wolfslag, W., & Ruina, A. (2018). The Boundaries of Walking Stability: Viability and Controllability of Simple Models. *IEEE Transactions on Robotics*, *34*(2). https://doi.org/10.1109/TRO.2017.2782818

Zehr, E. P., & Stein, R. B. (1999). What functions do reflexes serve during human locomotion? *Progress in Neurobiology*, *58*(2). https://doi.org/10.1016/S0301-0082(98)00081-1

Zelik, K. E., & Adamczyk, P. G. (2016). A unified perspective on ankle push-off in human walking. *Journal of Experimental Biology*, *219*(23). https://doi.org/10.1242/jeb.140376

CoM states

positive foot
placement error

negative foot
placement error

zero foot placement error:
$FP = \beta_{pos} \cdot CoM_{pos} + \beta_{vel} \cdot CoM_{vel}$