Sexually antagonistic co-evolution can explain female display signals and male sensory adaptations.

R. Axel W. Wiberg[1,3*], Rosalind L. Murray[1,2], Elizabeth Herridge[1,4], Varpu Pärssinen[5], Darryl T. Gwynne[2], Luc F. Bussière[1,5,6*]

1. Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, United Kingdom

2. Department of Biology, University of Toronto at Mississauga, Ontario, L5L 1C6, Canada

3. Department of Zoology, Stockholm University, Svante Arrhenius väg 18B, Stockholm, Sweden, 106 91

4. Health and Social Care Analytical Services, St Andrews House, Regent Road, Edinburgh, EH1 3DG, United Kingdom

5. Institut för Biologi och Miljövetenskap, University of Gothenburg, Box 451, Göteborg, Sweden, 405 30

6. Gothenburg Global Biodiversity Centre, University of Gothenburg, Box 451, Göteborg, Sweden, 405 30

* Corresponding authors: **R. Axel W. Wiberg**; raw.wiberg@protonmail.com, **Luc F. Bussière**; luc.bussiere@bioenv.gu.se

Keywords: sexual selection, sensory exploitation, sexual dimorphism, male mate choice, sex roles, female ornamentation, dagger flies, empididae, empidinae

Abstract

The prevalence and diversity of female ornaments poses a challenge to evolutionary theory because males should prefer mates that spend resources on offspring rather than ornaments. Among dance flies, there is extraordinary variation in sexual dimorphism. Females of many species have conspicuous ornaments (leg scales and inflatable abdominal sacs). Meanwhile males of some species have exaggerated regions of their eyes with larger ommatidial facets that allow for regionally elevated photosensitivity and/or acuity. Here, we conduct a comparative study of these traits using both species descriptions available from the literature, as well as quantitative measures of eyes and ornaments from wild-caught flies. We show a conspicuous covariance across species between exaggerated male dorsal eye regions and the extent of female ornaments: species with highly ornamented females have males with more exaggerated eyes. We discuss this pattern in the context of competing hypotheses for the evolution of these traits and advance a role for sexual conflict.

Introduction

Female secondary sexual traits, such as ornaments, remain poorly understood in evolutionary biology. This is, in part, because female fecundity is often resource-limited, and males should avoid females that divert resources away from offspring towards extravagant

morphologies (Fitzpatrick *et al.,* 1995; Clutton-Brock 2009). Although some examples of such traits exist (Tobias et al. 2012; Dougherty 2021), and to greater extents in some clades (e.g. Kraaijeveld et al. 2007), female ornamentation remains generally rare in nature. Efforts

to explain the evolution of female ornamentation have focused largely on their evolution as a result of male preferences for more ornamented females, whereby the ornaments are inter- or intra- sexual indicators of fecundity, fertility, or quality (Schlupp 2018). This theoretical

framing also informs much of the empirical work in this field (e.g. Weiss and Dubin 2018; Yong et al. 2018; Kopena et al. 2018; Higham et al. 2021; Nolazco et al. 2022). By contrast, theory developed for male-biased display traits allows for more varied processes including

direct or indirect benefit signalling, but also exploitation of sensory biases, and sexual conflict (e.g. Sakaluk 2000; Macías Garcia and Ramirez 2005; Nakano et al. 2010; Kolm et al. 2012; Macías Garcia et al. 2012; Amcoff and Kolm 2015; Nakano et al. 2013)..

The dance flies of the subfamily Empidinae (Diptera: Empididae) are a speciose group with a global distribution, and exhibit remarkable diversity in mating systems and sexual dimorphisms among both sexes (Collin 1961; Cumming 1994; Chvala 2005; Murray et

al. 2022). The ancestral mating behaviour appears to have involved the aerial transfer of a nuptial gift (typically, an exogenous oral nuptial gift, *sensu* Lewis *et al.,* 2014) from the male upon which the female feeds during copulation (Downes 1970; Cumming 1994). Females

generally do not hunt as adults (Newkirk 1970), and, in many species, contests among females

for access to nuptial gifts are so intense that the operational sex ratio (OSR; Emlen and Oring

1977; Ahnesjö *et al.,* 2002) in aerial mating swarms is heavily female-biased. Intense contests

60   for nuptial gifts seem to have promoted the evolution of several secondary sexual characters

in females (Cumming 1994), including enlarged or coloured wings, inflatable abdominal sacs

and, most frequently, pinnate scales on one or more pairs of legs (see figure 1) which females

63   often wrap around their abdomens in flight and serve to attract and secure a male with a

nuptial gift (Murray *et al.,* 2018).

Meanwhile, males of some dance fly species have dichoptic compound eyes, with

66   dorsal eye regions showing enlarged ommatidia in comparison to females (Collin 1961; see

figure 1). These areas can enhance certain visual capacities including photosensitivity

(referred to as "bright zones"), or, when paired with a less curved eye surface, improve visual

69   acuity ("acute zones"; Land 1989; 1997). These adaptations can locally fine-tune optimal

visual capabilities, but come at the cost of visual ability in other regions of the eye. Such

trade-offs are due to constraints imposed on the resolving power of the compound eye by an

72   upper limit on eye size (Land 1989; 1997). Previous hypotheses to explain variation in male

eye morphology have explicitly or implicitly invoked selection among males for faster mate

acquisition or choice of more ornamented females (Downes 1970; Thornhill and Alcock

75   1983; Cumming 1994). For example, selection might favour specialised eyes as adaptations to

"scramble-like" male competition as in other systems (e.g. Eichorn *et al.,* 2017).

Alternatively, they have been proposed as adaptations to assess honest female ornaments

78   (LeBas et al., 2003).

Although a link between male eye morphology and female ornamentation in dance

flies has been proposed (e.g. LeBas et al., 2003), the degree of variation in eye dimorphism

81 and whether these traits covary across species has not been investigated. Here we conduct a comparative study investigating the co-evolution of female ornaments and male eye morphology in dance flies. We first make use of a survey of species descriptions on

84 morphology to assess broad patterns of association between eye dimorphism and ornament dimorphism among many dance flies. We then quantify associations between continuous measurements of male and female eye morphology and the degree of female ornamentation

87 for a subset of species. We discuss our findings in the context of competing hypotheses on the evolution of female ornamentation and male eye adaptations in these species.

90 **Methods**

Data collection and field sampling

We examined species descriptions in Collin (1961) to document possible associations

93 between female ornaments and eye sexual dimorphism among the British dance fly species of the subfamily Empidinae. Our use of descriptions in this key reflects both our focus on local fauna but also a bias resulting from the fact that comprehensive descriptions are largely only

96 available for species from Northern Europe and the British Isles. However, we note that there is extraordinary diversity from other geographic regions (e.g. Daugeron et al. 2011), and our results must be interpreted with this in mind. We noted the presence or absence of four main

99 categories of sexually dimorphic traits in females that are likely to be sexually selected ornaments: enlarged wing size, darker wing colouration, pinnate scales on legs, and inflatable abdominal sacs. We also noted any descriptions of sexual dimorphism in eye facets. We

102 collected information for all 95 species in the genera *Empis* and *Rhamphomyia*, and filtered the data to 93 species for which we could obtain information of female leg pinnation and/or

other ornamentation, and descriptions of male eye morphology. We included information from one outgroup in the genus *Hilara* for phylogenetically corrected analyses (see below).

In order to assess the covariance between continuous measures of ornaments and eye morphology, we collected dance fly samples during the summers 2009, 2010 and 2011 on the eastern side of Loch Lomond between the Scottish Centre for Ecology and the Natural Environment (SCENE) and Rowardennan, Scotland, and during the summer of 2012 near Glen Williams in Ontario, Canada. We caught flies from mating swarms or when resting on vegetation using sweep nets (Murray *et al.,* 2017). Of the sampled taxa, we selected eight species, *R. longicauda*, *R. longipes*, *R. stigmosa*, *R. crassirostris*, *E. aestiva*, *E. tessellata*, *E. nigripes*, and *H. chorica*, that represent a range of variation in female ornaments, male eye morphologies, and mating systems (e.g. OSR, type of nuptial gift; see also figure 1 and table S2 for final sample sizes).

Morphological Measures

Differences in eye morphology were quantified from nail varnish casts (see figure 1; Ribi *et al.,* 1986; Narendra *et al.,* 2010). Varnish was painted onto the eyes of specimens, allowed to dry, and then carefully removed. Three small cuts (figure 1) were then made to flatten the varnish casts and allow micrography for digital measurements. The diameters of seven neighbouring ommatidia on the dorsal and ventral portions of the eyes were measured from micrographs using tpsUtil (v 1.58; Rholf 2017) and tpsDig2 (v 2.17; Rholf 2017) in a manner similar to Döring and Spaethe (2009). We selected ommatidia located as dorsally or as ventrally as possible while accounting for cutting locations, obvious distortions in the nail varnish casts, and ensuring entire units of one central ommatidia and six surrounding

ommatidia were measured. The difference between mean dorsal and ventral diameters was standardised by thorax length and taken as a measure of exaggeration. Thus, a positive value

129   of exaggeration indicates larger dorsal ommatidia and a negative value indicates larger ventral ommatidia.

Female ornamentation was quantified by first computing the total leg area of the four

132   posterior legs of females for each species using the software ImageJ (v 1.48; Rasband 2016). While these ornament indices underestimate ornament expression for some species (*e.g.,* for species where females have multiple ornaments, such as inflated abdominal sacs and/or darker

135   wings in addition to pinnate scales), they nevertheless allowed us to measure trait expression in an analogous way across multiple taxa. Because the most heavily adorned species also had elaborate pinnate scales, ignoring the other ornament types makes our assessment of

138   covariance between eyes and ornament expression conservative. Moreover, our results from the literature search of Collin's (1961) key indicate that pinnation itself may be an especially strong predictor (see Results below). These measures were scaled by the total length of the

141   femora and tibia of all four legs. Finally, the mean scaled male leg area was subtracted from each individual female measure for each species to give individual measures of ornamentation for each female.

Sequence data collection and phylogenetics

We obtained sequence data for the carbamoylphosphate synthase domain of the

147   rudimentary locus (CAD) and the mitochondrial locus *Cytochrome oxidase I* (COI) from the literature (Watts et al. 2016; Wahlberg & Johanson 2018; Murray et al. 2020), and from the BOLD database (Ratnasingham et al. 2007; https://v3.boldsystems.org). After curating the

150 data to a single sequence per species, favouring longer sequences with fewer gaps for each

species, the final dataset comprised 63 species from the genera *Empis*, *Rhamphomyia*, and

Hilara as well as a further outgroup *Heterophlebus versabilis*, used also to root the tree in

153 Murray et al. 2020 (see table S3 for sequences and sources).

We then aligned the sequences for each locus separately with MAFFT (v. 7.505;

Katoh et al. 2013) and concatenated the alignments. Note that not all species had a

156 representative sequence for both loci (table S3). Where this was the case, the species received

a sequence of gap characters in the concatenated alignment. We then performed maximum

likelihood tree construction with IQ-Tree (v. 2.0.7; Nguyen et al. 2015) setting different gene

159 partitions for the two loci. We allowed IQ-Tree to test and pick the best substitution model ("-

m TEST"). Comparing the subset of this new phylogeny to that of Murray et al. (2020) finds a

Robinson-Fould's distance of 0.29, indicating that the majority of the splits are in agreement.

Statistical analyses

We tested for an association between the categorical information on eye dimorphism

165 and female ornamentation collected for a total of 93 species from Collin (1961) using several

Chi-squared tests to evaluate whether ornamental traits were associated with eye dimorphism

(figure S1). The first test considered the presence of ornament, regardless of type (i.e.

168 enlarged or darkened wings, inflatable abdominal sacs, pinnate leg scales). We also tested for

association between eye exaggeration and either a) pinnate leg scales or b) all other ornaments

excluding pinnate leg scales. This final contingency table has expected frequencies < 5 in

171 some cells (figure S3), necessitating the use of a Fisher's Exact Test.

For a subset of the data from the taxonomic descriptions, we could perform more

robust testing using phylogenetically corrected mixed effects models using our new

phylogeny (30 species, or 32% of the full data). We therefore pruned the tree to contain only these species, and fit a model with the presence/absence of male dorsal facet exaggerations as the response variable, and female pinnate scales as a predictor. Since male eye morphology was coded as 0 (no enlarged dorsal facets) or 1 (enlarged dorsal facets), we applied a Bernoulli error model (family=bernoulli(), using a logit link function).

For the eight species (*R. longicauda*, *R. longipes*, *R. stigmosa*, *R. crassirostris*, *E. aestiva*, *E. tessellata*, *E. nigripes*, and *H. chorica*) from which we obtained continuous morphological measures, the relationship between ornamentation, sex and eye exaggeration was also tested in a mixed model framework, fitting ornamentation, sex, and their interaction to predict eye exaggeration. The species studied here are from two genera, but recent molecular phylogenetic work shows that these genera are not monophyletic, and that female ornamental traits have evolved independently several times (Watts et al., 2016; Murray et al., 2020). To include phylogenetic information, we used the consensus tree from Murray et al., (2020), pruned to contain the eight species of interest. The results are the same whether we use the phylogeny of Murray et al. (2020) or the new one from this study. We report results using the phylogenetic tree from Murray et al. (2020) in this analysis because there is an associated posterior distribution of trees. Thus, in addition to the main analysis on the consensus tree, we also marginalised our analyses over a random sample of trees from the. Previous studies have suggested similar approaches to account for uncertainties in the topology of the tree (Hadfield and Nakagawa 2010; Hadfield 2010; Ross et al., 2013; Murray et al., 2020). Although simulations have suggested that as few as 50 trees from the posterior distribution may be required to adequately account for phylogenetic uncertainty (Nakagawa & de Villemereuil 2019), we sampled 200. We did this by fitting several models, each with a

different input tree (with the function brm_multiple()), and merging posterior samples. In all

198   models we also included a second random effect to account for multiple measures from each

species. Continuous predictors were scaled (the mean subtracted from each value) prior to

analysis (Schielzeth et al., 2010).

201   All analyses were performed in R (v.4.3.3; R Core Team 2024) using the packages

"ape" (v. 5.4; Paradis et al., 2004), "brms" (v. 2.20.4; Bürkner 2017), "loo" (v. 2.7.0; Vehtari

et al. 2024) and "coda" (v. 0.19.4; Plummer et al. 2006). Figures were plotted with "ggplot2"

204   (v. 3.3.1; Wickham 2016). We follow Muff et al. (2022) and report results using the

"language of evidence" for results of frequentist statistics and evaluate Bayesian results in

terms of the entire posterior distributions and references to the 89% highest posterior density

207   intervals (HPDIs; McElreath 2018).

Results

210   We found strong evidence for positive associations between ornament expression and

eye exaggeration from the 93 taxonomic descriptions of female ornaments and male eyes

(figures S1-S3). Dimorphism in eye morphology was more strongly associated with pinnate

213   leg scales than other ornament characters (figure S1, all ornaments together: Chi-squared =

11.63, d.f. = 1, $p < 0.001$; figure S2, pinnate scales only: Chi-squared = 17.02, d.f. = 1, $p <$

0.001; table S1C). In contrast, there was no evidence for an association with other ornaments

216   excluding pinnate scales (Fisher's Exact Test: Odds Ratio (95% CI) = 0.76 (0.18, 2.93), p =

0.77; figure S3).

These same patterns are also reflected in the phylogenetically corrected analysis of

219   the subset of 30 species. Although the 89% highest posterior density interval (89% HPDI)

includes 0 (figure 2), 92% of the full posterior distribution lies above 0. The weight of the evidence thus indicates a decidedly positive association between female leg pinnation and

male eye exaggerations. Importantly, all effects are in the same direction; species with female ornamentation or leg pinnation are more likely to also have exaggerated eye morphology (figures S1-S3 and figure 2). The posterior mean of phylogenetic heritability was 0.24.

Although the 89% HPDI did not include 0, it was extremely wide (figure S4). Moreover, Leave-One-Out (LOO) model comparisons indicate that a model without phylogenetic effects is a better fit to the data (ELPD$_{phylogeny}$ = -21.4, ELPD$_{no-hylogeny}$ = -20.2; SE$_{diff}$ = 0.6).

For the subset of eight species for which we collected wild specimens, we found strong evidence for an association between continuous measures of female ornamentation and eye exaggeration, with more exaggerated eyes occurring in taxa with more ornate females

(figures 3 and 4). Posterior distributions for the sex by female ornamentation interaction term never overlapped 0, and the 89% HDPI ranged from 0.0921 to 0.108 (figure 3D). Intriguingly, female eyes in ornamented species tended to show the opposite pattern to males (89% HDPI =

[-0.055, -0.0128]; figure 3B); females had larger ventral facets than dorsal ones, and more ornamented species tended to have more exaggerated ventral facets (partial slopes for females were negative; figures 3 and 4). The evidence for these associations remained strong in

models that controlled for phylogeny while accounting for phylogenetic uncertainty by marginalising over a sample of the posterior distribution of trees (figures S7 and S8). Posterior means of phylogenetic heritability were low to moderate (0.41-0.45), but there was

considerable uncertainty in these estimates for both models (figure S9). Indeed, LOO model comparison of the model including the consensus phylogenetic tree to a model without the tree only marginally preferred including the phylogeny (ELPD$_{phylogeny}$ = 663.9, ELPD$_{no-hylogeny}$

$= 663.8$; $SE_{diff} = 0.2$).

Discussion

246   A number of hypotheses involving conventional sexual selection or sexual conflict

have been proposed for female ornamentation and male eye exaggerations in Empidine dance

flies. We show that species with more heavily ornamented females also tend to have males

249 with more exaggerated dorsal eye facets. These findings are consistent with a process of co-

evolution between these two traits.

Previous hypotheses to explain variation in male eye morphology have explicitly or

252 implicitly invoked selection *among males* for faster mate acquisition or choice of more

ornamented females (Downes 1970; Thornhill and Alcock 1983; Cumming 1994). First,

exaggerated eye morphologies could be adaptations to "scramble-like" male competition as in

255 other systems (Eichorn et al., 2017). However, this hypothesis predicts more exaggeration in

species with strong competition among males. Because female ornaments are more prevalent

in dance fly species where females are more mate-limited than males (Murray et al. 2020), we

258 would expect a negative association between eye dimorphism and female ornament

extravagance under this hypothesis. This is the opposite of our finding of positive covariance

between ornament expression and male eye dimorphism (figure 3).

261   Alternatively, eye adaptations could result from competition among males to facilitate

appraisal and identification of the highest quality females. Under this scenario, exaggerated

eye morphologies facilitate assessments of female quality. However, the coincidence of both

264 extensive female ornamentation and heightened visual ability would be surprising under this

hypothesis. There would be little need for males to invest in more exaggerated eye

morphologies if elaborate female ornaments were an honest signal of quality (in fact such

267   exaggerations could be costly due to trade-offs in visual adaptations; e.g. Land [1989]). Conversely, females would have no need to invest in honest ornaments if males had heightened visual acuity to assess female reproductive value (e.g., fecundity).

270   Another alternative is that eye exaggerations might be required as part of a male preference trait for female ornaments that honestly provide signals of reproductive value (e.g. Bloch 2015). Rather than playing a role in male scramble contests, in this case the

273   exaggerated eye morphology is the means by which males exercise preference for mates, and is expected to covary positively with the degree of female ornamentation. This pattern of positive covariation is a standard prediction of sexual selection theory where preferences and

276   ornamental traits coevolve (Andersson 1994). However, this hypothesis relies on female ornaments being an honest indicator of either direct benefits (more fecund females) or indirect benefits, such as offspring viability (good genes) or attractiveness (sexy daughters), both of

279   which are problematic on several grounds (see below).

Hypotheses to explain the evolution of female ornaments among dance flies either invoke honest signalling of direct benefits (LeBas et al., 2003) or deception (Funk and

282   Tallamy 2000; Hockham and Ritchie 2000). Conventional sexual selection on honest signals predicts a negative association between female ornament expression and the proportion of males in mating swarms (OSR; Murray et al., 2020), because choice is most likely when

285   receptive females outnumber males (Bonduriansky 2001). Although the OSR may not be a perfect index of sexual selection, it is nevertheless a reliable guide to the differences in the opportunity for sexual selection (in males), as well as sex-differences in the strength of sexual

288   selection among animals (Janicke et al. 2018). Ornamented females could for example be

advertising direct (material) fitness benefits to mates, or indirect (genetic) gains for pairing with them. Direct benefits are much more likely than indirect ones to explain male mating

291   preferences; they can act immediately rather than awaiting expression in offspring, and do not depend on strong inter-sexual genetic correlations (Kirkpatrick and Barton 2001). Furthermore, variation in reproductive value among females should almost always exceed

294   variation in genetic quality (Kirkpatrick and Barton 2001).

In dance flies, there are two especially likely sources of variation in female quality representing direct benefits. Males might prefer females with ornaments that signal (1) greater

fecundity or (2) a later stage of ovarian development (i.e., the imminence of oviposition). A female that is heavily gravid and soon to lay eggs may be especially valuable among insects with sperm storage, because such a female is less likely to mate again prior to oviposition, and

remating generally decreases a focal male's paternity by displacing his ejaculate within sperm stores (*e.g.* Boorman and Parker 1976). However, dance fly ornaments seem to indicate ovarian status especially inconsistently. While some studies suggest that males prefer females

with larger ornaments, the presence and intensity of sexual selection is remarkably variable across dance fly species and episodes of mating (Funk and Tallamy 2000; LeBas et al.*,* 2003; Murray et al., 2018). In some species, there is no evidence that larger or more ornamented

females acquire more mates (Hockham and Ritchie 2000; Bussière et al.*,* 2008; Wheeler et al., 2012; Herridge 2016; Murray et al., 2019). Such inconsistency is not predicted by hypotheses of honest signalling, even if the intensity of selection can sometimes vary across populations,

or through time (e.g. due to the availability of prey). Moreover, although there are usually significant and positive covariances between female ornamentation and fecundity or egg size, they can be low compared to covariances between fecundity or ovarian maturity and other

indices of body size (*e.g.* thorax length; LeBas et al., 2003; Bussière et al., 2008; Murray et

al., 2019; Murray et al., 2020; but see Funk and Tallamy 2000 and Wheeler 2008). In fact,

regardless of the covariance between cross-sectional measures of fecundity and ornament

315   expression, female ornaments are inherently unable to track changes in reproductive value

related to ovarian development because ornaments like pinnate leg scales and wing colour are

fixed at eclosion, while the number and developmental maturity of eggs changes throughout

318   adult life. For example, in some ornamented species, ovarian development appears to be

entirely arrested prior to an initial mating and nuptial feeding event (Hunter & Bussière 2019),

which means that signals of fecundity from virgin females are by necessity deceptive.

321   Together, these facts suggest that ornaments may sometimes degrade rather than improve

male assessments of female reproductive quality.

What if these ornaments were instead indicators of genetic benefits? Addressing this

324   possibility requires knowledge of the relative costs (e.g., in paternity share) and benefits to

males (in enhanced fitness of offspring) for mating with attractive females. There is some

evidence of increased sperm competition among males in more ornamented species (Murray

327   et al., 2020), but we altogether lack information on heritable benefits of female attractiveness.

Given equivocal support for indirect benefits for mating with attractive males in many

systems, the presence of substantial variation in direct benefits weakens the probability that

330   indirect benefits could favour male preferences that incur substantial direct costs (Kirkpatrick

and Barton 1997). Consequently, neither indirect nor direct benefits of honest signalling seem

able to satisfyingly explain the diversity of dance fly ornaments.

333   As an alternative to honest signalling, female ornaments in dance flies could be

generally deceptive adaptations that exploit pre-existing male preferences for larger females.

Deceptive ornaments would allow females to obtain nuptial gifts regardless of their current

336 fecundity or stage of egg development (Funk and Tallamy 2000; Hockham and Ritchie 2000; Arnqvist and Rowe 2005; Arnqvist 2006), even if males prefer highly gravid females to avoid increased sperm competition risk (Herridge et al., 2016). If female ornaments are in fact

339 deceptive signals designed to exploit male preferences for large females, we might expect resistance traits to evolve in some well-adorned species (Arnqvist and Rowe 2005). Although no resistance traits have ever been previously described among male dance flies, adaptations

342 that augment visual perception and acuity, such as the exaggerated male eye regions observed here, are exactly the sorts of traits that are expected.

Sexual conflict and sexually antagonistic coevolution (SAC) are relatively recent

345 models proposed to explain the origin and exaggeration of sexually dimorphic traits (Parker 1979; Holland and Rice 1998; Arnqvist and Rowe 2005). Sensory exploitation of one sex by the other can play a key role in this model, with deceptive signal traits selecting for resistance

348 in the receiver sex; the resulting arms race drives the diversification of both signal and receiver traits (Holland and Rice 1998; Arnqvist and Rowe 2005). When sensory exploitation via a seductive signal induces a suboptimal mating rate in the receiver, it increases the

351 receiver's exposure to costs of mating (Arnqvist 2006). To avoid costs, receivers should evolve increased discriminatory power to distinguish manipulative signals from honest ones, or to reduce susceptibility to seductive stimuli altogether. Repeated bouts of exploitation and

354 resistance should lead to cycles of antagonistic co-evolution that can lead to a proliferation of diverse seduction and resistance adaptations (Holland and Rice 1998). SAC could explain seductive female traits and male resistance in rare species possessing strong male choice

357 (Funk and Tallamy 2000; Murray et al., 2018). In such species male preference evolution

should be constrained because female fecundity is often resource-limited, and males should avoid females that divert resources away from offspring towards extravagant morphologies 360 (Fitzpatrick et al., 1995; Clutton-Brock 2009). Additionally, if females store their mates' sperm, heavily ornamented females might acquire more mates, and consequently provide each male with a lower paternity share than less ornamented (and less polyandrous) females 363 (Servedio and Lande 2006; Herridge et al., 2016). There is good evidence that sexually antagonistic co-evolution drives diversification in traits that mediate physical mating interactions (Arnqvist and Rowe 2002; Rowe and Arnqvist 2002; Rönn et al. 2007; Bergsten 366 and Miller 2007; Tatarnic and Cassis 2010). Sexually antagonistic sensory exploitation has also been offered as an explanation for the origins and diversification of nuptial gifts and some signal traits, and may drive the coevolution of male display traits and female 369 behavioural responses (Sakaluk 2000; Macías Garcia and Ramirez 2005; Nakano et al. 2010; Kolm et al. 2012; Macías Garcia et al. 2012; Amcoff and Kolm 2015; Nakano et al. 2013). However, where these examples demonstrate counter adaptations in females they tend to 372 result in dissociation between the sexual signal and the ecological signal (i.e. escape from a sexually antagonistic sensory trap), rather than co-evolving exaggerations of signal and receiver traits (e.g. Macías Garcia and Ramirez 2005; Nakano et al. 2013). To our knowledge, 375 no previous studies have shown comparative evidence for escalating sexually antagonistic co-evolution between signal and primary sensory traits, let alone in systems where females are the ornamented sex.

378   SAC, therefore, offers a compelling alternative hypothesis that overcomes several difficulties with the evolution of female ornaments, and can at once explain variation in both eye and ornamental traits across the dance flies. Female ornaments might exploit pre-existing

381 male preferences for larger but unadorned females (because large females tend to be more

gravid or more fecund; Funk and Tallamy 2000; Bonduriansky 2001; Arnqvist 2006), so that

females can obtain nutritious nuptial gifts regardless of their current state of ova development.

384 Males could then respond to female deception by developing resistance traits (increased

visual perception) to avoid females with low fecundity. According to SAC, exaggerations in

male eye morphology would reflect selection for increased discriminatory power to "see past"

387 increasingly deceptive ornaments of hungry females, and identify the most gravid or fecund

mates despite female disguises. This hypothesis predicts a positive association between

exaggerations in eye morphology and ornamentation across species, as is seen in our data.

390 This positive covariation would arise because exploitative adaptations that increase mating

frequency for females (increased ornamentation) should be countered by increased investment

in discriminatory power by males (exaggerated dorsal facet morphologies). The SAC

393 hypothesis also predicts a negative association between ornamentation and the proportion of

males in the swarm (the swarm sex ratio) because large deceptive ornaments should be most

favoured in environments where females compete more intensely for resources and

396 experience stronger selection to deceive males (Murray *et al.,* 2020).

Moreover, our results from the literature survey suggest that exaggerations of dorsal

eye morphologies in males may be specific responses to deception *via* leg pinnation. The

399 association between ornaments and male eye exaggerations were strongest when

ornamentation was defined as the presence of leg pinnation alone. However, the prevalence of

pinnate scales relative to other kinds of ornaments makes strong conclusions difficult (in our

402 sample of taxonomic descriptions, 33/93 species have females with pinnate leg scales; while

only 13/93 species have other forms of female ornamentation but lack pinnation). These

results hold in phylogenetically corrected analyses of a subset of these species for which we

405   were able to collect phylogenetic information. The possibility that leg ornaments in particular

select for dichoptic eyes is intriguing because the heightened photosensitivity of large eye

facets may be especially effective at detecting stray photons that betray the difference

408   between deceptively-positioned scaly legs that exaggerate abdomen size and egg-inflated

large abdomens. In this context it is also noteworthy that the most extremely ornamented

species, with the most exaggerated male dorsal acute zones, also have other ornamental

411   features. *R. longicauda,* for example, has inflatable abdominal sacs that can serve to further

increase a female's apparent size to the male (Funk and Tallamy 2000; Murray *et al.,* 2018;

Murray *et al.,* 2020). These inflatable sacs might also be more difficult for the male to resist

414   by increased photosensitivity or visual acuity. In this way, additional ornamental features,

such as abdominal sacs or darkened wings may represent more recent or sequential

innovations in the historical arms race between seductive female characters and

417   discriminatory male sensory structures.

Naturally, visual perception is an important part of dance fly ecology in many other

contexts apart from mate assessment. For example, hunting ability probably depends heavily

420   on visual acuity. The sex-difference in hunting nuptial gifts could therefore conceivably lead

to differences in eye dimorphism. However, there are no obvious cross-species differences in

prey choice that align with variation in ornament expression or eye exaggeration (almost all

423   empidines present nuptial gifts, including the eight species we measured quantitatively;

Laurence 1952; Collin 1961; Chvala 2005). Furthermore, if hunting ability did drive the

evolution of eye exaggerations, it might be strongest and most obvious in species where male

426   competition for females is intense (i.e. high OSR, low ornamentation), to favour rapid return

to mating swarms, whereas we found the converse (figure 4). We lack any evidence that hunting efficiency covaries strongly with ornament expression, and a more systematic survey of dance fly foraging ecology would help shed light on these questions.

The light environment also selects strongly on visual systems (Land 1989; 1997; Eichorn *et al.,* 2017), and some theory suggests that female ornaments are more likely in environments where male discrimination is difficult (Chenoweth *et al.,* 2006). However, no great differences in light environment (*e.g.* time of day, canopy cover) are obvious among these species from field observations (Downes 1970; Cumming 1994; Murray *et al.,* 2017; Murray *et al.,* 2018; our own observations). It is also not clear why the light environment would select for both better visual perception *and* large ornaments in females because either trait on its own would presumably improve discrimination. Nevertheless, subtle differences might play a role in adaptations toward greater visual acuity.

Our study establishes a strong signal of covariation between male eye-exaggeration and female ornamentation. We suggest that this signal is inconsistent with sexual selection in male biased swarms, and that explanations relying on honest female ornaments are problematic. We outline a plausible and intriguing alternative hypothesis involving sexual conflict and a co-evolutionary diversification of female ornaments and male visual systems. These traits provide several exciting opportunities to study the emergence of novel phenotypes, as it may be possible to use female (in the case of eye morphology) or phylogenetically "basal" morphologies as ancestral states or starting points. They also provide a novel system in which to study the inherent sexual conflicts that arise in producing sexually dimorphic phenotypes from a common genome (Pennell and Morrow 2013).

Funding

This work was supported by a Nuffield Undergraduate Summer Scholarship [URB/39500] to RAWW, a Royal Society Research Grant [2010 R1] to LFB, an NSERC PGS-D and a University of Stirling Horizon Graduate Studentship to RLM.

Acknowledgements

The authors would also like to thank Adrian Plant and David Price for their invaluable assistance in species identification and natural history, NSERC, the University of Stirling, and the Scottish Centre for Ecology and the Natural Environment (SCENE) for support and accommodation during field work, and Mike Ritchie and Locke Rowe for helpful comments and discussion.

Data and analysis code availability

All data, including phylogenetic trees, and scripts used for the statistical analyses and for producing the figures in this manuscript have been deposited in a dryad repository (https://doi.org/10.5061/dryad.rr4xgxd5z).

Competing interests

The authors declare no competing interests.

Author contributions

RAWW and LFB conceived the study. RAWW, RLM, EH, VP, DTG, and LFB all contributed to the collection of specimens and data. RAWW performed the data analysis with help from RLM and LFB. RAWW led the writing of the manuscript with input from RLM, EH, DTG and LFB. All authors have approved the publication of the final manuscript.

Ethics

Specimens of male and female Empidine dance flies were captured in the field and sacrificed

by freezing until morphological measurements were made. We obtained ethical approval for

all sampling and measurement procedures from the University of Stirling research ethics

474   committee.

References

Ahnesjö, I., Kvarnemo, C., & Merilaita, S. 2002. Using potential reproductive rates to predict

477   mating competition among individuals qualified to mate. Behavioral Ecology. 12: 397–

    401.

Amcoff, M., & Kolm, N. 2014. A test of sensory exploitation in the swordtail characin

480   (Corynopoma riisei) based on colour matching between female prey and a male

    ornament. Environmental Biology of Fishes. 97: 247–254.

Andersson, M. B. 1994. Sexual selection. Princeton University Press. Princeton, NJ.

483 Arnqvist, G. 2006. Sensory exploitation and sexual conflict. Philosophical Transactions of the

    Royal Society of London B: Biological Sciences. 361: 375–386.

Arnqvist, G. & Rowe, L. 2002. Correlated evolution of male and female morphologies in

486   water striders. Evolution. 56: 936–947.

Arnqvist, G. & Rowe, L., 2005. Sexual Conflict. (J.R. Krebs & T. Clutton-Brock, Eds.)

    Princeton University Press, Princeton, New Jersey.

489 Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. 2015. Fitting linear mixed-effects

    models using lme4. Journal of Statistical Software, 67

Bergsten, J. & Miller, K. 2007. Phylogeny of diving beetles reveals a coevolutionary arms

492   race between the sexes. PLoS ONE. 2: e522.

Bloch, N. I. 2015. Evolution of opsin expression in birds driven by sexual selection and

    habitat. Proceedings of the Royal Society: Biological Sciences. 282: 20142321.

495   Bonduriansky, R. .2001. The evolution of male mate choice in insects: a synthesis of ideas

    and evidence. Biological Reviews. 76: 305–339.

Boorman, E. and Parker, G. A. 1976. Sperm (ejaculate) competition in Drosophila

498   melanogaster, and the reproductive value of females to males in relation to female age

    and mating status. Ecological Entomology. 1: 145–155.

Bussière, L. F., Gwynne, D. T., & Brooks, R. 2008. Contrasting sexual selection on males and

501   females in a role-reversed swarming dance fly, Rhamphomyia longicauda Loew

    (Diptera: Empididae). Journal of Evolutionary Biology. 21: 1683–1691.

Bürkner, P-C. 2017. brms: An R Package for Bayesian Multilevel Models Using Stan.

504   *Journal of Statistical Software.* 80: 1-28.

Chenoweth, S., Doughty, P. & Kokko, H. 2006. Can non-directional male mating preferences

    facilitate honest female ornamentation? Ecology Letters. 9: 179–184.

507   Chvala, M. 2005. Fauna Entomologica Scandinavica: The Empidoidea (Diptera) of

    Fennoscandia and Denmark. III. New York. Leiden: E.J. Brill.

Clutton-Brock, T. 2009. Sexual selection in females. Animal Behaviour. 77: 3–11.

510   Collin, J. E. 1961. British Flies: Empididae. Cambridge University Press.

Cumming, J. M. 1994. Sexual selection and the evolution of dance fly mating systems

    (Diptera: Empididae: Empidinae). The Canadian Entomologist. 126: 907–920.

513   Dougherty, L. 2021. Meta-analysis reveals that animal sexual signalling behaviour is honest

and resource-based. Nature Ecology and Evolution. 5: 688 – 699.

Daugeron, C., Plant, A., Winkler, I. & Baylac, M. 2011. Extreme male leg polymorphic asymmetry in a new empidine dance fly (Diptera: Empididae). Biology Letters. 7: 11 – 14.

Downes, J. A. 1970. The feeding and behaviour of the specialized Epidinae (Diptera); observations on four species of Rhamphomyia in the high arctic and a general discussion. The Canadian Entomologist, 102: 769–791.

Döring, T. F., & Spaethe, J. 2009. Measurement of eye size and acuity in Aphids (Hemiptera: Aphididae). Entomologica Generalis. 32: 77–84.

Eichorn, C., Hrabar, M., Van Ryn, E. C., Brodie, B. S., Blake, A. J., & Gries, G. 2017. How flies are flirting on the fly. BMC Biology. 15:2

Emlen, S. T. & Oring, L. W. 1977. Ecology, sexual selection, and the evolution of mating systems. Science. 197: 215–223.

Fitzpatrick, S., Berglund, A., & Rosenqvist, G. 1995. Ornaments or offspring: costs to reproductive success restrict sexual selection process. Biological Journal of the Linnean Society. 55: 251–260.

Funk, D. H., & Tallamy, D. W. 2000. Courtship role reversal and deceptive signals in the long-tailed dance fly, Rhamphomyia longicauda. Animal Behaviour. 59: 411–421.

Hadfield, J. D. 2010. MCMC methods for multi-respoinse generalized linear mixed models: The MCMCglmm R package. Journal of Statistical Software. 33: 1–22.

Hadfield, J. D., & Nakagawa, S. 2010. General quantitative genetic methods for comparative

biology: Phylogenies, taxonomies and multi-trait models for continuous and categorical characters. Journal of Evolutionary Biology. 23: 494–508.

537 Halekoh, U. & Højsgaard, S. 2014. A Kenward-Roger approximation and parametric bootstrap methods for tests in linear mixed models - The R Package pbkrtest. Journal of Statistical Software. 59: 1–32.

540 Herridge, E. J., Murray, R. L., Gwynne, D. T., & Bussière, L. F. 2016. Mating and Parental Sex Roles, Diversity in. Pp 453 - 458 in Encyclopedia of Evolutionary Biology vol. 2, (R.M. Kliman, Ed.). Oxford Academic Press.

543 Herridge, EJ. 2016. The role of polyandry in sexual selection among dance flies. PhD Thesis, University of Stirling. http://hdl.handle.net/1893/25013.

Higham, J.P., Kimock, C.M., Mandalaywala, T.M., Heistermann, M., Cascio, J., Petersdorf, 546 M., Winters, S., Allen, W.L. & Dubuc, C. 2021. Female ornaments: is red skin color attractive to males and related to condition in rhesus macaques. Behavioral Ecology. 32: 236-247.

549 Hockham, L. R., & Ritchie, M. G. 2000. Female secondary sexual characteristics: appearance might be deceptive. Trends in Ecology and Evolution. 15: 436–437

Holland, B., & Rice, W. R. 1998. Perspective: Chase-away sexual selection: Antagonistic 552 seduction versus resistance. Evolution, 52: 1–7.

Hunter, F. D. L, & Bussière, L. F. 2019. Comparative evidence supports a role for reproductive allocation in the evolution of female ornament diversity. Ecological 555 Entomology. 44: 324–332.

Janicke, T. & Morrow, E.H. 2018. Operational sex ratio predicts the opportunity and direction of sexual selection across animals. Ecology Letters. 21: 384-391.

558   Katoh, K. & Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution. 30: 772-780.

561   Kirkpatrick, M. & Barton, N. H. 1997. The strength of indirect selection on female mating preferences. Proceedings of the National Academy of Sciences of the United States of America. 94: 1282–1286.

564   Kolm, N., Amcoff, M., Mann, R.P. & Arnqvist, G. 2012. Diversification of a food-mimicking male ornament via sensory drive. Current Biology. 22: 1440-1443.

Kopena, R., López, P., Majlathova, V. & Martín, J. 2018. Sexually dichromatic coloration of
567   female Iberian green lizards correlates with health state and reproductive investment. Behavioral Ecology and Sociobiology. 74: 131

Kraaijeveld, K., Kraaijeveld-Smit, Femmie J. L. & Komdeur, J. 2007. "The evolution of
570   mutual ornamentation." Animal Behaviour 74: 657-677.

Land, M. F. 1989. Facets of Vision. (D.G. Stavenga & R.C. Hardie, Eds.). Springer Verlag, Berlin.

573   Land, M. F. 1997. Visual acuity in insects. Annual Review of Entomology. 42: 146–177.

Laurence, B. R. 1952. The prey of some Empididae and Dolichopodidae (Dipt.). The Entomologist's Monthly Magazine 88: 156–157.

576   LeBas, N. R., Hockham, L. R. & Ritchie, M. G. 2003. Nonlinear and correlational sexual

selection on 'honest' female ornamentation. Proceedings of the Royal Society: Biological Sciences. 270: 2159–2165.

579   LeBas, N. R. & Hockham, L. R. 2005. An Invasion of Cheats: The evolution of worthless nuptial gifts. Current Biology. 15: 64–67.

Lewis, S. M., Vahed, K., Koene, J. M., Engqvist, L., Bussière, L. F., Perry, J. C., Gwynne, D.
582   T., & Lehmann, G. U. C. 2014. Emerging issues in the evolution of animal nuptial gifts. Biology Letters. 10: 20140336.

Macías Garcia, C. & Ramirez, E. 2005. Evidence that sensory traps can evolve into honest
585   signals. Nature. 434: 501-505.

Macías Garcia, C. & Saldívar Lemus, Y. 2012. Foraging costs drive female resistance to a sensory trap. Proceedings of the Royal Society: B. Biological Sciences. 279: 2262-2268.

588   McElreath, R. 2018. Statistical rethinking: A bayesian course with examples in r and stan. Chapman; Hall/CRC.

Muff, S., Nilsen, E.B., O'Hara, R.B. & Nater, C.R. 2022. Rewriting results sections in the
591   language of evidence. Trends in Ecology & Evolution. 8: 203-210.

Murray R. L, Herridge, E. J., Ness R. W., Wiberg R. A. W., Gwynne D. T. & Bussière L. F.
       2020. Competition for access to mates predicts female-specific ornamentation and male
594   investment in relative testis size. Evolution. 74: 1741–1754.

Murray, R. L., Gwynne, D. T. & Bussière, L. F. 2019. The role of functional constraints on non-random mating patterns for a dance fly with female ornaments. Journal of
597   Evolutionary Biology. 32: 984–993 .

Murray, R. L., Herridge E. J., Ness R. W. & Bussière L. F. 2017. Are sex ratio distorting endosymbionts responsible for mating system variation among dance flies (Diptera: Empidinae)? PLoS ONE 12: e0178364.

Murray, R. L., Wheeler, J., Gwynne, D. T. & Bussière, L. F. 2018. Sexual selection on multiple female ornaments in dance flies. Proceedings of the Royal Society: Biological Sciences. 285: 20181525.

Murray, R.L., Gwynne, D.T. & Bussière, L.F. 2022. Mating and Sexual Selection in Empidine Dance Flies (Empididae). Insects. 13: 839.

Nakagawa, S. & de Villemereuil, P. 2019. A general method for simultaneously accounting for phylogenetic and species sampling uncertainty via Rubin's rules in comparative analysis. Systematic Biology. 68: 632-641.

Nakano, R. Takanashi, T. Skals, N., Surlykke, A. & Ishikawa, Y. 2010. Ultrasonic courtship songs of male Asian corn borer moths assist copulation attempts by making the females motionless. Physiological Entomology. 35: 76-81.

Nakano, R., Takanashi, T., Surlykke, A., Skals, N. & Ishikawa, Y. 2013. Evolution of deceptive and true courtship songs in moths. Scientific Reports. 3: 2003.

Narendra, A., Reid, S. F., Greiner, B., Peters, R. A, Hemmi, J. M., Ribi, W. A, & Zeil, J. 2010. Caste-specific visual adaptations to distinct daily activity schedules in Australian Myrmecia ants. Proceedings of the Royal Society: Biological Sciences, 278: 1141–1149.

Newkirk M. R. 1970. Biology of the longtailed dance fly, Rhamphomyia longicauda (Diptera: Empididae). Annals of the Entomological Society of America. 63: 1407–1412.

Nguyen, L.T., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution. 32: 268–274.

Nolazco, S., Delhey, K., Nakagawa, S. & Peters, A. 2022. Ornaments are equally informative in male and female birds. Nature Communications. 13: 5917.

Paradis, E., Claude, J., & Strimmer, K. 2004. APE: Analyses of phylogenetics and evolution in R language. Bioinformatics, 20: 289–290.

Parker, G. 1979. Sexual selection and sexual conflict. In M.S. Blum and N. A. Blum (eds.) Sexual Selection and Reproductive Competition in Insects, Academic, New York, pp. 123–166.

Pennell, T. M. & Morrow, E. H. 2013. Two sexes, one genome: the evolutionary dynamics of intralocus sexual conflict. Ecology and Evolution. 3: 1819–1834.

Plummer, M., Best, N., Cowles, K. & Vines, K. 2006. CODA: Convergence Diagnosis and Output Analysis for MCMC, R News, 6: 7-11.

R Development Core Team. 2020. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at: https://www.R-project.org/

Rasband, W. S. 1997-2016 ImageJ, U.S. National Institutes of Health, Bethesda, Maryland available at: http://imagej.nih.gov/ij/

Ratnasingham S. & Hebert, P.D.N. 2007. BOLD: The Barcode of Life Data System (www.barcodinglife.org). Molecular Ecology Notes 7: 355-364.

Ribi, W. A., Engels, E., & Engels, W. 1989. Sex and caste specific eye structures in stingless bees and honey bees (Hymenoptera: Trigonidae, Apidae). Entomologica Generalis. 14: 233–242.

Rohlf, F. J. 2017. tpsDig2. Software. Available at: http://life.bio.sunysb.edu/morph/

Rohlf, F. J. 2017. tpsUtil. Software. Available at: http://life.bio.sunysb.edu/morph/

Ross, L., Gardner, A., Hardy, N., & West, S. A. 2013. Ecology, not the genetics of sex determination, determines who helps in eusocial populations. Curr. Biol. 23: 2383–2387.

Rowe, L., & Arnqvist, G. 2002. Sexually antagonistic coevolution in a mating system: combining experimental and comparative approaches to address evolutionary processes. Evolution. 56: 754–767.

Rönn, J., Katvala, M. & Arnqvist, G. 2007. Coevolution between harmful male genitalia and female resistance in seed beetles. *Proceedings of the National Academy of Sciences of the United States of America* 104: 10921-10925.

Sakaluk, S. K. 2000. Sensory exploitation as an evolutionary origin to nuptial food gifts in insects. Proceedings of the Royal Society: Biological Sciences. 267: 339–343.

Servedio, M.R. & Lande, R. 2006. Population genetic models of male and mutual mate choice. *Evolution*. 60: 647–685.

Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients. Methods in Ecology and Evolution. 1: 103–113.

Schlupp, I. 2018. Male mate choice, female competition, and female ornaments as components of sexual selection. Current Zoology. 64: 321-322.

Tatarnic, N. & Cassis, G. 2010. Sexual coevolution in the traumatically inseminating plant
bug genus Coridromius. Journal of Evolutionary Biology. 23: 1321–1326.

663   Thornhill, R., & Alcock, J. 1983. The evolution of insect mating systems. Harvard University
Press, Cambridge, Massachusetts.

Tobias, J.A., Montgomerie, R. & Lyon, B.E. 2012. The evolution of female ornaments and
666   weaponry: social selection, sexual selection and ecological competition.

Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Bürkner, P-C., Paananen T. & Gelman A.
2024. loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models. R
669   package version 2.7.0, available from: https://mc-stan.org/loo/.

Wahlberg, E. & Johanson, K.A. 2018. Molecular phylogenetics reveals novel relationships
within Empidoidea. Systematic Entomology. 43: 619-636.

672   Watts, M., Winkler, I. S., Daugeron, C., de Carvalho, C. J.B., Turner, S. P., & Wiegmann, B.
M. 2016. Where do the Neotropical Empidini lineages (Diptera: Empididae: Empidinae)
fit in a worldwide context? Molecular Phylogenetics and Evolution. 95: 67–78.

675   Weiss, S.L. & Dubin, M. 2018. Male mate choice as differential investment in contest
competition is affected by female ornament expression. Current Zoology. 64. 335-344.

Wheeler, J. 2008. Sexual Selection and Female Ornamentation in a Role-reversed Dance Fly.
678   M.Sc. thesis. University of Toronto, Toronto, Canada

Wheeler, J., Gwynne, D. T., & Bussière, L. F. 2012. Stabilizing sexual selection for female
ornaments in a dance fly. Journal of Evolutionary Biology. 25: 1233–1242.

681   Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York

Yong, L., Lee, B. & McKinnon, J.S. 2018. Variation in female aggression in 2 three-spined stickleback populations with female throat and spine coloration. Current Zoology. 64: 345-350.

Figure 1. Micrographs of nail varnish eye casts and female front, middle and hind legs. Each of the 8 species included in the comparative analysis is represented. Images are taken at different magnifications; black scale bars to the right of the respective female legs represent 1 mm, while grey scale bars to the right of the respective eye casts represent 0.1 mm.

Figure 2. Posterior distributions of fixed effects from a phylogenetically controlled model of exaggerations of male dorsal acute zones as a function of presence or absence of female pinnate scales in Empidine dance flies. **A)** posterior for the intercept term and **B)** posterior for the effect of the species having pinnate scales. Shaded areas under the curve depict the proportions of the distribution < 0 (in panel A) or > 0 (in panel B). Horizontal error bars show the 89% HPD intervals. Inset text above each panel give the interval ranges to three significant figures. Trace plots of MCMC chains are shown in figure S5 along with traces of phylogenetic effects.

Figure 3. Posterior distributions of fixed effects from a phylogenetically controlled model of exaggerations of dorsal acute zones in males and females as a function of the degree of female ornamentation and sex. **A)** posterior for the intercept term, **B)** posterior for the effect of species level ornamentation, **C)** posterior for the effect of sex, and **D)** posterior for the interaction effect of species level ornamentation and sex. Shaded areas under the curve depict the proportions of the distribution < 0 (**A** and **B**) or > 0 (**C** and **D**). Horizontal error bars show the 89% HPD intervals. Inset text above each panel give the interval ranges to three significant figures. Trace plots of MCMC chains are shown in figure S6 along with traces of phylogenetic effects.

Figure 4. The relationship between eye exaggeration (the standardised difference between dorsal facet sizes and ventral facet sizes), for males and females, and scaled species level mean female ornamentation (as measured by standardised leg area) for 8 species of dance flies (Diptera: Empididae). Red points with error bars represent response estimates and their error from phylogenetically controlled linear models. Dashed lines represent fitted lines through the response estimates for each species and reflect the effects in figure 3. The points represent individual values for male eyes from each species. To visualise the standard error around the mean of female ornamentation for each species, we have horizontally spread points for each species in proportion to the uncertainty in species mean.