An integrated single-cell atlas of the limb skeleton from development through adulthood

Tim Herpelinck †[1,2], Liesbeth Ory †[1,2], Tom Verbraeken[1,2], Gabriele Nasello[1,2], Mojtaba
Barzegari[3], Johanna Bolander[4], Frank P. Luyten[1,2], Przemko Tylzanowski #[1,2,5] and Liesbet
Geris*#[1,2,3,6]

[1] Skeletal Biology and Engineering Research Center, KU Leuven, Leuven, Belgium

[2] Prometheus, Division of Skeletal Tissue Engineering, KU Leuven, Leuven, Belgium

[3] Biomechanics Section, Department of Mechanical Engineering, KU Leuven, Leuven, Belgium

[4] Wake Forest Institute for Regenerative Medicine, Wake Forest School of Medicine, Winston-Salem,
7 NC, USA

[5] Department of Biomedical Sciences, Laboratory of Molecular Genetics, Medical University of
Lublin, Lublin, Poland

[6] Biomechanics Research Unit, GIGA In Silico Medicine, University of Liège, Liège, Belgium

† These authors contributed equally to this work and share first authorship

# These authors share senior authorship

*Correspondence:
Liesbet Geris
liesbet.geris@kuleuven.be

Abstract

The recent growth of single-cell transcriptomics has turned single-cell RNA sequencing (scRNA-seq)
into a near-routine experiment. Breakthroughs in improving scalability have led to the creation of
organism-wide transcriptomic datasets, aiming to comprehensively profile the cell types and states
within an organism throughout its lifecycle. To date, however, the skeleton remains a majorly
underrepresented organ system in organism-wide atlases. Considering how the skeleton not only serves
as the central framework of the vertebrate body but is also the home of the hematopoietic niche and a
central player in major metabolic and homeostatic processes, this presents a major deficit in current
reference atlas projects. To address this issue, we integrated ten separate scRNA-seq datasets
containing limb skeletal cells and their developmental precursors, generating an atlas of 133 332 cells.
This limb skeletal cell atlas describes cells across the mesenchymal lineage from the induction of the
limb to the adult bone and encompasses 39 different cell states. Furthermore, expanding the repertoire
of available time points and cell types within a single dataset allowed for more complete analyses of
cell-cell communication or *in silico* perturbation studies. Taken together, we present a missing piece
in the current atlas mapping efforts, which will be of value to researchers in the fields of skeletal
biology, hematopoiesis, metabolism and regenerative medicine.

Introduction

34   The skeleton is a highly advanced organ with a wide variety of functions, ranging from protection of
35   the internal organs and supporting locomotion to calcium homeostasis, housing the hematopoietic

system and serving an endocrine function. In addition, some skeletal tissues possess remarkable regenerative properties, with bone being able to spontaneously regenerate after fracture with minimal scar formation. While complex in its functions, the skeletal system is composed of a surprisingly low number of cell types, suggesting that its functional variety is a trait obtained through extensive intrinsic heterogeneity of cell states allowing advanced regionalized specialization.

Advancement towards ultra-high throughput single-cell RNA sequencing (scRNA-seq) platforms and the concomitant development of computational algorithms required to analyze the data, permit the generation of organism-wide transcriptome maps, resolved in time and space (1,2). If a reference atlas would be available, new datasets could be annotated automatically permitting a fast, data-driven and consistent cell annotation (3–5). Unfortunately, the skeleton is under-represented in most of these atlases, often with insufficiently detailed annotation of the skeletal lineage. Additionally, the skeletal biology research community generated scRNA-seq datasets but focused on very specific cell subpopulations. As a result, we find a significant discrepancy in the transcriptional characterization of the skeletal system. Large atlases contain a wealth of data but are left largely unexplored due to the insufficiently precise annotation while the specialized datasets remain confined to their domain of study. To bridge this gap, we reannotated and integrated specialized skeletal datasets (Supplementary Table 1) into a single framework comprising 133 332 cells (6–15). This reannotation was performed to ensure consistency across all datasets, using the original labels and enhancing them with more detailed annotations through the use of marker genes, resulting in the Limb Skeletal Cell Atlas (LSCA).

Following the generation of the LSCA, we explored its applicability and predictive value. We demonstrated the potential of the LSCA in pseudotemporal trajectory inference and intercellular communication analysis. The predictive capacity of the LSCA was further tested by simulation of Sox9 inactivation and the analysis of its consequences, which were in line with the *in vivo* phenotype. These results support the notion that our LSCA can be used, amongst others, as a reliable reference for the automated annotation of new skeletal scRNA-seq datasets. To facilitate accessibility, all notebooks required to build the atlas are available online.

Results

Dataset annotation and integration to produce a limb skeletal cell atlas

To build a comprehensive reference atlas of the limb skeleton, we selected ten publicly available mouse scRNA-seq datasets containing cells from the onset of limb development to mature bone. The individual datasets were manually reannotated based on the original labels and canonical marker gene expression (Supplementary Table 2). A comprehensive analysis was conducted on a total of 133 332 individual cells that met the quality control filters and led to the identification of 39 distinct cell clusters (Fig. 1 a). Twenty-six clusters originated from a mesenchymal lineage and encompassed mesenchyme, chondrocyte, osteoblast/osteocyte or fibroblast cell states. A further four states were associated with the muscle lineage and other clusters included hematopoietic (n=3), endothelial (n=2), epithelial (n=2), ectodermal (n=1) and neuronal (n=1) cell states.

During single-cell extraction, all spatial data is lost. A pseudospatial context within the cells of the early limb bud was reintroduced utilizing distinct gene expression patterns. Specifically, we used the expression of the Hox gene family to delineate cells along the proximal-distal axis (Hoxa/d9-13) (16). A subset of ectodermal cells situated at the distal tip of the limb bud, called Apical Ectodermal Ridge (AER), is a signaling center associated with the proximal-distal limb growth. It could be identified by

the mRNA expression of Fgf8 (17). Cells at the distal-posterior margin of the limb bud exhibited the
expression of Shh, designating it as the zone of polarizing activity (ZPA) (18).

Following annotation, we applied both an unsupervised (single-cell Variational Inference; scVI)
(19,20) and a semi-supervised (single-cell Annotation using Variational Inference; scANVI) (19,21–
24) integration model. The scVI model allowed us to infer the underlying structure of the data without
prior labels, while scANVI leveraged both labeled and unlabeled data to improve annotation accuracy.
To assess the effectiveness of these integration methods, we employed metrics focused on batch
correction and biological conservation (scIB) (19,25) (Supplementary Fig. 2). These metrics helped us
evaluate how well the integration methods corrected for batch effects while preserving the biological
variability inherent in the data. Notably, scANVI demonstrated the most favorable overall
performance, effectively integrating the datasets without introducing cross-laboratory batch effects
(Supplementary Fig. 3). This robust performance of scANVI ensured that the integration was cell-
based, accurately representing 39 distinct cell states (Fig. 1a-b). This comprehensive integration
provides a reliable foundation for downstream analyses and applications.

Virtual reconstruction of the growth plate

Long bones are formed through the process of endochondral bone formation. Initially, mesenchymal
cells condense and then undergo a series of differentiation steps going from resting chondrocytes over
proliferating to pre-hypertrophic and hypertrophic chondrocytes. These cells form columnar structures
that together make up the growth plate   (26). The hypertrophic chondrocytes then either
transdifferentiate into osteoblasts or undergo apoptosis and the space is replaced by the osteoblasts
producing bone tissue.

As a part of the validation of our atlas, we virtually restored the growth plate. To accomplish that, we
used pseudotime analysis to reconstruct the transcriptional trajectories from resting to hypertrophy
zone *in silico* (Fig. 2a). Cells were then binned by pseudotime and for each bin, we performed
dimensionality reduction by t-distributed stochastic neighborhood embedding (t-SNE). Importantly,
we imposed a circle as a boundary condition for each t-SNE. Alignment of these circular projections
then recreated the cylindrical shape of the growth plate, while also visualizing transcriptional
heterogeneity within the growth plate across bins of pseudotime (Fig. 2b). As such, plotting gene
expression along the pseudotime axis combined with expression in pseudospace artificially restores, to
a certain extend, the tissue architecture (Fig. 2c).

Intercellular communication inference across the skeleton

Intercellular communication driven by ligand-receptor interactions is pivotal for cellular
differentiation. Therefore, a wide variety of tools to infer intercellular signaling from scRNA-seq data
has been developed (27). Here, we used CellPhoneDB (28) as it considers multimeric receptor
complexes when inferring ligand-receptor interactions. We analyzed Bone Morphogenetic Protein
(BMP), Growth Differentiation Factor 5 (GDF5) and Sonic Hedgehog (SHH) signaling taking place
within the limb bud, the Apical Ectodermal Ridge (AER) and the Zone of Polarizing Activity (ZPA).
BMPs are members of the Transforming Growth Factor ligand superfamily and signal through a
tetrameric receptor complex. Type I and Type II BMP receptors are expressed across the limb
mesenchyme (29,30) and AER (29), which also acts as a source of BMPs (31). However, due to the
multimeric character of the receptor complexes resulting in many different combinations, it is
challenging to experimentally determine which ligand-receptor pair is predominantly used. Screening
with CellPhoneDB narrowed down the number of possible combinations. The analysis revealed that
the AER mainly signals to the limb bud through BMP4 and BMP7 (Fig. 3, Supplementary Fig. 4-6)

and that they appear to favor binding a receptor complex containing BMR1A. BMP signaling towards
the AER from the limb mesenchyme is known to be limited (31), as we confirm in our data. Noteworthy
is that the favored ligand-receptor combination is invariable over time. In addition, the AER is
predicted to be capable of autocrine signaling, which decreases over time (Fig. 3, Supplementary Fig.
4-6).

Previous experiments show that GDF5 binds to different sets of type I receptors. This interaction is
essential for the differentiation of digit cartilage and synovial joints (32). Our data shows significant
expression from Joint Precursors towards the Mesenchyme and the Chondroprogenitors (Fig. 3).

Antero-posterior outgrowth and proximodistal polarization are interconnected to each other by an SHH
epithelial-mesenchymal feedback loop (33). SHH expression decreases as the limb bud grows which
explains the decrease in cell-cell communication from E10.5 to E11.5 and no SHH expression in E12.5.
SHH signaling is regulated during limb development at many levels and one of them is a negative
feedback loop involving binding of SHH to HHIP which acts as a decoy receptor to inhibit HH
signaling. As a part of the LSCA validation, we investigated that regulatory loop and showed that this
cell-cell communication is mostly limited to intermediate limb bud mesenchyme and
chondroprogenitors (Fig. 3).

Simulation of transcription factor perturbation

Developmental biology was one of the key fields of research for exploring gene regulatory networks
(GRNs). Traditionally, this involved a series of experiments where transcription factor activity was
altered by gain- or loss-of-function and the resulting *in vivo* effects were analyzed. With the advances
in bioinformatics, CellOracle (34) algorithm was designed to study GRN inference from scRNA-seq
data. In addition, it allows to explore the *in silico* effects of the perturbation of transcription factor
expression on target gene expression. To test if our dataset was amenable to this type of analysis, we
decided to compare an *in silico* prediction to published *in vivo* data (35).

Sox9 is a transcription factor essential for maintaining the proliferation of columnar chondrocytes and
preventing their premature transdifferentiation into osteoblasts. It also plays a crucial role in inducing
chondrocyte hypertrophy, making it indispensable for the proper formation and maintenance of
functional growth plates. Inactivation of Sox9 leads to a shortening of the columnar and hypertrophic
zones in the growth plate and accelerates ossification due to premature pre-hypertrophy and matrix
mineralization (11,36). To assess the usefulness of our atlas for functional studies, we applied the
CellOracle algorithm to create a virtual Sox9 knockout and compared our *in silico* predictions to data
from an *in vivo* knockout study (Fig. 4a,b). We identified regions within the dataset that are
developmentally accessible or inaccessible depending on Sox9's regulatory influence. Accessible cell
states are those that cells can develop into under the current regulatory conditions, while inaccessible
states cannot be reached due to the absence or inactivation of essential transcription factors.

The simulations show that upon removal of Sox9, most mesoderm-derived cells fail to contribute to
cartilage formation, while myogenesis, osteogenesis, and endothelial differentiation remain largely
unaffected (Fig. 4c, Supplementary Fig. 7). These results align with *in vivo* findings, where Sox9
inactivation prevents mesodermal cells from differentiating into chondrocytes, leading to an
accumulation of cartilage precursors and osteogenic cells (Fig. 4d). For example, in the virtual
knockout model, cell type 11—pre-hypertrophic chondrocytes—is represented as a blue region,
indicating that mesenchymal cells cannot access this cell state. This observation is consistent with *in*
vivo data, where this cell type is exclusively present in the wild-type (blue) and absent in the knockout.

A limitation of the computational approach to create a virtual knockout is that it accounts only for
developmental flow and not lineage dependency. As a result, it will mark a region as an inaccessible
cell state if its development depends on Sox9, however, it will mark regions relying on this
(inaccessible) cell type as accessible states. For instance, cell type 12—hypertrophic chondrocytes—is
shown as a red region in the virtual knockout, indicating that this cell state is predicted to be present in
Sox9-perturbed situations. However, *in vivo* data show that this cell type is absent in the knockout
(blue) because it originates from pre-hypertrophic cells.

Discussion

We have generated a Limb Skeletal Cell Atlas representing a manually curated compendium of 133
332 cells across ten datasets and explored its applicability in both data- and hypothesis-driven analyses.
First, we presented the unsupervised construction of a spatiotemporal map of the growth plate, which
correctly recapitulates the molecular cascades of growth plate development, by imposing boundary
conditions on the gradient descent of t-SNE dimensionality reduction. Similar results have been
obtained previously by warping the principal component space (37).

Next, analysis of intercellular communication by BMP signaling was able to shed light on a
longstanding question in the field of limb development. It is known that BMP signaling is required for
AER regression, but not whether the AER, the mesenchyme or both act as the source of those BMPs
(31). Based on CellPhoneDB's cell-cell communication inference, considering the subunit architecture
of both ligands and heteromeric receptors (27,28), our case study suggests the AER to be the dominant
source. This finding does, however, require further *in vivo* validation.

Finally, we demonstrated that the *in silico* inactivation of Sox9 corroborates *in vivo* results confirming
the inability of Sox9-negative cells to differentiate into chondrocytes (11,38). Given that we can assess
the developmental outcome of computationally simulated transcription factor knockouts, the LSCA
represents a valuable resource for both limb developmental biologists and skeletal biology researchers.

The current release is restricted to the healthy murine skeleton. While our study provides valuable
insights, it is important to acknowledge certain limitations. Specifically, there are gaps in the limb sites
and stages that were sampled, as well as differences in genetic background, sex, and the enrichment of
sub-populations. These factors may influence the census and its potential downstream applications.
However, these limitations also highlight opportunities for future research to build upon our findings,
addressing these gaps and further enhancing our understanding of the subject. We will continuously
update and extend the LSCA, as data availability permits, toward the entire limb skeleton in health and
disease. The goal of this study was to provide an initial reference of the musculoskeletal system of the
limb. In future work, this first version will serve as the basis of a collaborative effort to complete the
atlas. All data, analyses and code to replicate the figures are freely available. Our web portal is designed
to be intuitive and to allow browsing of the data at a glance. In short, the Limb Skeletal Cell Atlas
provides a framework with characterization of most known cell populations in the limb skeleton and
represents a foundation for future studies in a wide variety of disciplines.

Materials and Methods

Data preprocessing

204 Publicly available datasets that were not aligned, were first mapped with Cell Ranger (39). Initial
205 Quality control of the raw count matrices was then performed for each dataset using Scater (40).
206 Filtering was performed based on the library size and number of expressed genes per cell. Cells more

than three standard deviations away from the median of either metric were filtered out. Subsequently, cells where the mitochondrial fraction of reads exceeded a proportion higher than three standard deviations from the median mitochondrial fraction were removed. Finally, low-abundance genes (expression lower than 1E-3) were filtered out and duplicate rows were removed if present.

Clustering and dimensionality reduction

All datasets were analyzed individually prior to integration. The filtered count matrices were imported into Seurat v4 (4). Normalization was performed using the LogNormalize parameter and a scale factor of 1E4. Subsequently, the data was centered and scaled using all genes followed by a calculation of principal components (PCs) using the top 2000 highly variable genes selected by the "vst" method. The optimal number of PCs to construct the Shared Nearest Neighbor (SNN) graph was visually determined based on the elbow plot and varied for each dataset. Clustering was performed using the Louvain algorithm. The resolution was adapted to the individual dataset, where we defined the optimal number of clusters as the maximum number of cell states that could confidently be labeled based on marker gene expression. These marker genes were obtained from the FindMarkers function using the default settings.

Integration with scvi-tools

The integrated atlas was constructed using scVI (20) and scANVI (21–24). For preprocessing the datasets, we filtered for the 5000 most variable genes. First, we used scVI as an unsupervised tool to find common axes of variation between the datasets. The parameters were used as described in the scvi-tools (19) tutorial of '*Atlas-level integration of lung data*'. Next, to compute a more accurate integration result, we used scANVI as a semi-supervised tool to integrate the data. scANVI used the trained scVI model to initialize the integration.

Integration metrics

To calculate integration metrics, the scIB-metrics package was used (25), which contains scalable implementations of the metrics used in the scIB benchmarking suite (41–44).

Virtual growth plate reconstruction

We first subsetted the atlas for growth plate chondrocytes: Resting zone chondrocytes, Proliferative chondrocytes, Pre-hypertrophic chondrocytes and Hypertrophic chondrocytes. This subset within the latent space of the scANVI integration was then passed to Monocle 3 (45). Clustering was performed on the resulting CellDataSet (CDS) object using the default parameters. The trajectory graph was learned on the monocle-derived clusters by calling learn_graph. Roots were manually chosen with order_cells. The resulting pseudotime was added to the metadata of the growth plate subset within the latent space of scANVI and used as input for the growth plate reconstruction. Cells were binned by pseudotime using the cut function from- pandas (46,47). Upon each bin, we performed dimensionality reduction by t-distributed stochastic neighborhood embedding (t-SNE). We imposed a circle as a boundary condition on the gradient descent function of each t-SNE. With the concatenation function, we were able to stack the circular projections in an ordered way to recreate the cylindrical shape of the growth plate.

Cell-cell interaction prediction

Prediction of cell-cell communication by ligand-receptor interactions between cell types was performed using CellPhoneDB v5 (28). The atlas was subsetted for each developmental time point

248 before downstream cell-cell communication analysis. For each subset, the translate function was used
249 to humanize the data and normalized. Differentially expressed genes were computed for each cell type
250 and used as input for ligand-receptor pair calculations.

Knockout simulation

For *in silico* knockout experiments in CellOracle (34) the atlas was subsetted to only include time
points E10.5-P21 and relevant cell types (Periosteal progenitors, Periosteum, Endosteum, Resting zone
chondrocytes, Proliferative chondrocytes, Pre-hypertrophic chondrocytes, Hypertrophic chondrocytes,
Chondrocytes, Chondroprogenitors, Osteoprogenitors, Pre-osteoblasts, Osteoblasts, Osteocytes,
Vascular endothelial cells, Lymphatic endothelial cells, Proximal limb bud mesenchyme, Intermediate
limb bud mesenchyme, Distal limb bud mesenchyme, Myogenic stem cells, Muscle progenitors and
Skeletal muscle cell). This subset within the latent space of the scANVI integration of the atlas was
then passed to Monocle 3 (45). Clustering was performed on the resulting CellDataSet (CDS) object
using the default parameters. The trajectory graph was learned on the monocle-derived clusters by
calling learn graph. Roots were manually chosen with order cells. The resulting pseudotime was added
to the metadata of the knockout subset within the latent space of scANVI and used as input for the
virtual knockout. To reduce the amount of computational time and resources required by a large
dataset, 20,000 cells were randomly selected and only highly variable genes (n=5000) were included.
For gene regulatory network (GRN) inference, we used the built-in base GRN made from the mouse
sci-ATAC-seq atlas (48) Following k nearest neighbors (KNN) imputation based on the first 27 PCs,
GRNs were imputed for each cluster. To simulate the knockout of a transcription factor, its expression
was set to 0. After this knockout, GRN inference was performed again. Signal perturbation propagation
and transition probabilities were calculated using the standard settings. Visualization of the pseudotime
gradient, simulation vector field and their inner product was performed as described in the CellOracle
online documentation.

***In vivo* Sox9 knockout and WT data analysis and integration**

Both wild type (WT) and Sox9 knockout (KO) datasets were preprocessed separately to ensure high-
quality data for subsequent analyses. The preprocessing steps included quality control measures to
remove low-quality cells, as described in section 4.1. Following preprocessing, cells from both WT
and Sox9 KO datasets were clustered and labeled independently. Clustering was conducted to identify
distinct cell populations within each condition, and labels were assigned based on known cell markers,
as described in section 4.2.
279
To integrate the WT and Sox9 KO datasets, the Harmony package (41) was employed. Harmony
facilitates the integration of multiple datasets by correcting batch effects and aligning data across
different conditions. This integration enabled a comprehensive comparison of cell populations and
gene expression profiles between WT and Sox9 KO conditions.

Data and Code Availability Statement

The datasets analyzed for this study can be found in Supplementary Table 1. The Skeletal Cell Atlas
can be downloaded or interactively explored at www.skeletalcellatlas.org. All code used to perform
the analyses and notebooks to generate the figures is available at
https://github.com/TElabSBE/LimbSkeletalCellAtlas.

Figure Legends

Figure 1. An integrated compendium of skeletal cell types with detailed annotation. a UMAP visualization of the scANVI latent space of 133 332 murine limb mesenchyme- and skeleton-derived cells, colored by annotation. b, Dot plot showing the expression of one selected marker gene per cluster. The color of the dot represents the mean expression level and its size represents the percentage of cells within the cluster in which that gene was detected. BMSCs: Bone Marrow-derived Stem Cells, ZPA: Zone of Polarizing Activity, AER: Apical Ectodermal Ridge.

Figure 2. Pseudospatiotemporal reconstruction of the transcriptional dynamics within the growth plate. a, UMAP visualization of the scANVI latent space of the integrated growth plate data subset from the main atlas, colored by annotation (left) and Monocle 3 pseudotime value (right). b, Growth plate chondrocytes were grouped in 50 bins based on similar pseudotime values. t-SNE dimensional reduction was performed on each bin, using a circle with a radius of 20 as the boundary condition for gradient descent, thus recreating the cylindrical shape of the growth plate upon stacking the bins. c, Known marker gene expression for different zones of the growth plate in pseudotime and - space.

Figure 3. BMP, GDF5 and SHH signaling case study between AER, ZPA and mesenchyme. Dotplot of predicted ligand-receptor complex interactions in developmental timepoints E10.5, E11.5 and E12.5. The sender/receiver cell types are represented on the y-axis, while the x-axis represents the ligand/receptor pairs. The color of the dot represents the mean gene expression, while the size of the dot is proportional to the percentage of cells of each cell type expressing the gene. Translucency represents the interaction specificity and if the gene is differentially expressed an outer red ring is shown. DEG: differentially expressed genes, AER: Apical Ectodermal Ridge, PLBM: Proximal limb bud mesenchyme, ILBM: Intermediate limb bud mesenchyme, DLBM: Distal limb bud mesenchyme, CP: Chondroprogenitors, RZC: Resting zone chondrocytes, PC: Proliferative chondrocytes, PHC: Pre-hypertrophic chondrocytes, JP: Joint precursors, ZPA: Zone of Polarizing Activity.

Figure 4. Sox9 knockout (KO) *in silico* predictions compared to *in vivo* data. a, Reference annotation of the subsetted atlas used for knockout analysis. b, a UMAP plot integrating WT and Sox9 KO *in vivo* data. c, Inner vector product of cell trajectories following *in silico* Sox9 perturbation and the pseudotime gradient from unperturbed conditions. Red regions indicate accessible cell states, while blue regions represent inaccessible cell states. d, UMAP of integrated Sox9 KO and WT *in vivo* data, colored by condition. Red corresponds to Sox9 KO data, while blue represents WT data. WT: Wild-type, KO: knockout.

Supplementary Data Figure Legends

Supplementary Figure 1. Individual datasets after reannotation. a-j, UMAP visualization of the scANVI latent space colored by cell type for the separate datasets with database ID: a, 34260921 b, 36175067, 37873464 c, 30250253 d, 31874220 e, 33297480 f, 33597301 g, 31543445 h, 38479598 i, 31130381 j, 36777346

Supplementary Figure 2. Assessment of integration method. a, UMAP visualization of the scANVI latent space (left) and the scVI latent space (right). b, Results of the metrics calculated with scIB. Bio conservation was captured with the use of classical label conservation metrics, which assess local neighborhoods (graph cLISI, extended from cLISI), global cluster matching (Adjusted Rand Index: ARI, normalized mutual information: NMI and a metric evaluating rare cell identity annotations (isolated label scores). Batch correction scores were measured via the k-nearest-neighbor batch effect test (kBET), k-nearest-neighbor (kNN) graph connectivity and the average silhouette

width (ASW) across batches. Independently of cell identity labels, batch removal is measured using
the graph integration local inverse Simpson's Index (graph iLISI, extended from iLISI) and PCA
regression. Aggregate scores represent the mean score of the Bio conservation and Batch correction
metrics.

Supplementary Figure 3. Assessment of batch effects influencing integration. a-b, UMAP
visualization of the scANVI latent space colored by study (a) and developmental stage or age (b).

Supplementary Figure 4. BMP, GDF5 and SHH signaling case study between AER, ZPA and
mesenchyme for developmental timepoint E10.5. Dotplot of predicted ligand-receptor complex
interactions in developmental timepoints E10.5. The sender/receiver cell types are represented on the
y-axis, while the x-axis represents the ligand/receptor pairs. The color of the dot represents the mean
gene expression, while the size of the dot is proportional to the percentage of cells of each cell type
expressing the gene. Translucency represents the interaction specificity and if the gene is
differentially expressed an outer red ring is shown. DEG: differentially expressed genes, AER:
Apical Ectodermal Ridge, PLBM: Proximal limb bud mesenchyme, ILBM: Intermediate limb bud
mesenchyme, DLBM: Distal limb bud mesenchyme, CP: Chondroprogenitors, RZC: Resting zone
chondrocytes, PC: Proliferative chondrocytes, PHC: Pre-hypertrophic chondrocytes, ZPA: Zone of
Polarizing Activity.

Supplementary Figure 5. BMP, GDF5 and SHH signaling case study between AER, ZPA and
mesenchyme for developmental timepoint E11.5. Dotplot of predicted ligand-receptor complex
interactions in developmental timepoints E11.5. The sender/receiver cell types are represented on the
y-axis, while the x-axis represents the ligand/receptor pairs. The color of the dot represents the mean
gene expression, while the size of the dot is proportional to the percentage of cells of each cell type
expressing the gene. Translucency represents the interaction specificity and if the gene is
differentially expressed an outer red ring is shown. DEG: differentially expressed genes, AER:
Apical Ectodermal Ridge, PLBM: Proximal limb bud mesenchyme, ILBM: Intermediate limb bud
mesenchyme, DLBM: Distal limb bud mesenchyme, CP: Chondroprogenitors, RZC: Resting zone
chondrocytes, PC: Proliferative chondrocytes, PHC: Pre-hypertrophic chondrocytes, JP: Joint
precursors.

Supplementary Figure 6. BMP, GDF5 and SHH signaling between AER and mesenchyme for
developmental timepoint E12.5. Dotplot of predicted ligand-receptor complex interactions in
developmental timepoints E12.5. The sender/receiver cell types are represented on the y-axis, while
the x-axis represents the ligand/receptor pairs. The color of the dot represents the mean gene
expression, while the size of the dot is proportional to the percentage of cells of each cell type
expressing the gene. Translucency represents the interaction specificity and if the gene is
differentially expressed an outer red ring is shown. DEG: differentially expressed genes, AER:
Apical Ectodermal Ridge, PLBM: Proximal limb bud mesenchyme, ILBM: Intermediate limb bud
mesenchyme, DLBM: Distal limb bud mesenchyme, CP: Chondroprogenitors, RZC: Resting zone
chondrocytes, PC: Proliferative chondrocytes, PHC: Pre-hypertrophic chondrocytes, JP: Joint
precursors, ZPA: Zone of Polarizing Activity.

Supplementary Figure 7. **Sox9 knockout (KO***) in silico* **predictions**. a, Vector field graph
indicating cell trajectories after Sox9 *in silico* knockout. b, Gradient of pseudotime in the absence of
perturbation. c, Inner vector product of cell trajectories following *in silico* Sox9 perturbation and the
pseudotime gradient from an unperturbed condition.

Supplementary Table Legends

Supplementary Table 1: scRNA-seq datasets used to construct the Limb Skeletal Cell Atlas.

Supplementary Table 2: List of genes used to annotate clusters in the Limb Skeletal Cell Atlas.

Author Contributions

TH, JB, FL, PT and LG conceptualized the study; TH, PT and LG designed the study; TH, LO and TV provided adapted code for the analysis of the datasets and downstream analysis of the LSCA; TH, LO and TV contributed to the analysis of the datasets and the LSCA; GN provided a docker work environment; MB adapted and optimized the code to create an interactive web application, TH and LO wrote the first draft of the manuscript and all the other authors contributed to the article. All authors approved the submitted version.

Acknowledgments

We would like to thank Inge Van Hoven for assistance and Ronny Moreas for technical support in the deployment of the web app. Research was funded by the Research Foundation Flanders (FWO Vlaanderen) T.H.: 1S80021N, G.N.: 12C5923N, MatheMorphosis: G0D3420N, and FWO-INSITE: G085018N; by the European Union's Horizon 2020 Framework Program (H2020/2014-2021) ERC INSITE (772418) and SC1-DTH ISW (101016503); by the European Union's Horizon Europe Framework Program (HEU/2022-2027) ERC INSTant CARMA (101088919), as well as by the special research fund of the KU Leuven (C24/17/077). The funders had no role in study design, data collection and analysis, the decision to publish, or the preparation of the manuscript. This work is part of Prometheus, the KU Leuven R&D division for skeletal tissue engineering (http://www.kuleuven.be/prometheus).

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Asp M, Giacomello S, Larsson L, Wu C, Fürth D, Qian X, et al. A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing Human Heart. Cell. 2019;179 (7).

2. Elmentaite R, Kumasaka N, Roberts K, Fleming A, Dann E, King HW, et al. Cells of the human intestinal tract mapped across space and time. Nature. 2021;597(7875).

3. Fischer DS, Dony L, König M, Moeed A, Zappia L, Heumos L, et al. Sfaira accelerates data and model reuse in single cell genomics. Genome Biol. 2021;22(1).

4. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. Cell. 2021;184(13).

5. Pasquini G, Rojo Arias JE, Schäfer P, Busskamp V. Automated methods for cell type annotation on scRNA-seq data. Vol. 19, Computational and Structural Biotechnology Journal. 2021.

6.   Agoro R, Nookaew I, Noonan ML, Marambio YG, Liu S, Chang W, et al. Single cell cortical
bone transcriptomics define novel osteolineage gene sets altered in chronic kidney disease.
Front Endocrinol (Lausanne). 2023;14.

7.   Baryawno N, Przybylski D, Kowalczyk MS, Kfoury Y, Severe N, Gustafsson K, et al. A
Cellular Taxonomy of the Bone Marrow Stroma in Homeostasis and Leukemia. Cell.
2019;177(7).

8.   Böhm AM, Dirckx N, Tower RJ, Peredo N, Vanuytven S, Theunis K, et al. Activation of
Skeletal Stem and Progenitor Cells for Bone Regeneration Is Driven by PDGFRβ Signaling.
Dev Cell. 2019;51(2).

9.   Debnath S, Yallowitz AR, McCormick J, Lalani S, Zhang T, Xu R, et al. Discovery of a
periosteal stem cell mediating intramembranous bone formation. Nature. 2018;562(7725).

10. Desanlis I, Paul R, Kmita M. Transcriptional trajectories in mouse limb buds reveal the
transition from anterior-posterior to proximal-distal patterning at early limb bud stage. J Dev
Biol. 2020;8(4).

11. Haseeb A, Kc R, Angelozzi M, de Charleroy C, Rux D, Tower RJ, et al. SOX9 keeps growth
plates and articular cartilage healthy by inhibiting chondrocyte dedifferentiation/ osteoblastic
redifferentiation. Proc Natl Acad Sci U S A. 2021;118(8).

12. Kelly NH, Huynh NPT, Guilak F. Single cell RNA-sequencing reveals cellular heterogeneity
and trajectories of lineage specification during murine embryonic limb development. Matrix
Biology. 2020;89.

13. Knights AJ, Farrell EC, Ellis OM, Lammlin L, Junginger LM, Rzeczycki PM, et al. Synovial
fibroblasts assume distinct functional identities and secrete R-spondin 2 in osteoarthritis. Ann
Rheum Dis. 2022;82(2).

14. Nookaew I, Xiong J, Onal M, Bustamante-Gomez C, Wanchai V, Fu Q, et al. Refining the
identity of mesenchymal cell types associated with murine periosteal and endosteal bone.
Journal of Biological Chemistry. 2024;300(4).

15. Sivaraj KK, Jeong HW, Dharmalingam B, Zeuschner D, Adams S, Potente M, et al. Regional
specialization and fate specification of bone stromal cells in skeletal development. Cell Rep.
2021;36(2).

16. Zakany J, Duboule D. The role of Hox genes during vertebrate limb development. Vol. 17,
Current Opinion in Genetics and Development. 2007.

17. Lewandoski M, Sun X, Martin GR. Fgf8 signalling from the AER is essential for normal limb
development. Nat Genet. 2000;26(4).

18. Panman L, Zeller R. Patterning the limb before and after SHH signalling. Vol. 202, Journal of
Anatomy. 2003.

19. Gayoso A, Lopez R, Xing G, Boyeau P, Valiollah Pour Amiri V, Hong J, et al. A Python
library for probabilistic analysis of single-cell omics data. Vol. 40, Nature Biotechnology.
2022.

20. Lopez R, Regier J, Cole MB, Jordan MI, Yosef N. Deep generative modeling for single-cell
transcriptomics. Nat Methods. 2018;15(12).

21. Consortium TTS, Quake SR. The Tabula Sapiens: a single cell transcriptomic atlas of
multiple organs from individual human donors. bioRxiv. 2021;

22. Kingma DP, Rezende DJ, Mohamed S, Welling M. Semi-supervised learning with deep
generative models. In: Advances in Neural Information Processing Systems. 2014.

23. Kingma DP, Welling M. Auto-encoding variational bayes. In: 2nd International Conference
on Learning Representations, ICLR 2014 - Conference Track Proceedings. 2014.

24. Xu C, Lopez R, Mehlman E, Regier J, Jordan MI, Yosef N. Probabilistic harmonization and
annotation of single-cell transcriptomics data with deep generative models. Mol Syst Biol.
2021;17(1).

25. Luecken MD, Büttner M, Chaichoompu K, Danese A, Interlandi M, Mueller MF, et al.
Benchmarking atlas-level data integration in single-cell genomics. Nat Methods. 2022;19(1).

26. Kronenberg HM. Developmental regulation of the growth plate. Vol. 423, Nature. 2003.

27. Armingol E, Officer A, Harismendy O, Lewis NE. Deciphering cell–cell interactions and
communication from gene expression. Vol. 22, Nature Reviews Genetics. 2021.

28. Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. CellPhoneDB: inferring
cell–cell communication from combined expression of multi-subunit ligand–receptor
complexes. Nat Protoc. 2020;15(4).

29. Kawakami Y, Ishikawa T, Shimabara M, Tanda N, Enomoto-lwamoto M, Iwamoto M, et al.
BMP signaling during bone pattern determination in the developing limb. Development.
1996;122(11).

30. Zou H, Wieser R, Massagué J, Niswander L. Distinct roles of type I bone morphogenetic
protein receptors in the formation and differentiation of cartilage. Genes Dev. 1997;11(17).

31. Pizette S, Niswander L. BMPs negatively regulate structure and function of the limb apical
ectodermal ridge. Development. 1999;126(5).

32. Baur ST, Mai JJ, Dymecki SM. Combinatorial signaling through BMP receptor IB and GDF5:
Shaping of the distal mouse limb and the genetics of distal limb diversity. Development.
2000;127(3).

33. Carballo GB, Honorato JR, De Lopes GPF, Spohr TCLDSE. A highlight on Sonic hedgehog
pathway. Vol. 16, Cell Communication and Signaling. 2018.

34. Kamimoto K, Hoffmann CM, Morris SA. CellOracle: Dissecting cell identity via network
inference and in silico gene perturbation. bioRxiv. 2020;

35.   Akiyama H, Chaboissier MC, Martin JF, Schedl A, De Crombrugghe B. The transcription
484   factor Sox9 has essential roles in successive steps of the chondrocyte differentiation pathway
485   and is required for expression of Sox5 and Sox6. Genes Dev. 2002;16(21).

486   36.   Dy P, Wang W, Bhattaram P, Wang Q, Wang L, Ballock RT, et al. Sox9 Directs
487   Hypertrophic Maturation and Blocks Osteoblast Differentiation of Growth Plate
488   Chondrocytes. Dev Cell. 2012;22(3).

489   37.   Li J, Luo H, Wang R, Lang J, Zhu S, Zhang Z, et al. Systematic Reconstruction of Molecular
490   Cascades Regulating GP Development Using Single-Cell RNA-Seq. Cell Rep. 2016;15(7).

491   38.   Bi W, Deng JM, Zhang Z, Behringer RR, De Crombrugghe B. Sox9 is required for cartilage
492   formation. Nat Genet. 1999;22(1).

493   39.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel
494   digital transcriptional profiling of single cells. Nat Commun. 2017;8.

495   40.   McCarthy DJ, Campbell KR, Lun ATL, Wills QF. Scater: Pre-processing, quality control,
496   normalization and visualization of single-cell RNA-seq data in R. Bioinformatics. 2017;33(8).

497   41.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and
498   accurate integration of single-cell data with Harmony. Nat Methods. 2019;16(12).

499   42.   Hubert L, Arabie P. Comparing partitions. J Classif. 1985;2(1).

500   43.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
501   Machine learning in Python. Journal of Machine Learning Research. 2011;12.

502   44.   Büttner M, Miao Z, Wolf FA, Teichmann SA, Theis FJ. A test metric for assessing single-cell
503   RNA-seq batch correction. Nat Methods. 2019;16(1).

504   45.   Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell
505   transcriptional landscape of mammalian organogenesis. Nature. 2019;566(7745).

506   46.   McKinney W. Data Structures for Statistical Computing in Python. In: Proceedings of the 9th
507   Python in Science Conference. 2010.

508   47.   The pandas development team. pandas-dev/pandas: Pandas. Zenodo; 2020.

509   48.   Cusanovich DA, Hill AJ, Aghamirzaie D, Daza RM, Pliner HA, Berletch JB, et al. A Single-
510   Cell Atlas of In Vivo Mammalian Chromatin Accessibility. Cell. 2018;174(5).

511   **Figure 1**

A

Figure 2

515 **Figure 3**

Figure 4

A

B

● 2 - Proximal limb bud mesenchyme	● 8 - Chondrocytes	● 17 - Osteoprogenitors	● 21 - Osteocytes	● 28 - Muscle progenitors
● 3 - Intermediate limb bud mesenchyme	● 9 - Resting zone chondrocytes	● 19 - Osteoblasts	● 22 - Periosteal progenitors	● 29 - Skeletal muscle cells
● 4 - Distal limb bud mesenchyme	● 10 - Proliferative chondrocytes	● 12 - Hypertrophic chondrocytes	● 23 - Periosteum	● 31 - Vascular endothelial cells
● 7 - Chondroprogenitors	● 11 - Pre-hypertrophic chondrocytes	● 13 - Articular chondrocytes	● 27 - Myogenic stem cells	● 32 - Lymphatic endothelial cells

C

Virtual Sox9 knock-out

accesible states

inaccesible states

D

In vivo conditional Sox9 knock-out

WT

Sox9 KO

519 **Supplementary Figure 1**

521 **Supplementary Figure 2**

523 **Supplementary Figure 3**

525 **Supplementary Figure 4**

Supplementary Figure 5

 Supplementary Figure 6

531 **Supplementary Figure 7**

533 **Supplementary Table 1**

Study (Database ID)	Study (PMID)	Study (First, Last Author)	Technology	Developmental stage
SRP278393	34260921	Sivaraj, Adams	10x	P21
PRJNA870944	36175067, 37873464	Maerz, Farrell	10x	12-16w
SRP122545	30250253	Debnath, Landau & Greenblatt	CEL-seq2	P7
SRP238355	31874220	Kelly, Guilak	10x	E11.5, E13.5, E15.5, E18,5
SRP285918	33297480	Desanlis, Kmita	10x	E10.5, E11.5, E12.5
SRP293998	33597301	Lefebvre	10x	P13
PRJEB33444	31543445	Bohm, Maes	Smart-seq2	E15.5
PRJNA1080020	38479598	OBrien	10x	8-14w
SRP188674	31130381	Baryawno, Regev & Scadden	10x	8-10w
PRJNA858479	36777346	White	10x	8w

534 **Supplementary Table 2**

Cell Type/ Cell State	Gene names	Gene ID
BMSCs	Lepr, Cxcl12, Pdgfra	16847, 20315, 18595

Proximal limb bud mesenchyme	Prrx1, Hoxa9, Hoxd9	18933, 15405, 15438
Intermediate limb bud mesenchyme	Prrx1, Hoxa11, Hoxd11	18933, 15396, 15431
Proximal limb bud mesenchyme	Prrx1, Hoxa13, Hoxd13	18933, 15398, 15433
ZPA	Shh	20423
Fast proliferating cells	Cdk1, Cdc20	12534, 107995
Chondroprogenitors	Prrx1, Sox9	18933, 20682
Chondrocytes	Sox9, Col2a1,	20682, 12824
Resting zone chondrocytes	Col2a1, Ucma, Pthlh	12824, 68527, 19227
Proliferative chondrocytes	Col2a1, Top2a, Pth1r	12824, 21973, 19228
Pre-hypertrophic chondrocytes	Col2a1, Ihh	12824, 16147
Hypertrophic chondrocytes	Col2a1, Col10a1	12824, 12813
Articular chondrocytes	Col2a1, Prg4	12824, 96875
Joint precursors	Gdf5, Wnt9a, (Sox9-)	14563, 216795, 20682
Tenocyte precursors	Prrx1, Tnmd, Scx	18933, 64103, 20289
Tenocytes	Tnmd, Scx	64103, 20289
Osteoprogenitors	Runx2, Sp7	12393, 170574
Pre-osteoblasts	Runx2, Sp7, Spp1	12393, 170574, 20750
Osteoblasts	Runx2, Bglap	12393, 12096
Fast proliferating osteoblasts	Cdk1, Cdc20, Bglap	12534, 107995, 12096

Osteocytes	Dmp1	13406
Periosteal progenitors	Prrx1, Postn, Ctsk	18933, 50706, 13038
Periosteum	Postn, Wnt16	50706, 93735
Endosteum	Postn, Wnt16	50706, 93735
Osteoclasts	Cd68, Ctsk	12514, 13038
Fibroblasts	S100a4, Dcn	20198, 13179
Myogenic stem cells	Pax3, Pdgfc	18505, 54635
Muscle progenitors	Pax7, Itm2a	18509, 16431
Skeletal muscle cells	Myhc, Tnnt1, Tnnt3	111671, 21955, 21957
Perivascular cells	Acta2, Rgs5	11475, 19737
Vascular endothelial cells	Emcn, Pecam1, Cdh5	59308, 18613, 12562
Lymphatic endothelial cells	Pecam1, Lyve1	18613, 114332
AER	Fgf8	14179
Epithelial cells	Epcam	17075
Immune cells	Ptprc	19264
Erythroid cells	Gypa, Car2	14934, 12349
Platelets	Pf4, Vwf	56744, 22371
Neurons	Mpz, Ednrb	17528, 13618
Keratinocytes	Krt5, Krt10	110308, 16661

A

Legend:

- 1 - BMSCs
- 2 - Proximal limb bud mesenchyme
- 3 - Intermediate limb bud mesenchyme
- 4 - Distal limb bud mesenchyme
- 5 - ZPA
- 6 - Fast proliferating cells
- 7 - Chondroprogenitors
- 8 - Chondrocytes
- 9 - Resting zone chondrocytes
- 10 - Proliferative chondrocytes
- 11 - Pre-hypertrophic chondrocytes
- 12 - Hypertrophic chondrocytes
- 13 - Articular chondrocytes
- 14 - Joint precursors
- 15 - Tenocyte precursors
- 16 - Tenocytes
- 17 - Osteoprogenitors
- 18 - Pre-osteoblasts
- 19 - Osteoblasts
- 20 - Fast proliferating osteoblasts

- 21 - Osteocytes
- 22 - Periosteal progenitors
- 23 - Periosteum
- 24 - Endosteum
- 25 - Osteoclasts
- 26 - Fibroblasts
- 27 - Myogenic stem cells
- 28 - Muscle progenitors
- 29 - Skeletal muscle cells
- 30 - Perivascular cells
- 31 - Vascular endothelial cells
- 32 - Lymphatic endothelial cells
- 33 - AER
- 34 - Epithelial cells
- 35 - Immune cells
- 36 - Erythroid cells
- 37 - Platelets
- 38 - Neurons
- 39 - Keratinocytes

B

A

B

| Growth plate | Resting zone chondrocytes | Proliferative chondrocytes | Pre-hypertrophic chondrocytes | Hypertrophic chondrocytes |

C

Average expression

Pthlh | Pth1r | Ihh | Col10a1

A

- 2 - Proximal limb bud mesenchyme
- 3 - Intermediate limb bud mesenchyme
- 4 - Distal limb bud mesenchyme
- 7 - Chondroprogenitors
- 8 - Chondrocytes
- 9 - Resting zone chondrocytes
- 10 - Proliferative chondrocytes
- 11 - Pre-hypertrophic chondrocytes
- 17 - Osteoprogenitors
- 19 - Osteoblasts
- 12 - Hypertrophic chondrocytes
- 13 - Articular chondrocytes
- 21 - Osteocytes
- 22 - Periosteal progenitors
- 23 - Periosteum
- 27 - Myogenic stem cells
- 28 - Muscle progenitors
- 29 - Skeletal muscle cells
- 31 - Vascular endothelial cells
- 32 - Lymphatic endothelial cells

C Virtual Sox9 knock-out

accesible states

inaccesible states

D *In vivo* conditional Sox9 knock-out

WT

Sox9 KO