Angiotensin blockade enhances motivational reward learning via enhancing striatal prediction error signaling and frontostriatal communication

Authors

Ting Xu[1,2], Xinqi Zhou[1,2], Jonathan W. Kanen[3,4], Lan Wang[1,2], Jialin Li[5], Zhiyi Chen[6,7], Ran Zhang[1,2], Guojuan Jiao[1,2], Feng Zhou[6,7], Weihua Zhao[2], Shuxia Yao[2], Benjamin Becker[1,2*]

Affiliations

[1] The Center of Psychosomatic Medicine, Sichuan Provincial Center for Mental Health, Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China, Chengdu, China

[2] MOE Key Laboratory for Neuroinformation, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China

[3] Department of Psychology, University of Cambridge, Cambridge, UK

[4] Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge, UK

[5] Max Planck School of Cognition, Leipzig, Germany

[6] Faculty of Psychology, Southwest University, Chongqing, China

[7] Key Laboratory of Cognition and Personality, Ministry of Education, China

*Correspondence to

Benjamin Becker: https://orcid.org/0000-0002-9014-9671

University of Electronic Science and Technology

Xiyuan Avenue 2006, 611731 Chengdu, China

mail: ben_@gmx.de

ABSTRACT

Adaptive human learning utilizes reward prediction errors (RPEs) that scale the differences between expected and actual outcomes to optimize future choices. Depression has been linked with biased RPE signaling and an exaggerated impact of negative outcomes on learning which may promote amotivation and anhedonia. The present proof-of-concept study combined computational modelling and multivariate decoding with neuroimaging to determine the influence of the selective competitive angiotensin II type 1 receptor antagonist losartan on learning from positive or negative outcomes and the underlying neural mechanisms in healthy humans. In a double-blind, between-subjects, placebo-controlled pharmaco-fMRI experiment, 61 healthy male participants (losartan, n=30; placebo, n=31) underwent a probabilistic selection reinforcement learning task incorporating a learning and transfer phase. Losartan improved choice accuracy for the hardest stimulus pair via increasing expected value sensitivity towards the rewarding stimulus relative to the placebo group during learning. Computational modelling revealed that losartan reduced the learning rate for negative outcomes and increased exploitatory choice behaviors while preserving learning for positive outcomes. These behavioral patterns were paralleled on the neural level by increased RPE signaling in orbitofrontal-striatal regions and enhanced positive outcome representations in the ventral striatum (VS) following losartan. In the transfer phase, losartan accelerated response times and enhanced VS functional connectivity with left dorsolateral prefrontal cortex when approaching maximum rewards. These findings elucidate the potential of losartan to reduce the impact of negative outcomes during learning and subsequently facilitate motivational approach towards maximum rewards in the transfer of learning. This may indicate a promising therapeutic mechanism to normalize distorted reward learning and fronto-striatal functioning in depression.

INTRODUCTION

Human learning is driven by reward prediction errors (RPEs) that signal the discrepancy between expected and actual outcomes. Computational approaches have closely linked RPEs to dopaminergic signaling in the midbrain-striatum circuitry and to motivation and reward seeking [1, 2]. Deficits in these domains, in particular amotivation and anhedonia, represent key symptoms of unipolar depression and dysregulated RPE signaling has been proposed as a potential underlying neurocomputational candidate mechanism [3]. Specifically, within a computational reinforcement learning (RL) framework, depressed individuals showed enhanced sensitivity to negative information while they concomitantly discounted positive feedback leading to reduced learning from positive events [4, 5]. On the neural level this learning bias was often accompanied by blunted RPE signaling in the ventral striatum and reduced fronto-striatal connectivity during reward feedback [6, 7]. These neural dysregulations have been associated with depressive symptom load, specifically anhedonia and persistent negative mood [8], and could predict anti-depressive treatment responses [9]. As such, distorted learning from negative and positive outcomes may play a key role in the pathophysiology of depression and may represent a promising target for novel antidepressive treatments.

Accumulating evidence suggests that the renin-angiotensin system (RAS) plays a key role in learning. Preclinical work in rodents and humans has utilized the selective competitive angiotensin II type 1 receptor (AT1R) antagonist losartan (LT) - an approved treatment for hypertension with an excellent safety record [10, 11] - to modulate learning from negative or positive events [12, 13]. Recent human studies have demonstrated that a single dose of LT selectively suppressed memory encoding of threatening materials [14] and accelerated threat extinction learning [15, 16]. Moreover, LT specifically affected probabilistic learning from negative outcomes by reducing the degree to which participants learned from loss feedback, while leaving learning from positive outcomes unaffected [12]. An initial neuroimaging study moreover reported modulatory effects of LT on mesocorticolimbic functional connectivity during social reward and punishment processing [17].

Given the pivotal role of dopamine (DA) in RPE signaling and modulation of the mesocorticolimbic circuits [18], these results may indicate a downstream effect of LT on DA signaling and in turn on learning from positive and negative outcomes. Support for a potential DA-mediated mechanism of action is provided by studies suggesting an important role of the AT1R in regulating central dopaminergic neurotransmission [19], high co-expression of AT1R and DA receptors [20] and evidence for functionally interactions between AT1R and DA receptors in the striatum [21].

Against this background, the present proof-of-concept study combined computational modelling and functional MRI, with a preregistered between-subjects randomized double-blind placebo-controlled pharmaco-fMRI design in n = 61 [Male] healthy participants to determine modulatory effects of LT-induced AT1R blockade on RL model parameters and the underlying neural mechanisms. We utilized a validated probabilistic selection RL paradigm with two stages: a learning phase in which participants learned to make better choices for fixed pairs of stimuli according to reward or loss feedback, and a subsequent transfer phase in which participants applied the learned optimal choices to novel combinations of stimuli without feedback. Behavioral responses during learning were fit using a computational RL

model to describe the dynamic learning process. Effects of LT on learning were examined by
means of comparing RL model parameters and neural activity related to model-derived
estimates of RPE. Effects of LT on learning transfer were examined by comparing choices and
functional connectivity in cortico-striatal pathways when participants approached the best
or avoided the worst stimulus. Based on previous literature [14, 16, 17], we predicted that LT
would: 1) reduce learning from negative outcomes, increase the RPE-associated signaling in
the ventral striatum (VS) and its neural expression for positive outcomes during the learning
phase and 2) increase selection of the best stimulus in the context of increased fronto-
striatal coupling during learning transfer.
133
METHODS
Participants
Seventy right-handed healthy male Chinese participants were screened according to
evaluated enrollment criteria (see **Supplemental Methods**, sample size based on previous
studies [15-17]). The study focused on male individuals to control for sex differences in
response to RAS blockade [22] and menstrual cycle-dependent variations in reward
processing [23]. Nine participants exhibited excessive head movement (LT, n=1) or poor
learning (LT, n=4; Placebo, n=4, **Fig. 1**) and thus failed to reach the choice accuracy criteria
(i.e., choosing 65% A in AB, 55% C in CD, 50% E in EF, in line with Frank et al., 2007), leading
to a final sample of n = 61 (mean±SD, age=20.89±2.32 years).
All participants provided written informed consent, protocols were pre-registered on
Clinical Trials.gov (https://clinicaltrials.gov/ct2/show/NCT04604938) and approved by the
ethics committee at the University of Electronic Science and Technology of China (Approval
355).
Using a double-blind randomized, placebo-controlled, between-subjects pharmacological
fMRI design, participants were administered either a single oral dose of LT (50mg) or
placebo (PLC) packed in identical capsules. Capsules were dispensed by an independent
researcher based on a computer-generated randomization sequence to implement double
blinding. Investigators involved in data acquisition and analyses were blinded for group
allocation. Prior studies suggest that single-dose LT effects on cardiovascular activity in
healthy subjects were observed after 3 hours [24]. Effects of the blood-brain barrier
permeable LT on brain activity and cognitive domains were observed in the timeframe
ranging from 1.5-2.5h after single-dose administration (which overlaps with peak plasma
levels after 90 minutes and an elimination half-life between 1.5-2.5h [25-27]). Consistent
with this, LT treatment was administered 90min before fMRI acquisition. Participants first
performed a reinforcement learning task (duration 30min) followed by an emotional
memory task (reported in Xu et al., 2021 [14]). To control for nonspecific effects of LT,
assessments of mood, attention and memory were incorporated at baseline and after the
experiment, while cardiovascular activity (i.e., blood pressure, heart rate) was measured at
baseline, after drug administration and after the experiment (**Fig. 1a** and **Supplemental
Methods**). To ensure double blinding participants were asked to guess treatment after the
experiment (treatment guess c^2=0.40, p=0.53; confirming successful double-blinding).
166
Experimental design

Probabilistic selection reinforcement learning paradigm
A validated probabilistic selection reinforcement learning paradigm was employed [28-30].
This paradigm consisted of two stages: an initial reinforcement learning phase and a
subsequent transfer phase. During the learning phase, participants were presented with one
of three different pairs of six shape stimuli (denoted as AB, CD and EF, **Fig. 1c**) on each trial in
a randomized order. Participants were instructed to learn to choose the better option of
each stimulus pair based on feedback (**Fig .1b**). Learning difficulty varied for the stimulus
pairs in terms of reward contingency (80:20, 70:30 or 60:40, for AB, CD or EF, respectively). A
total of 240 trials – dispersed across two fMRI runs with 120 trials each (40 trials per
stimulus pair, trial mean duration 4s) – were presented during the learning phase. Each trial
began with a fixation cross presented for a jittered interval of 0, 500, 1000, or 1500ms (**Fig.
1b**) followed by the presentation of two shapes displayed to the left and right of the fixation
cross (side was counterbalanced). Stimuli were presented until participants made a response
or 1700ms elapsed. The choice was visually confirmed by highlighting the chosen shape in
yellow for 300ms, followed by 400ms feedback presentation ('correct' or 'wrong'). Then, the
fixation cross was displayed again until the whole trial duration was reached. In addition, 12
null trials without stimulus presentation of the same duration were randomly interspersed in
each fMRI run to improve the model fitting of the rapid event-related fMRI design.
For the transfer phase, the six shape stimuli were recombined to constitute fifteen
stimulus pairs. Each stimulus pair permutation was presented 8 times (side was
counterbalanced) leading to 120 trials in the transfer phase, also with 12 null trials
interspersed in a random order. Duration of each trial was 1700ms and no feedback was
provided (**Fig. 1b**). Participants were told to choose the better option in each stimulus pair
according to what they had learned in the learning phase.
192

193
Fig. 1 Experimental timeline and paradigm. **(a)** Experimental timeline. **(b)**. The
reinforcement learning paradigm consisted of two subsequent phases. During the learning
phase participants were presented with one of three different pairs of six stimuli (denoted as
AB, CD and EF) on each trial in a randomized order. Participants were instructed to learn to
choose the best option within each stimulus pair based on the feedback presented (i.e.,
'correct' or 'wrong' presented as text, which indicated that 0.5 RMB or nothing were added

to the total payment). To avoid choice preference or reward associations with one particular
stimulus, stimulus pairs were presented in a counterbalanced order across subjects. During
the transfer phase, participants were presented with all permutations of combinations with
A and B - corresponding to the stimuli with the highest or lowest reward probability,
respectively - and were instructed to choose the better option according to their previous
learning experience. **(c)** The probabilities of acquiring reward for pairs AB, CD and EF were
80:20, 70:30 and 60:40, respectively, during the learning phase. The easiest condition was
therefore the AB pair while the EF pair was the hardest one to learn because of the relatively
equivalent reward probabilities between the two stimuli. A performance criterion (i.e.,
choosing 65% A in AB, 55% C in CD, 50% E in EF, a similar approach to that used by Frank et
al., 2007) was initially used to ensure successful learning in the subjects. N = 4 subjects in
each treatment group did not fulfill that criterion and were excluded from the further
analysis. The analysis in the transfer phase was conducted on trials in which A was correctly
chosen or B was avoided when being paired with another stimulus.
Abbreviations: BP, blood pressure; HR, heart rate; LT, losartan; PLC, placebo; FMRI, functional
magnetic resonance imaging; RL, reinforcement learning. License: Image in 1a were
designed by DinosoftLabs and obtained from Flaticon.com under the free use license.
217

Computational modeling of learning behavior
We explored the learning rate in terms of choice behavior by using a Q-learning algorithm
[31]. The Q-learning algorithm has been widely employed to model learning behavior and
serves to model the change in choice behavior based on trial-by-trial updates of the
expected value of choice options [30, 32, 33]. The corresponding model contains three free
parameters: learning rate for positive (αGain) and negative (αLoss) RPEs and estimation of
explore-exploit tendency (β). For details of modeling procedures, model estimation and
comparison see **Supplemental Methods and Fig. S1**.
226

Statistical analyses on the behavioral level
All analyses were performed in R (R development core and team, 2017). For the learning
phase, we employed a multilevel Bayesian linear model to analyze trial-by-trial choice
behavior using the Bayesian regression model in Stan (brms) R package [34]. Main effects of
treatment, stimulus pair and fMRI run, as well as the interaction of stimulus pair and
treatment on choice accuracy (proportion of choosing better stimulus in one stimulus pair,
e.g., choosing A in AB pair) were considered as credibly different when more than 95% of
the posterior distribution was above/below zero. Treatment effects on computational
modeling indices of choice behavior (learning rate, explore-exploit tendency) were examined
by using permutation-based two sample t tests via functions in the user contributed
packages permuco (version 1.1.0) for the R environment.
In the transfer phase, we performed similar analyses for trials including stimuli with the
highest and lowest reward probability (A, 80% or B, 20%) (**Fig. 1c**) to examine LT effects on
choosing the best and avoiding the worst option. An exploratory model additionally
examined effects of LT on choice times with choice (choose A, avoid choosing B) and
treatment (LT, PLC) as fixed factors and subject as random factor. Main effects of treatment,
choice behavior and their interaction were considered significant using the same 95%

posterior distribution criterion (details see **Supplemental Methods**). We additionally
explored the effects of treatment and rewarding probability (value) difference (e.g., AB-60,
AD-50, AF-40, CF-30, AE-20, AC-10) on choice accuracy and reaction time during learning
transfer (details see **Supplemental Methods**) to control for a potential confounding impact
of probability difference levels on treatment effects. We did not find significant main effects
of treatment or an interaction effect between treatment and rewarding probability
difference (**Fig. S2, Table S1-S2**), suggesting rather specific treatment effects on easily-
learned stimuli (i.e., A, B) during learning transfer.
252
MRI acquisition, preprocessing and first level analysis
MRI data were acquired on a 3.0-T GE Discovery MR system (General Electric Medical
System, Milwaukee, WI, USA) and preprocessed using standard procedures in SPM 12
(Statistical Parametric Mapping; http://www.fil.ion.ucl.ac.uk/spm/; Wellcome Trust Centre
for Neuroimaging) (see **Supplemental Methods**).Separate general linear models (GLM) were
designed for the learning and transfer phase.
For the learning phase, we established a GLM model that incorporated separate outcome
onsets for positive and negative feedback, each modulated by the corresponding RPE
estimated from the computational model. The highlight-period and six head motion
parameters were all included as covariates of no interest. Given that the choice accuracy in
both treatment groups rapidly approached a ceiling effect (such that irrespective of
treatment participants rapidly showed optimal choice accuracy, i.e., >80%, across stimulus
pairs with different rewarding probability in the second run, **Fig. S3**), the first and second
fMRI run were modeled separately, and analyses focused on the first run to increase the
sensitivity for learning-associated treatment effects.
During the transfer phase, approach A and avoid B were modeled as separate conditions,
and the six head motion parameters were included as nuisance regressors.
270
Examining neural effects of LT on RPE signaling during early learning
To examine the effects of LT on RPE signaling during early learning, the corresponding first
level contrasts (i.e., Positive+Negative RPE) were subjected to voxel-wise two sample t tests.
Whole brain analyses thresholded at cluster level family-wise error (FWE) corrected p<0.05
were employed (initial cluster threshold, p<0.001 uncorrected; see recommendations in
Slotnick, 2017 [35]).
277
Effects of LT on feedback-sensitive neural expressions in the VS during early learning
Given the higher sensitivity of multivariate neurofunctional representations for a given
mental process including reward and RPEs in the VS [36, 37], multi-voxel pattern analysis
(MVPA) was employed (see **Supplemental Methods**). We initially developed a decoder on
the whole brain neural pattern that differentiated positive versus negative outcomes during
early learning and tested it in an independent sample to validate brain systems strongly
involved in differentiating reward versus loss. Next, treatment effects on the corresponding
expression in the VS were examined (details see **Supplemental Methods**). The VS region of
interest included the ventral caudate and nucleus accumbens, defined from the
brainnetome atlas [38], which was functionally validated in our previous work [39, 40].

Examining neural effects of LT on optimal choice behavior during learning transfer

Effects of LT on the transfer of optimal choice behavior were examined by means of separate voxel-wise two sample t-tests for choosing A (choosing the best option) or avoiding B (avoiding the worst option), respectively. We employed the whole brain analyses thresholded at cluster level family-wise error (FWE) corrected $p<0.05$ (initial cluster threshold, $p<0.001$ uncorrected).

Functional connectivity analysis during learning transfer

Given that animal and human studies indicate that reinforcement learning is critically mediated by the functional communication between the VS and frontal regions [41, 42], we explored whether LT treatment could affect frontal-VS functional connectivity when subjects approach maximum (approach A) or avoid worst rewards (avoid B) during transfer phase. Treatment effects on frontal-VS functional networks were determined by performing two sample t tests on choosing A or avoid choosing B events. Within the brainnetome atlas-defined prefrontal cortex, results were thresholded at $p<0.05$ FWE corrected at peak level with small volume correction (SVC). In addition, we also explored the role of the VS in perceiving rewarding probability (value) difference by exploratory two sample t test with treatment as independent variable on VS activation for each value difference level.

RESULTS

Demographics and potential confounders

The LT (n=30) and PLC (n=31) groups were comparable with respect to sociodemographics and mood and cardiovascular indices arguing against nonspecific treatment effects (**Table 1**; all ps>0.10).

Table 1. Demographics and potential confounders in two groups

	Time	LT(M±SD)	PLC(M±SD)	t	p
Age		20.63±2.17	21.13±2.47	0.83	0.41
Body Mass Index, kg/m²		21.64±2.21	21.15±2.12	-0.87	0.39
Systolic blood pressure	Before	119.87±7.28	118.29±6.64	-0.88	0.38
	Drug peak	115.07±9.36	116.16±7.11	0.52	0.61
	After experiment	119.20±6.73	119.45±5.89	0.16	0.88
Diastolic blood pressure	Before	70.43±6.83	70.77±7.17	0.19	0.85
	Drug peak	69.77±6.92	69.23±7.61	-0.29	0.77
	After experiment	69.63±6.05	72.26±6.54	1.63	0.11
Heart rate	Before	75.00±11.09	76.10±10.51	0.40	0.69
	Drug peak	68.03±9.69	70.10±8.62	0.88	0.38
	After experiment	68.27±9.54	69.52±10.28	0.49	0.63
PANAS-P	Before	27.37±4.10	27.77±6.88	0.28	0.78

		LT	PLC		
	After experiment	24.5±4.98	26.84±7.23	-1.30	0.20
PANAS-N	Before	17.20±6.85	15.19±5.10	1.47	0.15
	After experiment	14.33±5.93	13.13±5.04	-0.86	0.40
SAI	Before	41.77±9.62	38.58±9.80	-1.28	0.21
	After experiment	39.9±8.56	37.39±8.12	-1.18	0.24
TAI	Before	41.73±9.47	40.13±8.02	-0.72	0.48
	After experiment	41.33±8.46	39.26±8.40	-0.96	0.34
BDI II	Before	9.13±7.16	8.61±8.82	-0.25	0.80
	After experiment	6.83±6.06	6.39±7.21	-0.26	0.80
D-CAT1	Before	39.83±8.81	38.06±5.77	-0.93	0.36
	After experiment	45.47±6.67	47.16±6.20	1.03	0.31
D-CAT2	Before	55.1±8.19	56.16±8.96	0.48	0.63
	After experiment	58.03±8.29	61.45±9.65	1.48	0.14
D-CAT3	Before	63.93±12.03	65.35±12.48	0.45	0.65
	After experiment	68.10±9.30	70.71±12.34	0.93	0.36
WM	Before	0.91±0.11	0.92±0.04	0.55	0.58
	After experiment	0.94±0.04	0.96±0.03	1.31	0.20

Values are presented as mean ± standard deviation.
Abbreviations: PANAS, Positive and Negative Affect Schedule; STAI, Spielberger State-Trait Anxiety Inventory; BDI, Beck Depression Inventory II; D-CAT 1-3, Digit cancellation test 1-3; WM, working memory; LT, Losartan, PLC, placebo.

LT increases choice accuracy for the hardest stimulus pair via increasing value sensitivity during early learning

The choice accuracy indicates the proportion of trials on which subjects chose the option with higher probability for reward in a stimulus pair (e.g., choose A in AB). Here we observed a significant main effect of stimulus pair (β=0.19, 95% highest-density interval (HDI), [0.16, 0.22], **Fig. S4**) such that participants exhibited the highest choice accuracy for the easy stimulus pair. The main effect of treatment did not reach significance (β=0.02, 95% HDI, [-0.02, 0.06], **Fig. S4**), but the main effect of fMRI run (β=0.12, 95% HDI, [0.11, 0.13], **Fig. S4**) was significant. Further inspection revealed that participants in both treatment groups showed an increased learning trend with high (all>85.3%) choice accuracy across stimulus pairs with different rewarding probability (**Fig. S3**). This may reflect that factors not related to probabilistic learning per se such as an understanding of the task structure or the reward probabilities may have led to an improved learning performance in the second run irrespective of treatment. The ceiling effect will lead to biased estimates and could critically reduce the sensitivity of detecting treatment effects on the trial-wise feedback-dependent learning process under examination (see also [43] for treatment effects on between run learning performance). This pattern was confirmed by a robust treatment × stimulus pair

interaction effect in the first (β=-0.05, 95% HDI, [-0.07,-0.03], **Fig. 2a**) but not the second
fMRI run (β=0.01, 95% HDI, [-0.01, 0.03], **Fig. 2b**), with further analyses indicating that
compared to PLC – LT increased choice accuracy for the most difficult stimulus pair (EF,
β=0.09, 95% HDI, [0.01, 0.18], **Fig. 2c**), especially during the early learning period for that
pair (trial 1-10, $t_{(59)}$=3.55, p=0.01, trial 11-20, $t_{(59)}$=7.29, p<0.001, **Fig. 2d**), but not the easier
pairs (AB, β=-0.01, 95% HDI, [-0.07,0.05]; CD, β=0.03, 95% HDI, [-0.05, 0.10], **Fig. 2c**) during
the first run. Our exploratory mediation analyses further indicated that Q value sensitivity
only played a mediating role in the effects of LT on choice accuracy for the hardest stimulus
pair (details see **Supplemental Methods and Fig. S5**). To increase the sensitivity to
determine learning-related treatment effects, all subsequent behavioral and neural analysis
consequently focused on the early learning phase (run 1).
348

Fig. 2 Behavioral effects of Losartan on choice accuracy and choice time. (a-b) A significant
interaction between treatment and stimulus pair was observed in the first but not the
second fMRI run of the learning phase. **(c)** During the first fMRI run of the learning phase,
losartan-treated participants increased the choice accuracy for the hardest stimulus pair (EF)
relative to the placebo group. **(d)** Specifically, in the first fMRI run, losartan group learned
faster relative to placebo group during the initial 20 trials for the hardest stimulus pair. PLC,
placebo; LT, losartan; *p<0.05, **p<0.01, ***p<0.001.
357

LT reduces learning rate for negative outcomes during early learning
In line with our hypothesis, LT significantly reduced learning rate from negative outcomes
($t_{(59)}$=-2.40, p=0.02, d=-0.61, **Fig. 3b**) but did not affect learning from positive outcomes
($t_{(59)}$=-1.84,p=0.07,d=-0.47, **Fig. 3a**). Moreover, LT enhanced the explore-exploit parameter
($t_{(59)}$=3.83, p<0.01, d=0.98, **Fig. 3c**), reflecting increased exploitatory choice behavior in

terms of more consistent selection of options with higher expected reward values.
364

365
366 **Fig. 3 Losartan effects on computational model parameters. (a)** Losartan and placebo
367 groups showed equivalent learning rate for positive outcome. **(b)** Compared to the placebo
368 group, losartan-treated individuals exhibited a reduced learning rate for negative outcomes.
369 **(c)** Moreover, losartan enhanced exploitatory decisions relative to placebo group. The error
370 bars denote standard error of the mean. n.s, non-significant, p<0.05, *** p<0.01.
371

LT increases RPE signaling during early learning

372
We initially examined brain regions that scaled positive and negative RPEs independent of
treatment. A corresponding one sample t-test confirmed previous studies suggesting that
activity in striatal and frontal regions linearly increased with the strength of the RPEs (**Fig.
S6, Table S3**). Examining treatment effects by means of a two sample t-test revealed that LT
enhanced RPE associated neural responses in the left VS (peak Montreal Neurological
Institute (MNI): x,y,z=-8,0,8, $t_{(59)}$=4.24,k=243, $P_{FWE-cluster}$<0.05, **Fig. 4a**) and bilateral
orbitofrontal cortex (left OFC, peak MNI: x,y,z =-40,54,-16, $t_{(59)}$=4.35, k=655, $P_{FWE-cluster}$<0.05;
right OFC, peak MNI, x,y,z=44,40,-14, $t_{(59)}$=4.24,k=326, $P_{FWE-cluster}$<0.05, **Fig. 4a**). Examination
of extracted parameter estimates (spherical masks, radius: 6mm) revealed that these regions
signaled positive but not negative RPEs under PLC whereas the LT-induced increase further
enhanced positive RPE and instated negative RPE signaling in these regions (**Fig. 4b**). To
further explore whether treatment differentially affected the RPEs in the VS we employed an
independent mask from the Brainnetome atlas [38]. The use of an independent mask
alleviates a potential bias of post hoc statistics [44] and the VS was chosen due to its critical
involvement in RPEs. Analyzing extracted estimates from the VS further confirmed that
losartan increased activation in the VS for both positive and negative RPEs (details **Fig. S7**).
389

a LT > PLC (p<0.05, FWE corrected)

left OFC

left VS

3.20 — *t - value* — 4.51

right OFC

b

390
Fig. 4 Losartan modulation on RPE-related neural response. (a) Comparison of losartan and
placebo groups in RPE-related response suggested a losartan-triggered increased activation
of bilateral orbitofrontal cortex and left ventral striatum. **(b)** For illustration purpose,
parameter estimates extraction from spherical masks (radius: 6 mm) of identified left or
right orbitofrontal cortex as well as left ventral striatum showed that losartan enhanced
activation in these regions to RPE for both positive and negative outcomes. The error bars
denoted standard error of the mean. LT, losartan; PLC, placebo. FWE-family-wise error, OFC-
orbitofrontal cortex, VS-ventral striatum, PRPE/N-positive reward prediction error, NRPE-
negative reward prediction error.
400

LT sharpens differential neural representations for positive vs negative outcomes in the VS
during early learning
We initially established an accurate whole brain multivariate predictive pattern for
classifying positive and negative outcomes (accuracy, 89.34%, sensitivity and specificity,
88.52% and 90.16%, respectively **Fig. 5b**). Applying thresholding (bootstrapped 10,000
samples) and multiple comparisons correction (false discovery rate [FDR] corrected,
p<0.001) revealed that a network including the VS, ventromedial prefrontal cortex,
dorsomedial prefrontal cortex and middle frontal gyrus strongly contributed to the
prediction of positive or negative outcomes during early learning (**Fig. 5a**, for a validation in
an independent dataset see **Fig. S8**).
Based on our a priori regional hypothesis about the crucial role of VS in reward learning
we examined effects of LT on VS neural representations for positive outcomes. Our results
suggested that only following LT - but not PLC - the VS expression accurately differentiated
positive from negative outcomes (accuracy=78.33%, p<0.001, sensitivity=0.83,
specificity=0.73, AUC=0.88; PLC, accuracy=56.45%, p=0.37, sensitivity=0.55, specificity=0.58,
AUC=0.68), with a direct comparison between the treatment groups suggesting that LT
specifically enhanced the VS representation for positive outcomes
($t_{(59)}$=9.92,p<0.001,d=1.29, **Fig. 5c**). A group comparison on classification accuracy using
permutation-inference further confirmed a significant treatment effect (t=-15.53, p<0.001,
95% CI, [-0.013, -0.010], **Fig. 5d**).

Fig. 5 Multivariate neural predictive pattern results. (a) Neural predictive pattern consists of voxels in which activity reliably predicted positive versus negative outcomes during early phase of learning. This map shows weights that exceed a threshold (p<0.001, FDR corrected based on bootstrapped 10,000 samples) for display only. Hot color indicates positive weights and cold color indicates negative weights. **(b)** ROC plot. This neural predictive pattern yields a classification accuracy of 89.34% in a leave-one-subject-out cross validation procedure. **(c)** Losartan treatment increases the neural pattern of the ventral striatum for positive outcome. **(d)** In the losartan group a better classification performance relative to the placebo group was observed across all permutations. The histograms of classification accuracy for the ventral striatum neural expression for classifying positive and negative outcome from permutation tests are presented. The red line shows the true classification accuracy in the losartan and placebo group separately. The error bars denotes standard error of the mean and the black line shows the mean value. VS-ventral striatum; vmPFC-ventromedial prefrontal cortex; dmPFC-dorsomedial prefrontal cortex; MFG-middle frontal gyrus; AUC- area under curve; FDR-False discovery rate; LT-losartan; PLC-placebo; ***p<0.001.

LT accelerates choices and increased VS-dlPFC coupling for approaching maximum rewards during the transfer phase

In the transfer phase, both choosing A and avoiding B were significantly higher than chance level (50%) regardless of treatment (choose A: t=15.07, p<0.001, avoid B: t=15.16, p<0.001), which indicated most participants could successfully apply their previous learning experience in the transfer phase. The main effect of treatment on choice accuracy was not significant (β=-0.03, 95% HDI, [-0.13, 0.06]). However, analyzing choice reaction times revealed a significant main effect of choice behavior (β=-12.70, 95% HDI, [-23.18, -1.94]) and

an interaction effect of treatment with choice behaviors (β=-27.36, 95% HDI, [-42.41, -
12.41]). Relative to PLC, LT accelerated choice times for choosing A (β=-83.52, 95% HDI, [-
147.03, -18.08], **Fig. 6a**), reflecting facilitated approach of the previously learned best option
following LT. Furthermore, we explored how previous learning experience supported the
optimal choice behaviors by conducting correlation analyses for learning rate and choice
behaviors of transfer. We found that learning rate for negative outcomes was not
significantly correlated with choice accuracy or response time of choose A or avoid B choices
across treatment groups, yet in the LT group, subjects with high learning rate for positive
outcomes during initial learning showed faster response towards maximum reward and
avoiding worst stimulus in learning transfer (**Fig. 6b**).

On the neural activation level, we did not observe treatment effects of LT on learning
transfer related regional activation, as well as on VS activity for each value difference level
(all ps>0.05, **Fig. S9**). However, on the level of functional connectivity LT increased functional
connectivity between the VS and left dorsolateral prefrontal cortex (left dlPFC, peak MNI:
x,y,z=-48,22,28, $t_{(59)}$=5.15, k=197, $P_{svc-FWEpeak}$=0.01, **Fig. 6c**), reflecting an LT-induced
enhancement of VS-dlPFC communication when approaching maximum rewards during
learning transfer.

Fig. 5 Behavioral and neural results for transfer phase.
(a) In the transfer phase, all participants responded quickly when choosing stimulus A or
avoiding stimulus B, and relative to the placebo group, losartan group exhibited faster
responses for approaching stimulus A in a novel environment. **(b)** Moreover, for losartan-
treated participants, higher learning rate for positive outcomes in initial learning phase was
related to the accelerated responses toward maximum reward or avoiding worst option in
transfer phase, while this association was abolished in placebo group. **(c)** For illustration

purpose parameter estimates were extracted from a spherical (radius: 6 mm) ROI in the
identified left dorsolateral prefrontal cortex (dlPFC) region. Losartan increased functional
coupling between ventral striatum and left dlPFC when participants choose A stimuli in
transfer phase. The statistical map of the left dlPFC was thresholded at p<0.001 uncorrected
(whole-brain level) for display purpose. The error bars denoted standard error of the mean.
LT-losartan; PLC-placebo.
480
DISCUSSION
The present pharmacological study utilized computational modeling in combination with
fMRI to examine the effects of transient LT-induced AT1R blockade on reinforcement
learning and the underlying neural mechanism in healthy individuals. On the behavioral level
LT facilitated choice accuracy in the most difficult reward condition while it specifically
reduced learning from negative outcomes and enhanced exploitatory choice behaviors. On
the neural level, the behavioral effects were paralleled by regional-specific effects on ventral
striatal-orbitofrontal reward systems, such that LT increased RPE signaling in these regions
and sharpened the fine-grained neurofunctional distinction between positive and negative
outcomes in the VS. During learning transfer, LT facilitated approach of the maximum
rewarding option and enhanced VS-dlPFC functional connectivity. Overall, these findings
indicate that LT-attenuated learning from negative feedback in the context of general
positive outcome learning and a subsequent increased motivation to obtain maximum
rewards during learning transfer, which on the neural level was accompanied by enhanced
RPE and functional communication in fronto-striatal circuits.
We found that LT specifically enhanced choice accuracy for the most difficult condition
suggesting that LT specifically improved learning under a low reinforcement probability.
Computational modeling additionally allowed a more fine-grained examination of the
behavioral effects by fitting trial-by-trial learning behavior and revealed that LT specifically
reduced the learning rate for negative outcomes and enhanced exploitatory choices. The
optimal learning ability could be understood when learning rate and other free parameters
are considered simultaneously (e.g., explore-exploit tendency) in the RL model and the
reward schedule [45]. Effects of LT on learning rate were outcome dependent, such that LT
specifically decreased learning from negative outcomes, indicating an attenuated influence
of negative information on reinforcement learning. Within the context of a stable
reinforcement schedule, it is adaptive for an agent to ignore relatively rare and potentially
misleading negative feedback given that an oversensitivity to negative outcomes would
cause suboptimal choice behaviors. Therefore, decreased negative learning rate may signal
the relatively high approach for positive outcomes in a stable reward contingency, which in
turn may facilitate an exploitatory choice tendency in terms of consistent decisions for
options with a higher expected reward [33]. Previous studies demonstrated that enhancing
central dopaminergic activity increases choices towards monetary gains [1, 46]. The current
pattern of results may reflect modulatory effects of RAS blockade on dopaminergic
neurotransmission given that LT has been shown to induce stronger D1 receptor expression
[47] which has been associated with better reward-associative learning [48]. These findings
resonate with recent studies reporting an LT-induced enhancement of learning from positive
relative to negative events [12] as well as an LT-induced shift from preferential social

punishment towards social reward processing [17]. Together, this pattern of effects suggests
that LT can attenuate the impact of negative information thus promoting motivation to
select rewarding options.
On the neural level LT increased orbitofronto-striatal RPE-signaling and induced a more
distinct neural expression for positive outcomes in the VS. The VS dopamine neurons are
critically involved in RPE signaling and reward seeking [49], while the OFC is strongly
implicated in computation of expected reward values and RPEs [50]. An LT-induced
enhancement of the neural RPE signal and the representation of rewarding outcomes may
reflect the potential for the RAS to modulate central dopaminergic neurotransmission during
reinforcement learning. The AT1R is expressed densely in dopamine-rich brain areas [20],
particularly in the striatum [51] and plays a key role in dopaminergic function [52].
Administering an antagonist of AT1R could mediate D1 density in the striatum [47] and block
the functional response of the D2 receptor [21] - both of these receptors exhibit dense
expression in ventral striatal and prefrontal regions crucially involved in reward learning [53,
54]. This may indicate a potential downstream effect of LT-induced AT1R blockade on DA
signaling, in turn modulating reward learning within orbitofronto- striatal circuits, thus
enhancing RPE encoding and reward representation in these regions.
During subsequent learning transfer, LT facilitated approach of the maximum reward in
terms of accelerated decisions. This effect was not linked with the role of VS in value
difference processing but was observed in the context of enhanced functional coupling of
the VS with left dlPFC. Faster decisions for choosing the best options following LT may reflect
an increased motivation to focus on maximizing rewards after reinforcement learning. The
findings partly align with early research on dopaminergic modulation of reinforcement
learning, which reported improved motivation for the highest-rewarding option during
transfer, an effect that was explained as dopamine-dependent enhancement of learning
signals [55]. The important role of fronto-striatal connectivity in reinforcement learning has
been extensively documented [41], indicating that reward associations initially formed in the
striatum are subsequently used to guide learning and decisions engaging the dlPFC [56]. To
be specific, the dlPFC plays an important role in integrating and transmitting reward
representations to the mesolimbic and mesocortical dopamine systems to initiate reward-
motivated behaviors [57]. Reduced striato-dlPFC connectivity has been observed in in
disorders characterized by a dysfunctional DA system [58] and linked with impaired
reinforcement learning [59]. As such, the present findings of an LT-induced increase in VS-
dlPFC connectivity when approaching rewards might reflect a modulatory role of angiotensin
signaling on fronto-striatal communication via effects on dopaminergic circuits.
Given the repeatedly observed hypersensitivity for negative information and an increased
impact of negative information on learning in depression [60], the current pattern of
behavioral effects may reflect potential for LT to normalize biased processing in depression
and in turn improve motivational deficits. The therapeutic potential in depression is further
supported by early animal models suggesting a crucial role of the RAS in depression [61, 62]
and documenting potential antidepressant behavioral effects of LT [63, 64]. Initial studies
aimed at targeting reward processing and reinforcement learning impairments in depression
via directly targeting the dopaminergic system [65, 66]. These studies revealed initially
promising evidence for a therapeutic potential of DA agonist in depression including

normalized neural functioning in fronto-striatal reward systems [65, 66] and anhedonia improvement [5]. However, effects on impaired reward learning were not observed and the clinical utility of DA agonist is limited by adverse effects such as triggering impulsive behaviors [67] and abuse [68]. The current pattern of results may point to a possibility that LT may represent a safe and potentially behavioral relevant strategy to modulate deficient reward learning and fronto-striatal functioning in depression.

While the current study found some evidence for a novel pathway to modulate reward learning, future studies are required to: (1) determine the potential of LT to influence reward learning and associated fronto-striatal deficits in depression, and (2) uncover the detailed interaction mechanism between the RAS with DA systems during reward learning such as incorporating receptor maps in combination with molecular imaging. In addition, future studies are required to demonstrate whether the observed effects generalize to women.

Taken together, we demonstrated that AT1R blockade via LT decreased negative learning rate but did not affect learning from positive outcomes, while increasing RPE signaling in orbitofronto-striatal regions and improving neural expression of positive outcomes in the VS. During the subsequent transfer, LT accelerated choices for maximizing rewards and increased VS-dlPFC functional coupling. Together, this pattern may reflect a promising mechanism of LT as a potential treatment to normalize impaired reward learning and fronto-striatal functioning in depression.

ACKNOWLEDGMENTS

We truly thank all colleagues for the impressive discussions that help us to improve the manuscript and the help during the FMRI data acquisition, and all volunteers that participated in our study. This work was supported by the National Natural Science Foundation of China (Grant No. 32250610208), China MOST2030 Brain Project (Grant No. 2022ZD0208500) and the National Key Research and Development Program of China (Grant No. 2018YFA0701400).

AUTHOR CONTRIBUTIONS

TX and BB designed the study. TX, LW, GJ, RZ conducted the experiment and collected the data. TX, LJ, XZ, FZ performed the data analysis. TX and BB wrote the manuscript draft. JK, CZ, FZ, WZ, YS and BB critically revised the manuscript draft.

COMPETING INTERESTS

The authors report no biomedical financial interests or potential conflicts of interest.

DATA AND MATERIALS AVAILABILITY

Unthresholded group-level statistical maps are available on NeuroVault (https://neurovault.org/collections/12001/). Additional data related to study is available from the corresponding author upon reasonable request. A preprint of the manuscript was archived on the biorxiv.org repository (doi: https://doi.org/10.1101/2022.03.14.484364).The present study was pre-registered on Clinical Trials.gov (Trial name: The effects of losartan on reward reinforcement learning; Registration number: NCT04604938; URL: https://clinicaltrials.gov/ct2/show/NCT04604938).

REFERENCES

1. Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD. Dopamine-dependent prediction errors underpin reward-seeking behaviour in humans. *Nature* 2006; **442**(7106)**:** 1042-1045.

2. Schultz W. Getting formal with dopamine and reward. *Neuron* 2002; **36**(2)**:** 241-263.

3. Husain M, Roiser JP. Neuroscience of apathy and anhedonia: a transdiagnostic approach. *Nature Reviews Neuroscience* 2018; **19**(8)**:** 470-484.

4. Chen C, Takahashi T, Nakagawa S, Inoue T, Kusumi I. Reinforcement learning in depression: a review of computational research. *Neuroscience & Biobehavioral Reviews* 2015; **55:** 247-267.

5. Whitton AE, Reinen JM, Slifstein M, Ang Y-S, McGrath PJ, Iosifescu DV *et al.* Baseline reward processing and ventrostriatal dopamine function are associated with pramipexole response in depression. *Brain* 2020; **143**(2)**:** 701-710.

6. Kumar P, Goer F, Murray L, Dillon DG, Beltzer ML, Cohen AL *et al.* Impaired reward prediction error encoding and striatal-midbrain connectivity in depression. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 2018; **43**(7)**:** 1581-1588.

7. Reinen JM, Whitton AE, Pizzagalli DA, Slifstein M, Abi-Dargham A, McGrath PJ *et al.* Differential reinforcement learning responses to positive and negative information in unmedicated individuals with depression. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 2021; **53:** 89-100.

8. Brown VM, Zhu L, Solway A, Wang JM, McCurry KL, King-Casas B *et al.* Reinforcement Learning Disruptions in Individuals With Depression and Sensitivity to Symptom Change Following Cognitive Behavioral Therapy. *JAMA psychiatry* 2021.

9. Queirazza F, Fouragnan E, Steele JD, Cavanagh J, Philiastides MG. Neural correlates of weighted reward prediction error during reinforcement learning classify response to cognitive behavioral therapy in depression. *Science advances* 2019; **5**(7)**:** eaav4962.

10. Goldberg AI, Dunlay MC, Sweet CS. Safety and tolerability of losartan potassium, an angiotensin II receptor antagonist, compared with hydrochlorothiazide, atenolol, felodipne ER, and angiotensin-converting enzyme inhibitors for the treatment of systemic hypertension. *The American journal of cardiology* 1995; **75**(12)**:** 793-795.

11. Khoury NM, Marvar PJ, Gillespie CF, Wingo A, Schwartz A, Bradley B *et al.* The renin-angiotensin pathway in posttraumatic stress disorder: angiotensin-converting enzyme inhibitors and angiotensin receptor blockers are associated with fewer traumatic stress symptoms. *The Journal of clinical psychiatry* 2012; **73**(6)**:** 17451.

12. Pulcu E, Shkreli L, Holst CG, Woud ML, Craske MG, Browning M *et al.* The effects of the angiotensin II receptor antagonist losartan on appetitive versus aversive learning: a randomized controlled trial. *Biological psychiatry* 2019; **86**(5)**:** 397-404.

13. Ranjbar H, Aghaei I, Moosazadeh M, Shabani M. Angiotensin II type 1 receptor blocker losartan attenuates locomotor, anxiety-like behavior, and passive avoidance learning deficits in a sub-chronic stress model. *Iranian journal of basic medical sciences* 2018; **21**(8)**:** 856.

14. Xu T, Zhou X, Jiao G, Zeng Y, Zhao W, Li J *et al.* Angiotensin antagonist inhibits preferential negative memory encoding via decreasing hippocampus activation and its coupling with amygdala. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* 2022.

15. Stout DM, Risbrough VB. Angiotensin II signaling and fear extinction: translational evidence and novel receptor targets. *Biological psychiatry* 2019; **86**(12)**:** 874-876.

651   16.   Zhou F, Geng Y, Xin F, Li J, Feng P, Liu C *et al.* Human extinction learning is accelerated by an
652   angiotensin antagonist via ventromedial prefrontal cortex and its connections with basolateral
653   amygdala. *Biological psychiatry* 2019; **86**(12)**:** 910-920.
654   17.   Zhou X, Xu T, Zeng Y, Zhang R, Qi Z, Zhao W *et al.* The angiotensin antagonist Losartan modulates social
655   reward motivation and punishment sensitivity via modulating midbrain-striato-frontal circuits. *Journal*
656   *of Neuroscience* 2023; **43**(3)**:** 472-483.
657   18.   Glimcher PW. Understanding dopamine and reinforcement learning: the dopamine reward prediction
658   error hypothesis. *Proceedings of the National Academy of Sciences* 2011; **108**(supplement_3)**:** 15647-
659   15654.
660   19.   Kobiec T, Otero-Losada M, Chevalier G, Udovin L, Bordet S, Menéndez-Maissonave C *et al.* The Renin–
661   Angiotensin System Modulates Dopaminergic Neurotransmission: A New Player on the Scene.
662   *Frontiers in Synaptic Neuroscience* 2021; **13:** 16.
663   20.   Saavedra J. Beneficial effects of Angiotensin II receptor blockers in brain disorders. *Pharmacological*
664   *research* 2017; **125:** 91-103.
665   21.   Martinez-Pinilla E, Rodriguez-Perez A, Navarro G, Aguinaga D, Moreno E, Lanciego J *et al.* Dopamine
666   D2 and angiotensin II type 1 receptors form functional heteromers in rat striatum. *Biochemical*
667   *pharmacology* 2015; **96**(2)**:** 131-142.
668   22.   Miller JA, Cherney DZ, Duncan JA, Lai V, Burns KD, Kennedy CR *et al.* Gender differences in the renal
669   response to renin-angiotensin system blockade. *Journal of the American Society of Nephrology* 2006;
670   **17**(9)**:** 2554-2560.
671   23.   Dreher J-C, Schmidt PJ, Kohn P, Furman D, Rubinow D, Berman KF. Menstrual cycle phase modulates
672   reward-related neural function in women. *Proceedings of the National Academy of Sciences* 2007;
673   **104**(7)**:** 2465-2470.
674   24.   Mechaeil R, Gard P, Jackson A, Rusted J. Cognitive enhancement following acute losartan in
675   normotensive young adults. *Psychopharmacology* 2011; **217**(1)**:** 51-60.
676   25.   Getyala A, V Gangadharappa H, Sarat Chandra Prasad M, Praveen Kumar Reddy M, M Pramod Kumar
677   T. Formulation and evaluation of non-effervescent floating tablets of losartan potassium. *Current drug*
678   *delivery* 2013; **10**(5)**:** 620-629.
679   26.   Samyuktha M, Vasanth P, Suresh K, Ramesh T, Ramesh M. Formulation and evaluation of
680   gastroretentive floating tablets of losartan potassium. *Int J Biopharm* 2013; **4:** 18-26.
681   27.   Sica DA, Gehr TW, Ghosh S. Clinical pharmacokinetics of losartan. *Clinical pharmacokinetics* 2005;
682   **44**(8)**:** 797-814.
683   28.   Frank MJ, Moustafa AA, Haughey HM, Curran T, Hutchison KE. Genetic triple dissociation reveals
684   multiple roles for dopamine in reinforcement learning. *Proceedings of the National Academy of*
685   *Sciences* 2007; **104**(41)**:** 16311-16316.
686   29.   Frank MJ, Woroch BS, Curran T. Error-related negativity predicts reinforcement learning and conflict
687   biases. *Neuron* 2005; **47**(4)**:** 495-501.
688   30.   Jahfari S, Ridderinkhof KR, Collins AGE, Knapen T, Waldorp LJ, Frank MJ. Cross-Task Contributions of
689   Frontobasal Ganglia Circuitry in Response Inhibition and Conflict-Induced Slowing. *Cerebral cortex*
690   2019; **29**(5)**:** 1969-1983.
691   31.   Sutton RS, Barto AG. *Reinforcement learning: An introduction*. MIT press2018.
692   32.   Grogan JP, Tsivos D, Smith L, Knight BE, Bogacz R, Whone A *et al.* Effects of dopamine on reinforcement
693   learning and consolidation in Parkinson's disease. *Elife* 2017; **6:** e26801.
33.   Van Slooten JC, Jahfari S, Knapen T, Theeuwes J. How pupil responses track value-based decision-

making during and after reinforcement learning. *PLoS computational biology* 2018; **14**(11)**:** e1006632.

34.   Bürkner P-C. brms: An R package for Bayesian multilevel models using Stan. *Journal of statistical*
software 2017; **80**(1)**:** 1-28.

35.   Slotnick SD. Cluster success: fMRI inferences for spatial extent have acceptable false-positive rates.
Cognitive neuroscience 2017; **8**(3)**:** 150-155.

36.   Kahnt T. A decade of decoding reward-related fMRI signals and where we go from here. *Neuroimage*
2018; **180:** 324-333.

37.   Zhou F, Zhao W, Qi Z, Geng Y, Yao S, Kendrick KM *et al.* A distributed fMRI-based signature for the
subjective experience of fear. *Nature communications* 2021; **12**(1)**:** 1-16.

38.   Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L *et al.* The human brainnetome atlas: a new brain atlas
based on connectional architecture. *Cerebral cortex* 2016; **26**(8)**:** 3508-3526.

39.   Zhao Z, Ma X, Geng Y, Zhao W, Zhou F, Wang J *et al.* Oxytocin differentially modulates specific dorsal
and ventral striatal functional connections with frontal and cerebellar regions. *Neuroimage* 2019; **184:**
781-789.

40.   Zhou X, Zimmermann K, Xin F, Zhao W, Derckx RT, Sassmannshausen A *et al.* Cue reactivity in the
ventral striatum characterizes heavy cannabis use, whereas reactivity in the dorsal striatum mediates
dependent use. *Biological psychiatry: cognitive neuroscience and neuroimaging* 2019; **4**(8)**:** 751-762.

41.   Averbeck   B,   O'Doherty   JP.   Reinforcement-learning   in   fronto-striatal   circuits.
Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology
2021.

42.   Lowet AS, Zheng Q, Matias S, Drugowitsch J, Uchida N. Distributional Reinforcement Learning in the
Brain. *Trends in neurosciences* 2020; **43**(12)**:** 980-997.

43.   Becker B, Klein EM, Striepens N, Mihov Y, Schlaepfer TE, Reul J *et al.* Nicotinic acetylcholine receptors
contribute to learning-induced metaplasticity in the hippocampus. *Journal of cognitive neuroscience*
2013; **25**(7)**:** 986-997.

44.   Kriegeskorte N, Simmons WK, Bellgowan PS, Baker CI. Circular analysis in systems neuroscience: the
dangers of double dipping. *Nature neuroscience* 2009; **12**(5)**:** 535-540.

45.   Zhang L, Lengersdorff L, Mikus N, Gläscher J, Lamm C. Using reinforcement learning models in social
neuroscience: frameworks, pitfalls and suggestions of best practices. *Social Cognitive and Affective*
Neuroscience 2020; **15**(6)**:** 695-707.

46.   Chowdhury R, Guitart-Masip M, Lambert C, Dayan P, Huys Q, Düzel E *et al.* Dopamine restores reward
prediction errors in old age. *Nature neuroscience* 2013; **16**(5)**:** 648-653.

47.   Schlenker EH. In dystrophic hamsters losartan affects control of ventilation and dopamine D1 receptor
density. *Respiratory physiology & neurobiology* 2010; **173**(1)**:** 71-78.

48.   Beninger RJ, Miller R. Dopamine D1-like receptors and reward-related incentive learning. *Neuroscience*
& biobehavioral reviews 1998; **22**(2)**:** 335-345.

49.   Cox J, Witten IB. Striatal circuits for reward learning and decision-making. *Nature Reviews*
Neuroscience 2019; **20**(8)**:** 482-494.

50.   Rolls ET. The roles of the orbitofrontal cortex via the habenula in non-reward and depression, and in
the responses of serotonin and dopamine neurons. *Neuroscience & Biobehavioral Reviews* 2017; **75:**
331-334.

51.   Garrido-Gil P, Rodriguez-Perez AI, Fernandez-Rodriguez P, Lanciego JL, Labandeira-Garcia JL. Expression
of angiotensinogen and receptors for angiotensin and prorenin in the rat and monkey striatal neurons
and glial cells. *Brain structure & function* 2017; **222**(6)**:** 2559-2571.

52. Rodriguez-Perez AI, Valenzuela R, Villar-Cheda B, Guerra MJ, Labandeira-Garcia JL. Dopaminergic
neuroprotection of hormonal replacement therapy in young and aged menopausal rats: role of the
brain angiotensin system. *Brain* 2012; **135**(1)**:** 124-138.

53. Higa KK, Young JW, Ji B, Nichols DE, Geyer MA, Zhou X. Striatal dopamine D1 receptor suppression
impairs reward-associative learning. *Behavioural brain research* 2017; **323:** 100-110.

54. Onge JRS, Abhari H, Floresco SB. Dissociable contributions by prefrontal D1 and D2 receptors to risk-
based decision making. *Journal of Neuroscience* 2011; **31**(23)**:** 8625-8633.

55. Jocham G, Klein TA, Ullsperger M. Dopamine-mediated reinforcement learning signals in the striatum
and ventromedial prefrontal cortex underlie value-based choices. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience 2011; **31**(5)**:** 1606-1613.

56. Pasupathy A, Miller EK. Different time courses of learning-related activity in the prefrontal cortex and
striatum. *Nature* 2005; **433**(7028)**:** 873-876.

57. Ballard IC, Murty VP, Carter RM, MacInnes JJ, Huettel SA, Adcock RA. Dorsolateral prefrontal cortex
drives mesolimbic dopaminergic regions to initiate motivated behavior. *Journal of Neuroscience* 2011;
31(28)**:** 10340-10346.

58. Becker A, Kirsch M, Gerchen MF, Kiefer F, Kirsch P. Striatal activation and frontostriatal connectivity
during non-drug reward anticipation in alcohol dependence. *Addiction biology* 2017; **22**(3)**:** 833-843.

59. Park SQ, Kahnt T, Beck A, Cohen MX, Dolan RJ, Wrase J *et al.* Prefrontal cortex fails to learn from
reward prediction errors in alcohol dependence. *Journal of Neuroscience* 2010; **30**(22)**:** 7749-7753.

60. Noworyta K, Cieslik A, Rygula R. Neuromolecular Underpinnings of Negative Cognitive Bias in
Depression. *Cells* 2021; **10**(11)**:** 3157.

61. Gironacci MM, Vicario A, Cerezo G, Silva MG. The depressor axis of the renin–angiotensin system and
brain disorders: A translational approach. *Clinical science* 2018; **132**(10)**:** 1021-1038.

62. Vian J, Pereira C, Chavarria V, Kohler C, Stubbs B, Quevedo J *et al.* The renin-angiotensin system: a
possible new target for depression. *BMC medicine* 2017; **15**(1)**:** 144.

63. Gard PR. The role of angiotensin II in cognition and behaviour. *European journal of pharmacology*
2002; **438**(1-2)**:** 1-14.

64. Gard PR. Angiotensin as a target for the treatment of Alzheimer's disease, anxiety and depression.
Expert opinion on therapeutic targets 2004; **8**(1)**:** 7-14.

65. Admon R, Kaiser RH, Dillon DG, Beltzer M, Goer F, Olson DP *et al.* Dopaminergic enhancement of
striatal response to reward in major depression. *American Journal of Psychiatry* 2017; **174**(4)**:** 378-
386.

66. Schneier FR, Slifstein M, Whitton AE, Pizzagalli DA, Reinen J, McGrath PJ *et al.* Dopamine release in
antidepressant-naive major depressive disorder: A multimodal [11C]-(+)-PHNO positron emission
tomography and functional magnetic resonance imaging study. *Biological psychiatry* 2018; **84**(8)**:** 563-
573.

67. Voon V, Reynolds B, Brezing C, Gallea C, Skaljic M, Ekanayake V *et al.* Impulsive choice and response in
dopamine agonist-related impulse control behaviors. *Psychopharmacology* 2010; **207**(4)**:** 645-659.

68. Gendreau KE, Potenza MN. Publicity and reports of behavioral addictions associated with dopamine
agonists. *Journal of Behavioral Addictions* 2016; **5**(1)**:** 140-143.

779
780
781