A pathogenicity locus of *Streptococcus gallolyticus* subspecies *gallolyticus*

John Culver Taylor[1], Ritesh Kumar[1#], Juan Xu[1], and Yi Xu[1,2,3]*

[1]Center for Infectious and Inflammatory Diseases, Texas A&M Health Science Center Institute

of Biosciences of Technology, Houston, Texas, United States of America

[2]Department of Microbiology and Molecular Genetics, McGovern Medical School, UT Health,

Houston, Texas, United States of America

[3]Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas A&M

Health Science Center, Texas, United States of America

[#]Current affiliation: IFF Health and Biosciences

*Corresponding author: Yi Xu

E-mail: yi-xu1@tamu.edu

Key words: Colorectal cancer, colon cancer, microbiome, *Streptococcus gallolyticus*,

Streptococcus bovis, cell proliferation, colonization.

Abstract

Streptococcus gallolyticus subspecies *gallolyticus* (*Sgg*) is known to be strongly

associated with colorectal cancer (CRC). Recent functional studies further demonstrated that *Sgg*

actively stimulates CRC cell proliferation and promotes the development of colon tumors.

However, the *Sgg* factors important for the pro-proliferative and pro-tumor activities of *Sgg*

remain unclear. Here, we identified a chromosomal locus in *Sgg* strain TX20005. Deletion of this

locus significantly reduced *Sgg* adherence to CRC cells and abrogated the ability of *Sgg* to

stimulate CRC cell proliferation. Thus, we designate this locus as the <u>S</u>gg <u>p</u>athogenicity-

<u>a</u>ssociated <u>r</u>egion (*SPAR*). More importantly, we found that *SPAR* is important for *Sgg*

pathogenicity *in vivo*. In a gut colonization model, mice exposed to the *SPAR* deletion mutant

showed significantly reduced *Sgg* load in the colonic tissues and fecal materials, suggesting that

SPAR contributes to the colonization capacity of *Sgg*. In a mouse model of CRC, deletion of

SPAR abolished the ability of *Sgg* to promote the development of colon tumors growth. Taken

together, these results highlight *SPAR* as a critical pathogenicity determinant of *Sgg*.

Introduction

Streptococcus gallolyticus subspecies *gallolyticus* (*Sgg*) is a group D streptococcal

bacterium, and an opportunistic pathogen belonging to the *Streptococcus bovis/Streptococcus*

equinis complex (SBSEC) (1). *Sgg* causes bacteremia and infective endocarditis (IE), and has a

long-standing clinical association with CRC, supported by extensive studies and case reports

dating back to the 1950s (2-15). Patients with *Sgg* IE and/or bacteremia have concomitant colon

adenomas or adenocarcinomas at a much higher rate (~60%) than that of the general population

(8). Moreover, a prospective study found that 45% of patients with *Sgg* IE developed colonic

neoplastic lesions within 5 years of IE diagnosis compared to 21% of patients with IE due to

closely-related enterococci, further supporting the association of *Sgg* with CRC (6). In CRC

patients with no symptoms of IE or bacteremia, *Sgg* was found to preferentially associate with

tumor tissues compared to normal adjacent tissues or normal tissues in multiple studies around

the world (4, 16-19).

Functionally, *Sgg* stimulates the proliferation of human CRC cells in a β-catenin

dependent manner *in vitro* and in a xenograft model (4). Exposure to *Sgg* also led to larger and

more advanced tumors in an azoxymethane (AOM)-induced CRC model and in a colitis-

associated CRC model (4, 15). These results indicate that *Sgg* possesses pro-proliferation and

pro-tumor activities. Previous work also indicated that there are variations among *Sgg* strains in

their ability to stimulate human CRC cell proliferation. Some *Sgg* strains such as ATCC43143

showed significantly reduced capacity to adhere to HT29 cells, to stimulate the proliferation of

these cells and to promote tumor development *in vivo* (2), suggesting that there are specific *Sgg*

factors contributing to the pro-proliferative and pro-tumor activities of *Sgg*. The identity of these

specific *Sgg* factors was unknown.

Here, we describe work characterizing a chromosomal locus of *Sgg* strain TX20005.

Deletion of this locus significantly impaired the ability of *Sgg* to adhere to cultured CRC cells

and abrogated the ability of *Sgg* to stimulate CRC cell proliferation. *In vivo*, the deletion mutant

displayed significantly reduced capacity to colonize normal mouse colons. More importantly, the

mutant lost the ability to promote the development of colon tumors. Given the role of this locus

in *Sgg* pathogenesis, we have coined this locus as the <u>S</u>gg <u>p</u>athogenicity-<u>a</u>ssociated <u>r</u>egion

(*SPAR*).

61   **Materials and Methods**

Bacterial strains, cell lines and growth conditions. *Sgg* strains were routinely grown in

brain heart infusion (BHI) broth or tryptic soy broth at 37°C with shaking, or on BHI or TSB

agar plates at 37°C overnight (Teknova). For co-culture experiments and animal studies,

stationary phase bacterial cultures were pelleted, washed with phosphate-buffered saline (PBS),

pH 7.4, resuspended in PBS containing 15% glycerol, aliquoted and stored at -80°C. Aliquoted

stocks were thawed, washed with PBS, diluted in appropriate media to obtain the desired

concentration and directly added to cells or administered to mice. Human colon cancer cell lines

HT29, HCT116, and SW480, as well as the HEK293 cell line were grown in Dulbecco's

Modified Eagle's Medium F-12 50/50 (DMEM/F-12, GIBCO) supplemented with 10% FBS at

37°C with 5% CO_2 in a humidified chamber. Cells from less than 30 passages were used in the

experiments.

Preparation of culture supernatants (CSs). Supernatants from stationary phase *Sgg*

cultured in BHI were filtered-sterilized through a 0.2μm filter and concentrated approximately

10 – 20-fold using centrifugal concentrators (3kD molecular weight cut off). The concentrated

CSs were aliquoted and stored at -80°C. A vial is thawed and diluted in the appropriate media for

adherence or cell proliferation assays and used immediately.

Adherence assay. This was performed as described previously (3). Briefly, HT29 cells

were seeded at a density of $1x10^6$ cells/well in a 24 well plate and allowed to attach overnight.

Cells were incubated with bacteria at a multiplicity of infection (MOI) of 10 in the absence or

presence of CSs (diluted to 1x) for 1 hour at 37°C with 5% CO_2. Cells were washed thrice in

PBS to remove unattached bacteria, lysed with 1mL of 0.01% Triton X-100 (Sigma) and dilution

plated onto BHI or TSB agar plates. Adherence was calculated as the percentage of adhered

bacteria vs. total bacteria added.

Cell proliferation assay. This was performed as described previously (3) with slight

modifications. Briefly, cells were seeded in 96 well plates at a concentration of $\sim 1x10^4$ cells/well

and incubated overnight. Cells were then incubated in fresh DMEM/F-12 containing 10% FBS in

the absence or presence of *Sgg* bacteria (MOI = 1) or *Sgg* CSs (1x) for a total of 24 hours.

Trimethoprim was added after 6 hours of incubation (1 μg/mL final concentration) to inhibit

bacterial growth, as previously described. The number of viable cells was determined using the

cell counting kit (CCK)-8 kit following the instructions of the supplier (Apex Bio).

Western Blot. Detection of β-catenin and PCNA was carried out as described previously

(3). Briefly, HT29 cells were seeded at a density of $1x10^6$ cells per well in 6-well plates and

incubated overnight. The cells were then incubated with media only or media containing bacteria

(MOI=1) for 9 hours. Total cell lysates were subjected to SDS-PAGE, transferred, and probed

with antibodies against β-catenin (1:1000), PCNA (1:1000), and β-actin (1:1000) (Cell Signaling

Technology), followed by incubation with HRP-conjugated secondary antibodies (1:3000). Band

intensities were measured using FIJI ImageJ and normalized to those of β-actin.

Deletion of *SPAR*. This was performed following a procedure described previously (3).

Briefly, the ~ 1kb region upstream of *sparA* and the ~1kb region downstream of *sparL* were

synthesized by Genscript and cloned into pUC57 (Genscript). The insert was then subcloned into

a temperature sensitive conjugative plasmid pG1-*oriT*$_{TnGBS1}$. The insert sequence was verified by

Sanger sequencing. The construct was introduced into *S. agalactiae* NEM316 by electroporation

and then into *Sgg* strain TX20005 by conjugation under the permissive temperature. PCR was

used to screen for double cross-over deletion mutants. Deletion was further confirmed by PCR

amplification of the regions spanning the deleted fragment and DNA sequencing of the PCR

product.

RNA extraction, cDNA synthesis and PCR. This was performed following the method

described previously (3). Briefly, RNA was extracted from TX20005 cultured in the presence of

HT29, as well as from the colonic tissues of mice that had been orally gavaged with TX20005.

RNA was treated with DNase and synthesized using a ProtoScript II First Strand cDNA

Synthesis Kit (NEB). Primers used in PCR amplification are: *sparA*, forward

5'GCAAGCTGGTCGAACAGAAC and reverse 5'GCTTCTATGGTTGGGGCTAGA; *sparD*,

forward 5' GGAGGTGGATCCAACAAGGG and reverse 5'CAGGTTCCTCGATAGCCAGC;

and *sparG*, forward 5'TCAGTTGTTAGCGGATGCGT and reverse 5'

CCCTTTATTGCTTGTGCTCCC.

Growth curves. Overnight cultures of *Sgg* were inoculated into fresh BHI broth at 1:100

dilution and grown at 37°C with shaking. Samples were taken at 0, 3, 6, 9, 12, and 24 hours,

dilution plated onto agar plates, incubated for 24 hours and colonies enumerated.

Animal experiments. Animal studies were performed in accordance with protocols

approved by the Institutional Animal Care and Use Committee at the Texas A&M Health

Science Center, Institute of Biosciences and Technology. Mice were fed with standard ProLab

IsoPro RMH3000 (LabDiet). **1) Colonization.** This was performed as previously described (3)

with slight modifications. Briefly, 6-week-old A/J mice, sex matched (Jackson Laboratory), were

treated with ampicillin at a concentration of 1 g/L in drinking water for 3 days and switched to

antibiotic-free water 24 hours prior to the administration of bacteria. *Sgg* was orally gavaged at a

dose of ~1 x 10^9 CFU/mouse. Colons and fecal materials were collected at day 1, 3, and 7 post-

gavage. Samples were weighed, homogenized in sterile PBS in a TissueLyser (Qiagen), dilution

plated onto Enterococcus Selective Agar (ESA) plates and incubated at 37°C for 24–48 hours to

enumerate *Sgg* colonies. **2) AOM-induced model of CRC.** This was performed as previously

described (3) with slight modifications. Briefly, 6-week-old A/J mice, sex matched were treated

with 4 weekly i.p. injections of AOM (10 mg/kg body weight), followed by ampicillin in

drinking water (1 g/L) for 7 days and switched to antibiotic-free water 24 hours prior to the first

oral gavage with *Sgg*. Mice were orally gavaged with bacteria at ~ 1x10^9 CFU/mouse or saline (n

= 11-12 per group) once per week for 12 weeks. Mice were euthanized one week after the last

oral gavage by CO_2 inhalation followed by cervical dislocation. Colon and fecal pellets were

collected. The number and size of macroscopic were recorded. Visual evaluation of colons was

carried out by a blinded observer. A random subset of fecal pellets were weighed, homogenized

in sterile PBS and dilution plated onto ESA plates.

Statistical analysis. GraphPad Prism 9 was used for statistical analyses. Two-tailed

unpaired *t*-test was used for pairwise comparisons to assess the significance of differences

between two groups in cell proliferation assays, western blot analysis, and adherence assays. The

non-parametric Mann-Whitney test was used to assess the significance of differences of results

between groups in animal studies. Two-way ANOVA was used to analyze the bacterial growth

curves. Ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Ethics statement. Animal studies were performed in accordance with protocols

(IACUC#2017-0420-IBT) approved by the Institutional Animal Care and Use Committee at the

Texas A&M Health Science Center, Institute of Biosciences and Technology. The Texas A&M

University Health Science Center—Institute of Biosciences and Technology is registered with the

Office of Laboratory Animal Welfare per Assurance A4012-01. It is guided by the PHS Policy on

Human Care and Use of Laboratory Animals (Policy), as well as all applicable provisions of the

Animal Welfare Act.

Results

The *Sgg* SPAR locus. We compared the genome sequence of TX20005

(NZ_CP077423.1) with that of ATCC 43143 (NC_017576.1), an *Sgg* strain that is defective in

stimulating CRC cell proliferation or promoting the development of colon tumors, using the

multiple genome alignment tool MAUVE (20). The SPAR locus is one of the regions in the

chromosome of TX20005 that display differences from ATCC 43143. The locus is comprised of

12 genes (*sparA* to *sparL*) (Fig. 1A). The protein sequences encoded by these genes were

analyzed for secretion signals using SignalP - 6.0 (21). None of them contains typical secretion

signals recognized by the Sec or Tat translocon. SparA, SparK and SparL are predicted to

contain transmembrane helices (TMHMM2.0) (22) and are thus putative transmembrane

proteins. Homology search using Protein BLAST showed that SparA to C are hypothetical

proteins of unknown function (Table 1). SparD to SparF exhibit homology to predicted ATP-

dependent endonuclease of the OLD family, GIY-YIG endonuclease/PcrA/UvrD helicase, and

GntR family transcriptional regulator, respectively. SparG to SparJ display features of effectors

secreted by the type VII secretion system (T7SS). SparG, H and J are homologous to T7SS

effectors, LXG domain-containing protein and T7SS effector LXG polymorphic toxin. SparG, H

and I are of 91, 102 and 137 amino acids in length, in keeping with T7SS effectors belonging to

the WXG100 family. Analysis using HHpred (23) showed that SparG and SparI fold into a four-

helical bundle structure typical of ESAT-6-like WXG100 proteins, subfamily of T7SS effectors,

with a probability of 96.48% and 97.97%, respectively. Spar H, I and J also contain a conserved

sequence motif HxxxD/ExxhxxxH (H denotes highly conserved and h less conserved

hydrophobic residues) that is considered to be important for the secretion of WXG100 proteins

by T7SS (24) (Supplemental Fig. S1). Immediately downstream of the *sparJ* gene are two genes

that encode putative TipC family immunity proteins. Genes encoding T7SS effector LXG

polymorphic toxins are typically clustered together with genes encoding their cognate immunity

proteins (25-27). The arrangement of *sparJ*, *K* and *L* further suggests that SparJ is a putative

T7SS effector LXG polymorphic toxin and SparK and L are likely the cognate immunity

proteins.

We next tested if the *SPAR* genes were expressed *in vitro* and *in vivo* by RT-PCR (Fig.

1B). We focused on the expression of *sparA*, *D* and *G*, under conditions that are relevant to the

pro-proliferation and pro-tumor activity of *Sgg*. All three genes were expressed when the

bacteria were co-cultured with HT29 cells and in the colonic tissues collected from mice orally

gavaged with TX20005 (Fig. 1B).

SPAR is important for *Sgg* adherence. To investigate the role of the *SPAR* locus in the

pathogenic potential of *Sgg*, we generated a mutant in the TX20005 background, in which all 12

genes were deleted (TX20005Δ*SPAR*). We verified that the deletion does not affect bacterial

growth (Fig. 2A). We sought to determine if deletion of the *SPAR* locus had any effect on the

adherence capacity of *Sgg*. The results showed that the mutant strain exhibited a significantly

reduced adherence to HT29 cells compared to the wild type (WT) parent strain TX20005 (~2%

vs ~9%) (Fig. 2B), suggesting that the *SPAR* locus is important for *Sgg* adherence to host cells.

Previous work showed that T7SS-secreted factors of *Sgg* significantly enhanced the adherence

capacity of the WT bacteria (3). We tested whether filter sterilized culture supernatants (CS)

from TX20005Δ*SPAR* can enhance the adherence capacity of TX20005. Addition of CS from the

WT TX20005 significantly enhanced the adherence of WT bacteria, whereas addition of the

mutant CS did not cause any increase in the adherence of WT bacteria (Fig. 2B, indicated by the

arrow), suggesting that *SPAR*-dependent secreted factors contribute to *Sgg* adherence .

SPAR is required for *Sgg* to stimulate CRC cell proliferation. *Sgg* has been previously

shown to stimulate cell proliferation in human CRC cell lines HT29 and HCT116, but not in

SW480 or HEK293 cell lines (2-4). We examined the ability of TX20005Δ*SPAR* to stimulate

cell proliferation in these cell lines. HT29, HCT116, SW480, and HEK293 cells were co-

cultured with TX20005 or TX20005Δ*SPAR*. As anticipated, the WT strain significantly

stimulated HT29 and HCT116 cell proliferation compared to untreated cells (Fig. 3A and 3B). In

contrast, HT29 and HCT116 cells co-cultured with the mutant did not exhibit any significant

increase in cell proliferation, indicating that *SPAR* is required for *Sgg*-stimulated CRC cell

proliferation. Neither the WT nor the mutant had any effect on the proliferation of SW480 or

HEK293 cells (Fig. 3C and 3D), as expected.

To further validate the results, we investigated the effect of the mutant on cell

proliferation markers β-catenin and proliferating cell nuclear antigen (PCNA) in HT29 cells (Fig.

3E-G). Cells co-cultured with TX20005 exhibited a significant increase in the level of both β-

catenin and PCNA compared to cells cultured in media only, as expected. In contrast, cells co-

cultured with the *SPAR* mutant showed no significant increase in the level of either marker

compared to cells cultured in media only, further confirming that the *SPAR* locus is essential for

Sgg to stimulate cell proliferation.

Previous work demonstrated that filter sterilized CSs from WT *Sgg* strains induce CRC

cell proliferation (3, 28). Therefore, we examined the effect of CS from TX20005Δ*SPAR* on cell

proliferation (Fig. 2H). HT29 cells treated with the WT CS exhibited a significant increase in

cell proliferation, as expected, whereas cells treated with the mutant CS did not exhibit any

significant increase in cell proliferation. These results suggest that the defect in the *SPAR* mutant

to stimulate host cell proliferation is due, at least in part, to the absence of certain secreted

factors in the CS. Taken together, these results indicate that *SPAR*, and particularly secreted

factors dependent on *SPAR*, is required for *Sgg* to stimulate CRC cell proliferation *in vitro*.

***Sgg* gut colonization is impaired by *SPAR* deletion.** The ability of *Sgg* to colonize the

host gut is an important aspect of its pathogenic potential (2). We sought to determine if the

SPAR locus is involved in *Sgg* gut colonization. Mice were orally gavaged with WT or the *SPAR*

deletion mutant. Colon tissues as well as fecal pellets were collected at day 1, 3, and 7 post-

bacterial gavage to determine the bacterial load in the samples. At day 1, the *Sgg* bacterial load

in the colonic tissues from mice gavage with the WT and the mutant did not significantly differ.

At day 3 and 7, the bacterial load of the mutant was significantly decreased compared to that of

the WT (Fig. 4A). In the fecal pellets, a similar patten was observed, such that at day 1, the

bacterial load of mutant and WT strains did not significantly differ, while At days 3 and 7, the

bacterial load of the mutant was decreased significantly compared to that of the WT (Fig. 4B).

Taken together, these results indicate that the *SPAR* locus contributes to the colonization capacity

of *Sgg* in the normal colon.

237   **SPAR is essential for the ability of *Sgg* to promote the development of colon tumors**.

Next, we investigated if *SPAR* is important for *Sgg* to promote the development of colon tumors

using an AOM-induced CRC model (Fig. 5A). Mice gavaged with the WT strain exhibited a

significant increase in the colon tumor burden (Fig. 5B) compared to saline-gavaged mice,

consistent with previous results (3, 4). In contrast, mice exposed to the *SPAR* mutant showed no

increase in the tumor burden compared to the saline control and a significant reduction compared

to the WT-treated mice, suggesting that *SPAR* is important for *Sgg* to promote colon tumors. We

also determined the bacterial load in the fecal pellets isolated at the end of the experiment and

found no difference in the *Sgg* burden between WT or mutant-treated groups (Fig. 5C). We note

that this result could be due to the repeated gavages with bacteria during the experiment and does

not necessarily reflect the ability of the strains to colonize tumor-bearing colons in this model.

These results together suggest that *SPAR* plays a critical functional role in promoting the

development of colon tumors.

Discussion

In this study, we describe an *Sgg* chromosomal locus, *SPAR*, that plays a key role in *Sgg*

pathogenesis. Deletion of *SPAR* led to striking phenotypes *in vitro* and *in vivo*, including a

reduced capacity to adhere to CRC cells, decreased colonization of the normal colon, and a

complete loss of the ability to stimulate CRC cell proliferation *in vitro* and to promote the

development of colon tumors *in vivo*. These results highlight *SPAR* as a critical pathogenicity

determinant of *Sgg*.

Our data showed that deletion of the *SPAR* locus resulted in significantly reduced

adherence of *Sgg* to HT29 cells. Previous studies showed that T7SS-secreted factors enhance *Sgg*

adherence (3). Interestingly, our results revealed that deletion of *SPAR* eliminated the adherence-

enhancing activity in the CS, suggesting that the deletion might have affected the T7-secreted

factors involved in adherence. In terms of cell proliferation, we showed that the *SPAR* locus is

essential for *Sgg* to stimulate certain CRC cell proliferation. Furthermore, the results indicate that

SPAR is also important for the production of secreted factors responsible for stimulating host cell

proliferation. In this aspect, the *SPAR* mutant exhibits a very similar phenotype as that described

for a T7SS defective mutant of *Sgg* (TX20005Δ*esx*) with respect to adherence and stimulation of

cell proliferation (3). One possible reason for the phenotypic similarity is that the *SPAR* locus

encodes several putative T7SS effectors (SparG to SparJ). Examination of the TX20005 genome

reveals that the previously reported T7SS locus (*Sgg*T7SS^{T05}) (3) is the only locus that encodes a

set of proteins comprising the T7SS secretion apparatus. Thus, SparG to SparJ are likely secreted

by *Sgg*T7SS^{T05}. This would imply that one or more of the *SPAR*-encoded putative T7SS effectors

are involved in the adherence and the pro-proliferative activity of *Sgg*. On the other hand, the

sparF gene is predicted to encode a GntR family transcriptional regulator (29). A recent

publication indicated that a GntR family transcriptional regulator (OG1RF_11099) of

Enterococcus faecalis controls the expression of T7SS genes (30). Using Global Align (NCBI),

we found that SparF is highly homologous to OG1RF_11099, showing an overall 74% similarity

at the amino acid sequence level. Hence, a second possible reason for the phenotypic similarity

between TX20005*Δspar* and TX20005*Δesx* is that SparF regulates the expression of genes in the

*Sgg*T7SS^{T05} locus.

In a gut colonization model, we observed that deletion of *SPAR* resulted in reduced *Sgg*

bacterial load in the normal colon. This could be due to the involvement of *SPAR* in *Sgg*

adherence to the colonic epithelium. Additionally, SparJ is a putative T7SS-secreted

polymorphic toxin. This family of toxins are often involved in interbacterial competition (26).

Thus, SPAR may contribute to *in vivo* colonization by antagonizing other gut commensal

bacteria through SparJ.

In the AOM model of CRC, we observed a striking difference between the *SPAR* mutant

and the WT bacteria in promoting the development of colon tumors, such that the mutant has

completely lost the ability to promote the development of colon tumors. This phenotype is much

stronger than that previously reported for the T7SS mutant (3), raising the possibility that in

addition to T7SS genes, SparF also regulates the expression of other genes that contribute tumor

development *in vivo*. Studies to further investigate the biological activities of SparF, and SparG

to SparL are needed to delineate the specific contribution of the *SPAR* locus to the observed

phenotypes.

It is also possible that other genes within the *SPAR* locus, SparA-SparE, also play a role,

however, their predicted function based on homology search does not provide clues regarding

their respective contribution. SparA-C are putative hypothetical proteins. SparD is homologous

to OLD family endonucleases. OLD family endonucleases are widely present among bacteria

and archaea (31). While their biological function is not completely understood, they contain a C-

terminal ATPase domain, as well as an N-terminal Toprim domain, which is believed to be

important in DNA replication, recombination, and repair (32, 33). SparE is homologous to GIY-

YIG endonuclease (34-38) and PcrA/UvrD helicase. The PcrA/UvrD helicase is an essential

helicase in gram-positive bacteria and has been shown to play a role in DNA repair, as well as

the replication of small drug-resistance plasmids in *Staphylococcus aureus* and *Bacillus subtilis*

(39-45).

In summary, we have identified a chromosomal locus in *Sgg*, *SPAR*, that is critical to the

pathogenicity of *Sgg*. We report that deletion of *SPAR* significantly reduces the capacity of *Sgg*

to adhere to CRC cells and to colonize the gut. Furthermore, deletion of *SPAR* abrogates the

ability for *Sgg* to stimulate CRC cell proliferation and to accelerate colon tumor development.

Examination of genes with the locus highlighted several potential candidates responsible for the

observed phenotypes. The results suggest a connection between the *SPAR* locus and the T7SS of

Sgg in terms of additional T7SS effectors encoded by *SPAR* genes and a potential regulator of

T7SS expression. Further investigations into the activities of specific *SPAR* proteins will be

important for dissecting the intricate *Sgg* pathogenic mechanisms and will open up the path

ahead to identify biomarker candidates and targets for clinical prevention and intervention

strategies.

Acknowledgements

This study was supported by funding from the Hamill Foundation, the Cancer Prevention

Research Institute of Texas (RP170653) and NIAID (1R21AI151914) to Y. Xu and

(T32AI055449-16) to J. Taylor. The content of this manuscript is also available at bioRxiv.org

(https://doi.org/10.1101/2022.03.15.484266).

References

1.   Schlegel L, Grimont F, Ageron E, Grimont PAD, Bouvet A. Reappraisal of the taxonomy

of the Streptococcus bovis/Streptococcus equinus complex and related species: description of

Streptococcus gallolyticus subsp. gallolyticus subsp. nov., S. gallolyticus subsp. macedonicus

subsp. nov. and S. gallolyticus subsp. pasteurianus subsp. nov. Int J Syst Evol Microbiol.

2003;53(Pt 3):631-45.

2.   Kumar R, Herold JL, Taylor J, Xu J, Xu Y. Variations among Streptococcus gallolyticus

subsp. gallolyticus strains in connection with colorectal cancer. Scientific reports.

2018;8(1):1514.

3.   Taylor JC, Gao X, Xu J, Holder M, Petrosino J, Kumar R, et al. A type VII secretion

system of Streptococcus gallolyticus subsp. gallolyticus contributes to gut colonization and the

development of colon tumors. PLoS Pathog. 2021;17(1):e1009182.

4.   Kumar R, Herold JL, Schady D, Davis J, Kopetz S, Martinez-Moczygemba M, et al.

Streptococcus gallolyticus subsp. gallolyticus promotes colorectal tumor development. PLoS

Pathog. 2017;13(7):e1006440.

5.   Marmolin ES, Hartmeyer GN, Christensen JJ, Nielsen XC, Dargis R, Skov MN, et al.

Bacteremia with the bovis group streptococci: species identification and association with

infective endocarditis and with gastrointestinal disease. Diagn Microbiol Infect Dis.

2016;85(2):239-42.

6.   Corredoira J, Garcia-Pais MJ, Coira A, Rabunal R, Garcia-Garrote F, Pita J, et al.

Differences between endocarditis caused by Streptococcus bovis and Enterococcus spp. and their

association with colorectal cancer. Eur J Clin Microbiol Infect Dis. 2015;34(8):1657-65.

7.   Boleij A, Tjalsma H. Gut bacteria in health and disease: a survey on the interface

between intestinal microbiology and colorectal cancer. Biol Rev Camb Philos Soc.

2012;87(3):701-30.

8.   Boleij A, van Gelder MM, Swinkels DW, Tjalsma H. Clinical Importance of

Streptococcus gallolyticus infection among colorectal cancer patients: systematic review and

meta-analysis. Clin Infect Dis. 2011;53(9):870-8.

9.   Abdulamir AS, Hafidh RR, Abu Bakar F. The association of Streptococcus

bovis/gallolyticus with colorectal tumors: the nature and the underlying mechanisms of its

etiological role. J Exp Clin Cancer Res. 2011;30:11.

10.   Gupta A, Madani R, Mukhtar H. Streptococcus bovis endocarditis, a silent sign for

colonic tumour. Colorectal Dis. 2010;12(3):164-71.

11.   Waisberg J, Matheus CeO, Pimenta J. Infectious endocarditis from Streptococcus bovis

associated with colonic carcinoma: case report and literature review. Arq Gastroenterol.

2002;39(3):177-80.

12.   Alazmi W, Bustamante M, O'Loughlin C, Gonzalez J, Raskin JB. The association of

Streptococcus bovis bacteremia and gastrointestinal diseases: a retrospective analysis. Dig Dis

Sci. 2006;51(4):732-6.

13.   Gold JS, Bayar S, Salem RR. Association of Streptococcus bovis bacteremia with colonic

neoplasia and extracolonic malignancy. Arch Surg. 2004;139(7):760-5.

14.   Corredoira J, Grau I, Garcia-Rodriguez JF, Alonso-Garcia P, Garcia-Pais MJ, Rabunal R,

et al. The clinical epidemiology and malignancies associated with Streptococcus bovis biotypes

in 506 cases of bloodstream infections. J Infect. 2015;71(3):317-25.

15.   Zhang Y, Weng Y, Gan H, Zhao X, Zhi F. Streptococcus gallolyticus conspires myeloid

cells to promote tumorigenesis of inflammatory bowel disease. Biochem Biophys Res Commun.

2018;506(4):907-11.

16.   Thomas AM, Manghi P, Asnicar F, Pasolli E, Armanini F, Zolfo M, et al. Metagenomic

analysis of colorectal cancer datasets identifies cross-cohort microbial diagnostic signatures and

a link with choline degradation. Nat Med. 2019;25(4):667-78.

17.   Perichon B, Lichtl-Hafele J, Bergsten E, Delage V, Trieu-Cuot P, Sansonetti P, et al.

Detection of Streptococcus gallolyticus and Four Other CRC-Associated Bacteria in Patient

Stools Reveals a Potential "Driver" Role for Enterotoxigenic Bacteroides fragilis. Frontiers in

cellular and infection microbiology. 2022;12:794391.

18.   Kwong TNY, Wang X, Nakatsu G, Chow TC, Tipoe T, Dai RZW, et al. Association

Between Bacteremia From Specific Microbes and Subsequent Diagnosis of Colorectal Cancer.

Gastroenterology. 2018;155(2):383-90 e8.

19.   Abdulamir AS, Hafidh RR, Bakar FA. The association of Streptococcus

bovis/gallolyticus with colorectal tumors: The nature and the underlying mechanisms of its

etiological role. J Exp Clin Cancer Res. 2011;30:11.

20.   Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved

genomic sequence with rearrangements. Genome Res. 2004;14(7):1394-403.

21.   Teufel F, Almagro Armenteros JJ, Johansen AR, Gislason MH, Pihl SI, Tsirigos KD, et

al. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat

Biotechnol. 2022.

22.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein

topology with a hidden Markov model: application to complete genomes. J Mol Biol.

2001;305(3):567-80.

23.   Hildebrand A, Remmert M, Biegert A, Soding J. Fast and accurate automatic structure

prediction with HHpred. Proteins. 2009;77 Suppl 9:128-32.

24.   Poulsen C, Panjikar S, Holton SJ, Wilmanns M, Song YH. WXG100 protein superfamily

consists of three subfamilies and exhibits an α-helical C-terminal conserved residue pattern.

PLoS One. 2014;9(2):e89313.

25.   Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L. Polymorphic toxin

systems: Comprehensive characterization of trafficking modes, processing, mechanisms of

action, immunity and ecology using comparative genomics. Biol Direct. 2012;7:18.

26.   Whitney JC, Peterson SB, Kim J, Pazos M, Verster AJ, Radey MC, et al. A broadly

distributed toxin family mediates contact-dependent antagonism between gram-positive bacteria.

Elife. 2017;6.

27.   Klein TA, Pazos M, Surette MG, Vollmer W, Whitney JC. Molecular Basis for Immunity

Protein Recognition of a Type VII Secretion System Exported Antibacterial Toxin. J Mol Biol.

2018;430(21):4344-58.

28.   Taddese R, Garza DR, Ruiter LN, de Jonge MI, Belzer C, Aalvink S, et al. Growth rate

alterations of human colorectal cancer cells by 157 gut bacteria. Gut Microbes. 2020;12(1):1-20.

29.   Liu GF, Wang XX, Su HZ, Lu GT. Progress on the GntR family transcription regulators

in bacteria. Yi Chuan. 2021;43(1):66-73.

30.   Chatterjee A, Willett JLE, Dunny GM, Duerkop BA. Phage infection and sub-lethal

antibiotic exposure mediate Enterococcus faecalis type VII secretion system dependent inhibition

of bystander bacteria. PLoS Genet. 2021;17(1):e1009204.

31.   Schiltz CJ, Lee A, Partlow EA, Hosford CJ, Chappie JS. Structural characterization of

Class 2 OLD family nucleases supports a two-metal catalysis mechanism for cleavage. Nucleic

Acids Res. 2019;47(17):9448-63.

32.   Aravind L, Leipe DD, Koonin EV. Toprim--a conserved catalytic domain in type IA and

II topoisomerases, DnaG-type primases, OLD family nucleases and RecR proteins. Nucleic

Acids Res. 1998;26(18):4205-13.

33.   Koonin EV, Gorbalenya AE. The superfamily of UvrA-related ATPases includes three

more subunits of putative ATP-dependent nucleases. Protein Seq Data Anal. 1992;5(1):43-5.

34.   Kowalski JC, Belfort M, Stapleton MA, Holpert M, Dansereau JT, Pietrokovski S, et al.

Configuration of the catalytic GIY-YIG domain of intron endonuclease I-TevI: coincidence of

computational and molecular findings. Nucleic Acids Res. 1999;27(10):2115-25.

35.   Van Roey P, Meehan L, Kowalski JC, Belfort M, Derbyshire V. Catalytic domain

structure and hypothesis for function of GIY-YIG intron endonuclease I-TevI. Nat Struct Biol.

2002;9(11):806-11.

36.   Truglio JJ, Rhau B, Croteau DL, Wang L, Skorvaga M, Karakas E, et al. Structural

insights into the first incision reaction during nucleotide excision repair. EMBO J.

2005;24(5):885-94.

37.   Dunin-Horkawicz S, Feder M, Bujnicki JM. Phylogenomic analysis of the GIY-YIG

nuclease superfamily. BMC Genomics. 2006;7:98.

38.   Ibryashkina EM, Sasnauskas G, Solonin AS, Zakharova MV, Siksnys V. Oligomeric

structure diversity within the GIY-YIG nuclease family. J Mol Biol. 2009;387(1):10-6.

39.   Bruand C, Ehrlich SD. UvrD-dependent replication of rolling-circle plasmids in

Escherichia coli. Mol Microbiol. 2000;35(1):204-10.

40.   Chang TL, Naqvi A, Anand SP, Kramer MG, Munshi R, Khan SA. Biochemical

characterization of the Staphylococcus aureus PcrA helicase and its role in plasmid rolling circle

replication. J Biol Chem. 2002;277(48):45880-6.

41.   del Solar G, Giraldo R, Ruiz-Echevarría MJ, Espinosa M, Díaz-Orejas R. Replication and

control of circular bacterial plasmids. Microbiol Mol Biol Rev. 1998;62(2):434-64.

42.   Naqvi A, Tinsley E, Khan SA. Purification and characterization of the PcrA helicase of

Bacillus anthracis. J Bacteriol. 2003;185(22):6633-9.

43.   Khan SA. Rolling-circle replication of bacterial plasmids. Microbiol Mol Biol Rev.

1997;61(4):442-55.

44.   Khan SA. Plasmid rolling-circle replication: recent developments. Mol Microbiol.

2000;37(3):477-84.

45.   Soultanas P, Dillingham MS, Papadopoulos F, Phillips SE, Thomas CD, Wigley DB.

Plasmid replication initiator protein RepD increases the processivity of PcrA DNA helicase.

Nucleic Acids Res. 1999;27(6):1421-8.

Table 1. Proteins encoded by the *SPAR* genes.

Gene	Length (aa)[a]	Predicted Gene Function[b]
sparA	424	Hypothetical Protein
sparB	53	Hypothetical Protein
sparC	582	Hypothetical Protein
sparD	656	OLD-Family Endonuclease
sparE	609	GIY-YIG Endonuclease/PcrA/UvrD Helicase
sparF	232	GntR-Family Transcriptional Regulator
sparG	91	TIGR04197-Family T7SS Effector
sparH	102	Hypothetical Protein
sparI	137	LXG domain-containing protein
sparJ	417	T7SS effector LXG polymorphic toxin
sparK	199	TipC Family Immunity Protein
sparL	197	TipC Family Immunity Protein

451 [a]Based on the annotated genome of TX20005 (NZ_CP077423.1)

452 [b]The amino acid sequence of each protein was used to search the non-redundant protein

453 sequences database (updated on 04/05/2022) at NCBI using Protein BLAST.

 Figure Legends

Fig 1. *SPAR* **genetic organization and expression. A.** The *SPAR* locus. Genes encoding

proteins with homology to proteins of known functions or containing characteristics of certain

protein families are colored in blue, while genes encoding hypothetical proteins of unknown

function are colored in gray. **B. RT-PCR.** cDNA synthesized from RNA extracted from HT29

cells co-cultured with TX20005 and the colonic tissues of three mice that had been orally

gavaged with TX20005 were used as a template for the RT-PCR. cDNA from HT29 cells only

and from mice orally gavaged with saline were used as controls to show the specificity of the

PCR primers. RNA samples that had not been treated with reverse transcriptase (RT) were used

as controls to show the lack of DNA contamination.

Fig 2. **SPAR is important for** *Sgg* **adherence but does not impact growth. A. Growth Curve**.

A single colony of WT or mutant bacteria was inoculated into 1mL of fresh BHI and incubated

at 37ºC overnight with shaking. The overnight culture was then diluted 1:100 with fresh BHI and

incubated at 37ºC with shaking. 100μL of the culture were taken at the 0, 3, 6, 9, 12, and 24

hours for serial dilution and plating. Results were combined from three biological replicates. **B.**

Adherence. HT29 cells were seeded at a density of $1x10^6$ cells per well and incubated with

TX20005 or TX20005*ΔSPAR* (MOI=10) in media only or in the presence of CS from TX20005

or TX20005*ΔSPAR* for 1 hour. Percentage adherence was calculated as the percentage of

adhered bacteria vs total bacteria added. Results were combined from three biological replicates.

Unpaired two-tailed *t* tests were used. **, $p<0.01$; NS, not significant.

Fig 3. *SPAR* **is essential for** *Sgg* **to stimulate CRC cell proliferation. A-D. Cell proliferation**

assay. HT29 (**A**), HCT116 (**B**), SW480 (**C**), and HEK293 (**D**) cells were incubated in the

presence or absence of bacteria (MOI=1) or CSs for 24 hours, as described in the Materials and

Methods section. Cell proliferation was measured using a CCK-8 kit. Cell-free wells containing

only the culture media were served as a blank control to which the absorbance values were

normalized. **E-G. Western blot**. Cell lysates from HT29 cells co-cultured in the presence or

absence of bacteria (MOI=1) were analyzed by western blot, probed with antibodies against β-

catenin, PCNA, and β-actin (**E**). Band intensity was quantified and normalized to β-actin. Fold

change in β-catenin (**F**) and PCNA (**G**) were calculated against cells incubated in media only.

Data was combined from three biological replicates. **H. Cell proliferation in the presence of**

Sgg **CSs.** HT29 cells were cultured in the presence or absence of WT or mutant *Sgg* (MOI=1) or

CSs from WT or mutant *Sgg* for 24 hours. Cell proliferation was measured by using a CCK-8 kit.

Results were combined from at least three biological replicates. Unpaired two-tailed *t* tests were

used for the comparisons. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$; ns, not

significant. Significance in panel H indicates comparison to untreated cells.

Fig 4. *SPAR* **deletion reduced the colonization capacity of** *Sgg*. This was performed as

described in the Materials and Methods section. Colon tissues (**A**) and fecal pellets (**B**) were

weighed, homogenized in sterile PBS and dilution plated onto ESA plates. Mann-Whitney test

was used for the comparisons. *, $p<0.05$; **, $p<0.01$; ****, $p<0.0001$; NS, not significant.

Fig 5. *SPAR* **is critical for** *Sgg* **to promote the development of colon tumors**. This was

performed as described in the Materials and Methods section. The procedure is illustrated in **A**.

499   **B.** The number and size of macroscopic tumors were recorded by blinded observers. Tumor

500   burden is calculated as the sum of tumor volumes per mouse. **C.** *Sgg* load in the fecal pellets

501   collected at the experimental endpoint was determined by dilution plating of homogenized

502   pellets onto ESA plates. Mann-Whitney test was used for all comparisons. *, $p<0.05$; **, $p<0.01$;

503   ***, $p<0.001$; ****, $p, <0.0001$; NS, not significant.

Fig 1

A

B

Fig 2

Fig 3

Fig 4

Fig 5

Fig S1

```
SPAR_H     GETDEALAELKKQEDDLEDALVDLLRKEAQDEH
SPAR_I     DQQQQCLNVLEEAELQLSLISVMSLERKTSLHQ
SPAR_J     EVKTTIFSSLVPKEVVAINPVTGAPIYASVSEDEAMAYLQENYEGSYLFIQSLEEEENLFGEDN
Consensus           HxxxDxxhxxxH
                        E
```

Supplemental Figure S1: The conserved T7SS sequence in SparH-J. H denotes highly conserved and h less conserved hydrophobic residues.