A pathogenicity locus of *Streptococcus gallolyticus* subspecies *gallolyticus*

John Culver Taylor[1], Ritesh Kumar[1#], Juan Xu[1], and Yi Xu[1,2,3]*

[1]Center for Infectious and Inflammatory Diseases, Texas A&M Health Science Center Institute

of Biosciences of Technology, Houston, Texas, United States of America

[2]Department of Microbiology and Molecular Genetics, McGovern Medical School, UT Health,

Houston, Texas, United States of America

[3]Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas A&M

Health Science Center, Texas, United States of America

[#2]Current affiliation: IFF Health and Biosciences

*Corresponding author: Yi Xu

E-mail: yi-xu1@tamu.edu

Key words: Colorectal cancer, colon cancer, microbiome, *Streptococcus gallolyticus*,

Streptococcus bovis, cell proliferation, colonization.

16   **Abstract**

Streptococcus gallolyticus subspecies *gallolyticus* (*Sgg*) is a pathobiont of the gut with

clinical significance. *Sgg* adheres to the colonic epithelium and stimulates colorectal cancer

(CRC) cell proliferation. In pre-clinical models of CRC, *Sgg* promotes the development of colon

tumors. However, the molecular determinants of *Sgg* virulence remain unclear. Here, we

identified a 12-gene locus in *Sgg* strain TX20005. Deletion of this locus resulted significantly

reduced capacity to adhere to CRC cells and abrogated the ability of *Sgg* stimulate CRC cell

proliferation. Thus, we designate this locus as the <u>S</u>gg <u>p</u>athogenicity-<u>a</u>ssociated <u>r</u>egion (SPAR).

In a gut colonization model, mice exposed to the SPAR mutant showed significantly reduced *Sgg*

load in the colon tissues and fecal materials. In a mouse mode of CRC, deletion of *SPAR*

abolished the ability of *Sgg* to promote the development of colon tumors growth, while the fecal

Sgg load in the CRC model does not show a significant difference between the wild type and the

SPAR mutant. The results also show a connection between the *SPAR* locus and a previously

described type VII secretion system of *Sgg*. Taken together, these results indicate that SPAR

genes are critical pathogenicity determinants of *Sgg*. Studies to elucidate the specific activities of

individual SPAR genes will be important.

Introduction

Streptococcus gallolyticus subspecies *gallolyticus* (*Sgg*) is a group D streptococcal bacterium, and an opportunistic pathogen belonging to the *Streptococcus bovis*/*Streptococcus equinis* complex[1]. *Sgg* primarily causes bacteremia and infective endocarditis (IE), and has had a long-standing clinical association with CRC, supported by extensive studies and case reports dating back to the 1950s[2-16]. Patients with *Sgg* IE and/or bacteremia have concomitant colon adenomas or adenocarcinomas at a much higher rate (~60%) than that of the general population[8]. Moreover, a prospective study found that 45% of patients with *Sgg* IE (45%) developd colonic neoplastic lesions within 5 years compared to 21% of patients with IE due to closely-related enterococci, further supporting the association of *Sgg* with CRC[6]. A recent metagenomic study of datasets from multiple countries across the continents confirmed that *Sgg* is enriched in colon tumors and can be a biomarker for CRC[17].

Functionally, *Sgg* stimulates the proliferation of human CRC cells in a β-catenin dependent manner *in vitro* and in a xenograft model[4]. Exposure to *Sgg* led to larger and more advanced tumors in an azoxymethane (AOM)-induced CRC model and in a colitis-associated CRC model[4,15]. These results highlight a pro-proliferation and pro-tumor activity of *Sgg*. Recent studies showed that there are variations among *Sgg* strains in their ability to stimulate human CRC cell proliferation. Some *Sgg* strains such as ATCC43143 showed significantly reduced capacity to adhere to HT29 cells, to stimulate the proliferation of these cells and to promote tumor development in the AOM model[2], suggesting that there are specific *Sgg* factors contributing to the pro-proliferative and pro-tumor activity of *Sgg*. The identity of these specific *Sgg* factors was unknown.

Here, we describe work characterizing a 12-gene locus in the chromosome of *Sgg* strain

TX20005. Deletion of this locus significantly impaired the ability of *Sgg* to adhere to cultured

CRC cells and to stimulate CRC cell proliferation. *In vivo*, the deletion mutant displayed

significantly reduced capacity to colonize normal mouse colon. More importantly, the mutant

lost the ability to promote the development of colon tumors. Given the role of this locus in *Sgg*

pathogenesis, we have coined this locus as the *Sgg* pathogenicity-associated region (*SPAR*).

61   **Results**

The *Sgg* SPAR locus. The SPAR locus was identified by comparing the genome

sequence of TX20005 (NZ_CP077423.1) with that of ATCC 43143 (NC_017576.1) using the

multiple genome alignment tool MAUVE[18]. The locus is situated in a region displaying

substantial differences between the strains. The locus is comprised of 12 genes (*sparA* to *sparL*),

organized into 3 putative operons, *SPAR 1, 2*, and *3*, and one standalone gene, *sparC* (Fig. 1A).

The majority of the *spar* genes except for *sparD*, *E*. and *F* encode hypothetical proteins of

unknown function (Table 1). SparD to SparF exhibits homology to Predicted ATP-dependent

endonuclease of the OLD family, GIY-YIG endonuclease/PcrA/UvrD helicase, and GntR family

transcriptional regulator, respectively.

We next sought to determine if the *SPAR* genes were expressed *in vitro* and *in vivo* by

RT-PCR (Fig 1B). We focused on the expression of the first gene of the three putative operons,

sparA, *D* and *G*, under conditions that are relevant to the pro-proliferation and pro-tumor activity

of *Sgg*. All three genes were expressed when the bacteria were co-cultured with HT29 cells (Fig.

1B). We further analyzed the expression of these genes in the colonic tissues collected from mice

orally gavaged with TX20005. The result showed that these genes were also expressed in the

colon of mice (Fig. 1B).

SPAR is important for *Sgg* adherence. To investigate the role of the *SPAR* locus in the

pathogenic potential of *Sgg*, we generated a mutant in the TX20005 background, in which all 12

genes were deleted (TX20005Δ*SPAR*). We verified that the deletion does not affect bacterial

growth (Fig. 2A).

We sought to determine if deletion of the *spar* locus had any effect on the adherence

capability of *Sgg*. The results showed that the mutant strain exhibited a significantly decreased

capacity for adherence to HT29 cells compared to the WT parent strain (Fig 2B). Previous work

indicated that type VII-secreted factors of *Sgg* are involved in adherence by significantly

enhancing the adherence capacity of the WT bacteria[3]. We tested whether filter-sterilized culture

supernatants (CS) from TX20005Δ*SPAR* can enhance the adherence capacity of TX20005.

Addition of the WT CS significantly enhanced the adherence of WT bacteria, but not that of the

mutant. Further, addition of the mutant supernatant did not render any increase in WT adherence,

nor did it enhance mutant adherence. These results suggest that the *SPAR* locus is important for

Sgg to adhere to CRC cells. Furthermore, the results also suggest that the contribution of the

SPAR locus to adherence involves secreted factors and surface adhesins.

SPAR is required for *Sgg* to stimulate CRC cell proliferation. *Sgg* has been previously

shown to stimulate cell proliferation in human CRC cell lines HT29 and HCT116, but not in

HEK293 or SW480 cell lines[2-4]. We examined the ability of TX20005Δ*SPAR* to stimulate cell

proliferation in these cell lines. HT29, HCT116, SW480, and HEK293 cells were co-cultured

with the wild type (WT) strain TX20005 or TX20005Δ*SPAR* (Fig 3A-D). As anticipated, the WT

strain significantly stimulated HT29 and HCT116 cell proliferation compared to untreated cells.

In contrast, HT29 and HCT116 cells exposed to the mutant did not exhibit any significant

increase in cell proliferation, indicating that *SPAR* is required for *Sgg*-stimulated CRC cell

proliferation. Neither the WT or the mutant had any effect on the proliferation of SW480 or

HEK293 cells, as expected.

To further validate the results, we investigated the effect of the mutant on cell

proliferation markers β-catenin and proliferating cell nuclear antigen (PCNA) in HT29 cells (Fig.

3E-G). Cells co-cultured with WT TX20005 exhibited a significant increase in the levels of both

β-catenin and PCNA compared to cells cultured in media only, as expected. Notably, cells co-

cultured with the *SPAR* mutant showed no difference in the level of β-catenin or PCNA

compared to cells cultured in media only, further confirming that the *spar* locus is essential for

Sgg to stimulate cell proliferation.

Previous work demonstrated that filter-sterilized CS from *Sgg* is able to induce cell

proliferation[3,19]. Therefore, we examined the effect of CS from TX20005Δ*SPAR* (Fig. 2H). Cells

treated with the WT CS exhibited a significant increase in cell proliferation, as expected, while

cells treated with the mutant CS did not exhibit any significant increase in cell proliferation.

These results suggest that the defect in the *SPAR* mutant to stimulate host cell proliferation is due

to, at least in part, the absence of secreted factors in the CS. Taken together, these results indicate

that *SPAR*, and particularly secreted factors dependent on *SPAR* is required for *Sgg* to stimulate

CRC cell proliferation *in vitro*.

Sgg gut colonization is impaired by SPAR deletion. The ability of *Sgg* to colonize the

host gut is an important aspect of its pathogenic potential[2]. We sought to determine if the *SPAR*

locus is involved in *Sgg* gut colonization (Fig 4, A and B). Mice were orally gavaged with WT

and the mutant bacteria. Colon tissues as well as fecal materials were collected at days 1, 3, and

7 post gavage to determine the bacterial load in the samples. In the colon tissues, the bacterial

load of the WT and mutant strains did not significantly differ at day 1. However, the bacterial

load of the mutant was significantly decreased compared to that of the WT at days 3 and 7 (Fig.

4A). In the fecal pellets, a similar patten was observed, such that at the 1-day mark, the bacterial

loads of mutant and WT strains did not significantly differ. At days 3 and 7, the bacterial load of

the mutant was decreased significantly compared to that of the WT (Fig. 4B). Taken together,

these results indicate that the *SPAR* locus is an important contributor to the colonization capacity

of *Sgg* in the normal colon.

SPAR is essential for the ability of *Sgg* to promote the development of colon tumors.

Next, we investigated if *SPAR* is important for the ability of *Sgg* to promote the development of

colon tumors. We examined the effect of TX20005Δ*SPAR* on macroscopic tumors in the AOM-

induced CRC model (Fig 5A). Mice gavaged with the WT strain exhibited a significant increase

in tumor incidence (Fig. 5B), tumor multiplicity (Fig. 5C), tumor size (Fig. 5D) and total tumor

burden (Fig 5E) compared to saline-gavaged mice. In contrast, mice exposed to the mutant

showed no significant changes in incidence, multiplicity or size compared to saline treated mice,

and significant reduction compared to WT-treated mice (Fig. 5B-E). We also determined the

bacterial load in the fecal pellets isolated at collection and found no difference in the *Sgg* burden

between WT or mutant-treated groups (Fig. 5F). We note that this result could be due to the

repeated gavage with bacteria during the experiment and is not a definitive measure of the ability

of the strains to colonize tumor-bearing colons. Taken together, these results suggest that the

SPAR locus plays a critical role in *Sgg*-induced of colon tumor development.

Discussion

In this study, we describe a 12-gene *Sgg* chromosomal locus, *SPAR*, that plays a key role

to *Sgg* pathogenicity in the gut. Deletion of *SPAR* renders *Sgg* deficient in adherence to CRC

cells and colonization of the colon, abrogates the ability for *Sgg* to stimulate CRC cell

proliferation *in vitro* and to promote the development of colon tumors *in vivo*. Furthermore,

results from this study also suggested that multiple factors within the *SPAR* locus may contribute

to its phenotype in a direct or indirect manner.

Our data showed that deletion of the *SPAR* locus resulted in significantly reduced

adherence of *Sgg* to HT29 cells and colonization of the colon. Previous studies showed that

T7SS-secreted factors enhance *Sgg* adherence[3]. Interestingly, our results revealed that *SPAR*

deletion eliminated the adherence-enhancing activity in the CS, suggesting that the deletion

might have affected the T7-secreted factors involved in adherence. Furthermore, previous work

showed that T7SS-dependent surface anchored factor(s) is required for adherence[3]. The *SPAR*

mutant also exhibited a similar deficiency in the surface anchored factor(s), in that the CS from

WT bacteria could not enhance the adherence of the *SPAR* mutant. In terms of cell proliferation,

we showed that the *SPAR* locus is essential for *Sgg* to stimulate certain CRC cell proliferation.

Furthermore, the results suggest that *SPAR* is also important for the production of secreted

factors that stimulate host cell proliferation. Thus, the *SPAR* mutant exhibits a very similar

phenotype as that described for a T7SS defective mutant of *Sgg* (TX20005Δ*esx*) with respect to

adherence and induction of cell proliferation[3]. In the AOM model of CRC, we observed a

striking difference between the *SPAR* mutant and the WT bacteria in promoting the development

of colon tumors. This phenotype is much stronger than previously reported for the T7SS mutant[3],

suggesting that the *SPAR* locus has a more extensive tumor promoting activity than the T7SS of

in vivo.

Further examination of genes within the *SPAR* locus showed that *sparG – sparK* in the

third putative operon display some characteristics of proteins secreted by T7SS. SparG, H and I

are of 91, 102 and 137 amino acids in length. Analysis with HHpred showed that SparG and

SparI fold into a four-helical bundle structure typical of ESAT-6-like WXG100 proteins with a

probability of 96.48% and 97.97%, respectively. SparH to SparK contain a conserved sequence

motif HxxxD/ExxhxxxH (H denotes highly conserved and h less conserved hydrophobic

residues) that is considered to be important for the secretion of WXG100 proteins by T7SS[20]

(Supplemental Fig. S1). SparH – SparK also contain an LXG motif present in WXG100 proteins

or T7SS-secreted LXG-toxins. Thus, SparG to SparK are potential T7SS secreted proteins.

Genes encoding the T7 secretion machinery are not identified within the *SPAR* locus. In fact, the

previously identified T7SS (*Sgg*T7SS[T05]) is the only T7SS locus in TX20005 that contains a

complete set of secretion machinery genes. It is likely that SparG to SparK are secreted through

*Sgg*T7SS[T05].

The *sparF* gene is predicted to encode a GntR family transcriptional regulator. This

family of regulators contain a conserved N-terminal DNA binding domain and a C-terminal

effector-binding/oligomerization domain which varies between different GntR family members[21].

A recent publication indicated that a GntR family transcriptional regulator (OG1RF_11099) of

Enterococcus faecalis played an important role in the activation of T7SS gene transcription[22].

Using Global Align (NCBI), we found that SparF is highly homologous to the OG1RF_11099,

showing a 74% positivity at the amino acid sequence level. This indicates that in addition to the

putative T7SS secreted proteins in the *SPAR* locus, SparF can potentially regulate the expression

of T7SS genes. Further studies are needed to reveal the specific activities of SparF to SparK in

the context of adherence and cell proliferation. However, the presence of these genes within the

SPAR locus is likely a reason for the phenotypic similarity between the *SPAR* and T7SS mutants.

191   Our data also showed that the *SPAR* mutant has a more pronounced phenotype than the

T7SS mutant *in vivo*, suggesting that in addition to *Sgg*T7SS[T05]-mediated activities, the *SPAR*

locus may encode or regulate functions that faciliate tumor development. One possibility is that

SparF can regulate the expression of genes in addition to T7SS genes. Those genes may

contribute to the more severe defects observed for the *SPAR* mutant. It is also possible that other

genes within the *SPAR* locus, SparA-SparE, play a role, however, their predicted function based

on homology search does not provide clues regarding their possible contribution. SparA-C are

putative hypothetical proteins. SparD is homologous to OLD family endonucleases. OLD family

endonucleases are widely present among bacteria and archaea[23]. While their biological function

is not completely understood, they contain a C-terminal ATPase domain, as well as an N-

terminal Toprim domain, which is believed to be important in DNA replication, recombination,

and repair[24,25]. SparE is homologous to GIY-YIG endonucleas[26-30] and PcrA/UvrD helicase. The

PcrA/UvrD helicase is an essential helicase in gram-positive bacteria and has been shown to play

a role in DNA repair, as well as the replication of small drug-resistance plasmids in

Staphylococcus aureus and *Bacillus subtilis*[31-37]. This raises the possibility that the *SPAR* locus

may be a remnant of a plasmid or mobile genetic element that had been integrated into the

chromosome.

208   In summary, we have identified a chromosomal locus in *Sgg*, *SPAR*, that is vital to the

pathogenicity of *Sgg*. We report that deletion of *SPAR* significantly reduces the capacity of *Sgg*

to adhere to CRC cells and to colonize the gut. Further, *SPAR* deletion abrogates the ability for

Sgg to stimulate CRC cell proliferation and to accelerate colon tumor development. Examination

of genes with the locus provided potential candidates responsible for the observed phenotypes

and suggested a connection between the *SPAR* locus and T7SS in terms of additional T7SS

effectors and a possible regulator of T7SS expression. Further investigation into the activities of

specific *SPAR* proteins will be important for unveiling the intricate *Sgg* virulence mechanisms

and will open up the path ahead to identify biomarker candidates and targets for clinical

prevention and intervention strategies.

Materials and Methods

Bacterial strains, cell lines and growth conditions *Sgg* strains were routinely grown in

brain heart infusion (BHI) broth or tryptic soy broth with shaking, or on BHI or TSB agar plates

at 37 ˚C overnight (Teknova). For co-culture experiments and animal studies, stationary phase

bacterial cultures were pelleted, washed with phosphate-buffered saline (PBS), pH 7.4,

resuspended in PBS containing 15% glycerol, aliquoted and stored at -80 ˚C. Aliquoted stocks

were thawed and washed with PBS, and diluted in appropriate media to obtain the desired

concentration. Human colon cancer cell lines HT29, HCT116, and SW480, as well as the

HEK293 cell line (ATCC), were grown in Dulbecco's Modified Eagle's Medium F-12 50/50

(DMEM/F-12, GIBCO) supplemented with 10% FBS at 37 ˚C with 5% CO_2 in a humidified

chamber. Cells from less than 30 passages were used in the experiments.

Adherence assay. This was performed as described previously[3]. Briefly, HT29 cells

were seeded at a density of 1×10^6 cells/well in a 24 well plate and allowed to attach overnight.

Cells were incubated with bacteria at a multiplicity of infection (MOI) of 10 for 1 hour at 37 ˚C

with 5% CO_2. Cells were then washed with PBS, lysed with PBS containing 0.025% Triton-X

100 and dilution plated. Adherence was calculated as the percentage of adherent bacteria vs. total

bacteria added.

Cell proliferation assay. This was performed as described previously[3] with slight

modifications. Briefly, cells were seeded in 96 well plates at a concentration of ~1×10^4 cells/well

and incubated overnight. Cells were then incubated in fresh DMEM/F-12 with 10% FBS and *Sgg*

bacteria (MOI = 1) for a total of 24 hours. Trimethoprim was added after 6 hours of incubation

(1 μg/mL final concentration) to inhibit bacterial growth, as previously described. The number of

viable cells was determined using the cell counting kit (CCK)-8 kit following the instructions of

the supplier (Apex Bio).

Western Blot. Detection of β-catenin and PCNA was carried out as described

previously[3]. Briefly, total cell lysates from HT29 cells cultured in the presence or absence of

bacteria were subjected to SDS-PAGE, transferred, and probed with antibodies against β-catenin

(1:1000), PCNA (1:1000), and β-actin (1:1000) (Cell Signaling Technology), followed by

incubation with HRP-conjugated secondary antibodies (1:3000). Band intensities were measured

using FIJI (ImageJ) and normalized to those of β-actin.

Deletion of *SPAR*. This was performed as described previously[3]. Briefly, the ~ 1kb

region upstream of *SPAR_A* and the ~1kb region downstream of *SPAR_L* were synthesized by

Genscript and cloned into pUC57 (Genscript). The insert was then subcloned into a temperature

sensitive conjugative plasmid pG1-*oriT*TnGBS1. The insert sequence was verified by Sanger

sequencing. The construct was introduced into *S. agalactiae* NEM316 by electroporation and

then into *Sgg* strain TX20005 by conjugation under the permissive temperature. Erythromycin-

resistant transconjugants were then cultured under a non-permissive temperature to select for

single cross-over recombinants, followed by serial passages in antibiotic-free BHI and screening

for double cross-over deletion mutants by PCR. Deletion was confirmed by PCR amplification of

the regions spanning the deleted fragment and DNA sequencing of the PCR product.

RNA extraction, cDNA synthesis and PCR. This was performed following the method

described previously [Taylor]. Primers used in PCR amplification are: *sparA*, forward

5'GCAAGCTGGTCGAACAGAAC and reverse 5'GCTTCTATGGTTGGGGCTAGA; *sparD*,

forward 5' GGAGGTGGATCCAACAAGGG and reverse 5'CAGGTTCCTCGATAGCCAGC;

and *sparG*, forward 5'TCAGTTGTTAGCGGATGCGT and reverse 5'

CCCTTTATTGCTTGTGCTCCC.

Growth curve. Overnight cultures of *Sgg* were inoculated into fresh BHI broth at 1:100

dilution and grown at 37 ˚C with shaking. Samples were taken at 0, 3, 6, 9, 12, and 24 hours,

dilution plated, incubated for 24 hours and colonies enumerated.

Preparation of culture supernatants. Supernatants from stationary phase Sgg cultured

in BIH were filtered-sterilized through a 0.2μm filter and concentrated approximately 10 – 20

fold using centrifugal concentrators (3kD molecular weight cut off). The concentrated CSs were

aliquoted and stored at -80°C. A vial is thawed and diluted in the appropriate media for

adherence or cell proliferation assays.

Animal experiments. Animal studies were performed in accordance with protocols

approved by the Institutional Animal Care and Use Committee at the Texas A&M Health

Science Center, Institute of Biosciences and Technology. Mice were fed with standard ProLab

IsoPro RMH3000 (LabDiet). **Colonization experiments.** This was performed as previously

described[3] with slight modifications. Briefly, 6-week old A/J mice, sex matched (Jackson

Laboratory), were treated with ampicillin at a concentration of 1 g/L in drinking water for 3 days

and switched to antibiotic-free water 24 hours prior to administration of bacteria. *Sgg* was orally

gavaged at a dose of ~1 x 10^9 CFU/mouse. Colons and fecal materials were collected at day 1, 3,

and 7 post-gavage. Samples were weighed, homogenized in PBS in a TissueLyser (Qiagen),

dilution plated onto Enterococcus Selective Agar (ESA) plates and incubated at 37 ˚C for 24–48

hours to enumerate *Sgg* colonies, as previously described. **AOM-induced model of CRC.** This

was performed as previously described[3] with slight modifications. Briefly, A/J mice were treated

with 4 weekly i.p. injections of AOM (10 mg/kg body weight), followed by ampicillin in

drinking water (1 g/L) for 7 days and switched to antibiotic-free water 24 hours prior to the first

oral gavage with *Sgg*. Mice were orally gavaged with bacteria at $\sim 1\text{x}10^9$ CFU/mouse or saline

once per week for 12 weeks. Colons of mice were collected and cut opened longitudinally. The

number and size of macroscopic were recorded. Tumor burden was calculated as the sum of all

tumor volumes per mouse. Visual evaluation of colons was carried out by a blinded observer.

290   **Statistical analysis**. GraphPad Prism 8 was used for statistical analyses. Two-tailed

unpaired *t*-test was used for pairwise comparisons to assess the significance of differences

between two groups in cell proliferation assays, western blot analysis, and adherence assays. The

non-parametric Mann-Whitney test was used to assess the significance of differences in the

bacterial load and tumor burden in animal studies. Two-way ANOVA was used to analyze the

bacterial growth curves. Ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p <$

0.0001.

Acknowledgements

This study was supported by funding from the Hamill Foundation, the Cancer Prevention

Research Institute of Texas and NIAID.

References

1   Schlegel, L., Grimont, F., Ageron, E., Grimont, P. A. D. & Bouvet, A. Reappraisal of the

taxonomy of the Streptococcus bovis/Streptococcus equinus complex and related species:

description of Streptococcus gallolyticus subsp. gallolyticus subsp. nov., S. gallolyticus

subsp. macedonicus subsp. nov. and S. gallolyticus subsp. pasteurianus subsp. nov. *Int J*

Syst Evol Microbiol **53**, 631-645, doi:10.1099/ijs.0.02361-0 (2003).

2   Kumar, R., Herold, J. L., Taylor, J., Xu, J. & Xu, Y. Variations among Streptococcus

gallolyticus subsp. gallolyticus strains in connection with colorectal cancer. *Scientific*

reports **8**, 1514, doi:10.1038/s41598-018-19941-7 (2018).

3   Taylor, J. C. *et al*. A type VII secretion system of Streptococcus gallolyticus subsp.

gallolyticus contributes to gut colonization and the development of colon tumors. *PLoS*

Pathog **17**, e1009182, doi:10.1371/journal.ppat.1009182 (2021).

4   Kumar, R. *et al*. Streptococcus gallolyticus subsp. gallolyticus promotes colorectal tumor

development. *PLoS Pathog* **13**, e1006440, doi:10.1371/journal.ppat.1006440 (2017).

5   Marmolin, E. S. *et al*. Bacteremia with the bovis group streptococci: species

identification and association with infective endocarditis and with gastrointestinal

disease. *Diagn Microbiol Infect Dis* **85**, 239-242,

doi:10.1016/j.diagmicrobio.2016.02.019 (2016).

6   Corredoira, J. *et al*. Differences between endocarditis caused by Streptococcus bovis and

Enterococcus spp. and their association with colorectal cancer. *Eur J Clin Microbiol*

Infect Dis **34**, 1657-1665, doi:10.1007/s10096-015-2402-1 (2015).

7   Boleij, A. & Tjalsma, H. Gut bacteria in health and disease: a survey on the interface

322   between intestinal microbiology and colorectal cancer. *Biol Rev Camb Philos Soc* **87**,

323   701-730, doi:10.1111/j.1469-185X.2012.00218.x (2012).

324   8   Boleij, A., van Gelder, M. M., Swinkels, D. W. & Tjalsma, H. Clinical Importance of

325   Streptococcus gallolyticus infection among colorectal cancer patients: systematic review

326   and meta-analysis. *Clin Infect Dis* **53**, 870-878, doi:10.1093/cid/cir609

327 cir609 [pii] (2011).

328   9   Abdulamir, A. S., Hafidh, R. R. & Abu Bakar, F. The association of Streptococcus

329   bovis/gallolyticus with colorectal tumors: the nature and the underlying mechanisms of

330   its etiological role. *J Exp Clin Cancer Res* **30**, 11, doi:10.1186/1756-9966-30-11 (2011).

331   10   Gupta, A., Madani, R. & Mukhtar, H. Streptococcus bovis endocarditis, a silent sign for

332   colonic tumour. *Colorectal Dis* **12**, 164-171, doi:CDI1814 [pii]

333 10.1111/j.1463-1318.2009.01814.x (2010).

334   11   Waisberg, J., Matheus, C. e. O. & Pimenta, J. Infectious endocarditis from Streptococcus

335   bovis associated with colonic carcinoma: case report and literature review. *Arq*

336   *Gastroenterol* **39**, 177-180, doi:10.1590/s0004-28032002000300008 (2002).

337   12   Alazmi, W., Bustamante, M., O'Loughlin, C., Gonzalez, J. & Raskin, J. B. The

338   association of Streptococcus bovis bacteremia and gastrointestinal diseases: a

339   retrospective analysis. *Dig Dis Sci* **51**, 732-736, doi:10.1007/s10620-006-3199-7 (2006).

340   13   Gold, J. S., Bayar, S. & Salem, R. R. Association of Streptococcus bovis bacteremia with

341   colonic neoplasia and extracolonic malignancy. *Arch Surg* **139**, 760-765,

342   doi:10.1001/archsurg.139.7.760 (2004).

14 Corredoira, J. *et al*. The clinical epidemiology and malignancies associated with

Streptococcus bovis biotypes in 506 cases of bloodstream infections. *J Infect* **71**, 317-

325, doi:10.1016/j.jinf.2015.05.005 (2015).

15 Zhang, Y., Weng, Y., Gan, H., Zhao, X. & Zhi, F. Streptococcus gallolyticus conspires

myeloid cells to promote tumorigenesis of inflammatory bowel disease. *Biochem Biophys*

Res Commun **506**, 907-911, doi:10.1016/j.bbrc.2018.10.136 (2018).

16 Kwong, T. N. Y. *et al*. Association Between Bacteremia From Specific Microbes and

Subsequent Diagnosis of Colorectal Cancer. *Gastroenterology* **155**, 383-390.e388,

doi:10.1053/j.gastro.2018.04.028 (2018).

17 Thomas, A. M. *et al*. Metagenomic analysis of colorectal cancer datasets identifies cross-

cohort microbial diagnostic signatures and a link with choline degradation. *Nat Med* **25**,

667-678, doi:10.1038/s41591-019-0405-7 (2019).

18 Darling, A. C., Mau, B., Blattner, F. R. & Perna, N. T. Mauve: multiple alignment of

conserved genomic sequence with rearrangements. *Genome Res* **14**, 1394-1403,

doi:10.1101/gr.2289704 (2004).

19 Corredoira, J. *et al*. Colorectal neoplasm in cases of Clostridium septicum and

Streptococcus gallolyticus subsp. gallolyticus bacteraemia. *Eur J Intern Med* **41**, 68-73,

doi:10.1016/j.ejim.2017.02.009 (2017).

20 Poulsen, C., Panjikar, S., Holton, S. J., Wilmanns, M. & Song, Y. H. WXG100 protein

superfamily consists of three subfamilies and exhibits an α-helical C-terminal conserved

residue pattern. *PLoS One* **9**, e89313, doi:10.1371/journal.pone.0089313 (2014).

21 Liu, G. F., Wang, X. X., Su, H. Z. & Lu, G. T. Progress on the GntR family transcription

regulators in bacteria. *Yi Chuan* **43**, 66-73, doi:10.16288/j.yczz.20-245 (2021).

22   Chatterjee, A., Willett, J. L. E., Dunny, G. M. & Duerkop, B. A. Phage infection and sub-

lethal antibiotic exposure mediate Enterococcus faecalis type VII secretion system

dependent inhibition of bystander bacteria. *PLoS Genet* **17**, e1009204,

doi:10.1371/journal.pgen.1009204 (2021).

23   Schiltz, C. J., Lee, A., Partlow, E. A., Hosford, C. J. & Chappie, J. S. Structural

characterization of Class 2 OLD family nucleases supports a two-metal catalysis

mechanism for cleavage. *Nucleic Acids Res* **47**, 9448-9463, doi:10.1093/nar/gkz703

(2019).

24   Aravind, L., Leipe, D. D. & Koonin, E. V. Toprim--a conserved catalytic domain in type

IA and II topoisomerases, DnaG-type primases, OLD family nucleases and RecR

proteins. *Nucleic Acids Res* **26**, 4205-4213, doi:10.1093/nar/26.18.4205 (1998).

25   Koonin, E. V. & Gorbalenya, A. E. The superfamily of UvrA-related ATPases includes

three more subunits of putative ATP-dependent nucleases. *Protein Seq Data Anal* **5**, 43-

45 (1992).

26   Kowalski, J. C. *et al*. Configuration of the catalytic GIY-YIG domain of intron

endonuclease I-TevI: coincidence of computational and molecular findings. *Nucleic*

Acids Res **27**, 2115-2125, doi:10.1093/nar/27.10.2115 (1999).

27   Van Roey, P., Meehan, L., Kowalski, J. C., Belfort, M. & Derbyshire, V. Catalytic

domain structure and hypothesis for function of GIY-YIG intron endonuclease I-TevI.

Nat Struct Biol **9**, 806-811, doi:10.1038/nsb853 (2002).

28   Truglio, J. J. *et al*. Structural insights into the first incision reaction during nucleotide

excision repair. *EMBO J* **24**, 885-894, doi:10.1038/sj.emboj.7600568 (2005).

29   Dunin-Horkawicz, S., Feder, M. & Bujnicki, J. M. Phylogenomic analysis of the GIY-

YIG nuclease superfamily. *BMC Genomics* **7**, 98, doi:10.1186/1471-2164-7-98 (2006).

30   Ibryashkina, E. M., Sasnauskas, G., Solonin, A. S., Zakharova, M. V. & Siksnys, V.

Oligomeric structure diversity within the GIY-YIG nuclease family. *J Mol Biol* **387**, 10-

16, doi:10.1016/j.jmb.2009.01.048 (2009).

31   Bruand, C. & Ehrlich, S. D. UvrD-dependent replication of rolling-circle plasmids in

Escherichia coli. *Mol Microbiol* **35**, 204-210, doi:10.1046/j.1365-2958.2000.01700.x

(2000).

32   Chang, T. L. *et al*. Biochemical characterization of the Staphylococcus aureus PcrA

helicase and its role in plasmid rolling circle replication. *J Biol Chem* **277**, 45880-45886,

doi:10.1074/jbc.M207383200 (2002).

33   del Solar, G., Giraldo, R., Ruiz-Echevarría, M. J., Espinosa, M. & Díaz-Orejas, R.

Replication and control of circular bacterial plasmids. *Microbiol Mol Biol Rev* **62**, 434-

464, doi:10.1128/MMBR.62.2.434-464.1998 (1998).

34   Naqvi, A., Tinsley, E. & Khan, S. A. Purification and characterization of the PcrA

helicase of Bacillus anthracis. *J Bacteriol* **185**, 6633-6639, doi:10.1128/JB.185.22.6633-

6639.2003 (2003).

35   Khan, S. A. Rolling-circle replication of bacterial plasmids. *Microbiol Mol Biol Rev* **61**,

442-455, doi:10.1128/mmbr.61.4.442-455.1997 (1997).

36   Khan, S. A. Plasmid rolling-circle replication: recent developments. *Mol Microbiol* **37**,

477-484, doi:10.1046/j.1365-2958.2000.02001.x (2000).

37   Soultanas, P. *et al*. Plasmid replication initiator protein RepD increases the processivity

of PcrA DNA helicase. *Nucleic Acids Res* **27**, 1421-1428, doi:10.1093/nar/27.6.1421

(1999).

413   **Tables**

414   **Table 1. Features of predicted SPAR Proteins.**

Gene	Length (aa)	Predicted Gene Function
sparA	424	Hypothetical
sparB	53	Hypothetical
sparC	582	Hypothetical
sparD	656	OLD-Family Endonuclease
sparE	609	GIY-YIG Endonuclease/PcrA/UvrD Helicase
sparF	232	GntR-Family Transcriptional Regulator
sparG	91	Hypothetical, potential T7SS effector
sparH	102	Hypothetical, potential T7SS effector
sparI	137	Hypothetical, potential T7SS effector
sparJ	417	Hypothetical, potential T7SS effector
sparK	199	Hypothetical, potential T7SS effector
sparL	197	Hypothetical

Fig 1. *SPAR* **Genetic Organization and Expression. A.** The *SPAR* locus. Genes encoding

proteins with homology to proteins of known functions or containing characteristics of certain

protein families are colored in blue, while genes encoding hypothetical proteins of unknown

function are colored in gray. **B. RT-PCR.** RNA was extracted from TX20005 cultured in the

presence of HT29, as well as from the colon tissues of mice that had been orally gavaged with

TX20005. RNA was treated with DNase and synthesized using a ProtoScript II First Strand

cDNA Synthesis Kit (NEB). Newly synthesized cDNA was then used as PCR template to detect

sparA, D, and *G*.

Fig 2. **SPAR is important for *Sgg* adherence but does not impact growth. A. Adherence**.

HT29 cells were seeded at a density of $1x10^6$ cells per well and incubated with TX20005 or

TX20005Δ*SPAR* (MOI=10) in media only or in the presence of CS from TX20005 or

TX20005Δ*SPAR* for 1 hour. Cells were washed thrice in PBS to remove unattached bacteria,

then lysed with 1mL of 0.01% Triton X-100 (Sigma). The lysate was collected, serially diluted,

and plated on tryptic soy agar (TSA) plates (Sigma) to be incubated for 24 hours. Colonies were

enumerated and percentage adherence was calculated as the percentage of adhered bacteria vs

total bacteria added. Results were combined from three biological replicates. Unpaired two-tailed

t tests were used. **, $p<0.01$; NS, not significant. **B. Growth Curve**. Single colonies of either

bacteria were inoculated into 1mL of BHI (Teknova) and incubated at 37℃ overnight with

shaking. The overnight culture was then diluted 1:100 with fresh BHI and incubated at 37℃ with

shaking. 100μL of the culture were taken at the 0, 3, 6, 9, 12, and 24 hours for serial dilution and

platin. Results were combined from three biological replicates.

Fig 3. *SPAR* **is essential for** *Sgg* **to stimulate CRC cell proliferation. A-D**. **Cell Proliferation**

assay. HT29 (**A**), HCT116 (**B**), SW480 (**C**), and HEK293 (**D**) cells were seeded at a density of

1×10^4 cells per well and co-cultured in a 37ºC humidified incubator with 5% CO_2 for 24 hours in

the presence or absence of bacteria (MOI=1), as described in the Materials and Methods section.

Cell proliferation was measured using a CCK-8 kit. Cell-free wells containing only the culture

media were served as a baseline control to which the raw values were normalized. **E-G. Western**

blot. HT29 cells were seeded at a density of 1×10^6 cells per well, and co-cultured in the presence

or absence of bacteria (MOI=1) for 9 hours. Cell lysates were analyzed by western blot, probed

with antibodies against β-catenin, PCNA, and β-actin (**E**). Band intensity was quantified and

normalized to β-actin. Fold change in β-catenin (**F**) and PCNA (**G**) were calculated against

untreated cells. Data was combined from three biological replicates. **H. Cell proliferation in the**

presence of *Sgg* **CS.** HT29 cells were cultured in the presence or absence of bacteria (MOI=1)

or with CSs from WT or mutant *Sgg* for 24 hours. Cell proliferation was measured by using a

CCK-8 kit. Unpaired two-tailed *t* tests were used for all comparisons. *, $p<0.05$; **, $p<0.01$; ***,

$p<0.001$; ****, $p<0.0001$; ns, not significant.

Fig 4. *SPAR* **deletion reduced the colonization capacity of** *Sgg*. This was performed as

described in the Materials and Methods section. Colon tissues (**A**) and fecal pellets (**B**) were

weighed, homogenized in sterile PBS and dilution plated onto selective agar plates. The non-

parametric Mann-Whitney test was used for the comparisons. *, $p<0.05$; **, $p<0.01$; ****,

$p<0.0001$; NS, not significant.

Fig 5. *SPAR* is important for *Sgg* to promote the development of colon tumors. This was

performed as described in the Materials and Methods section. The procedure is illustrated in **A**.

B-D. Macroscopic tumors were observed by blinded observers. Tumor incidence (**B**) is shown as

the ratio of tumor-bearing mice vs. total mice, presented as a fraction of the total. Tumor number

per mouse and average tumor size are shown in **C** and **D**. Tumor burden (**E**) is calculated as sum

of tumor volumes per mouse. **F.** *Sgg* load in the fecal pellets collected at the experimental

endpoint was determined by dilution plating onto selective agar plates. Mann-Whitney test was

used for all comparisons. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, p, <0.0001; NS, not

significant.

Fig 1

A

SPAR-1   SPAR-2   SPAR-3

A B C — Hypothetical

D OLD Family Endonuclease

E GIY-YIG PrcA/UvrD Helicase

F GntR Family Transcriptional Regulator

G Hypothetical

H I J K — LXG

L Hypothetical

B

Fig 2

Fig 3

Fig 4

Fig 5

Fig S1

```
sparH       GETDEALAELKKQEDDLEDALVDLLRKEAQDEH
sparI       DQQQQCLNVLEEAELQLSLISVMSLERKTSLHQ
sparJ       EVKTTIFSSLVPKEVVAINPVTGAPIYASVSEDEAMAYLQENYEGSYLFIQSLEEEENLFGEDN
sparK       GDDLLTQLSAYGKDLDWLQTTSQTVLEDDILGLWFKKGSHRYSLNNLGDLTITYDDVLNDN
Consensus            HxxxDxxhxxxH
                         E
```

Fig S1: The conserved T7SS sequence in SparH-K. H denotes highly conserved and h less conserved hydrophobic residues