Defining the genes required for survival of *Mycobacterium bovis* in the bovine host offers

novel insights into the genetic basis of survival of pathogenic mycobacteria

Amanda J Gibson[1,5&*], Jennifer Stiens[2*], Ian J Passmore[3], Valwynne Faulkner[1,6&], Josephous

Miculob[1], Sam Willcocks[3], Michael Coad[4], Stefan Berg[4], Dirk Werling[1], Brendan W Wren[3],

Irene Nobeli[2], Bernardo Villarreal-Ramos[4,5&], Sharon L Kendall[1**]

[1]Centre for Emerging, Endemic and Exotic Diseases, Pathobiology and Population Sciences,

Royal Veterinary College, Hawkshead Lane, North Mymms, Hatfield, AL9 7TA, United

Kingdom.

[2]Institute of Structural and Molecular Biology, Biological Sciences, Birkbeck, University of

London, Malet Street, London, WC1E 7HX, United Kingdom.

[3]London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, United

Kingdom.

[4]Animal and Plant Health Agency, Woodham Ln, Addlestone, Surrey, KT15 3NB, United

Kingdom.

[5]Centre of Excellence for Bovine Tuberculosis, IBERS, Aberystwyth University, Penglais,

Aberystwyth, Ceredigion, SY23 3EE, United Kingdom.

[6]Systems Chemical Biology of Infection and Resistance Laboratory, The Francis Crick

Institute, 1 Midland Road, London 15 NW1 1AT, United Kingdom.

[&]current address

*the authors contributed equally to the work

**corresponding author

Abstract

Tuberculosis has severe impacts in both humans and animals. Understanding the genetic

basis of survival of both *Mycobacterium tuberculosis*, the human adapted species, and

Mycobacterium bovis, the animal adapted species is crucial to deciphering the biology of

both pathogens. There are several studies that identify the genes required for survival of *M.*

tuberculosis in vivo using mouse models, however, there are currently no studies probing

the genetic basis of survival of *M. bovis in vivo.* In this study we utilise transposon insertion

sequencing in *M. bovis* to determine the genes required for survival in cattle. We identify

genes encoding established mycobacterial virulence functions such as the ESX-1 secretion

system, PDIM synthesis, mycobactin synthesis and cholesterol catabolism that are required

in vivo. We show that, as in *M. tuberculosis, phoPR* is required by *M. bovis in vivo* despite

the known defect in signalling through this system. Comparison to studies performed in

glycerol adapted species such as *M. bovis* BCG and *M. tuberculosis* suggests that there are

differences in the requirement genes involved in cholesterol import (*mce4* operon),

oxidation (*hsd*) and detoxification (*cyp125*). We report good correlation with existing

mycobacterial virulence functions, but also find several novel virulence factors, including

genes involved in protein mannosylation, aspartate metabolism and glycerol-phosphate

metabolism. These findings further extend our knowledge of the genetic basis of survival *in*

vivo in bacteria that cause tuberculosis and provide insight for the development of novel

diagnostics and therapeutics.

Importance

This is the first report of the genetic requirements of an animal adapted member of the MTBC in a natural host. *M. bovis* has devastating impacts in cattle and bovine tuberculosis is a considerable economic, animal welfare and public health concern. The data highlight the importance of mycobacterial cholesterol catabolism and identifies several new virulence factors. Additionally, the work informs the development of novel differential diagnostics and therapeutics for TB in both human and animal populations.

Introduction

Bacteria belonging to the *Mycobacterium tuberculosis* complex (MTBC) have devastating impacts in both animal and human populations. *Mycobacterium bovis,* an animal adapted member of the MTBC and one of the main causative agents of bovine tuberculosis (bTB), remains endemic in some high-income settings despite the implementation of a test and slaughter policy. In low- and middle-income settings, the presence of bTB in livestock combined with the absence of rigorous control measures contributes to the risk of zoonotic transmission (1, 2). Control measures based on cattle vaccination utilise the live attenuated vaccine *M. bovis* BCG but the efficacy of this vaccine still remains low in field situations (3, 4). In addition to vaccines, the development of diagnostic tools for the identification of infected individuals is crucial for the management of transmission. Currently, vaccination with *M. bovis* BCG sensitises animals to the diagnostic tuberculin skin test, therefore, sensitive and specific differentiating diagnostic strategies are a current imperative (5, 6).

The increased accessibility of whole genome fitness screens has allowed the

assessment of the impacts of the loss of gene function on bacterial survival (7). Such screens

have been invaluable in identifying novel drug targets or candidates for the generation of

new live attenuated vaccines in a number of bacterial pathogens, including *Mycobacterium*

tuberculosis (8–13). Studies utilising whole genome transposon mutagenesis screens to

examine gene fitness *in vivo* in *M. tuberculosis* have been limited to mouse models (8, 9,

13). These models do not faithfully replicate the granulomatous pathology associated with

TB, nor do mice contain the same repertoire of CD1 molecules expressed by bovine T cells

required to present mycobacterial lipid antigens (14). Whole genome transposon

mutagenesis screens utilising non-human primates are limited because screening is

restricted to smaller mutant pools (15). To date, Tn-seq based studies in the context of bTB

in cattle have only been performed using *M. bovis* BCG strains (16, 17).

In this study we use transposon insertion sequencing (Tn-seq) to determine the

genes required for survival of *M. bovis* directly in cattle. We show that genes involved in the

biosynthesis of phthiocerol dimycocerosates (PDIMs), the ESX-1 secretion system,

cholesterol catabolism, and mycobactin biosynthesis are essential for survival *in vivo* in

cattle, corroborating current knowledge of gene essentiality in members of the MTBC (8, 9,

13, 16, 17). We identify differences in the requirement for genes involved in cholesterol

transport and oxidation in the fully virulent *M. bovis* strain. We also identify several novel

genes required for survival *in vivo* that have not been previously described in members of

the MTBC.

Results and Discussion

Generation and sequencing of the input library.

We generated a transposon library in *M. bovis* AF2122/97 using the MycomarT7 phagemid system as previously described (18, 19). Sequencing of the input library showed that transposon insertions were evenly distributed around the genome and 27,751 of the possible 73,536 thymine–adenine dinucleotide (TA) sites contained an insertion representing a saturation density of ~38% (Supplementary Figure S1 and Supplementary Table S1 -input library). The *M. bovis* AF2122/97 genome has 3,989 coding sequences and insertions were obtained in 3,319 of these, therefore the input library contained insertions in 83% of the total coding sequences.

Mycobacterium bovis **specific immune responses were observed in cattle**

Twenty-four clinically healthy calves of approximately 6 months of age were inoculated with the library through the endobronchial route. Infection was monitored by IFN-γ release assay (IGRA) at the time of inoculation and 2 weeks post infection. *M. bovis* specific immune responses were observed for all study animals at 2 weeks post infection (Figure 1A and B). Each animal presented very low background of circulating IFN-γ together with a statistically significant increase in IFN-γ release in response to PPD-B compared to PPD-A antigens (Figure 1C; *** $p \leq 0.001$). This indicates that infection with the library was successfully established in the cattle.

Pathology associated with infection was greater in the lung and thoracic lymph nodes

Animals were culled at 6 weeks post infection. Lung sections and upper (head and neck) and

lower (thoracic) respiratory tract associated lymph nodes were examined for gross lesions.

Lesions typical of *M. bovis* infection were observed in the tissues examined. Pathology

scores are shown in Figure 2A. Greater pathology was observed in lung and thoracic lymph

nodes compared to the head and neck lymph nodes.

Higher bacterial loads were associated with the lung and thoracic lymph nodes

Bacterial counts were highest in lesions derived from the lung compared to those from the

thoracic lymph nodes and head and neck lymph nodes (Figure 2B). The lowest bacterial

counts were observed within the head and neck lymph nodes. However, this was not

significant when compared to thoracic lymph nodes. The volume of each macerate varied

depending on lesion size. Considering macerate volume, average bacterial loads of 10^7, 10^6

and 10^5 were recovered from lesions from samples of the lungs, thoracic lymph nodes and

head and neck lymph nodes, respectively.

Recovery and sequencing of *in vivo* selected transposon libraries

In order to recover the Tn library from harvested tissue ~ 10^5 - 10^6 CFU from lungs and

thoracic lymph nodes were plated onto several 140 mm modified 7H11 plates containing

kanamycin to minimize competition between mutants. Samples from 4 cattle were lost due

to fungal contamination, therefore the samples processed represent samples from 20 cattle.

Lung samples were plated from all 20 animals and thoracic lymph nodes samples were

plated from 6 cattle. Bacteria were grown for 4-6 weeks before harvesting for genomic DNA

extraction and subsequent sequencing (see Supplementary Table S1 for assignation of

sequencing files to samples). The diversities of the output libraries were compared to the

input library for each sample (Supplementary Figure S2 and Table S1). On average, libraries

recovered from lung lesions from 20 different cattle contained 14,456 unique mutants and

those recovered from the thoracic lymph nodes contained an average of 16,210 unique

mutants. Given that the input library contained 27,751 unique mutants this represented a

loss of diversity of ~ 40-50%. Good coverage of coding sequences (CDSs) was maintained as

the output libraries still contained insertions in (on average) 68-70% of the open reading

frames.

Comparison of the read counts between the input and output libraries allowed a

measurement of the impact of the insertion on the survival of mutants in cattle. The results

are represented as a mean $\log_2$ fold-change in the output compared with the input for each

gene. The entire dataset is shown in supplementary Table S4 and a volcano plot from the

lungs and thoracic lymph node of two representative animals is shown in Supplementary

Figure S3. Comparison of the mean $\log_2$ fold-change between lung and lymph node samples

showed good correlation (Spearman's rho = 0.88, p-value <2.2e-16) (Supplementary Figure

S4). TRANSIT resampling was performed to compare the composition of the mutant

population in the lungs and thoracic lymph nodes of paired cattle, it was also applied to

compare all the thoracic lymph nodes with the lungs of all cattle samples. No statistically

significant differences were observed indicating that there were no differences in mutant

composition between the tissue sites.

No insertion mutants were significantly over-represented in the output library in any

of the animals. Although interestingly, insertions in *MB0025,* a gene that is unique to *M.*

bovis appeared to improve growth in cattle as mean $\log_2$ fold-changes of +3.9 (lungs) and

+4.2 (lymph nodes) were observed; however, significance criteria were not met in any of the

animals.

In order to define a list of attenuating mutations, we used a similar approach to that

used in a previous study with an *M. bovis* BCG library in cattle (16). Insertions in genes were

defined as attenuating if they had $\log_2$ fold-change of -1.5 or below and an adjusted p-value

of <0.05 in at least half of the animals (Table S4, significant in 50% of cattle tab). When using

these criteria, there were 141 genes where insertions caused significant attenuation in the

lungs or the thoracic lymph nodes, 20 genes that reached significance only in the lungs

(shown in red) and 16 genes that reached significance only in the thoracic lymph nodes

(shown in green). Of the 141 genes, 109 had been previously described as being required *in*

vivo in *M. tuberculosis* H37Rv in mouse models through the use of whole genome Tn

screens representing ~77% overlap with the previous literature (8, 9, 13).

Comparison with mutations known to cause attenuation in the MTBC

Insertions in the RD1 encoded ESX-1 type VII secretion system secreting virulence

factors and immunodominant antigens EsxA (CFP-10) and EsxB (ESAT-6) are expected to

cause attenuation (20). The impacts of insertions in this region are summarised in Figure 3

but are also available in Supplementary Table S4 (RD regions tab) and Supplementary Figure

S5. Insertions in genes encoding the structural components of the apparatus (*eccB1, eccCa1,*

eccCb1, eccD1) were severely attenuating ($\log_2$ fold-change -6 to -9). Insertions in *eccA1,*

which also codes for a structural component of the apparatus, were less impactful ($\log_2$ fold-

changes of -2 to -3) despite good insertion saturation in this gene. This is supported by the

work of others who have shown that deletion of *eccA1* in *Mycobacterium marinum* leads to

only a partial secretion defect (21). There were no impacts seen due to insertions in

accessory genes *espJ, espK* and *espH*. The lack of attenuation seen in *espK* mutants is

supported by other studies showing that this gene is dispensable for secretion through the

apparatus and is not required for virulence of *M. bovis* in guinea pigs (22, 23). Insertions in

esxA and *esxB* resulted in severe attenuation (log$_2$ fold-change of -6) but this did not reach

significance cut-offs (adj. p=<0.05) in any of the cattle. This is likely to be due to the small

number of TAs in these genes which makes it challenging to measure mutant frequency.

The highest levels of attenuation seen were in genes involved in the synthesis of the

cell wall virulence lipids PDIMs (*ppsABCDE* and *mas* with log$_2$ fold-changes of ~-10

commonly seen). PDIM synthesis is well known to be required for the survival of *M.*

tuberculosis and *M. bovis* in mice and guinea pigs (24, 25). Insertions in genes involved in

the synthesis of PDIMs were the most under-represented (log$_2$ fold-changes of -8 to -10) in

the output library (Figure 4, Supplementary Table S4, mycolipids tab). MmpL7 is involved in

PDIM transport and there is evidence that it is phosphorylated by the serine-threonine

kinase PknD (26). PknD-MmpL7 interactions are thought to be perturbed in *M. bovis* as *pknD*

is split into two coding sequences in the bovine pathogen by a frameshift mutation (27). The

data presented here suggest that MmpL7 still functions despite the frameshift mutation.

Iron restriction is thought to be a mechanism by which the host responds to

mycobacterial infection, although different cellular compartments may be more restrictive

than others (28). Insertion in many of the genes involved in mycobactin synthesis *(Mb2406-*

Mb2398, mbtJ-mbtH) were attenuating in cattle (Figure 4, Supplementary Table S4,

mycobactin synthesis tab). As mycobactin is required for the acquisition of iron, this

confirms that, like other members of the MTBC, needs to scavenge iron from the host for

survival (13, 16).

The role of the cholesterol catabolism in *M. tuberculosis* is well documented and it is

required for both energy generation and manipulation of the immune response (29–31).

Cholesterol uptake is mediated by the Mce4 transporter coded by the *mce4* operon

Rv3492c-Rv3501c (*Mb3522c-MB3531c*) (32, 33). It has been suggested that an alternative

cholesterol acquisition pathway operates in *M. bovis* BCG Danish as, unlike insertions in

genes in the down-stream catabolic pathway, insertions in the *mce4* operon do not result in

attenuation in this strain (16). In contrast, our study shows that cholesterol transport via the

Mce4 transporter is required in *M. bovis* (Figure 4, Supplementary Table S4 -cholesterol

catabolism tab, Figure 5). This corroborates work performed in *M. tuberculosis*, where Mce4

has been shown to be required for growth in chronically infected mice (9, 32). Propionyl-coA

generated from the catabolism of cholesterol is toxic and detoxification mechanisms include

incorporation into PDIMs (34, 35). The observation that BCG Danish contains a lower

amount of PDIMs compared to BCG Pasteur (16) suggests a correlation between Mce4

mediated cholesterol transport and PDIM synthesis and previous studies have

demonstrated an increase in PDIM biosynthesis as a result of *mce4* over-expression (36).

PDIMs biosynthesis genes are over-expressed in *M. bovis* compared to *M. tuberculosis* (27)

and comparison of our dataset with Tn-seq studies performed in *M. tuberculosis* (9)

indicates an over-reliance of *M. bovis* on cholesterol transport through the Mce4

transporter (Figure 5).

Early stages of cholesterol catabolism involve the oxidation of cholesterol to

cholestenone, a reaction catalysed by the 3β-hydroxysteroid dehydrogenase (*hsd*) encoded

by *Rv1106c/Mb1136c* (37). The cytochrome P450 Cyp125 (*Mb3575c/Rv3545c*) is required

for the subsequent detoxification of cholestenone (38). Insertions in both *hsd* and *cyp125* in

M. bovis were severely attenuating with $\log_2$ fold-changes of ~-5 to -7 (Supplementary Table

S4 - cholesterol catabolism tab, Figure 5). Previous studies have shown that these genes are

not required for the survival of *M. tuberculosis* in macrophages or in guinea pigs and this is

thought to be due to the availability of other carbon sources, including glycolytic substrates,

in vivo (37, 39–43). *M. bovis* is more restricted in metabolic capabilities and is unable to

generate energy from glycolytic intermediates, largely due to a disrupted pyruvate kinase

encoded by *pykA* (44, 45). The essentiality of *hsd* and *cyp125* during infection for *M. bovis*

but not *M. tuberculosis* supports the hypothesis of an over-reliance of *M. bovis* on

cholesterol. Given the potential for the use of host cholesterol metabolites as diagnostic

biomarkers, this observation might have applications in the development of differential

diagnostics (46).

Genes that are differentially expressed between *Mycobacterium bovis* and

Mycobacterium tuberculosis.

Several studies have identified key expression differences between *M. bovis* and *M.*

tuberculosis (27, 47, 48). We examined the dataset for insights on the role of differentially

expressed genes and transcriptional regulators during infection. One important regulatory

system in *M. tuberculosis* is the two-component regulatory system PhoPR and deletions in

the *phoPR* genes alongside *fadD26* are attenuating mutations in the live vaccine MTBVAC

(49–51). Our data show that insertions in both *phoPR* and *fadD26* were severely attenuating

with $\log_2$ fold-changes of -6 to -9 (Figure 6, Supplementary Table S4, *phoPR* regulon tab and

mycolipids tab). This reinforces the role of this system in virulence, despite the presence of a

single nucleotide polymorphism (SNP) in the sensor kinase *phoR* that impacts signalling

through the system in *M. bovis* (52). Signal potentiation via *phoR* is required for secretion of

ESAT-6 through the ESX-1 secretory system and *M. bovis* is known to have compensatory

mutations elsewhere in the genome, e.g. in the *espACD* operon, that restores ESAT-6

secretion in the face of a deficient signalling system (49, 52, 53). Our data also show that Tn

insertions in *espA* of the *espACD* operon (required for ESAT-6 secretion) and in *mprA*, a

transcriptional regulator of that operon (54) were severely attenuating (log2 fold-changes -7

to -9), emphasising the relevance of ESAT-6 as a virulence factor.

Studies comparing differences in expression during *in vitro* growth between *M. bovis*

and *M. tuberculosis* show that genes involved in sulfolipid (SL-1) biosynthesis are expressed

at lower levels in *M. bovis* compared to *M. tuberculosis* (27, 47). Interestingly, insertions in

genes involved in SL-1 biosynthesis (*Mb3850-Mb3856*) are not attenuating *in vivo*

(Supplementary File S4, mycolipids tab), reinforcing the lack of importance of SL-1 for *M.*

bovis in vivo, at least at the stages of infection studied here.

One of the most highly attenuating insertions occurred in *Mb0222/Rv0216* (log_2 fold

change –8 to -9). This gene has been shown to be highly (> 10-fold) over-expressed in *M.*

bovis compared with *M. tuberculosis* but the physiological function of this gene is not

currently known. The secreted antigens MPB70 and MPB83, encoded by *Mb2900* and

Mb2898 are also over-expressed in *M. bovis* and play a role in host-specific immune

responses, however, insertions in these genes did not cause attenuation *in vivo* in our

dataset (55).

Novel attenuating mutations

We identified 32 genes that were required for survival of *M. bovis* in cattle that had not

been previously described as being essential *in vivo* through transposon mutagenesis

screens of *M. tuberculosis* in mouse models (8, 9, 13) (see Supplementary Table 4,

Significant in 50% of cattle tab). While writing this publication, a large scale Tn-seq study

that utilised over 120 *M. tuberculosis* libraries and several diverse mouse genotypes (the

collaborative cross mouse panel (56)) showed that the panel of genes required for the

survival of *M. tuberculosis in vivo* is much larger than previously reported (57). A direct

comparison of our dataset with the study by Smith et al., revealed that a further 13 genes

were shown to be required in at least one mouse strain in that study. A summary set of the

remaining 19 genes is given in Supplementary Table 4, Not in Mtb Tn-seqs tab. Some of

these genes have been shown to be attenuated in the mouse model in *M. tuberculosis*

through the use of single mutants (58–61).

Included in this list are genes required for phenolic glycolipid synthesis. Insertions in

Mb2971c/Rv2947c (*pks15/1*) and in *Mb2972c/Rv2948c* (*fadD22*) were attenuating in *M.*

bovis but these genes are not required *in vivo* in *M. tuberculosis,* including in the extended

panel of mouse genotypes (8, 9, 13, 57). Both *pks15/1* and *fadD22* are involved in the early

stages of synthesis of phenolic glycolipids (PGLs) and are involved in virulence (62). The

requirement for these genes in *M. bovis* but not in *M. tuberculosis* is consistent with the

observation that the Tn-seq studies in *M. tuberculosis* are often carried out using lineage 4

strains (H37Rv and CDC1551) that harbour a frameshift mutation in the *pks15/1* gene, which

renders them unable to synthesise PGLs. This removes the requirement for these genes *in*

vivo in lineage 4 strains of *M. tuberculosis. pks15/1* has been previously reported to be

required for survival of *M. bovis* in a guinea pig model of infection (63).

The list also includes genes that are involved in post-translational modifications such

as glycosylation. *Rv1002c* is thought to add mannose groups to secreted proteins and over-

expression of this protein in *M. smegmatis* was recently shown to enhance survival *in vivo*

and inhibit pro-inflammatory cytokine production (64). The substrates of this protein

mannosyltransferase are thought to be several secreted lipoproteins, including LpqW which

is involved in the insertion of the virulence lipid LAM at the mycobacterial cell surface (64,

65).

Finally, this list includes two genes (*aspC* and *glpD2*) that are essential *in vitro* in *M.*

tuberculosis but not in *M. bovis* (10, 11, 18, 66). Information regarding *aspC* and *glpD2* from

Tn-seq approaches is likely to be lacking in *M. tuberculosis* because Tn mutants will not be

represented in the input pool. The absence of insertion mutants in these genes in the most

recent large-scale *M. tuberculosis* Tn-seq study supports this (57). One of these genes

MB0344c/Rv0337c (*aspC*) is an aspartate aminotransferase involved in the utilisation of

amino acids (aspartate) as a nitrogen source (67). The other gene *Mb3303c/Rv3302c* (*glpD2*)

is a membrane bound glycerol-phosphate dehydrogenase. In *Escherichia coli*, *glpD2* is an

essential enzyme, functioning at the central junction of respiration, glycolysis, and

phospholipid biosynthesis and catalyses the oxidation of dihydroxyacetone phosphate

(DHAP) from glycerol-3-phosphate resulting in the donation of electrons to the electron

transport chain (68). Its essentiality *in vitro* in *M. tuberculosis* might be explained by the

usage of glycerol during *in vitro* growth in this species. The contribution of the membrane

bound *glpD2* in donation of electrons to the electron transport chain, has been suggested

but not yet explored in the MTBC (69). Given the interest in the electron transport chain as a

chemotherapeutic target in *M. tuberculosis*, the data presented here suggests that

inhibition of *glpD2* might be a fruitful approach in the development of new drugs for the

treatment of TB in humans (70). The role of this gene in *M. bovis in vivo* is perhaps

surprising, given the disruptions in glycerol phosphate uptake and pathways that

phosphorylate glycerol in *M. bovis* AF2122/97 (71). However, *M. tuberculosis* is thought to

engage in catabolism of membrane derived glycerophospholipids which may be a potential

source of glycerol-3-phosphate in members of the complex (72).

Materials and Methods

Bacterial strains and culture methods

M. bovis strain AF2122/97 was maintained on modified Middlebrook 7H11 (BD Difco^{TM})

medium (73). Liquid cultures of *M. bovis* were grown in Middlebrook 7H9 media (BD

Difco^{TM}) containing 75 mM sodium pyruvate, 0.05% v/v Tween$^{®}$80 and 10% Middlebrook

albumin-dextrose-catalase (ADC) (BBL BD Biosciences). Kanamycin at 25 µg/ml was used for

selection where appropriate.

Generation of input transposon mutant library and preparation of the inoculum

Transposon libraries in *M. bovis* were generated as previously described using the

MycomarT7 phagemid system as per Majumdar et al with modifications (19). Approximately

66,000 kanamycin resistant transductants were scraped and homogenised in 7H9 medium

and stored frozen at -80°C in 1 ml aliquots. CFU counting was performed on the

homogenised culture to inform inoculum dosage.

Cattle Infection

Experiments were carried out according to the UK Animal (Scientific Procedures) Act 1986

under project license PPL70/7737. Ethical permission was obtained from the APHA Animal

Welfare Ethical Review Body (AWERB) (UK Home Office PCD number 70/6905). All animal

infections were carried out within the APHA large animal biocontainment level 3 facility.

Twenty-four Holstein-Friesian crosses of 6 months of age were sourced from an officially TB-

free herd. An infectious dose of 7 x 10^4 CFU was targeted for the "input" library, allowing

each mutant to be represented in the library ~ 2.5-fold. Retrospective counting of the

inoculum revealed the actual inoculum for infection contained 4 x 10^4 CFU. The inoculum

was delivered endobronchially in 2 ml of 7H9 medium. In brief, animals were sedated with

xylazine (Rompun® 2%, Bayer, France) according to the manufacturer's instructions (0.2

mL/100 kg, IV route) prior to the insertion of an endoscope through the nasal cavity into the

trachea for delivery of the inoculum through a 1.8 mm internal diameter cannula

(Veterinary Endoscopy Services, U.K.) above the bronchial opening to the cardiac lobe and

the main bifurcation between left and right lobes.

Infection Monitoring with the IFN-γ release Assay (IGRA)

Blood was collected by jugular venepuncture from animals on the day of the infectious

challenge and two weeks after infection. Heparinized whole blood (250 μl) was incubated

with purified protein derivative (PPD) from *M. avium* (PPD-A) or PPD from *M. bovis* (PPD-B)

(Prionics™) respectively at 25 IU and 30 IU final. Pokeweed mitogen was used as the positive

control at 10 μg/mL and a medium-only negative control. After 24 h incubation in 5% (v/v)

CO_2, 95% humidity, 37 °C atmosphere bloods were centrifuged (400 × g for 5 min); 120 μl

of supernatant was removed and stored at −80°C for subsequent IFN-γ quantification using

the BOVIGAM® kit (Prionics™) in accordance with the manufacturer's instructions.

Collection of tissues and gross pathology scores

Six weeks after the initial infection animals were subjected to post-mortem examination.

Initially the experiment was designed with two time points; an early time point (6 weeks)

and a later time point of 8 weeks. However, due to the unexpected high-levels of pathology

seen at the earlier time-points all animals were culled at 6 weeks. Gross pathology and

evidence of TB-like granulomas lesions was scored using a modified methodology to that

previously described in (74). Tissue from head and neck lymph nodes (from the right and left

sub-mandibular lymph nodes, the right and left medial retropharyngeal lymph nodes),

thoracic lymph nodes (the right and left bronchial lymph nodes, the cranial tracheobronchial

lymph nodes, the cranial and caudal mediastinal lymph nodes) and from lung lesions, was

collected into sterile containers and frozen at −80°C until further processing. Frozen tissues

were thawed and homogenised in PBS using a Seward Stomacher Paddle Blender.

Recovery of the output transposon mutant library from tissues

Tissue macerates collected from study animals were thawed at room temperature, diluted

in PBS and plated on modified 7H11 agar to determine bacterial loads. Colony counts were

performed after 3-4 weeks growth. For recovery of the library from tissue macerates ~10^5-

10^6 CFU were plated from lung lesions and thoracic lymph node lesions onto modified 7H11

media containing 25 µg/ml kanamycin. The colonies were plated over several 140 mm petri

dishes to minimise competition between mutants. The colonies were harvested after 4-6

weeks growth and genomic DNA extracted.

Genomic DNA extraction

Genomic DNA from the input and recovered libraries was isolated by an extended bead

beating procedure with detergent-based lysis, phenol-chloroform DNA extraction and

precipitation as previously described (18). DNA quality was assessed by nano-spectrometry

(DeNovix) and gel electrophoresis and quantified by Qubit analysis using the Broad Range

Assay Kit (ThermoScientific).

Library preparation for transposon directed insertion sequencing

DNA (2 µg) was resuspended in 50 µL distilled water and sheared to approximately 550 bp

fragments using a S220 focussed-ultrasonicator (Covaris), according to the manufacturer's

protocol. Fragmented DNA was repaired using NEBNext blunt-end repair kit (New England

Biolabs) and purified using Monarch PCR clean-up kit (NEB). Blunted DNA was A-tailed using

NEBNext dA-tailing kit (NEB) and column purified. Custom transposon sequencing adaptors

(Supplementary Table S3) were generated by heating an equimolar mix of Com_AdaptorPt1

primer and Com_AdaptorPt2 (P7+index) primers to 95°C for 5 min, followed by cooling by

1°C every 40 s to a final temperature of 4°C in a thermocycler. Adaptors were ligated to A-

tailed library fragments using NEBNext quick ligase kit. Transposon-containing fragments

were enriched by PCR using ComP7 primer (10 µM) and an equimolar mix of primers P5-

IR2a-d primer (10 µM) in a reaction with 50 ng of adaptor ligated template and Phusion DNA

polymerase (NEB) in a thermocycler with the following program 98°C 3 min; 4 cycles of 98°C

20s, 70°C 20s, 72°C 1 min; 20 cycles of 98°C 20s, 67°C 20s, 72°C 1 min; 72°C 3 min.

Transposon-enriched libraries were subsequently purified with AMPureXP beads

(Beckman), pooled together and further purified using AMPure XP beads.

Data analysis

Indexed libraries were combined, spiked with 20% PhiX, and sequenced on the Illumina

Hiseq 3000 platform, using v2 chemistry, generating single-end reads of 250 bp. Raw .fastq

sequencing files were analysed for quality and pre-processed using the TRANSIT TPP tool

(75) set to default 'Sassetti' protocol, in order to remove transposon tags and adapter

sequences, and to map reads using BWA-mem to TA sites to the *M. bovis* AF2122/97

genome (NC_002945.3). The TRANSIT 'tnseq_stats' tool was run on each sample to assess

insertion density, skew, kurtosis and potential amplification bias.

The *M. bovis* AF2122/97 genome was scanned for the non-permissive Himar1

transposon insertion motif ('SGNTANCS', where S is either G or C and N is any base) as

previously described [10]. 6605 sites were identified as non-permissive (approximately 9%

of total TA sites) and excluded from resampling analysis. A custom annotation, '.prot-table'

for TRANSIT, was created from the *M. bovis* AF2122/97 annotation file (NCBI Accession

Number LT708304, version LT708304.1). TRANSIT HMM was run on the input library using

the default normalisation (TTR) with LOESS correction for genomic position bias. Each TA

site was assigned an essentiality state and genes were assigned an essentiality call based on

the assigned state of the TA sites within annotated gene boundaries.

Resampling between the input library and each of the output sample libraries was

performed independently using the TRANSIT resampling algorithm and the complete prot-

table. TTR normalisation was used for 23 of the samples, and betageom normalisation for

the three samples with skew of greater than 50. The initial resampling output files were

evaluated to identify genes with very few, or no, reads at any TA site within the gene

boundaries in both the input library and output sample libraries. Genes with no read counts

greater than 4 at any TA site, in any sample, and with a sum of all reads at any TA site across

the 26 samples less than 55, were flagged. Essential and unchanged genes were removed

from the prot-table prior to further evaluation. Resampling was further limited to protein-

coding genes. Resampling was re-run for each sample using the attenuated prot-table and

an edited TRANSIT resampling script to return the left-tail p-value, as the data were

expected to reflect attenuation. All p-values were corrected for multiple testing with FDR

adjustment.

All analysis and plots were performed using R and R packages, tidyverse and circlize

(76–78). Orthologous TB genes were obtained from supplementary data files published by

Malone et al, 2018 (27). All scripts, prot-tables and insertion files are available at

https://github.com/jenjane118/Mbovis_in-vivo_Tnseq,   DOI:10.5281/zenodo.6354151.

Sequencing files (.fastq) are deposited in SRA (Bioproject ID: PRJNA816175, Submission ID:

SUB11067380)

Funding

This work was funded by the BBSRC Grant Ref: BB/N004590/1 [awarded to SK (PI), DW (Co-

I), BW (Co-I), and SE3314 to BV-R as part of the joint BBSRC-DEFRA EradbTB consortium. AG,

IP, and SW were supported by the funding. VF was in receipt of an RVC PhD studentship. AG

currently holds a Sêr Cymru II Lectureship funded by the European Research Development

Fund and Welsh Government. BV-R is a Ser Cymru II Professor of Immunology at

Aberystwyth University. JS is supported by a Bloomsbury Colleges PhD Studentship (LIDo

program).

458   AUTHOR CONTRIBUTIONS

459   SK, DW, BW, BV-R and SB undertook funding acquisition and designed the study. AJG, VF,

460   JM, SW, IP, MC carried out the experimental work. Data analysis was done by IN and JS.

461   AJG, JS and SLK wrote the first draft of the manuscript. All authors contributed to the

462   manuscript revision, read, and approved the submitted version.

464   Acknowledgements

465   The authors would like to acknowledge the help and support of APHA colleagues from the

466   BAC4 workgroup and the Pathology Unit, and in particular acknowledge the care and

467   support provided to animals under experimentation by members of APHA's Animals

468   Sciences Unit. We would like to thank Dr Dany Beste for useful discussions surrounding the

469   role of *glpD2* in mycobacterial metabolism.

References

1. Müller B, Dürr S, Alonso S, Hattendorf J, Laisse CJM, Parsons SDC, van Helden PD,

Zinsstag J. 2013. Zoonotic Mycobacterium bovis-induced tuberculosis in humans.

Emerg Infect Dis 19:899–908.

2. Mableson HE, Okello A, Picozzi K, Welburn SC. 2014. Neglected Zoonotic Diseases—

The Long and Winding Road to Advocacy. PLoS Negl Trop Dis 8:e2800.

3. Bayissa B, Sirak A, Worku A, Zewude A, Zeleke Y, Chanyalew M, Gumi B, Berg S,

Conlan A, Hewinson RG, Wood JLN, Vordermeier HM, Ameni G. 2021. Evaluation of

the Efficacy of BCG in Protecting Against Contact Challenge With Bovine Tuberculosis

in Holstein-Friesian and Zebu Crossbred Calves in Ethiopia. Front Vet Sci 8.

4. Srinivasan S, Conlan AJK, Easterling LA, Herrera C, Dandapat P, Veerasami M, Ameni

G, Jindal N, Raj GD, Wood J, Juleff N, Bakker D, Vordermeier M, Kapur V. 2021. A

Meta-Analysis of the Effect of Bacillus Calmette-Guérin Vaccination Against Bovine

Tuberculosis: Is Perfect the Enemy of Good? Front Vet Sci 0:116.

5. Vordermeier HM, Whelan A, Cockle PJ, Farrant L, Palmer N, Hewinson RG. 2001. Use

of synthetic peptides derived from the antigens ESAT-6 and CFP-10 for differential

diagnosis of bovine tuberculosis in cattle. Clin Diagn Lab Immunol 8:571–578.

6. Whelan AO, Clifford D, Upadhyay B, Breadon EL, McNair J, Hewinson GR, Vordermeier

MH. 2010. Development of a skin test for bovine tuberculosis for differentiating

infected from vaccinated animals. J Clin Microbiol 48:3176–3181.

7. Cain AK, Barquist L, Goodman AL, Paulsen IT, Parkhill J, van Opijnen T. 2020. A decade

of advances in transposon-insertion sequencing. Nat Rev Genet. Nat Rev Genet.

8.   Zhang YJ, Reddy MC, Ioerger TR, Rothchild AC, Dartois V, Schuster BM, Trauner A,

Wallis D, Galaviz S, Huttenhower C, Sacchettini JC, Behar SM, Rubin EJ. 2013.

Tryptophan biosynthesis protects mycobacteria from CD4 T-cell-mediated killing. Cell

155:1296–1308.

9.   Bellerose MM, Proulx MK, Smith CM, Baker RE, Ioerger TR, Sassetti CM. 2020. Distinct

Bacterial Pathways Influence the Efficacy of Antibiotics against Mycobacterium

tuberculosis. mSystems 5.

10. Dejesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, Rubin EJ, Schnappinger D,

Ehrt S, Fortune SM, Sassetti CM, Ioerger TR. 2017. Comprehensive essentiality

analysis of the Mycobacterium tuberculosis genome via saturating transposon

mutagenesis. MBio 8.

11. Griffin JE, Gawronski JD, Dejesus MA, Ioerger TR, Akerley BJ, Sassetti CM. 2011. High-

resolution phenotypic profiling defines genes essential for mycobacterial growth and

cholesterol catabolism. PLoS Pathog 7:e1002251.

12. Patil S, Palande A, Lodhiya T, Pandit A, Mukherjee R. 2021. Redefining genetic

essentiality in Mycobacterium tuberculosis. Gene 765.

13. Sassetti CM, Rubin EJ. 2003. Genetic requirements for mycobacterial survival during

infection. Proc Natl Acad Sci U S A 100:12989–12994.

14. Rhijn I Van, Branch Moody D. CD1 and mycobacterial lipids activate human T cells

https://doi.org/10.1111/imr.12253.

15. Dutta NK, Mehra S, Didier PJ, Roy CJ, Doyle LA, Alvarez X, Ratterree M, Be NA,

Lamichhane G, Jain SK, Lacey MR, Lackner AA, Kaushal D. 2010. Genetic Requirements

for the Survival of Tubercle Bacilli in Primates. J Infect Dis 201:1743–1752.

16.   Mendum TA, Chandran A, Williams K, Vordermeier HM, Villarreal-Ramos B, Wu H,

Singh A, Smith AA, Butler RE, Prasad A, Bharti N, Banerjee R, Kasibhatla SM, Bhatt A,

Stewart GR, McFadden J. 2019. Transposon libraries identify novel Mycobacterium

bovis BCG genes involved in the dynamic interactions required for BCG to persist

during in vivo passage in cattle. BMC Genomics 20:431.

17.   Smith AA, Villarreal-Ramos B, Mendum TA, Williams KJ, Jones GJ, Wu H, McFadden J,

Vordermeier HM, Stewart GR. 2020. Genetic screening for the protective antigenic

targets of BCG vaccination. Tuberculosis 124:101979.

18.   Gibson AJ, Passmore IJ, Faulkner V, Xia D, Nobeli I, Stiens J, Willcocks S, Clark TG,

Sobkowiak B, Werling D, Villarreal-Ramos B, Wren BW, Kendall SL. 2021. Probing

Differences in Gene Essentiality Between the Human and Animal Adapted Lineages of

the Mycobacterium tuberculosis Complex Using TnSeq. Front Vet Sci 8.

19.   Majumdar G, Mbau R, Singh V, Warner DF, Dragset MS, Mukherjee R. 2017. Genome-

wide transposon mutagenesis in Mycobacterium tuberculosis and Mycobacterium

smegmatis, p. 321–335. In Methods in Molecular Biology. Humana Press, New York,

NY.

20.   Lewis KN, Liao R, Guinn KM, Hickey MJ, Smith S, Behr MA, Sherman DR. 2003.

Deletion of RD1 from Mycobacterium tuberculosis Mimics Bacille Calmette-Guérin

Attenuation . J Infect Dis 187:117–123.

21.   Phan TH, van Leeuwen LM, Kuijl C, Ummels R, van Stempvoort G, Rubio-Canalejas A,

Piersma SR, Jiménez CR, van der Sar AM, Houben ENG, Bitter W. 2018. EspH is a

hypervirulence factor for Mycobacterium marinum and essential for the secretion of

the ESX-1 substrates EspE and EspF. PLoS Pathog 14:e1007247.

22. Inwald J, Jahans K, Hewinson RG, Gordon S V. 2003. Inactivation of the

Mycobacterium bovis homologue of the polymorphic RD1 gene Rv3879c (Mb3909c)

does not affect virulence. Tuberculosis (Edinb) 83:387–393.

23. Bold TD, Davis DC, Penberthy KK, Cox LM, Ernst JD, de Jong BC. 2012. Impaired fitness

of Mycobacterium africanum despite secretion of ESAT-6. J Infect Dis 205:984–90.

24. Hotter GS, Wards BJ, Mouat P, Besra GS, Gomes J, Singh M, Bassett S, Kawakami P,

Wheeler PR, De Lisle GW, Collins DM. 2005. Transposon mutagenesis of Mb0100 at

the ppe1-nrp locus in Mycobacterium bovis disrupts phthiocerol dimycocerosate

(PDIM) and glycosylphenol-PDIM biosynthesis, producing an avirulent strain with

vaccine properties at least equal to those of M. bovis BCG. J Bacteriol 187:2267–2277.

25. Cox JS, Chess B, McNeil M, Jacobs WR. 1999. Complex lipid determines tissue-specific

replication of Mycobacterium tuberculosis in mice. Nature 402:79–83.

26. Pérez J, Garcia R, Bach H, de Waard JH, Jacobs WR, Av-Gay Y, Bubis J, Takiff HE. 2006.

Mycobacterium tuberculosis transporter MmpL7 is a potential substrate for kinase

PknD. Biochem Biophys Res Commun 348:6–12.

27. Malone KM, Rue-Albrecht K, Magee DA, Conlon K, Schubert OT, Nalpas NC, Browne

JA, Smyth A, Gormley E, Aebersold R, Machugh DE, Gordon S V. 2018. Comparative

'omics analyses differentiate mycobacterium tuberculosis and mycobacterium bovis

and reveal distinct macrophage responses to infection with the human and bovine

tubercle bacilli. Microb Genomics 4.

28.   Pisu D, Huang L, Grenier JK, Russell DG. 2020. Dual RNA-Seq of Mtb-Infected

559   Macrophages In Vivo Reveals Ontologically Distinct Host-Pathogen Interactions. Cell

560   Rep 30:335-350.e4.

561   29.   Van Der Geize R, Yam K, Heuser T, Wilbrink MH, Hara H, Anderton MC, Sim E,

562   Dijkhuizen L, Davies JE, Mohn WW, Eltis LD. 2007. A gene cluster encoding cholesterol

563   catabolism in a soil actinomycete provides insight into Mycobacterium tuberculosis

564   survival in macrophages. Proc Natl Acad Sci U S A 104:1947–1952.

565   30.   Nesbitt NM, Yang X, Fontán P, Kolesnikova I, Smith I, Sampson NS, Dubnau E. 2010. A

566   thiolase of Mycobacterium tuberculosis is required for virulence and production of

567   androstenedione and androstadienedione from cholesterol. Infect Immun 78:275–

568   282.

569   31.   Bandyopadhyay U, Chadha A, Gupta P, Tiwari B, Bhattacharyya K, Popli S, Raman R,

570   Brahamachari V, Singh Y, Malhotra P, Natarajan K. 2017. Suppression of Toll-like

571   receptor 2-mediated proinflammatory responses by Mycobacterium tuberculosis

572   protein Rv3529c. J Leukoc Biol 102:1249–1259.

573   32.   Pandey AK, Sassetti CM. 2008. Mycobacterial persistence requires the utilization of

574   host cholesterol. Proc Natl Acad Sci U S A 105:4376–4380.

575   33.   Mohn WW, Van Der Geize R, Stewart GR, Okamoto S, Liu J, Dijkhuizen L, Eltis LD.

576   2008. The actinobacterial mce4 locus encodes a steroid transporter. J Biol Chem

577   283:35368–35374.

578   34.   Lee W, VanderVen BC, Fahey RJ, Russell DG. 2013. Intracellular Mycobacterium

579   tuberculosis exploits host-derived fatty acids to limit metabolic stress. J Biol Chem

288:6788–6800.

35.   Jain M, Petzold CJ, Schelle MW, Leavell MD, Mougous JD, Bertozzi CR, Leary JA, Cox

JS. 2007. Lipidomics reveals control of Mycobacterium tuberculosis virulence lipids via

metabolic coupling. Proc Natl Acad Sci U S A 104:5133.

36.   Singh P, Sinha R, Tyagi G, Sharma NK, Saini NK, Chandolia A, Prasad AK, Varma-Basil

M, Bose M. 2018. PDIM and SL1 accumulation in Mycobacterium tuberculosis is

associated with mce4A expression. Gene 642:178–187.

37.   Yang X, Gao J, Smith I, Dubnau E, Sampson NS. 2011. Cholesterol Is Not an Essential

Source of Nutrition for Mycobacterium tuberculosis during Infection. J Bacteriol

193:1473.

38.   Ouellet H, Guan S, Johnston JB, Chow ED, Kells PM, Burlingame AL, Cox JS, Podust LM,

De Montellano PRO. 2010. Mycobacterium tuberculosis CYP125A1, a steroid C27

monooxygenase that detoxifies intracellularly generated cholest-4-en-3-one. Mol

Microbiol 77:730–742.

39.   Serafini A, Tan L, Horswell S, Howell S, Greenwood DJ, Hunt DM, Phan MD, Schembri

M, Monteleone M, Montague CR, Britton W, Garza-Garcia A, Snijders AP, VanderVen

B, Gutierrez MG, West NP, de Carvalho LPS. 2019. Mycobacterium tuberculosis

requires glyoxylate shunt and reverse methylcitrate cycle for lactate and pyruvate

metabolism. Mol Microbiol 112:1284–1307.

40.   Safi H, Gopal P, Lingaraju S, Ma S, Levine C, Dartois V, Yee M, Li L, Blanc L, Liang HPH,

Husain S, Hoque M, Soteropoulos P, Rustad T, Sherman DR, Dick T, Alland D. 2019.

Phase variation in Mycobacterium tuberculosis glpK produces transiently heritable

drug tolerance. Proc Natl Acad Sci U S A 116:19665–19674.

41.   De Carvalho LPS, Fischer SM, Marrero J, Nathan C, Ehrt S, Rhee KY. 2010.

Metabolomics of Mycobacterium tuberculosis Reveals Compartmentalized Co-

Catabolism of Carbon Substrates. Chem Biol 17:1122–1131.

42.   Carroll P, Parish T. 2015. Deletion of cyp125 Confers Increased Sensitivity to Azoles in

Mycobacterium tuberculosis. PLoS One 10:e0133129.

43.   Borah K, Mendum TA, Hawkins ND, Ward JL, Beale MH, Larrouy-Maumus G, Bhatt A,

Moulin M, Haertlein M, Strohmeier G, Pichler H, Forsyth VT, Noack S, Goulding CW,

McFadden J, Beste DJ V. 2021. Metabolic fluxes for nutritional flexibility of

Mycobacterium tuberculosis. Mol Syst Biol 17.

44.   Keating LA, Wheeler PR, Mansoor H, Inwald JK, Dale J, Hewinson RG, Gordon S V.

2005. The pyruvate requirement of some members of the Mycobacterium

tuberculosis complex is due to an inactive pyruvate kinase: implications for in vivo

growth. Mol Microbiol 56:163–174.

45.   Lofthouse EK, Wheeler PR, Beste DJV, Khatri BL, Wu H, Mendum TA, Kierzek AM,

McFadden J. 2013. Systems-based approaches to probing metabolic variation within

the Mycobacterium tuberculosis complex. PLoS One 8.

46.   Chandra P, Coullon H, Agarwal M, Goss CW, Philips JA. 2022. Macrophage global

metabolomics identifies cholestenone as host/pathogen cometabolite present in

human Mycobacterium tuberculosis infection. J Clin Invest 132.

47.   Golby P, Hatch KA, Bacon J, Cooney R, Riley P, Allnutt J, Hinds J, Nunez J, Marsh PD,

Hewinson RG, Gordon S V. 2007. Comparative transcriptomics reveals key gene

expression differences between the human and bovine pathogens of the

Mycobacterium tuberculosis complex. Microbiology 153:3323–3336.

48.   Rehren G, Walters S, Fontan P, Smith I, Zárraga AM. 2007. Differential gene

expression between Mycobacterium bovis and Mycobacterium tuberculosis.

Tuberculosis (Edinb) 87:347–359.

49.   Gonzalo-Asensio J, Malaga W, Pawlik A, Astarie-Dequeker C, Passemar C, Moreau F,

Laval F, Daffé M, Martin C, Brosch R, Guilhot C. 2014. Evolutionary history of

tuberculosis shaped by conserved mutations in the PhoPR virulence regulator. Proc

Natl Acad Sci U S A 111:11491–11496.

50.   White AD, Sibley L, Sarfas C, Morrison A, Gullick J, Clark S, Gleeson F, McIntyre A,

Arlehamn CL, Sette A, Salguero FJ, Rayner E, Rodriguez E, Puentes E, Laddy D,

Williams A, Dennis M, Martin C, Sharpe S. 2021. MTBVAC vaccination protects rhesus

macaques against aerosol challenge with M. tuberculosis and induces immune

signatures analogous to those observed in clinical studies. NPJ Vaccines 6.

51.   Gonzalo-Asensio J, Marinova D, Martin C, Aguilo N. 2017. MTBVAC: Attenuating the

Human Pathogen of Tuberculosis (TB) Toward a Promising Vaccine against the TB

Epidemic. Front Immunol 8:1803.

52.   Altes HK, Dijkstra F, Lugnèr A, Cobelens F, Wallinga J. 2009. Targeted bcg vaccination

against severe tuberculosis in low-prevalence settings epidemiologic and economic

assessment. Epidemiology 20:562–568.

53.   Frigui W, Bottai D, Majlessi L, Monot M, Josselin E, Brodin P, Garnier T, Gicquel B,

Martin C, Leclerc C, Cole ST, Brosch R. 2008. Control of M. tuberculosis ESAT-6

secretion and specific T cell recognition by PhoP. PLoS Pathog 4.

54. Pang X, Samten B, Cao G, Wang X, Tvinnereim AR, Chen XL, Howard ST. 2013. MprAB

regulates the espA operon in Mycobacterium tuberculosis and modulates ESX-1

function and host cytokine response. J Bacteriol 195:66–75.

55. Queval CJ, Fearns A, Botella L, Smyth A, Schnettger L, Mitermite M, Wooff E,

Villarreal-Ramos B, Garcia-Jimenezi W, Heunis T, Trosti M, Werlingi D, Salguero FJ,

Gordon S V., Gutierrez MG. 2021. Macrophage-specific responses to human- and

animal-adapted tubercle bacilli reveal pathogen and host factors driving

multinucleated cell formation. PLoS Pathog 17.

56. Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, Beavis WD, Belknap JK,

Bennett B, Berrettini W, Bleich A, Bogue M, Broman KW, Buck KJ, Buckler E,

Burmeister M, Chesler EJ, Cheverud JM, Clapcote S, Cook MN, Cox RD, Crabbe JC,

Crusio WE, Darvasi A, Deschepper CF, Doerge RW, Farber CR, Forejt J, Gaile D, Garlow

SJ, Geiger H, Gershenfeld H, Gordon T, Gu J, Gu W, de Haan G, Hayes NL, Heller C,

Himmelbauer H, Hitzemann R, Hunter K, Hsu HC, Iraqi FA, Ivandic B, Jacob HJ, Jansen

RC, Jepsen KJ, Johnson DK, Johnson TE, Kempermann G, Kendziorski C, Kotb M, Kooy

RF, Llamas B, Lammert F, Lassalle JM, Lowenstein PR, Lu L, Lusis A, Manly KF,

Marcucio R, Matthews D, Medrano JF, Miller DR, Mittleman G, Mock BA, Mogil JS,

Montagutelli X, Morahan G, Morris DG, Mott R, Nadeau JH, Nagase H, Nowakowski

RS, O'Hara BF, Osadchuk A V., Page GP, Paigen B, Paigen K, Palmer AA, Pan HJ,

Peltonen-Palotie L, Peirce J, Pomp D, Pravenec M, Prows DR, Qi Z, Reeves RH, Roder J,

Rosen GD, Schadt EE, Schalkwyk LC, Seltzer Z, Shimomura K, Shou S, Sillanpää MJ,

Siracusa LD, Snoeck HW, Spearow JL, Svenson K, Tarantino LM, Threadgill D, Toth LA,

Valdar W, Pardo-Manuel de Villena F, Warden C, Whatley S, Williams RW, Wiltshire T,

Yi N, Zhang D, Zhang M, Zou F. 2004. The Collaborative Cross, a community resource

for the genetic analysis of complex traits. Nat Genet 2004 3611 36:1133–1137.

57.   Smith CM, Baker RE, Proulx MK, Mishra BB, Long JE, Park SW, Lee H-N, Kiritsy MC,

Bellerose MM, Olive AJ, Murphy KC, Papavinasasundaram K, Boehm FJ, Reames CJ,

Meade RK, Hampton BK, Linnertz CL, Shaw GD, Hock P, Bell TA, Ehrt S, Schnappinger

D, Pardo-Manuel de Villena F, Ferris MT, Ioerger TR, Sassetti CM. 2022. Host-

pathogen genetic interactions underlie tuberculosis susceptibility in genetically

diverse mice. Elife 11.

58.   Cowley S, Ko M, Pick N, Chow R, Downing KJ, Gordhan BG, Betts JC, Mizrahi V, Smith

DA, Stokes RW, Av-Gay Y. 2004. The Mycobacterium tuberculosis protein

serine/threonine kinase PknG is linked to cellular glutamate/glutamine levels and is

important for growth in vivo. Mol Microbiol 52:1691–1702.

59.   Liu CF, Tonini L, Malaga W, Beau M, Stella A, Bouyssié D, Jackson MC, Nigou J, Puzo G,

Guilhot C, Burlet-Schiltz O, Rivière M. 2013. Bacterial protein-O-mannosylating

enzyme is crucial for virulence of Mycobacterium tuberculosis. Proc Natl Acad Sci U S

A 110:6560–6565.

60.   Maksymiuk C, Balakrishnan A, Bryk R, Rhee KY, Nathan CF. 2015. E1 of☐?-

ketoglutarate dehydrogenase defends Mycobacterium tuberculosis against glutamate

anaplerosis and nitroxidative stress. Proc Natl Acad Sci U S A 112:E5834–E5843.

61.   Venugopal A, Bryk R, Shi S, Rhee K, Rath P, Schnappinger D, Ehrt S, Nathan C. 2011.

Virulence of Mycobacterium tuberculosis depends on lipoamide dehydrogenase, a

member of three multienzyme complexes. Cell Host Microbe 9:21–31.

62.   Constant P, Perez E, Malaga W, Lanéelle MA, Saurel O, Daffé M, Guilhot C. 2002. Role

of the pks15/1 Gene in the Biosynthesis of Phenolglycolipids in the Mycobacterium

tuberculosisComplex: EVIDENCE THAT ALL STRAINS SYNTHESIZE GLYCOSYLATEDp-

HYDROXYBENZOIC METHYL ESTERS AND THAT STRAINS DEVOID OF

PHENOLGLYCOLIPIDS HARBOR A FRAMESHIFT MU. J Biol Chem 277:38148–38158.

63.   Samanta S, Singh A, Biswas P, Bhatt A, Visweswariah SS. 2017. Mycobacterial phenolic

glycolipid synthesis is regulated by cAMP-dependent lysine acylation of FadD22.

Microbiol (United Kingdom) 163:373–382.

64.   Yang S, Sui S, Qin Y, Chen H, Sha S, Liu X, Deng G, Ma Y. 2022. Protein O-

mannosyltransferase Rv1002c contributes to low cell permeability, biofilm formation

in vitro, and mycobacterial survival in mice. APMIS 130:181–192.

65.   Crellin PK, Kovacevic S, Martin KL, Brammananth R, Morita YS, Billman-Jacobe H,

McConville MJ, Coppel RL. 2008. Mutations in pimE restore lipoarabinomannan

synthesis and growth in a Mycobacterium smegmatis lpqW mutant. J Bacteriol

190:3690–3699.

66.   Butler RE, Smith AA, Mendum TA, Chandran A, Wu H, Lefrançois L, Chambers M,

Soldati T, Stewart GR. 2020. Mycobacterium bovis uses the ESX-1 Type VII secretion

system to escape predation by the soil-dwelling amoeba Dictyostelium discoideum.

ISME J 14:919.

67.   Agapova A, Serafini A, Petridis M, Hunt DM, Garza-Garcia A, Sohaskey CD, de Carvalho

LPS. 2019. Flexible nitrogen utilisation by the metabolic generalist pathogen

713   Mycobacterium tuberculosis. Elife 8.

714 68. Yeh JI, Chinte U, Du S. 2008. Structure of glycerol-3-phosphate dehydrogenase, an

715   essential monotopic membrane enzyme involved in respiration and metabolism. Proc

716   Natl Acad Sci U S A 105:3280.

717 69. Cook GM, Hards K, Dunn E, Heikal A, Nakatani Y, Greening C, Crick DC, Fontes FL,

718   Pethe K, Hasenoehrl E, Berney M. 2017. Drug Discovery & Development: State-of-the-

719   Art and Future Directions" on the topic of "Targets": OXPHOS as a target space for

720   tuberculosis: success, caution, and future directions. Microbiol Spectr 5.

721 70. Bald D, Villellas C, Lu P, Koul A. 2017. Targeting energy metabolism in Mycobacterium

722   tuberculosis, a new paradigm in antimycobacterial drug discovery. MBio 8.

723 71. Garnier T, Eiglmeier K, Camus J-C, Medina N, Mansoor H, Pryor M, Duthoy S, Grondin

724   S, Lacroix C, Monsempe C, Simon S, Harris B, Atkin R, Doggett J, Mayes R, Keating L,

725   Wheeler PR, Parkhill J, Barrell BG, Cole ST, Gordon S V., Hewinson RG. 2003. The

726   complete genome sequence of Mycobacterium bovis. Proc Natl Acad Sci U S A

727   100:7877.

728 72. Larrouy-Maumus G, Biswas T, Hunt DM, Kelly G, Tsodikov O V., De Carvalho LPS.

729   2013. Discovery of a glycerol 3-phosphate phosphatase reveals glycerophospholipid

730   polar head recycling in Mycobacterium tuberculosis. Proc Natl Acad Sci U S A

731   110:11320–11325.

732 73. Gallagher J, Horwill DM. 1977. A selective oleic acid albumin agar medium for the

733   cultivation of Mycobacterium bovis. J Hyg (Lond) 79:155.

734 74. Vordermeier HM, Chambers MA, Cockle PJ, Whelan AO, Simmons J, Hewinson RG.

2002. Correlation of ESAT-6-specific gamma interferon production with pathology in

cattle following Mycobacterium bovis BCG vaccination against experimental bovine

tuberculosis. Infect Immun 70:3026–3032.

75. DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. 2015. TRANSIT--A

Software Tool for Himar1 TnSeq Analysis. PLoS Comput Biol 11.

76. Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. circlize Implements and enhances

circular visualization in R. Bioinformatics 30:2811–2812.

77. R Core Team. R: A Language and Environment for Statistical Computing.

78. Wickham H, Averick M, Bryan J, Chang W, D' L, Mcgowan A, François R, Grolemund G,

Hayes A, Henry L, Hester J, Kuhn M, Lin Pedersen T, Miller E, Bache SM, Müller K,

Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K,

Yutani H. 2019. Welcome to the Tidyverse. J Open Source Softw 4:1686.

 Figure Legends

Figure 1. bTB specific IFN-gamma release in cattle infected with the *M. bovis* Tn-library.

Blood was collected from all 24 animals on the day of infection and 2 weeks later. No

response was detected to either PPD-A or PPD-B antigen stimulation prior to infection

(Figure 1A and Figure 1B, week 0). All animals presented a significant and specific response

to PPD-B compared to PPD-A as determined by a paired T-test using GraphPad Prism (Figure

1C). *** $p \leq 0.001$

Figure 2. Tissue pathology and bacterial load in tissue sites. Six weeks after infection

animals were subjected to post-mortem examination. Gross pathology and evidence of TB-

like granulomas lesions were scored. Data presented is the mean across animals of the total

scores for each tissue group from 24 animals +/- the standard deviation. Lung and thoracic

lymph nodes were observed to contain the highest pathology compared to head and neck

lymph nodes (Figure 2A). For bacterial load estimation, aliquots of macerates were plated

onto modified 7H11 agar containing kanamycin. Colonies were counted after 3-4 weeks

growth. Data are presented as mean CFU/ml per collected tissue group +/- standard

deviation. Lung tissue contained the highest bacterial burden compared to thoracic and

head and neck lymph nodes as determined by one-way ANOVA analysis using GraphPad

Prism (Figure 2b). *** $p \leq 0.001$, ** $p = 0.002$, *$p=0.01$

Figure 3. Fold-changes caused by transposon insertions in the ESX-1 secretion system in

the lungs and lymph nodes of infected cattle. Asterisks indicate that genes had an adjusted

p-value of <0.05 in at least half of the animals. The genes are grouped according to function

as indicated by the colour scheme. The $\log_2$ fold-change are indicated on a yellow to red

scale and present as a dot in the centre of the gene.

Figure 4. Violin plot of normalised $\log_2$ fold changes in gene insertions recovered from

bovine lung or thoracic lymph node tissue samples in selected gene groups. Black bars

indicate overall median of normalized $\log_2$ fold-change among genes in grouping. White bars

indicate mean $\log_2$ fold-change for each gene in the group across all samples in either lung

or lymph node tissue

Figure 5. Comparison of reported $\log_2$ fold-change in *M. bovis*, *M. bovis* BCG and Mtb

transposon insertion sequencing experiments for orthologous genes in the cholesterol

catabolic pathway. Greatest attenuation (most negative $\log_2$ fold-change) is coloured by

darkest red. Studies used for comparison include Mendum et al., (24) and Bellarose et al.,

(9). Grey bars represent genes for which there is no information as they were either ES or

GD in input library or had less than 5 insertions in any TA site in any sample (input and all

output).

Figure 6. Fold-changes caused by transposon insertions in *phoP, phoR* and *fadD26* in the

lungs and lymph nodes of infected cattle. Samples with adjusted p-values (BH-fdr

corrected) < 0.05 are indicated with purple points.

(A)

PPD-A

Week 0   Week 1

OD 490nm

(B)

PPD-B

Week 0   Week 1

OD 490nm

(C)

Post infection PPD IFN-γ Release

***

PPD-A   PPD-B

O.D (490nm)

ESX-1 locus (Mb3894-3913)