Endocytosis is required for access of surface-bound cargo to the flagellar pocket of

trypanosomes

Running title: Characterisation of TbSmee1 (27)

Authors:

Daja Schichler[1], Eva-Maria Spath*[2], Antonia Konle*, Sina Riegler*, Alexandra Klein°, Anna

Seleznev°, Sisco Jung°, Tim Wuppermann[#], Noah Wetterich[#], Alyssa Borges, Elisabeth

Meyer-Natus, Katharina Havlicek[3], Sonia Pérez Cabrera[4], Korbinian Niedermüller, Sara

Sajko[5], Maximilian Dohn, Xenia Malzer, Emily Riemer, Tuguldur Tumurbaatar, Kristina

Djinovic-Carugo[5,6], Gang Dong[7], Christian J. Janzen, Brooke Morriswood.

*/°/[#] These authors contributed equally

Affiliations:

Unless stated otherwise: Department of Cell & Developmental Biology, Biocenter, University

of Würzburg, Würzburg, Germany

1 present address: Cell Biology and Biophysics Unit, EMBL, Heidelberg, Germany

2 present address: Department of Internal Medicine 1, Friedrich-Alexander-University

Erlangen-Nuremberg, Erlangen, Germany

3 Department of Biochemistry and Cell Biology, Max Perutz Labs, University of Vienna, Vienna

Biocenter (VBC) Vienna, Austria

4 Francisco de Vitoria University, Madrid, Spain

5 Department of Structural and Computational Biology, Max Perutz Labs, University of Vienna,

Vienna Biocenter (VBC), Vienna, Austria

6 Department of Biochemistry, Faculty of Chemistry and Chemical Technology, University of

Ljubljana, Ljubljana, Slovenia

7 Center for Medical Biochemistry, Max Perutz Labs, Medical University of Vienna, Vienna

Biocenter (VBC) Vienna, Austria

Email for correspondence: brooke.morriswood@uni-wuerzburg.de

Keywords: Trypanosoma brucei, cytoskeleton, hook complex, TbSmee1, flagellar pocket

Summary: Characterisation of the essential trypanosome protein TbSmee1 suggests that

endocytosis is required for flagellar pocket access of surface-bound cargo

Abstract

All endo- and exocytosis in the African trypanosome *Trypanosoma brucei* occurs at a single subdomain of the plasma membrane. This subdomain, the flagellar pocket, is a small vase-shaped invagination containing the root of the cell's single flagellum. Several cytoskeleton-associated multiprotein complexes are coiled around the neck of the flagellar pocket on its cytoplasmic face. One of these, the hook complex, may affect macromolecule entry into the flagellar pocket lumen. In previous work, knockdown of the hook complex component TbMORN1 resulted in larger cargo being unable to enter the flagellar pocket. In this study, the hook complex component TbSmee1 was characterised in bloodstream form *Trypanosoma brucei* and was found to be essential for cell viability. TbSmee1 knockdown resulted in flagellar pocket enlargement, and impaired access to the pocket membrane by surface-bound cargo. Inhibition of endocytosis by knockdown of clathrin phenocopied TbSmee1 knockdown, suggesting that endocytic activity itself is a prerequisite for the entry of surface-bound cargo into the flagellar pocket.

Introduction

The flagellated parasitic protist *Trypanosoma brucei* lives in the bloodstream of its mammalian hosts, in continuous exposure to the immune system. Endo- and exocytosis are therefore critical to the parasite's survival, as they are used to remove bound antibodies from the cell surface and scavenge macromolecular nutrients from the surroundings (Borst and Fairlamb, 1998; Engstler et al., 2007). Remarkably, all endo- and exocytosis in *T. brucei* occurs at just a single subdomain of the plasma membrane, despite the overall rate of endocytosis being comparable to a professional phagocyte (Engstler et al., 2004; Grunfelder et al., 2003). This subdomain is a small vase-shaped invagination that houses the root of the cell's single flagellum. It is called the flagellar pocket (Halliday et al., 2021; Lacomble et al., 2009).

Fluid-phase or surface-bound cargo entering the flagellar pocket is rapidly internalised by clathrin-mediated endocytosis, and then routed to the endosomal/lysosomal system (Link et al., 2021; Overath and Engstler, 2004). Depletion of clathrin results in a gross enlargement of the flagellar pocket due to an imbalance between endocytosis and exocytosis, and causes cargo accumulation inside the flagellar pocket (Allen et al., 2003). Although endocytosis in *T. brucei* is relatively well-characterised, the exact mechanism(s) by which fluid-phase and surface-bound cargo enter the flagellar pocket is less well-understood.

Coiled around the neck of the flagellar pocket on its cytoplasmic face are a number of poorly-characterised cytoskeleton-associated complexes, which might contribute to these processes

(Esson et al., 2012; Halliday et al., 2019). The flagellar pocket collar demarcates the boundary
between the flagellar pocket and the flagellar pocket neck, and contains the protein TbBILBO1
(Bonhivers et al., 2008; Florimond et al., 2015). The centrin arm is a centrin-containing kinked
rod which appears to be involved in the biogenesis of various cytoskeleton-associated
structures (Selvapandiyan et al., 2007; Shi et al., 2008). The hook complex is a hook-shaped
multiprotein complex that sits atop the flagellar pocket collar and alongside the centrin arm. It
contains the protein TbMORN1 (Morriswood et al., 2009). Previous characterisation of
TbMORN1 revealed that it is essential for the viability of bloodstream form *T. brucei* cells
(Morriswood and Schmidt, 2015). Cells depleted of TbMORN1 have enlarged flagellar
pockets, indicative of an endocytosis defect. In addition, although small endocytic reporters
were still capable of entering the enlarged flagellar pocket, the access of larger
macromolecules such as the lectin concanavalin A (ConA) or bovine serum albumin (BSA)
conjugated to 5 nm gold particles was blocked. On this basis, it was proposed that the hook
complex might be regulating the passage of large macromolecules through the flagellar pocket
neck, and thereby mediating cargo entry into the flagellar pocket.
90
In this study, the hook complex protein TbSmee1 was characterised in bloodstream form *T.*
brucei. TbSmee1 (Tb927.10.8820) was initially identified in a proximity labelling screen using
TbMORN1 as bait, and shown to localise to the shank part of the hook complex (Morriswood
et al., 2013). It was subsequently characterised in procyclic form *T. brucei*, the life cycle stage
found within the tsetse fly vector (Perry et al., 2018). In bloodstream form *T. brucei,* depletion
of TbSmee1 phenocopies TbMORN1, with ConA and BSA unable to access an enlarged
flagellar pocket. Surprisingly, clathrin-depleted cells also exhibited this flagellar pocket access
phenotype, suggesting that endocytosis is required for the entry of surface-bound cargo into
the flagellar pocket.
100
101
Results
TbSmee1 is a cytoskeleton-associated phosphoprotein.
Bioinformatic analysis predicted that TbSmee1 is composed of three structured domains
separated by linker regions (Fig. 1A). Specifically, a multiple sequence alignment of the
TbSmee1 protein with orthologues from other trypanosome species indicated the presence of
three blocks of highly-conserved sequence (Fig. S1). These blocks of conserved sequence
also corresponded to regions of predicted secondary structure (Fig. S2).
109
Two polyclonal antibodies (303, 304) were raised against a recombinant TbSmee1(1-400)
truncation, and a third (508) was raised against two peptides (Fig. 1A, green lines). The

specificity of the antibodies was confirmed using immunoblots of whole-cell lysates obtained

from TbSmee1 RNAi cells and 3xTy1-TbSmee1 endogenous replacement cells. In situ tagging

of the *SMEE1* gene in the 3xTy1-TbSmee1 cells was confirmed by PCR analysis of genomic

DNA (Fig. S3A,B). All three antibodies detected a band of ~ 85 kDa in the immunoblots, which

was strongly depleted in the RNAi (Tet +) samples (Fig. 1B, upper panels. An extra band

corresponding to 3xTy1-TbSmee1 was seen in lysates from the endogenous replacement

(End. repl.) cells). This extra band was also detected using anti-Ty1 antibodies, confirming its

identity (Fig 1B, lower panels).

To investigate TbSmee1 association with the cytoskeleton, cells were fractionated using

nonionic detergent (Fig. 1C). TbSmee1 co-fractionated with the cytoskeleton markers PFR1,2,

confirming that it was cytoskeleton-associated (Fig. 1D). Interestingly, the fuzzy appearance

of the TbSmee1 band observed in immunoblots of whole-cell lysates was reduced or absent

in the fractionation blots (compare Fig. 1B and Fig. 1D). TbSmee1 is heavily phosphorylated

in vivo, and is a substrate and potential binding partner of the mitotic kinase TbPLK (Benz and

Urbaniak, 2019; McAllaster et al., 2015). The existence of different phosphoforms of a protein

is known to cause a fuzzy appearance of bands in gels, so one explanation for this was that

exposure of TbSmee1 to endogenous phosphatases during the extraction step was causing

dephosphorylation.

To investigate whether the band collapse could be attributed to dephosphorylation of

TbSmee1, extracted cytoskeletons were incubated with exogenous phosphatase at various

timepoints prior to immunoblotting. Incubation with exogenous phosphatase resulted in a

progressive collapse of the TbSmee1 band over a 20 min period (Fig. 1E). This band collapse

was not seen when the samples were in the presence of phosphatase inhibitors either at RT

or on ice, or kept on ice (Fig. 1E, last three lanes). The fuzzy appearance of TbSmee1 in

immunoblots could therefore be attributed exclusively to phosphorylation.

TbSmee1 localises to the shank part of the hook complex.

The localisation of endogenous and tagged TbSmee1 protein was analysed in bloodstream

form *T. brucei* cells using immunofluorescence microscopy (Fig. 1F). All three main stages of

the cell division cycle (1K1N, 2K1N, 2K2N) were analysed. In 1K1N cells (i.e. those with a

single kinetoplast, K, and nucleus, N), TbSmee1 localised to a single structure near the

flagellum base, consistent with the position of the hook complex (Fig. 1F, arrow). Interestingly,

in 2K1N and 2K2N cells there was a third subpopulation of TbSmee1 at varying distances

along the cell body in addition to the replicated hook complex (Fig. 1F, arrowheads). This

subpopulation was fainter in 2K2N cells than 2K1N ones. The same TbSmee1 distributions
were seen using all three anti-TbSmee1 antibodies, and also with anti-Ty1 antibodies in
3xTy1-TbSmee1 cells (Fig. S3C). In the endogenous replacement cells, the anti-TbSmee1
and anti-Ty1 signals colocalised (Fig. S3D).
153
To confirm the localisaiton of TbSmee1 at the hook complex, colabelling experiments were
carried out. As expected, TbSmee1 strongly overlapped with the shank part of the hook
complex as labelled by TbMORN1 and TbLRRP1 (Fig. 2A,B). The signals were strongly
correlated (graphs at right). TbSmee1 also strongly overlapped with the hook complex protein
Tb927.10.3010. This 133 kDa protein was one of the top hits in the BioID screen using
TbMORN1 (Morriswood et al., 2013). Consistent with the convention started with TbSmee1,
Tb927.10.3010 was named TbStarkey1 after another one of Captain Hook's pirates. Two anti-
peptide antibodies were generated against TbStarkey1, and their specificity was confirmed by
immunoblotting and immunofluorescence imaging of TbStarkey1 RNAi cells (Fig. S4A,B).
Strong overlap seen between TbSmee1 and TbStarkey1 was seen on the shank part of the
hook complex, and the signals were again strongly correlated (Fig. S4C).
165
As expected, TbSmee1 showed only a partial overlap with TbCentrin4, a marker of the centrin
arm, and little overlap with TbBILBO1, a marker of the flagellar pocket collar (Fig. 2C, D).
Despite the low level of co-occurrence between TbSmee1, TbCentrin4, and TbBILBO1, the
signals were strongly correlated. The high levels of global correlation even when the level of
co-occurrence was low probably reflects the fact that all these structures are closely
juxtaposed in a small volume of intracellular space. In summary, TbSmee1 was localised to
the shank part of the hook complex, and showed strong correlation with both strongly and
partially overlapping markers (Fig. 2E).
174
175
TbSmee1 is transiently associated with the flagellum attachment zone (FAZ) tip.
The flagellum attachment zone (FAZ) is a cytoskeleton-associated apparatus that adheres the
flagellum of *T. brucei* to the cell body (Sunter and Gull, 2016). During replication, a new FAZ
is assembled and grows from its initiation point very close to the hook complex towards the
anterior end of the cell (Sunter et al., 2015; Zhou et al., 2015). It was previously shown that a
tagged version of TbSmee1 is transiently localised to the tip of the new FAZ in replicating
insect-stage *T. brucei* cells, in addition to its hook complex localisation (Perry et al., 2018).
This dynamic localisation contrasts with other hook complex proteins such as TbMORN1,
which are localised exclusively to the hook complex.
185

To investigate whether the endogenous TbSmee1 subpopulation seen in 2K1N and 2K2N
BSF cells corresponded to the FAZ tip, TbSmee1 and the FAZ marker FAZ1 were analysed
at different stages of the cell division cycle. As expected, the TbSmee1 present at the hook
complex showed a partial overlap with the posterior end of the FAZ (Fig. 3A, arrow). The extra
TbSmee1 subpopulation in 2K1N and 2K2N cells overlapped with the tip of the new FAZ (Fig.
3A, arrowheads). The TbSmee1 signal got fainter as the new FAZ elongated, and was quite
faint in the 2K2N stage. TbSmee1 was sometimes faintly observed along the full length of the
new FAZ in replicating cells, and upon careful observation, a TbSmee1 signal could also be
detected at the tip of the old FAZ as well, even in 1K1N cells.
195
Using structured illumination microscopy, TbSmee1 was observed to be slightly in front of the
tip of the new FAZ filament (Fig. 3B). This suggested that TbSmee1 might be present at the
"groove". The groove is a structure involved in remodelling of the microtubule cytoskeleton
during cell replication in bloodstream form *T. brucei* (Hughes et al., 2013). It can be detected
using the monoclonal antibody DOT1. TbSmee1 appeared to be just behind and slightly
enveloped by the DOT1 labelling, confirming its presence at the groove (Fig. 3C). In summary,
in replicating cells TbSmee1 is present at the groove in addition to the hook complex, and
travels in front of the newly-assembling FAZ filament.
204
205
206
The C-terminal part of TbSmee1 is required for targeting to the hook complex.
To determine what parts of TbSmee1 primary structure are responsible for targeting to the
hook complex, a series of truncation constructs based on the predicted domain architecture
of TbSmee1 were designed (Fig. 4A). Cell lines that could inducibly express each of these
truncations with an N-terminal Ty1 tag were generated. The presence of the ectopic
transgenes in the genomes of these cells was confirmed by PCR analysis of genomic DNA
(Fig. S5A). To determine the localisation of each construct, and whether it could associate
with the cytoskeleton, detergent-extracted cells were analysed using immunofluorescence
microscopy.
216
TbSmee1(161-766) correctly localised to the hook complex, indicating that Domain1 is not
necessary for targeting (Fig. 4B). TbSmee1(265-766) also localised correctly, indicating that
the predicted linker region between Domain1 and Domain2 is not necessary for localisation
(Fig. 4B). No other truncations were observed to localise to the hook complex, making
TbSmee1(265-766) the smallest construct to localise correctly. Expression of all TbSmee1
truncations was confirmed by immunoblotting with anti-TbSmee1 and anti-Ty1 antibodies

(Figs 4C, S5A). Quantification of the immunoblots indicated that most were expressed at
around the level of the endogenous protein (Fig. 4D).
225
226   Interestingly, TbSmee1(161-766) also localised to the FAZ tip in replicating cells (Fig. S5B).
227   TbSmee1(265-766) did not localise to the FAZ tip, suggesting that the linker region between
228   domain1 and Domain2 (aa 161-264) might mediate targeting to this structure. In support of
229   this hypothesis, TbSmee1(2-400) was found to localise to the FAZ tip despite not localising to
230   the hook complex (FigS5C).
231
232
TbSmee1 is essential for the viability of BSF *T. brucei*, and its depletion causes gross
enlargement of the flagellar pocket.
The effects of TbSmee1 depletion were analysed using tetracycline-inducible RNAi. Induction
of RNAi (Tet+) resulted in a rapid cessation of population growth after around 24 h (Fig. 5A).
Visual inspection of the stalled populations at 48 h and 72 h post-induction showed widespread
lysis, confirming that TbSmee1 is essential for the viability of bloodstream form cells in vitro.
A shorter timecourse with higher sampling frequency indicated that the growth defect began
after around 20 h of RNAi, and was already clear at 24 h (Fig. 5A, inset).
241
242   TbSmee1 depletion was confirmed at the single-cell level using immunofluorescence
243   microscopy (Fig. 5B). TbSmee1 signal was lost from both the hook complex and FAZ tip. The
244   kinetics of TbSmee1 protein depletion either side of the onset of the growth defect were
245   assessed by immunoblotting. TbSmee1 protein levels were reduced to ~ 20-25% at 15 h and
246   18 h post-induction, with a further reduction to around 10-15% from 21 h onwards (Fig. 5C).
247
The effect of TbSmee1 knockdown on cell cycle progression was assessed by quantifying the
numbers of 1K1N, 2K1N, 2K2N, and abnormal cell types in the same time window (Fig. S6A).
An increase in 2K1N and a decrease in 1K1N cells was visible from 21 h post-induction
onwards, followed by an increase in 2K2N cells at the 24 h timepoint. This indicated that cell
cycle progression was inhibited from around 21 h after induction of RNAi, correlating with the
onset of the growth defect. To summarise, TbSmee1 protein was already significantly depleted
at 15 h, with subsequent effects on population growth and cell cycle progression visible from
around 21 h onwards.
256
The effects of TbSmee1 depletion on other hook complex proteins and also components of
the centrin arm and flagellar pocket collar were systematically evaluated (Fig. S7A-E).
TbSmee1 depletion resulted in a weak effect on TbLRRP1 (Fig. S7C), but no other structural

consequences were observed. Conversely, depletion of TbMORN1 resulted in reductions in
the levels of TbSmee1, TbStarkey1, and potentially also TbLRRP1 (Fig. S7A, F-H). The lack
of observed structural changes to the hook complex, flagellar pocket collar, and centrin arm
following TbSmee1 depletion implied that the lethal phenotype might be due to a loss of
function.
265
The most obvious morphological effect of TbSmee1 depletion was the gradual accumulation
of cells with enlarged flagellar pockets (Fig. 5D). Interestingly, depletion of TbStarkey1 also
frequently resulted in the generation of cells with enlarged flagellar pockets, despite causing
no growth defect (Fig. S6B-D). This means that depletion of three separate hook complex
proteins - TbMORN1, TbSmee1, TbStarkey1 - results in flagellar pocket enlargement, though
with varying magnitudes of effect.
272
273
274 **Flagellar pocket enlargement is an early consequence of TbSmee1 depletion.**
275 Flagellar pocket enlargement is not an uncommon phenotype in bloodstream form RNAi cells,
276 and can be both a direct or an indirect consequence of protein depletion (Ali et al., 2014; Allen
277 et al., 2003; Hall et al., 2004; Hall et al., 2005; Price et al., 2007).
278
To determine whether flagellar pocket enlargement was an early (and therefore probably
direct) consequence of TbSmee1 depletion, a fluorescent, fixable 10 kDa dextran reporter was
used. Dextran is a polysaccharide that traffics in the fluid phase and is well-established as an
endocytic marker. Control (-Tet) and TbSmee1-depleted (+Tet) cells were incubated on ice in
order to block endocytosis (Brickman et al., 1995). The cells were then incubated with the
labelled dextran for 15 min to allow it to enter the flagellar pocket, and then fixed afterwards
(Fig. 6A). The magnitude of the dextran signal should therefore be proportional to flagellar
pocket volume. Visual analysis of the cells confirmed that dextran visibly labelled the flagellar
pocket, with a much greater signal seen for cells with an enlarged flagellar pocket (Fig. 6B).
288
Flow cytometry was then used for high-throughput, unbiased, and quantitative analysis of the
cells (Fig. 6C). At 15 h post-induction, no difference between TbSmee1-depleted cells and
controls was observed. At 18 h and 21 h post-induction a slight "shoulder" on the +Tet traces
became visible, indicating the emergence of cells with higher fluorescence values than seen
in controls. By 24 h post-induction there was a clear hump visible in the traces, indicating a
subpopulation with fluorescence values sometimes two orders of magnitude greater than
those seen in control cells (Fig. 6C, arrow). Quantification of the flow cytometry data from
multiple experiments showed that average fluorescence intensity of the whole + Tet population

was noticeably higher than controls at 21 h and 24 h post-induction (Fig. 6D). Even at 18 h
post-induction, i.e. before the onset of the growth defect, there was already a clear increase
in the average fluorescence intensity in the + Tet population (Fig. 6D, blue arrow). This strongly
suggests that flagellar pocket enlargement is an early and probably direct consequence of
TbSmee1 depletion.
302
303
TbSmee1 depletion results in impaired flagellar pocket access of surface-bound cargo.
It was previously shown that the ability of large cargo to enter the flagellar pocket is affected
after knockdown of the hook complex protein TbMORN1 (Morriswood and Schmidt, 2015).
Specifically, the fluid phase marker 10 kDa dextran accumulates in the enlarged flagellar
pocket of TbMORN1-depleted cells, while larger fluid phase cargo such as BSA-5 nm gold
and large surface-bound cargo such as ConA (which binds to surface glycoproteins) do not
access the flagellar pocket lumen.
311
To test whether TbSmee1 knockdown also impairs flagellar pocket access, TbSmee1 RNAi
cells were incubated with both dextran and ConA to simultaneously monitor the uptake of fluid
phase and surface-bound cargo. The cells were first incubated on ice to block endocytosis,
and then with the reporters (also on ice) to allow ConA to bind and dextran to enter the flagellar
pocket. The cells were then shifted to 37ºC to reactivate endocytosis, and subsequently fixed
and imaged (Fig. 7A).
318
In control cells, both dextran and ConA were observed to strongly overlap in the part of the
cell corresponding to the endosomal/lysosomal system (Fig. 7B, -Tet, arrow). To confirm that
the internalised material was being trafficked to the lysosome, the cells were labelled with
antibodies specific for the lysosomal enzyme p67 (Kelley et al., 1999). The dextran reporter
was not compatible with immunolabelling, but the ConA signal clearly overlapped with the
lysosome marker p67 (Fig. S8, -Tet cells). In TbSmee1-depleted cells, dextran filled the
enlarged flagellar pocket while ConA was restricted to one or two small foci that appeared to
be on the cell surface (Fig. 7B, +Tet, arrow). No overlap was seen between the dextran and
ConA labels, suggesting that ConA was not able to enter the flagellar pocket. In addition, no
overlap was observed between ConA and the lysosome marker p67 (Fig. S8, +Tet cells).
329
To quantify these observations, the cells were grouped into 4 categories: 1, complete overlap
between the two labels; 2, partial overlap between the two labels; 3, no overlap between the
two labels; 4, whole-cell labelling. Control (-Tet) cells all showed complete overlap between

the dextran and ConA, while cells depleted of TbSmee1 for 24 h showed >25% of cells with
partial or no overlap between the cargoes (Fig. 7C).

To further validate these observations, the experiments were repeated using fluorescently-
labelled BSA (Fig. 7D). BSA traffics in the fluid phase and is a physiological cargo, unlike
ConA (Coppens et al., 1987). As expected, in control cells there was strong overlap between
dextran and BSA from the endosomal/lysosomal system (Fig. 7E, -Tet, arrow). In TbSmee1-
depleted cells, there was once again no overlap between the reporters, indicating that BSA
was unable to access the enlarged flagellar pocket. Surprisingly, despite BSA being fluid-
phase cargo there was a punctate signal analogous to that seen with ConA, suggesting that
it could at least partly bind to the cell surface (Fig. 7E, +Tet, arrow). Quantification of the data
showed that the effect on BSA was more pronounced than that seen for ConA, with >55% of
TbSmee1-depleted cells showing no or only partial overlap between the two reporters after 24
h of RNAi (Fig. 7F).

Endocytosis is required for flagellar pocket access of surface-bound cargo.
The inability of surface-bound cargo to enter the flagellar pocket of TbMORN1- and TbSmee1-
depleted cells could either be due to a defect in endocytosis (indicated by the enlargement of
the flagellar pocket), or a secondary effect.

To distinguish between these possibilities, the assays were repeated following clathrin
depletion (Fig. 8). Clathrin is an essential endocytic coat protein and its loss results in a block
in endocytosis. As expected, in control cells there was strong overlap between the dextran
and ConA cargoes (Fig. 8B, -Tet, arrow). In clathrin-depleted cells however, there was again
no overlap observed between the two reporters and ConA did not appear able to access the
enlarged flagellar pocket (Fig. 8B, +Tet, arrow). Quantification of the data showed that >50%
of clathrin-depleted cells showed either partial or no overlap between the two reporters after
19 h of RNAi (Fig. 8C). The same effect was seen in assays using dextran and BSA, with little
to no overlap between the reporters in clathrin-depleted cells (Fig. 8D-F). These results
indicated that inhibition of endocytosis by itself resulted in the failure of either ConA or BSA to
enter the enlarged flagellar pocket.

Discussion
In this study, the hook complex component TbSmee1 was characterised in bloodstream form
T. brucei. It is only the second hook complex protein to be characterised in this life cycle stage

after TbMORN1, and like TbMORN1 it is essential for the viability of the cells in vitro. The
localisation of TbSmee1 was found to be identical to that previously documented in the
procyclic life cycle stage of *T. brucei* (Figs 2, 3). It is constitutively present at the shank part of
the hook complex and probably also the FAZ tip, and additionally localises to the tip of the
growing new FAZ in replicating cells.
375
The analysis of truncation constructs suggested that different parts of the TbSmee1 primary
structure are responsible for targeting to the two locations (Figs 4, S5). Targeting to the hook
complex is mediated by the second and third predicted domains of TbSmee1. It is currently
not clear whether the linker region between these two domains (aa 401-630) is also required,
and this would be an obvious hypothesis to test in future work. Targeting to the FAZ tip appears
to be mediated by the linker region (aa 161-265) between the first and second predicted
domains. This is somewhat surprising, as there are few highly-conserved residues in this
region (Fig. S1). It should be noted that while aa 161-265 have been shown to be necessary
for FAZ tip targeting, they have not yet been shown to be sufficient - this again would be an
obvious future experiment should this line of enquiry be pursued further.
386
Phenotypic characterisation of TbSmee1 showed that its depletion results in the enlargement
of the flagellar pocket with concomitant effects on the entry of macromolecular cargo (Figs 5-
7). A clear limitation of all the RNAi experiments was the inability to specifically isolate cells
with reduced levels of TbSmee1. As the cells were being assayed very soon after the onset
of the growth defect, this meant that all assays were being conducted against a background
of cells exhibiting different and often much lower levels of TbSmee1 depletion. This could be
addressed in future experiments either by using a fluorescent tag on TbSmee1 to enable
sorting of the subpopulation of depleted cells, or by synchronising the cells prior to the
induction of RNAi. Nonetheless, there was still extremely good correlation between the
immunoblotting, growth curve, and cell cycle analysis data in these experiments.
397
At 18 h post-induction of RNAi, TbSmee1 protein levels were reaching their minimum but there
was as yet no observable effect on cell cycle progression or population growth. From around
20/21 h post-induction, when TbSmee1 levels reached 10-15% of controls, cell cycle
progression slowed, resulting in an increase in 2K1N and 2K2N cells. This was mirrored by a
slowing of population growth. Systematic analysis of a number of marker proteins for the hook
complex, the centrin arm, and the flagellar pocket collar showed that none were strongly
affected in terms of their protein levels or localisation by the depletion of TbSmee1. This
argues that the effects seen upon TbSmee1 depletion at these early timepoints are probably
due to a loss of function, rather than due to structural perturbations.

Flow cytometry analysis indicated that the flagellar pocket enlargement began around 18 h
post-induction, just prior to the effects on population growth and cell cycle progression. This
therefore suggests that flagellar pocket enlargement is an early and likely direct consequence
of TbSmee1 depletion. One plausible hypothesis to account for these observations is that the
geometric problems entailed by flagellar pocket enlargement may lead to the subsequent
delay in cell cycle progression and population growth.

The enlargement of the flagellar pocket was accompanied by a cargo access defect previously
documented for TbMORN1 - small fluid-phase cargo such as 10 kDa dextran fills the enlarged
flagellar pocket, while larger cargo such a ConA and BSA is unable to (Morriswood and
Schmidt, 2015). In TbSmee1-depleted cells, BSA was observed in one or more small foci next
to the flagellar pocket. This was surprising, as BSA is known to traffic in the fluid phase and
should not therefore be binding to the cell surface. BSA was however also observed to
associate with the flagellar pocket membrane in an electron microscopy-based study of cargo
uptake (Gadelha et al., 2009). In a classic radiochemical study of cargo uptake in *T. brucei,* it
was observed that "at low concentration, a small adsorbtive component may become
prevailing" in assays using BSA (Coppens et al., 1987). As the experiments conducted here
and in the 2009 paper were using BSA well below physiological concentrations, it seems likely
that these foci represent the small adsorbtive component that was indirectly observed in the
biochemical assays. Therefore, in these assays, the ConA and BSA probes both report on the
behaviour of surface-bound cargo, while the dextran reports on fluid-phase uptake.

Analysis of clathrin-depleted cells indicated that impairment of endocytosis is most likely
responsible for the defect in entry of surface-bound cargo. These data contradict previously-
published observations, where ConA was claimed to accumulate inside the enlarged flagellar
pocket of clathrin-depleted cells (Allen et al., 2003). Re-examination of previously-published
data from clathrin RNAi cells suggests however that the effect has always been present, with
ConA signal (and indeed other cargoes) absent from the flagellar pocket interior and
predominantly concentrated in one or more foci just outside it (see for example Fig. S4 in
(Allen et al., 2003), and Fig. S2 in (Zoltner et al., 2015)). This oversight is understandable,
given that the focus in these clathrin papers was on whether or not cargo endocytosis was
occurring, and not whether the cargo was able to access the flagellar pocket (Mark Field,
personal communication). Importantly, it also means that similar observations have been
made by (at least) two independent groups.

It has been known for some time that surface-bound cargo in the lumen of the flagellar pocket
is associated with the membrane underlying the microtubule quartet (Brickman and Balber,
1990; Gadelha et al., 2009). It has also previously been shown that upon inhibition of
endocytosis, tomato lectin, wheat germ agglutinin, and BSA accumulate on this part of the
flagellar pocket membrane (Gadelha et al., 2009). What is intriguing about the observations
with TbSmee1 and clathrin RNAi cells here is that they show that fluid phase cargo (dextran)
is still able to enter the flagellar pocket even when endocytosis is inhibited, so the effects are
specific to surface-bound cargo. Given that the ConA and BSA material becomes
concentrated into one or more foci that do not overlap with the dextran signal (Figs 7,8), this
suggests that the majority of cargo is not able to enter the flagellar pocket lumen at all when
endocytosis is inhibited.
454
This begs two important questions: why is it that an inhibition of endocytosis apparently
hinders the entry of surface-bound cargo into the flagellar pocket? And why does the depletion
of hook complex proteins - which are spatially removed from the actual sites of clathrin-coated
vesicle formation - cause an endocytosis defect? The prevailing model for several years has
been that hydrodynamic flow, driven by flagellar motility, is responsible for sorting surface-
bound cargo to the posterior end of the cell where the flagellar pocket is located (Engstler et
al., 2007). Once there, the surface-bound material enters the flagellar pocket through a narrow
channel where the flagellar membrane and the flagellar pocket neck membrane are not as
closely apposed (Gadelha et al., 2009).The results here suggest that the posterior sorting
mechanism still functions when endocytosis is inhibited, but that the transit of surface-bound
material through the channel is impeded. The fact that fluid phase cargo still enters the
enlarging flagellar pockets demonstrates that the channel lumen must still be open.
467
An earlier hypothesis for hook complex function was that it was somehow maintaining the
integrity of the channel in the flagellar pocket neck. This remains possible - and determining
the exact site of cargo blockade, as well as imaging the microtubule quartet, will be key goals
for future work - but the fact that the same phenotype is obtained following depletion of clathrin
suggests that the hook complex is instead indirectly affecting endocytosis. This could
potentially be by affecting either the localisation or activity of a number of lipid kinases, at least
two of which are known to localise to the hook complex (Dean et al., 2017; Demmel et al.,
2014). It appears therefore that while hydrodynamic flow may be responsible for concentrating
cargo at the entrance to the flagellar pocket, endocytic activity is required for the entry of the
cargo into the flagellar pocket itself, and that endocytosis may assist or be responsible for
pulling material in through the channel. Exploring these mechanisms is likely to be a
fascinating area for future enquiry.

Materials and Methods

Recombinant protein expression and purification

The TbSmee1(1-400) open reading frame was amplified from *Trypanosoma brucei brucei* strain Lister 427 genomic DNA by PCR. The PCR product was ligated into the p3NH expression vector, which encodes an N-terminal His6 tag, using sequence and ligation-independent cloning (Li and Elledge, 2012). The plasmid was used to transform *E. coli* strain Rosetta II (DE3)pLysS by heat shock, and individual colonies were subsequently grown at 37 °C in the presence of 100 µg/ml kanamycin to an OD600 ~ 0.8–1.0. Recombinant protein expression was induced by the addition of 50 µM IPTG, and the cells were then incubated overnight at 20 °C with shaking. Cells were harvested by centrifugation (5000× g for 30 min). The pooled pellet from 6 L of cell culture was resuspended in 300 ml of lysis buffer (50 mM Hepes pH 7.0, 500 mM NaCl, 20 mM imidazole, 5% glycerol, 0.5% Triton-X, 1 mM TCEP, 200 mM PMSF and protease inhibitor cocktail). Pellet emulsions were first homogenised by mixing on ice using a T 10 basic Ultra-Turrax dispersing instrument (IKA). Final lysis was then achieved with sonication on ice, using 3 cycles of 3 min at 50% strength. Lysates were clarified by centrifugation (18,000× g, 45 min, 4 °C). The lysates were added to a HiTrap Chelating HP 5 ml column (GE Healthcare) equilibrated with buffer A (20 mM Hepes pH 7, 300 mM NaCl, 40 mM imidazole, 2% glycerol, 1 mM TCEP), and eluted with a 100% step gradient of buffer B1 (20 mM Hepes pH 7, 300 mM NaCl, 400 mM imidazole, 2% glycerol, 1 mM TCEP). Selected peak fractions were examined by SDS-PAGE for protein content and purity. Fractions containing a dominant band at approximately 46 kDa were pooled and concentrated using Amicon Ultra centrifugal filter units with 10K pore size (MerckMillipore) according to the manufacturer's instructions. The His6 tag was removed by 3C protease during overnight dialysis in dialysis buffer (20 mM Tris-HCl pH 7, 300 mM NaCl, 2% glycerol, 1 mM DTT). Significant losses were incurred during this step. The TbSmee1(1-400) was applied to a previously equilibrated HiTrap Chelating HP 5 ml column charged with 50 mM $CoCl_2$ and coupled to a GSTrap HP 1 ml column (both GE Healthcare). Buffer A was used for equilibration. TbSmee1(1-1400) was mostly collected from the flow-through and a few initial collected fractions. Selected peak fractions were examined by 15% SDS-PAGE for protein content and purity. Fractions containing a dominant band at approximately 44 kDa were pooled and concentrated in Amicon Ultra centrifugal filter units (10K pore size) according to the manufacturer's instructions. Finally, TbSmee1(1-400) concentrates were applied to a previously equilibrated HiLoad 16/600 Superdex 200 pg column (GE Healthcare) pre-equilibrated in dialysis buffer. Flow speed was adjusted to 1 ml/min and fractions of 1.5 ml

were collected. Fractions corresponding to the targeted chromatographic peak were examined

for protein content by 15% SDS-PAGE, pooled accordingly to their purity, concentrated, and

stored at -80 °C.

Antibody generation and affinity purification

Purified recombinant TbSmee1(1-400) was used for the generation of two polyclonal rabbit

antisera (Eurogentec). Antisera were initially affinity-purified against the TbSmee1(1-400)

antigen, but the neat antisera were later found to show high specificity and were also used. A

third polyclonal antibody was generated against two TbSmee1 peptides (Eurogentec) and

affinity purified using the peptide antigens immonilised on a Sulfolink affinity column

(ThermoFisher). The anti-Starkey1 rabbit polyclonal antibodies were generated and affinity

purified in the same way.

Antibodies

The following primary antibodies have been described previously: rabbit anti-TbMORN1

(Morriswood et al., 2013), rabbit anti-TbBILBO1 (Esson et al., 2012), mouse anti-Ty1 ("BB2")

(Bastin et al., 1996), mouse anti-TbLRRP1 (Zhou et al., 2010), mouse anti-TbCentrin4 ("6C5"

(Ikeda and de Graffenried, 2012)), mouse anti-TbCentrin2 ("2B2H1") (de Graffenried et al.,

2013), mouse anti-TbFAZ1 ("L3B2") (Kohl et al., 1999), mouse anti-PFR1,2 ("L13D6") (Kohl

et al., 1999), mouse anti-FAZ filament ("DOT1")(Woods et al., 1989). The following antibodies

came from commercial sources: goat anti-rabbit(IRDye800CW) (LI-COR), goat anti-mouse

(IRDye680LT) (LI-COR), goat anti-rabbit and anti-mouse antibodies conjugated to AlexaFluor

dyes (Molecular Probes).

Cell culture

Wildtype 427 (monomorphic) BSF cells were cultured in HMI-9 medium (Hirumi and Hirumi,

1989) supplemented with 10 % foetal bovine serum (FBS), 100 U/ml penicillin and 0.1 mg/ml

streptomycin at 37°C and 5% CO_2. The SM (single marker) cells (Wirtz et al., 1999) were

cultured in the presence of G418 (2.5 µg/ml). Population density was monitored using a Z2

Coulter Counter (Beckman Coulter), and kept below $2x10^6$ cells/ml.

Generation of transgenic cell lines

The Ty1-TbSmee1 endogenous replacement cell line was generated by transfection of 427

cells with a targeting fragment containing 285 bp of the TbSmee1 5' untranslated region

(UTR), a blasticidine resistance gene, the intergenic region of alpha- and beta-tubulin, the

sequence for a triple Ty1-tag and the first 399 bp of the TbSmee1 open reading frame (ORF)

without the start codon. Clones were selected by growth in medium containing 5 µg/ml

blasticidine. RNAi target sequences were chosen using RNAit (Redmond et al., 2003). TbSmee1 RNAi cells were generated by cloning the RNAi target sequence into the pGL2084 plasmid (Jones et al., 2014) and then transfecting 2T1 cells with the linearised plasmid (Alsford et al., 2005). Clones were selected by growth in medium containing phleomycin (2.5 μg/ml) and hygromycin (5 μg/ml). TbStarkey1 RNAi cells and TbCHC RNAi cells were generated by cloning the relevant RNAi target sequence into the p2T7_TAblue plasmid (Alibu et al., 2005) and then transfecting SM cells with the linearised plasmid. Clones were selected by growth in medium containing G418 (2.5 μg/ml) and hygromycin (5 μg/ml). Ty1-TbSmee1 truncation constructs were cloned into the pLew100_v5-Hyg plasmid using in vivo assembly (Watson and Garcia-Nafria, 2019). SM cells were transfected with the linearised plasmid. Clones were selected by growth in medium containing G418 (2.5 μg/ml) and hygromycin (5 μg/ml). For transfection >2.5×10^7 cells of the parental strain were washed and resuspended in 100 μl transfection buffer (90 mM Na_2PO_4, 5 mM KCl, 0.15 mM $CaCl_2$, 50 mM HEPES [4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid]-NaOH, pH 7.3) containing 10 μg DNA and transfected by electroporation using an AMAXA Nucleofector® Device (Lonza) with program "X-001 free choice". Transfected cells were incubated in 50 ml HMI-9 medium without selection overnight. The next day, drug selection was applied and clones were selected by limiting dilution. At least three separate clones of all cell lines were isolated to control for biological variability. Integration of targeting fragments at endogenous loci or the presence of Ty1-TbSmee1 truncation constructs in the genome were confirmed by PCR analysis of genomic DNA. Genomic DNA was isolated using a DNeasy Bloody & Tissue kit (QIAGEN) and relevant products were amplified by PCR. All cloned constructs used for cell line generation had their DNA sequence confirmed by sequencing.

Immunoblotting

For generation of dephosphorylated whole-cell lysates, cells were harvested by centrifugation (1000xg, 10 min) and the cell pellet was resuspended in 1 ml vPBS (PBS, 46 mM sucrose, 10 mM glucose) containing EDTA-free protease inhibitors (Roche). The washed cells were pelleted by centrifugation (750xg, 4 min). The cells were then resuspended in lysis buffer (0.5% IGEPAL, 0.1M PIPES-NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor cocktail) to a final concentration of $4x10^5$ cells/μl and incubated for 15 min at RT on an orbital mixer to allow dephosphorylation to occur. SDS-loading buffer was then added to a final concentration of $2x10^5$ cells/μl, the samples were further denatured by boiling (100°C, 10 min), and then stored at -20°C. SDS-PAGE was carried out using a Mini-Protean Tetra Cell (Bio-Rad), and protein transfer to nitrocellulose membranes using a Mini-Trans blot cell (Bio-Rad). Protein transfer and equal loading was confirmed using REVERT

total protein stain (LI-COR) according to the manufacturer's instructions. Membranes were

blocked using blocking buffer (PBS, 0.3% Tween 20, 10% milk)(30 min, RT, rocker). The

membranes were then incubated in primary antibodies diluted in blocking buffer (1 h, RT,

roller). After three washes in immunoblot buffer (PBS, 0.3% Tween 20) the membranes were

incubated with IRDye-conjugated secondary antibodies diluted in immunoblot buffer (1 h, RT,

roller). After another three washes in immunoblot buffer, the membranes were visualised using

an Odyssey CLx (LI-COR). Background subtraction, qualitative analysis, normalisation

relative to total protein stain, and quantification were carried out using Image Studio Lite 5.2

and Empiria Studio 1.1 (LI-COR). Dot plots were generated using the web app PlotsOfData

(Postma and Goedhart, 2019).

In vitro phosphatase assays

427 BSF cells were grown to approximately 1.5×10^6 cells/ml and harvested by centrifugation

(1,000xg, 10 min, 4 °C). The cells were washed in 1 ml ice-cold wash buffer (0.1M PIPES-

NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, phosphatase Inhibitor Cocktail 2)

and pelleted by centrifugation (750xg, 3 min, 4 °C). For extraction of the cytoskeletons, the

cell pellet was resuspended in 1ml ice-cold extraction buffer (0.5% IGEPAL, 0.1M PIPES-

NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor

cocktail, phosphatase Inhibitor Cocktail 2) and incubated for 15 min on ice. The cell

suspension was inverted every 5 min. The cytoskeletons were separated from the cytoplasm

by centrifugation (750xg, 3 min, 4 °C). The cytoskeletons were washed with 0.5 ml ice-cold 1x

NEBuffer for PMP supplemented with 1 mM $MnCl_2$ and pelleted by centrifugation (750xg, 3

min, 4 °C). After resuspension in 375 µl ice-cold 1x NEBuffer for PMP supplemented with 1

mM $MnCl_2$, all following samples were taken from this stock. An input sample (0 min) of 40 µl

was taken and added to 20 µl of SDS-loading buffer. Two control samples were taken, 0.4 µl

Phosphatase Inhibitor Cocktail 2 was added, and incubated on ice and at 26°C, respectively,

for 20 min. 20 µl of SDS-loading buffer was then added. To assay for dephosphorylation, 130

µl were taken from the stock and 1 µl lambda phosphatase (400 U) was added. This sample

and the remainder from the stock were incubated at 26 °C. After 1/2/3/5/10/20 min 20 µl

samples were taken from each and were added to 10 µl SDS-loading buffer. All the samples

were boiled at 104 °C for 10 min and stored at -20 °C. Samples were analysed by

immunoblotting.

Fractionation

50 ml 427 BSF cells were grown to approximately 1.5×10^6 cells/ml and then harvested by

centrifugation (1000xg, 10 minutes, 4°C). The cell pellet was resuspended in 1 ml vPBS,

transferred to a microfuge tube, and the cells again pelleted by centrifugation (750xg, 4 min,
4°C). The supernatant was discarded and the centrifugation step was repeated to remove
remaining supernatant. The cell pellet was resuspended in 200 µl extraction buffer (0.5%
IGEPAL, 0.1M PIPES-NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free
protease inhibitor cocktail) and incubated for 15 minutes at RT in an orbital mixer. A 5% input
sample was taken (10µl), put into a new microfuge tube and left on ice. The extracted cells
were fractionated by centrifugation (3400xg, 2 minutes, 4°C) and the supernatant transferred
to a fresh microfuge tube and the exact volume noted. The tube containing the extracted cells
was centrifuged again at the same settings and this second residual supernatant discarded.
The cytoskeleton pellet was then resuspended in 200 µl extraction buffer. 5% samples of
supernatant and pellet fractions were taken and analysed by immunoblotting. Equal fractions
of I, SN, P were loaded in each lane (I ~ 1.4x10^6 cells).

Preparation of samples for immunofluorescence microscopy

Coverslips were washed in 70% ethanol and then incubated with 0.01% poly-L-lysine in a 24-
well plate (>20 min, RT) and left to dry. 2x10^6 cells were taken per coverslip and transferred
to 15 ml Falcon tubes. The cells were pelleted by centrifugation (1000xg, 1 min per ml of liquid,
RT) in a swing-bucket centrifuge. The supernatant was removed, and the cell pellet was gently
resuspended in 1 ml ice-cold vPBS + Complete. The cells were again pelleted by
centrifugation (1000xg, 2 min, RT) and subsequently resuspended in 1 ml ice-cold vPBS +
Complete and directly added to the coverslips. The cells were attached to the coverslips by
centrifugation (1000xg, 1 min, RT), and attachment was confirmed visually. The attached cells
were then incubated in 1 ml ice-cold extraction buffer (0.5% IGEPAL 0.1M PIPES-NaOH pH
6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor cocktail) (5 min,
on ice). The extracted cells were washed two times with 1 ml vPBS + Complete and then fixed
in 1 ml ice-cold 99.9% methanol (30 min, -20°C). The fixed cells were rehydrated using 1 ml
PBS. The coverslips were blocked in 1 ml 3% BSA in PBS (30 min, RT), and sequentially
incubated with clarified primary and secondary antibodies diluted in PBS (1 h, RT, humidified
chamber for each) with three PBS washing steps (3 x 5 min, RT, rocker) after each incubation.
After the final wash, glass slides were cleaned with 70% ethanol and a spot of DAPI-
Fluoromount G (Southern Biotech) was placed on the surface. The coverslips were rinsed in
ddH$_2$O, carefully dried, and then mounted. For preparation of samples for SIM imaging,
different coverslips (high precision, No. 1.5 H) were used. Cells were harvested as above,
washed, and fixed in 4% paraformaldehyde solution (10 min, ice then 30 min, RT). The fixed
cells were washed, attached to coverslips, permeabilised (0.25% Triton X-100 in PBS; 5 min,
RT), and then labelled and mounted as described above.

Fluorescence microscopy

Images were acquired using a DMI6000B widefield microscope (Leica Microsystems, Germany) with a HCX PL APO CS objective (100x, NA = 1.4, Leica Microsystems, Germany) and Type F Immersion Oil (refractive index = 1.518, Leica Microsystems, Germany). The microscope was controlled using LAS-X software (Leica). Samples were illuminated with an EL6000 light source (Leica) containing a mercury short-arc reflector lamp (HXP-R120W/45C VIS, OSRAM, Germany). Excitation light was selected by using Y3 (545/25 nm), GFP (470/40 nm), and A4 (360/40 nm) bandpass filter cubes (Leica Microsystems, Germany). The power density, measured at the objective focal plane with a thermal sensor head (S175C, Thorlabs), was respectively 0.749 ± 0.086, 0.557 ± 0.069, 0.278 ± 0.076 W/cm^2 for the three filters. Emitted light was collected at ranges of 605/70 (Y3), 525/50 nm (GFP), and 470/40 nm (DAPI) respectively. The individual exposure times and camera gains were adjusted according to the different samples. RNAi samples (control and depleted) were imaged using identical settings. Differential interference contrast (DIC) was used to visualise cell morphology. 3D recording of each field of view was obtained using 40 Z-slices (step size = 0.21 μm). Fields of view were selected in the DIC channel in order to blind the user to the fluorescence signal and subjectively select for cells with optimum morphology. Images were captured using a DFC365 FX monochrome CCD camera (Leica, 6.45 μm pixel size). SIM images were acquired using an Elyra S.1 SIM microscope (Zeiss) and ZEN software (Zeiss).

Fluorescence microscopy image processing and analysis

Processing was carried out using FIJI (Schindelin et al., 2012) and a custom macro for the generation of maximum-intensity z-projections with single DIC z-slices overlaid. Overlaps between two proteins were confirmed in individual z-slices (thickness: 210 nm). The plugin ScientiFig was utilised to create the collage and adding the inserts (Aigouy and Mirouse, 2013). For global correlation analysis between TbSmee1 and other flagellar pocket collar and/or hook complex associated proteins, 1K1N cells were selected using both DIC and DAPI channels. 2D sum slices projections were prepared for each stack of both green and red channels. The projections were clipped to 8-bit depth and a convoluted background subtraction was applied. All resultant 2D images were analysed individually, without channel overlay, and each selected cell had the intensity values of both channels acquired from the histogram. In order to check for data distribution and to determine the suitable correlation method (Aaron et al., 2018; Dunn et al., 2011), the intensity values were further analyzed in R version 4.1.2 (Team, 2021) using the packages ggplot2 (H, 2016) and openxlsx (Schauberger, 2021). The normality was checked by visual inspection of q-q plots and the correlation was tested using Spearman's rank correlation. A scatter plot using the intensity values (green x red) was generated to each protein against Smee1. To visualise the variation of the correlation

within the sample population, a scatter plot of the correlation coefficient obtained per individual
cell was prepared.

Growth curves
RNAi cells were seeded at the required starting concentration in a volume of 22 ml and divided
into two 10 ml aliquots in separate flasks. Tetracycline was added to a final concentration of 1
µg/ml in one flask to induce RNAi, and refreshed every 24h. The population density of the
control and induced cells was measured every 24h over a time course of 72h, or every hour
over a time course of 8h, using a Z2 Coulter Counter (Beckman Coulter). Depletion of the
target protein was confirmed in every experiment by immunoblotting of whole-cell lysates.

Cell division cycle analysis
An aliquot of 10^6 cells was taken, and the cells were fixed directly in media by addition of
isothermal 25% glutaraldehyde to a final concentration of 2.5% (10 min, 37°C, gentle mixing).
The cells were then pelleted by centrifugation (750xg, 10 min). The cell pellet was
resuspended in 0.5 ml 2.5% glutaraldehyde in PBS, transferred to a microfuge tube, and
incubated at RT (30 min, gentle mixing). The cells were pelleted again by centrifugation
(750xg, 4 min), and the cell pellet resuspended in 500 µl PBS. The cells were then added to
the coverslips inside the 24-well plate and attached by centrifugation (1000xg, 1 min, RT). The
coverslips were mounted on poly-L-lysine-coated slides using DAPI-Fluoromount G (Southern
Biotech). Imaging was as described for fluorescence microscopy above, using DAPI and DIC
channels only. Cell division cycle stages (1K1N, 2K1N, 2K2N) were manually quantified from
maximum intensity projections of the DAPI signal overlaid with single DIC z-slices. Depletion
of the target protein at each timepoint was confirmed by immunoblotting of whole-cell lysates
from the same experiments.

Preparation and imaging of electron microscopy samples
Induced and uninduced RNAi cells were grown for 24 h to a density of 1-2x10^6 cells/ml in 50
ml medium and harvested by centrifugation (1000xg, 10 min, RT). The supernatant was
removed to 4 ml and 4 ml FBS was added. The cells were pelleted again (1000xg, 10 min,
RT) and the supernatant was removed to 200 µl. The cells were resuspended in the
supernatant, and the suspension was then transferred to a PCR tube and pelleted by
centrifugation (1,600xg, 10s, RT). The cells were then transferred into a carrier with a closed
lid to avoid air inclusions. High pressure freezing (HPF) was started immediately (Leica EM
HPM100). After HPF the samples were transferred to an AFS (Leica EM AFS2) for freeze
substitution and progressive lowering of temperature. Low temperature embedding and
polymerisation of Epon raisin (DDSA, MNA, Epon812, 2,4,6

Tris(dimethylaminomethyl(phenol))) were then carried out. Ultra-thin cuts (60 nm) were carried

out with an ultramicrotome (Leica EM UC7/FC7) and were placed on slotted grids. For

contrasting they were incubated in 2 % uranyl acetate for 8 min. Afterwards the grids were

washed 3x in ddH$_2$O (boiled to remove CO$_2$) and incubated for 5 min on 50 % Reynold's lead

citrate in a petri dish with NaOH tablets. The grids were again washed 2x in ddH$_2$O. A 200 kV

transmission electron microscope (Jeol, JEM-2100) with a TemCam F416 4k x 4k camera

(Tietz Video and Imaging Processing Systems) and EMMenu 4.0.9.31 software (Tietz) were

used. Uninduced control cells were viewed at a magnification of 12,000x, induced cells were

viewed at a magnification of 8,000x.

Measurement of flagellar pocket enlargement

Induced and uninduced RNAi cells at a concentration of ~5×10^6 cells/ml were harvested by

centrifugation (1000xg, 4 ºC). The cells were resuspended in 45 µl ice-cold vPBS + protease

inhibitors and incubated on ice to block endocytosis (10 min, ice). 5 µl labelled dextran (10

kDa, 50 mg/ml stock) was added and mixed by flicking. The mixture was then incubated to

allow dextran to enter and fill the flagellar pocket (15 min, on ice, dark). At the end of the

incubation, 1 ml ice-cold vPBS + protease inhibitors was added and the cells pelleted by

centrifugation (750 x*g*, 2 min, 4 ºC). The cell pellet was resuspended in 0.5 ml vPBS + protease

inhibitors, and the cells fixed by addition of 0.5 ml 8 % pfa and 0.1 % glutaraldehyde solution

in vPBS + protease inhibitors (20 min, on ice then 60 min, RT). The fixed cells were pelleted

by centrifugation (750 x*g*, 2 min) and washed twice with 1 ml vPBS + protease inhibitors. The

dextran signal was immediately measured by flow cytometry using a FACSCalibur (BD

Biosciences) running CellQuest ProTM Software (BD Biosciences). Later processing and

analysis was carried out using FlowJo 10.8.1. Dot plots were created using Plots of Data

(CITATION). After flow cytometry, the remaining cells were attached to coverslips by

centrifugation, mounted on clean glass slides using DAPI-Fluoromount G (Southern biotech),

and imaged using fluorescence microscopy on the same day to confirm flagellar pocket

labelling. Depletion of TbSmee1 was confirmed in every experiment by immunoblotting of

whole-cell lysates.

Cargo uptake assays

2x10^6 cells per sample were harvested by centrifugation (1000 xg, 10 min, 4 ºC) and washed

in 1 ml ice-cold vPBS. The washed cells were pelleted by centrifugation (750 xg, 2 min, 4 ºC),

and resuspended in 100 µl ice-cold vPBS. The cells were then incubated at low temperature

(20 min, ice) to block endocytosis. During this incubation, the labelled cargoes were prepared.

The 50 mg/ml dextran aliquot (10 kDa AF488 conjugate, Molecular Probes) was mixed using

a sonicator bath (10 min, 37 Hz) and then vortexed. The 0.5 mg/ml ConA aliquot (TMR

conjugate, Molecular Probes) was clarified by centrifugation (11,000 xg, 10 min, 4 ºC). Both
cargoes were kept on ice and in the dark. At the end of the incubation, 10 µl of the 50 mg/ml
dextran and 2 µl of the 0.5 mg/ml ConA were added to the chilled cells, mixed by flicking, and
incubated at low temperature (15 min, on ice, in the dark). T=0 samples were quenched and
fixed at this point; t=30 samples had an additional incubation to allow endocytosis of cargo (30
min, 37 ºC, in the dark) before quenching and fixing. To quench samples, 1 ml ice-cold vPBS
was added, and the cells were pelleted by centrifugation (750 xg, 2 min, 4 ºC). The pelleted
cells were resuspended in 50 µl vPBS by flicking, and then fixed by addition of 0.5 ml ice-cold
fix solution (4% paraformaldehyde solution, 0.1% glutaraldehyde in vPBS) (20 min, on ice, in
the dark, then 30 min, RT, in the dark). The fixed cells were pelleted by centrifugation (750 xg,
2 min, RT), washed in 2 ml vPBS, resuspended in 1 ml vPBS, and then attached to poly-L-
lysine-coated coverslips by centrifugation. Coverslips were mounted on glass slides using
DAPI-Fluoromount G (Southern Biotech) and imaged immediately. Assays using BSA
(AlexaFluor555 conjugate, Molecular Probes) were done in the same way. The 20 mg/ml BSA
aliquot was clarified by centrifugation (750 xg, 1 min, RT)), and 3 µl was added to the
cells/dextran mixture. Depletion of TbSmee1 was confirmed by immunoblotting of whole-cell
lysates in every experiment.
786

Acknowledgements
Antibodies were obtained from the following sources: Cynthia He (National University of
Singapore) provided the anti-Ty1 (BB2) and anti-TbLRRP1 antibodies; Chris de Graffenried
(Brown University) provided the anti-TbCentrin2 (2B2H1) and anti-TbCentrin4 (6C5)
antibodies, Keith Gull (University of Oxford) provided the anti-FAZ1 (L3B2), anti-PFR1,2
(L13D6) and DOT1 antibodies. Nadine Weisert provided assistance with cloning. Markus
Engstler provided support and advice. Balázs Szöőr (University of Edinburgh) provided input
on the phosphatase assays. Mark Field (University of Dundee) provided extremely helpful
advice and context on earlier clathrin RNAi work. Tim Krüger provided input and assistance
with imaging. Helena Jambor (TU Dresden) advised on data visualisation. Graham Warren
(Max Perutz Laboratories) hosted the preliminary experiments for this project. Susanne
Kramer and Manfred Alsheimer provided feedback on the manuscript.
799

800 **Competing interests**
801 There are no competing interests to be declared.
802

Funding
Funding in the initial phase was from the Austrian Science Fund (FWF) grant P27016. Later
funding was from the German Research Foundation (DFG) grant Mo 3188/2-1.

References

Aaron, J. S., Taylor, A. B. and Chew, T. L. (2018). Image co-localization - co-occurrence
versus correlation. *Journal of cell science* **131**.

Aigouy, B. and Mirouse, V. (2013). ScientiFig: a tool to build publication-ready scientific
figures. *Nat Methods* **10**, 1048.

Ali, M., Leung, K. F. and Field, M. C. (2014). The ancient small GTPase Rab21 functions in
intermediate endocytic steps in trypanosomes. *Eukaryotic cell* **13**, 304-19.

Alibu, V. P., Storm, L., Haile, S., Clayton, C. and Horn, D. (2005). A doubly inducible system
for RNA interference and rapid RNAi plasmid construction in Trypanosoma brucei. *Molecular*
and biochemical parasitology **139**, 75-82.

Allen, C. L., Goulding, D. and Field, M. C. (2003). Clathrin-mediated endocytosis is essential
in Trypanosoma brucei. *The EMBO journal* **22**, 4991-5002.

Alsford, S., Kawahara, T., Glover, L. and Horn, D. (2005). Tagging a T. brucei RRNA locus
improves stable transfection efficiency and circumvents inducible expression position effects.
Molecular and biochemical parasitology **144**, 142-8.

Bastin, P., Bagherzadeh, Z., Matthews, K. R. and Gull, K. (1996). A novel epitope tag
system to study protein targeting and organelle biogenesis in Trypanosoma brucei. *Molecular*
and biochemical parasitology **77**, 235-9.

Benz, C. and Urbaniak, M. D. (2019). Organising the cell cycle in the absence of
transcriptional control: Dynamic phosphorylation co-ordinates the Trypanosoma brucei cell
cycle post-transcriptionally. *PLoS Pathog* **15**, e1008129.

Bonhivers, M., Nowacki, S., Landrein, N. and Robinson, D. R. (2008). Biogenesis of the
trypanosome endo-exocytotic organelle is cytoskeleton mediated. *PLoS biology* **6**, e105.

Borst, P. and Fairlamb, A. H. (1998). Surface receptors and transporters of Trypanosoma
brucei. *Annu Rev Microbiol* **52**, 745-78.

Brickman, M. J. and Balber, A. E. (1990). Trypanosoma brucei rhodesiense bloodstream
forms: surface ricin-binding glycoproteins are localized exclusively in the flagellar pocket and
the flagellar adhesion zone. *The Journal of protozoology* **37**, 219-24.

Brickman, M. J., Cook, J. M. and Balber, A. E. (1995). Low temperature reversibly inhibits
transport from tubular endosomes to a perinuclear, acidic compartment in African
trypanosomes. *Journal of cell science* **108 (Pt 11)**, 3611-21.

Coppens, I., Opperdoes, F. R., Courtoy, P. J. and Baudhuin, P. (1987). Receptor-mediated
endocytosis in the bloodstream form of Trypanosoma brucei. *J Protozool* **34**, 465-73.

de Graffenried, C. L., Anrather, D., Von Raussendorf, F. and Warren, G. (2013). Polo-like

kinase phosphorylation of bilobe-resident TbCentrin2 facilitates flagellar inheritance in

Trypanosoma brucei. *Molecular biology of the cell* **24**, 1947-63.

Dean, S., Sunter, J. D. and Wheeler, R. J. (2017). TrypTag.org: A Trypanosome Genome-

wide Protein Localisation Resource. *Trends Parasitol* **33**, 80-82.

Demmel, L., Schmidt, K., Lucast, L., Havlicek, K., Zankel, A., Koestler, T., Reithofer, V.,

de Camilli, P. and Warren, G. (2014). The endocytic activity of the flagellar pocket in

Trypanosoma brucei is regulated by an adjacent phosphatidylinositol phosphate kinase.

Journal of cell science **127**, 2351-64.

Dunn, K. W., Kamocka, M. M. and McDonald, J. H. (2011). A practical guide to evaluating

colocalization in biological microscopy. *Am J Physiol Cell Physiol* **300**, C723-42.

Engstler, M., Pfohl, T., Herminghaus, S., Boshart, M., Wiegertjes, G., Heddergott, N. and

Overath, P. (2007). Hydrodynamic flow-mediated protein sorting on the cell surface of

trypanosomes. *Cell* **131**, 505-15.

Engstler, M., Thilo, L., Weise, F., Grunfelder, C. G., Schwarz, H., Boshart, M. and

Overath, P. (2004). Kinetics of endocytosis and recycling of the GPI-anchored variant surface

glycoprotein in Trypanosoma brucei. *Journal of cell science* **117**, 1105-15.

Esson, H. J., Morriswood, B., Yavuz, S., Vidilaseris, K., Dong, G. and Warren, G. (2012).

Morphology of the trypanosome bilobe, a novel cytoskeletal structure. *Eukaryotic cell* **11**, 761-

72.

Florimond, C., Sahin, A., Vidilaseris, K., Dong, G., Landrein, N., Dacheux, D., Albisetti,

A., Byard, E. H., Bonhivers, M. and Robinson, D. R. (2015). BILBO1 Is a Scaffold Protein

of the Flagellar Pocket Collar in the Pathogen Trypanosoma brucei. *PLoS pathogens* **11**,

e1004654.

Gadelha, C., Rothery, S., Morphew, M., McIntosh, J. R., Severs, N. J. and Gull, K. (2009).

Membrane domains and flagellar pocket boundaries are influenced by the cytoskeleton in

African trypanosomes. *Proceedings of the National Academy of Sciences of the United States*

of America **106**, 17425-30.

Grunfelder, C. G., Engstler, M., Weise, F., Schwarz, H., Stierhof, Y. D., Morgan, G. W.,

Field, M. C. and Overath, P. (2003). Endocytosis of a glycosylphosphatidylinositol-anchored

protein via clathrin-coated vesicles, sorting by default in endosomes, and exocytosis via

RAB11-positive carriers. *Molecular biology of the cell* **14**, 2029-40.

H, W. (2016). ggplot2: Elegant Graphics for Data Analysis: Springer-Verlag New York.

Hall, B., Allen, C. L., Goulding, D. and Field, M. C. (2004). Both of the Rab5 subfamily small

GTPases of Trypanosoma brucei are essential and required for endocytosis. *Molecular and*

biochemical parasitology **138**, 67-77.

Hall, B. S., Smith, E., Langer, W., Jacobs, L. A., Goulding, D. and Field, M. C. (2005).
Developmental variation in Rab11-dependent trafficking in Trypanosoma brucei. *Eukaryotic*
cell **4**, 971-80.

Halliday, C., Billington, K., Wang, Z., Madden, R., Dean, S., Sunter, J. D. and Wheeler, R.
J. (2019). Cellular landmarks of Trypanosoma brucei and Leishmania mexicana. *Molecular*
and biochemical parasitology **230**, 24-36.

Halliday, C., de Castro-Neto, A., Alcantara, C. L., Cunha, E. S. N. L., Vaughan, S. and
Sunter, J. D. (2021). Trypanosomatid Flagellar Pocket from Structure to Function. *Trends*
Parasitol **37**, 317-329.

Hirumi, H. and Hirumi, K. (1989). Continuous cultivation of Trypanosoma brucei blood stream
forms in a medium containing a low concentration of serum protein without feeder cell layers.
J Parasitol **75**, 985-9.

Hughes, L., Towers, K., Starborg, T., Gull, K. and Vaughan, S. (2013). A cell-body groove
housing the new flagellum tip suggests an adaptation of cellular morphogenesis for parasitism
in the bloodstream form of Trypanosoma brucei. *Journal of cell science* **126**, 5748-57.

Ikeda, K. N. and de Graffenried, C. L. (2012). Polo-like kinase is necessary for flagellum
inheritance in Trypanosoma brucei. *Journal of cell science* **125**, 3173-84.

Jones, N. G., Thomas, E. B., Brown, E., Dickens, N. J., Hammarton, T. C. and Mottram,
J. C. (2014). Regulators of Trypanosoma brucei cell cycle progression and differentiation
identified using a kinome-wide RNAi screen. *PLoS Pathog* **10**, e1003886.

Kelley, R. J., Alexander, D. L., Cowan, C., Balber, A. E. and Bangs, J. D. (1999). Molecular
cloning of p67, a lysosomal membrane glycoprotein from Trypanosoma brucei. *Molecular and*
biochemical parasitology **98**, 17-28.

Kohl, L., Sherwin, T. and Gull, K. (1999). Assembly of the paraflagellar rod and the flagellum
attachment zone complex during the Trypanosoma brucei cell cycle. *J Eukaryot Microbiol* **46**,
105-9.

Lacomble, S., Vaughan, S., Gadelha, C., Morphew, M. K., Shaw, M. K., McIntosh, J. R.
and Gull, K. (2009). Three-dimensional cellular architecture of the flagellar pocket and
associated cytoskeleton in trypanosomes revealed by electron microscope tomography.
Journal of cell science **122**, 1081-90.

Li, M. Z. and Elledge, S. J. (2012). SLIC: a method for sequence- and ligation-independent
cloning. *Methods Mol Biol* **852**, 51-9.

Link, F., Borges, A. R., Jones, N. G. and Engstler, M. (2021). To the Surface and Back:
Exo- and Endocytic Pathways in Trypanosoma brucei. *Front Cell Dev Biol* **9**, 720521.

McAllaster, M. R., Ikeda, K. N., Lozano-Nunez, A., Anrather, D., Unterwurzacher, V.,
Gossenreiter, T., Perry, J. A., Crickley, R., Mercadante, C. J., Vaughan, S. et al. (2015).
Proteomic identification of novel cytoskeletal proteins associated with TbPLK, an essential

regulator of cell morphogenesis in Trypanosoma brucei. *Molecular biology of the cell* **26**, 3013-
29.

Morriswood, B., Havlicek, K., Demmel, L., Yavuz, S., Sealey-Cardona, M., Vidilaseris, K.,
Anrather, D., Kostan, J., Djinovic-Carugo, K., Roux, K. J. et al. (2013). Novel Bilobe
Components in Trypanosoma brucei Identified Using Proximity-Dependent Biotinylation.
Eukaryotic cell **12**, 356-67.

Morriswood, B., He, C. Y., Sealey-Cardona, M., Yelinek, J., Pypaert, M. and Warren, G.
(2009). The bilobe structure of Trypanosoma brucei contains a MORN-repeat protein.
Molecular and biochemical parasitology **167**, 95-103.

Morriswood, B. and Schmidt, K. (2015). A MORN Repeat Protein Facilitates Protein Entry
into the Flagellar Pocket of Trypanosoma brucei. *Eukaryot Cell* **14**, 1081-93.

Overath, P. and Engstler, M. (2004). Endocytosis, membrane recycling and sorting of GPI-
anchored proteins: Trypanosoma brucei as a model system. *Mol Microbiol* **53**, 735-44.

Perry, J. A., Sinclair-Davis, A. N., McAllaster, M. R. and de Graffenried, C. L. (2018).
TbSmee1 regulates hook complex morphology and the rate of flagellar pocket uptake in
Trypanosoma brucei. *Mol Microbiol* **107**, 344-362.

Postma, M. and Goedhart, J. (2019). PlotsOfData-A web app for visualizing data together
with their summaries. *PLoS biology* **17**, e3000202.

Price, H. P., Stark, M. and Smith, D. F. (2007). Trypanosoma brucei ARF1 plays a central
role in endocytosis and golgi-lysosome trafficking. *Molecular biology of the cell* **18**, 864-73.

Redmond, S., Vadivelu, J. and Field, M. C. (2003). RNAit: an automated web-based tool for
the selection of RNAi targets in Trypanosoma brucei. *Molecular and biochemical parasitology*
128, 115-8.

Schauberger, P. (2021). openxlsx, pp. Simplifies the creation of Excel .xlsx files by providing
a
high level interface to writing, styling and editing worksheets.
Through the use of 'Rcpp', read/write times are comparable to the
'xlsx' and 'XLConnect' packages with the added benefit of removing the dependency on Java.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B. et al. (2012). Fiji: an open-source
platform for biological-image analysis. *Nat Methods* **9**, 676-82.

Selvapandiyan, A., Kumar, P., Morris, J. C., Salisbury, J. L., Wang, C. C. and Nakhasi,
H. L. (2007). Centrin1 is required for organelle segregation and cytokinesis in Trypanosoma
brucei. *Molecular biology of the cell* **18**, 3290-301.

Shi, J., Franklin, J. B., Yelinek, J. T., Ebersberger, I., Warren, G. and He, C. Y. (2008).
Centrin4 coordinates cell and nuclear division in T. brucei. *Journal of cell science* **121**, 3062-
70.

Sunter, J. D. and Gull, K. (2016). The Flagellum Attachment Zone: 'The Cellular Ruler' of

Trypanosome Morphology. *Trends Parasitol* **32**, 309-324.

Sunter, J. D., Varga, V., Dean, S. and Gull, K. (2015). A dynamic coordination of flagellum

and cytoplasmic cytoskeleton assembly specifies cell morphogenesis in trypanosomes.

Journal of cell science **128**, 1580-94.

Team, R. C. (2021). R: A Language and Environment for Statistical Computing.

Watson, J. F. and Garcia-Nafria, J. (2019). In vivo DNA assembly using common laboratory

bacteria: A re-emerging tool to simplify molecular cloning. *J Biol Chem* **294**, 15271-15281.

Wirtz, E., Leal, S., Ochatt, C. and Cross, G. A. (1999). A tightly regulated inducible

expression system for conditional gene knock-outs and dominant-negative genetics in

Trypanosoma brucei. *Molecular and biochemical parasitology* **99**, 89-101.

Woods, A., Baines, A. J. and Gull, K. (1989). Evidence for a Mr 88,000 glycoprotein with a

transmembrane association to a unique flagellum attachment region in Trypanosoma brucei.

Journal of cell science **93 (Pt 3)**, 501-8.

Zhou, Q., Gheiratmand, L., Chen, Y., Lim, T. K., Zhang, J., Li, S., Xia, N., Liu, B., Lin, Q.

and He, C. Y. (2010). A comparative proteomic analysis reveals a new bi-lobe protein required

for bi-lobe duplication and cell division in Trypanosoma brucei. *PloS one* **5**, e9660.

Zhou, Q., Hu, H., He, C. Y. and Li, Z. (2015). Assembly and maintenance of the flagellum

attachment zone filament in Trypanosoma brucei. *Journal of cell science* **128**, 2361-72.

Zoltner, M., Leung, K. F., Alsford, S., Horn, D. and Field, M. C. (2015). Modulation of the

Surface Proteome through Multiple Ubiquitylation Pathways in African Trypanosomes. *PLoS*

Pathog **11**, e1005236.

Figure legends

Figure 1. TbSmee1 is a cytoskeleton-associated phosphoprotein. (A) Schematic

representation of TbSmee1 (766 amino acids long) with the three predicted domains shown

in purple, orange, and teal. Approximate amino acid ranges of each domain are indicated. The

regions used as antigens for the generation of the three anti-TbSmee1 antibodies (303, 304,

508) are shown in light and dark green bars below the schematic. (B) Validation of the

specificity of the three anti-TbSmee1 antibodies. Immunoblots were carried out against whole

cell lysates from uninduced TbSmee1 RNAi cells (Tet -), induced TbSmee1 RNAi cells (Tet

+), and Ty1-TbSmee1 endogenous replacement cells (End. repl.) using the three anti-

TbSmee1 antibodies (304, 303, 508). All three antibodies detected an ~ 80 kDa protein in the

Tet - samples (upper panels). The protein was strongly depleted after induction of TbSmee1

RNAi (Tet +). A second heavier band was detected in the endogenous replacement cells. This

heavier band could also be detected using anti-Ty1 (BB2) antibodies (lower panels).

Exemplary results from multiple (n>3) independent experiments are shown. (C) Fractionation

scheme. Bloodstream form (BSF) cells were extracted with the non-ionic detergent IGEPAL

and separated into cytoplasm and cytoskeleton fractions by centrifugation. (D) TbSmee1 is

associated with the cytoskeleton. Immunoblot of whole-cell input (I), cytoplasmic supernatant

(SN) and cytoskeletal pellet (P) fractions with anti-TbSmee1 antibodies. Equal fractions were

loaded in each lane. Exemplary results from multiple (n>3) experiments are shown. (E)

TbSmee1 is phosphorylated in vivo. Trypanosome cells were lysed and the lysates incubated

with λ phosphatase for the indicated times (0-20 min), followed by TbSmee1 detection by

immunoblotting. The fuzzy appearance of the TbSmee1 was gradually lost over the

timecourse. This was not observed if the lysates were kept in the presence of phosphatase

inhibitors (+ inhibit), either at 24 ºC or on ice. Equal loading of samples was confirmed using

total protein stain (magenta). Exemplary results from multiple (n>3) independent experiments

are shown. (F) TbSmee1 localisation through the cell cycle. Bloodstream form trypanosomes

were extracted using non-ionic detergent, fixed, and labelled with anti-TbSmee1 antibodies.

DNA was stained using DAPI. In 1K1N cells, TbSmee1 localised to a bar-shaped structure

(arrow) close to the kinetoplast. In 2K1N cells the structure had replicated (arrows). TbSmee1

was additionally present on a third structure farther along the cell body (arrowhead). In 2K2N

cells the signal intensity of the third structure was much weaker. Exemplary images from

multiple (n>3) independent experiments are shown.

1010

Figure 2. TbSmee1 is localised to the shank part of the hook complex. Bloodstream form

trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated

antibodies. Either wild-type or Ty1-TbSmee1 cells were used. Correlation plots are shown

next to the immunofluorescence data. (A) TbSmee1 overlaps with the shank part of the hook

complex protein TbMORN1 (arrow). (B) TbSmee1 overlaps with the shank part of the hook

complex protein TbLRRP1 (arrow). (C) TbSmee1 partially overlaps with TbCen4. TbCen4 is

present at the basal and probasal bodies (yellow arrow), centrin arm (arrowhead), and a small

additional projection (white arrow). (D) TbSmee1 does not overlap with the flagellar pocket

collar protein TbBILBO1 (arrow). (E) Summary of measured correlation coefficients for each

of the colabelling experiments. Each dot represents a single cell. All fluorescence images are

maximum intensity projections, and an overlay with a single DIC section is shown. Overlap

was manually confirmed in single z-slices. Results were obtained from multiple (n>3)

independent experiments; exemplary images are shown.

1024

Figure 3. TbSmee1 is associated with the tip of the elongating new FAZ in replicating
cells. (A) Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and
labelled with the indicated antibodies. DNA was stained with DAPI. Maximum intensity
projections are shown, with a single DIC z-slice overlaid. In 1K1N cells, TbSmee1 partially
overlapped (arrow) with the posterior end of the FAZ. In 2K1N and 2K2N cells, the additional
TbSmee1 structure (arrowhead) was present at the tip of the elongating new FAZ. Exemplary
images from multiple (n>3) independent experiments are shown. (B) SIM image of the same
preparations. The TbSmee1 structure (inset) lay just ahead of the elongating FAZ tip. (C) SIM
image of a detergent-extracted cell colabelled for TbSmee1 and TbDOT1. TbDOT1 appeared
to envelop and partially overlap with TbSmee1.

1035

Figure 4. TbSmee1 domains 2 and 3 are required for targeting to the hook complex. (A)
Schematics of the 9 TbSmee1 truncation constructs tested, along with details of amino acid
ranges, localisation, and size in kDa. (B) TbSmee1 domains 2 and 3 are required for
localisation to the hook complex. Stably-transfected cells that inducibly expressed the
indicated TbSmee1 truncations were used. Ty1-TbSmee1 was detected using anti-Ty1
antibodies. In the absence of induction (-Tet), no signal overlapping with endogenous
TbMORN1 was seen. After induction of expression (+Tet), the Ty1-TbSmee1 truncations
overlapped with the shank part of TbMORN1 (insets). Images shown are maximum intensity
projections of the fluorescence channels overlaid with a single DIC z-slice. Exemplary images
are shown. (C) Confirmation of Ty1-TbSmee1 truncation expression. Whole-cell lysates from
control (-Tet) cells and cells expressing Ty1-TbSmee1 truncations (+Tet) were immunoblotted
using anti-TbSmee1 (left panel) and anti-Ty1 (right panel) antibodies. Total protein stain was
used for normalisation. In the exemplary blot shown, the Ty1-TbSmee1 was detected at ~ 44
kDa, as expected. (D) Quantification of immunoblotting data. Anti-TbSmee1 signals in the
immunoblots were normalised relative to total protein staining. TbSmee1 levels in control (T-)
cells were expressed relative to the mean value of all control samples. Ty1-TbSmee1 levels
were expressed relative to the levels of endogenous TbSmee1 in each clone. All data in the
experiments shown in this figure were obtained from two independent experiments with each
Ty1-TbSmee1 truncation; each experiment used three separate clones.

1055

Figure 5. TbSmee1 is essential for the viability of bloodstream form *T. brucei*. (A)
Depletion of TbSmee1 causes a strong growth defect. The population density of uninduced
control (Tet-) and TbSmee1-depleted (Tet+) cells were measured at regular intervals after
induction of RNAi over a 72 h timecourse. The insert shows data from a more high-resolution
experiments with readings taken at hourly intervals. A growth defect was evident from 21 h
post-induction onwards. (B) Confirmation of TbSmee1 depletion at the single-cell level.

Uninduced control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were extracted, fixed, and
labelled with anti-TbSmee1 antibodies (green). DNA was labelled with DAPI (magenta).
Maximum intensity projections of the fluorescence channels are shown overlaid with a single
DIC z-slice. Insets show the TbSmee1 signal. Exemplary 1K1N and 2K2N cells from each
condition are shown. (C) Confirmation and quantification of TbSmee1 depletion. Whole-cell
lysates from control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were immunoblotted
using antibodies against TbSmee1 (green). Total protein stain (TPS, magenta) was used for
signal normalisation. An exemplary immunoblot is shown. Normalised TbSmee1 signals in
+Tet cells were expressed relative to the -Tet signal for each sample. TbSmee1 -Tet signals
were expressed relative to the mean of all TbSmee1 -Tet values in the dataset. TbSmee1
depletion was quantified at the indicated timepoints post-induction. (D) Depletion of TbSmee1
results in an enlargement of the flagellar pocket. Electron microscopy images of control (Tet-
) and TbSmee1-depleted (Tet+) RNAi cells are shown. The cells were fixed 24 h post-
induction. All data in the experiments shown in this figure were obtained from multiple (n≥3)
independent experiments, each using three separate clones.

1077

Figure 6. Flagellar pocket enlargement is an early consequence of TbSmee1 depletion.
(A) Summary of dextran uptake protocol. The use of temperature blocks inhibits endocytosis,
allowing dextran to enter the flagellar pocket but not be internalised. (B) The intensity of the
dextran signal reports on flagellar pocket volume. Cells were incubated with fluorophore-
conjugated dextran (green) prior to being fixed and imaged using fluorescence microscopy.
DNA was labelled with DAPI (magenta). Exemplary cells from control (- Tet) and TbSmee1-
depleted (+ Tet) conditions are shown. The + Tet cells exhibited various degrees of flagellar
pocket enlargement and progressive morphological aberration. Insets show the dextran signal
with a log-scale LUT. (C) Flow cytometry analysis of control (- Tet) and TbSmee1-depleted (+
Tet) cells incubated with fluorescent dextran at various timepoints after induction of RNAi. At
later timepoints there is a clear emergence of a subpopulation of cells with much greater
fluorescence intensity (arrow). Exemplary traces from a single experiment are shown. (D)
Quantification of flow cytometry data. The geometric mean of the fluorescence intensity in
control (- Tet) and TbSmee1-depleted (+ Tet) cells was measured at various timepoints after
induction of RNAi. Bars indicate median values. The population mean of the + Tet cells was
clearly higher at the later timepoints; a visible shift was visible as early as 18 h post-induction
(arrow). All data in the experiments shown in this figure were obtained from multiple (n≥2)
independent experiments; each experiment used three separate clones.

1096

Figure 7. Surface-bound reporters cannot enter the flagellar pocket of TbSmee1-
depleted cells. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to

enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, the

dextran and ConA signals strongly overlapped (arrow, upper panels). In TbSmee1-depleted

cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower panels).

Maximum intensity projections of the fluorescence channels are shown overlaid with a single

DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet)

and TbSmee1-depleted (+Tet) samples were manually classified into four categories (1-4)

based on the degree of overlap between the ConA and dextran reporters. Results are shown

as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are

shown. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the

enlarged flagellar pocket of TbSmee1-depleted cells. (F) Quantification of dextran+BSA

uptake experiments. Data were obtained from multiple (n>3) independent experiments; each

experiment used three separate clones.

1111

Figure 8. Blockade of surface reporters is due to an inhibition of endocytosis. (A)

Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged

flagellar pocket of clathrin-depleted cells. In control (-Tet) cells, the dextran and ConA signals

strongly overlapped (arrow, upper panels). In clathrin-depleted cells (+Tet), little to no overlap

between the two reporters was observed (arrow, lower-panels). Maximum intensity projections

of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification

of dextran+ConA uptake experiments. Cells from control (-Tet) and clathrin-depleted (+Tet)

samples were manually classified into four categories (1-4) based on the degree of overlap

between the ConA and dextran reporters. Results are shown as a slope chart (left) and

stacked bar chart (right); exemplary cells for each category are shown. (D) Summary of paired

dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of

clathrin-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Data were

obtained from multiple (n>3) independent experiments; each experiment used three separate

clones.

1126

1127

Figure S1. The TbSmee1 primary structure contains three blocks of highly-conserved

sequence. Multiple sequence alignment of *Trypanosoma brucei* Smee1 (TbSmee1) and

homologous proteins from other trypanosome species, generated using MultAlin (Corpet,

1988) with default parameters. Amino acid numbers are indicated in black numerals above

the alignment. Moderately (50-90%) conserved residues are highlighted in blue. Highly-

conserved (>90%) or completely conserved residues are highlighted in red. The consensus

sequence is shown below the alignment. Abbreviations: TbSmee1, *Trypanosoma brucei*

Smee1; T.cong, *Trypanosoma congolense*; T.vivax, *Trypanosoma vivax*; T.grayi,

Trypanosoma grayi. T.conorhini, *Trypanosoma conorhini*; T.rangelii, *Trypanosoma rangelii*;
T.cruzi, *Trypanosoma cruzi*; T.theileri, *Trypanosoma theileri*. The % sequence identity of each
homologue to TbSmee1 is indicated after the name.

1139

Figure S2. TbSmee1 contains three predicted structured domains. (A) Schematic
representation of TbSmee1 with the three predicted domains (Dom1/2/3) indicated in blue,
magenta, and green. The approximate amino acid coordinates in the primary structure are
indicated in black above each domain. The isoelectric point (pI) of each domain is indicated in
bold text above the domain. (B) TbSmee1 contains three regions of predicted secondary
structure. Secondary structure prediction generated using the PSIPRED server (Buchan &
Jones, 2019) with default parameters. Abbreviations and schematics are defined in the legend
at the bottom right. The three predicted structured domains are outlined as blue, magenta,
and green boxes.

1149

Figure S3. Generation of Ty1-TbSmee1 endogenous replacement cell line. (A) Schematic
showing the in situ tagging procedure and annealing sites for PCR primers. The two
endogenous alleles of TbSmee1 in the gDNA are shown in yellow, flanked by 5'UTR and
3'UTR sequences. *T.* brucei cells were transfected with a targeting fragment with homology
arms for recombination with the 3' end of the 5'UTR and 5' end of the ORF. The targeting
fragment contained a blasticidin resistance gene (bla^R), the intergenic region from the
alpha/beta tubulin locus ($\alpha\beta$) and a 3xTy1 epitope tag preceded by an ATG start codon.
Homologous recombination removed the endogenous ATG start codon of the TbSmee1 ORF.
(B) Confirmation of targeting fragment integration at the endogenous TbSmee1 locus by PCR
analysis of genomic DNA. Genomic DNA from wild-type (wt) and candidate Ty1-TbSmee1
clones (c1,c2,c3) was analysed by PCR. Left panel: positive control using 8820_ORF_F and
8820_gDNA-R primers; 490 bp product expected. Right panel: integration test using BB2-F
and 8820_gDNA-R primers. A product is only expected if the 3xTy1 sequence has integrated
upstream of the TbSmee1 ORF (see primer annealing sites in panel A). Two independent
experiments were carried out, each using all three separate clones. (C) Ty1-TbSmee1
displays the same localisations through the cell cycle as endogenous TbSmee1. Detergent-
extracted cells were fixed with methanol and labelled with anti-Ty1 antibodies; DAPI was
stained using DAPI. Exemplary cells from the three main cell cycle states (1K1N, 2K1N, 2K2N)
are shown. Maximum-intensity z-projections of the fluorescence channels are shown, together
with a single DIC z-slice overlay. The Ty1-TbSmee1 signal is shown in magenta in the overlay
and highlighted with arrows. Arrowheads indicate the Ty1-TbSmee1 present at the tip of the
new FAZ. Insets show an enlarged view of the TbSmee1 signal from the hook complex or FAZ
tip. Multiple (n>3) independent experiments were carried out using three separate clones. (D)

The anti-Ty1 signal is specific for Ty1-TbSmee1. Wild-type (wt) and a Ty1-TbSmee1 clone
(c1) were extracted with detergent, fixed with methanol, and labelled with anti-TbSmee1 and
anti-Ty1 antibodies. Hook complex (arrows) and FAZ tip (arrowheads) localisations are
indicated. No anti-Ty1 signal was seen in wild-type cells; strong overlap between the anti-Ty1
and anti-TbSmee1 signals was seen in the Ty1-TbSmee1 cells. Maximum-intensity z-
projections of the fluorescence channels are shown, together with a single DIC z-slice overlay.
1179

Figure S4. Validation of anti-TbStarkey1 antibodies. (A) Validation of anti-TbStarkey1
antibody specificity by immunoblotting. Three separate TbStarkey1 RNAi clones (c8, c9, c10)
were analysed by immunoblotting. Both whole-cell lysates (WCL) and detergent-extracted
cytoskeleton (CK) samples were obtained at the indicated timepoints (24 h, 48 h) from control
(Tet-) and induced (Tet+) cells. The samples were immunoblotted using two separate anti-
TbStarkey1 affinity-purified antibodies (antibody 476, 477). Both antibodies recognised a >130
kDa protein whose abundance was depleted after 48 h of RNAi. The 476 antibody additionally
showed a significant cross-reaction with a protein of <100 kDa in WCL but not CK samples. A
section of the total protein stain of each membrane is shown as an inset (magenta). (B)
Validation of anti-TbStarkey1 antibody specificity by immunofluorescence microscopy. Control
(Tet-) and TbStarkey1-depleted (Tet+) RNAi cells were extracted with nonionic detergent,
fixed with methanol, and labelled with anti-TbStarkey1 antibodies (green); DNA was stained
using DAPI (magenta). TbStarkey1 localised to the hook complex, and signal was lost upon
depletion. The boxed areas are shown enlarged in the bottom panels. Identical results were
obtained using both anti-TbStarkey1 antibodies; exemplary images using the 477 antibodies
are shown. (C) TbSmee1 overlaps with the shank part of the hook complex protein TbStarkey1
(arrow). (D) The TbSmee1 and TbStarkey1 pixel values are correlated. (E) Summary of
measured correlation coefficients. Each dot represents a single cell. All fluorescence images
are maximum intensity projections, and an overlay with a single DIC section is shown. Overlap
was manually confirmed in single z-slices. Results were obtained from multiple (n>3)
independent experiments; exemplary images are shown.
1201

1202

Figure S5. Validation of TbSmee1 truncations. (A) Validation of Ty1-TbSmee1 truncation
cell lines by PCR analysis of genomic DNA and immunoblotting. The results for all 8 TbSmee1
truncations are shown, separated by dotted lines. Upper panels: PCR analysis of genomic
DNA to confirm the presence of the truncation construct. PCR was used to amplify DNA from
clones (c1, c2, c3) and wild-type control (wt) genomic DNA. Primers annealing to the sequence
encoding Ty1 epitope and within the truncation were used. Lower panels: confirmation of Ty1-
TbSmee1 truncation construct expression by immunoblotting. Whole-cell lysates from

uninduced (-Tet) and induced (+Tet) cells were analysed by immunoblotting with anti-TbSmee1 (green) and anti-Ty1 (cyan) antibodies. Both antibodies detected proteins corresponding to the predicted size of the Ty1-TbSmee1 truncations; arrows indicate the target protein in the anti-Ty1 blots. A portion of the total protein staining of the membranes is shown in magenta. (B) Ty1-TbSmee1(161-766) localises to both the hook complex and the FAZ tip. Cells expressing the Ty1-TbSmee1(161-766) construct were extracted with nonionic detergent, fixed with methanol, and labelled with anti-TbMORN1 and anti-Ty1 antibodies. DNA was stained with DAPI (cyan). Ty1-TbSmee1(161-766) was observed at both the hook complex and the FAZ tip (arrow). (C) Ty1-TbSmee1(2-400) localises to the FAZ tip but not the hook complex. Cells expressing the Ty1-TbSmee1(2-400) construct were extracted, fixed, and labelled as above. Ty1-TbSmee1(2-400) was observed exclusively at the FAZ tip (arrow). Images in panels B and C are maximum intensity z-projections with a single DIC z-slice overlay. Multiple (n>2) independent experiments using three separate clones for each construct were carried out; exemplary cells are shown.

Figure S6. Additional results for TbSmee1 and TbStarkey1 RNAi experiments. (A) TbSmee1 depletion causes changes to cell cycle state distribution. Control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells from 18 h, 21 h, 24 h post-induction were fixed using glutaraldehyde; DNA was stained using DAPI. The various cell cycle states (1K1N, etc) were manually quantified from images taken of the fixed cells. Data were obtained from three independent experiments, each using three separate clones; at least 240 cells were quantified for each timepoint. (B) Depletion of TbStarkey1 has no effect on population cell growth. Control (-Tet) and TbStarkey1-depleted (+Tet) cells were followed over a 72 h timecourse, and population density (cells/ml) was measured every 24 h. Data were obtained from three independent experiments, each using 3 separate clones. (C, D) Depletion of TbStarkey1 causes morphological abnormality. TbStarkey1-depleted cells were prepared for electron microscopy using high-pressure freezing and imaged. Cells with enlarged flagellar pockets could readily be observed.

Figure S7. Depletion of TbSmee1 affects other cytoskeleton-associated proteins. TbSmee1 RNAi cells were induced for 24 h and analysed by immunoblotting of whole-cell lysates and immunofluorescence imaging of detergent-extracted cells. (A) Exemplary immunoblots showing the effect of 24 h TbSmee1 depletion on marker proteins for the hook complex, flagellar pocket collar, and centrin arm. For comparison, the effect of 16 h TbMORN1 depletion was also analysed. (B) Quantification of immunoblot data. TbSmee1 depletion did not affect any of the proteins analysed. Total protein staining was used for signal normalisation. Normalised TbSmee1 signals in +Tet cells were expressed relative to the -Tet

signal for each sample. TbSmee1 -Tet signals were expressed relative to the mean of all

TbSmee1 -Tet values in the dataset. Immunoblots from TbMORN1 RNAi experiments were

quantified in the same way; TbMORN1 depletion resulted in a loss of both TbSmee1 and

TbStarkey1 signal. Data were obtained from multiple (n>2) independent experiments, each

using three (TbSmee1 RNAi) or two (TbMORN1 RNAi) separate clones. (C-F) TbSmee1

depletion affects TbLRRP1 but not other marker proteins analysed. Control (-Tet) and

TbSmee1-depleted (+Tet) cells were extracted with detergent, fixed with methanol, and

labelled with the indicated antibodies; DNA was stained using DAPI. The position of the hook

complexes in exemplary 1K1N and 2K2N cells are shown with arrows; panel E also indicates

basal bodies. Data were obtained from 2 independent experiments for each labelling

combination, each using 3 separate clones. (G) TbSmee1 and TbLRRP1 in TbMORN1-

depleted cells. (H) Summary of the observed effects on the abundance of marker proteins

caused by either TbMORN1 or TbSmee1 depletion, based on immunoblotting and

immunofluorescence data.

1261

1262

Figure S8. Depletion of TbSmee1 prevents trafficking of ConA to the lysosome. Control

(-Tet) and TbSmee1-depleted cells (+Tet; 24 h timepoint) were incubated on ice with ConA

(magenta). The cells were then shifted to 37 ºC for 30 min to allow internalisation. The cells

were then fixed and labelled with antibodies specific for the lysosome marker p67 (green).

Control (-Tet) cells showed strong overlap between the two labels, indicating that ConA had

been trafficked to the lysosome. +Tet cells showed no overlap between the two labels.

Maximum intensity projections of the fluorescence channels are shown overlaid with a single

DIC z-slice. Overlap in -Tet cells was confirmed in single z-slices. Single channels from the

boxed area in each image are shown as insets. Data obtained from multiple (n>2) independent

experiments each using three separate clones; exemplary cells are shown.

1273

1274

1275

1276

1277

Figure 1. TbSmee1 is a cytoskeleton-associated phosphoprotein. (A) Schematic representation of TbSmee1 (766 amino acids long) with the three predicted domains shown in purple, orange, and teal. Approximate amino acid ranges of each domain are indicated. The regions used as antigens for the generation of the three anti-TbSmee1 antibodies (303, 304, 508) are shown in light and dark green bars below the schematic. (B) Validation of the specificity of the three anti-TbSmee1 antibodies. Immunoblots were carried out against whole cell lysates from uninduced TbSmee1 RNAi cells (Tet -), induced TbSmee1 RNAi cells (Tet +), and Ty1-TbSmee1 endogenous replacement cells (End. repl.) using the three anti-TbSmee1 antibodies (304, 303, 508). All three antibodies detected an ~ 80 kDa protein in the Tet - samples (upper panels). The protein was strongly depleted after induction of TbSmee1 RNAi (Tet +). A second heavier band was detected in the endogenous replacement cells. This heavier band could also be detected using anti-Ty1 (BB2) antibodies (lower panels). Exemplary results from multiple (n>3) independent experiments are shown. (C) Fractionation scheme. Bloodstream form (BSF) cells were extracted with the non-ionic detergent IGEPAL and separated into cytoplasm and cytoskeleton fractions by centrifugation. (D) TbSmee1 is associated with the cytoskeleton. Immunoblot of whole-cell input (I), cytoplasmic supernatant (SN) and cytoskeletal pellet (P) fractions with anti-TbSmee1 antibodies. Equal fractions were loaded in each lane. Exemplary results from multiple (n>3) experiments are shown. (E) TbSmee1 is phosphorylated in vivo. Trypanosome cells were lysed and the lysates incubated with λ phosphatase for the indicated times (0-20 min), followed by TbSmee1 detection by immunoblotting. The fuzzy appearance of the TbSmee1 was gradually lost over the timecourse. This was not observed if the lysates were kept in the presence of phosphatase inhibitors (+ inhibit), either at 24 ºC or on ice. Equal loading of samples was confirmed using total protein stain (magenta). Exemplary results from multiple (n>3) independent experiments are shown. (F) TbSmee1 localisation through the cell cycle. Bloodstream form trypanosomes were extracted using non-ionic detergent, fixed, and labelled with anti-TbSmee1 antibodies. DNA was stained using DAPI. In 1K1N cells, TbSmee1 localised to a bar-shaped structure (arrow) close to the kinetoplast. In 2K1N cells the structure had replicated (arrows). TbSmee1 was additionally present on a third structure farther along the cell body (arrowhead). In 2K2N cells the signal intensity of the third structure was much weaker. Exemplary images from multiple (n>3) independent experiments are shown.

Figure 2. TbSmee1 is localised to the shank part of the hook complex. Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated antibodies. Either wild-type or Ty1-TbSmee1 cells were used. Correlation plots are shown next to the immunofluorescence data. (A) TbSmee1 overlaps with the shank part of the hook complex protein TbMORN1 (arrow). (B) TbSmee1 overlaps with the shank part of the hook complex protein TbLRRP1 (arrow). (C) TbSmee1 partially overlaps with TbCen4. TbCen4 is present at the basal and probasal bodies (yellow arrow), centrin arm (arrowhead), and a small additional projection (white arrow). (D) TbSmee1 does not overlap with the flagellar pocket collar protein TbBILBO1 (arrow). (E) Summary of measured correlation coefficients for each of the colabelling experiments. Each dot represents a single cell. All fluorescence images are maximum intensity projections, and an overlay with a single DIC section is shown. Overlap was manually confirmed in single z-slices. Results were obtained from multiple (n>3) independent experiments; exemplary images are shown.

Figure 3. TbSmee1 is associated with the tip of the elongating new FAZ in replicating cells. (A) Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated antibodies. DNA was stained with DAPI. Maximum intensity projections are shown, with a single DIC z-slice overlaid. In 1K1N cells, TbSmee1 partially overlapped (arrow) with the posterior end of the FAZ. In 2K1N and 2K2N cells, the additional TbSmee1 structure (arrowhead) was present at the tip of the elongating new FAZ. Exemplary images from multiple (n>3) independent experiments are shown. (B) SIM image of the same preparations. The TbSmee1 structure (inset) lay just ahead of the elongating FAZ tip. (C) SIM image of a detergent-extracted cell colabelled for TbSmee1 and TbDOT1. TbDOT1 appeared to envelop and partially overlap with TbSmee1.

Figure 4. TbSmee1 domains 2 and 3 are required for targeting to the hook complex. (A) Schematics of the 9 TbSmee1 truncation constructs tested, along with details of amino acid ranges, localisation, and size in kDa. (B) TbSmee1 domains 2 and 3 are required for localisation to the hook complex. Stably-transfected cells that inducibly expressed the indicated TbSmee1 truncations were used. Ty1-TbSmee1 was detected using anti-Ty1 antibodies. In the absence of induction (-Tet), no signal overlapping with endogenous TbMORN1 was seen. After induction of expression (+Tet), the Ty1-TbSmee1 truncations overlapped with the shank part of TbMORN1 (insets). Images shown are maximum intensity projections of the fluorescence channels overlaid with a single DIC z-slice. Exemplary images are shown. (C) Confirmation of Ty1-TbSmee1 truncation expression. Whole-cell lysates from control (-Tet) cells and cells expressing Ty1-TbSmee1 truncations (+Tet) were immunoblotted using anti-TbSmee1 (left panel) and anti-Ty1 (right panel) antibodies. Total protein stain was used for normalisation. In the exemplary blot shown, the Ty1-TbSmee1 was detected at ~ 44 kDa, as expected. (D) Quantification of immunoblotting data. Anti-TbSmee1 signals in the immunoblots were normalised relative to total protein staining. TbSmee1 levels in control (T-) cells were expressed relative to the mean value of all control samples. Ty1-TbSmee1 levels were expressed relative to the levels of endogenous TbSmee1 in each clone. All data in the experiments shown in this figure were obtained from two independent experiments with each Ty1-TbSmee1 truncation; each experiment used three separate clones.

Figure 5. TbSmee1 is essential for the viability of bloodstream form T. brucei. (A) Depletion of TbSmee1 causes a strong growth defect. The population density of uninduced control (Tet-) and TbSmee1-depleted (Tet+) cells were measured at regular intervals after induction of RNAi over a 72 h timecourse. The insert shows data from a more high-resolution experiments with readings taken at hourly intervals. A growth defect was evident from 21 h post-induction onwards. (B) Confirmation of TbSmee1 depletion at the single-cell level. Uninduced control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were extracted, fixed, and labelled with anti-TbSmee1 antibodies (green). DNA was labelled with DAPI (magenta). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. Insets show the TbSmee1 signal. Exemplary 1K1N and 2K2N cells from each condition are shown. (C) Confirmation and quantification of TbSmee1 depletion. Whole-cell lysates from control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were immunoblotted using antibodies against TbSmee1 (green). Total protein stain (TPS, magenta) was used for signal normalisation. An exemplary immunoblot is shown. Normalised TbSmee1 signals in +Tet cells were expressed relative to the -Tet signal for each sample. TbSmee1 -Tet signals were expressed relative to the mean of all TbSmee1 -Tet values in the dataset. TbSmee1 depletion was quantified at the indicated timepoints post-induction. (D) Depletion of TbSmee1 results in an enlargement of the flagellar pocket. Electron microscopy images of control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells are shown. The cells were fixed 24 h post-induction. All data in the

Figure 6. Flagellar pocket enlargement is an early consequence of TbSmee1 depletion. (A) Summary of dextran uptake protocol. The use of temperature blocks inhibits endocytosis, allowing dextran to enter the flagellar pocket but not be internalised. (B) The intensity of the dextran signal reports on flagellar pocket volume. Cells were incubated with fluorophore-conjugated dextran (green) prior to being fixed and imaged using fluorescence microscopy. DNA was labelled with DAPI (magenta). Exemplary cells from control (- Tet) and TbSmee1-depleted (+ Tet) conditions are shown. The + Tet cells exhibited various degrees of flagellar pocket enlargement and progressive morphological aberration. Insets show the dextran signal with a log-scale LUT. (C) Flow cytometry analysis of control (- Tet) and TbSmee1-depleted (+ Tet) cells incubated with fluorescent dextran at various timepoints after induction of RNAi. At later timepoints there is a clear emergence of a subpopulation of cells with much greater fluorescence intensity (arrow). Exemplary traces from a single experiment are shown. (D) Quantification of flow cytometry data. The geometric mean of the fluorescence intensity in control (- Tet) and TbSmee1-depleted (+ Tet) cells was measured at various timepoints after induction of RNAi. Bars indicate median values. The population mean of the + Tet cells was clearly higher at the later timepoints; a visible shift was visible as early as 18 h post-induction (arrow). All data in the experiments shown in this figure were obtained from multiple (n≥2) independent experiments; each experiment used three separate clones.

Figure 7. Surface-bound reporters cannot enter the flagellar pocket of TbSmee1-depleted cells. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In TbSmee1-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and TbSmee1-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Data were obtained from multiple (n>3) independent experiments; each experiment used three separate clones.

Figure 8. Blockade of surface reporters is due to an inhibition of endocytosis. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In clathrin-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower-panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and clathrin-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Data were obtained from multiple (n>3) independent experiments; each experiment used three separate clones.