1 **Characterisation of TbSmee1 indicates that endocytosis is required for access of**
2 **surface-bound cargo to the trypanosome flagellar pocket**
3
4 **Running title:** Characterisation of TbSmee1
5

Authors:
Daja Schichler[1]*, Antonia Konle*, Eva-Maria Spath**[2], Sina Riegler**, Alexandra Klein[0], Anna
Seleznev[0], Sisco Jung[0], Timothy Wuppermann[#], Noah Wetterich[#], Alyssa Borges, Elisabeth
Meyer-Natus, Katharina Havlicek[3], Sonia Pérez Cabrera[4], Korbinian Niedermüller, Sara
Sajko[5], Maximilian Dohn, Xenia Malzer, Emily Riemer, Tuguldur Tumurbaatar, Kristina
Djinovic-Carugo[5,6,7], Gang Dong[8], Christian J. Janzen, Brooke Morriswood.
12
13 */**/[0]/[#] These authors contributed equally
14
15 **Affiliations:**
16 Unless stated otherwise: Department of Cell & Developmental Biology, Biocenter, University
17 of Würzburg, Würzburg, Germany
18
1 present address: Cell Biology and Biophysics Unit, EMBL, Heidelberg, Germany
2 present address: Department of Internal Medicine 1, Friedrich-Alexander-University
Erlangen-Nuremberg, Erlangen, Germany
3 Department of Biochemistry and Cell Biology, Max Perutz Labs, University of Vienna, Vienna
Biocenter (VBC) Vienna, Austria
4 Francisco de Vitoria University, Madrid, Spain
5 Department of Structural and Computational Biology, Max Perutz Labs, University of Vienna,
Vienna Biocenter (VBC), Vienna, Austria
6 Department of Biochemistry, Faculty of Chemistry and Chemical Technology, University of
Ljubljana, Ljubljana, Slovenia
7 European Molecular Biology Laboratory (EMBL) Grenoble, Grenoble, France
8 Center for Medical Biochemistry, Max Perutz Labs, Medical University of Vienna, Vienna
Biocenter (VBC) Vienna, Austria
32
33 **Email for correspondence:** brooke.morriswood@gmail.com

35 **Keywords:** Trypanosoma brucei, cytoskeleton, hook complex, TbSmee1, flagellar pocket,
36 endocytosis

38   **Summary:** Characterisation of the essential trypanosome protein TbSmee1 suggests that
39   endocytosis is required for flagellar pocket access of surface-bound cargo.

Abstract

All endo- and exocytosis in the African trypanosome *Trypanosoma brucei* occurs at a single subdomain of the plasma membrane. This subdomain, the flagellar pocket, is a small vase-shaped invagination containing the root of the cell's single flagellum. Several cytoskeleton-associated multiprotein complexes are coiled around the neck of the flagellar pocket on its cytoplasmic face. One of these, the hook complex, was proposed to affect macromolecule entry into the flagellar pocket lumen. In previous work, knockdown of the hook complex component TbMORN1 resulted in larger cargo being unable to enter the flagellar pocket. In this study, the hook complex component TbSmee1 was characterised in bloodstream form *Trypanosoma brucei* and was found to be essential for cell viability. TbSmee1 knockdown resulted in flagellar pocket enlargement and impaired access to the flagellar pocket membrane by surface-bound cargo, similar to depletion of TbMORN1. Unexpectedly, inhibition of endocytosis by knockdown of clathrin phenocopied TbSmee1 knockdown, suggesting that endocytic activity itself is a prerequisite for the entry of surface-bound cargo into the flagellar pocket.

Introduction

The flagellated parasitic protist *Trypanosoma brucei* lives in the bloodstream of its mammalian hosts, in continuous exposure to the immune system. Endo- and exocytosis are therefore critical to the parasite's survival, as they are used to remove bound antibodies from the cell surface and scavenge macromolecular nutrients from the surroundings (Borst and Fairlamb, 1998; Engstler et al., 2007; Webster et al., 1990). Remarkably, all endo- and exocytosis in *T. brucei* occurs at just a single subdomain of the plasma membrane, despite the overall rate of endocytosis being comparable to a professional phagocyte (Engstler et al., 2004; Grunfelder et al., 2003). This subdomain is a small vase-shaped invagination that houses the root of the cell's single flagellum, and is called the flagellar pocket (Halliday et al., 2021; Lacomble et al., 2009).

Molecules that enter the flagellar pocket, either in the fluid phase or when bound to the surface of the parasite (here "surface-bound"), are rapidly internalised by clathrin-mediated endocytosis, and then routed to the endosomal/lysosomal system (Link et al., 2021; Overath and Engstler, 2004). Depletion of clathrin results in a gross enlargement of the flagellar pocket due to an imbalance between endocytosis and exocytosis, and causes cargo accumulation inside the flagellar pocket (Allen et al., 2003). Although endocytosis of fluid phase and surface-bound cargo is relatively well-characterised in *T. brucei*, the exact mechanism(s) by which cargo initially enters the flagellar pocket are less well-understood.

Coiled around the neck of the flagellar pocket on its cytoplasmic face are a number of poorly-
characterised cytoskeleton-associated complexes, which might contribute to these processes
(Esson et al., 2012; Halliday et al., 2019)(Fig. 1A). The flagellar pocket collar is a multiprotein
complex shaped like a cuff bracelet, and demarcates the boundary between the flagellar
pocket and the flagellar pocket neck. It contains the protein TbBILBO1, and a number of its
other components such as FPC4 and BILBO2 have recently been characterised (Albisetti et
al., 2017; Bonhivers et al., 2008; Florimond et al., 2015; Isch et al., 2021). The centrin arm is
a centrin-containing kinked rod which appears to be involved in the biogenesis of various
cytoskeleton-associated structures (Selvapandiyan et al., 2007; Shi et al., 2008). The hook
complex is a hook-shaped multiprotein complex that sits atop the flagellar pocket collar and
alongside the centrin arm. Characterised components include the proteins TbMORN1, BOH1,
BOH2, and Bhalin (Broster Reix et al., 2021; Morriswood et al., 2009; Pham et al., 2020; Pham
et al., 2019). Previous characterisation of TbMORN1 revealed that it is essential for the viability
of bloodstream form *T. brucei* cells (Morriswood and Schmidt, 2015). Cells depleted of
TbMORN1 have enlarged flagellar pockets, indicative of an endocytosis defect. In addition,
although small endocytic reporters such as 10 kDa dextran (hydrodynamic radius ~1.86 nm)
were still capable of entering the enlarged flagellar pocket, the access of larger
macromolecules such as the lectin concanavalin A (ConA, hydrodynamic radius ~4.2 nm) or
bovine serum albumin (BSA, hydrodynamic radius ~ 3.51 nm) conjugated to 5 nm gold
particles was blocked (Ahmad et al., 2007; Armstrong et al., 2004; Morriswood and Schmidt,
2015). This phenotype was also observed upon depletion of the hook complex protein Bhalin
(Broster Reix et al., 2021). On this basis, it was proposed that the hook complex might be
regulating the passage of large macromolecules through the flagellar pocket neck, and thereby
mediating cargo entry into the flagellar pocket (Morriswood and Schmidt, 2015).

In this study, the hook complex protein TbSmee1 was characterised in bloodstream form *T.*
brucei. TbSmee1 (Tb927.10.8820) was initially identified in a proximity labelling screen using
TbMORN1 as bait, and shown to localise to the shank part of the hook complex (Morriswood
et al., 2013). It was subsequently characterised in procyclic form *T. brucei*, the life cycle stage
found within the tsetse fly vector (Perry et al., 2018). In bloodstream form *T. brucei,* depletion
of TbSmee1 phenocopied TbMORN1 and Bhalin, with ConA, BSA and surface-bound
antibodies unable to access the enlarged flagellar pocket. Surprisingly, clathrin-depleted cells
also exhibited this flagellar pocket access phenotype, suggesting that the observed effects
following depletion of the hook complex proteins are due to impaired endocytosis, and
therefore that endocytosis is required for the entry of surface-bound cargo into the flagellar
pocket.

Results

TbSmee1 is a cytoskeleton-associated phosphoprotein.

Bioinformatic analysis predicted that TbSmee1 is composed of three structured domains separated by linker regions (Fig. 1B). Specifically, a multiple sequence alignment of the TbSmee1 protein with orthologues from other trypanosome species indicated the presence of three blocks of highly-conserved sequence (Fig. S1). These blocks of conserved sequence also corresponded to regions of predicted secondary structure (Fig. S2). Three-dimensional structural prediction by AlphaFold suggested a close association between the first two folded domains, and showed a good match with the secondary structural predictions (https://alphafold.ebi.ac.uk/entry/Q38A92) (Jumper et al., 2021; Wheeler, 2021).

Two polyclonal antibodies (303, 304) were raised against a recombinant TbSmee1(1-400) truncation, and a third (508) was raised against two peptides (Fig. 1B, green lines). The specificity of the antibodies was confirmed using immunoblots of whole-cell lysates obtained from TbSmee1 RNAi cells and 3xTy1-TbSmee1 endogenous replacement cells. In situ tagging of the *SMEE1* gene in the 3xTy1-TbSmee1 cells was confirmed by PCR analysis of genomic DNA (Fig. S3A,B). All three antibodies detected a protein of ~ 85 kDa in the immunoblots (Fig. 1C, white arrows). This protein was strongly depleted in the RNAi (Tet +) samples (Fig. 1C). An extra band corresponding to 3xTy1-TbSmee1 was seen in lysates from the endogenous replacement (End. repl.) cells (Fig. 1C). This extra band was also detected using anti-Ty1 (BB2) antibodies, confirming its identity as Ty1-TbSmee1 (Fig 1C, lower panels).

To investigate TbSmee1 association with the cytoskeleton, cells were fractionated using non-ionic detergent (Fig. 1D). The fractions were analysed by immunoblotting, using antibodies against PFR1,2 as markers for the cytoskeleton, and against the ER chaperone BiP as a marker for the cytoplasm. TbSmee1 co-fractionated with PFR1,2, confirming that it was cytoskeleton-associated (Fig. 1E). Interestingly, the fuzzy appearance of the TbSmee1 band observed in immunoblots of whole-cell lysates was reduced or absent in the fractionation blots (compare Fig. 1C and Fig. 1E). TbSmee1 is heavily phosphorylated in vivo, and is a substrate and potential binding partner of the mitotic kinase TbPLK (Benz and Urbaniak, 2019; McAllaster et al., 2015; Nett et al., 2009; Urbaniak et al., 2013). The existence of different phosphoforms of a protein is known to cause a fuzzy appearance of bands in gels, so it seemed possible that exposure of TbSmee1 to endogenous phosphatases during the extraction step was causing dephosphorylation.

To investigate whether the band collapse could be attributed to dephosphorylation of
TbSmee1, extracted cytoskeletons were incubated with exogenous phosphatase at various
timepoints prior to immunoblotting. Incubation with exogenous phosphatase resulted in a
progressive collapse of the TbSmee1 band over a 20 min period (Fig. 1F). This band collapse
was not seen when the samples were in the presence of phosphatase inhibitors either at RT
or on ice, or kept on ice (Fig. 1F, last three lanes). The fuzzy appearance of TbSmee1 in
immunoblots could therefore be attributed exclusively to phosphorylation.
158
159

TbSmee1 localises to the shank part of the hook complex.
The localisation of endogenous and tagged TbSmee1 protein was analysed in bloodstream
form *T. brucei* cells using immunofluorescence microscopy (Fig. 1G). All three main stages of
the cell division cycle (1K1N, 2K1N, 2K2N) were analysed. In 1K1N cells (i.e. those with a
single kinetoplast, K, and nucleus, N), TbSmee1 localised to a single structure near the
flagellum base, consistent with the position of the hook complex (Fig. 1G, arrow). Interestingly,
in 2K1N and 2K2N cells there was a third subpopulation of TbSmee1 at varying distances
along the cell body in addition to the replicated hook complex (Fig. 1G, arrowheads). This
subpopulation was fainter in 2K2N cells than 2K1N ones. The same TbSmee1 distributions
were seen using all three anti-TbSmee1 antibodies; anti-Ty1 antibody labelling of 3xTy1-
TbSmee1 cells also produced the same labelling pattern (Fig. S3C). In the endogenous
replacement cells, the anti-TbSmee1 and anti-Ty1 signals colocalised (Fig. S3D).
172

To confirm the localisation of TbSmee1 at the hook complex, colabelling experiments were
carried out. As expected, TbSmee1 strongly overlapped with the shank part of the hook
complex as labelled by TbMORN1 and TbLRRP1 (Fig. 2A, B). TbSmee1 also strongly
overlapped with the hook complex protein Tb927.10.3010. This 133 kDa protein was one of
the top hits in the BioID screen using TbMORN1 (Morriswood et al., 2013). Consistent with
the convention started with TbSmee1, Tb927.10.3010 was named TbStarkey1 after another
of Captain Hook's pirates. Two anti-peptide antibodies were generated against TbStarkey1,
and their specificity was confirmed by immunoblotting and immunofluorescence imaging of
TbStarkey1 RNAi cells (Fig. S4A, B). Strong overlap was seen between TbSmee1 and
TbStarkey1 on the shank part of the hook complex (Fig. S4C). As expected, TbSmee1 showed
only a partial overlap with TbCentrin4, a marker of the centrin arm, and little overlap with
TbBILBO1, a marker of the flagellar pocket collar (Fig. 2C, D).
185

186 To quantify the degree of colocalisation between TbSmee1 and the various hook complex and
187 flagellar pocket collar marker proteins, correlation coefficients from the pairwise labelling

188 experiments were calculated. Moderate correlation was seen between TbSmee1 and
189 TbMORN1 and TbLRRP1, with lower values being measured for TbCentrin4 and TbBILBO1
190 (Fig. 2E).
191
192

TbSmee1 is transiently associated with the flagellum attachment zone (FAZ) tip.
The flagellum attachment zone (FAZ) is a cytoskeleton-associated apparatus that adheres the
flagellum of *T. brucei* to the cell body (Sunter and Gull, 2016). During replication, a new FAZ
is assembled and grows from its initiation point very close to the hook complex towards the
anterior end of the cell (Sunter et al., 2015b; Zhou et al., 2015). It was previously shown that
a tagged version of TbSmee1 is transiently localised to the tip of the new FAZ in replicating
insect-stage *T. brucei* cells, in addition to its hook complex localisation (Perry et al., 2018).
This dynamic localisation contrasts with other hook complex proteins such as TbMORN1,
which are localised exclusively to the hook complex.
202

To investigate whether the endogenous TbSmee1 subpopulation seen in 2K1N and 2K2N
BSF cells corresponded to the FAZ tip, TbSmee1 and the FAZ marker FAZ1 were analysed
at different stages of the cell division cycle. As expected, the TbSmee1 present at the hook
complex showed a partial overlap with the posterior end of the FAZ (Fig. 3A, arrow). The extra
TbSmee1 subpopulation in 2K1N and 2K2N cells overlapped with the tip of the new FAZ (Fig.
3A, arrowheads). The TbSmee1 signal got fainter as the new FAZ elongated, and was quite
faint in the 2K2N stage. TbSmee1 was sometimes faintly observed along the full length of the
new FAZ in replicating cells, and upon careful observation, a TbSmee1 signal could also be
detected at the tip of the old FAZ as well, even in 1K1N cells.
212

Using structured illumination microscopy, TbSmee1 was observed to be slightly in front of the
tip of the new FAZ filament (Fig. 3B). This suggested that TbSmee1 might be present at the
"groove". The groove is a structure involved in remodelling of the microtubule cytoskeleton
during cell replication in bloodstream form *T. brucei* (Hughes et al., 2013; Smithson et al.,
2022). It can be detected using the monoclonal antibody DOT1. TbSmee1 appeared to be
enveloped by the DOT1 labelling, confirming its presence at the groove (Fig. 3C). In summary,
in replicating cells TbSmee1 is present at the groove in addition to the hook complex, and
travels in front of the newly-assembling FAZ filament.
221
222
223

224 **The C-terminal part of TbSmee1 is required for targeting to the hook complex.**

To determine what parts of TbSmee1 primary structure are responsible for targeting to the hook complex, a series of truncation constructs based on the predicted domain architecture of TbSmee1 were designed (Fig. 4A). Cell lines were generated that could inducibly express each of these truncations with an N-terminal Ty1 tag. The presence of the ectopic transgenes in the genomes of these cells was confirmed by PCR analysis of genomic DNA (Fig. S5A). To determine the localisation of each construct, and whether it could associate with the cytoskeleton, detergent-extracted cells were analysed using immunofluorescence microscopy. It should be noted that both endogenous *SMEE1* alleles were still present, meaning that targeting of the truncation constructs was being assayed in the presence of the endogenous Smee1 protein.

TbSmee1(161-766) correctly localised to the hook complex, indicating that Domain1 is not necessary for targeting (Fig. 4B). Of note, no dominant negative effects on cell growth were observed upon overexpression of the TbSmee1(161-766) construct. TbSmee1(265-766) also localised correctly, indicating that the predicted linker region between Domain1 and Domain2 is not necessary for localisation (Fig. 4B). No other truncations were observed to localise to the hook complex, making TbSmee1(265-766) the smallest construct to localise correctly. Expression of all TbSmee1 truncations was confirmed by immunoblotting with anti-TbSmee1 and anti-Ty1 antibodies (Figs 4C, S5A). Quantification of the immunoblots indicated that most were expressed at around the level of the endogenous protein (Fig. 4D).

Interestingly, TbSmee1(161-766) also localised to the FAZ tip in replicating cells (Fig. S5B). TbSmee1(265-766) did not localise to the FAZ tip, suggesting that the linker region between Domain1 and Domain2 (aa 161-264) might be required for targeting to this structure. In support of this hypothesis, TbSmee1(2-400) was found to localise to the FAZ tip despite not localising to the hook complex (Fig. S5C).

TbSmee1 is essential for the viability of BSF *T. brucei*, and its depletion causes gross enlargement of the flagellar pocket.

The effects of TbSmee1 depletion were analysed using tetracycline-inducible RNAi. Induction of RNAi (Tet+) resulted in a rapid cessation of population growth after around 24 h (Fig. 5A). Visual inspection of the stalled populations at 48 h and 72 h post-induction showed widespread lysis, confirming that TbSmee1 is essential for the viability of bloodstream form cells in vitro. A shorter timecourse with higher sampling frequency indicated that the growth defect began after around 20 h of RNAi, and was already clear at 24 h (Fig. 5A, inset).

TbSmee1 depletion was confirmed at the single-cell level using immunofluorescence microscopy (Fig. 5B). TbSmee1 signal was lost from both the hook complex and FAZ tip. The kinetics of TbSmee1 protein depletion on either side of the onset of the growth defect were assessed by immunoblotting. TbSmee1 protein levels were reduced to ~ 20-25% at 15 h and 18 h post-induction, with a further reduction to around 10-15% from 21 h onwards (Fig. 5C).

The effect of TbSmee1 knockdown on cell cycle progression was assessed by quantifying the numbers of 1K1N, 2K1N, 2K2N, and abnormal cell types in the same time window (Fig. S6A). An increase in 2K1N and a decrease in 1K1N cells was visible from 21 h post-induction onwards, followed by an increase in 2K2N cells at the 24 h timepoint. This indicated that cell cycle progression was inhibited from around 21 h after induction of RNAi, correlating with the onset of the growth defect. To summarise, TbSmee1 protein was already significantly depleted at 15 h, with subsequent effects on population growth and cell cycle progression visible from around 21 h onwards.

The effects of TbSmee1 depletion on a panel of hook complex proteins and components of the centrin arm and flagellar pocket collar were systematically evaluated (Fig. S7A-E). TbSmee1 depletion for 24 h did not result in observable effects on the expression levels or localisation of any of the candidates. No flagellum detachment was observed either. Conversely, depletion of TbMORN1 for just 16 h resulted in reductions in the levels of TbSmee1 and TbStarkey1 (Fig. S7B, F-H). The lack of observable structural changes to the hook complex, flagellar pocket collar, and centrin arm following TbSmee1 depletion implied that the lethal phenotype might be due to a loss of a specific cellular function, rather than destabilisation of structural complexes.

To determine the ultrastructural changes caused by TbSmee1 depletion, the cells were imaged by electron microscopy after high-pressure freezing. High-pressure freezing was used, as this approach gives better morphological preservation than chemically fixing the cells in the growth media. The most obvious morphological effect of TbSmee1 depletion was the gradual accumulation of cells with enlarged flagellar pockets (Fig. 5D). This presumably was the cause of the lethality phenotype, as depletion of TbMORN1 was also shown to result in progressive enlargement of the flagellar pocket until cells rounded up and lysed (Morriswood and Schmidt, 2015). Interestingly however, depletion of TbStarkey1 also frequently resulted in the generation of cells with enlarged flagellar pockets, despite causing no growth defect (Fig. S6B-D). This, together with the recently-published characterisation of Bhalin (Broster Reix et al., 2021), means that depletion of four separate hook complex proteins - TbMORN1,

TbSmee1, TbStarkey1, Bhalin - results in flagellar pocket enlargement, though with varying

magnitudes of effect.

Flagellar pocket enlargement is an early consequence of TbSmee1 depletion.

Although flagellar pocket enlargement appears to be a consistent phenotype resulting from

hook complex protein depletion, it is not an uncommon phenotype in bloodstream form RNAi

cells. Importantly, it can be either a direct or an indirect consequence of protein depletion (Ali

et al., 2014; Allen et al., 2003; Hall et al., 2004; Hall et al., 2005; Price et al., 2007).

To determine whether flagellar pocket enlargement was an early/direct consequence of

TbSmee1 depletion (i.e. occurring very soon after or even before the onset of the growth

defect), a fluorescent, fixable 10 kDa dextran reporter was used. Dextran is a polysaccharide

that traffics in the fluid phase and is well-established as an endocytic marker. Control (- Tet)

and TbSmee1-depleted (+ Tet) cells were incubated on ice in order to block endocytosis

(Brickman et al., 1995). The cells were then incubated with the labelled dextran for 15 min to

allow it to enter the flagellar pocket, and fixed afterwards (Fig. 6A). The magnitude of the

dextran signal should therefore be proportional to flagellar pocket volume. Visual analysis of

the cells confirmed that dextran visibly labelled the flagellar pocket, with a much greater signal

seen for cells with an enlarged flagellar pocket (Fig. 6B).

Flow cytometry was then used for high-throughput, unbiased, and quantitative analysis of the

cells (Fig. 6C). At 15 h post-induction, no difference between TbSmee1-depleted cells and

controls was observed. At 18 h and 21 h post-induction a slight "shoulder" on the +Tet traces

became visible, indicating the emergence of cells with higher fluorescence values than seen

in controls. By 24 h post-induction there was a clear hump visible in the traces, indicating a

subpopulation with fluorescence values sometimes two orders of magnitude greater than

those seen in control cells (Fig. 6C, arrow). Quantification of the flow cytometry data from

multiple experiments showed that average fluorescence intensity of the whole + Tet population

was noticeably higher than controls at 21 h and 24 h post-induction (Fig. 6D). Even at 18 h

post-induction, i.e. before the onset of the growth defect, there was already a clear increase

in the average fluorescence intensity in the + Tet population (Fig. 6D, blue arrow). This strongly

suggests that flagellar pocket enlargement is an early and probably direct consequence of

TbSmee1 depletion.

TbSmee1 depletion results in impaired flagellar pocket access of surface-bound cargo.

It was previously shown that the ability of large cargo to enter the flagellar pocket is affected

after knockdown of the hook complex protein TbMORN1 (Morriswood and Schmidt, 2015).

Specifically, the fluid phase marker 10 kDa dextran accumulates in the enlarged flagellar

pocket of TbMORN1-depleted cells, while larger fluid phase cargo such as BSA-5 nm gold

and large surface-bound cargo such as ConA (which binds to surface glycoproteins) do not

access the flagellar pocket lumen.

To test whether TbSmee1 knockdown also impairs flagellar pocket access, TbSmee1 RNAi

cells were incubated with both dextran and ConA to simultaneously monitor the uptake of fluid

phase and surface-bound cargo. The cells were first incubated on ice to block endocytosis,

and then with the reporters (also on ice) to allow ConA to bind and dextran to enter the flagellar

pocket. The cells were then shifted to 37°C to reactivate endocytosis, and subsequently fixed

and imaged (Fig. 7A).

In control (- Tet) cells, both dextran and ConA strongly overlapped in the part of the cell

corresponding to the endosomal/lysosomal system (Fig. 7B, -Tet, arrow). To confirm that the

internalised material was being trafficked to the lysosome, the cells were labelled with

antibodies specific for the lysosomal enzyme p67 (Kelley et al., 1999). The dextran reporter

was not compatible with immunolabelling, but the ConA signal clearly overlapped with the

lysosome marker p67 (Fig. S8, -Tet cells). In TbSmee1-depleted (+ Tet) cells, dextran filled

the enlarged flagellar pocket while ConA was restricted to one or two small foci that appeared

to be on the cell surface (Fig. 7B, +Tet, arrow). No overlap was seen between the dextran and

ConA labels, suggesting that ConA was not able to enter the flagellar pocket. In addition, no

overlap was observed between ConA and the lysosome marker p67 (Fig. S8, +Tet cells).

To quantify these observations, the cells were grouped into 4 categories: 1, complete overlap

between the two labels; 2, partial overlap between the two labels; 3, no overlap between the

two labels; 4, whole-cell labelling. Whole-cell labelling occurs when the cell has lost integrity,

and labelling is found throughout the cytoplasm. Control (-Tet) cells all showed complete

overlap between the dextran and ConA, while cells depleted of TbSmee1 for 24 h showed

>25% of cells with partial or no overlap between the cargoes (Fig. 7C).

To further validate these observations, the experiments were repeated using fluorescently-

labelled BSA (Fig. 7D). BSA traffics in the fluid phase and is a physiological cargo, unlike

ConA (Coppens et al., 1987). As expected, in control cells there was strong overlap between

dextran and BSA from the endosomal/lysosomal system (Fig. 7E, -Tet, arrow). In TbSmee1-

depleted cells, there was once again no overlap between the reporters, indicating that BSA

was unable to access the enlarged flagellar pocket. Surprisingly, despite BSA being fluid-
phase cargo there was a punctate signal analogous to that seen with ConA, suggesting that
it could at least partly bind to the cell surface (Fig. 7E, +Tet, arrow). Quantification of the data
showed that the effect on BSA was more pronounced than that seen for ConA, with >55% of
TbSmee1-depleted cells showing no or only partial overlap between the two reporters after 24
h of RNAi (Fig. 7F).
378
379
380 **Endocytosis is required for flagellar pocket access of surface-bound cargo.**
381 The inability of surface-bound cargo to enter the flagellar pocket of TbMORN1- and TbSmee1-
382 depleted cells could either be due to a defect in endocytosis (indicated by the enlargement of
383 the flagellar pocket), or a secondary effect.
384
To distinguish between these possibilities, the assays were repeated following clathrin
depletion (Fig. 8). Clathrin is an essential endocytic coat protein and its loss results in a block
in endocytosis. This results in the expected enlargement of the flagellar pocket and ultimately
cell lysis, and in fact this phenotype was first described as a result of clathrin depletion (Allen
et al., 2003). Studies of clathrin in trypanosomes have however focused on whether or not
cargo is endocytosed, but not whether or not it can enter the flagellar pocket prior to
encountering the block in internalisation. In control cells, as expected, there was strong
overlap between the dextran and ConA cargoes in the region of the endolysosomal system
(Fig. 8B, -Tet, arrow). In clathrin-depleted cells however, there was again no overlap observed
between the two reporters, and ConA did not appear able to access the enlarged flagellar
pocket (Fig. 8B, +Tet, arrow). Quantification of the data showed that >50% of clathrin-depleted
cells showed either partial or no overlap between the two reporters after 19 h of RNAi (Fig.
8C). The same effect was seen in assays using dextran and BSA, with little to no overlap
between the reporters in clathrin-depleted cells (Fig. 8D-F). These results indicated that
inhibition of endocytosis by itself resulted in the failure of either ConA or BSA to enter the
enlarged flagellar pocket.
401
Endocytosis is required for flagellar pocket access of surface-bound antibodies.
One of the main functions of endocytosis in trypanosomes is to remove any antibodies bound
to the surface glycoprotein coat. To determine whether the effects observed above using ConA
and BSA reporters also apply to antibodies, another set of uptake assays were carried out. As
the uptake of surface-bound antibodies occurs in a matter of seconds, a slightly modified
assay protocol was used (Fig. 9A). The cells were first incubated on ice to block endocytosis,
and then incubated with antiserum specific for the variant surface glycoprotein (VSG), which

accounts for the overwhelming majority of the surface glycoprotein coat. A sample of cells
were fixed at this timepoint (t=0) to confirm surface binding of the antibodies. The remaining
cells were washed (at low temperature), after which a second sample was fixed (t=1). The
remaining cells were shifted to 37 °C to reactivate endocytosis and incubated for 2 minutes to
allow the antibodies to enter the flagellar pocket and then be internalised by endocytosis.
414
The distribution of anti-VSG antibodies in control and TbSmee1-depleted cells was analysed
using microscopy (Fig. 9B). In control (-Tet) cells, anti-VSG signals were observed at the cell
surface (t=0), then at the location of the flagellar pocket (t=1) and finally at the
endosomal/lysosomal system after allowing endocytosis (t=2) (arrows, upper panels). In
TbSmee1-depleted cells (+Tet), no anti-VSG signals were observed in the enlarged flagellar
pockets (t=1, t=2) (arrows, lower panels). It should be noted that there was considerable loss
of anti-VSG signal between the t=0 and t=2 timepoints, and half of the TbSmee1-depleted
cells displaying a phenotype had no observable signal at all. This obviously makes the
conclusions from these experiments somewhat tentative.
424
To quantify the results, the incidence of morphologically abnormal cells was counted, and then
the various distributions of anti-VSG antibodies were classified into three categories (Fig. 9C).
Of the 35 % of cells in the TbSmee1-depleted population that displayed a phenotype after 24
h of RNAi, 26 % were classified as being completely rounded up ("BigEye") while 7 % had the
less-developed "enlarged pocket" state. The anti-VSG antibody distributions were grouped
into three categories, respectively either a dot signal, another signal (usually a dot with other
weaker labelling), and no signal. Although most of the morphologically-abnormal cells had no
anti-VSG signal at all, TbSmee1-depleted cells with clearly internalised anti-VSG of the kind
observed in control cells were never seen.
434
435 In summary, the inhibition of endocytosis caused by depletion of TbSmee1 prevents the
436 internalisation of surface-bound anti-VSG antibodies, and appears to also prevent their entry
437 into the enlarged flagellar pocket, consistent with the results obtained using ConA and BSA.
438
439
440
441
Discussion
In this study, the hook complex component TbSmee1 was characterised in bloodstream form
T. brucei. It is only the third hook complex protein to be characterised in this life cycle stage
after TbMORN1 and Bhalin, and like them is essential for the viability of the cells in vitro

(Broster Reix et al., 2021; Morriswood and Schmidt, 2015). The localisation of TbSmee1 was
found to be identical to that previously documented in the procyclic life cycle stage of *T. brucei*
(Figs 2, 3) (Perry et al., 2018). It is constitutively present at the shank part of the hook complex
and probably also the FAZ tip, and additionally localises to the tip of the growing new FAZ in
replicating cells. At the time of the preprint of this work being published, TbSmee1 was the
first protein besides the DOT1 antigen and FLAM3 to be shown to localise to the groove
structure associated with the tip of the growing new FAZ (Sunter et al., 2015a). A number of
additional proteins present at the groove have since been identified (Smithson et al., 2022).
454
The analysis of truncation constructs suggested that different parts of the TbSmee1 primary
structure are responsible for targeting to the two locations (Figs 4, S5). Targeting to the hook
complex is mediated by the second and third predicted domains of TbSmee1. It is currently
not clear whether the linker region between these two domains (aa 401-630) is also required,
although the AlphaFold prediction for this region shows only unstructured loops. Nonetheless,
this would be an obvious hypothesis to test in future work. Targeting to the FAZ tip appears to
be mediated by the linker region (aa 161-265) between the first and second predicted
domains. This is somewhat surprising, as there are few highly-conserved residues in this
region (Fig. S1), and AlphaFold again predicts an unstructured loop for this region. It should
be noted that while aa 161-265 have been shown to be necessary for FAZ tip targeting, they
have not yet been shown to be sufficient - this again would be an obvious future experiment
should this line of enquiry be pursued further. The very low observed expression levels for
TbSmee1(2-160) and TbSmee1(265-640) also mean that the negative results obtained with
these constructs should be treated with some caution. Given that it is unknown whether
Smee1 can self-interact, the presence of the endogenous protein in these targeting
experiments is an additional factor that could be addressed in follow-up work. It is unclear
what role phosphorylation of TbSmee1 plays in its localisation, but given that it is both a
substrate and putative direct binding partner of the mitotic kinase TbPLK, it is possible that its
phosphorylation status may be related to its localisation to the groove structure.
474
Phenotypic characterisation of TbSmee1 showed that its depletion results in the enlargement
of the flagellar pocket with concomitant effects on the entry of macromolecular cargo (Figs 5-
7). The effects on the entry of macromolecular cargo are identical to those seen following
TbMORN1 and Bhalin depletion (Broster Reix et al., 2021; Morriswood and Schmidt, 2015). A
clear limitation of all the TbSmee1 RNAi experiments was the inability to specifically isolate
cells with reduced levels of TbSmee1. As the cells were being assayed very soon after the
onset of the growth defect, this meant that all assays were being conducted against a
background of cells exhibiting different and often much lower levels of TbSmee1 depletion.

This could be addressed in future experiments either by using a fluorescent tag on TbSmee1
to enable sorting of the subpopulation of depleted cells, or by synchronising the cells prior to
the induction of RNAi. Nonetheless, there was still extremely good agreement between the
immunoblotting, growth curve, and cell cycle analysis data.
487
At 18 h post-induction of RNAi, TbSmee1 protein levels were reaching their minimum but there
was as yet no observable effect on cell cycle progression or population growth. From around
20-21 h post-induction, when TbSmee1 levels reached 10-15% of controls, cell cycle
progression slowed, resulting in an increase in 2K1N and 2K2N cells. This was mirrored by a
slowing of population growth. Systematic analysis of a number of marker proteins for the hook
complex, the centrin arm, and the flagellar pocket collar showed that none were strongly
affected in terms of their protein levels or localisation by the depletion of TbSmee1. This
argues that the effects seen upon TbSmee1 depletion at these early timepoints are probably
due to a loss of function, rather than due to structural perturbations in the hook complex.
497
In the previous characterisation of TbSmee1 in the procyclic form life cycle stage, depletion of
TbSmee1 was found to cause structural changes to the hook complex and altered TbMORN1
distribution patterns (Perry et al., 2018). These analyses were however done after 6 days of
RNAi induction, when TbSmee1 levels were already undetectable from 1 day onwards (a
growth defect was observable from 3 days of RNAi onwards). It is very likely that similar
changes would be observed in the bloodstream form stage at later RNAi timepoints, although
the essential nature of TbSmee1 in bloodstream form cells might make this analysis difficult.
Regardless, the focus in this study was the immediate and early effects of TbSmee1 depletion,
which appear to be focused on the flagellar pocket.
507
Flow cytometry analysis indicated that the flagellar pocket enlargement began around 18 h
post-induction, prior to the effects on population growth and cell cycle progression. This
therefore suggests that flagellar pocket enlargement is an early and likely direct consequence
of TbSmee1 depletion, with the effects on cell cycle progression coming afterwards. One
plausible hypothesis to account for these observations is that the spatial problems produced
by flagellar pocket enlargement impair cell cycle progression. The relative lack of cell cycle
checkpoints means that "monster" cells with re-duplicated organelles are then produced. The
internal pressure caused by continuing enlargement of the flagellar pocket forces the cells into
an increasingly spherical shape and eventually cell viability is lost. In this context, the lack of
a strong phenotype following TbStarkey1 depletion, despite evidence from electron
microscopy that there is some enlargement of the flagellar pocket (Fig. S6), is puzzling. Cells
with enlarged flagellar pockets in TbStarkey1-depleted populations were however not

immediately obvious at the light microscopy level, so the prevalence of this phenotype might
have been overestimated during image acquisition on the electron microscope.

The enlargement of the flagellar pocket was accompanied by a cargo access defect previously
documented for TbMORN1 and Bhalin - small fluid-phase cargo such as 10 kDa dextran fills
the enlarged flagellar pocket, while larger cargo such a ConA and BSA is unable to (Broster
Reix et al., 2021; Morriswood and Schmidt, 2015). In TbSmee1-depleted cells, BSA was
observed in one or more small foci next to the flagellar pocket. This was surprising, as BSA is
known to traffic in the fluid phase and should not therefore be binding to the cell surface. BSA
was however also observed to associate with the flagellar pocket membrane in an electron
microscopy-based study of cargo uptake (Gadelha et al., 2009). In a classic radiochemical
study of cargo uptake in *T. brucei,* it was observed that "at low concentration, a small
adsorbtive component may become prevailing" in assays using BSA (Coppens et al., 1987).
As the experiments conducted here and in the 2009 paper were using BSA well below
physiological concentrations, it seems likely that these foci represent the small adsorbtive
component that was indirectly observed in the biochemical assays. Therefore, in these assays,
the ConA and BSA probes both report on the behaviour of surface-bound cargo, while the
dextran reports on fluid-phase uptake.

As ConA is not a physiological cargo, the entry of anti-VSG antibodies into the flagellar pocket
was also investigated. These assays turned out to be unexpectedly difficult, owing to much of
the signal being lost during the wash steps. This may be related to the recently-reported high
rate of VSG shedding (Garrison et al., 2021). Thus, the conclusions obtained here are
necessarily tentative. Nonetheless, the t=0 and t=2 timepoints would appear to match
expectations: in control cells, the anti-VSG antibodies are taken up by endocytosis, while in
TbSmee1-depleted cells any remaining signal was found outside the area of the enlarged
flagellar pocket and apparently still on the cell surface. It would be informative to repeat these
experiments using a defined amount of affinity-purified antibody in order to minimise the need
for washing, but this is outside the scope of this study.

When the TbMORN1 phenotype was characterised, this was the first time that a protein had
been shown to play a role in the entry of surface-bound cargo into the flagellar pocket.
Subsequent characterisation of Bhalin and the work here on TbSmee1 shows that this cargo
entry defect appears to be a consistent effect after enlargement of the flagellar pocket.
Furthermore, the fact that the same cargo entry defect is seen upon depletion of clathrin
suggests that this is a previously-unobserved feature of all flagellar pocket enlargement

("BigEye") phenotypes. Therefore, active endocytosis appears to be required for the entry of
surface-bound cargo into the flagellar pocket.

It should be noted that the results obtained in the clathrin RNAi experiments contradict
previously-published observations, where ConA was claimed to accumulate inside the
enlarged flagellar pocket of clathrin-depleted cells (Allen et al., 2003). Re-examination of
previously-published data from clathrin RNAi cells suggests however that the effect has
always been present, with ConA signal (and indeed other cargoes) absent from the flagellar
pocket interior and predominantly concentrated in one or more foci just outside it (see for
example Fig. S4 in (Allen et al., 2003), and Fig. S2 in (Zoltner et al., 2015)). This oversight is
understandable, given that the focus in these clathrin papers was on whether or not cargo
endocytosis was occurring, and not whether the cargo was able to access the flagellar pocket
(Mark Field, personal communication). Importantly, it also means that similar observations
have been made by (at least) two independent groups.

It has been known for some time that surface-bound cargo in the lumen of the flagellar pocket
is associated with the membrane underlying the microtubule quartet (Brickman and Balber,
1990; Gadelha et al., 2009). It has also previously been shown that upon inhibition of
endocytosis, tomato lectin, wheat germ agglutinin, and BSA accumulate on this part of the
flagellar pocket membrane (Gadelha et al., 2009). What is intriguing about the observations
with TbSmee1 and clathrin RNAi cells here is that they show that fluid phase cargo (dextran)
is still able to enter the flagellar pocket even when endocytosis is inhibited, so the effects are
specific to surface-bound cargo. Given that the ConA and BSA material becomes
concentrated into one or more foci that do not overlap with the dextran signal (Figs 7,8), this
suggests that the majority of cargo is not able to enter the flagellar pocket lumen at all when
endocytosis is inhibited.

This begs two important questions: why is it that an inhibition of endocytosis apparently
hinders the entry of surface-bound cargo into the flagellar pocket? And why does the depletion
of hook complex proteins - which are spatially removed from the actual sites of clathrin-coated
vesicle formation - cause an endocytosis defect? The prevailing model for several years has
been that hydrodynamic flow, driven by flagellar motility, is responsible for sorting surface-
bound cargo to the posterior end of the cell where the flagellar pocket is located (Engstler et
al., 2007). Once there, the surface-bound material enters the flagellar pocket through a narrow
channel where the flagellar membrane and the flagellar pocket neck membrane are not as
closely apposed (Gadelha et al., 2009). The results here suggest that the posterior sorting

mechanism still functions when endocytosis is inhibited, but that the transit of surface-bound
material through the channel and into the flagellar pocket might be impeded.

In this context, it is worth considering the somewhat confusing anti-VSG uptake results
obtained at the t=1 timepoint in control cells, where the anti-VSG signal came from a single
spot adjacent to the kinetoplast. As a temperature block was in place to prevent endocytosis,
this would at first sight suggest that the anti-VSG antibodies can enter the flagellar pocket in
the absence of endocytosis, and that the entire hypothesis proposed here is invalid. Given
however that the flagellar pocket is less than 1 µm across, it is possible that there is insufficient
resolution to distinguish the entrance to the flagellar pocket from the flagellar pocket itself in
control/wild-type cells. In this interpretation, it is only when the flagellar pocket is expanded
due to the BigEye phenotype that these two localisations can be resolved at normal widefield
resolution. Higher-resolution imaging of these cells is therefore an important target for follow-
up work.

An earlier hypothesis for hook complex function was that it was somehow maintaining the
integrity of the channel in the flagellar pocket neck. This remains possible, and determining
the exact site of cargo blockade, as well as imaging the microtubule quartet, will be key goals
for future work. Nevertheless, the fact that the same phenotype is obtained following depletion
of clathrin suggests that the hook complex is instead indirectly affecting endocytosis. This
could potentially be by affecting either the localisation or activity of a number of lipid kinases,
at least two of which are known to localise to the hook complex (Dean et al., 2017; Demmel
et al., 2014). The activity of these enzymes could then be licensing the flagellar pocket
membrane for endocytosis, for instance by generating the essential endocytic cofactor
phosphatidylinositol-(4,5)-bisphosphate. Subsequent internalisation of membrane by
endocytosis would then assist to pull in more membrane currently at the flagellar pocket
entrance. How this would be integrated with the activity of the exocytic pathway is, however,
unclear.

Thus, while hydrodynamic flow may be responsible for concentrating cargo at the entrance to
the flagellar pocket, endocytic activity seems required for the entry of the cargo into the
flagellar pocket itself, and endocytosis may assist or be responsible for pulling material in
through the channel. Exploring these mechanisms is likely to be a fascinating area for future
enquiry.

Materials and Methods

Recombinant protein expression and purification

The TbSmee1(1-400) open reading frame was amplified from *Trypanosoma brucei brucei* strain Lister 427 genomic DNA by PCR. The PCR product was ligated into the p3NH expression vector, which encodes an N-terminal His6 tag, using sequence and ligation-independent cloning (Li and Elledge, 2012). The plasmid was used to transform *E. coli* strain Rosetta II (DE3)pLysS by heat shock, and individual colonies were subsequently grown at 37 °C in the presence of 100 µg/ml kanamycin to an OD600 ~ 0.8–1.0. Recombinant protein expression was induced by the addition of 50 µM IPTG, and the cells were then incubated overnight at 20 °C with shaking. Cells were harvested by centrifugation (5000× g for 30 min). The pooled pellet from 6 L of cell culture was resuspended in 300 ml of lysis buffer (50 mM Hepes pH 7.0, 500 mM NaCl, 20 mM imidazole, 5% glycerol, 0.5% Triton-X, 1 mM TCEP, 200 mM PMSF and protease inhibitor cocktail). Pellet emulsions were first homogenised by mixing on ice using a T 10 basic Ultra-Turrax dispersing instrument (IKA). Final lysis was then achieved with sonication on ice, using 3 cycles of 3 min at 50% strength. Lysates were clarified by centrifugation (18,000× g, 45 min, 4 °C). The lysates were added to a HiTrap Chelating HP 5 ml column (GE Healthcare) equilibrated with buffer A (20 mM Hepes pH 7, 300 mM NaCl, 40 mM imidazole, 2% glycerol, 1 mM TCEP), and eluted with a 100% step gradient of buffer B1 (20 mM Hepes pH 7, 300 mM NaCl, 400 mM imidazole, 2% glycerol, 1 mM TCEP). Selected peak fractions were examined by SDS-PAGE for protein content and purity. Fractions containing a dominant band at approximately 46 kDa were pooled and concentrated using Amicon Ultra centrifugal filter units with 10K pore size (MerckMillipore) according to the manufacturer's instructions. The His6 tag was removed by 3C protease during overnight dialysis in dialysis buffer (20 mM Tris-HCl pH 7, 300 mM NaCl, 2% glycerol, 1 mM DTT). Significant losses were incurred during this step. The TbSmee1(1-400) was applied to a previously equilibrated HiTrap Chelating HP 5 ml column charged with 50 mM $CoCl_2$ and coupled to a GSTrap HP 1 ml column (both GE Healthcare). Buffer A was used for equilibration. TbSmee1(1-1400) was mostly collected from the flow-through and a few initial collected fractions. Selected peak fractions were examined by 15% SDS-PAGE for protein content and purity. Fractions containing a dominant band at approximately 44 kDa were pooled and concentrated in Amicon Ultra centrifugal filter units (10K pore size) according to the manufacturer's instructions. Finally, TbSmee1(1-400) concentrates were applied to a previously equilibrated HiLoad 16/600 Superdex 200 pg column (GE Healthcare) pre-equilibrated in dialysis buffer. Flow speed was adjusted to 1 ml/min and fractions of 1.5 ml were collected. Fractions corresponding to the targeted chromatographic peak were examined

for protein content by 15% SDS-PAGE, pooled accordingly to their purity, concentrated, and
stored at -80 °C.

Antibody generation and affinity purification

Purified recombinant TbSmee1(1-400) was used for the generation of two polyclonal rabbit
antisera (Eurogentec). Antisera (303, 304) were initially affinity-purified against the
TbSmee1(1-400) antigen, but the neat antisera were later found to show high specificity and
were also occasionally used. A third polyclonal antibody (508) was generated against two
TbSmee1 peptides (Eurogentec) and affinity purified using the peptide antigens immobilised
on a Sulfolink affinity column (ThermoFisher). Results shown were predominantly obtained
using affinity-purified "303" and "304" anti-TbSmee1 antibodies; most immunoblotting data
were generated using the "304" affinity-purified antibodies, as these had the lowest
background in this application, while labelling was identical for "303" and "304" in
immunofluorescence experiments. The results obtained with all three antibodies were
consistent. The anti-Starkey1 anti-peptide rabbit polyclonal antibodies were generated and
affinity purified in the same way.

Antibodies

The following primary antibodies have been described previously: rabbit anti-TbMORN1
(Morriswood et al., 2013), rabbit anti-TbBILBO1 (Esson et al., 2012), mouse anti-Ty1 ("BB2")
(Bastin et al., 1996), mouse anti-TbLRRP1 (Zhou et al., 2010), mouse anti-TbCentrin4 ("6C5"
(Ikeda and de Graffenried, 2012)), mouse anti-TbCentrin2 ("2B2H1") (de Graffenried et al.,
2013), mouse anti-TbFAZ1 ("L3B2") (Kohl et al., 1999), mouse anti-PFR1,2 ("L13D6") (Kohl
et al., 1999), mouse anti-FAZ filament ("DOT1")(Woods et al., 1989), rabbit anti-VSG(221)
(Batram et al., 2014). The following antibodies came from commercial sources: goat anti-
rabbit(IRDye800CW) (LI-COR), goat anti-mouse(IRDye680LT) (LI-COR), goat anti-rabbit and
anti-mouse antibodies conjugated to AlexaFluor dyes (Molecular Probes).

Cell culture

Wildtype Lister 427 (monomorphic) BSF cells were cultured in HMI-9 medium (Hirumi and
Hirumi, 1989) supplemented with 10 % foetal bovine serum (FBS), 100 U/ml penicillin and 0.1
mg/ml streptomycin at 37°C and 5% CO_2. The SM (single marker) cells (Wirtz et al., 1999)
were cultured in the presence of G418 (2.5 µg/ml). Population density was monitored using a
Z2 Coulter Counter (Beckman Coulter), and kept below $2x10^6$ cells/ml.

Generation of transgenic cell lines

The Ty1-TbSmee1 endogenous replacement cell line was generated by transfection of 427
cells with a targeting fragment containing 285 bp of the TbSmee1 5' untranslated region
(UTR), a blasticidin resistance gene, the intergenic region of alpha- and beta-tubulin, the
sequence for a triple Ty1-tag and the first 399 bp of the TbSmee1 open reading frame (ORF)
without the start codon. Clones were selected by growth in medium containing 5 µg/ml
blasticidin. RNAi target sequences were chosen using RNAit (Redmond et al., 2003).
TbSmee1 RNAi cells were generated by cloning the RNAi target sequence into the pGL2084
plasmid (Jones et al., 2014) and then transfecting 2T1 cells with the linearised plasmid (Alsford
et al., 2005). Clones were selected by growth in medium containing phleomycin (2.5 µg/ml)
and hygromycin (5 µg/ml). TbStarkey1 RNAi cells and TbCHC RNAi cells were generated by
cloning the relevant RNAi target sequence into the p2T7_TAblue plasmid (Alibu et al., 2005)
and then transfecting SM cells with the linearised plasmid. Clones were selected by growth in
medium containing G418 (2.5 µg/ml) and hygromycin (5 µg/ml). Ty1-TbSmee1 truncation
constructs were cloned into the pLew100_v5-Hyg plasmid using in vivo assembly (Watson
and Garcia-Nafria, 2019). SM cells were transfected with the linearised plasmid. Clones were
selected by growth in medium containing G418 (2.5 µg/ml) and hygromycin (5 µg/ml). For
transfection >2.5×10^7 cells of the parental strain were washed and resuspended in 100 µl
transfection buffer (90 mM Na_2PO_4, 5 mM KCl, 0.15 mM $CaCl_2$, 50 mM HEPES [4-(2-
hydroxyethyl)-1-piperazineethanesulfonic acid]-NaOH, pH 7.3) containing 10 µg DNA and
transfected by electroporation using an AMAXA Nucleofector® Device (Lonza) with program
"X-001 free choice". Transfected cells were incubated in 50 ml HMI-9 medium without
selection overnight. The next day, drug selection was applied and clones were selected by
limiting dilution. At least three separate clones of all cell lines were isolated to control for
biological variability. Integration of targeting fragments at endogenous loci or the presence of
Ty1-TbSmee1 truncation constructs in the genome were confirmed by PCR analysis of
genomic DNA. Genomic DNA was isolated using a DNeasy Bloody & Tissue kit (QIAGEN)
and relevant products were amplified by PCR. All cloned constructs used for cell line
generation had their DNA sequence confirmed by sequencing.
730

Immunoblotting
For generation of dephosphorylated whole-cell lysates, cells were harvested by centrifugation
(1000xg, 10 min) and the cell pellet was resuspended in 1 ml vPBS (PBS, 46 mM sucrose, 10
mM glucose) containing EDTA-free protease inhibitors (Roche). The washed cells were
pelleted by centrifugation (750xg, 4 min). The cells were then resuspended in lysis buffer
(0.5% IGEPAL, 0.1M PIPES-NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-
free protease inhibitor cocktail) to a final concentration of $4x10^5$ cells/µl and incubated for 15
min at RT on an orbital mixer to allow dephosphorylation to occur. SDS-loading buffer was

then added to a final concentration of 2×10^5 cells/µl, the samples were further denatured by
boiling (100°C, 10 min), and then stored at -20°C. SDS-PAGE was carried out using a Mini-
Protean Tetra Cell (Bio-Rad), and protein transfer to nitrocellulose membranes using a Mini-
Trans blot cell (Bio-Rad). Protein transfer and equal loading was confirmed using REVERT
total protein stain (LI-COR) according to the manufacturer's instructions. Membranes were
blocked using blocking buffer (PBS, 0.3% Tween 20, 10% milk)(30 min, RT, rocker). The
membranes were then incubated in primary antibodies diluted in blocking buffer (1 h, RT,
roller). After three washes in immunoblot buffer (PBS, 0.3% Tween 20) the membranes were
incubated with IRDye-conjugated secondary antibodies diluted in immunoblot buffer (1 h, RT,
roller). After another three washes in immunoblot buffer, the membranes were visualised using
an Odyssey CLx (LI-COR). Background subtraction, qualitative analysis, normalisation
relative to total protein stain, and quantification were carried out using Image Studio Lite 5.2
and Empiria Studio 1.1 (LI-COR). Dot plots were generated using the web app PlotsOfData
(Postma and Goedhart, 2019).
753
In vitro phosphatase assays
427 BSF cells were grown to approximately 1.5×10^6 cells/ml and harvested by centrifugation
(1,000xg, 10 min, 4 °C). The cells were washed in 1 ml ice-cold wash buffer (0.1M PIPES-
NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, phosphatase Inhibitor Cocktail 2)
and pelleted by centrifugation (750xg, 3 min, 4 °C). For extraction of the cytoskeletons, the
cell pellet was resuspended in 1 ml ice-cold extraction buffer (0.5% IGEPAL, 0.1M PIPES-
NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor
cocktail, phosphatase Inhibitor Cocktail 2) and incubated for 15 min on ice. The cell
suspension was inverted every 5 min. The cytoskeletons were separated from the cytoplasm
by centrifugation (750xg, 3 min, 4 °C). The cytoskeletons were washed with 0.5 ml ice-cold 1x
NEBuffer for PMP supplemented with 1 mM MnCl$_2$ and pelleted by centrifugation (750xg, 3
min, 4 °C). After resuspension in 375 µl ice-cold 1x NEBuffer for PMP supplemented with 1
mM MnCl$_2$, all following samples were taken from this stock. An input sample (0 min) of 40 µl
was taken and added to 20 µl of SDS-loading buffer. Two control samples were taken, 0.4 µl
Phosphatase Inhibitor Cocktail 2 was added, and incubated on ice and at 26°C, respectively,
for 20 min. 20 µl of SDS-loading buffer was then added. To assay for dephosphorylation, 130
µl were taken from the stock and 1 µl lambda phosphatase (400 U) was added. This sample
and the remainder from the stock were incubated at 26 °C. After 1/2/3/5/10/20 min 20 µl
samples were taken from each and were added to 10 µl SDS-loading buffer. All the samples
were boiled at 104 °C for 10 min and stored at -20 °C. Samples were analysed by
immunoblotting.
775

Fractionation

50 ml 427 BSF cells were grown to approximately 1.5×10^6 cells/ml and then harvested by centrifugation (1000xg, 10 minutes, 4°C). The cell pellet was resuspended in 1 ml vPBS, transferred to a microfuge tube, and the cells again pelleted by centrifugation (750xg, 4 min, 4°C). The supernatant was discarded and the centrifugation step was repeated to remove remaining supernatant. The cell pellet was resuspended in 200 µl extraction buffer (0.5% IGEPAL, 0.1M PIPES-NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor cocktail) and incubated for 15 minutes at RT in an orbital mixer. A 5% input sample was taken (10 µl), put into a new microfuge tube and left on ice. The extracted cells were fractionated by centrifugation (3400xg, 2 minutes, 4°C) and the supernatant transferred to a fresh microfuge tube and the exact volume noted. The tube containing the extracted cells was centrifuged again at the same settings and this second residual supernatant discarded. The cytoskeleton pellet was then resuspended in 200 µl extraction buffer. 5% samples of supernatant and pellet fractions were taken and analysed by immunoblotting. Equal fractions of I, SN, P were loaded in each lane (I ~ 1.4×10^6 cells).

Preparation of samples for immunofluorescence microscopy

Coverslips were washed in 70% ethanol and then incubated with 0.01% poly-L-lysine in a 24-well plate (>20 min, RT) and left to dry. 2×10^6 cells were taken per coverslip and transferred to 15 ml Falcon tubes. The cells were pelleted by centrifugation (1000xg, 1 min per ml of liquid, RT) in a swing-bucket centrifuge. The supernatant was removed, and the cell pellet was gently resuspended in 1 ml ice-cold vPBS + Complete. The cells were again pelleted by centrifugation (1000xg, 2 min, RT) and subsequently resuspended in 1 ml ice-cold vPBS + Complete and directly added to the coverslips. The cells were attached to the coverslips by centrifugation (1000xg, 1 min, RT), and attachment was confirmed visually. The attached cells were then incubated in 1 ml ice-cold extraction buffer (0.5% IGEPAL 0.1M PIPES-NaOH pH 6.9, 2 mM EGTA, 1 mM MgCl2, 0.1 mM EDTA, EDTA-free protease inhibitor cocktail) (5 min, on ice). The extracted cells were washed two times with 1 ml vPBS + Complete and then fixed in 1 ml ice-cold 99.9% methanol (30 min, -20°C). The fixed cells were rehydrated using 1 ml PBS. The coverslips were blocked in 1 ml 3% BSA in PBS (30 min, RT), and sequentially incubated with clarified primary and secondary antibodies diluted in PBS (1 h, RT, humidified chamber for each) with three PBS washing steps (3 x 5 min, RT, rocker) after each incubation. After the final wash, glass slides were cleaned with 70% ethanol and a spot of DAPI-Fluoromount G (Southern Biotech) was placed on the surface. The coverslips were rinsed in ddH_2O, carefully dried, and then mounted. For preparation of samples for SIM imaging, different coverslips (high precision, No. 1.5 H) were used. Cells were harvested as above, washed, and fixed in 4% paraformaldehyde solution (10 min, ice then 30 min, RT). The fixed

cells were washed, attached to coverslips, permeabilised (0.25% Triton X-100 in PBS; 5 min,

RT), and then labelled and mounted as described above.

Fluorescence microscopy

Images were acquired using a DMI6000B widefield microscope (Leica Microsystems,

Germany) with a HCX PL APO CS objective (100x, NA = 1.4, Leica Microsystems, Germany)

and Type F Immersion Oil (refractive index = 1.518, Leica Microsystems, Germany). The

microscope was controlled using LAS-X software (Leica). Samples were illuminated with an

EL6000 light source (Leica) containing a mercury short-arc reflector lamp (HXP-R120W/45C

VIS, OSRAM, Germany). Excitation light was selected by using Y3 (545/25 nm), GFP (470/40

nm), and A4 (360/40 nm) bandpass filter cubes (Leica Microsystems, Germany). The power

density, measured at the objective focal plane with a thermal sensor head (S175C, Thorlabs),

was respectively 0.749 ± 0.086, 0.557 ± 0.069, 0.278 ± 0.076 W/cm^2 for the three filters.

Emitted light was collected at ranges of 605/70 (Y3), 525/50 nm (GFP), and 470/40 nm (DAPI)

respectively. The individual exposure times and camera gains were adjusted according to the

different samples. RNAi samples (control and depleted) were imaged using identical settings.

Differential interference contrast (DIC) was used to visualise cell morphology. 3D recording of

each field of view was obtained using 40 Z-slices (step size = 0.21 μm). Fields of view were

selected in the DIC channel in order to blind the user to the fluorescence signal and

subjectively select for cells with optimum morphology. Images were captured using a DFC365

FX monochrome CCD camera (Leica, 6.45 μm pixel size). SIM images were acquired using

an Elyra S.1 SIM microscope (Zeiss) and ZEN software (Zeiss).

Fluorescence microscopy image processing and analysis

Processing was carried out using FIJI (Schindelin et al., 2012) and a custom macro for the

generation of maximum-intensity z-projections with single DIC z-slices overlaid. Overlaps

between two proteins were confirmed in individual z-slices (thickness: 210 nm). The plugin

ScientiFig was utilised to create the collage and adding the inserts (Aigouy and Mirouse,

2013). For correlation analysis between TbSmee1 and other flagellar pocket collar and/or hook

complex associated proteins, 1K1N cells were selected using both DIC and DAPI channels.

2D sum slices projections were prepared for each stack of both green and red channels. The

projections were clipped to 8-bit depth and a convoluted background subtraction was applied.

All resultant individual 2D images, without channel overlay, were analysed pairwise to check

for intensity-based correlation based on Zhang & Cordelières (Zhang and Cordelières, 2016).

The Spearman's rank correlation results were further computed into Microsoft Excel sheets

and analysed using R version 4.1.2 (R Core Team, 2022) in the environment RStudio

2021.09.1.372 (RStudio Team, 2020). The packages used for all descriptive analysis and plot

generation were ggplot2 (Wickham, 2016), openxlsx (Schauberger and Walker, 2022), and
psych (Revelle, 2023). All Fiji/ImageJ macros and R scripts for correlation analysis were
written by Alyssa Borges and are available upon request.
853

Growth curves
RNAi cells were seeded at the required starting concentration in a volume of 22 ml and divided
into two 10 ml aliquots in separate flasks. Tetracycline was added to a final concentration of 1
µg/ml in one flask to induce RNAi, and refreshed every 24 h. The population density of the
control and induced cells was measured every 24 h over a time course of 72h, or every hour
over a time course of 8 h, using a Z2 Coulter Counter (Beckman Coulter). Depletion of the
target protein was confirmed in every experiment by immunoblotting of whole-cell lysates.
861

Cell division cycle analysis
An aliquot of 10^6 cells was taken, and the cells were fixed directly in media by addition of
isothermal 25% glutaraldehyde to a final concentration of 2.5% (10 min, 37°C, gentle mixing).
The cells were then pelleted by centrifugation (750xg, 10 min). The cell pellet was
resuspended in 0.5 ml 2.5% glutaraldehyde in PBS, transferred to a microfuge tube, and
incubated at RT (30 min, gentle mixing). The cells were pelleted again by centrifugation
(750xg, 4 min), and the cell pellet resuspended in 500 µl PBS. The cells were then added to
the coverslips inside the 24-well plate and attached by centrifugation (1000xg, 1 min, RT). The
coverslips were mounted on poly-L-lysine-coated slides using DAPI-Fluoromount G (Southern
Biotech). Imaging was as described for fluorescence microscopy above, using DAPI and DIC
channels only. Cell division cycle stages (1K1N, 2K1N, 2K2N) were manually quantified from
maximum intensity projections of the DAPI signal overlaid with single DIC z-slices. Depletion
of the target protein at each timepoint was confirmed by immunoblotting of whole-cell lysates
from the same experiments.
876

Preparation and imaging of electron microscopy samples
Induced and uninduced RNAi cells were grown for 24 h to a density of 1-2x10^6 cells/ml in 50
ml medium and harvested by centrifugation (1000xg, 10 min, RT). The supernatant was
removed to 4 ml and 4 ml FBS was added. The cells were pelleted again (1000xg, 10 min,
RT) and the supernatant was removed to 200 µl. The cells were resuspended in the
supernatant, and the suspension was then transferred to a PCR tube and pelleted by
centrifugation (1,600xg, 10s, RT). The cells were then transferred into a carrier with a closed
lid to avoid air inclusions. High pressure freezing (HPF) was started immediately (Leica EM
HPM100). After HPF the samples were transferred to an AFS (Leica EM AFS2) for freeze
substitution and progressive lowering of temperature. Low temperature embedding and

polymerisation of Epon raisin (DDSA, MNA, Epon812, 2,4,6
Tris(dimethylaminomethyl(phenol))) were then carried out. Ultra-thin cuts (60 nm) were carried
out with an ultramicrotome (Leica EM UC7/FC7) and were placed on slotted grids. For
contrasting they were incubated in 2 % uranyl acetate for 8 min. Afterwards the grids were
washed 3x in ddH$_2$O (boiled to remove CO$_2$) and incubated for 5 min on 50 % Reynold's lead
citrate in a petri dish with NaOH tablets. The grids were again washed 2x in ddH$_2$O. A 200 kV
transmission electron microscope (Jeol, JEM-2100) with a TemCam F416 4k x 4k camera
(Tietz Video and Imaging Processing Systems) and EMMenu 4.0.9.31 software (Tietz) were
used. Uninduced control cells were viewed at a magnification of 12,000x, induced cells were
viewed at a magnification of 8,000x.
897

Measurement of flagellar pocket enlargement

Induced and uninduced RNAi cells at a concentration of ~5×10^6 cells/ml were harvested by
centrifugation (1000xg, 4 ºC). The cells were resuspended in 45 µl ice-cold vPBS + protease
inhibitors and incubated on ice to block endocytosis (10 min, ice). 5 µl labelled dextran (10
kDa, 50 mg/ml stock) was added and mixed by flicking. The mixture was then incubated to
allow dextran to enter and fill the flagellar pocket (15 min, on ice, dark). At the end of the
incubation, 1 ml ice-cold vPBS + protease inhibitors was added and the cells pelleted by
centrifugation (750 xg, 2 min, 4 ºC). The cell pellet was resuspended in 0.5 ml vPBS + protease
inhibitors, and the cells fixed by addition of 0.5 ml 8 % pfa and 0.1 % glutaraldehyde solution
in vPBS + protease inhibitors (20 min, on ice then 60 min, RT). The fixed cells were pelleted
by centrifugation (750 x*g*, 2 min) and washed twice with 1 ml vPBS + protease inhibitors. The
dextran signal was immediately measured by flow cytometry using a FACSCalibur (BD
Biosciences) running CellQuest ProTM Software (BD Biosciences). Later processing and
analysis was carried out using FlowJo 10.8.1. Dot plots were created using Plots of Data
(Postma and Goedhart, 2019). After flow cytometry, the remaining cells were attached to
coverslips by centrifugation, mounted on clean glass slides using DAPI-Fluoromount G
(Southern biotech), and imaged using fluorescence microscopy on the same day to confirm
flagellar pocket labelling. Depletion of TbSmee1 was confirmed in every experiment by
immunoblotting of whole-cell lysates.
917

Cargo (ConA, BSA) uptake assays

2x10^6 cells per sample were harvested by centrifugation (1000 xg, 10 min, 4 ºC) and washed
in 1 ml ice-cold vPBS. The washed cells were pelleted by centrifugation (750 xg, 2 min, 4 ºC),
and resuspended in 100 µl ice-cold vPBS. The cells were then incubated at low temperature
(20 min, ice) to block endocytosis. During this incubation, the labelled cargoes were prepared.
The 50 mg/ml dextran aliquot (10 kDa AF488 conjugate, Molecular Probes) was mixed using

a sonicator bath (10 min, 37 Hz) and then vortexed. The 0.5 mg/ml ConA aliquot (TMR
conjugate, Molecular Probes) was clarified by centrifugation (11,000 xg, 10 min, 4 ºC). Both
cargoes were kept on ice and in the dark. At the end of the incubation, 10 µl of the 50 mg/ml
dextran and 2 µl of the 0.5 mg/ml ConA were added to the chilled cells, mixed by flicking, and
incubated at low temperature (15 min, on ice, in the dark). T=0 samples were quenched and
fixed at this point; t=30 samples had an additional incubation to allow endocytosis of cargo (30
min, 37 ºC, in the dark) before quenching and fixing. To quench samples, 1 ml ice-cold vPBS
was added, and the cells were pelleted by centrifugation (750 xg, 2 min, 4 ºC). The pelleted
cells were resuspended in 50 µl vPBS by flicking, and then fixed by addition of 0.5 ml ice-cold
fix solution (4% paraformaldehyde solution, 0.1% glutaraldehyde in vPBS) (20 min, on ice, in
the dark, then 30 min, RT, in the dark). The fixed cells were pelleted by centrifugation (750 xg,
2 min, RT), washed in 2 ml vPBS, resuspended in 1 ml vPBS, and then attached to poly-L-
lysine-coated coverslips by centrifugation. Coverslips were mounted on glass slides using
DAPI-Fluoromount G (Southern Biotech) and imaged immediately. Assays using BSA
(AlexaFluor555 conjugate, Molecular Probes) were done in the same way. The 20 mg/ml BSA
aliquot was clarified by centrifugation (750 xg, 1 min, RT)), and 3 µl was added to the
cells/dextran mixture. Depletion of TbSmee1 was confirmed by immunoblotting of whole-cell
lysates in every experiment.
942
Anti-VSG uptake assay
2×10^6 cells per sample were harvested by centrifugation (1000 xg, 1 min per ml of liquid + 1,
4 °C) and washed in 1 ml ice-cold vPBS. The washed cells were pelleted by centrifugation
(1000 xg, 3 min, 4 °C), resuspended in 50 µl ice-cold vPBS and incubated at low temperature
to block endocytosis (20 min, ice). During incubation, the primary and secondary antibodies
were clarified by centrifugation (10,000 xg, 10 min, 4 °C). The primary antibody (anti-VSG
antiserum) was diluted 1:50 in ice-cold vPBS. The secondary antibody was diluted to required
concentration (1:3000) in 3 % BSA in PBS. Both were kept on ice in the dark until used. At the
end of the incubation, 50 µl of diluted anti-VSG was added to the cells and quickly mixed by
flicking, followed by an incubation to allow binding (15 min, ice). To fix cells for t=0, 0.5 ml ice-
cold vPBS and 0.5 ml ice-cold 8 % pfa in vPBS were sequentially added at the end of the
incubation and kept on ice. For timepoints t=1 & t=2, 1 ml ice-cold vPBS was added. The cells
were washed by centrifugation (1000 xg, 2 min, 4 °C) and resuspended in 1 ml ice-cold vPBS.
To fix t=1, 350 µl of ice-cold 16% pfa was added and the cells were kept on ice. The cells for
t=2 were incubated (2 min, 37°C) to allow redistribution, flagellar pocket entry, and endocytosis
of the anti-VSG. 350 µl of ice-cold 16 % pfa was then added to fix the cells. All three timepoints
were incubated on ice (20 min) after addition of pfa, followed by a subsequent incubation at
RT (30 min). The fixed cells were pelleted by centrifugation (1000 xg, 2 min, RT), resuspended

in 1 ml RT vPBS, and attached to poly-L-lysine-coated coverslips by centrifugation (1000 xg,
1 min, RT). The attached cells were then permeabilised in 1 ml 0.25 % Tx100 in PBS (5 min,
RT). The permeabilised cells were washed two times in 1 ml PBS and the coverslips with cells
were placed on a drop of clarified and diluted secondary antibody (70 μl) in a humidified
chamber and incubated (1h, RT, dark). The coverslips were washed 3 times in 1 ml PBS (5
min, RT, rocker, dark). After rinsing the cells in ddH$_2$O, they were mounted onto a drop of
DAPI-Fluoromount G (Southern Biotech) on a coverslip that was cleaned with 70% EtOH.
968
969

Acknowledgements
Antibodies were obtained from the following sources: Cynthia He (National University of
Singapore) provided the anti-Ty1 (BB2) and anti-TbLRRP1 antibodies; Chris de Graffenried
(Brown University) provided the anti-TbCentrin2 (2B2H1) and anti-TbCentrin4 (6C5)
antibodies; Markus Engstler provided the anti-VSG antibodies; Keith Gull (University of
Oxford) provided the anti-FAZ1 (L3B2), anti-PFR1,2 (L13D6) and DOT1 antibodies. Nadine
Weisert provided assistance with cloning. Philip Kollmannsberger assisted the correlation
analysis. Markus Engstler provided support and advice. Balázs Szöőr (University of
Edinburgh) and Mick Urbaniak (Lancaster University) provided input on the phosphatase
assays. Mark Field (University of Dundee) provided extremely helpful advice and context on
earlier clathrin RNAi work. Jay Bangs (University at Buffalo) provided helpful advice on the
interpretation of the anti-VSG uptake data. Tim Krüger provided input and assistance with
imaging. Helena Jambor (TU Dresden) advised on data visualisation. Graham Warren (Max
Perutz Laboratories) hosted the preliminary experiments for this project. Susanne Kramer and
Manfred Alsheimer provided feedback on the initial manuscript. Various members of the
trypanosome research community provided feedback during the revision of the manuscript.
986

987 **Competing interests**
988 There are no competing interests to be declared.
989

990 **Funding**
991 Funding in the initial phase was from the Austrian Science Fund (FWF) grant P27016. Later
992 funding was from the German Research Foundation (DFG) grant Mo 3188/2-1. Alyssa Borges
993 was funded by the Brazilian agency CAPES (program: CAPES/DAAD - Call No. 22/2018;
994 process 8881.199683/2018-01)

 Figure legends

Figure 1. TbSmee1 is a cytoskeleton-associated phosphoprotein. (A) Schematic
representation of a trypanosome cell with flagellar pocket (FP), flagellum (F), and flagellum
attachment zone (FAZ) indicated. Inset shows the cytoskeleton-associated structures in the
flagellar pocket neck area: hook complex (HC); flagellar pocket collar (FPC); centrin arm
(Cen4). (B) Schematic representation of TbSmee1 (766 amino acids long) with the three
predicted domains shown in purple, orange, and teal. Approximate amino acid ranges of each
domain are indicated. The regions used as antigens for the generation of the three anti-
TbSmee1 antibodies (303, 304, 508) are shown in light/dark green bars below the schematic.
(C) Validation of the specificity of the three anti-TbSmee1 antibodies. Immunoblots against
whole cell lysates from uninduced TbSmee1 RNAi cells (Tet -), induced TbSmee1 RNAi cells
(Tet +), and Ty1-TbSmee1 endogenous replacement cells (End. repl.) using the three anti-
TbSmee1 antibodies (304, 303, 508) are shown. All three antibodies detected an ~ 85 kDa
protein in the Tet - samples (upper panels, white arrows). The protein was strongly depleted
after induction of TbSmee1 RNAi (Tet +). A second heavier band was detected in the
endogenous replacement cells. This heavier band could also be detected using anti-Ty1 (BB2)
antibodies (lower panels). Exemplary results from multiple (n>3) independent experiments are
shown. (D) Fractionation scheme. Bloodstream form (BSF) cells were extracted with the non-
ionic detergent IGEPAL and separated into cytoplasm and cytoskeleton fractions by
centrifugation. (E) TbSmee1 is associated with the cytoskeleton. Immunoblot of whole-cell
input (I), cytoplasmic supernatant (SN) and cytoskeletal pellet (P) fractions with anti-TbSmee1
antibodies. PFR1,2 (upper panel) and BiP (lower panel) were used as markers for the
cytoskeleton and cytoplasm respectively. Equal fractions were loaded in each lane. Exemplary
results from multiple (n>3) experiments are shown. (F) TbSmee1 is phosphorylated in vivo.
Trypanosome cell lysates were incubated with λ phosphatase for the indicated times (0-20
min), followed by TbSmee1 detection by immunoblotting. The fuzzy appearance of the
TbSmee1 was gradually lost over the timecourse. This was not observed in the presence of
phosphatase inhibitors (+ inhibit), both at 24 ºC and on ice, or when samples were kept on ice
for 20 min. Equal loading of samples was confirmed using total protein stain (magenta).
Exemplary results from multiple (n>3) independent experiments are shown. (G) TbSmee1
localisation through the cell cycle. Bloodstream form trypanosomes were extracted using non-
ionic detergent, fixed, and labelled with anti-TbSmee1 antibodies. DNA was stained using
DAPI. In 1K1N cells, TbSmee1 localised to a bar-shaped structure (arrow) close to the
kinetoplast. In 2K1N cells the structure had replicated (arrows). TbSmee1 was additionally
present on a third structure farther along the cell body (arrowhead). In 2K2N cells the signal

intensity of the third structure was much weaker. Exemplary images from multiple (n>3)
independent experiments are shown.

1036

Figure 2. TbSmee1 is localised to the shank part of the hook complex. Bloodstream form
trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated
antibodies. Either wild-type or Ty1-TbSmee1 cells were used. Insets show an enlarged view
of the hook complex region. (A) TbSmee1 overlaps with the shank part of the hook complex
protein TbMORN1 (arrow). (B) TbSmee1 overlaps with the shank part of the hook complex
protein TbLRRP1 (arrow). (C) TbSmee1 partially overlaps with TbCen4. TbCen4 is present at
the basal and probasal bodies (yellow arrow), centrin arm (arrowhead), and a small additional
projection (white arrow). (D) TbSmee1 does not overlap with the flagellar pocket collar protein
TbBILBO1 (arrow). (E) Summary of measured correlation coefficients for each of the
colabelling experiments; red bars show median values. TbSmee11 showed a moderate
correlation with TbMORN1 (0.41) and TbLRRP1 (0.55) and a weak correlation with TbCen4
(0.37) TbBILBO1 (0.3). Each dot represents a single cell in the 1K1N stage (N = 30). All
fluorescence images are maximum intensity z-projections, and an overlay with a single DIC
section is shown. Overlap was manually confirmed in single z-slices. Results were obtained
from multiple (n>3) independent experiments; exemplary images are shown.

1052

Figure 3. TbSmee1 is associated with the tip of the elongating new FAZ in replicating
cells. (A) Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and
labelled with the indicated antibodies. DNA was stained with DAPI. Maximum intensity
projections are shown, with a single DIC z-slice overlaid. In 1K1N cells, TbSmee1 partially
overlapped (arrow) with the posterior end of the FAZ. In 2K1N and 2K2N cells, the additional
TbSmee1 structure (arrowhead) was present at the tip of the elongating new FAZ. Exemplary
images from multiple (n>3) independent experiments are shown. (B) SIM image of the same
preparations. The TbSmee1 structure (inset) lay just ahead of the elongating FAZ tip. (C) SIM
image of a detergent-extracted cell colabelled for TbSmee1 and DOT1. DOT1 appeared to
envelop and partially overlap with TbSmee1.

1063

Figure 4. TbSmee1 domains 2 and 3 are required for targeting to the hook complex. (A)
Schematics of the 8 TbSmee1 truncation constructs tested, along with details of amino acid
ranges, localisation, and size in kDa. (B) TbSmee1 domains 2 and 3 are required for
localisation to the hook complex. Stably-transfected cells inducibly expressing the indicated
TbSmee1 truncations were used. Ty1-TbSmee1 was detected using anti-Ty1 antibodies. In
the absence of induction (-Tet), no signal overlapping with endogenous TbMORN1 was seen.
After induction of expression (+Tet), the Ty1-TbSmee1 truncations overlapped with the shank

part of TbMORN1 (insets). Images shown are maximum intensity projections of the
fluorescence channels overlaid with a single DIC z-slice. Exemplary images are shown. (C)
Confirmation of Ty1-TbSmee1 truncation expression. Whole-cell lysates from control (-Tet)
cells and cells expressing Ty1-TbSmee1 truncations (+Tet) were immunoblotted using anti-
TbSmee1 (left panel) and anti-Ty1 (right panel) antibodies. Total protein stain (magenta) was
used as a loading control. In the exemplary blot shown, the Ty1-TbSmee1 was detected at ~
44 kDa, as expected. (D) Quantification of immunoblotting data. Anti-TbSmee1 signals in the
immunoblots were normalised relative to total protein staining. TbSmee1 levels in uninduced
control cells (TbSmee1) were expressed relative to the mean value of all control samples for
each clone in each experiment. Ty1-TbSmee1 levels were expressed relative to the levels of
endogenous TbSmee1 for each clone in each experiment. The data shown were obtained
from two independent experiments with each Ty1-TbSmee1 truncation; each experiment used
three separate clones.

1084

Figure 5. TbSmee1 is essential for the viability of bloodstream form *T. brucei*. (A)
Depletion of TbSmee1 causes a strong growth defect. The population density of uninduced
control (Tet-) and TbSmee1-depleted (Tet+) cells were measured at regular intervals after
induction of RNAi over a 72 h timecourse. The insert shows data from experiments with
readings taken at hourly intervals. A growth defect was evident from 21 h post-induction
onwards. (B) Confirmation of TbSmee1 depletion at the single-cell level. Uninduced control
(Tet-) and TbSmee1-depleted (Tet+) RNAi cells were extracted, fixed, and labelled with anti-
TbSmee1 antibodies (green). DNA was labelled with DAPI (magenta). Maximum intensity
projections of the fluorescence channels are shown overlaid with a single DIC z-slice. Insets
show the TbSmee1 signal. Exemplary 1K1N and 2K2N cells from each condition are shown.
(C) Confirmation and quantification of TbSmee1 depletion. Whole-cell lysates from control
(Tet-) and TbSmee1-depleted (Tet+) RNAi cells were immunoblotted using antibodies against
TbSmee1 (green). Total protein stain (TPS, magenta) was used for signal normalisation. An
exemplary immunoblot is shown. Normalised TbSmee1 signals in +Tet cells were expressed
relative to the -Tet signal for each clone in each experiment. TbSmee1 -Tet signals for each
clone in each experiment were expressed relative to the mean of all TbSmee1 -Tet values in
the dataset. TbSmee1 depletion was quantified at the indicated timepoints post-induction. (D)
Depletion of TbSmee1 results in an enlargement of the flagellar pocket. Electron microscopy
images of control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells are shown. The cells were
fixed 24 h post-induction. All data in the experiments shown in this figure were obtained from
multiple (n≥3) independent experiments, each using three separate clones.

1106

Figure 6. Flagellar pocket enlargement is an early consequence of TbSmee1 depletion.
(A) Summary of dextran uptake protocol. The use of temperature blocks inhibits endocytosis, allowing dextran to enter the flagellar pocket but not be internalised. (B) The intensity of the dextran signal reports on flagellar pocket volume. Cells were incubated with fluorophore-conjugated dextran (green) prior to being fixed and imaged using fluorescence microscopy. DNA was labelled with DAPI (magenta). Exemplary cells from control (- Tet) and TbSmee1-depleted (+ Tet) conditions are shown. The + Tet cells (18, 21, or 24 h of induction) exhibited various degrees of flagellar pocket enlargement and progressive morphological aberration. Insets show the dextran signal with a log-scale LUT. (C) Flow cytometry analysis of control (-Tet) and TbSmee1-depleted (+ Tet) cells incubated with fluorescent dextran at various timepoints after induction of RNAi. At later timepoints there is a clear emergence of a subpopulation of cells with much greater fluorescence intensity (arrow). Exemplary traces from a single experiment are shown. (D) Quantification of flow cytometry data. The geometric mean of the fluorescence intensity in control (- Tet) and TbSmee1-depleted (+ Tet) cells was measured at various timepoints after induction of RNAi. Bars indicate median values. The population mean of the + Tet cells was clearly higher at the later timepoints; a visible shift was visible as early as 18 h post-induction (arrow). All data in the experiments shown in this figure were obtained from multiple (n≥2) independent experiments; each experiment used three separate clones.

Figure 7. Surface-bound reporters cannot enter the flagellar pocket of TbSmee1-depleted cells. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In TbSmee1-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and TbSmee1-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. Percentages represent total cell counts (442 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Percentages represent total cell counts (103 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure 8. Blockade of surface reporters is due to an inhibition of endocytosis. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In clathrin-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower-panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and clathrin-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. Percentages represent total cell counts (84 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Percentages represent total cell counts (58 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure 9. Anti-VSG does not enter the flagellar pocket of TbSmee1-depleted cells. (A) Summary of anti-VSG uptake protocol. (B) Anti-VSG is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, anti-VSG signals were observed at the cell surface (t=0), then at the location of the flagellar pocket (t=1) and finally at the endosomal/lysosomal system after allowing endocytosis (t=2) (arrows, upper panels). In TbSmee1-depleted cells (+Tet), no anti-VSG signals were observed in the enlarged flagellar pocket or in the endosomal/lysosomal system (t=1, t=2) (arrows, lower panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of anti-VSG uptake experiments. Cells from TbSmee1-depleted samples (+Tet, t=2) were manually classified into three categories (Normal phenotype, BigEye, and Enlarged pocket). The anti-VSG signal could not be observed in enlarged flagellar pockets in BigEye cells or cells with slightly enlarged flagellar pockets. Signals in these cells were classified into three categories (Dot, Other signal or No signal). Results are shown as a stacked bar chart; exemplary cells for each category are shown. Percentages represent total cell counts (408 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure S1. The TbSmee1 primary structure contains three blocks of highly-conserved
sequence. Multiple sequence alignment of *Trypanosoma brucei* Smee1 (TbSmee1) and
homologous proteins from other trypanosome species, generated using MultAlin (Corpet,
1988) with default parameters. Amino acid numbers are indicated in black numerals above
the alignment. Moderately (50-90%) conserved residues are highlighted in blue. Highly-
conserved (>90%) or completely conserved residues are highlighted in red. The consensus
sequence is shown below the alignment. Abbreviations: TbSmee1, *Trypanosoma brucei*
Smee1; T.cong, *Trypanosoma congolense*; T.vivax, *Trypanosoma vivax*; T.grayi,
Trypanosoma grayi. T.conorhini, *Trypanosoma conorhini*; T.rangelii, *Trypanosoma rangelii*;
T.cruzi, *Trypanosoma cruzi*; T.theileri, *Trypanosoma theileri*. The % sequence identity of each
homologue to TbSmee1 is indicated after the name.

1191

Figure S2. TbSmee1 contains three predicted structured domains. (A) Schematic
representation of TbSmee1 with the three predicted domains (Dom1/2/3) indicated in blue,
magenta, and green. The approximate amino acid coordinates in the primary structure are
indicated in black above each domain. The isoelectric point (pI) of each domain is indicated
above the domain. (B) TbSmee1 contains three regions of predicted secondary structure.
Secondary structure prediction generated using the PSIPRED server (Buchan & Jones, 2019)
with default parameters. Abbreviations and schematics are defined in the legend at the bottom
right. The three predicted structured domains are outlined as blue, red, and green boxes.

1200

Figure S3. Generation of Ty1-TbSmee1 endogenous replacement cell line. (A) Schematic
showing the in situ tagging procedure and annealing sites for PCR primers. The two
endogenous alleles of TbSmee1 in the gDNA are shown in yellow, flanked by 5'UTR and
3'UTR sequences. *T.* brucei cells were transfected with a targeting fragment with homology
arms for recombination with the 3' end of the 5'UTR and 5' end of the ORF. The targeting
fragment contained a blasticidin resistance gene (bla^R), the intergenic region from the
alpha/beta tubulin locus ($\alpha\beta$) and a 3xTy1 epitope tag preceded by an ATG start codon.
Homologous recombination removed the endogenous ATG start codon of the TbSmee1 ORF.
(B) Confirmation of targeting fragment integration at the endogenous TbSmee1 locus by PCR
analysis of genomic DNA. Genomic DNA from wild-type (wt) and candidate Ty1-TbSmee1
clones (c1, c2, c3) was analysed by PCR. Left panel: positive control using 8820_ORF_F and
8820_gDNA_R primers; 490 bp product expected. Right panel: integration test using BB2_F
and 8820_gDNA_R primers. A product is only expected if the 3xTy1 sequence has integrated
upstream of the TbSmee1 ORF (see primer annealing sites in panel A). Two independent
experiments were carried out, each using all three separate clones. (C) Ty1-TbSmee1
displays the same localisations through the cell cycle as endogenous TbSmee1. Detergent-

extracted cells were fixed with methanol and labelled with anti-Ty1 antibodies; DNA was
stained using DAPI. Exemplary cells from the three main cell cycle states (1K1N, 2K1N, 2K2N)
are shown. Maximum-intensity z-projections of the fluorescence channels are shown, together
with a single DIC z-slice overlay. The Ty1-TbSmee1 signal is shown in magenta in the overlay
and highlighted with arrows. Arrowheads indicate the Ty1-TbSmee1 present at the tip of the
new FAZ. Insets show an enlarged view of the TbSmee1 signal from the hook complex or FAZ
tip. Multiple (n>3) independent experiments were carried out using three separate clones. (D)
The anti-Ty1 signal is specific for Ty1-TbSmee1. Wild-type (wt) and Ty1-TbSmee1 cells were
extracted with detergent, fixed with methanol, and labelled with anti-TbSmee1 and anti-Ty1
antibodies. Hook complex (arrows) and FAZ tip (arrowheads) localisations are indicated. No
anti-Ty1 signal was seen in wild-type cells; strong overlap between the anti-Ty1 and anti-
TbSmee1 signals was seen in the Ty1-TbSmee1 cells. Maximum-intensity z-projections of the
fluorescence channels are shown, together with a single DIC z-slice overlay.

1230

Figure S4. Validation of anti-TbStarkey1 antibodies. (A) Validation of anti-TbStarkey1
antibody specificity by immunoblotting. Three separate TbStarkey1 RNAi clones (C8, C9, C10)
were analysed by immunoblotting. Both whole-cell lysates (WCL) and detergent-extracted
cytoskeleton (CK) samples were obtained at the indicated timepoints (24 h, 48 h) from control
(Tet-) and induced (Tet+) cells. The samples were immunoblotted using two separate anti-
TbStarkey1 affinity-purified antibodies (antibody 476, 477). Both antibodies recognised a >130
kDa protein whose abundance was depleted after 48 h of RNAi. The 476 antibody additionally
showed a significant cross-reaction with a protein of <100 kDa in WCL but not CK samples. A
section of the total protein stain of each membrane is shown as an inset (magenta). (B)
Validation of anti-TbStarkey1 antibody specificity by immunofluorescence microscopy. Control
(Tet-) and TbStarkey1-depleted (Tet+) RNAi cells were extracted with non-ionic detergent,
fixed with methanol, and labelled with anti-TbStarkey1 antibodies (green); DNA was stained
using DAPI (magenta). TbStarkey1 localised to the hook complex, and signal was lost upon
depletion. The boxed areas are shown enlarged in the bottom panels. Identical results were
obtained using both anti-TbStarkey1 antibodies; exemplary images using the 477 antibodies
are shown. (C) TbSmee1 overlaps with the shank part of the hook complex protein TbStarkey1
(arrow). All fluorescence images are maximum intensity projections, and an overlay with a
single DIC section is shown. Overlap was manually confirmed in single z-slices. Results were
obtained from multiple (n>3) independent experiments; exemplary images are shown. (D)
Summary of measured correlation coefficients. Each dot represents a single cell; red lines
show median values.

1252

1253

Figure S5. Validation of TbSmee1 truncations. (A) Validation of Ty1-TbSmee1 truncation
cell lines by PCR analysis of genomic DNA and immunoblotting. The results for all 8 TbSmee1
truncations are shown, separated by dotted lines. Upper panels: PCR analysis of genomic
DNA to confirm the presence of the truncation construct. PCR was used to amplify DNA from
clones (c1, c2, c3) and wild-type control (wt) genomic DNA. Primers annealing to the sequence
encoding Ty1 epitope and within the truncation were used. Lower panels: confirmation of Ty1-
TbSmee1 truncation construct expression by immunoblotting. Whole-cell lysates from
uninduced (-Tet) and induced (+Tet) cells were analysed by immunoblotting with anti-
TbSmee1 (green) and anti-Ty1 (cyan) antibodies. Both antibodies detected proteins
corresponding to the predicted size of the Ty1-TbSmee1 truncations; arrows indicate the
target protein in the anti-Ty1 blots. A portion of the total protein staining of the membranes is
shown in magenta. (B) Ty1-TbSmee1(161-766) localises to both the hook complex and the
FAZ tip. Cells expressing the Ty1-TbSmee1(161-766) construct were extracted with non-ionic
detergent, fixed with methanol, and labelled with anti-TbMORN1 and anti-Ty1 antibodies. DNA
was stained with DAPI (cyan). Ty1-TbSmee1(161-766) was observed at both the hook
complex and the FAZ tip (arrow). (C) Ty1-TbSmee1(2-400) localises to the FAZ tip but not the
hook complex. Cells expressing the Ty1-TbSmee1(2-400) construct were extracted, fixed, and
labelled as above. Ty1-TbSmee1(2-400) was observed exclusively at the FAZ tip (arrow).
Images in panels B and C are maximum intensity z-projections with a single DIC z-slice
overlay. Multiple (n>2) independent experiments using three separate clones for each
construct were carried out; exemplary cells are shown.

1275

Figure S6. Additional results for TbSmee1 and TbStarkey1 RNAi experiments. (A)
TbSmee1 depletion causes changes to cell cycle state distribution. Control (-Tet) and
TbSmee1-depleted (+Tet) RNAi cells from 18 h, 21 h, 24 h post-induction were fixed using
glutaraldehyde; DNA was stained using DAPI. The various cell cycle states (1K1N, etc) were
manually quantified from images taken of the fixed cells. Data were obtained from three
independent experiments, each using three separate clones; at least 240 cells were quantified
for each timepoint. (B) Depletion of TbStarkey1 has no effect on population cell growth. Control
(-Tet) and TbStarkey1-depleted (+Tet) cells were followed over a 72 h timecourse, and
population density (cells/ml) was measured every 24 h. Data were obtained from three
independent experiments, each using 3 separate clones. (C, D) Depletion of TbStarkey1
causes morphological abnormality. TbStarkey1-depleted cells were prepared for electron
microscopy using high-pressure freezing and imaged. Cells with enlarged flagellar pockets
could readily be observed.

1289

Figure S7. Depletion of TbSmee1 affects other cytoskeleton-associated proteins.
TbSmee1 RNAi cells were induced for 24 h and analysed by immunoblotting of whole-cell
lysates and immunofluorescence imaging of detergent-extracted cells. (A) Exemplary
immunoblots showing the effect of 24 h TbSmee1 depletion on marker proteins for the hook
complex, flagellar pocket collar, and centrin arm. For comparison, the effect of 16 h TbMORN1
depletion was also analysed. (B) Quantification of immunoblot data. TbSmee1 depletion did
not affect any of the proteins analysed. Total protein staining was used for signal
normalisation. Normalised TbSmee1 signals in +Tet cells were expressed relative to the -Tet
signal for each sample. TbSmee1 -Tet signals were expressed relative to the mean of all
TbSmee1 -Tet values in the dataset. Immunoblots from TbMORN1 RNAi experiments were
quantified in the same way; TbMORN1 depletion resulted in a loss of both TbSmee1 and
TbStarkey1 signal. The data shown were obtained from multiple (n>2) independent
experiments, each using three (TbSmee1 RNAi) or two (TbMORN1 RNAi) separate clones.
(C-F) TbSmee1 depletion does not affect the localisation of the marker proteins. Control (-Tet)
and TbSmee1-depleted (+Tet) cells were extracted with detergent, fixed with methanol, and
labelled with the indicated antibodies; DNA was stained using DAPI. The position of the hook
complexes in exemplary 1K1N and 2K2N cells are shown with arrows; panel E also indicates
basal bodies. Data were obtained from 2 independent experiments for each labelling
combination, each using 3 separate clones. (G) TbSmee1 and TbLRRP1 in TbMORN1-
depleted cells. (H) Summary of the observed effects (arrow = depletion, squiggles = no
change) on the abundance of marker proteins caused by either TbMORN1 or TbSmee1
depletion, based on immunoblotting and immunofluorescence data.

1312

1313

Figure S8. Depletion of TbSmee1 prevents trafficking of ConA to the lysosome. Control
(-Tet) and TbSmee1-depleted cells (+Tet; 24 h timepoint) were incubated on ice with ConA
(magenta). The cells were then shifted to 37 ºC for 30 min to allow internalisation. The cells
were then fixed and labelled with antibodies specific for the lysosome marker p67 (green).
Control (-Tet) cells showed strong overlap between the two labels, indicating that ConA had
been trafficked to the lysosome. +Tet cells showed no overlap between the two labels.
Maximum intensity projections of the fluorescence channels are shown overlaid with a single
DIC z-slice. Overlap in -Tet cells was confirmed in single z-slices. Single channels from the
boxed area in each image are shown as insets. Data obtained from multiple (n>2) independent
experiments each using three separate clones; exemplary cells are shown.

1324

1325

1326

References

Ahmad, E., A. Naeem, S. Javed, S. Yadav, and R.H. Khan. 2007. The minimal structural requirement of concanavalin A that retains its functional aspects. *J Biochem*. 142:307-315.

Aigouy, B., and V. Mirouse. 2013. ScientiFig: a tool to build publication-ready scientific figures. *Nat Methods*. 10:1048.

Albisetti, A., C. Florimond, N. Landrein, K. Vidilaseris, M. Eggenspieler, J. Lesigang, G. Dong, D.R. Robinson, and M. Bonhivers. 2017. Interaction between the flagellar pocket collar and the hook complex via a novel microtubule-binding protein in Trypanosoma brucei. *PLoS Pathog*. 13:e1006710.

Ali, M., K.F. Leung, and M.C. Field. 2014. The ancient small GTPase Rab21 functions in intermediate endocytic steps in trypanosomes. *Eukaryotic cell*. 13:304-319.

Alibu, V.P., L. Storm, S. Haile, C. Clayton, and D. Horn. 2005. A doubly inducible system for RNA interference and rapid RNAi plasmid construction in Trypanosoma brucei. *Mol Biochem Parasitol*. 139:75-82.

Allen, C.L., D. Goulding, and M.C. Field. 2003. Clathrin-mediated endocytosis is essential in Trypanosoma brucei. *Embo J*. 22:4991-5002.

Alsford, S., T. Kawahara, L. Glover, and D. Horn. 2005. Tagging a T. brucei RRNA locus improves stable transfection efficiency and circumvents inducible expression position effects. *Mol Biochem Parasitol*. 144:142-148.

Armstrong, J.K., R.B. Wenby, H.J. Meiselman, and T.C. Fisher. 2004. The hydrodynamic radii of macromolecules and their effect on red blood cell aggregation. *Biophys J*. 87:4259-4270.

Bastin, P., Z. Bagherzadeh, K.R. Matthews, and K. Gull. 1996. A novel epitope tag system to study protein targeting and organelle biogenesis in Trypanosoma brucei. *Mol Biochem Parasitol*. 77:235-239.

Batram, C., N.G. Jones, C.J. Janzen, S.M. Markert, and M. Engstler. 2014. Expression site attenuation mechanistically links antigenic variation and development in Trypanosoma brucei. *Elife*. 3:e02324.

Benz, C., and M.D. Urbaniak. 2019. Organising the cell cycle in the absence of transcriptional
control: Dynamic phosphorylation co-ordinates the Trypanosoma brucei cell cycle post-
transcriptionally. *PLoS Pathog*. 15:e1008129.

Bonhivers, M., S. Nowacki, N. Landrein, and D.R. Robinson. 2008. Biogenesis of the
trypanosome endo-exocytotic organelle is cytoskeleton mediated. *PLoS Biol*. 6:e105.

Borst, P., and A.H. Fairlamb. 1998. Surface receptors and transporters of Trypanosoma
brucei. *Annu Rev Microbiol*. 52:745-778.

Brickman, M.J., and A.E. Balber. 1990. Trypanosoma brucei rhodesiense bloodstream forms:
surface ricin-binding glycoproteins are localized exclusively in the flagellar pocket and the
flagellar adhesion zone. *The Journal of protozoology*. 37:219-224.

Brickman, M.J., J.M. Cook, and A.E. Balber. 1995. Low temperature reversibly inhibits
transport from tubular endosomes to a perinuclear, acidic compartment in African
trypanosomes. *J Cell Sci*. 108 (Pt 11):3611-3621.

Broster Reix, C.E., C. Florimond, A. Cayrel, A. Mailhe, C. Agnero-Rigot, N. Landrein, D.
Dacheux, K. Havlicek, M. Bonhivers, B. Morriswood, and D.R. Robinson. 2021. Bhalin, an
Essential Cytoskeleton-Associated Protein of Trypanosoma brucei Linking TbBILBO1 of the
Flagellar Pocket Collar with the Hook Complex. *Microorganisms*. 9.

Coppens, I., F.R. Opperdoes, P.J. Courtoy, and P. Baudhuin. 1987. Receptor-mediated
endocytosis in the bloodstream form of Trypanosoma brucei. *J Protozool*. 34:465-473.

de Graffenried, C.L., D. Anrather, F. Von Raussendorf, and G. Warren. 2013. Polo-like kinase
phosphorylation of bilobe-resident TbCentrin2 facilitates flagellar inheritance in Trypanosoma
brucei. *Mol Biol Cell*. 24:1947-1963.

Dean, S., J.D. Sunter, and R.J. Wheeler. 2017. TrypTag.org: A Trypanosome Genome-wide
Protein Localisation Resource. *Trends Parasitol*. 33:80-82.

Demmel, L., K. Schmidt, L. Lucast, K. Havlicek, A. Zankel, T. Koestler, V. Reithofer, P. de
Camilli, and G. Warren. 2014. The endocytic activity of the flagellar pocket in Trypanosoma
brucei is regulated by an adjacent phosphatidylinositol phosphate kinase. *J Cell Sci*. 127:2351-
2364.

Engstler, M., T. Pfohl, S. Herminghaus, M. Boshart, G. Wiegertjes, N. Heddergott, and P. Overath. 2007. Hydrodynamic flow-mediated protein sorting on the cell surface of trypanosomes. *Cell*. 131:505-515.

Engstler, M., L. Thilo, F. Weise, C.G. Grunfelder, H. Schwarz, M. Boshart, and P. Overath. 2004. Kinetics of endocytosis and recycling of the GPI-anchored variant surface glycoprotein in Trypanosoma brucei. *J Cell Sci*. 117:1105-1115.

Esson, H.J., B. Morriswood, S. Yavuz, K. Vidilaseris, G. Dong, and G. Warren. 2012. Morphology of the trypanosome bilobe, a novel cytoskeletal structure. *Eukaryotic cell*. 11:761-772.

Florimond, C., A. Sahin, K. Vidilaseris, G. Dong, N. Landrein, D. Dacheux, A. Albisetti, E.H. Byard, M. Bonhivers, and D.R. Robinson. 2015. BILBO1 Is a Scaffold Protein of the Flagellar Pocket Collar in the Pathogen Trypanosoma brucei. *PLoS pathogens*. 11:e1004654.

Gadelha, C., S. Rothery, M. Morphew, J.R. McIntosh, N.J. Severs, and K. Gull. 2009. Membrane domains and flagellar pocket boundaries are influenced by the cytoskeleton in African trypanosomes. *Proc Natl Acad Sci U S A*. 106:17425-17430.

Garrison, P., U. Khan, M. Cipriano, P.J. Bush, J. McDonald, A. Sur, P.J. Myler, T.K. Smith, S.L. Hajduk, and J.D. Bangs. 2021. Turnover of Variant Surface Glycoprotein in Trypanosoma brucei Is a Bimodal Process. *mBio*. 12:e0172521.

Grunfelder, C.G., M. Engstler, F. Weise, H. Schwarz, Y.D. Stierhof, G.W. Morgan, M.C. Field, and P. Overath. 2003. Endocytosis of a glycosylphosphatidylinositol-anchored protein via clathrin-coated vesicles, sorting by default in endosomes, and exocytosis via RAB11-positive carriers. *Mol Biol Cell*. 14:2029-2040.

Hall, B., C.L. Allen, D. Goulding, and M.C. Field. 2004. Both of the Rab5 subfamily small GTPases of Trypanosoma brucei are essential and required for endocytosis. *Mol Biochem Parasitol*. 138:67-77.

Hall, B.S., E. Smith, W. Langer, L.A. Jacobs, D. Goulding, and M.C. Field. 2005. Developmental variation in Rab11-dependent trafficking in Trypanosoma brucei. *Eukaryotic cell*. 4:971-980.

Halliday, C., K. Billington, Z. Wang, R. Madden, S. Dean, J.D. Sunter, and R.J. Wheeler. 2019. Cellular landmarks of Trypanosoma brucei and Leishmania mexicana. *Mol Biochem Parasitol*. 230:24-36.

Halliday, C., A. de Castro-Neto, C.L. Alcantara, E.S.N.L. Cunha, S. Vaughan, and J.D. Sunter.
2021. Trypanosomatid Flagellar Pocket from Structure to Function. *Trends Parasitol*. 37:317-
329.

Hirumi, H., and K. Hirumi. 1989. Continuous cultivation of Trypanosoma brucei blood stream
forms in a medium containing a low concentration of serum protein without feeder cell layers.
J Parasitol. 75:985-989.

Hughes, L., K. Towers, T. Starborg, K. Gull, and S. Vaughan. 2013. A cell-body groove
housing the new flagellum tip suggests an adaptation of cellular morphogenesis for parasitism
in the bloodstream form of Trypanosoma brucei. *J Cell Sci*. 126:5748-5757.

Ikeda, K.N., and C.L. de Graffenried. 2012. Polo-like kinase is necessary for flagellum
inheritance in Trypanosoma brucei. *J Cell Sci*. 125:3173-3184.

Isch, C., P. Majneri, N. Landrein, Y. Pivovarova, J. Lesigang, F. Lauruol, D.R. Robinson, G.
Dong, and M. Bonhivers. 2021. Structural and functional studies of the first tripartite protein
complex at the Trypanosoma brucei flagellar pocket collar. *PLoS Pathog*. 17:e1009329.

Jones, N.G., E.B. Thomas, E. Brown, N.J. Dickens, T.C. Hammarton, and J.C. Mottram. 2014.
Regulators of Trypanosoma brucei cell cycle progression and differentiation identified using a
kinome-wide RNAi screen. *PLoS Pathog*. 10:e1003886.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool,
R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S.A.A. Kohl, A.J. Ballard, A. Cowie,
B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy,
M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O.
Vinyals, A.W. Senior, K. Kavukcuoglu, P. Kohli, and D. Hassabis. 2021. Highly accurate
protein structure prediction with AlphaFold. *Nature*. 596:583-589.

Kelley, R.J., D.L. Alexander, C. Cowan, A.E. Balber, and J.D. Bangs. 1999. Molecular cloning
of p67, a lysosomal membrane glycoprotein from Trypanosoma brucei. *Mol Biochem*
Parasitol. 98:17-28.

Kohl, L., T. Sherwin, and K. Gull. 1999. Assembly of the paraflagellar rod and the flagellum
attachment zone complex during the Trypanosoma brucei cell cycle. *J Eukaryot Microbiol*.
46:105-109.

Lacomble, S., S. Vaughan, C. Gadelha, M.K. Morphew, M.K. Shaw, J.R. McIntosh, and K.
Gull. 2009. Three-dimensional cellular architecture of the flagellar pocket and associated

cytoskeleton in trypanosomes revealed by electron microscope tomography. *J Cell Sci.*
122:1081-1090.

Li, M.Z., and S.J. Elledge. 2012. SLIC: a method for sequence- and ligation-independent
cloning. *Methods Mol Biol.* 852:51-59.

Link, F., A.R. Borges, N.G. Jones, and M. Engstler. 2021. To the Surface and Back: Exo- and
Endocytic Pathways in Trypanosoma brucei. *Front Cell Dev Biol.* 9:720521.

McAllaster, M.R., K.N. Ikeda, A. Lozano-Nunez, D. Anrather, V. Unterwurzacher, T.
Gossenreiter, J.A. Perry, R. Crickley, C.J. Mercadante, S. Vaughan, and C.L. de Graffenried.
2015. Proteomic identification of novel cytoskeletal proteins associated with TbPLK, an
essential regulator of cell morphogenesis in Trypanosoma brucei. *Mol Biol Cell.* 26:3013-3029.

Morriswood, B., K. Havlicek, L. Demmel, S. Yavuz, M. Sealey-Cardona, K. Vidilaseris, D.
Anrather, J. Kostan, K. Djinovic-Carugo, K.J. Roux, and G. Warren. 2013. Novel Bilobe
Components in Trypanosoma brucei Identified Using Proximity-Dependent Biotinylation.
Eukaryotic cell. 12:356-367.

Morriswood, B., C.Y. He, M. Sealey-Cardona, J. Yelinek, M. Pypaert, and G. Warren. 2009.
The bilobe structure of Trypanosoma brucei contains a MORN-repeat protein. *Mol Biochem*
Parasitol. 167:95-103.

Morriswood, B., and K. Schmidt. 2015. A MORN Repeat Protein Facilitates Protein Entry into
the Flagellar Pocket of Trypanosoma brucei. *Eukaryot Cell.* 14:1081-1093.

Nett, I.R., D.M. Martin, D. Miranda-Saavedra, D. Lamont, J.D. Barber, A. Mehlert, and M.A.
Ferguson. 2009. The phosphoproteome of bloodstream form Trypanosoma brucei, causative
agent of African sleeping sickness. *Mol Cell Proteomics.* 8:1527-1538.

Overath, P., and M. Engstler. 2004. Endocytosis, membrane recycling and sorting of GPI-
anchored proteins: Trypanosoma brucei as a model system. *Mol Microbiol.* 53:735-744.

Perry, J.A., A.N. Sinclair-Davis, M.R. McAllaster, and C.L. de Graffenried. 2018. TbSmee1
regulates hook complex morphology and the rate of flagellar pocket uptake in Trypanosoma
brucei. *Mol Microbiol.* 107:344-362.

Pham, K.T.M., H. Hu, and Z. Li. 2020. Maintenance of hook complex integrity and centrin arm
assembly facilitates flagellum inheritance in Trypanosoma brucei. *J Biol Chem.* 295:12962-
12974.

Pham, K.T.M., Q. Zhou, Y. Kurasawa, and Z. Li. 2019. BOH1 cooperates with Polo-like kinase
to regulate flagellum inheritance and cytokinesis initiation in Trypanosoma brucei. *J Cell Sci*.
132.

Postma, M., and J. Goedhart. 2019. PlotsOfData-A web app for visualizing data together with
their summaries. *PLoS Biol*. 17:e3000202.

Price, H.P., M. Stark, and D.F. Smith. 2007. Trypanosoma brucei ARF1 plays a central role in
endocytosis and golgi-lysosome trafficking. *Mol Biol Cell*. 18:864-873.

R Core Team. 2022. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. Available at: https://www.r-project.org/index.html
(Accessed: 24 July 2023).

Redmond, S., J. Vadivelu, and M.C. Field. 2003. RNAit: an automated web-based tool for the
selection of RNAi targets in Trypanosoma brucei. *Mol Biochem Parasitol*. 128:115-118.

Revelle, W. 2023. psych: Procedures for Psychological, Psychometric, and Personality
Research. Available at: https://cran.r-project.org/web/packages/psych/index.html (Accessed:
24 July 2023).
1491
RStudio Team. 2020. RStudio: Integrated Development Environment for R. RStudio, PBC.,
Boston, USA. Available at: https://www.r-project.org/conferences/useR-
2011/abstracts/180111-allairejj.pdf (Accessed: 24 July 2023).

Schauberger, P., and A. Walker. 2022. openxlsx: Read, Write and Edit xlsx Files. Available
at: https://cran.r-project.org/web/packages/openxlsx/openxlsx.pdf (Accessed: 24 July 2023).

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,
C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis.
Nat Methods. 9:676-682.

Selvapandiyan, A., P. Kumar, J.C. Morris, J.L. Salisbury, C.C. Wang, and H.L. Nakhasi. 2007.
Centrin1 is required for organelle segregation and cytokinesis in Trypanosoma brucei. *Mol
Biol Cell*. 18:3290-3301.

Shi, J., J.B. Franklin, J.T. Yelinek, I. Ebersberger, G. Warren, and C.Y. He. 2008. Centrin4
coordinates cell and nuclear division in T. brucei. *J Cell Sci*. 121:3062-3070.

Smithson, L., P. Ihuoma Akazue, L. Findlater, T.M. Gwira, S. Vaughan, and J.D. Sunter. 2022.
Diversity in new flagellum tip attachment in bloodstream form African trypanosomes. *Mol*
Microbiol. 118:510-525.

Sunter, J.D., C. Benz, J. Andre, S. Whipple, P.G. McKean, K. Gull, M.L. Ginger, and J. Lukes.
2015a. Modulation of flagellum attachment zone protein FLAM3 and regulation of the cell
shape in Trypanosoma brucei life cycle transitions. *J Cell Sci*. 128:3117-3130.

Sunter, J.D., and K. Gull. 2016. The Flagellum Attachment Zone: 'The Cellular Ruler' of
Trypanosome Morphology. *Trends Parasitol*. 32:309-324.

Sunter, J.D., V. Varga, S. Dean, and K. Gull. 2015b. A dynamic coordination of flagellum and
cytoplasmic cytoskeleton assembly specifies cell morphogenesis in trypanosomes. *J Cell Sci*.
128:1580-1594.

Urbaniak, M.D., D.M. Martin, and M.A. Ferguson. 2013. Global quantitative SILAC
phosphoproteomics reveals differential phosphorylation is widespread between the procyclic
and bloodstream form lifecycle stages of Trypanosoma brucei. *J Proteome Res*. 12:2233-
2244.

Watson, J.F., and J. Garcia-Nafria. 2019. In vivo DNA assembly using common laboratory
bacteria: A re-emerging tool to simplify molecular cloning. *J Biol Chem*. 294:15271-15281.

Webster, P., D.C. Russo, and S.J. Black. 1990. The interaction of Trypanosoma brucei with
antibodies to variant surface glycoproteins. *J Cell Sci*. 96 (Pt 2):249-255.

Wheeler, R.J. 2021. A resource for improved predictions of Trypanosoma and Leishmania
protein three-dimensional structure. *PLoS One*. 16:e0259871.

Wickham, H. 2016. ggplot2 : Elegant Graphics for Data Analysis. *In* Use R!,. Springer
International Publishing : Imprint: Springer,, Cham. 1 online resource (XVI, 260 pages 232
illustrations, 140 illustrations in color.

Wirtz, E., S. Leal, C. Ochatt, and G.A. Cross. 1999. A tightly regulated inducible expression
system for conditional gene knock-outs and dominant-negative genetics in Trypanosoma
brucei. *Mol Biochem Parasitol*. 99:89-101.

Woods, A., A.J. Baines, and K. Gull. 1989. Evidence for a Mr 88,000 glycoprotein with a
transmembrane association to a unique flagellum attachment region in Trypanosoma brucei.
J Cell Sci. 93 (Pt 3):501-508.

Zhang, C., and F.P. Cordelières. 2016. 3D Quantitative colocalization analysis. *In* Bioimage

Data Analysis K. Miura, editor. Wiley-VCH. 237-266

.

Zhou, Q., L. Gheiratmand, Y. Chen, T.K. Lim, J. Zhang, S. Li, N. Xia, B. Liu, Q. Lin, and C.Y.

He. 2010. A comparative proteomic analysis reveals a new bi-lobe protein required for bi-lobe

duplication and cell division in Trypanosoma brucei. *PLoS One*. 5:e9660.

Zhou, Q., H. Hu, C.Y. He, and Z. Li. 2015. Assembly and maintenance of the flagellum

attachment zone filament in Trypanosoma brucei. *J Cell Sci*. 128:2361-2372.

Zoltner, M., K.F. Leung, S. Alsford, D. Horn, and M.C. Field. 2015. Modulation of the Surface

Proteome through Multiple Ubiquitylation Pathways in African Trypanosomes. *PLoS Pathog*.

11:e1005236.

1547

Figure 1. TbSmee1 is a cytoskeleton-associated phosphoprotein. (A) Schematic representation of a trypanosome cell with flagellar pocket (FP), flagellum (F), and flagellum attachment zone (FAZ) indicated. Inset shows the cytoskeleton-associated structures in the flagellar pocket neck area: hook complex (HC); flagellar pocket collar (FPC); centrin arm (Cen4). (B) Schematic representation of TbSmee1 (766 amino acids long) with the three predicted domains shown in purple, orange, and teal. Approximate amino acid ranges of each domain are indicated. The regions used as antigens for the generation of the three anti-TbSmee1 antibodies (303, 304, 508) are shown in light/dark green bars below the schematic. (C) Validation of the specificity of the three anti-TbSmee1 antibodies. Immunoblots against whole cell lysates from uninduced TbSmee1 RNAi cells (Tet -), induced TbSmee1 RNAi cells (Tet +), and Ty1-TbSmee1 endogenous replacement cells (End. repl.) using the three anti-TbSmee1 antibodies (304, 303, 508) are shown. All three antibodies detected an ~ 85 kDa protein in the Tet - samples (upper panels, white arrows). The protein was strongly depleted after induction of TbSmee1 RNAi (Tet +). A second heavier band was detected in the endogenous replacement cells. This heavier band could also be detected using anti-Ty1 (BB2) antibodies (lower panels). Exemplary results from multiple (n>3) independent experiments are shown. (D) Fractionation scheme. Bloodstream form (BSF) cells were extracted with the non-ionic detergent IGEPAL and separated into cytoplasm and cytoskeleton fractions by centrifugation. (E) TbSmee1 is associated with the cytoskeleton. Immunoblot of whole-cell input (I), cytoplasmic supernatant (SN) and cytoskeletal pellet (P) fractions with anti-TbSmee1 antibodies. PFR1,2 (upper panel) and BiP (lower panel) were used as markers for the cytoskeleton and cytoplasm respectively. Equal fractions were loaded in each lane. Exemplary results from multiple (n>3) experiments are shown. (F) TbSmee1 is phosphorylated in vivo. Trypanosome cell lysates were incubated with λ phosphatase for the indicated times (0-20 min), followed by TbSmee1 detection by immunoblotting. The fuzzy appearance of the TbSmee1 was gradually lost over the timecourse. This was not observed in the presence of phosphatase inhibitors (+ inhibit), both at 24 °C and on ice, or when samples were kept on ice for 20 min. Equal loading of samples was confirmed using total protein stain (magenta). Exemplary results from multiple (n>3) independent experiments are shown. (G) TbSmee1 localisation through the cell cycle. Bloodstream form trypanosomes were extracted using non-ionic detergent, fixed, and labelled with anti-TbSmee1 antibodies. DNA was stained using DAPI. In 1K1N cells, TbSmee1 localised to a bar-shaped structure (arrow) close to the kinetoplast. In 2K1N cells the structure had replicated (arrows). TbSmee1 was additionally present on a third structure farther along the cell body (arrowhead). In 2K2N cells the signal intensity of the third structure was much weaker. Exemplary images from multiple (n>3) independent experiments are shown.

Figure 2. TbSmee1 is localised to the shank part of the hook complex. Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated antibodies. Either wild-type or Ty1-TbSmee1 cells were used. Insets show an enlarged view of the hook complex region. (A) TbSmee1 overlaps with the shank part of the hook complex protein TbMORN1 (arrow). (B) TbSmee1 overlaps with the shank part of the hook complex protein TbLRRP1 (arrow). (C) TbSmee1 partially overlaps with TbCen4. TbCen4 is present at the basal and probasal bodies (yellow arrow), centrin arm (arrowhead), and a small additional projection (white arrow). (D) TbSmee1 does not overlap with the flagellar pocket collar protein TbBILBO1 (arrow). (E) Summary of measured correlation coefficients for each of the colabelling experiments; red bars show median values. TbSmee11 showed a moderate correlation with TbMORN1 (0.41) and TbLRRP1 (0.55) and a weak correlation with TbCen4 (0.37) TbBILBO1 (0.3). Each dot represents a single cell in the 1K1N stage (N = 30). All fluorescence images are maximum intensity z-projections, and an overlay with a single DIC section is shown. Overlap was manually confirmed in single z-slices. Results were obtained from multiple (n>3) independent experiments; exemplary images are shown.

Figure 3. TbSmee1 is associated with the tip of the elongating new FAZ in replicating cells. (A) Bloodstream form trypanosomes were extracted with non-ionic detergent, fixed, and labelled with the indicated antibodies. DNA was stained with DAPI. Maximum intensity projections are shown, with a single DIC z-slice overlaid. In 1K1N cells, TbSmee1 partially overlapped (arrow) with the posterior end of the FAZ. In 2K1N and 2K2N cells, the additional TbSmee1 structure (arrowhead) was present at the tip of the elongating new FAZ. Exemplary images from multiple (n>3) independent experiments are shown. (B) SIM image of the same preparations. The TbSmee1 structure (inset) lay just ahead of the elongating FAZ tip. (C) SIM image of a detergent-extracted cell colabelled for TbSmee1 and DOT1. DOT1 appeared to envelop and partially overlap with TbSmee1.

Figure 4. TbSmee1 domains 2 and 3 are required for targeting to the hook complex. (A) Schematics of the 8 TbSmee1 truncation constructs tested, along with details of amino acid ranges, localisation, and size in kDa. (B) TbSmee1 domains 2 and 3 are required for localisation to the hook complex. Stably-transfected cells inducibly expressing the indicated TbSmee1 truncations were used. Ty1-TbSmee1 was detected using anti-Ty1 antibodies. In the absence of induction (-Tet), no signal overlapping with endogenous TbMORN1 was seen. After induction of expression (+Tet), the Ty1-TbSmee1 truncations overlapped with the shank part of TbMORN1 (insets). Images shown are maximum intensity projections of the fluorescence channels overlaid with a single DIC z-slice. Exemplary images are shown. (C) Confirmation of Ty1-TbSmee1 truncation expression. Whole-cell lysates from control (-Tet) cells and cells expressing Ty1-TbSmee1 truncations (+Tet) were immunoblotted using anti-TbSmee1 (left panel) and anti-Ty1 (right panel) antibodies. Total protein stain (magenta) was used as a loading control. In the exemplary blot shown, the Ty1-TbSmee1 was detected at ~ 44 kDa, as expected. (D) Quantification of immunoblotting data. Anti-TbSmee1 signals in the immunoblots were normalised relative to total protein staining. TbSmee1 levels in uninduced control cells (TbSmee1) were expressed relative to the mean value of all control samples for each clone in each experiment. Ty1-TbSmee1 levels were expressed relative to the levels of endogenous TbSmee1 for each clone in each experiment. The data shown were obtained from two independent experiments with each Ty1-TbSmee1 truncation; each experiment used three separate clones.

Figure 5. TbSmee1 is essential for the viability of bloodstream form T. brucei. (A) Depletion of TbSmee1 causes a strong growth defect. The population density of uninduced control (Tet-) and TbSmee1-depleted (Tet+) cells were measured at regular intervals after induction of RNAi over a 72 h timecourse. The insert shows data from experiments with readings taken at hourly intervals. A growth defect was evident from 21 h post-induction onwards. (B) Confirmation of TbSmee1 depletion at the single-cell level. Uninduced control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were extracted, fixed, and labelled with anti-TbSmee1 antibodies (green). DNA was labelled with DAPI (magenta). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. Insets show the TbSmee1 signal. Exemplary 1K1N and 2K2N cells from each condition are shown. (C) Confirmation and quantification of TbSmee1 depletion. Whole-cell lysates from control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells were immunoblotted using antibodies against TbSmee1 (green). Total protein stain (TPS, magenta) was used for signal normalisation. An exemplary immunoblot is shown. Normalised TbSmee1 signals in +Tet cells were expressed relative to the -Tet signal for each clone in each experiment. TbSmee1 -Tet signals for each clone in each experiment were expressed relative to the mean of all TbSmee1 -Tet values in the dataset. TbSmee1 depletion was quantified at the indicated timepoints post-induction. (D) Depletion of TbSmee1 results in an enlargement of the flagellar pocket. Electron microscopy images of control (Tet-) and TbSmee1-depleted (Tet+) RNAi cells are shown. The cells were fixed 24 h post-induction. All data in the experiments shown in this figure were obtained from multiple (n≥3) independent experiments, each using three separate clones.

Figure 6. Flagellar pocket enlargement is an early consequence of TbSmee1 depletion. (A) Summary of dextran uptake protocol. The use of temperature blocks inhibits endocytosis, allowing dextran to enter the flagellar pocket but not be internalised. (B) The intensity of the dextran signal reports on flagellar pocket volume. Cells were incubated with fluorophore-conjugated dextran (green) prior to being fixed and imaged using fluorescence microscopy. DNA was labelled with DAPI (magenta). Exemplary cells from control (- Tet) and TbSmee1-depleted (+ Tet) conditions are shown. The + Tet cells (18, 21, or 24 h of induction) exhibited various degrees of flagellar pocket enlargement and progressive morphological aberration. Insets show the dextran signal with a log-scale LUT. (C) Flow cytometry analysis of control (- Tet) and TbSmee1-depleted (+ Tet) cells incubated with fluorescent dextran at various timepoints after induction of RNAi. At later timepoints there is a clear emergence of a subpopulation of cells with much greater fluorescence intensity (arrow). Exemplary traces from a single experiment are shown. (D) Quantification of flow cytometry data. The geometric mean of the fluorescence intensity in control (- Tet) and TbSmee1-depleted (+ Tet) cells was measured at various timepoints after induction of RNAi. Bars indicate median values. The population mean of the + Tet cells was clearly higher at the later timepoints; a visible shift was visible as early as 18 h post-induction (arrow). All data in the experiments shown in this figure were obtained from multiple (n≥ 2) independent experiments; each experiment used three separate clones.

Figure 7. Surface-bound reporters cannot enter the flagellar pocket of TbSmee1-depleted cells. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In TbSmee1-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and TbSmee1-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. Percentages represent total cell counts (442 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Percentages represent total cell counts (103 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure 8. Blockade of surface reporters is due to an inhibition of endocytosis. (A) Summary of paired dextran+ConA uptake protocol. (B) ConA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. In control (-Tet) cells, the dextran and ConA signals strongly overlapped (arrow, upper panels). In clathrin-depleted cells (+Tet), little to no overlap between the two reporters was observed (arrow, lower-panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of dextran+ConA uptake experiments. Cells from control (-Tet) and clathrin-depleted (+Tet) samples were manually classified into four categories (1-4) based on the degree of overlap between the ConA and dextran reporters. Results are shown as a slope chart (left) and stacked bar chart (right); exemplary cells for each category are shown. Percentages represent total cell counts (84 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones. (D) Summary of paired dextran+BSA uptake protocol. (E) BSA is unable to enter the enlarged flagellar pocket of clathrin-depleted cells. (F) Quantification of dextran+BSA uptake experiments. Percentages represent total cell counts (58 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure 9. Anti-VSG does not enter the flagellar pocket of TbSmee1-depleted cells. (A) Summary of anti-VSG uptake protocol. (B) Anti-VSG is unable to enter the enlarged flagellar pocket of TbSmee1-depleted cells. In control (-Tet) cells, anti-VSG signals were observed at the cell surface (t=0), then at the location of the flagellar pocket (t=1) and finally at the endosomal/lysosomal system after allowing endocytosis (t=2) (arrows, upper panels). In TbSmee1-depleted cells (+Tet), no anti-VSG signals were observed in the enlarged flagellar pocket or in the endosomal/lysosomal system (t=1, t=2) (arrows, lower panels). Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. (C) Quantification of anti-VSG uptake experiments. Cells from TbSmee1-depleted samples (+Tet, t=2) were manually classified into three categories (Normal phenotype, BigEye, and Enlarged pocket). The anti-VSG signal could not be observed in enlarged flagellar pockets in BigEye cells or cells with slightly enlarged flagellar pockets. Signals in these cells were classified into three categories (Dot, Other signal or No signal). Results are shown as a stacked bar chart; exemplary cells for each category are shown. Percentages represent total cell counts (408 cells) that were obtained from multiple independent experiments (n>3); each experiment included three separate clones.

Figure S1. The TbSmee1 primary structure contains three blocks of highly-conserved sequence. Multiple sequence alignment of Trypanosoma brucei Smee1 (TbSmee1) and homologous proteins from other trypanosome species, generated using MultAlin (Corpet, 1988) with default parameters. Amino acid numbers are indicated in black numerals above the alignment. Moderately (50-90%) conserved residues are highlighted in blue. Highly-conserved (>90%) or completely conserved residues are highlighted in red. The consensus sequence is shown below the alignment. Abbreviations: TbSmee1, Trypanosoma brucei Smee1; T.cong, Trypanosoma congolense; T.vivax, Trypanosoma vivax; T.grayi, Trypanosoma grayi. T.conorhini, Trypanosoma conorhini; T.rangelii, Trypanosoma rangelii; T.cruzi, Trypanosoma cruzi; T.theileri, Trypanosoma theileri. The % sequence identity of each homologue to TbSmee1 is indicated after the name.

Figure S2. TbSmee1 contains three predicted structured domains. (A) Schematic representation of TbSmee1 with the three predicted domains (Dom1/2/3) indicated in blue, magenta, and green. The approximate amino acid coordinates in the primary structure are indicated in black above each domain. The isoelectric point (pI) of each domain is indicated above the domain. (B) TbSmee1 contains three regions of predicted secondary structure. Secondary structure prediction generated using the PSIPRED server (Buchan & Jones, 2019) with default parameters. Abbreviations and schematics are defined in the legend at the bottom right. The three predicted structured domains are outlined as blue, red, and green boxes.

Figure S3. Generation of Ty1-TbSmee1 endogenous replacement cell line. (A) Schematic showing the in situ tagging procedure and annealing sites for PCR primers. The two endogenous alleles of TbSmee1 in the gDNA are shown in yellow, flanked by 5'UTR and 3'UTR sequences. T. brucei cells were transfected with a targeting fragment with homology arms for recombination with the 3' end of the 5'UTR and 5' end of the ORF. The targeting fragment contained a blasticidin resistance gene (blaR), the intergenic region from the alpha/beta tubulin locus (αβ) and a 3xTy1 epitope tag preceded by an ATG start codon. Homologous recombination removed the endogenous ATG start codon of the TbSmee1 ORF. (B) Confirmation of targeting fragment integration at the endogenous TbSmee1 locus by PCR analysis of genomic DNA. Genomic DNA from wild-type (wt) and candidate Ty1-TbSmee1 clones (c1, c2, c3) was analysed by PCR. Left panel: positive control using 8820_ORF_F and 8820_gDNA_R primers; 490 bp product expected. Right panel: integration test using BB2_F and 8820_gDNA_R primers. A product is only expected if the 3xTy1 sequence has integrated upstream of the TbSmee1 ORF (see primer annealing sites in panel A). Two independent experiments were carried out, each using all three separate clones. (C) Ty1-TbSmee1 displays the same localisations through the cell cycle as endogenous TbSmee1. Detergent-extracted cells were fixed with methanol and labelled with anti-Ty1 antibodies; DNA was stained using DAPI. Exemplary cells from the three main cell cycle states (1K1N, 2K1N, 2K2N) are shown. Maximum-intensity z-projections of the fluorescence channels are shown, together with a single DIC z-slice overlay. The Ty1-TbSmee1 signal is shown in magenta in the overlay and highlighted with arrows. Arrowheads indicate the Ty1-TbSmee1 present at the tip of the new FAZ. Insets show an enlarged view of the TbSmee1 signal from the hook complex or FAZ tip. Multiple (n>3) independent experiments were carried out using three separate clones. (D) The anti-Ty1 signal is specific for Ty1-TbSmee1. Wild-type (wt) and Ty1-TbSmee1 cells were extracted with detergent, fixed with methanol, and labelled with anti-TbSmee1 and anti-Ty1 antibodies. Hook complex (arrows) and FAZ tip (arrowheads) localisations are indicated. No anti-Ty1 signal was seen in wild-type cells; strong overlap between the anti-Ty1 and anti-TbSmee1 signals was seen in the Ty1-TbSmee1 cells. Maximum-intensity z-projections of the fluorescence channels are shown, together with a single DIC z-slice overlay.

Figure S4. Validation of anti-TbStarkey1 antibodies. (A) Validation of anti-TbStarkey1 antibody specificity by immunoblotting. Three separate TbStarkey1 RNAi clones (C8, C9, C10) were analysed by immunoblotting. Both whole-cell lysates (WCL) and detergent-extracted cytoskeleton (CK) samples were obtained at the indicated timepoints (24 h, 48 h) from control (Tet-) and induced (Tet+) cells. The samples were immunoblotted using two separate anti-TbStarkey1 affinity-purified antibodies (antibody 476, 477). Both antibodies recognised a >130 kDa protein whose abundance was depleted after 48 h of RNAi. The 476 antibody additionally showed a significant cross-reaction with a protein of <100 kDa in WCL but not CK samples. A section of the total protein stain of each membrane is shown as an inset (magenta). (B) Validation of anti-TbStarkey1 antibody specificity by immunofluorescence microscopy. Control (Tet-) and TbStarkey1-depleted (Tet+) RNAi cells were extracted with non-ionic detergent, fixed with methanol, and labelled with anti-TbStarkey1 antibodies (green); DNA was stained using DAPI (magenta). TbStarkey1 localised to the hook complex, and signal was lost upon depletion. The boxed areas are shown enlarged in the bottom panels. Identical results were obtained using both anti-TbStarkey1 antibodies; exemplary images using the 477 antibodies are shown. (C) TbSmee1 overlaps with the shank part of the hook complex protein TbStarkey1 (arrow). All fluorescence images are maximum intensity projections, and an overlay with a single DIC section is shown. Overlap was manually confirmed in single z-slices. Results were obtained from multiple (n>3) independent experiments; exemplary images are shown. (D) Summary of measured correlation coefficients. Each dot represents a single cell; red lines show median values.

Figure S5. Validation of TbSmee1 truncations. (A) Validation of Ty1-TbSmee1 truncation cell lines by PCR analysis of genomic DNA and immunoblotting. The results for all 8 TbSmee1 truncations are shown, separated by dotted lines. Upper panels: PCR analysis of genomic DNA to confirm the presence of the truncation construct. PCR was used to amplify DNA from clones (c1, c2, c3) and wild-type control (wt) genomic DNA. Primers annealing to the sequence encoding Ty1 epitope and within the truncation were used. Lower panels: confirmation of Ty1-TbSmee1 truncation construct expression by immunoblotting. Whole-cell lysates from uninduced (-Tet) and induced (+Tet) cells were analysed by immunoblotting with anti-TbSmee1 (green) and anti-Ty1 (cyan) antibodies. Both antibodies detected proteins corresponding to the predicted size of the Ty1-TbSmee1 truncations; arrows indicate the target protein in the anti-Ty1 blots. A portion of the total protein staining of the membranes is shown in magenta. Note that the immunoblot for Ty1-TbSmee1(161-400) is the same data as shown in Figure 3C. (B) Ty1-TbSmee1(161-766) localises to both the hook complex and the FAZ tip. Cells expressing the Ty1-TbSmee1(161-766) construct were extracted with non-ionic detergent, fixed with methanol, and labelled with anti-TbMORN1 and anti-Ty1 antibodies. DNA was stained with DAPI (cyan). Ty1-TbSmee1(161-766) was observed at both the hook complex and the FAZ tip (arrow). (C) Ty1-TbSmee1(2-400) localises to the FAZ tip but not the hook complex. Cells expressing the Ty1-TbSmee1(2-400) construct were extracted, fixed, and labelled as above. Ty1-TbSmee1(2-400) was observed exclusively at the FAZ tip (arrow). Images in panels B and C are maximum intensity z-projections with a single DIC z-slice overlay. Multiple (n>2) independent experiments using three separate clones for each construct were carried out; exemplary cells are shown.

Figure S6. Additional results for TbSmee1 and TbStarkey1 RNAi experiments. (A) TbSmee1 depletion causes changes to cell cycle state distribution. Control (-Tet) and TbSmee1-depleted (+Tet) RNAi cells from 18 h, 21 h, 24 h post-induction were fixed using glutaraldehyde; DNA was stained using DAPI. The various cell cycle states (1K1N, etc) were manually quantified from images taken of the fixed cells. Data were obtained from three independent experiments, each using three separate clones; at least 240 cells were quantified for each timepoint. (B) Depletion of TbStarkey1 has no effect on population cell growth. Control (-Tet) and TbStarkey1-depleted (+Tet) cells were followed over a 72 h timecourse, and population density (cells/ml) was measured every 24 h. Data were obtained from three independent experiments, each using 3 separate clones. (C, D) Depletion of TbStarkey1 causes morphological abnormality. TbStarkey1-depleted cells were prepared for electron microscopy using high-pressure freezing and imaged. Cells with enlarged flagellar pockets could readily be observed.

Figure S7. Depletion of TbSmee1 affects other cytoskeleton-associated proteins. TbSmee1 RNAi cells were induced for 24 h and analysed by immunoblotting of whole-cell lysates and immunofluorescence imaging of detergent-extracted cells. (A) Exemplary immunoblots showing the effect of 24 h TbSmee1 depletion on marker proteins for the hook complex, flagellar pocket collar, and centrin arm. For comparison, the effect of 16 h TbMORN1 depletion was also analysed. (B) Quantification of immunoblot data. TbSmee1 depletion did not affect any of the proteins analysed. Total protein staining was used for signal normalisation. Normalised TbSmee1 signals in +Tet cells were expressed relative to the -Tet signal for each sample. TbSmee1 -Tet signals were expressed relative to the mean of all TbSmee1 -Tet values in the dataset. Immunoblots from TbMORN1 RNAi experiments were quantified in the same way; TbMORN1 depletion resulted in a loss of both TbSmee1 and TbStarkey1 signal. The data shown were obtained from multiple (n>2) independent experiments, each using three (TbSmee1 RNAi) or two (TbMORN1 RNAi) separate clones. (C-F) TbSmee1 depletion does not affect the localisation of the marker proteins. Control (-Tet) and TbSmee1-depleted (+Tet) cells were extracted with detergent, fixed with methanol, and labelled with the indicated antibodies; DNA was stained using DAPI. The position of the hook complexes in exemplary 1K1N and 2K2N cells are shown with arrows; panel E also indicates basal bodies. Data were obtained from 2 independent experiments for each labelling combination, each using 3 separate clones. (G) TbSmee1 and TbLRRP1 in TbMORN1-depleted cells. (H) Summary of the observed effects (arrow = depletion, squiggles = no change) on the abundance of marker proteins caused by either TbMORN1 or TbSmee1 depletion, based on immunoblotting and immunofluorescence data.

Figure S8. Depletion of TbSmee1 prevents trafficking of ConA to the lysosome. Control (-Tet) and TbSmee1-depleted cells (+Tet; 24 h timepoint) were incubated on ice with ConA (magenta). The cells were then shifted to 37 °C for 30 min to allow internalisation. The cells were then fixed and labelled with antibodies specific for the lysosome marker p67 (green). Control (-Tet) cells showed strong overlap between the two labels, indicating that ConA had been trafficked to the lysosome. +Tet cells showed no overlap between the two labels. Maximum intensity projections of the fluorescence channels are shown overlaid with a single DIC z-slice. Overlap in -Tet cells was confirmed in single z-slices. Single channels from the boxed area in each image are shown as insets. Data obtained from multiple (n>2) independent experiments each using three separate clones; exemplary cells are shown.