The use of pigs vocalisation structure to assess the quality of human-pig relationship

Authors:

Avelyne S Villain* [1,2], Carole Guérin [1], Camille Noûs [3], Céline Tallet* [1]

Affiliation:

[1] PEGASE, INRAE, Institut Agro, 35590 Saint Gilles, France.

[2] Behavioural Ecology Group, Section for Ecology & Evolution, Department of Biology, University of Copenhagen, 2100, Copenhagen Ø, Denmark

[3] Cogitamus laboratory, 75000 Paris, France

* Corresponding authors : avelyne.s.villain@protonmail.com, celine.tallet@inrae.fr

Key words:

Positive handling, Acoustic communication, Emotions, Mood, Behaviour, Welfare, Interspecific interactions.

Abstract:

Studying human-animal interactions in domestic species and how they affect the establishment of a positive Human-Animal Relationship (HAR) may help us improve animal welfare and better understand the evolution of interspecific interactions associated with the domestication process. Understanding and describing the quality of an HAR requires information on several aspects of the animal biology and emotional states (social, spatial and postural behaviours, physiological and cognitive states). Growing evidence shows that acoustic features of animal vocalisations may be indicators of emotional states. Here, we tested the hypothesis that the quality of vocal expression may indicate the quality of HAR. At weaning, 30 piglets were positively handled by an experimenter who talked to and physically interacted with them three times a day, while 30 other piglets only received the contact necessary for proper husbandry. After two weeks, we recorded the behaviours and vocalisations produced in the presence of the static experimenter for five minutes. We repeated this test two weeks later, after a conditioning period during which human presence with

additional positive contact was used as a reward for all piglets. We hypothesized this conditioning
period would lead to a positive human-piglet relationship for all piglets. As expected, piglets that
were positively handled at weaning expressed a higher attraction toward the experimenter, and, after
the conditioning, piglets that were not positively handled at weaning expressed a similar level of
attraction than the positively handled ones. Piglets positively handled at weaning generally produced
shorter grunts than the other ones. However the latter expressed more flexibility in call structure
when vocalising close to a human, with a decrease of grunt duration and an increase in pitch,
frequency range and noisiness in their grunt. This differential effect of proximity between groups of
piglets was attenuated after the conditioning during a standard reunion with a static human but
remained over time when the human was providing additional positive contacts. Results suggest that
first, changes in vocal structure are consistent with indicators of positive states in the presence of a
human. Second, increasing familiarity and proximity between a human and a pig may induce
changes in the acoustic structure of its grunts. Third, a human providing additional positive contacts
triggers more changes in vocalisation structure than by their presence only. We show that
vocalisation structure may allow us to assess the quality of human-pig relationship.

Introduction

The process of domestication was conducted to shape physiology and morphology of domestic
animal species, but also their behaviour. It notably has shaped interspecific interactions between
human and non-human animals, by improving animals' capacity to use human signals to adapt their
behaviour both decreasing fearfulness toward humans and increasing attention toward humans
(Mignon-Grasteau et al., 2005). In farms, the relationship that domestic animals form with humans is
important for animal welfare. Therefore, studying human-animal interactions and their consequences
to understand the mechanisms of emergence and maintenance of a positive human-animal
relationship (HAR) directly applies to welfare (Rault et al., 2020). Animal welfare consists of three
major aspects: the ability of an animal to control its mental and physiological stability (Broom,
2011), the decrease of experiencing negatively perceived contexts and the increase in experiencing
positively perceived contexts and species-specific behaviors (Peterson et al., 1995; Weerd & Day,
2009). A positive HAR is thought to be established through repeated positive interactions between
the human and the non-human animal. Some of the mechanisms involved in this process are:

accumulation of positive experiences through positive associative learning, modifications of
cognitive biases, shaping expectations from the non-human animal toward the human. A positive
HAR can be appreciated through behavioural and physiological measures, for example by assessing
the expression of positive emotions [reviewed in (Rault et al. 2020)]. Several behavioural measures
may help to define a positive HAR such as: short latency to approach and spatial proximity (Boivin
et al., 2000; Schmied et al., 2008), body postures (Villain et al., 2020b) or play behaviour
(Jerolmack, 2009). Contacts from a human such as stroking, may induce changes in body postures
and exposition of body areas by the animal to the human, supposedly vulnerable [central neck area in
cattle (Schmied et al. 2008), abdominal area in pigs (Rault et al., 2019)]. Such grooming solicitation
may be markers of engagement, trust and motivation to interact with the human. In most cases, these
behaviours are similar to those shown during intraspecific socio positive interactions, although there
are some species specific behaviours [e.g., dog vs.. wolf (Gácsi et al., 2005)]. Vocal behaviour may
also help defining the quality of an HAR. First, vocalisations are known to carry markers of the
emotional states in several bird and mammal species (Briefer, 2012, 2020). Markers of emotional
valence (positive versus negative) has been studied in domestic farm animals [reviewed in Laurijs et
al. (2021)]. Second, some vocalisations have been associated with positive interactions with humans,
for example the cat – human communication : purring is thought to be derived from mother pup
communication during nursing and is observed associated with care solicitation from humans;
meowing, which is not observed during intra specific interactions is thought to emerge from
associative learning during cat – human interactions (Brown & Bradshaw, 2014). This shows that
HAR may elicit specific vocalisations from the non human animal toward the human.

In pigs, diversified evidence attest the possibility of a positive HAR. Animals may be handled by
humans providing regular additional positive contacts, leading to the expression of a positive
perception of humans, with evidence from behavioural and physiological studies. Cognitive bias tests
showed a positive judgment bias in piglets that had received gentle contacts with humans (Brajon et
al., 2015b). Pigs may recognise a human providing positive contacts compared to an unfamiliar one
and adapt their behaviour accordingly (Brajon et al., 2015c). Pigs may be sensitive to human voice
and respond accordingly (Bensoussan et al., 2019, 2020). Pigs vocalisations are diverse and linked to
their emotional states, attested by the use of positive or negative call types (Briefer et al., 2019, 2022;
Tallet et al., 2013). In addition, even within a call type, spectro-temporal changes are closely related
to the valence of a situation or the arousal a situation may trigger for the animal. For example,
grunts, that are among of the most used vocal signals and various situations is now known to be a
flexible call: shorter grunts have been associated with positive situations (Briefer et al., 2019, 2022;

Friel et al., 2019), as well as higher formants (which are frequency peaks containing more energy

than others) and a lower fundamental frequency during positive situations (Briefer et al., 2019,

2022). Grunt structure may also change according to the arousal of a situation, with a higher

frequency range and a higher bandwidth when the arousal increases (Linhart et al., 2015). In order to

determine if vocalisations may be used as non invasive indicators of the quality of human-pig

relationship by themselves, we tested whether they could encode the quality of the human-pig

relationship, through the vocal expression of emotional state. Because they are used in contexts of

different valence and arousal and in most pigs, we studied the spectro-temporal structure of grunts.

We predicted that if grunts carry information on the quality of the human-pig relationship, then 1. A

period of positive handling given by a human should modulate vocal quality of piglets when in

presence of the human, leading to grunts exhibiting markers of positive states (shorter grunts), 2.

spatial proximity toward the human should influence the spectro-temporal structure of grunts (higher

pitched grunts as the arousal increases).

Methods
109

Ethical note
110

The study was approved by the ethic committee CREEA and received the authorization no.

APAFIS#17071-2018101016045373_V3 from the French Ministry of Higher Education, Research

and Innovation. UE3P, where the experiment was carried out, is an experimental unit authorized by

the French Ministry of Agriculture to breed animals for experimentation under the number D35-275-

32. This authorization includes a derogation to follow the directive 2008/120/EC relative to the

protection of piglets and its regulations.

Subjects and housing conditions
118

Sixty weaned female pigs (in two replicates from January to April 2019), *Sus scrofa domesticus*,

bred from crosses between Large White and Landrace females and Piétrain males were used for this

study from 28 to 62 days after birth. Animal housing and experiments took place at the experimental

unit UE3P (UE 1421, INRAE France).

One piglet had to be excluded from our sample size to receive care/medication due to health issues

independent from the experiment. From weaning at 28 days of age, piglets from the same litter and

having similar weight (<1 kg difference) were housed by three in a 1.2 x 1.3m pen on plastic

duckboard. Wooden panels were used to visually isolate pens. One metal chain per pen was used for

enrichment. Food and water were available *ad libitum*. Artificial lights were turned on from 8:00 to

17:00 and temperature was maintained between 26 and 27 ºC. The experiment was carried out in two

replicates and two identical rearing rooms were used (5 pens per room per replicate).

Treatment: positive handling et weaning

From day 28 (day of weaning) to day 39 of life, piglets were separated into two groups that

experienced a different post-weaning period as follows:

- <u>Non positively handled piglets (H piglets)</u>: Control piglets from 10 rearing pens, housed in the

same room, received the minimal amount of daily contact with a stockperson (a 1.70m tall male who

did the feeding, cleaning and health checkups). The stockperson wore a dark green shirt and pants

and brown shoes.

- <u>Positively handled piglets piglets (H+ piglets)</u>: Experimental piglets from the 10 other rearing pens,

housed in another room, received the same daily care given by the same stockperson as for H piglets.

They additionally received repeated sessions of additional human contacts. Each pen of three piglets

received 29 sessions of 10 minutes, from day 28 (weaning) until day 39, occurring five days a week.

Three sessions per day were performed (except on the day of weaning during which only two were

done with a two-hour break in between). Each session took place in the rearing pen and the order of

the interventions in the pens was balanced across days. The handling procedure, using gentle tactile

contacts is described in supplementary material of Villain et al. (2020) and was similar to Tallet et al.

(2014). Two experimenters performed these sessions (both women, both between 1.70-1.73 m tall,

with a balanced number of pens attributed to each of them). The experimenters wore the same blue

overalls and green boots each time they interacted with the piglets. The experimenters tried to imitate

each others behaviours (remote video monitoring) to decrease variability.

This intense period of additional positive contacts for half of the piglets after weaning constituted the

treatment of positive handling at weaning: positively handled piglets are referred to as H+ piglets and

non positively handled piglets are referred to as H piglets to describe the early experimental

treatment they experienced regarding a human, prior to the conditioning.

Conditioning: sessions of additional positive contacts with (un)familiar human

The conditioning took place between day 42 and 62 of age and lasted twelve days, with two trials per

day and at least three hours between trials on the same day. Piglets were habituated to the test room

for 10 minutes, by pen, two days before the start of the conditioning. All piglets (H and H+) were

subjected to the same conditioning. The experimental design of the conditioning is already published

in an article dedicated to the study of anticipatory behaviour (Villain et al., 2020).

Briefly, all piglets were individually trained to learn to associate two different stimuli with the arrival

of two different (pseudo)-social partners: either two pen mates (partner = Conspecifics) or a familiar

human (partner = Human). When entering the room, the piglets and the partner(s) would remain in

the room for two minutes. Specifically, when the human was the partner, the human entered, sat on a

bucket and positively interacted with the piglet for two minutes, in the same manner as additional

contacts was provided to the H+ piglets during the previous period (see above section) (figure 1).

Therefore, at the beginning of the conditioning phase, H+ piglets were already familiar with the

human, whereas H piglets were unfamiliar with the human and only became familiar during the

conditioning.

The same sessions occurred in both treatment groups (H and H+). It was thus excepted that, at the

end of the conditioning, all piglets would be familiar with the human, but with a different degree in

H+ and H- piglets, due to a different time of exposure (H+: period of positive handling at weaning +

conditioning, H: conditioning only). Sessions of reunions with social partners were not studied and

only served as reward during the conditioning in a previous analysis of vocal expression of positive

anticipation (Villain et al., 2020). Indeed, first the two (pseudo) social contexts would have been

difficult to compare (reunion between three piglets vs. reunion between one piglet and one human).

Second, regarding the vocal behaviour, the caller among the group of three piglets would not have

been identified reliably, making it difficult to study within individual vocal flexibility.

For every second trial, the two-minute reunions with the human were analysed by the same person:

trials number 2, 4, 6, 8, 10 and 11 (see behavioural analyses section).

Standard Isolation/Reunion Tests

At 40 or 41 (before conditioning) and then 63 or 64 (after conditioning) days of age, piglets were

subjected to a standard Isolation/Reunion test in order to assess their perception of the human. The

test consisted of two phases. The piglet was brought individually in a trolley to the experimental

room. It was left alone for five minutes, which defined the 'Isolation' phase. Then, the human

entered the room, remained stand up for 30 seconds and they sat on a bucket, remaining silent and

not moving for 4.5 minutes (figure 1).

↔ 65.5cm

■ Location of the microphone

⬚ Virtual zones for behavioural scoring

🟠 Location of the human

▢ Proximal area   ⬚ Distal area

188   *Figure 1: Design of the room used during the Isolation/Reunion tests and the additional positive contacts sessions of the*
189   *conditioning. The room was split into 16 virtual zones. A proximal area (zones 10, 11, 14, 15) and a distal area (zones 1,*
190   *2, 3, 4) were defined, suing the location of the human as reference.*

Behavioural monitoring and analyses

Sessions and tests were recorded using a camera (Bosh, Box 960H-CDD) and behaviours were

scored *a posteriori* on videos using *The Observer XT 14.0* (Noldus, The Netherlands) software. The

room was split into 16 virtual equally-dimensioned zones to assess the mobility and exploratory

behaviour of the piglet. A proximal area, around the human was defined by merging four zones, a

distal area was defined merging the four most distant zones from the human (figure 1).

The behaviours scored during the reunion of the Isolation/Reunion test and the sessions of additional

positive contacts of the conditioning are available in table 1. Every time the shoulders of the piglet

crossed a zone, a zone change was scored. Looks and watching behaviours were scored as point

events, all other behaviours were scored as state events. Behavioural scores were then calculated to

quantify global responses (see below).

Table 1: Ethogram.

Behaviour	Description
Nb zones crossed [1,2]	The number of times the piglet crossed a virtual zone
Nb approaches H [1]	Number of times the piglets entered the proximal area
Time watching H [1,2]	The amount of time the piglet spent watching the human
Latency to contact H [1,2]	The latency to the first contact of the human by the piglet

Nb looks toward H [1,2]	The number of times the piglet turned its head toward the human
Nb looks other than H [1]	The number of times the piglet looked at other parts of the room than the human or the floor (walls, doors)
Time watching room [1]	The number of times the piglet watched other parts of the room than the human or the floor (walls, doors)
Time in proximal area [1,2]	The amount of time the piglet spent in the proximal area
Time in distal area [1,2]	The amount of time the piglet spent in the distal area
Time in contact H [1,2]	The amount of time the piglet investigated the human
Time investigating floor [1,2]	The amount of time the piglet investigated the floor
Nb contacts H [2]	Number of times the piglet was in contact with the human (initiated by the piglet or the human)

[1]: Scored during reunions of Isolation/Reunion tests. [2]: Scored during reunions of conditioning sessions

Acoustic monitoring and analyses

Vocalisations were recorded with an AKG C314 microphone placed in the center of the room and

one meter above the ground, connected to a Marantz MD661MK2 recorder. Vocalisations produced

during each phase of the trial were manually annotated according to vocal type (grunt, squeal, bark,

scream and mixed calls (Kiley, 1972)), after visual inspection of spectrograms using the 'Annotate'

function of the Praat software (Boersma & Paul, 2001), version 6.0 from http://www.praat.org/.

Checking the occurence of each call type in the several contexts of the study, we confirmed that

'grunt' was the call type used in all contexts and by most of the piglets in each context. So only the

spectro-temporal structure of grunts was further analysed. For information, a table of the number of

each call types recorded in each context as well as the number of individuals involved in the count is

presented in the electronic supplementary material. We could not conduct a robust statistical analysis

on call type utterance, due to the rarity (per subject and tests) of other vocalisations than grunt. (table

S5).

A spectro-temporal analysis was performed with custom-written codes using the Seewave R package

(Sueur et al., 2008) implemented in R (R Core Team, 2015). We first studied the spectral properties

of the remaining background noise of the experimental room (electric noises and remaining low

frequency noises from the rest of the building), using 20 examples of 0.5 second fragments. Since the

first quartile (Q25) of the normalized spectrum of the background noise was 250Hz and the grunts

are low frequency vocalisations, we decided to remove all frequencies below 200Hz in order to focus

on the most relevant frequencies, using a 0.2-8 kHz bandpass filtering ('fir' function). As a

consequence, all results presented in this study are on a 0.2-8kHz frequency range, and no

conclusions on possible frequency components of grunts below this 200Hz threshold can be drawn

here. To measure grunt duration, a 5% to maximal amplitude threshold was used ('timer' function).

After normalisation, the following spectral parameters were calculated using the 'specprop' function
(FFT with Hamming window, window length = 512, overlap = 50%): mean (Q50), first (Q25) and
third (Q75) quartiles, interquartile range (IQR), centroid and standard deviation (all in Hz). The grunt
dominant frequency (in kHz) was also calculated ('dfreq', 50% overlapping FFTs, window length =
512), which is the mean over the grunt duration of the frequencies of highest energy of each window.
Frequency peaks were detected and the minimal and maximal peaks were kept as descriptors
('fpeaks' function, window length = 512, peak detection threshold = 10% of the normalized
amplitude). Measures of noisiness and entropy of the grunts were assessed using: Shannon entropy
(sh), Spectral Flatness (Wiener entropy, sfm) and Entropy (H) [combining both Shannon and
Temporal envelop entropy, length = 512, Hilbert envelop). Two vocal scores were used: the
logarithm of grunt duration and a built-in spectral vocal score with all spectral parameters (see
below). A table describing mean and range of variation of each acoustic parameter in the relevant
contexts of the study is available in the supplementary material (table S4).

Statistical analyses
<u>Behavioural and vocal response scores</u>
The symmetrical distribution of all behavioural parameters on the one hand and all acoustic
parameters on the other hand was visually inspected, and linear transformations were computed
when necessary to reach symmetrical distribution (see tables 2, 3, 4). Two Principal Component
Analyses (PCA, one for the behavioural analysis and one for the spectral acoustic analysis) were
performed using all parameter having a symmetrical distribution ('dudi.pca' function from 'ade4' R
package (Dray & Dufour, 2007) and 'inertia.dudi' function to extract the loadings). Indeed, PCAs are
generally used to reduce the number of variables used in statistical models. It also generates
quantifiable global descriptors of behaviours or acoustic parameters, since correlated parameters
usually load on the same PC (McGregor, 1992). All PCs having an eigenvalue above one were kept
and constituted response scores of behavioural ('ReuPCs' and 'CondPCs' in table 2 and 3
respectively) and vocal ('VocPCs', table 4) parameters. Only the duration of grunts was kept
separated from the spectral parameters to keep it as a temporal parameter.

Table 2: Percentage of explained variance and relative loadings of parameters on PCs, following the
Principal Component Analysis computed on the behaviours scored during the reunion of the
Isolation/Reunion test. The first three PCs, having an eigenvalue above 1, constituted three
behavioural scores: ReuPC1, ReuPC2, ReuPC3. Parameters that explain the most each PC are
bolded (|loading| >0.4).

	ReuPC1	ReuPC2	ReuPC3
Cumulative variance explained %	38.3	60.8	74
Nb of zones crossed	24.177	**-55.843**	-0.435
Nb approaches H	**47.748**	-30.163	0.578
Time watching H	**-52.914**	-7.422	25.585
Latency to contact H	**-64.232**	-0.464	1.688
Nb looks toward H	-7.787	**-43.721**	31.633
Time watching room	-32.048	-13.581	-6.238
Nb looks other than H	3.524	**-72.408**	-2.027
Time in proximal area	**69.96**	-0.156	9.584
Time in distal area	**-46.416**	-12.437	-1.215
Time in contact H	**61.041**	3.586	24.183
Time spent investigating floor	11.868	-7.503	**-42.265**

Table 3: Percentage of explained variance and relative loadings of parameters on PCs, following the Principal Component Analysis computed on the behaviours scored during the sessions of additional positive contacts of the conditioning. The first three PCs, having an eigenvalue above 1 constituted three behavioural scores: CondPC1, CondPC2, CondPC3. Parameters that explain the most each PC are bolded (|loading| >0.4).

	CondPC1	CondPC2	CondPC3
Cumulative variance explained %	41	68.5	80.7
Time in proximal area	**80.23**	2.542	-0.112
Time in distal area	-33.826	8.547	30.789
Number of contacts H	**78.55**	6.476	2.288
Time in contact H	**86.625**	0.715	-0.369
Nb looks toward H	-2	**79.898**	-0.745
Time watching H	-6.757	**65.67**	-10.325
Nb of zones crossed	0.129	33.599	**48.457**
Time spent investigating floor	0.006	**-49.286**	14.205
Latency to contact H	**-81.01**	-0.248	-2.83

Table 4: Percentage of explained variance and relative loadings of parameters on PCs following a Principal Component Analysis on spectral parameters of the grunts recorded in the entire dataset (including both types of tests, N=17 546 grunts). The transformations used to reach symmetrical distribution before the PCA are indicated in parentheses. The first three PCs, having an eigenvalue above 1 constituted three vocal response scores: VocPC1, VocPC2, VocPC3. Parameters that explain the most each PC are bolded (|loading| >0.4) .

	VocPC1	VocPC2	VocPC3
Cumulative variance explained %	59.769	76.807	87.712
Mean Dominant Frequency[1]	-13.558	**53.557**	2.220
Min frequency peak[1] (log)	-0.349	**58.758**	24.236
Max frequency peak[1]	**-43.023**	8.760	-9.537
Mode[2] (log)	-0.522	**66.248**	19.268
Mean[2] (log)	**-95.092**	-2.295	2.028
Q50[2] (log)	**-85.278**	0.280	-0.093
Q25[2] (log)	**-52.360**	19.327	0.985
Q75[2] (sqrt)	**-88.925**	-4.645	2.309
Centroid[2] (log)	**-95.092**	-2.295	2.028
Sd[2]	**-64.484**	-11.303	7.680
IQR[2]	**-87.981**	-5.851	2.640
Sfm[3] (sqrt)	**-94.344**	-3.189	0.962
Sh[3] (sqrt)	**-96.087**	-0.785	-0.175
H[3]	**-88.205**	-1.059	-1.063
Skewness[4]	28.032	-18.010	**48.652**
Kurtosis[4]	22.973	-16.241	**50.615**

[1]: parameters related to the pitch of the vocalisation; [2]: parameters related to the frequency distribution descriptors; [3]: parameters related to the noise component of the vocalisation; [4]: parameters related to the shape of the frequency distribution

Statistical models

All statistics were carried out on R (R Core Team, 2015). Linear mixed effect models ('lmer'
function, 'lme4' R package (Bates et al., 2014)) were built when studied variables were linear
(behavioural and vocal scores, grunt duration) and one binomial generalized mixed effect model was
built for binary parameters (occurrence of missed contacts initiated by human during the
conditioning). The following subsections describe how models were built for each type of tests. In all
models described below, the identity of the replicate ('1' or '2') was used as an interacting fixed
factor, since the experiment was run in two identical replicates on two independent groups. The
identity of the human ('AH' or 'AV') was used as interacting fixed factor in all models described
below, since two experimenters were involved in the positive handling at weaning and in the session

of additional positive contacts of the conditioning (but always the same human was attributed to a

given piglet). The piglet was used as random factor to take into account the within-subject design.

Isolation/Reunion tests

The aim of this part was to test the effect of the positive handling at weaning treatment (H vs. H+

piglets) and additional human contacts during sessions of the conditioning on the piglet's reaction to

human presence. Since the same Isolation/Reunion test was repeated before and after the

conditioning, we used the variable 'Conditioning time' as a two level interacting factor ('before' or

'after' the conditioning) to test the effect of the conditioning.

```
292  Model1 <- lmer (Response variable ~ Treatment*Time + Treatment*Replicate +
293  Treatment*HumanID + Time*Replicate + Time*HumanID + (1 | pigletID), data=
294  dataBehaviourReunion).
```

Concerning the analysis of vocal behaviour, the isolation phase represents a negative social context

for the piglets and may be used as a negative control when monitoring the effect of human presence

on vocal expression of emotional states (Villain et al. 2020a). So, the two phases of the test were

used to study the three way interaction between the treatment (H vs.. H+), the phase of the test

(isolation vs.. reunion) and the time of the conditioning (before vs.. after). The following model was

computed :

```
301  Model2 <- lmer (Vocal response score ~ Treatment*Phase*Time + Treatment*HumanID
302  + Time*HumanID + Treatment*Replicate + Time*Replicate + (1 |
303  pigletID/Time/Phase) , data= dataVocalIsolation + dataVocalReunion).
```

304   To go further, only the reunion phase was kept and a proximity variable was added. Indeed, the

305   piglet could vocalise either when near the human or away from them and this spatial proximity was

306   demonstrated as an important factor of changes of vocal features (Villain et al. 2020b). Thus, a two

307   level proximity factor was built: either '1' when the piglet was in the proximal area (figure 1) or '0'

308   when it was elsewhere in the room.

```
309  Model3 <- lmer (Vocal response score ~ Treatment*Time*InProxArea +
310  Treatment*HumanID + InProxArea*HumanID + Treatment*Replicate +
311  InProxArea*Replicate + Time*Replicate + Time*HumanID + (1 | pigletID/Time), data
312  = dataVocalReunion).
```

Conditioning trials

The aim was to study the evolution of human-piglet relationship over the conditioning [the variable

'Trial number', used as a continuous variable], depending on the previous experience piglets had

with the human [either already familiar (H+ group) or unfamiliar (H group) at the beginning of the

conditioning]. Trial number (Trial) was also used as a random slope to take into account individual

trajectories (Schielzeth and Forstmeier 2009). The following model was built to test the behavioural

response scores (lmer) and the occurrence of missed contact initiated by the human during a session

(presence/absence, binomial model, glmer):

```
321   Model4  <— (g)lmer (Behavioural Response score ~ Trial*Treatment + Trial*HumanID
322   + Trial*Replicate + Treatment*Replicate + Treatment* umanID + (1+ Trial |
323   pigletID), (family=Binomial), data= dataBehaviourConditioning).
```

324   For the analysis of vocal response scores, similarly to the Isolation/Reunion test, the piglet could

325   vocalise either when near the human or away from them. We thus added the proximity factor in the

326   analysis of vocal response variables. The following model was built :

```
327   Model5      <—   lmer   (Vocal   Response   score   ~   Trial*Treatment*InProxArea+
328   Trial*HumanID + Trial*Replicate + Treatment*Replicate + Treatment*HumanID +
329   HumanID*InProxArea  +  Replicate*InProxArea   +  (1+  Trial  |  pigletID),  data=
330   dataVocalConditioning).
```

Model validation and statistical tests

All linear models were validated by visual inspection of the symmetrical and normal distribution of

the residuals. Anovas ('car' R package (Fox & Weisberg, 2011)) were computed on models to test

for significant effects of explanatory variables. Following the Anova, when interactions were found

significant, post hoc test were run on model interactions, correcting for multiple testing with Tukey

contrasts ('emmeans' or 'lstrends' functions from 'emmeans' R package (Lenth, 2016), for

categorical or continuous variables respectively). Results of the Anova, model estimates and pairwise

post hoc comparisons are reported in the supplementary material (tables S1 and S2 for tests, table S3

for model estimates).

Results

Effect of the conditioning process on piglets' reaction to human presence (Isolation/Reunion tests)

Piglets that were not positively handled at weaning express a similar behavioural proximity to a human after a positive conditioning as positively handled ones.

Figure 2: Effect of conditioning and treatment on spatial behaviour and proximity toward the human during the reunion of the Isolation/Reunion test. Mean ± SE per group is indicated, different letters indicates significantly different groups. Significant interaction between treatment (H : grey squares and H+ : black circles) and time (Before the conditioning: empty elements and After the conditioning: filled elements) on behavioural PC1 (letters a to c) and PC3 (letters z and y). Full statistical report is available as supplementary material (tables S1 S2 for statistical tests and S3 for model estimates)

The interaction between the treatment (positively handled piglets before the conditioning (H+) or not (H)) and the conditioning time (before or after the conditioning) was significant for both ReuPC1 and ReuPC3 ($\chi^2_1 = 28.0$, $p < 0.001$, and $\chi^2_1 = 3.7$, $p = 0.05$ respectively, figure 2) but not for ReuPC2 ($\chi^2_1 < 0.001$, $p = 0.99$, supplementary table S1). Post hoc tests on ReuPC1 showed that ReuPC1 was higher after the conditioning than before (H: after – before, t.ratio = 12.1, $p < 0.001$, H+: after – before t.ratio = 11.0, $p < 0.001$) and that before the conditioning, piglets that were positively handled at weaning had significantly higher ReuPC1 than non handled piglets (Before, H – H+: t.ratio = -2.1, $p < 0.001$), but not after (After, H – H+: t.ratio = 0.02, $p = 1.0$). According to the loadings, this

means that piglets that were positively handled at weaning had a lower latency to contact the human,

approached them more often and spent more time close to and investigating the human (ReuPC1)

than non handled piglets, before the conditioning. This score increased after the conditioning and no

evidence of a difference between treatments after the conditioning was found (figure 2). Post hoc

tests on ReuPC3 showed a significant effect of the conditioning time only in piglets that were

positively handled at weaning (H+: after – before, t.ratio = 5.2, p < 0.001, H: after – before, t.ratio =

2.6, p = 0.06). No difference in ReuPC3 was found between treatments before the conditioning

(Before: H – H+, t.ratio = -0.75, p = 0.87), whereas positively handled piglets had a higher -ReuPC3

after the conditioning than before (After : H – H+, t.ratio = -3.2, p = 0.009). According to the

loadings, this means that piglets that were positively handled at weaning expressed more

investigation of the room after the conditioning than before. No evidence of any effect on ReuPC2

was found (table S2).

Positive handling at weaning decreases grunt duration even when no human is present
with the piglet

Comparing the effect of the phase of the test (Isolation vs.. Reunion with the human), taking into

account the conditioning time and the treatment, no evidence of any effect of neither the three way

interaction (χ^2_1 < 0.62, p > 0.43) nor two way interactions of interest was found (treatment: phase,

conditioning time:phase, conditioning time: treatment interactions : χ^2_1 <3.5, p > 0.06, table S2) in

any of the scores. However, grunts produced by positively handled piglets were shorter than grunts

produced by non handled piglets (χ^2_1 = 5.5, p = 0.02, estimates of log(duration)[95% confidence

interval]: -1.25[-1.32;-1.19] and -1.12[-1.2;-1.1] respectively in H+ and H piglets, table S3). Single

effects of the phase of the test were significant for grunt duration and all AcPCs (χ^2_1 > 6.6, p < 0.01,

table S1). Grunts produced during the reunion phase with the human were shorter (estimates of

log(duration) : -1.32[-1.37;-1.26] vs.. -1.06[-1.12;-1.00]) and, according to the loadings, grunts

produced during the reunion phase had a higher frequency range, higher bandwidth and a higher

noise component (-VocPC1: 0.78[0.48;1.08] vs.. 0.34[0.03;0.66]), higher pitched (VocPC2: -0.18[-

0.36;0.01] vs.. -0.46[-0.65;-0.28]) and their spectrum had a higher skewness and kurtosis (VocPC3:

-0.25[-0.37;-0.14] vs.. -0.11[-0.23;0.01]), compared to the isolation phase.

The conditioning process attenuates the effect of proximity on grunts vocal features in
non handled piglets

Figure 3: Acoustic structure of grunt during the reunions with a silent and static human (Isolation/Reunion test). Effect
of conditioning (before or after), treatment (H or H+), and location of the piglet relatively to the human (close: dark blue
or away from them: light blue). Violin plots representing the median and the density of data distribution in the
considered groups. (A, B) Results of post hoc tests following the significant three way interaction between the treatment,
the conditioning time and the location on grunt duration (A) and on the first vocal score (-VocPC1, B). (C,D) Results of
post hoc tests following the significant two way interactions between conditioning time and location (C) and between
treatment and location (D) on the second vocal score (VocPC2). When involved in interaction, the conditioning time was
fixed (as it was relevant to consider difference affected only by time). It thus allowed pairwise comparisons of interacting

location and treatment (A, B) or levels of location (C). Letters represent significantly different groups (p < 0.05). When
no letters are present, no significant difference between groups was found. Stars (*) between two groups represent a
statistical trend (p< 0.10). Full statistical report is available as supplementary material (tables S1 S2 for statistical test
and S3 for model estimates).

During the five-minute reunion, the piglet was scored either as close to the human or away from

them. The three way interaction of the conditioning time, the treatment and the location was

significant for grunt duration, -VocPC1 and VocPC3 (χ^2_1 > 4.9, p < 0.03). Post hoc tests revealed

that grunts produced closer to the human were shorter than the ones produced further away, but only

in piglets that were not positively handled at weaning, effect being stronger before the conditioning

than after it (H piglets: away – close, z.ratio = 6.3, p < 0.001 before and z.ratio = 4.1 p < 0.001 after

the conditioning; H+ piglets: away – close z.ratio < 1.98 p > 0.19, figure 3A). -VocPC1 was higher,

i.e. grunts had a higher frequency range, bandwidth and were noisier when produced closer to the

human than further away, but only in non handled piglets and before the conditioning (H piglets:

away – close, z.ratio = -3.34, p = 0.005 before and z.ratio = -1.23 p = 0.61 after the conditioning; H+

piglets: away – close, z.ratio < 0.36 p > 0.21, figure 3B). For VocPC2, the three way interaction did

not reach significance (χ^2_1 = 3.3, p = 0.07), so only subsequent two way interactions were

considered (but post hoc tests on the three way interaction can be found in supplementary, tables S1

to S3). For VocPC2, significant two way interactions were found between the conditioning time and

the location (χ^2_1 = 10.3, p = 0.001) on the one hand, and between the location and the treatment (χ^2_1

= 4.2, p = 0.04) on the other hand. Post hoc tests revealed that grunts produced closer to the human

had a higher VocPC2, meaning they had a higher pitch, effect being stronger before the conditioning

than after (before: away – close, z.ratio = -6.12, p < 0.001; after: away – close, z.ratio = -2.88, p =

0.004, figure 3C). The increase in VocPC2 with the location was greater for non handled piglets than

positively handled piglets (H piglets: away – close, z.ratio = -5.54, p < 0.001; H+ piglets: away –

close, z.ratio = -3.82, p = 0.001, figure 3D). The last two-way interaction of interest between the

conditioning time and the treatment did not reach significant level (χ^2_1 = 0.80, p = 0.37). For

VocPC3, post hoc tests did not reach significant levels (|z.ratio| < 2.3 p > 0.09 for any comparison) .

Emergence of positive perception of human (effect of additional positive contacts sessions
over the conditioning)
The conditioning process increases behavioural proximity
429

Figure 4: Effect of trial number over the conditioning on spatial behaviour and proximity to the human during the 2min
sessions of additional positive contacts of the conditioning. (A, B) Mean ± SE per group. (A) Single effect of trial number
on behavioural scores CondPC1 and condPC2 according to treatment (H: grey, H+: black). (B) Single effect of
treatment on behavioural score CondPC3 and CondPC2. (C) Single effect of treatment on predicted probability of
occurrence of at least one missed contact by the human, mean estimates ± 95% confidence interval from the generalized
mixed effect model. Stars in the legend box represent significant effect of the trial number (A), different letters represent
significantly different groups (B and C). Full statistical report is available as supplementary material (tables S1 et S2 for
statistical tests, table S3 for model estimates).

No evidence of any effect of the interaction between the treatment [positively handled piglets before
the conditioning (H+) or not (H)] and the trial number was found for all behavioural scores
(CondPC1, CondPC2 and CondPC3, table 3). Independently from the treatment, the higher the trial
number the higher CondPC1 (χ^2_1 = 59.3, p < 0.001, slope estimate [95% confidence interval]: 0.20
[0.15 : 0.25]) and the lower CondPC2 was (χ^2_1 = 48.6, p < 0.001, slope estimate: -0.17 [-0.22 : -
0.12]). According to the loadings, over the conditioning, piglets decreased the latency to contact the
human, made more contacts, spent more time in the proximal area and in contact with the human
(condPC1), decreased the number of looks to the human, spent less watching the human and more
time investigating the room (CondPC2) (figure 4A). Independently from the trial number, positively
handled piglets had a lower CondPC2 and a higher CondPC3 than the non handled ones (χ^2_1 = 12.8,
p < 0.001 and χ^2_1 = 7.0, p = 0.008 respectively), meaning that piglets that were positively handled at
weaning expressed a fewer number of looks to the human, spent less time watching them and more
time investigating the room (CondPC2) and crossed more virtual zone during the test (CondPC3)
(figure 4B). The probability of having at least one missed contact by the human during a session was

lower for positively handled piglets than non handled ones ($\chi^2_1 = 9.57$, p = 0.002, figure 4C), with no

interaction with the trial number ($\chi^2_1 = 0.22$, p = 0.064).

Grunt acoustic features depends on spatial proximity to human

Figure 5: Evolution of vocal scores over the conditioning, during the 2min sessions of additional positive contacts. (A, B)
Violin plots representing the median and the density of data distribution in the group. Interacting effect of location (in
proximal area of the human 'close': dark blue) or elsewhere in the room ('away' from the human: light blue) and
treatment (H vs. H+ piglets) on grunt duration (A) and VocPC2 (B). (C) Mean ± SE per group, interacting effect of trial
number and location of piglets on VocPC2. Different letters in A and B represent significantly different groups, "" in C*
represents significant difference between the two slopes. Full statistical report is available as supplementary material
(tables S1-S3).

During the sessions of additional positive contacts of the conditioning, the three-way interaction

between the trial number, the treatment and the location was not significant for any of the vocal

scores ($\chi^2_1 < 0.18$, p > 0.67), allowing the analysis of the two way interactions of interest. The

interaction between treatment and the trial number was not significant for all vocal scores ($\chi^2_1 < 2.5$ p

> 0.11), however grunt duration decreased over the conditioning sessions (trial number:replicate

interaction, $\chi^2_1 < 5.3$ p = 0.02, slope estimate -0.03[-0.04;-0.01] for the lower slope, table S1 and S3).

Independently from the trial number, grunt duration was lower when piglets were located close to the

human and this effect was stronger in non handled piglets than positively handled piglets

(treatment:location interaction: $\chi^2_1 = 15.8$ p < 0.001, away vs.. close, H piglets: z.ratio = 10.2 p <

0.001, H+ piglets: z.ratio = 6.86 p < 0.001, figure 5A). VocPC2 was higher when piglets were close

to the human, but only in non handled piglets (treatment:location interaction, $\chi^2_1 = 7.6$ p = 0.005,

pairwise comparisons away vs. close, in H: z.ratio = -4.9 p z 0.001 and in H+: z.ratio = -2.0 p =

0.21), meaning that non handled piglets produced higher pitched grunts when closer to the human

(figure 5B). The effect of the location on -VocPC1 and VocPC2 depended on the trial number (trial

number : location interaction, χ^2_1 = 3.97 p = 0.05 and χ^2_1 = 6.1 p = 0.01 respectively): -VocPC1 and

VocPC2 were higher when closer to the human with a greater extent later in the conditioning than

earlier (slope comparison away – close, -VocPC1 : z.ratio = -1.80 p = 0.07, VocPC2 : z.ratio = -2.34

p = 0.02). According to the loadings, this means that the frequency range, bandwidth and noisiness of

grunts (-VocPC1) as well as the pitch (VocPC2) decreased over the conditioning when piglets were

located away from the human but remained high when piglets were close (figure 5C and 5D).

Impact of human identity on piglets behaviour and grunt structure
487

Figure 6: Effect of human identity on spatial behaviour and proximity during the reunion of the Isolation/Reunion test.
Violin plots representing the median and the density of data distribution in the group. Different letters represent
significantly different groups. Full statistical report is available as supplementary material (tables S1 and S2 for
statistical tests, table S3 for model estimates).

Since half of the piglets had been assigned to one human experimenter and the other half to another

one, the identity of the human was included in the model. This allowed to test interactions between

the identity of the human and the treatment of positive handling at weaning on the one hand and the

conditioning time on the other hand.

During the reunions of the Isolation/Reunion test, the interaction between treatment and human

identity was significant for the first behavioural proximity score (ReuPC1, χ^2_1 = 6.01, p = 0.01) but

not the others (ReuPC2 and ReuPC3 (χ^2_1 < 1.98, p > 0.16, table S1). The effect of treatment on

ReuPC1 was higher when piglets were handled by the human 'AH' (H vs. H+, AH: t.ratio = -4.77, p

< 0.001, figure 6). When the human 'AV' handled the piglets, for which ReuPC1 scores exhibited

intermediate values, treatment was not significant (AV, H vs. H+: t.ratio = -1.33, p = 0.56). These

interacting effects of the human identity and treatment on behaviour were not found when

considering the reunions of the conditioning ($\chi^2{}_1 < 1.32$, p > 0.25 for all CondPCs, table S1).

Interactions between the human identity and conditioning time were not significant, neither

considering the reunions of the Isolation/Reunion test (ReuPCs, $\chi^2{}_1 < 0.642$, p > 0.42, tables S1),

neither the trial number during the session of additional positive contacts of the conditioning

(CondPCs, $\chi^2{}_1 < 0.11$ p > 0.74, table S1).

Table 5: Significant effects of human identity on vocal response score (VocCP1 and VocPC2) during the reunion of the
Isolation/Reunion test and during the sessions of additional positive contacts of the conditioning. Only significant effect
are presented here but a full statistical report is available as supplementary material (tables S1 and S2 for statistical
tests, table S3 for model estimates). When single effects were interpretable, the Chi-squared statistic are reported. When
significant interactions were significant, post hoc pairwise comparisons were performed with Tukey corrected and are
thus reported. The estimates correspond either to the group estimate and comparisons of groups (categorical fixed effect)
or slope estimates and comparison of slopes (continuous fixed effect, 'Trial number').

Vocal response score	Fixed effect	Levels	Estimate	Lower.95% CI	Upper.95%CI	Statistic	P-value
Reunion of the Isolation/Reunion test							
VocPC2	humanID	AH	0.154	-0.119	0.427	$\chi^2{}_1 = 4.94$	P = 0.03
		AV	-0.292	-0.571	-0.012		
Sessions of additional positive contacts of the conditioning							
-VocPC1	humanID	AH	0.091	0.037	0.144	$\chi^2{}_1 = 4.69$	P = 0.03
		AV	0.076	0.021	0.132		
VocPC2	humanID*InProxArea	AH – away	0.317	0.110	0.524	Z-ratio = -1.23	P = 0.60
		AH – close	0.402	0.161	0.643		
		AV – away	0.027	-0.182	0.236	Z-ratio = -5.77	P < 0.001
		AV – close	0.462	0.212	0.712		
VocPC3	humanID * Trial number	AH – Trial number	-0.048	-0.070	-0.026		
		AV – Trial number	-0.007	-0.031	0.016	Z-ratio = -2.82	P = 0.005

Considering the vocal scores, no effect of human identity was found on VocPC1 during the

Isolation/Reunion tests but -VocPC1 was higher when the human 'AH' was in the room during the

reunion periods of the conditioning (table 5), meaning the frequency range and the bandwidth of the

grunt were higher when the human 'AH' interacted with the piglet compared to the human 'AV'.

VocPC2 was higher when the human 'AH' was in the room during the Isolation/Reunion tests (table

5), meaning that the pitch of grunts was higher and this effect was also found during the sessions of

additional positive contacts of the conditioning in interaction with the location of the piglet ($\chi^2_1 =$

11.9, p = 0.001): VocPC2 increased when piglets were located close to the human but this increase

was significant only for the human 'AV' and not for 'AH' (table 5). VocPC3 was not different

between humans during the reunions of the Isolation/Reunion tests but, over the conditioning,

VocPC3 changed differently when piglets were handled by the human 'AH' or 'AV', as showed by

the significant interaction between trial number and human identity ($\chi^2_1 = 8.0$, p = 0.005): the

skewness and kurtosis of grunts decreased over the conditioning when 'AH' was interacting with the

piglets, but not 'AV' (see slope estimates, table 5). No evidence of any effect of human identity was

found on grunt duration neither during the Isolation/Reunion tests nor during the sessions of

additional positive contacts of the conditioning (table S1).

Discussion

Behavioural evidence of a rapid establishment of interest and proximity toward a human providing additional positive contacts

The standard reunion test with the human before the conditioning showed that the treatment of

positive handling at weaning succeeded in creating two different levels of human-piglet relationship

(H and H+), as positively handled piglets expressed a higher attraction toward the human than non

handled piglets (ReuPC1), parameters considered as indicators of a positive HAR (Rault et al.,

2020). This test also showed that the conditioning modified the behaviour of non handled piglets so

that they finally expressed a similar attraction toward the human as positively handled piglets, after

the conditioning. These results are in line with the behavioural results of the sessions of additional

positive contacts. The analysis of piglets' behaviour every second sessions of the conditioning

showed that, although positively handled and non handled piglets started with different degree of

proximity toward the human (trials 2 and 4, CondPC1), then, over time and for both treatments (H

and H+), piglets expressed a higher attraction toward the human (CondPC1) and avoided less the

human when the latter attempted to interact with them. At the end of the conditioning, piglets from

both groups had similar level of proximity toward the human (trials 8, 10, 11 CondPC1). So it seems

that the conditioning process allowed non handled piglets to compensate the lack of positive

handling before the conditioning and develop a similar proximity toward the human. Two minute

daily sessions of additional positive contacts changed positively the perception of the human for the

piglets, and thus their willingness to interact with them. Since no evidence of any interaction

between time and treatment was found, no conclusion on differential developmental trajectories

between treatments can be drawn, but a parallel development of the human-piglet relationship in

both groups, when considering the proximity.

Beside behavioural proximity, piglets that were positively handled at weaning expressed more

exploratory behaviours than non handled piglets after the conditioning (ReuPC3). This was also

observed during the sessions of additional positive contacts of the conditioning: positive handled

piglets started with a higher score associated with investigation than non handled piglets (CondPC2)

and it held over the conditioning. Piglets that were positively handled at weaning also expressed a

higher mobility than non handled piglets (CondPC3). These observations may be interpreted as an

expression of natural foraging and disinterest from human contact, which may be a sign of positive

welfare (Weerd & Day, 2009). In addition, this could also be interpreted in terms of attachment to

the human. Indeed, attachment to a human may facilitate exploration of novel environments or

objects, as shown in dogs (Palmer & Custance, 2008). A period of positive handling at weaning may

provide an environment secure enough for the piglets to explore their environment in the presence of

the human. Attachment has also been hypothesised in the lambs-human relationship (Tallet et al.,

2009).

Overall, the behavioural monitoring showed that two minute sessions of positive additional contacts

per day are sufficient to increase proximity to a human to similar levels as when piglets were

previously familiarised for 2 weeks, even when piglets experienced social isolation. But it did not

allow the non handled piglets to express natural exploratory behaviours as the positively handled

piglets. We may be able to hypothesize a sequential establishment of a positive HAR over time:

firstly with a decrease of attentive state and an increase in proximity and accepted contacts, and

secondly with a disinterest of human contacts and the expression of natural foraging behaviour. The

latter may require a higher exposure time.

Links between vocal expression and positive HAR

In this study, piglets were subjected to two types of interactions with the human: one in a standard

reunion test during which, no movements nor speech was produced by the human. The second type

of reunion consisted of sessions of additional positive contacts, during which the human actively

interacted with the piglet, providing contacts and producing speech. These two types of interactions

had different effects on vocal expression, which allows us to evaluate the origin and functions of the

vocal flexibility expressed in grunt structure. In the next two paragraphs, we discuss the results of the

standard reunion test before and after conditioning. This enables us to raise two possible hypotheses.

We then use the results of the sessions of additional positive contacts of the conditioning to discuss

theses hypotheses.

<u>Human mere presence affects vocal expression according to previous experience</u>

Social isolation was associated with longer and lower pitched grunts with a downshifted frequency

spectrum. A reunion with a static human changed grunt structure to shorter, higher pitched with an

upshifted frequency spectrum and this was observed independently from the treatment (H or H+) and

the conditioning time (before or after). In terms of emotional indicators, similar changes in acoustic

features of grunts were already found in studies focusing on vocal markers of valence in pigs

(Briefer et al., 2019, 2022; Friel et al., 2019; Villain et al., 2020) and are also in line with previous

results in similar contexts in relation to the human presence (Villain et al., 2020b). These results,

combined with the behavioural results, may validate The symmetrical distribution of some vocal

parameters encoding of positive emotions (shorter and higher pitched grunts) in the context of a

reunion with a human, potentially releasing the piglets from the stress of isolation.

In addition, and surprisingly, positively handled piglets produced shorter grunts than non handled

piglets whatever the context of observation (with or without human presence). This was previously

shown in another context (anticipation of (pseudo)social events independently from the type of

partner) in the same groups of piglets (Villain et al. 2020). This may show that the period of positive

handling at weaning modulated general vocal production in the studied pigs in the long term. To our

knowledge, long-term effects of a period of positive handling at weaning on grunt duration has not

been studied to confront this finding to the literature.

We showed that the proximity to the human changed the structure of piglets grunts and that

proximity significantly interacted with the treatment and the conditioning time. Indeed, similarly to a

previous study (Villain et al. 2020a, 2020b), during the standard reunion test (no contact from the

human), piglets produced shorter and higher pitched grunts with an upshifted frequency spectrum

when closer to the human. It was especially the case in non handled piglets, before the conditioning.

These results may be linked to the reactivity to the human, non handled piglets being more reactive

to the presence of a human than positively handled piglets. Indeed, before the sessions of the

conditioning, positively handled piglets were habituated to a human interacting positively when

present whereas non handled piglets were not, hence, during the first standard reunion test, when the

human is present but do not interact with the piglet, positively handled and non handled piglets may

have diverging expectations regarding the presence of the static and silent human. As positively

handled piglets received positive contacts every time they were in the presence of the human, they
may have expected positive contacts when approaching and experienced an absence of reward during
the test. This has already been hypothesised in piglets deprived from human voice during interactions
after a period of habituation to it (Bensoussan et al. 2020). On the contrary, piglets that were not
positively handled at weaning never experienced additional positive contacts and being close to a
human, having the possibility to investigate them may be some kind of reward after the period of
total isolation. After the conditioning, piglets from both treatments were conditioned to receive
additional positive contacts and both groups had experienced a first standard reunion test, so they
may both experience an absence of reward during the test, which may explain a lower reaction to
proximity to the human, and thus fewer changes on grunt spectro-temporal features.

The results on vocal expression during the standard reunion test show that even in a context in which
the human does not provide positive contacts, the early experience the piglets had with them affected
their vocal expression when close to them and on the long-term. On the one hand, we know that a
positive HAR establishes through successive positive experiences (Rault et al. 2020) and, on the
other hand, HAR may have long term effects on behavioural expressions, as suggested by Brajon et
al. (2015) using cognitive bias tests. We can thus hypothesize this may also be reflected in the way
piglets vocalise. In that case, we may have evidence of expression of another category of affect,
moods, and not only emotional expression. Indeed, as suggested by Schnall (2010), although
emotions are short-term affects triggered by an external stimulus, moods, on the other hand, may be
experienced on a longer term and may not be attributable to a specific stimulus. Although emotions
and moods do not rely on the same time scale, they may interact with one another, and more studies
are needed to understand their effects on vocal expression.

This test may allow us to suggest two potential non-exclusive hypotheses to explain why the effect of
human proximity on grunt acoustic structure attenuates as the familiarity to the human increases. In a
first hypothesis, we could think that this attenuation of acoustic flexibility is due to a decrease in
reactivity to the human, which may be linked to a disinterest of human contacts and an increase in
natural foraging behaviours. In another hypothesis, this attenuation of acoustic flexibility may be due
to a violation of piglets expectations: because the human remains static during the test, this may
inhibit vocal reactions to the proximity. The interpretation of the second type of human-piglet
interactions below may allow us to address these hypotheses.

<u>Providing rewarding additional positive contacts changes the structure of grunts</u>

During the sessions of additional positive contacts of the conditioning, independently from the trial

number, the duration of grunts was lower when the piglets were located close to the human and

similarly short in both experimental treatments, but when away from the human, non handled piglets

had longer grunts than positively handled piglets. In addition, although the pitch (VocPC2) and the

frequency distribution (-VocPC1) increased over the conditioning, it remained higher when grunts

were produced closer to the human and especially in non handled piglets (VocPC2). These results are

in line with the behavioural results showing an increase in proximity to the human over the

conditioning (CondPC1). During this type of interaction, and contrary to the standard reunion test,

changes in acoustic features of grunts when close to the human were consistent over the

conditioning: piglets remained reactive to the proximity to the human over time. This result may not

be in line with the first hypothesis: in the case of a human interacting with a piglet, the effect of the

proximity does not seem to attenuate over time. Hence, the behaviour of the human during a session

impacts the way a piglet vocalises. In that case, we may raise two more hypotheses to explain this

vocal behaviour : either it is linked to the emotional state or it is linked to a specific human-animal

communication, the two explanations may not be exclusive. The first possible explanation may be

linked to the emotional state experienced by the piglets when approaching a human providing

additional positive contact. As a reminder, in the context of the session, the piglet can choose to

approach and stay close to the human, which will provide positive contacts systematically. So the

piglet may anticipate to receive positive contact and systematically being rewarded. When close to

the human, observed changes in frequency distribution of grunts (increased pitch and upshifted

frequency spectrum) are known to be markers of arousal (in multiple mamalian species (Briefer,

2012, 2020) and pigs (Linhart et al., 2015)). As a consequence, this may show that pigs enter a state

of higher arousal when being close to a carrying human. To go further, we show that these spectral

changes were also associated with shorter grunts. Although the duration of grunts is associated with

the valence of a situation, it is not clear whether the duration could also be an indicator of the

arousal; if our hypothesis is true then it would be the first demonstration that shorter grunts are also

indicators of higher arousal positive state in pigs. Last, we can raise the question whether changes in

grunt structure may also be associated with a specific human-pig communication. In other domestic

species, owner directed vocalisations has been shown (in cats, reviewed in (Turner, 2017); in dogs

(Gaunet et al., 2022)). In addition, studies have found similar socio-communicative behaviours

toward a human in socialized pigs and dogs (Gerencsér et al., 2019). Hence, we may profit from

testing the existence of human directed vocalisations in pigs, as consequences of their socio
communicative abilities.

Effect of human identity on piglets' perception: perspectives on HAR

We found that the identity of the human had effects on behavioural and vocal response scores.
Piglets that were handled by the human 'AH' had higher values of behavioural proximity (ReuPC1)
than piglets handled by the human 'AV' during reunion test after a period of isolation. This effect
was not found during conditioning sessions. The effect of the human did not interact with the
conditioning time, leading to the conclusion that the difference between the two experimenter may
have establishment during the period of positive handling at weaning, prior to the conditioning.
Additionally, when the human 'AH' was in the room, piglets produced grunts with a more upshifted
frequency spectrum and a higher pitch than when the human 'AV' was in the room. If upshifted
grunts may be a indicator of positive higher arousal, then we may conclude that 'AH' was more
likely to trigger higher positive states than 'AV'. Interestingly, the human identity and the spatial
proximity had different effects on piglets grunts during sessions of additional positive contacts but
not when the human was static during the standard reunion test. Hence, it is possible that the way one
human interacts (behavioural and vocally) with a piglet may be more or less effective at triggering
positive emotions and thus modifications of grunt structure. Several evidence exists in the literature
that pigs discriminate humans visual and auditory cues (Bensoussan et al., 2019; Brajon et al.,
2015c). Pigs may also show behavioural changes hearing human voice (Bensoussan et al., 2020). We
may question the efficiency of different human features to generate a positive HAR. In our study,
both humans that interacted with the piglets wear exactly the same clothes and standardized their
tactile interactions toward the piglets before starting the study, and agreed on the rhythm and types of
sounds (words, intonation) to use, to minimise generating variability although no systematic controls
of the human behaviour or spectral feature of voices were performed here. It thus remains unclear
whether experimenters interacted differently or if they were initially perceived differently by piglets.
Our results show that the identity of the human may modulate piglet proximity and vocal behaviour
but the design of this experiment does not allow to find the causes of these observations (behaviour,
voice characteristics, or even odour profile). Thus, more studies of human features that are most
likely to generate a positive HAR are needed and may be of interest regarding animal welfare. In
addition, studying human-piglet relationship in a more systematic way, as in other domestic species,
for example the play behaviour in dogs (Horowitz & Hecht, 2016) or the pet directed speech
(Jeannin et al., 2017; Lansade et al., 2021), may shed light on the evolution and converging strategies
of interspecific relationships. However, the influence of human identity did not modify the general

outcomes of our study, but only decreased some effects, suggesting that this variability does not

modify the main results, but should be considered in future studies.

To conclude, we showed that degrees of familiarity toward a human could be reflected in the way

piglets vocalise in their presence, and out of it. We also showed that the spatial proximity toward a

human providing additional care could change the acoustic structure of piglet grunts. These changes

are likely to be linked to positive and more intense emotional states than when piglets are further

away from the human. However, it is still unclear whether the changes in grunt structure could also

be linked to human-animal communication and more studies are needed to determine it. We did also

show that the identity of the human may be of importance, and may generate vocal changes during

additional positive contacts that were not associated with changes in behaviour of the human. More

systematic studies of human behaviour along with pig behaviour during the human-animal

interactions would be needed to have a better understanding of the evolution of HAR, especially

interactive interspecific communication as well as providing new procedures to promote positive

welfare. We suggest that analysing vocalisations structure may be a good tool to assess the quality of

human-pig relationship and help monitor the establishment of a positive HAR.

Authors contributions

728   Conceived and designed the experiment (A.V., C.T., C.N.). Performed the experiment (A.V., C.G.).

729   Collection and processing of the acoustic and behavioural data (A.V., C.G.). Statistical analyses

730   (A.V.). Contributed to the writing of the manuscript (A.V., C.T., C.N.).

Acknowledgments

We acknowledge all the technical staff at UEPR: especially Patrick Touanel and Marie-Hélène

Lohat, who largely participated in handling the piglets. We thank Eric Siroux who helped building

the acoustic chamber at the beginning of the experiment, Remi Resmond for great discussions about

statistics and Bliss Elizabeth Bagnato-Conlin for their carefully proof reading of the manuscript This

project is part of the SoundWel project in the framework of the Anihwa Eranet and funded by ANR

30001199.

Data availability

The datasets used for the study are available at (Villain et al., 2022). The folder contains all datasets

and a readme to match the type of analysis to the proper dataset. We have made sure to report in the

main text of the article which R libraries and which functions in these libraries we used. All formulas

of the statistical models are explicit in the text to facilitate transfer of information and replicate the

analysis. All libraries are open source as well.

References

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting Linear Mixed-Effects Models using lme4. *ArXiv:1406.5823 [Stat]*. http://arxiv.org/abs/1406.5823

Bensoussan, S., Tigeot, R., Lemasson, A., Meunier-Salaün, M.-C., & Tallet, C. (2019). Domestic piglets (Sus scrofa domestica) are attentive to human voice and able to discriminate some prosodic features. *Applied Animal Behaviour Science*, *210*, 38–45. https://doi.org/10.1016/j.applanim.2018.10.009

Bensoussan, S., Tigeot, R., Meunier-Salaün, M.-C., & Tallet, C. (2020). Broadcasting human voice to piglets (Sus scrofa domestica) modifies their behavioural reaction to human presence in the home pen and in arena tests. *Applied Animal Behaviour Science*, *225*, 104965. https://doi.org/10.1016/j.applanim.2020.104965

Boersma & Paul. (2001). Praat, a system for doing phonetics by computer. *Glot International*, *5*(9), 341–345.

Boivin, X., Tournadre, H., & Le Neindre, P. (2000). Hand-feeding and gentling influence early-weaned lambs' attachment responses to their stockperson. *Journal of Animal Science*, *78*(4), 879–884. https://doi.org/10.2527/2000.784879x

Brajon, S., Laforest, J.-P., Bergeron, R., Tallet, C., & Devillers, N. (2015c). The perception of humans by piglets: Recognition of familiar handlers and generalisation to unfamiliar humans. *Animal Cognition*, *18*(6), 1299–1316. https://doi.org/10.1007/s10071-015-0900-2

Brajon, S., Laforest, J.-P., Schmitt, O., & Devillers, N. (2015b). The Way Humans Behave Modulates the Emotional State of Piglets. *PLOS ONE*, *10*(8), e0133408. https://doi.org/10.1371/journal.pone.0133408

Briefer, E. F. (2012). Vocal expression of emotions in mammals: Mechanisms of production and evidence. *Journal of Zoology*, *288*(1), 1–20. https://doi.org/10.1111/j.1469-7998.2012.00920.x

Briefer, E. F. (2020). Coding for 'Dynamic' Information: Vocal Expression of Emotional Arousal and Valence in Non-human Animals. In T. Aubin & N. Mathevon (Eds.), *Coding Strategies*

in Vertebrate Acoustic Communication (pp. 137–162). Springer International Publishing. https://doi.org/10.1007/978-3-030-39200-0_6

Briefer, E. F., Sypherd, C. C.-R., Linhart, P., Leliveld, L. M. C., Padilla de la Torre, M., Read, E. R., Guérin, C., Deiss, V., Monestier, C., Rasmussen, J. H., Špinka, M., Düpjan, S., Boissy, A., Janczak, A. M., Hillmann, E., & Tallet, C. (2022). Classification of pig calls produced from birth to slaughter according to their emotional valence and context of production. *Scientific Reports*, *12*(1), 3409. https://doi.org/10.1038/s41598-022-07174-8

Briefer, E. F., Vizier, E., Gygax, L., & Hillmann, E. (2019). Expression of emotional valence in pig closed-mouth grunts: Involvement of both source- and filter-related parameters. *The Journal of the Acoustical Society of America*, *145*(5), 2895–2908. https://doi.org/10.1121/1.5100612

Broom, D. M. (2011). A history of animal welfare science. *Acta Biotheoretica*, *59*(2), 121–137. https://doi.org/10.1007/s10441-011-9123-3

Brown, S. L., & Bradshaw, J. W. S. (2014). Communication in the domestic cat: Within- and between-species. In P. P. G. Bateson (Ed.), *The domestic cat: The biology of its behaviour* (Third edition). Cambridge University Press.

Dray, S., & Dufour, A.-B. (2007). The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal of Statistical Software*, *22*, 1–20. https://doi.org/10.18637/jss.v022.i04

Fox, J., & Weisberg, S. (2011). *car-package: Companion to Applied Regression*. GitHub. https://rdrr.io/github/jonathon-love/car/man/car-package.html

Friel, M., Kunc, H. P., Griffin, K., Asher, L., & Collins, L. M. (2019). Positive and negative contexts predict duration of pig vocalisations. *Scientific Reports*, *9*(1), 2062. https://doi.org/10.1038/s41598-019-38514-w

Gácsi, M., Győri, B., Miklósi, Á., Virányi, Z., Kubinyi, E., Topál, J., & Csányi, V. (2005). Species-specific differences and similarities in the behavior of hand-raised dog and wolf pups in social situations with humans. *Developmental Psychobiology*, *47*(2), 111–122. https://doi.org/10.1002/dev.20082

Gaunet, F., Savalli, C., & Legou, T. (2022). An exploratory study on dogs' vocalizations towards their owner and food in an unsolvable task. *Applied Animal Behaviour Science*, *246*, 105529. https://doi.org/10.1016/j.applanim.2021.105529

Gerencsér, L., Pérez Fraga, P., Lovas, M., Újváry, D., & Andics, A. (2019). Comparing interspecific socio-communicative skills of socialized juvenile dogs and miniature pigs. *Animal Cognition*, *22*(6), 917–929. https://doi.org/10.1007/s10071-019-01284-z

Horowitz, A., & Hecht, J. (2016). Examining dog–human play: The characteristics, affect, and vocalizations of a unique interspecific interaction. *Animal Cognition*, *19*(4), 779–788. https://doi.org/10.1007/s10071-016-0976-3

Jeannin, S., Gilbert, C., Amy, M., & Leboucher, G. (2017). Pet-directed speech draws adult dogs' attention more efficiently than Adult-directed speech. *Scientific Reports*, *7*(1), 1–9. https://doi.org/10.1038/s41598-017-04671-z

Jerolmack, C. (2009). Humans, Animals, and Play: Theorizing Interaction When Intersubjectivity is Problematic. *Sociological Theory*, *27*(4), 371–389. https://doi.org/10.1111/j.1467-9558.2009.01353.x

Kiley, M. (1972). The Vocalizations of Ungulates, their Causation and Function. *Zeitschrift Für Tierpsychologie*, *31*(2), 171–222. https://doi.org/10.1111/j.1439-0310.1972.tb01764.x

Lansade, L., Trösch, M., Parias, C., Blanchard, A., Gorosurreta, E., & Calandreau, L. (2021). Horses are sensitive to baby talk: Pet-directed speech facilitates communication with humans in a pointing task and during grooming. *Animal Cognition*, *24*(5), 999–1006. https://doi.org/10.1007/s10071-021-01487-3

Laurijs, K. A., Briefer, E. F., Reimert, I., & Webb, L. E. (2021). Vocalisations in farm animals: A step towards positive welfare assessment. *Applied Animal Behaviour Science*, *236*, 105264. https://doi.org/10.1016/j.applanim.2021.105264

Lenth, R. V. (2016). *Least-Squares Means: The R   Package lsmeans*. *69*(1), 1–33. https://doi.org/doi:10.18637/jss.v069.i01

Linhart, P., Ratcliffe, V. F., Reby, D., & Špinka, M. (2015). Expression of Emotional Arousal in Two Different Piglet Call Types. *PLOS ONE*, *10*(8), e0135414. https://doi.org/10.1371/journal.pone.0135414

McGregor, P. K. (1992). Quantifying Responses to Playback: One, Many, or Composite Multivariate Measures? In P. K. McGregor (Ed.), *Playback and Studies of Animal Communication* (pp. 79–96). Springer US. http://link.springer.com/chapter/10.1007/978-1-4757-6203-7_6

Mignon-Grasteau, S., Boissy, A., Bouix, J., Faure, J.-M., Fisher, A. D., Hinch, G. N., Jensen, P., Le Neindre, P., Mormède, P., Prunet, P., Vandeputte, M., & Beaumont, C. (2005). Genetics of adaptation and domestication in livestock. *Livestock Production Science*, *93*(1), 3–14. https://doi.org/10.1016/j.livprodsci.2004.11.001

Palmer, R., & Custance, D. (2008). A counterbalanced version of Ainsworth's Strange Situation Procedure reveals secure-base effects in dog–human relationships. *Applied Animal Behaviour Science*, *109*(2), 306–319. https://doi.org/10.1016/j.applanim.2007.04.002

Peterson, V., Simonsen, H. B., & Lawson, L. G. (1995). The effect of environmental stimulation on the development of behavior in pigs. *Applied Animal Behaviour Science*, *45*(3–4), 215–224. https://doi.org/10.1016/0168-1591(95)00631-2

R Core Team. (2015). *R: A language and environment for statistical computing.* (3.1.1) [Computer software]. R Foundation for Statistical Computing. http://www.R-project.org/.

Rault, J.-L., Truong, S., Hemsworth, L., Le Chevoir, M., Bauquier, S., & Lai, A. (2019). Gentle abdominal stroking ('belly rubbing') of pigs by a human reduces EEG total power and increases EEG frequencies. *Behavioural Brain Research*, *374*, 111892. https://doi.org/10.1016/j.bbr.2019.04.006

Rault, J.-L., Waiblinger, S., Boivin, X., & Hemsworth, P. (2020). The Power of a Positive Human–Animal Relationship for Animal Welfare. *Frontiers in Veterinary Science*, *7*. https://www.frontiersin.org/article/10.3389/fvets.2020.590867

Schmied, C., Boivin, X., & Waiblinger, S. (2008). Stroking Different Body Regions of Dairy Cows: Effects on Avoidance and Approach Behavior Toward Humans. *Journal of Dairy Science*, *91*(2), 596–605. https://doi.org/10.3168/jds.2007-0360

Sueur, J., Aubin, T., & Simonis, C. (2008). Seewave, a free modular too for sound analysis ans synthesis. *Bioacoustics*, *18*(2), 213–226. https://doi.org/10.1080/09524622.2008.9753600

Tallet, C., Linhart, P., Policht, R., Hammerschmidt, K., Šimeček, P., Kratinova, P., & Špinka, M. (2013). Encoding of Situations in the Vocal Repertoire of Piglets (Sus scrofa): A Comparison of Discrete and Graded Classifications. *PLOS ONE*, *8*(8), e71841. https://doi.org/10.1371/journal.pone.0071841

Tallet, C., Sy, K., Prunier, A., Nowak, R., Boissy, A., & Boivin, X. (2014). Behavioural and physiological reactions of piglets to gentle tactile interactions vary according to their previous experience with humans. *Livestock Science*, *167*, 331–341. https://doi.org/10.1016/j.livsci.2014.06.025

Tallet, C., Veissier, I., & Boivin, X. (2009). How does the method used to feed lambs modulate their affinity to their human caregiver? *Applied Animal Behaviour Science*, *119*(1), 56–65. https://doi.org/10.1016/j.applanim.2009.03.012

Turner, D. C. (2017). A review of over three decades of research on cat-human and human-cat interactions and relationships. *Behavioural Processes*, *141*, 297–304. https://doi.org/10.1016/j.beproc.2017.01.008

Villain, A. S., Guérin, C., Noûs, C., & Tallet, C. (2022). *The use of pig vocalization structure to assess the quality of human-pig relationship*. Portail Data INRAE, V1. https://doi.org/10.15454/RTBO3O

Villain, A. S., Hazard, A., Danglot, M., Guérin, C., Boissy, A., & Tallet, C. (2020). Piglets vocally express the anticipation of pseudo-social contexts in their grunts. *Scientific Reports*, *10*(1), 18496. https://doi.org/10.1038/s41598-020-75378-x

Villain, A. S., Lanthony, M., Guérin, C., & Tallet, C. (2020b). Manipulable Object and Human Contact: Preference and Modulation of Emotional States in Weaned Pigs. *Frontiers in Veterinary Science*, *7*, 577433. https://doi.org/10.3389/fvets.2020.577433

Weerd, H. A. van de, & Day, J. E. L. (2009). *A review of environmental enrichment for pigs housed in intensive housing systems*. https://doi.org/10.1016/j.applanim.2008.08.001

13	14	15	16
9	10	11	12
5	6	7	8
1	2	3	4

↔ 65.5cm

■ Location of the microphone

⬚ Virtual zones for behavioural scoring

🔴 Location of the human

▭ Proximal zone   ⬜ Distal zone

A

Before conditioning | After conditioning

B

C

Location of piglet | away from human | close to human