Zika virus NS4A hijacks host ANKLE2 to promote viral replication

Adam T Fishburn[1], Matthew W Kenaston[1*], Nicholas J Lopez [1*], Vivian Hoang[1], Traci N Shiu[1],

Sophia T Haggard Arcé[1], Shahabal S Khan[1], Priya S Shah[1,2**]

[1] University of California, Davis, Department of Microbiology and Molecular Genetics

[2] University of California, Davis, Department of Chemical Engineering

* These authors equally contributed to this work

** Correspondence: prsshah@ucdavis.edu

Keywords: Zika virus, flavivirus, NS4A, ANKLE2, virus-host interaction, microcephaly

Abstract

Zika virus (ZIKV) is infamous among flaviviruses for its unique association with congenital birth defects, notably microcephaly. We previously mapped ZIKV-host protein interactions and identified the interaction between ZIKV NS4A and host ANKLE2, which itself has established ties to congenital microcephaly. In fruit flies, NS4A induces microcephaly phenotypes in an ANKLE2-dependent manner. This suggests that NS4A interacts with ANKLE2 to dysregulate cell behavior and contributes to abnormal host neurodevelopment. Here, we explore the role of ANKLE2 in ZIKV replication to understand the biological significance of the interaction from the viral perspective. We show that knockdown of ANKLE2 reduces replication of two ZIKV strains, across multiple MOIs and timepoints. We observe that localization of ANKLE2 is drastically shifted to sites of NS4A accumulation during infection. We investigate which domains of ANKLE2 mediate this behavior and the interaction with NS4A. Using co-immunoprecipitation, we show that deletion of either the transmembrane or LEM domain has little impact on the interaction, but deletion of both significantly reduces interaction with NS4A. We show that the C-terminal transmembrane domains of NS4A stabilize the interaction with ANKLE2. Finally, we explore this

interaction in other flaviviruses and observe ANKLE2 interacts with NS4A across four additional mosquito-borne flaviviruses. Together, these results suggest NS4A interacts with ANKLE2 through a combination of its transmembrane and LEM domains, bringing it to sites of ZIKV replication to promote replication through an unknown mechanism. Taken together with our previous results, our findings indicate that, in the process of hijacking ANKLE2 for replication, ZIKV disrupts its physiological function to cause disease.

Importance

The ZIKV epidemic led to the astonishing revelation that congenital ZIKV infection is associated with devastating birth defects, including microcephaly. Microcephaly is the condition in which head and brain size are severely reduced, and is often accompanied by intellectual disability. The molecular mechanisms by which ZIKV replicates and causes microcephaly are still incompletely understood. We previously identified the protein interaction between ZIKV NS4A and host ANKLE2, which is associated with congenital microcephaly. In flies, NS4A induces microcephaly in an ANKLE2-dependent manner, suggesting this interaction is crucial for ZIKV pathogenesis. Here, we explore the relevance of this physical interaction for virus replication. We find that ANKLE2 promotes ZIKV replication, concentrates at sites of NS4A accumulation during infection, and interacts with NS4A via its N-terminal domain. Thus, this represents a rare example of a ZIKV-host protein interaction that impacts both disease and virus replication.

Introduction

Zika virus (ZIKV) is a positive sense RNA virus belonging to the genus *Flavivirus* and can cause severe disease. ZIKV infection in otherwise healthy adults typically leads to mild symptoms and very rarely Guillain-Barré Syndrome (1). The primary concern surrounding ZIKV arises from the occurrence of Congenital Zika Syndrome (CZS) in individuals born from mothers

infected during pregnancy (2). CZS is a spectrum of disease and can be clinically characterized by multiple hallmark features, including congenital contractures, ocular anomalies, cortical calcifications, and in the most severe cases, microcephaly (3, 4).

Microcephaly is a condition in which the head and brain size are significantly reduced at birth and is associated with a wide range of complications, including developmental delays, intellectual disability, and predisposition to seizures (5). Beyond CZS, microcephaly can arise from genetic mutations (6), exposure to toxins during pregnancy (7–9), and congenital infections by pathogens such as rubella virus (10, 11), cytomegalovirus (12), and *Toxoplasma gondii* (13).

The specific molecular mechanisms by which ZIKV contributes to abnormal neurodevelopment and microcephaly are not entirely understood. ZIKV can infect placental tissue and compromise placental function (14–16), potentially providing ZIKV access to the fetal compartment. Gestational timing and type I interferon responses *in utero* are also critical in ZIKV susceptibility (17, 18). Upon fetal infection, ZIKV also breaches the blood-brain barrier and infects susceptible neuronal tissue (19–23). Infection of these neuronal tissues is associated with direct tissue damage through the cytopathic effect (24–26), molecular level pathogenesis (27), disruption of cell division (24, 28–30), and dysregulation of developmental pathways (31–37).

Previously, we used affinity-purification and mass-spectrometry to identify ZIKV-host protein-protein interactions that may contribute to the development of microcephaly in CZS (38). By searching for host proteins with known roles in neurodevelopment or associations with microcephaly, we identified the interaction between the ZIKV non-structural protein 4A (NS4A) and host Ankyrin-repeat and LEM domain containing protein 2 (ANKLE2). ANKLE2 has well established roles in human congenital microcephaly, which has been replicated *in vivo* using *Drosophila* (39–41). Mutations in *ankle2* result in small brain phenotypes and cellular defects in neuroblasts in 3rd instar larvae. These phenotypes are rescued by the expression of human ANKLE2, suggesting that human ANKLE2 and *Drosophila* Ankle2 are functionally conserved in

brain development (39, 40). Using this *Drosophila* model, we showed that transgenic expression of ZIKV NS4A induces similar microcephaly phenotypes which are also rescued by the expression of human ANKLE2. Overall, this suggests NS4A induces microcephaly *in vivo* in an ANKLE2-dependent manner (38, 40).

Flaviviruses physically hijack host factors for their own replication and use many virus-host protein interactions to reshape the host cell into a niche that maximizes virus replication (42). Whether NS4A inhibition of ANKLE2 function during development is simply an unfortunate coincidence, or if there is a functional role for this virus-host protein interaction in ZIKV replication is unknown. ANKLE2 is primarily considered a scaffolding protein, facilitating protein-protein interactions between kinases, phosphatases, and their substrates. ANKLE2 localizes to the ER and inner nuclear membrane where it mediates interactions with proteins, including barrier to autointegration factor (BANF1), vaccinia related kinase 1 (VRK1), and protein phosphatase 2A (PP2A), to control nuclear membrane disassembly during cell division (43, 44). NS4A itself is integral to the formation of flavivirus replication compartments in the ER (45, 46) and facilitates other aspects of flavivirus replication through interactions with host proteins (47–50). We hypothesize that NS4A interacts with ANKLE2 to hijack its scaffolding function in a way beneficial to virus replication.

In this study we explore the role of ANKLE2 in ZIKV replication and what structural domains mediate the biophysical interaction between ANKLE2 and NS4A. We show that knockdown of ANKLE2 reduces ZIKV replication across multiple ZIKV strains and MOIs. We find ANKLE2 is concentrated at sites of NS4A accumulation during infection. We identify that the transmembrane (TM) domain of ANKLE2 localizes it to the ER and nuclear lamina, but both the TM and LEM domains are necessary for its physical interaction with NS4A. Finally, we establish the domain of NS4A that interacts with ANKLE2 and show the conservation of the ANKLE2-NS4A interaction and this domain among other mosquito-borne flaviviruses. Together, we report the novel function of ANKLE2 in promoting ZIKV infection, providing evidence that NS4A

102   disruption of neurodevelopment through ANKLE2 may arise from an underlying virus replication

103   mechanism.

Results

Knockdown of ANKLE2 reduces ZIKV replication

To explore the role of ANKLE2 in ZIKV replication, we used a gene perturbation approach.

We depleted ANKLE2 using CRISPRi with two synthetic guide RNAs (gRNAs) (Supplementary

Figure 1A). Huh7 cells stably expressing dCas9 were generated using lentiviral transduction

(Supplementary Figure 1B). We chose Huh7 cells, derived from a hepatocarcinoma, since they

are readily infected by ZIKV *in vitro* (51, 52), and the liver is an established site of ZIKV replication

in vivo (53–55). CRISPRi gRNAs were delivered by reverse transfection. At 72 hours post

transfection (hpt) the expression of ANKLE2 in ANKLE2 knockdown cells was decreased to

~30% of negative control RNA (ncRNA) (Figure 1A and B). *ANKLE2* mRNA was also significantly

reduced by both gRNAs compared to ncRNA (Supplementary Figure 1C). We confirmed that

cell viability of ANKLE2 knockdown cells was comparable to that of cells transfected with ncRNA

(Supplementary Figure 1D).

To determine if ANKLE2 plays a role in ZIKV replication, we infected Huh7-dCas9 cells

with ZIKV after 72 hours of ANKLE2 knockdown. Infection with ZIKV PLCal (Thailand, 2013)

revealed modest (maximum 7-fold decrease) but consistent and statistically significant

decreases in virus titers between ncRNA and *ANKLE2* gRNA treated conditions at 24 hours post

infection (hpi) for a range of multiplicities of infection (MOIs) (Figure 1B). To further support the

role of ANKLE2 in ZIKV replication we repeated this experiment using the ZIKV strain MR766

(Uganda, 1947) and this time measured virus titers at multiple timepoint post-infection for a

single MOI. Western blot analysis of uninfected cells showed that ANKLE2 knockdown persists

for 6 days post transfection, which would span the entire experimental time course for this

experiment (Supplementary Figure 1E). Following knockdown and infection, we observed

consistent decreases in virus titers starting at 24 hpi, with a maximum 10-fold decrease at 72

hpi (Figure 1D). Together, this shows knockdown of *ANKLE2* reduces ZIKV replication in Huh7

cells.

Figure 1: Knockdown of *ANKLE2* reduces ZIKV replication across multiple MOIs. A)

Western blot of ANKLE2 following CRISPRi knockdown (72 hpt). **B)** Densitometry measurement

of ANKLE2 band intensity, normalized to GAPDH. Data shown from three biological replciates.

C) ZIKV PLCal titers at 24 hpi across three MOIs. **D)** ZIKV MR766 titers at MOI 0.1 over 72 hpi.

All error bars represent standard deviation. Assessment of ZIKV PLCal at MOI 0.1 and 1 were

done with five paired biological replicates. The remaining conditions were done with three

technical replicates. Student's two-tailed T-test, paired for PLCal MOI 0.1 and 1, unpaired for

remaining conditions. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

ANKLE2 colocalizes with NS4A and dsRNA during infection

ANKLE2 is known to localize to the ER and inner nuclear membrane where it facilitates

host protein-protein interactions (43). NS4A is essential for remodeling the host ER into a niche

suited for flavivirus replication. NS4A alone can induce membrane curvature which promotes the

formation of virus replication compartments (46). It can also hijack host proteins to further

facilitate ER remodeling (50). Given our previous proteomics that established the interaction

between ANKLE2 and NS4A (38) we speculated that the scaffolding function of ANKLE2 may

be hijacked by NS4A to facilitate replication. We therefore evaluated ANKLE2 subcellular

localization during ZIKV infection. Due to a lack of compatible immunofluorescence-grade

antibodies we generated HEK293T cell lines that express FLAG affinity tagged fusions of

ANKLE2 (ANKLE2-FLAG). As controls, we also generated cell lines that express GFP-FLAG as

a general non-specific control and ANKLE1-FLAG to distinguish between intrinsic localization in

this family of proteins, and localization unique to ANKLE2. We infected these cells with several

ZIKV strains (PLCal, MR766, and PRVABC59) and evaluated the localization of FLAG and

NS4A. Strikingly, we found that ANKLE2 distribution was concentrated to clusters that had near

perfect co-localization with clusters of NS4A during infection (Figure 2A). Conversely, we found

that the localization and behavior of GFP and ANKLE1 did not change following ZIKV infection

(Figure 2B-C). We measured colocalization between FLAG and NS4A signal using Pearson's

correlation and consistently found very high levels of colocalization between ANKLE2 and NS4A

(Pearson's R-value = 0.93 ± 0.04 across all 3 strains), which was significantly higher than GFP

or ANKLE1 with NS4A (R-value = 0.43 ± 0.10, 0.46 ± 0.15, respectively) (Figure 2D).

Figure 2: ANKLE2 localization pattern is altered during ZIKV infection and colocalizes with NS4A. A-C) HEK293T cells expressing ANKLE2, GFP, or ANKLE1 were infected with noted ZIKV strain for 48 hours. Proteins were visualized by immunofluorescence and confocal microscopy: FLAG (green) and NS4A (magenta) and Hoechst (blue). All scale bars = 5 μm. **D)** Pearson's correlation was measured between the FLAG and NS4A signal in infected cells. Five images and n = 16-24 cells measured per condition. One-way ANOVA with Šidák multiple comparisons test, **** $p < 0.0001$.

Given the colocalization of ANKLE2 and NS4A during infection, the positive role of ANKLE2 in the ZIKV replication cycle, and the known role of NS4A in forming ZIKV replication

factories, we next explored if ANKLE2 colocalizes with dsRNA generated in replication factories during viral genome replication. We and others observed that ZIKV MR766 had the highest replicative efficiency in HEK293T cells and therefore used it for our remaining immunofluorescence experiments (56). To evaluate the colocalization of ANKLE2 with sites of ZIKV replication we used antibodies against endogenous ANKLE2 and flavivirus envelope (E) protein (4G2) or dsRNA (rJ2) (Figure 3A-B). We observed high colocalization between ANKLE2 and E (R-value = 0.89 ± 0.05) that clustered similarly in cells to FLAG-tagged ANKLE2 when infected with the same ZIKV MR766 strain (Figure 3G). We observed that clusters of ANKLE2 signal in ZIKV infected cells partially overlapped with sites of dsRNA signal, although quantitative measurement revealed only modest colocalization (R-value = 0.44 ± 0.10), again, likely due to the punctate behavior of dsRNA. To confirm these were indeed sites of ZIKV genome replication we performed the same experiment evaluating the colocalization between NS4A and E or dsRNA, since NS4A is inherently involved in forming replication factories where dsRNA is formed (Figure 3C-D). This revealed very high correlation between NS4A and E (R-value = 0.96 ± 0.02), suggesting these large clusters maybe the location where viral protein accumulates during or after translation. Colocalization between NS4A and dsRNA was lower (R-value = 0.42 ± 0.12) and similar to that of ANKLE2 with dsRNA. In fact, there was no statistically significant difference between them (Figure 3H). As an additional control we compared E and dsRNA colocalization with GFP. GFP had a substantially and significantly lower colocalization than ANKLE2 or NS4A with both E and dsRNA (Figure 3E-H), suggesting that the modest correlation observed for ANKLE2 and NS4A with dsRNA is biologically meaningful. We performed the same analysis in Huh7 cells and observed similar clustering by ANKLE2 in infected cells, although quantitative measurements of colocalization were lower in these cells (Supplementary Figure 2). Collectively, these results suggest that during ZIKV infection, ANKLE2 is present at sites of viral dsRNA formation.

Figure 3: ANKLE2 colocalizes with ZIKV proteins and dsRNA. **A-F)** HEK293T cells were

infected with ZIKV MR766 (MOI 5) for 48 hours prior to fixation and antibody staining. All scale

bars = 5 μm. **G-H)** Pearson's R-value measured using ImageJ colocalization tool of n = 21-27

cells. One-way ANOVA with Tukey's multiple comparisons test, n.s. = not significant, *** p <

0.001, **** p < 0.0001.

NS4A interacts with ANKLE2 through its TM and LEM domains

To further understand the biophysical interaction between NS4A and ANKLE2, we sought

to determine which domains of ANKLE2 were necessary for the interaction. The structure of

ANKLE2 has not yet been resolved. Thus, we employed the structural prediction provided by

AlphaFold2 (57, 58). This revealed high-confidence structured regions corresponding to the

known TM, LEM, and ankyrin repeat (ANK) domains. Surprisingly, this also revealed three

previously uncharacterized high-confidence structures (Supplementary Figure 3). Using this

predicted structure as a template, we then generated seven C-terminal truncation mutants with progressively fewer of these domains or structures, each with FLAG affinity tags (3xF) (Supplementary Figure 4A). To characterize the biochemical behavior of these truncations we performed subcellular fractionation to enrich for cytosolic, membrane, and nuclear/nuclear lamina compartments. Western blotting analysis of these fractions showed the presence of each primarily in the membrane-bound (including ER, Golgi, etc.) and nuclear/nuclear-lamina fractions (Supplementary Figure 4B and C). This result is expected given the established localization of ANKLE2 to the ER and inner nuclear membrane (43, 44), and the role of the TM, which was not deleted, in facilitating ER localization (59). To evaluate interaction of these ANKLE2 truncations with NS4A we co-transfected each with C-terminally Strep-tagged NS4A-2K plasmid we generated previously (38) and then performed FLAG co-immunoprecipitation (co-IP). Intriguingly, FLAG co-IP revealed that each truncation maintained its interaction with NS4A-2K, although Δ54-938 only very weakly interacted (Supplementary Figure 4D). Together, these results showed the C-terminal deletions of ANKLE2 do not substantially impact its localization or interaction with ZIKV NS4A.

Given that C-terminal deletions did not impact ANKLE2 or its interaction with NS4A, we generated additional mutants with either N-terminal deletions of the TM domain only (Δ2-53), the TM and LEM domains (Δ2-157), the LEM domain only (Δ54-158), or the LEM through ANK domains (Δ54-474) (Figure 4A). We performed similar subcellular fractionation analysis which showed that Δ2-53 and Δ2-157 were enriched in the cytosolic fraction and depleted in the nuclear/nuclear lamina fraction, but maintained some membrane (ER) localization (Figure 4B-C). Confocal microscopy of Δ2-53 and Δ2-157 showed disperse signal in the cytosol with decreased overlap with the ER marker Calreticulin, whereas WT ANKLE2 and internal deletions Δ54-158 and Δ54-474 had high colocalization with the ER as measured by Pearson's correlation (Figure 4D-E). Thus, both ER and nuclear/nuclear lamina localization were disrupted in the Δ2-53 and Δ2-157 mutant.

Figure 4: Assessment of ANKLE2 N-terminal truncations and internal deletion mutants.

A) N-terminal ANKLE2 truncations and internal deletion mutants were designed based on a structural prediction by AlphaFold2. TM = transmembrane domain; LEM = LAP2, emerin, MAN1 domain; Unc = uncharacterized structure predicted by AlphaFold2; ANK = ankyrin repeat domain; 3xF = 3x FLAG affinity tag. **B)** Western blot analysis of subcellular fractions of cells transfected with the indicated ANKLE2 construct. SERCA2 and Lamin A/C antibodies were used to validate enrichment of membrane-bound and nuclear/nuclear lamina fractions, respectively, while GAPDH was used as a cytoplasmic marker. **C)** Densitometry analysis was performed on western blot images by measuring the intensity of FLAG/Strep bands relative to each fraction's

marker. These values were then plotted as a fractional distribution for relative intensity across all fractions. **D)** Confocal microscopy of HEK293T cell transfected with noted ANKLE2-3xFLAG truncations. The colocalization of each truncation (green) was compared with the ER marker, Calreticulin (magenta). **E)** Pearson's correlation was quantified from n = 21-41 cells across 5-11 images per condition. All scale bars = 10 μm. One-way ANOVA with Dunnett's multiple comparisons test compared to control (WT ANKLE2 1-938), n.s. = not significant, *** $p < 0.001$, **** $p < 0.0001$.

Co-transfection with NS4A-2K and FLAG co-IP of these ANKLE2 mutants revealed that deletion of both the TM and LEM domains (Δ2-157) strongly ablated interaction with NS4A-2K, whereas deletion of either the TM or LEM domain alone retained the interaction with NS4A-2K (Figure 5A, Supplementary Figure 4D). This was somewhat surprising since deletion of the TM substantially altered subcellular localization that could preclude the interaction with NS4A in cells. To corroborate this biochemical finding, we visualized the subcellular localization of these ANKLE2 truncations along with NS4A during infection. Here, we clearly observed similar clusters of NS4A that colocalized with wild-type ANKLE2, but not the Δ2-53 and Δ2-157 mutants. Instead, we observed our Δ2-53 and Δ2-157 ANKLE2 mutants appeared to be distinctly separated from sites of NS4A accumulation (Figure 5B-C, white arrows). However, this separation is not necessarily indicative of the ability to physically interact, as Δ2-53 still readily interacts with NS4A when assayed by co-IP since subcellular compartments can be mixed during cell lysis. Altogether, these results suggest that while the TM domain controls ER and nuclear lamina localization of ANKLE2, both the TM and LEM regions together contribute to the interaction with NS4A, with presence of at least one being sufficient for the biochemical interaction.

Figure 5: ANKLE2 truncations reveal that ER localization, and the LEM domain are important for interaction with NS4A-2K. **A)** Western blotting of lysate and FLAG-IP samples. Empty vector condition denotes transfection with NS4A-2K-2xStrep plasmid and an empty plasmid instead of one harboring an ANKLE2-3xF sequence to account for potential FLAG-IP background. ANKLE2-3xFLAG truncations were detected with both FLAG and ANKLE2 antibodies to ensure that lower molecular weight bands are degradation products of these truncations. Due to high transfection efficiency of our truncations, endogenous ANKLE2 is only observable with higher exposure and is only faintly visible. **B)** HEK293T cells expressing indicated ANKLE2 construct and infected with ZIKV were analyzed by immunofluorescence microscopy for ANKLE2-3xFLAG (green) and NS4A (magenta) localization, and Hoescht nuclear stain (blue). White arrows denote regions where truncated ANKLE2 is absent from clusters of NS4A. **C)** Pearson's correlation was quantified from n = 21-29 cells across 5 images

per condition. All scale bars = 5 µm. One-way ANOVA with Tukey's multiple comparisons test, n.s. = not significant, **** $p < 0.0001$.

NS4A-ANKLE2 interaction is dependent on NS4A TM2 and TM3

To further explore the determinants of the NS4A-ANKLE2 interaction, we sought to generate NS4A truncation mutants. Using a previous evaluation of dengue virus (DENV) NS4A as a template (46), we initially generated plasmids encoding one of three C-terminal truncations, each removing an additional TM domain (Figure 6A). As done previously with our ANKLE2 truncations, we characterized the localization of these NS4A truncations using subcellular fractionation and confocal microscopy. This confirmed that each was retained in the membrane-bound fraction (Figure 6B and C), and all had similar colocalization with the ER-marker Calnexin (Figure 6D and E). We then expressed these NS4A truncations in HEK293T cells and performed co-IP against the Strep affinity tag to determine interaction with endogenous ANKLE2 (Figure 7A). The first of these truncations, NS4A ΔTM4, had better expression and interaction with ANKLE2 than our NS4A-2K. This also represents a biologically relevant form of NS4A during ZIKV infection, since the 2K-peptide is first cleaved at the C-terminus of NS4A to separate NS4A from 2K-NS4B (60). The other truncations, NS4A ΔTM3-4 and ΔTM2-4, had significantly reduced to no visible interaction with ANKLE2, suggesting that at least TM3 is crucial for stabilizing the interaction with ANKLE2. To corroborate this result, as well as our previous results with ANKLE2 (Figure 5A), we co-transfected our ANKLE2-3xFLAG and NS4A-2xStrep truncation constructs and performed FLAG Co-IP (Figure 7B). Here, to display the importance of NS4A TM domains, we additionally generated a fourth NS4A truncation that does not express any TM domains (NS4A ΔTM1-4). In this experiment we observed that NS4A ΔTM3-4 had detectable but dramatically reduced interaction with ANKLE2. However, NS4A ΔTM2-4 had no detectable interaction. Additionally, ΔTM1-4 had very low expression and we could not draw

conclusions regarding its ability to interact with ANKLE2. Our co-IP results indicate that both

TM2 and TM3 contribute to the interaction with ANKLE2. Finally, we evaluated the effect of the

2K-peptide on the interaction between NS4A and ANKLE2 Δ2-158, which had strongly reduced

interaction with NS4A-2K previously (Figure 5A). We observed faint and reduced interaction

between ANKLE2 Δ2-157 and NS4A-2K, but no detectable interaction between ANKLE2 Δ2-

157 and NS4A ΔTM4, suggesting that the 2K-peptide may stabilize the interaction outside of the

TM and LEM domains, but that the processed form of NS4A requires these ANKLE2 domains

for interaction.

Figure 6: ZIKV NS4A C-terminal truncations are localized to the ER. A) Schematic of NS4A

with annotated amino acid number for consensus predicted transmembrane domains (above).

C-terminal truncations mutants were generated based on these domains, each with C-terminal

2xStrep affinity tags (below). TM = transmembrane domain, 2xS = 2xStrep affinity tag. **B)**

Western blot analysis of subcellular fractions of cells transfected with the indicated NS4A

construct. SERCA2 and Lamin A/C antibodies were used to validate enrichment of membrane-

bound and nuclear fractions, respectively, while GAPDH was used as a cytoplasmic marker. **C)**

Densitometry analysis was performed on western blot images by measuring the intensity of

Strep bands relative to each fraction's marker. These values were then plotted as a fractional

distribution for relative intensity across all fractions. **D)** Confocal microscopy of HEK293T cell

transfected with noted NS4A-2xS truncations. The colocalization of each truncation (green) was

compared with the ER marker, calnexin (magenta). **E)** Pearson's correlation was quantified from

n = 17-26 cells across 5 images per condition. All scale bars = 10 µm. One-way ANOVA, n.s. =

not significant.

Figure 7: TM2 and TM3 of NS4A mediate interaction with ANKLE2 in vitro. A) Western blotting of lysate and Strep IP samples with indicated NS4A. B) Western blotting of lysate and FLAG IP samples of cells co-transfected with indicated ANKLE2 and NS4A.

Conservation of flavivirus NS4A and its interaction with ANKLE2

Given that ANKLE2 promotes ZIKV replication, we next sought to explore the conservation of this interaction amongst other mosquito-borne flaviviruses. We generated NS4A Strep-tagged fusions for these four additional flaviviruses (dengue virus [DENV], West Nile virus [WNV], yellow fever virus [YFV], and Japanese encephalitis virus [JEV]) and evaluated their interaction with ANKLE2 using the reverse FLAG-IP we performed previously (Figure 7B). We observed that these NS4As expressed higher than ZIKV NS4A, and all were immunoprecipitated by ANKLE2-3xFLAG (Figure 8A). To further investigate this, we used Jensen-Shannon divergence to measure conservation at each amino acid position along NS4A (Figure 8B) (62). This showed regions of ANKLE2-interacting region (TM2 and TM3), including the C-terminus have high levels of conservation (Jensen-Shannon divergence score > 0.7). Sequence logo analysis of this region using a broader set of flaviviruses (all DENV serotypes, St. Louis encephalitis virus [SLEV], Powassan virus [POWV] and Langat virus [LGTV]) (63, 64) revealed several amino acids in the interaction domain that were highly conserved and could mediate the interaction with ANKLE2 (Figure 8C).

Figure 8: Conservation of flavivirus NS4A and its interaction with ANKLE2. A) Western

blot of lysate and FLAG IP samples for indicated flavivirus NS4A protein. ANKLE2-3xFLAG co-

transfected with an empty Strep plasmid was used as a control. **B)** Conservation of NS4A

measured by Jensen-Shannon divergence at each amino acid position (62). **C)** Logo analysis of

the ANKLE2 interaction domain of NS4A (TM2 and TM3, amino acids 73-127), compiled from

12 flaviviruses, with consensus TM domains from ZIKV NS4A overlayed (63, 64). Only a portion

of TM1 is shown as a reference.

Discussion

In this study we explore the relationship between ZIKV and host ANKLE2, with a focus

on virus replication. Knockdown of ANKLE2 led to reductions in ZIKV replication at multiple

timepoints across a range of MOIs and was present for multiple ZIKV strains. The observed

decreases in ZIKV replication are modest (maximum effect ~10-fold), likely due to the incomplete

nature of our knockdown. It is possible that full genetic knockout of ANKLE2 would yield more dramatic virus replication defects.

The supportive role of ANKLE2 in ZIKV replication raises exciting possibilities about ANKLE2 function at sites of replication. Beyond BANF1, PP2A, and VRK1 (43, 44), ANKLE2 also interacts with many other host proteins. For example, ANKLE2 also influences the cell cycle by interacting with Aurora-A and ERα to mediate ERα phosphorylation (65). Plentiful other ANKLE2-host protein interactions have been found in large proteomic screens (66–69). Presumably the bulk of these interactions are mediated through its ANK domain, which facilitates scaffolding functions in many proteins (70, 71), or by the LEM domain, which is known to mediate interaction with the inner nuclear membrane lamina and BANF1 (72–74). It is particularly appealing to speculate that ZIKV leverages the protein interaction/scaffolding function of ANKLE2 to facilitate protein interactions within replication compartments. Future mechanistic studies exploring ANKLE2 protein interactions in replication compartments will be valuable.

To our knowledge, the NS4A-ANKLE2 interaction is a rare example of a ZIKV-host protein interaction that dysregulates the host function to cause neuropathogenesis (38, 40) in the process of promoting ZIKV replication. Several other protein interactions impact aspects of neurodevelopment but appear distinct from virus replication. ZIKV NS3 cleaves host BMP2, inducing osteogenesis and intracranial calcification commonly seen in CZS (27). Expression of NS2A *in vivo* impacts neurodevelopment by disrupting adherens junctions in radial glial cells (31). Expression of NS4A and NS4B impairs the growth of neural stem cells *in vitro* and perturbs autophagy (29). Yet the host factors involved in pathogenesis were not linked to virus replication directly in these studies. A notable exception is the interaction between ZIKV Capsid and Dicer. Capsid interacts with Dicer and inhibits its antiviral activity to promote ZIKV replication while simultaneously inducing neurodevelopmental defects (35). These types of virus-host interactions that result in compounding losses for the host represent an exciting model system to simultaneously study virus replication and pathogenesis.

Our work provides molecular-level insight into the biophysical nature of the NS4A-ANKLE2 interaction. Co-IP studies revealed that both the TM and LEM domains of ANKLE2 together mediate the interaction. Interestingly, loss of the TM alone is sufficient to disrupt the concentration of ANKLE2 to sites of NS4A during infection, though the biochemical interaction is still possible following deletion of just the TM domain. The TM anchors ANKLE2 to the ER and nuclear envelope, and its loss reduces this localization in favor of cytoplasmic localization (59). This deletion separates ANKLE2 from NS4A, which would otherwise interact with the LEM domain. This does not imply an inherent necessity of the TM domain for the interaction with NS4A. The presence of this domain varies between organisms with no clear phylogenetic separation. Mammalian (human, primate, mouse, etc.) ANKLE2 and the *C. elegans* homolog LEM-4L all contain a N-terminal TM. *Drosophila* Ankle2 lacks a clear TM but does have structured elements at the N-terminus (Supplementary Figure 5). Interestingly, *Drosophila* Ankle2 still localizes to the ER and is inhibited by NS4A (40), suggesting this structured region may serve as a previously uncharacterized TM, or it has another mechanism of ER localization. Moreover, the interaction with NS4A could still be detected biochemically by co-IP for the TM-deleted ANKLE2. This is likely due to mixing of cytoplasmic and ER compartments during lysis, which enables the interaction through the LEM domain even in the absence of correct ER localization. Thus, ER localization and the TM and LEM domains are the major contributors to the interaction. Dissecting this interaction with amino acid resolution in the future has the potential to identify ANKLE2 mutants that do not interact with NS4A (or vice versa). This would enable direct testing of the importance of this protein interaction for ZIKV replication. Ultimately it could reveal protective ANKLE2 variants which are functional in brain development, but do not interact with NS4A and thus do not support replication. Thus, this compounded loss scenario for the host could be turned into a compounded win

Finally, we identified the NS4A determinants of the interactions and demonstrate that the NS4A-ANKLE2 protein interaction is highly conserved amongst mosquito-borne flaviviruses.

The conserved nature of the interactions suggests that while inhibition of ANKLE2 to cause microcephaly may be unique to ZIKV, coopting ANKLE2 to promote replication maybe a general feature of flaviviruses. By analyzing the sequence of this interaction domain (TM2 and TM3) for diverse flaviviruses, we were able to identify highly conserved amino acids in this region that may mediate the interaction with ANKLE2. Mutagenesis of this region will be critical to generating a ZIKV mutant that does not interact with ANKLE2. Such a mutant could be evaluated for replication, and if viable, neuropathogenesis.

In summary, we report that knockdown of *ANKLE2* reduces ZIKV replication. During infection, ANKLE2 becomes concentrated at sites of virus replication within the ER. The NS4A-ANKLE2 physical interaction is mediated by the N-terminal TM and LEM domains of ANKLE2, whereas NS4A requires TM2 and TM3. Future studies exploring how ANKLE2 function is hijacked by NS4A in virus replication compartments and which amino acids mediate this interaction will provide molecular mechanistic insight into this virus-host protein interaction involved in replication and pathogenesis.

Methods

Cells. HEK293T (gift of Dr. Sam Díaz-Muñoz), Huh7 (gift of Dr. Raul Andino), and Vero (ATCC) cell lines were maintained in Dulbecco's modified Eagle's medium (DMEM, Gibco ThermoFisher) supplemented with 10 % fetal bovine serum (FBS, Gibco ThermoFisher) at 37ºC, 5% CO_2. Cells we washed with Dulbecco's phosphate buffered saline (D-PBS, Life Technologies) and dissociated with 0.05% Trypsin-EDTA (Life Technologies). Cells were tested for *Mycoplasma spp.* monthly by PCR.

Plasmids. Lentiviral plasmid (pHR-UCOE-EF1α-KRAB-dCas9-P2A-Bls) encoding a catalytically dead Cas9 (dCas9) with a C-terminal hemagglutinin (HA) tag was a gift from Dr.

Sean Collins. For stable, inducible expression in HEK293T cells, ANKLE2 and GFP sequences were amplified by PCR and cloned into pLVX-TetOne-Puro, cut with EcoRI, using Gibson assembly. Codon-optimized ANKLE1 sequence was acquired from Twist BioSciences and similarly inserted into pLVX-TetOne-Puro, cut with EcoRI, with C-terminal APEX2 and 3xFLAG affinity tags. ANKLE2 truncations were designed based on AlphaFold2 structural prediction (57, 58) using NM_015114.3 accession sequence (searched August 8[th], 2021). Codon optimized DNA fragments with C-terminal 3xFLAG affinity-tags were acquired from Twist Bioscience and inserted into pcDNA4_TO, cut with KpnI and ApaI, using Gibson assembly. A full length pcDNA4_TO ZIKV NS4A-2xStrep plasmid was previously generated (38), and used as a template for generation of NS4A truncation sequences using PCR amplification. Products were inserted into pcDNA4_TO cut with BamHI and EcoRI upstream of 2xStrep using Gibson assembly. Flavivirus (DENV, YFV, WNV, JEV) NS4A sequences were acquired from Twist Bioscience and inserted into pcDNA4_TO, cut with BamHI and XhoI, with C-terminal 2xStrep tags using Gibson assembly. All plasmids were prepared in Stbl3 or DH5α using MiniPrep (Sigma-Aldrich) or MidiPrep (Macherey-Nagel) kits and verified using sequencing services provided by GeneWiz. All sequence accession numbers are available in Table S2. Primer sequences used for the generation of all our constructs are available in Table S3.

Lentiviral packaging, transduction, and cell selection. Lentiviral packaging and transduction were performed as previously described (75) using the calcium phosphate protocol (76). In short, 3.5 µg of cloning product plasmid were transfected into HEK293T with lentiviral packaging plasmids including 1.8 µg pMDLg/p-RRE, 1.25 µg pCMV-VSV-g and 1.5 µg pRSV-Rev. After 48 hours lentivirus particles were collected, and cell debris was removed by centrifugation (Eppendorf centrifuge 5810 R, Rotor S-4-104, 94 g, 5 minutes) and filtration through a 0.45 µm filter. The resulting lentiviral stocks were used to transduce HEK293T cells. Transduced cells were bulk selected for puromycin resistance (10 µg/ml, ThermoFisher). A control lentiviral

plasmid encoding GFP without a selection marker was used in tandem as a control to ensure

efficient packaging, transduction, and selection.

Viruses and stock preparation. All ZIKV stocks were propagated in Vero cells and monitored

for CPE. Supernatant was then harvested, and cell debris was removed by centrifugation

(Eppendorf centrifuge 5810 R, Rotor S-4-104, 211 g, 5 minutes, 4°C) Cleared supernatant was

then distributed into 500 µL aliquots and frozen at -80°C. Each aliquot was only used once to

prevent repetitive freeze-thaw. Aliquots were titered by plaque assay (method below). Strains

used were ZIKV PLCal/2013 (gift of Dr. Richard Wozniak), ZIKV PRVABC59 (gift of Dr. Lark

Coffey), and ZIKV MR766 (BEI Resources, NIAID, NIH, as part of the WRCEVA program: Zika

Virus, MR 766, NR-50065).

ANKLE2 **CRISPRi knockdown and ZIKV Infection.** Custom synthetic guide RNAs (gRNA)

were acquired from Sigma and resuspended to 3 µM in TE buffer (10 mM Tris Base, 1 mM

EDTA, pH 8.0). For Huh7-dCas9 knockdown in 12-well dishes 30 µL of each gRNA was

combined with 7 µL TransIT-CRISPR transfection reagent (Sigma) in 363 µL OPTI-MEM (Life

Technologies) and complexed at room-temperature for 20 minutes prior to being added to each

well (90nM final gRNA concentration). A total of 1.2×10^5 Huh7-dCas9 cells were then added

and grown overnight at 37°C. Additional DMEM was then added 24 hpt. Viability experiments

were done 72 hpt with ZombieGreen dye (BioLegend, gift of Dr. Scott Dawson) diluted 1:100 in

D-PBS and incubated on live cells for 5 minutes. Ten images were taken for each condition and

total and dead cells were then counted to determine viability. ZIKV replication after *ANKLE2*

knockdown was done by removing media from each well 72 hpt. 2 mL of fresh DMEM and

appropriate volume of ZIKV stock was then added to each well. Aliquots of supernatant were

harvested at 0, 18, 24, 48, and 72 hpi and frozen at –80°C.

Plaque assay. Vero cells were grown as a monolayer in 6-well dishes overnight. Virus aliquots were thawed on ice then subjected to 10-fold serial dilution. Media was removed from Vero cells and the monolayer was washed once with 2mL D-PBS. 500µL of each virus dilution was then added and incubated for 1 hour at 37ºC with periodic rocking. Virus was then removed and cells were overlayed with 3 mL of DMEM with 0.8% methylcellulose (Sigma), 1% FBS, 1% penicillin/streptomycin (Thomas Scientific) and incubated at 37ºC for 4 days. Cells were then fixed with 4% formaldehyde (Fisher) for 30 minutes at room temperature. Formaldehyde and media were then removed, and cells were stained with 0.23% crystal violet solution (Fisher) for 30 minutes. Solution was then removed, and plaques were counted.

Quantitative RT-PCR. RNA was harvested using *Quick*-RNA Miniprep kit per manufacturer's instructions (Zymo). Purified RNA (500ng) was then used to make cDNA using iScript™ cDNA synthesis kit (Bio-Rad). After cDNA synthesis each sample was resuspended to a total volume of 100µL using RNase-Free water. A total of 2 µL of cDNA was then used for each qPCR reaction (done in technical triplicates) using LightCycler 480 SYBR Master Mix (Fisher). Samples were run in a Roche LightCycler® 480 II Instrument. Changes in gene expression were calculated using the Livak method ($2^{\Delta\Delta Ct}$) (77).

Confocal immunofluorescence microscopy. HEK293T or Huh7 cells cultured on #1.5 coverslips were fixed with 4% paraformaldehyde (Fisher) for 15 minutes at room temperature. Cells were permeabilized with 0.1% Triton X-100 (Integra) for 10 minutes and blocked with 5% goat serum (Sigma) in PBS-Tween (0.1% Tween-20, Fisher). Coverslips were incubated with primary antibodies overnight at 4°C. Coverslips were then washed in PBS-Tween and incubated in secondary antibody at room temperature for 1 hour. Nuclei were visualized with Hoechst (Invitrogen). Confocal images were acquired using an Olympus FV1000 Spectral Scan point-scanning confocal fitted to an Olympus IX-81 inverted microscope using a PlanApo 60x/NA1.40

oil immersion lens or Zeiss Airyscan LSM800 with Axiocam using a 63x/NA1.40 oil immersion lens. Laser lines at 405, 488, and 543nm were employed sequentially for each image using optics and detector stock settings in the "Dye List" portion of the FluoView microscope-controlling software. All antibodies and dilutions are listed in Supplementary Table S1. Confocal images were analyzed using ImageJ (Fiji) software (78). Signal colocalization was quantified using Pearson's correlation coefficient (R-value) determined with the "Colocalization 2" analysis tool within Fiji after masking individual cells across at least 5 images.

ANKLE2 and NS4A co-transfection and FLAG affinity-purification. For transfection $5x10^6$ HEK293T cells were plated in 10 cm dishes and grown overnight. Transfection was performed by combining 3.5 µg of each corresponding plasmid DNA with 700 µL of serum-free DMEM. Next, 21 µL of PolyJet transfection reagent (SignaGen) was combined with 700 µL serum-free DMEM and added to each plasmid DNA tube. Samples were mixed and incubated at room temperature for 15 minutes prior to addition to cells. Cells were then grown for an additional 24 hours. Transfection efficiency was confirmed using a GFP encoding plasmid. Media was then removed from each plate. To dissociate cells, 5 mL of D-PBS supplemented with 10 mM EDTA was added and allowed to incubate for several minutes. Cells were resuspended in 5 mL of D-PBS and transferred to 15 mL conical tubes prior to centrifugation 94 g, 4ºC for 5 minutes (Eppendorf centrifuge 5810 R, Rotor S-4-104). Cell pellets were washed with 5 mL D-PBS and centrifugation was repeated. Supernatant was removed and pellets were then resuspended in 1 mL IP buffer (50 mM Tris Base, 150 mM NaCl, 0.5 M EDTA, pH 7.4) with Pierce™ protease (Thermo Scientific) and Pierce™ phosphatase (Fisher) inhibitor tablets supplemented with 0.5% NP-40 Substitute (Igepal™ CA-630, Affymetrix). Cells were lysed for 30 minutes at 4ºC, and lysate was then centrifugated at 845 g, 4ºC, for 20 minutes (Eppendorf centrifuge 5424 R, Rotor FA-45-24-11). A portion of each lysate (60-100 µL) was collected, normalized by BCA assay (Thermo Scientific), and saved for western blot analysis. Remaining lysate was added to 40 µL

of magnetic FLAG beads (Sigma) and incubated overnight at 4ºC with gentle rotation. Beads

were then washed 4 times with 1 mL IP buffer with 0.05% NP-40 and once with 1 mL IP buffer

without NP-40. Beads were then incubated in 40 µL of 100 ng/mL 3x FLAG peptide (APExBIO)

at 211 g for 1 hour at room temperature (Eppendorf ThermoMixerC). Eluate was then removed.

Eluate and lysate were resuspended in NuPAGE LDS sample buffer and bond-breaker TCEP

(Thermo Scientific) according to manufacturer's recommendation. Samples were boiled for 10

minutes at 95ºC prior to evaluation by western blot (below).

Subcellular fractionation. Proteins were isolated from the cytosol, membrane-bound

organelles, and the nucleus using a previously established protocol for cultured cells (79). In

brief, cells were sequentially lysed in buffer (3 M NaCl, 1 M HEPES, 1 M glycerol, 1X protease

inhibitor) containing increasingly stronger detergents. Following cell trypsinization, buffer with

digitonin (25 µg/mL) disrupted the plasma membrane over the course of gentle rotation at 4ºC.

Buffer with Igepal (1% v:v, NP-40 substitute) permeabilized membrane-bound organelles during

incubation on ice. The nuclear membrane was disrupted by buffer containing sodium

deoxycholate (0.5% w:v) and sodium dodecyl sulfate (0.1 % w:v), with subsequent sonification

to disrupt genomic DNA. All separations were performed by centrifugation (Eppendorf centrifuge

5424 R, Rotor FA-45-24-11, 4ºC). Cell fractions were evaluated by western blot.

Western blot. For whole cell lysates, cells were lysed in RIPA buffer (150 mM NaCl, 50 mM

Tris Base, 1% Triton X-100, 0.5% sodium deoxycholate) supplemented with protease inhibitors

for 5 minutes at room temperature. Cell lysate was incubated on ice for 30 minutes prior to

centrifugation (Eppendorf centrifuge 5424 R, Rotor FA-45-24-11, 13,500 g, 4ºC, 20 min). When

possible, the total protein concentration of each sample was normalized by BCA assay. Protein

samples (lysates or IP eluates) were resuspended in NuPAGE LDS sample buffer supplemented

with TCEP and boiled at 95ºC for 10 minutes. Samples were run on 7.5-12% polyacrylamide

gels for ~1 hour at 150V and transferred to PVDF membranes (VWR) for 1 hour at 330 mA on

ice. Membranes were then blocked in 5% milk solution for 1 hour prior to overnight incubation in

primary antibodies (Table S1) at 4⁰C. Membranes were washed 3x in Tris-buffered saline with

Tween-20 (TBS-T) (150 mM NaCl, 20 mM Tris Base, 0.1% Tween-20, Fisher) and incubated

with HRP-conjugated secondary antibodies in 5% milk for 1 hour at room temperature.

Membranes were again washed 3x in TBS-T and 1x in TBS (without Tween-20) prior to Pierce^{TM}

ECL activation (Fisher). Membranes were imaged using Amersham Imager 600 (GE). Western

blot images were analyzed using Fiji. Densitometry was calculated by measuring the band

intensity ratio of the experimental band to the loading control band.

Statistical analysis.

Statistical analysis and plotting were performed using GraphPad Prism 6 software (GraphPad

Prism 6.0; GraphPad Software Inc., La Jolla, CA, USA). Error bars represent standard

deviations. Data were considered statistically significant when a $p < 0.05$ was determined by

Student's T-test or one-way ANOVA with noted multiple-comparison test.

Acknowledgements

We would like to thank members of the Shah lab for their constant support and helpful feedback. Funding was provided by the University of California, Davis and the W. M. Keck foundation. ATF was partially supported by NIH T32 fellowships (5T32GM7377-42 and 2T32AI060555-16). STH was partially supported by NIH T34 (T34GM136469). The Olympus FV1000 confocal used in this study was purchased using NIH Shared Instrumentation Grant 1S10RR019266-01. We thank the MCB Light Microscopy Imaging Facility, which is a UC Davis Campus Core Research Facility, for the use of this microscope. We thank Dr. Neil Hunter's lab for the use of their AiryScan Confocal microscope.

Author Contributions

ATF and PSS conceived of work and designed experiments. ATF performed and analyzed all experiments. MWK, VH, NL, TNS, and STH assisted with experiments. STH, NL, TNS, and SK performed cloning and plasmid preparation. ATF and PSS wrote the manuscript. All authors revised the manuscript. PSS secured funding for work. All authors read and approved the final manuscript.

Supplementary Figures

Supplementary Figure 1: Knockdown of *ANKLE2* in Huh7-dCas9 cells using CRISPRi. **A)** Schematic of *ANKLE2* and the targeting guide RNAs (gRNA) that were used to knockdown *ANKLE2* expression. **B)** Western blot to evaluate the expression of dCas9-HA in Huh7 cells that were used for CRISPRi experiments. **C)** Western blot to evaluate ANKLE2 expression after CRIPSRi knockdown with three different gRNAs. The primary isoform is shown at 104kDa, and the knockdown of various other isoforms can be observed at 76 and 72kdA. **D)** Densitometry analysis to quantify ANKLE2 band intensity. ANKLE2 band intensity was normalized to its respective loading control and then normalized to ncRNA. Error bars represent the standard deviation between 3 biological replicates. **E)** *ANKLE2* gene expression after knockdown was quantified by qRT-PCR. Error bars represent the standard deviation between 3 biological replicates. **F)** Cell viability was evaluated using ZombieGreen dye. Data represent the average cell viability across 10 images for each condition. **D-F)** Error bars represent the standard deviation. Student's unpaired two-tailed t-test, compared to ncRNA, n.s., not significant, * p > 0.05, *** p > 0.001.

Supplementary Figure 2: ANKLE2 colocalizes with sites of ZIKV genome replication in

Huh7-dCas9 cells. **A-F)** Huh7-dCas9 cells plated on cover slips were infected with ZIKV for 24-

48 hours and subsequently immunostained for with noted antibodies and their target (in the case

of 4G2/Env and rJ2/dsRNA). Cover slips were imaged by confocal microscopy and analyzed

using ImageJ. All scale bars = 5 µm **G-H)** Pearson's R-value measured using ImageJ

colocalization tool. N = 22 to 28 cells across 5 images per condition. One-way ANOVA with

Tukey's multiple comparisons test, n.s. = not significant, ** p < 0.01, **** p < 0.0001.

A   Human ANKLE2 AlphaFold Prediction

Supplementary Figure 3: AlphaFold2 protein structure prediction reveals uncharacterized structural regions in human ANKLE2. AlphaFold2 was searched for human ANKLE2 and predicted structures were compared against known domains. AlphaFold2 confidence is shown as a gradient from very high (blue) to very low (orange). Known domains are highlighted, ankyrin repeats = dark blue, LEM = green, TM = yellow, uncharacterized = light blue.

A C-Terminal ANKLE2 Truncations

B Cytosolic / Membrane-Bound / Nuclear/Nuclear Lamina

α-FLAG (ANKLE2)
α-Strep (GFP)
α-SERCA2
α-Lamin A/C
α-GAPDH

C Relative FLAG/Strep Distribution

Nuclear/Nuclear-Lamina
Membrane-Bound
Cytoplasmic

D

α-FLAG (ANKLE2)
α-Strep (NS4A)
α-GAPDH

Lysate / FLAG-IP

Supplementary Figure 4: ANKLE2 C-terminal truncations interact with ZIKV NS4A and are retained primarily in the ER and Nuclear regions. A) C-terminal ANKLE2 truncations were designed based on a structural prediction by AlphaFold2. TM = transmembrane domain; LEM = LAP2, emerin, MAN1 domain; Unc = uncharacterized structure predicted by AlphaFold2; ANK = ankyrin repeat domain; 3xF = 3x FLAG affinity tag. **B)** Western blot analysis of subcellular fractions of cells transfected with the indicated ANKLE2 construct. SERCA2 and Lamin A/C antibodies were used to validate enrichment of membrane-bound and nuclear fractions, respectively, while GAPDH was used as a cytoplasmic marker. **C)** Densitometry analysis was performed on western blot images by measuring the intensity of FLAG/Strep bands relative to each fraction's marker. These values were then plotted as a fractional distribution for relative intensity across all fractions. **D)** Western blot of lysate and FLAG-IP samples for indicated ANKLE2 construct cotransfected with NS4A.

Supplementary Figure 5: Transmembrane domain prediction of ANKLE2 homologs and structural prediction of *Drosophila* Ankle2. A) Transmembrane domain prediction of human ANKLE2, *Drosophila melanogaster* Ankle2 (NM_001298446.1), and *C. elegans* LEM-4L. TM prediction was determined using SOSUI and TMHMM 2.0 modeling (80, 81). **B)** AlphaFold2 was searched for *Drosophila* Ankle2 and predicted structures were compared against known domains. AlphaFold2 confidence is shown as a gradient from very high (blue) to very low (orange). Known domains are highlighted, ankyrin repeats = dark blue, uncharacterized = light blue.

675 **Supplementary Table 1: Antibodies**

Antibody	Host Species	Dilution Used	Supplier (Catalog #)	RRID
GAPDH	Mouse	1:1000 (WB)	Fisher (PIMA515738)	AB_2537652
FLAG-M2	Mouse	1:200 (IF), 1:1000 (WB)	MilliporeSigma (F1804)	AB_262044
Strep	Mouse	1:1000 (IF, WB)	Qiagen (34850)	AB_2810987
4G2 (Flavivirus E)	Mouse	1:250 (IF)	ATCC (HB-112)	CVCL_J890
rJ2 (dsRNA)	Mouse	1:40 (IF)	MilliporeSigma (MABE1134)	AB_2819101
Lamin A/C (4C11)	Mouse	1:1000 (WB)	Cell Signaling (4777) Gift from Dr. Jodi Nunnari	AB_10545756
SERCA2	Mouse	1:100 (IF), 1:500 (WB)	Invitrogen (MA3919)	AB_325502
ANKLE2	Rabbit	1:80 (IF)	Atlas Antibodies (HPA074838)	N/A
ANKLE2	Rabbit	1:1000 (WB)	Atlas Antibodies (HPA003602)	AB_1858349
ZIKV NS4A	Rabbit	1:1000 (IF, WB)	GeneTex (GTX133704)	AB_2887067
Actin	Rabbit	1:5000 (WB)	Sigma (A2066)	AB_476693
Anti-Mouse IgG-HRP	Rabbit	1:5000 (WB)	SouthernBiotech (6170-05)	AB_2796243
Calreticulin	Rabbit	1:500 (IF)	Abcam (AB92516)	AB_10562796
Calnexin	Rabbit	1:250 (IF)	Proteintech (10427-2-AP) Gift from Dr. Jodi Nunnari	AB_2069033
HA	Rat	1:2000 (WB)	MilliporeSigma (3F10) Gift from Dr. Joanna Chiu	AB_2314622
Anti-Rabbit IgG-HRP	Goat	1:5000 (WB)	SouthernBiotech (4030-05)	AB_2687483
Anti-Mouse AlexaFlour-488	Goat	1:1000 (IF)	Invitrogen (A28175)	AB_2536161
Anti-Rabbit AlexaFlour-555	Goat	1:1000 (IF)	Invitrogen (A27039)	AB_2536100
Anti-Mouse AlexaFlour-680	Goat	1:1000 (IF)	Invitrogen (A21057)	AB_2535723

676 WB = Western blot, IF = Immunofluorescence

678 **Supplementary Table 2: Sequences**

Sequence	Accession Number	Relevant Figure(s)
Human ANKLE2	NM_015114.3	Figures 1, 2, 4, 5, 7 and 8 Supplementary Figures 1, 3, 4, and 5
D. melanogaster Ankle2	NM_001298446.1	Supplementary Figure 5
C. elegans LEM-4L	NM_001028343.3	Supplementary Figure 5
Human ANKLE1	NM_152363.6	Figure 2
eGFP	UDY80669.1	Figures 2, 3, 4, and 6 Supplementary Figures 2, and 4
Zika virus (ZIKV)	MK713748.1	Figures 5, 6, 7, and 8. Supplementary Figure 4
Dengue virus serotype 1 (DENV-1)	QZN14214.1	Figure 8
Dengue virus serotype 2 (DENV-2)	NC_001474.2	Figure 8
Dengue virus serotype 3 (DENV-3)	QXM02604.1	Figure 8
Dengue virus serotype 4 (DENV-4)	QTX92397.1	Figure 8
Yellow Fever virus (YFV)	KF769016.1	Figure 8
West Nile virus (WNV)	DQ211652.1	Figure 8
Japanese encephalitis virus (JEV)	NC_001437.1	Figure 8
Saint Louis encephalitis virus (SLEV)	QVN47903.1	Figure 8
Tick-borne encephalitis virus (TBEV)	JX498940.1	Figure 8
Powassan virus (POWV)	L06436.1	Figure 8
Langat virus (LGTV)	NC_003690.1	Figure 8

680 **Supplementary Table 3: Oligonucleotides**

Oligo Name	Sequence	Application
GAPDH Forward	ACATCGCTCAGACACCATG	qPCR Control
GAPDH Reverse	TGTAGTTGAGGTCAATGAAGGG	qPCR Control
ANKLE2 Forward	AAAGAGAACCAGGCTTCCATC	qPCR Analysis
ANKLE2 Reverse	CACGTTGACTACATCTGCATTTC	qPCR Analysis
NS4A Forward	CTTGGTACCGAGCTCGGATCCGCCACCATGGGAGCCG	Gibson Assembly of ZIKVpr NS4A C-terminal truncations
NS4A 1-54 Reverse	CATCCACCGCCTCCCTCGAGCGTTTCGAGAGTTTCCGGGAG	Gibson Assembly of ZIKVpr NS4A 1-54
NS4A 1-72 Reverse	GTGCTGGATATCTGCAGAATTCAAGGACGAAAAATATACCGAGAGAGACG	Gibson Assembly of ZIKVpr NS4A 1-72
NS4A 1-96 Reverse	GTGCTGGATATCTGCAGAATTCCCACATCAGCCATGCGC	Gibson Assembly of ZIKVpr NS4A 1-96
NS4A 1-127 Reverse	GTGCTGGATATCTGCAGAATTCGCGTTGCTTCTCCGGCT	Gibson Assembly of ZIKVpr NS4A 1-127
GFP Forward #1	TAAAGGCCGCCATGGTGAGCAA	Gibson Assembly preparation of GFP-APEX2-3xFLAG
GFP Reverse #1	CGCTCTTGTACAGCTCGTCCAT	Gibson Assembly preparation of GFP-APEX2-3xFLAG
GFP Forward #2	CTTATACCAACTTTCCGTACCACTTCCTACCCTCGTAAAGGCCGCCATGGTGAGCA	Gibson Assembly of GFP-APEX2-3xFLAG
GFP Reverse #2	CAGATCCACCTCCTGAACCACCTCCGCTACCGCCACCGCTCTTGTACAGCTCGTCCATGC	Gibson Assembly of GFP-APEX2-3xFLAG
ANKLE2 Forward #1	TAAAGGCCGCCATGCTGTGGCCGCGG	Gibson Assembly preparation of ANKLE2-APEX2-3xFLAG
ANKLE2 Reverse #1	GCTCAGGGCGGCAAGCTCAGCCAGG	Gibson Assembly preparation of ANKLE2-APEX2-3xFLAG
ANKLE2 Forward #2	ACTTCCTACCCTCGTAAAGGCCGCCATGCTG	Gibson Assembly of ANKLE2-APEX2-3xFLAG
ANKLE2 Reverse #2	CCGCTACCGCCACCGCTCAGGGCGGCAAG	Gibson Assembly of ANKLE2-APEX2-3xFLAG
ANKLE1 Forward #1	GCCGCCATGTGTTCCGAAGCCC	Gibson Assembly preparation of ANKLE1-APEX2-3xFLAG
ANKLE1 Reverse #1	CGCTTCCACGCGCTTGAATATCT	Gibson Assembly preparation of ANKLE1-APEX2-3xFLAG
ANKLE1 Forward #2	ACTTCCTACCCTCGTAAAGGCCGCCATGTGT	Gibson Assembly of ANKLE1-APEX2-3xFLAG
ANKLE1 Reverse #2	TCCGCTACCGCCACCGCTTCCACGCG	Gibson Assembly of ANKLE1-APEX2-3xFLAG
Mycoplasma Forward #1	TGCACCATCTGTCACTCTGTTAACCTC	For *Mycoplasma sp.* detection
Mycoplasma Reverse #1	GGGAGCAAACAGGATTAGATACCCT	For *Mycoplasma sp.* detection
Mycoplasma Forward #2	GGCGAATGGGTGAGTAACACG	For *Mycoplasma sp.* detection
Mycoplasma Reverse #2	CGGATAACGCTTGCGACCTATG	For *Mycoplasma sp.* detection

References

1. Cao-Lormeau V-M, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, Dub T, Baudouin L, Teissier A, Larre P, Vial A-L, Decam C, Choumet V, Halstead SK, Willison HJ, Musset L, Manuguerra J-C, Despres P, Fournier E, Mallet H-P, Musso D, Fontanet A, Neil J, Ghawché F. 2016. Guillain-Barré Syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. Lancet 387:1531–1539.

2. de Araújo TVB, Rodrigues LC, de Alencar Ximenes RA, de Barros Miranda-Filho D, Montarroyos UR, de Melo APL, Valongueiro S, de Albuquerque M de FPM, Souza WV, Braga C, Filho SPB, Cordeiro MT, Vazquez E, Di Cavalcanti Souza Cruz D, Henriques CMP, Bezerra LCA, da Silva Castanha PM, Dhalia R, Marques-Júnior ETA, Martelli CMT, investigators from the Microcephaly Epidemic Research Group, Brazilian Ministry of Health, Pan American Health Organization, Instituto de Medicina Integral Professor Fernando Figueira, State Health Department of Pernambuco. 2016. Association between Zika virus infection and microcephaly in Brazil, January to May, 2016: preliminary report of a case-control study. Lancet Infect Dis 16:1356–1363.

3. Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Fonseca EB da, Ribeiro EM, Ventura LO, Neto NN, Arena JF, Rasmussen SA. 2017. Characterizing the Pattern of Anomalies in Congenital Zika Syndrome for Pediatric Clinicians. JAMA Pediatr 171:288–295.

4. Campo M del, Feitosa IML, Ribeiro EM, Horovitz DDG, Pessoa ALS, França GVA, García-Alix A, Doriqui MJR, Wanderley HYC, Sanseverino MVT, Neri JICF, Pina-Neto JM, Santos ES, Verçosa I, Cernach MCSP, Medeiros PFV, Kerbage SC, Silva AA, Linden V van der, Martelli CMT, Cordeiro MT, Dhalia R, Vianna FSL, Victora CG, Cavalcanti DP, Schuler-Faccini L. 2017. The phenotypic spectrum of congenital Zika syndrome. American Journal of Medical Genetics Part A 173:841–857.

5. Boonsawat P, Joset P, Steindl K, Oneda B, Gogoll L, Azzarello-Burri S, Sheth F, Datar C, Verma IC, Puri RD, Zollino M, Bachmann-Gagescu R, Niedrist D, Papik M, Figueiro-Silva J, Masood R, Zweier M, Kraemer D, Lincoln S, Rodan L, Passemard S, Drunat S, Verloes A, Horn AHC, Sticht H, Steinfeld R, Plecko B, Latal B, Jenni O, Asadollahi R, Rauch A. 2019. Elucidation of the phenotypic spectrum and genetic landscape in primary and secondary microcephaly. Genetics in Medicine 21:2043–2058.

6. Jayaraman D, Bae B-I, Walsh CA. 2018. The Genetics of Primary Microcephaly. Annu Rev Genomics Hum Genet 19:177–200.

7. Jones KennethL, Smith DavidW, Ulleland ChristyN, Streissguth A. 1973. PATTERN OF MALFORMATION IN OFFSPRING OF CHRONIC ALCOHOLIC MOTHERS. The Lancet 301:1267–1271.

8. Ouellette EM, Rosett HL, Rosman NP, Weiner L. 1977. Adverse effects on offspring of maternal alcohol abuse during pregnancy. N Engl J Med 297:528–530.

9. Van den Eeden SK, Karagas MR, Daling JR, Vaughan TL. 1990. A case-control study of maternal smoking and congenital malformations. Paediatr Perinat Epidemiol 4:147–155.

726   10.   Giles JP, Cooper LZ, Krugman S. 1965. The rubella syndrome. The Journal of Pediatrics
727   66:434–437.

728   11.   Lambert N, Strebel P, Orenstein W, Icenogle J, Poland GA. 2015. Rubella. Lancet
729   385:2297–2307.

730   12.   Messinger CJ, Lipsitch M, Bateman BT, He M, Huybrechts KF, MacDonald S, Mogun H,
731   Mott K, Hernández-Díaz S. 2020. Association Between Congenital Cytomegalovirus and
732   the Prevalence at Birth of Microcephaly in the United States. JAMA Pediatr e203009.

733   13.   McLeod R, Boyer K, Karrison T, Kasza K, Swisher C, Roizen N, Jalbrzikowski J,
734   Remington J, Heydemann P, Noble AG, Mets M, Holfels E, Withers S, Latkany P, Meier
735   P, Toxoplasmosis Study Group. 2006. Outcome of Treatment for Congenital
736   Toxoplasmosis, 1981–2004: The National Collaborative Chicago-Based, Congenital
737   Toxoplasmosis Study. Clinical Infectious Diseases 42:1383–1394.

738   14.   Miner JJ, Cao B, Govero J, Smith AM, Fernandez E, Cabrera OH, Garber C, Noll M,
739   Klein RS, Noguchi KK, Mysorekar IU, Diamond MS. 2016. Zika virus infection during
740   pregnancy in mice causes placental damage and fetal demise. Cell 165:1081–1091.

741   15.   Quicke KM, Bowen JR, Johnson EL, McDonald CE, Ma H, O'Neal JT, Rajakumar A,
742   Wrammert J, Rimawi BH, Pulendran B, Schinazi RF, Chakraborty R, Suthar MS. 2016.
743   Zika Virus Infects Human Placental Macrophages. Cell Host Microbe 20:83–90.

744   16.   Tabata T, Petitt M, Puerta-Guardo H, Michlmayr D, Wang C, Fang-Hoover J, Harris E,
745   Pereira L. 2016. Zika Virus Targets Different Primary Human Placental Cells, Suggesting
746   Two Routes for Vertical Transmission. Cell Host Microbe 20:155–166.

747   17.   Jagger BW, Miner JJ, Cao B, Arora N, Smith AM, Kovacs A, Mysorekar IU, Coyne CB,
748   Diamond MS. 2017. Gestational stage and IFN-λ signaling regulate ZIKV infection in
749   utero. Cell Host Microbe 22:366-376.e3.

750   18.   Yockey LJ, Jurado KA, Arora N, Millet A, Rakib T, Milano KM, Hastings AK, Fikrig E,
751   Kong Y, Horvath TL, Weatherbee S, Kliman HJ, Coyne CB, Iwasaki A. 2018. Type I
752   interferons instigate fetal demise after Zika virus infection. Sci Immunol 3:eaao1680.

753   19.   Alimonti JB, Ribecco-Lutkiewicz M, Sodja C, Jezierski A, Stanimirovic DB, Liu Q,
754   Haqqani AS, Conlan W, Bani-Yaghoub M. 2018. Zika virus crosses an in vitro human
755   blood brain barrier model. Fluids and Barriers of the CNS 15:15.

756   20.   Clé M, Desmetz C, Barthelemy J, Martin M-F, Constant O, Maarifi G, Foulongne V,
757   Bolloré K, Glasson Y, De Bock F, Blaquiere M, Dehouck L, Pirot N, Tuaillon E, Nisole S,
758   Najioullah F, Van de Perre P, Cabié A, Marchi N, Gosselet F, Simonin Y, Salinas S.
759   2020. Zika Virus Infection Promotes Local Inflammation, Cell Adhesion Molecule
760   Upregulation, and Leukocyte Recruitment at the Blood-Brain Barrier. mBio 11:e01183-
761   20.

762   21.   Panganiban AT, Blair RV, Hattler JB, Bohannon DG, Bonaldo MC, Schouest B, Maness
763   NJ, Kim W-K. 2020. A Zika virus primary isolate induces neuroinflammation,
764   compromises the blood-brain barrier and upregulates CXCL12 in adult macaques. Brain
765   Pathol 30:1017–1027.

22. Chiu C-F, Chu L-W, Liao I-C, Simanjuntak Y, Lin Y-L, Juan C-C, Ping Y-H. 2020. The
Mechanism of the Zika Virus Crossing the Placental Barrier and the Blood-Brain Barrier.
Front Microbiol 11:214.

23. Shelton SM, Soucy AR, Kurzion R, Zeldich E, Connor JH, Haydar TF. 2021. Forebrain
Neural Precursor Cells Are Differentially Vulnerable to Zika Virus Infection. eNeuro
8:ENEURO.0108-21.2021.

24. Souza BSF, Sampaio GLA, Pereira CS, Campos GS, Sardi SI, Freitas LAR, Figueira CP,
Paredes BD, Nonaka CKV, Azevedo CM, Rocha VPC, Bandeira AC, Mendez-Otero R,
dos Santos RR, Soares MBP. 2016. Zika virus infection induces mitosis abnormalities
and apoptotic cell death of human neural progenitor cells. Scientific Reports 6:39775.

25. Goodfellow FT, Willard KA, Wu X, Scoville S, Stice SL, Brindley MA. 2018. Strain-
Dependent Consequences of Zika Virus Infection and Differential Impact on Neural
Development. Viruses 10:550.

26. Mazar J, Li Y, Rosado A, Phelan P, Kedarinath K, Parks GD, Alexander KA,
Westmoreland TJ. 2018. Zika virus as an oncolytic treatment of human neuroblastoma
cells requires CD24. PLOS ONE 13:e0200358.

27. Chen W, Foo S-S, Hong E, Wu C, Lee W-S, Lee S-A, Evseenko D, Moreira MEL, García-
Sastre A, Cheng G, Nielsen-Saines K, Brasil P, Avvad-Portari E, Jung JU. 2021. Zika
virus NS3 protease induces bone morphogenetic protein-dependent brain calcification in
human fetuses. Nat Microbiol 6:455–466.

28. Tang H, Hammack C, Ogden SC, Wen Z, Qian X, Li Y, Yao B, Shin J, Zhang F, Lee EM,
Christian KM, Didier RA, Jin P, Song H, Ming G. 2016. Zika Virus Infects Human Cortical
Neural Precursors and Attenuates Their Growth. Cell Stem Cell 18:587–590.

29. Liang Q, Luo Z, Zeng J, Chen W, Foo S-S, Lee S-A, Ge J, Wang S, Goldman SA,
Zlokovic BV, Zhao Z, Jung JU. 2016. Zika Virus NS4A and NS4B Proteins Deregulate
Akt-mTOR Signaling in Human Fetal Neural Stem Cells to Inhibit Neurogenesis and
Induce Autophagy. Cell Stem Cell 19:663–671.

30. Devhare P, Meyer K, Steele R, Ray RB, Ray R. 2017. Zika virus infection dysregulates
human neural stem cell growth and inhibits differentiation into neuroprogenitor cells. 10.
Cell Death & Disease 8:e3106–e3106.

31. Yoon K-J, Song G, Qian X, Pan J, Xu D, Rho H-S, Kim N-S, Habela C, Zheng L, Jacob
F, Zhang F, Lee EM, Huang W-K, Ringeling FR, Vissers C, Li C, Yuan L, Kang K, Kim
S, Yeo J, Cheng Y, Liu S, Wen Z, Qin C-F, Wu Q, Christian KM, Tang H, Jin P, Xu Z,
Qian J, Zhu H, Song H, Ming G-L. 2017. Zika-Virus-Encoded NS2A Disrupts Mammalian
Cortical Neurogenesis by Degrading Adherens Junction Proteins. Cell Stem Cell 21:349-
358.e6.

32. Chavali PL, Stojic L, Meredith LW, Joseph N, Nahorski MS, Sanford TJ, Sweeney TR,
Krishna BA, Hosmillo M, Firth AE, Bayliss R, Marcelis CL, Lindsay S, Goodfellow I,
Woods CG, Gergely F. 2017. Neurodevelopmental protein Musashi-1 interacts with the
Zika genome and promotes viral replication. Science 357:83–88.

33. Lossia OV, Conway MJ, Tree MO, Williams RJ, Goldthorpe SC, Srinageshwar B, Dunbar
GL, Rossignol J. 2018. Zika virus induces astrocyte differentiation in neural stem cells. J
Neurovirol 24:52–61.

34. Ferraris P, Cochet M, Hamel R, Gladwyn-Ng I, Alfano C, Diop F, Garcia D, Talignani L,
Montero-Menei CN, Nougairède A, Yssel H, Nguyen L, Coulpier M, Missé D. 2019. Zika
virus differentially infects human neural progenitor cells according to their state of
differentiation and dysregulates neurogenesis through the Notch pathway. Emerg
Microbes Infect 8:1003–1016.

35. Zeng J, Dong S, Luo Z, Xie X, Fu B, Li P, Liu C, Yang X, Chen Y, Wang X, Liu Z, Wu J,
Yan Y, Wang F, Chen J-F, Zhang J, Long G, Goldman SA, Li S, Zhao Z, Liang Q. 2020.
The Zika Virus Capsid Disrupts Corticogenesis by Suppressing Dicer Activity and miRNA
Biogenesis. Cell Stem Cell 27:618-632.e9.

36. Thulasi Raman SN, Latreille E, Gao J, Zhang W, Wu J, Russell MS, Walrond L, Cyr T,
Lavoie JR, Safronetz D, Cao J, Sauve S, Farnsworth A, Chen W, Shi P-Y, Wang Y, Wang
L, Rosu-Myles M, Li X. Dysregulation of Ephrin receptor and PPAR signaling pathways
in neural progenitor cells infected by Zika virus. Emerg Microbes Infect 9:2046–2060.

37. Yau C, Low JZH, Gan ES, Kwek SS, Cui L, Tan HC, Mok DZL, Chan CYY, Sessions
OM, Watanabe S, Vasudevan SG, Lee YH, Chan KR, Ooi EE. 2021. Dysregulated
metabolism underpins Zika-virus-infection-associated impairment in fetal development.
Cell Reports 37:110118.

38. Shah PS, Link N, Jang GM, Sharp PP, Zhu T, Swaney DL, Johnson JR, Von Dollen J,
Ramage HR, Satkamp L, Newton B, Hüttenhain R, Petit MJ, Baum T, Everitt A, Laufman
O, Tassetto M, Shales M, Stevenson E, Iglesias GN, Shokat L, Tripathi S,
Balasubramaniam V, Webb LG, Aguirre S, Willsey AJ, Garcia-Sastre A, Pollard KS,
Cherry S, Gamarnik AV, Marazzi I, Taunton J, Fernandez-Sesma A, Bellen HJ, Andino
R, Krogan NJ. 2018. Comparative Flavivirus-Host Protein Interaction Mapping Reveals
Mechanisms of Dengue and Zika Virus Pathogenesis. Cell 175:1931-1945.e18.

39. Yamamoto S, Jaiswal M, Charng W-L, Gambin T, Karaca E, Mirzaa G, Wiszniewski W,
Sandoval H, Haelterman NA, Xiong B, Zhang K, Bayat V, David G, Li T, Chen K, Gala
U, Harel T, Pehlivan D, Penney S, Vissers LELM, de Ligt J, Jhangiani SN, Xie Y, Tsang
SH, Parman Y, Sivaci M, Battaloglu E, Muzny D, Wan Y-W, Liu Z, Lin-Moore AT, Clark
RD, Curry CJ, Link N, Schulze KL, Boerwinkle E, Dobyns WB, Allikmets R, Gibbs RA,
Chen R, Lupski JR, Wangler MF, Bellen HJ. 2014. A drosophila genetic resource of
mutants to study mechanisms underlying human genetic diseases. Cell 159:200–214.

40. Link N, Chung H, Jolly A, Withers M, Tepe B, Arenkiel BR, Shah PS, Krogan NJ, Aydin
H, Geckinli BB, Tos T, Isikay S, Tuysuz B, Mochida GH, Thomas AX, Clark RD, Mirzaa
GM, Lupski JR, Bellen HJ. 2019. Mutations in ANKLE2, a ZIKA Virus Target, Disrupt an
Asymmetric Cell Division Pathway in Drosophila Neuroblasts to Cause Microcephaly.
Dev Cell https://doi.org/10.1016/j.devcel.2019.10.009.

41. Shaheen R, Maddirevula S, Ewida N, Alsahli S, Abdel-Salam GMH, Zaki MS, Tala SA,
Alhashem A, Softah A, Al-Owain M, Alazami AM, Abadel B, Patel N, Al-Sheddi T, Alomar
R, Alobeid E, Ibrahim N, Hashem M, Abdulwahab F, Hamad M, Tabarki B, Alwadei AH,
Alhazzani F, Bashiri FA, Kentab A, Şahintürk S, Sherr E, Fregeau B, Sogati S,
Alshahwan SAM, Alkhalifi S, Alhumaidi Z, Temtamy S, Aglan M, Otaify G, Girisha KM,

Tulbah M, Seidahmed MZ, Salih MA, Abouelhoda M, Momin AA, Saffar MA, Partlow JN, Arold ST, Faqeih E, Walsh C, Alkuraya FS. 2019. Genomic and phenotypic delineation of congenital microcephaly. Genet Med 21:545–552.

42. Fishburn AT, Pham OH, Kenaston MW, Beesabathuni NS, Shah PS. 2022. Let's Get Physical: Flavivirus-Host Protein–Protein Interactions in Replication and Pathogenesis. Frontiers in Microbiology 13.

43. Asencio C, Davidson IF, Santarella-Mellwig R, Ly-Hartig TBN, Mall M, Wallenfang MR, Mattaj IW, Gorjánácz M. 2012. Coordination of Kinase and Phosphatase Activities by Lem4 Enables Nuclear Envelope Reassembly during Mitosis. Cell 150:122–135.

44. Snyers L, Erhart R, Laffer S, Pusch O, Weipoltshammer K, Schöfer C. 2018. LEM4/ANKLE-2 deficiency impairs post-mitotic re-localization of BAF, LAP2α and LaminA to the nucleus, causes nuclear envelope instability in telophase and leads to hyperploidy in HeLa cells. Eur J Cell Biol 97:63–74.

45. Roosendaal J, Westaway EG, Khromykh A, Mackenzie JM. 2006. Regulated cleavages at the West Nile virus NS4A-2K-NS4B junctions play a major role in rearranging cytoplasmic membranes and Golgi trafficking of the NS4A protein. J Virol 80:4623–4632.

46. Miller S, Kastner S, Krijnse-Locker J, Bühler S, Bartenschlager R. 2007. The non-structural protein 4A of dengue virus is an integral membrane protein inducing membrane alterations in a 2K-regulated manner. J Biol Chem 282:8873–8882.

47. Teo CSH, Chu JJH. 2014. Cellular vimentin regulates construction of dengue virus replication complexes through interaction with NS4A protein. J Virol 88:1897–1913.

48. He Z, Zhu X, Wen W, Yuan J, Hu Y, Chen J, An S, Dong X, Lin C, Yu J, Wu J, Yang Y, Cai J, Li J, Li M. 2016. Dengue Virus Subverts Host Innate Immunity by Targeting Adaptor Protein MAVS. J Virol 90:7219–7230.

49. Zhang J, Lan Y, Li MY, Lamers MM, Fusade-Boyer M, Klemm E, Thiele C, Ashour J, Sanyal S. 2018. Flaviviruses Exploit the Lipid Droplet Protein AUP1 to Trigger Lipophagy and Drive Virus Production. Cell Host Microbe 23:819-831.e5.

50. Aktepe TE, Liebscher S, Prier JE, Simmons CP, Mackenzie JM. 2017. The Host Protein Reticulon 3.1A Is Utilized by Flaviviruses to Facilitate Membrane Remodelling. Cell Rep 21:1639–1654.

51. Chan JF-W, Yip CC-Y, Tsang JO-L, Tee K-M, Cai J-P, Chik KK-H, Zhu Z, Chan CC-S, Choi GK-Y, Sridhar S, Zhang AJ, Lu G, Chiu K, Lo AC-Y, Tsao S-W, Kok K-H, Jin D-Y, Chan K-H, Yuen K-Y. 2016. Differential cell line susceptibility to the emerging Zika virus: implications for disease pathogenesis, non-vector-borne human transmission and animal reservoirs. Emerg Microbes Infect 5:e93.

52. Vicenti I, Boccuto A, Giannini A, Dragoni F, Saladini F, Zazzi M. 2018. Comparative analysis of different cell systems for Zika virus (ZIKV) propagation and evaluation of anti-ZIKV compounds in vitro. Virus Research 244:64–70.

53. Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, Diamond MS. 2016. A Mouse Model of Zika Virus Pathogenesis. Cell Host & Microbe 19:720–730.

54.   Adams Waldorf KM, Stencel-Baerenwald JE, Kapur RP, Studholme C, Boldenow E,
Vornhagen J, Baldessari A, Dighe MK, Thiel J, Merillat S, Armistead B, Tisoncik-Go J,
Green RR, Davis MA, Dewey EC, Fairgrieve MR, Gatenby JC, Richards T, Garden GA,
Diamond MS, Juul SE, Grant RF, Kuller L, Shaw DWW, Ogle J, Gough GM, Lee W,
English C, Hevner RF, Dobyns WB, Gale M, Rajagopal L. 2016. Fetal brain lesions after
subcutaneous inoculation of Zika virus in a pregnant nonhuman primate. 11. Nature
Medicine 22:1256–1259.

55.   Wu Y, Cui X, Wu N, Song R, Yang W, Zhang W, Fan D, Chen Z, An J. 2017. A unique
case of human Zika virus infection in association with severe liver injury and coagulation
disorders. Sci Rep 7:11393.

56.   Ramos da Silva S, Cheng F, Huang I-C, Jung JU, Gao S-J. 2019. Efficiencies and
kinetics of infection in different cell types/lines by African and Asian strains of Zika virus.
J Med Virol 91:179–189.

57.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie
A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy
E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D,
Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate
protein structure prediction with AlphaFold. 7873. Nature 596:583–589.

58.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe
O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J,
Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt
G, Birney E, Hassabis D, Velankar S. 2022. AlphaFold Protein Structure Database:
massively expanding the structural coverage of protein-sequence space with high-
accuracy models. Nucleic Acids Research 50:D439–D444.

59.   Elkhatib RA, Paci M, Boissier R, Longepied G, Auguste Y, Achard V, Bourgeois P, Levy
N, Branger N, Mitchell MJ, Metzler-Guillemain C. 2017. LEM-domain proteins are lost
during human spermiogenesis but BAF and BAF-L persist. Reproduction 154:387–401.

60.   Lin C, Amberg SM, Chambers TJ, Rice CM. 1993. Cleavage at a novel site in the NS4A
region by the yellow fever virus NS2B-3 proteinase is a prerequisite for processing at the
downstream 4A/4B signalase site. J Virol 67:2327–2335.

61.   Needleman SB, Wunsch CD. 1970. A general method applicable to the search for
similarities in the amino acid sequence of two proteins. Journal of Molecular Biology
48:443–453.

62.   Capra JA, Singh M. 2007. Predicting functionally important residues from sequence
conservation. Bioinformatics 23:1875–1882.

63.   Schneider TD, Stephens RM. 1990. Sequence logos: a new way to display consensus
sequences. Nucleic Acids Res 18:6097–6100.

64.   Crooks GE, Hon G, Chandonia J-M, Brenner SE. 2004. WebLogo: a sequence logo
generator. Genome Res 14:1188–1190.

65. Gao A, Sun T, Ma G, Cao J, Hu Q, Chen L, Wang Y, Wang Q, Sun J, Wu R, Wu Q, Zhou J, Liu L, Hu J, Dong J-T, Zhu Z. 2018. LEM4 confers tamoxifen resistance to breast cancer cells by activating cyclin D-CDK4/6-Rb and ERα pathway. Nat Commun 9:4180.

66. Hein MY, Hubner NC, Poser I, Cox J, Nagaraj N, Toyoda Y, Gak IA, Weisswange I, Mansfeld J, Buchholz F, Hyman AA, Mann M. 2015. A Human Interactome in Three Quantitative Dimensions Organized by Stoichiometries and Abundances. Cell 163:712–723.

67. Gupta GD, Coyaud É, Gonçalves J, Mojarad BA, Liu Y, Wu Q, Gheiratmand L, Comartin D, Tkach JM, Cheung SWT, Bashkurov M, Hasegan M, Knight JD, Lin Z-Y, Schueler M, Hildebrandt F, Moffat J, Gingras A-C, Raught B, Pelletier L. 2015. A dynamic protein interaction landscape of the human centrosome-cilium interface. Cell 163:1484–1499.

68. Huttlin EL, Bruckner RJ, Navarrete-Perea J, Cannon JR, Baltier K, Gebreab F, Gygi MP, Thornock A, Zarraga G, Tam S, Szpyt J, Gassaway BM, Panov A, Parzen H, Fu S, Golbazi A, Maenpaa E, Stricker K, Guha Thakurta S, Zhang T, Rad R, Pan J, Nusinow DP, Paulo JA, Schweppe DK, Vaites LP, Harper JW, Gygi SP. 2021. Dual proteome-scale networks reveal cell-specific remodeling of the human interactome. Cell 184:3022-3040.e28.

69. Go CD, Knight JDR, Rajasekharan A, Rathod B, Hesketh GG, Abe KT, Youn J-Y, Samavarchi-Tehrani P, Zhang H, Zhu LY, Popiel E, Lambert J-P, Coyaud É, Cheung SWT, Rajendran D, Wong CJ, Antonicka H, Pelletier L, Palazzo AF, Shoubridge EA, Raught B, Gingras A-C. 2021. A proximity-dependent biotinylation map of a human cell. Nature 595:120–124.

70. Mosavi LK, Cammett TJ, Desrosiers DC, Peng Z. 2004. The ankyrin repeat as molecular architecture for protein recognition. Protein Sci 13:1435–1448.

71. Mosavi LK, Minor DL, Peng Z-Y. 2002. Consensus-derived structural determinants of the ankyrin repeat motif. Proc Natl Acad Sci U S A 99:16029–16034.

72. Wagner N, Krohne G. 2007. LEM-Domain Proteins: New Insights into Lamin-Interacting Proteins, p. 1–46. *In* International Review of Cytology. Academic Press.

73. Brachner A, Foisner R. 2011. Evolvement of LEM proteins as chromatin tethers at the nuclear periphery. Biochemical Society Transactions 39:1735–1741.

74. Burla R, La Torre M, Maccaroni K, Verni F, Giunta S, Saggio I. 2020. Interplay of the nuclear envelope with chromatin in physiology and pathology. Nucleus 11:205–218.

75. Petit MJ, Kenaston MW, Pham OH, Nagainis AA, Fishburn AT, Shah PS. 2021. Nuclear dengue virus NS5 antagonizes expression of PAF1-dependent immune response genes. PLoS Pathog 17:e1010100.

76. Yu JH, Schaffer DV. 2006. Selection of novel vesicular stomatitis virus glycoprotein variants from a peptide insertion library for enhanced purification of retroviral and lentiviral vectors. J Virol 80:3285–3292.

77. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25:402–408.

970   78.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
971   Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
972   Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
973   analysis. 7. Nat Methods 9:676–682.

974   79.   Baghirova S, Hughes BG, Hendzel MJ, Schulz R. 2015. Sequential fractionation and
975   isolation of subcellular proteins from tissue or cultured cells. MethodsX 2:440–445.

976   80.   Hirokawa T, Boon-Chieng S, Mitaku S. 1998. SOSUI: classification and secondary
977   structure prediction system for membrane proteins. Bioinformatics 14:378–379.

978   81.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane
979   protein topology with a hidden Markov model: application to complete genomes. J Mol
980   Biol 305:567–580.