Targeted Anatomical and Functional Identification of Antinociceptive and Pronociceptive Serotonergic Neurons that Project to the Spinal Dorsal Horn

Abbreviated title: Descending Serotonergic Projection Neurons

Robert P. Ganley[1], Marília Magalhaes de Sousa[1], Kira Werder[1], Tugce Öztürk[1], Raquel Mendes[1], Matteo Ranucci[1], Hendrik Wildner[1] & Hanns Ulrich Zeilhofer[1,2]

[1]Institute for Pharmacology and Toxicology, University of Zürich, CH-8057 Zürich, Switzerland

[2]Institute of Pharmaceutical Sciences, Swiss Federal Institute of Technology Zürich, CH-8090, Zürich, Switzerland

Corresponding author: zeilhofer@pharma.uzh.ch

Pages: 42

Figures: 6

Figure supplements: 13

Tables: 8

Number of words
    Abstract: 202
    Introduction: 601
    Discussion: 1910

Conflict of interest: none

Acknowledgements: We are grateful for funding provided by grants from the Swiss National Science Foundation (grant number 310030_197888) and by the clinical research priority programme (CRPP) "Pain – from phenotypes to mechanisms" of the Faculty of Medicine, University of Zurich, to HUZ, and a grant from the Olga Mayenfisch Stiftung to HW. We thank Louis Scheurer, Katharina Struckmeyer-Fichtel, and Isabelle Kellenberger for technical support.

Abstract

Spinally-projecting serotonergic neurons play a key role in controlling pain sensitivity and can either increase or decrease nociception depending on physiological context. It is currently unknown how serotonergic neurons mediate these opposing effects. Utilizing virus-based strategies, we identified two anatomically separated populations of serotonergic hindbrain neurons located in the lateral paragigantocellularis (LPGi) and the medial hindbrain, which respectively innervate the superficial and deep spinal dorsal horn and have contrasting effects on sensory perception. Our tracing experiments revealed that serotonergic neurons of the LPGi were much more susceptible to transduction with spinally injected AAV2retro vectors than medial hindbrain serotonergic neurons. Taking advantage of this difference, we employed intersectional chemogenetic approaches to demonstrate that activation of the LPGi serotonergic projections decreases thermal sensitivity, whereas activation of medial serotonergic neurons increases sensitivity to mechanical von Frey stimulation. Together these results suggest that there are functionally distinct classes of serotonergic hindbrain neurons that differ in their anatomical location in the hindbrain, their postsynaptic targets in the spinal cord, and their impact on nociceptive sensitivity. The LPGi neurons that give rise to rather global and bilateral projections throughout the rostrocaudal extent of the spinal cord appear to be ideally posed to contribute to widespread systemic pain control.

1 Introduction

Descending pain control is a critical endogenous mechanism of pain modulation that is

required for survival [47], and allows an organism to respond in an appropriate context-

dependent manner to external threats. Key brain structures involved in this system were

discovered through stimulation-produced analgesia experiments and include the

periaqueductal grey matter (PAG) and the rostroventromedial medulla (RVM) of the

hindbrain [9]. This descending system acts to inhibit the flow of nociceptive information

through the spinal dorsal horn and requires descending tracts within the dorsolateral

funiculus to mediate its effects [8; 10]. The RVM is also required for maintaining some

chronic pain states through a process of descending facilitation [54; 74], indicating that this

area has a bidirectional control over pain sensitivity. Both descending inhibition and

facilitation will ultimately require the activation of descending tracts to modulate the neuronal

activity within the dorsal horn. To understand the precise neuronal circuitry and transmitter

systems underlying descending pain inhibition and facilitation, an understanding of the

different projection neurons from the RVM to the spinal cord is needed.

Monoamines are important neurotransmitters involved in descending pain control, with

serotonin (5-HT) being able to inhibit nociception when injected intrathecally [64]. Spinal 5-

HT is also known to contribute to certain forms of endogenous pain suppression, such as

stress-induced analgesia [72], but is also required to maintain chronic pain in rodent models

of nerve injury [66]. This pain facilitation in pre-clinical neuropathy models is thought to be a

result of reduced tonic diffuse noxious inhibitory controls, due to increased spinal 5-HT$_3$

receptor activation [4; 6]. The variety and abundance of 5-HT receptors within the spinal

dorsal horn likely explains, at least in part, why these multiple effects are observed [7].

Clearly, spinal 5-HT and the serotonergic projections to this region have a complex and

bidirectional control over pain perception that require further study.

Functional studies of specific descending projections have been facilitated by the development of viral retrograde tracers, such as AAV2retro and canine adenoviruses [25; 35; 60]. However, it has been reported that the AAV2retro serotype, may not infect serotonergic hindbrain neurons as efficiently as other descending projections [60; 65]. Further, despite the dense serotonergic innervation of the spinal dorsal horn, studies employing transsynaptic neuronal circuit tracing from dorsal horn neurons with modified rabies viruses rarely found labeled hindbrain serotonergic neurons [25; 43]. This is surprising since both descending serotonergic neurons and dorsal horn interneurons are known to strongly influence nociception, and it is possible that these rabies virus-based tracers fail to detect functional connections between serotonergic neurons and starter populations within the spinal cord. Together these reports would suggest that descending serotonergic projections of the hindbrain are particularly challenging to study using the currently available tools.

In an attempt to address this discrepancy, we compared the labeling efficiency of viral and non-viral tracers. Specifically, we examined the susceptibility of serotonergic hindbrain neurons to retrograde spinal transduction by AAV2retro serotype vectors [60]. Additionally, we traced serotonergic neurons of the hindbrain with modified rabies viruses and tested whether they could be labelled from dorsal horn neuron starter populations with transsynaptic rabies tracing. To allow functional interrogation of descending serotonergic neurons, we assessed the specificity of the TPH2::Cre mouse line and used this together with the preferential transduction efficiency of AAV2retro to develop an intersectional system that allowed selective labelling and manipulation of serotonergic neurons in the lateral paragigantocellularis (LPGi), or the medial serotonergic neurons including the nucleus raphe magnus (NRM). We find that the lateral and medial serotonergic neurons of the RVM are distinct in terms of their anatomical organization, susceptibility to AAV2retro transduction, and influence on acute nociception.

Results

Spinal injection of AAV2retro serotype vectors preferentially transduce TPH2-containing

projection neurons in the lateral hindbrain

AAV vectors have been developed to permit the efficient retrograde labelling of projection neurons via their axon terminals [60]. However, it is debatable whether these vectors are capable of transducing serotonergic neurons that project to the spinal cord [65]. To test this, we used AAV2retro serotype vectors to retrogradely label hindbrain neurons that project to the spinal cord to assess their ability to transduce serotonergic projection neurons (Figure 1A). Following intraspinal injection of AAV2retro.GFP, many transduced neurons were labelled within the RVM (Figure 1B). When we inspected the location of neurons that expressed both TPH2 and eGFP (according to scheme depicted in Figure 1- figure supplement 1), we found that the majority of these cells were located within the lateral paragigantocellularis (LPGi) (Figure 1B). Quantification of the location of the eGFP-labelled TPH2-expressing neurons indicated that 74% were in the LPGi both ipsilateral and contralateral to the spinal cord injection site (left hand side). In agreement with [65], far fewer eGFP+TPH2+ neurons were found within the midline serotonergic nuclei (23/100 neurons in NRM, (n = 3 animals) (Figure 1C). However, most non-serotonergic projection neurons (eGFP+ TPH2-) were found close to the midline of the ventral hindbrain (412/439 neurons in midline, (n = 3 animals) (see Figure 1B and Figure 1E)

There is increasing evidence to suggest that some projections are resistant to retrograde transduction with AAV2retro vectors, and comparisons with independent tracing methods are required to demonstrate this resistance [26; 60]. To study the anatomical organization of serotonergic pathways using a non-viral method, we used Cholera toxin b subunit (CTb) retrograde tracing and compared the hindbrain labelling with our AAV2retro tracing experiments (Figure 1- figure supplement 2A).

Strikingly, many CTb-labeled TPH2-expressing hindbrain neurons were found in the midline

NRM, although CTb-containing TPH2-expressing neurons were also found in the LPGi

(Figure 1 - figure supplement 2B). The location of these CTb+TPH2+ cells was more evenly

divided between the NRM and LPGi that the eGFP+TPH2+ cells from the AAV2retro labelling

(Figure 1 - figure supplement 2C compared to Figure 1C). To directly test whether there were

differences in labelling efficiency between AAV2retro and CTb, AAV2retro.GFP and CTb

were co-injected into the spinal dorsal horn (Figure 1D). In total, more neurons in the ventral

hindbrain were retrogradely labelled with CTb than AAV2retro.eGFP (Figure 1 – figure

supplement 3A and table 1). When only TPH2-expressing retrogradely labelled neurons were

examined, far fewer were labelled with AAV2retro.GFP relative to the overall population of

traced neurons (Figure 1 – figure supplement 3 and table 1). However, when the retrogradely

labelled TPH2+ neurons were divided into separate areas, it was apparent that a lower

percentage of neurons in the medial serotonergic nuclei including the NRM were labelled

with eGFP compared to the LPGi, whereas the proportion of serotonergic neurons in the

LPGi labelled with AAV2retro, CTb, and both AAV2retro and CTb were similar to the general

population of retrogradely labelled hindbrain neurons (Figure 1 - figure supplement 3A and

B). Therefore, we conclude that the reduced labelling of hindbrain serotonergic neurons with

AAV2retro was due to a resistance of midline serotonergic neurons to AAV2retro-mediated

transduction. In contrast, laterally located serotonergic neurons were more amenable to

retrograde transduction with AAV2retro. A summary of these data can be found in table 1.

Serotonergic hindbrain neurons are rarely infected using transsynaptic rabies tracing from

the dorsal horn

Since many serotonergic neurons were largely resistant to AAV2retro transduction, we

considered that these may be resistant to other viruses used for retrograde and neuronal

circuit tracing. Transsynaptic rabies tracing is a commonly used technique for tracing

neuronal circuits from genetically defined populations of neurons and provides useful

information regarding the presynaptic inputs to a neuronal population. [69]. However, in the

case of serotonergic neurons, the results of previous studies have demonstrated that very

few neurons of the hindbrain are traced from dorsal horn starter populations, despite the

dense serotonergic innervation of the superficial dorsal horn and the important role of

serotonin in pain modulation [25; 43]. This suggests that serotonergic neurons are either

traced with only limited efficacy, or that there is highly specific serotonergic innervation of

some spinal neuron populations but not others.

To test whether serotonergic hindbrain neurons are susceptible to rabies virus infection, we

injected a rabies virus, whose genome lacked the glycoprotein-encoding sequence but was

pseudotyped with the SAD glycoprotein (SAD.RabiesΔG-eGFP (SAD-G)), into the spinal

dorsal horn (Figure 1 - figure supplement 4A). This virus can directly infect most neurons but

cannot be propagated beyond the initially infected neurons due to the lack of the rabies

glycoprotein required for transsynaptic spread [2; 68]. Five days after injection of

SAD.rabΔG-eGFP (SAD-G), 44% of the labelled hindbrain neurons contained detectable

TPH2 immunoreactivity (Figure 1 – figure supplement 4B and C). Notably, the percentage of

labeled neurons dropped to only 12% of labelled hindbrain neurons at day 7 post injection

(Figure 1 – figure supplement 4C). This reduction may be a result of rabies virus toxicity,

which could lead to a downregulation of cytoplasmic enzymes such as TPH2 to undetectable

levels, and hence an underestimation of labelled serotonergic neurons. For this reason, we

used a 5-day survival time for all subsequent experiments involving rabies viruses.

To determine whether serotonergic hindbrain neurons could be traced transsynaptically from

the spinal dorsal horn, we used Hoxb8-Cre mice to define a broad starter population of spinal

cord neurons. During development, Hoxb8-Cre is expressed transiently in almost all spinal

neurons and astrocytes caudal to C4. It can be used to define a starter population that

includes virtually all dorsal horn neurons [70]. We used Hoxb8-Cre; ROSA^{TVA} mice to induce

stable expression of TVA in most dorsal horn neurons during development, enabling their

infection in the adult with EnvA pseudotyped rabies viruses (SAD.Rabies∆G-GFP (EnvA)).

To allow transsynaptic spread from the starter population, a helper virus containing mCherry

and rabies glycoprotein was injected two weeks prior to injection of SAD.Rabies∆G-GFP

(EnvA) (Figure 1– figure supplement 4D).

Five days after rabies virus injection, neurons in the ventral hindbrain labelled with eGFP

were assessed for TPH2 expression by immunostaining. On average, the number of eGFP-

labelled hindbrain neurons counted per animal was similar between direct and transsynaptic

tracing from the spinal cord (8.3 and 8.25 respectively), both of which had far lower labelling

efficiency when compared to labelling with either CTb or AAV2retro labelling. We only

observed coexpression of eGFP and TPH2 once in a total of 33 inspected neurons,

corresponding to 3% of our total sample (Figure 1 – figure supplement 4E and F). A

summary of all rabies virus tracing experiments can be found in table 2. In the hindbrain, we

never observed colocalization of mCherry and eGFP, suggesting that all eGFP-expressing

neurons in the hindbrain were transsynaptically traced from the dorsal horn. Together these

data demonstrate that although TPH2-expressing hindbrain neurons can be directly infected

with modified rabies viruses, they are largely underrepresented with transsynaptic tracing

from the dorsal horn. This may partly explain their absence from many of the circuit tracing

studies that have used this approach.

Descending serotonergic projection neurons of the LPGi can be preferentially labelled using

AAV2retro vectors and TPH2::Cre mouse line

In order to study descending serotonergic neurons functionally, we must first identify a

suitable method to specifically influence those neurons. TPH2-containing neurons of the

LPGi were susceptible to retrograde transduction by AAV2retro in contrast to other

serotonergic nuclei of the hindbrain (Figure 2), which would permit selective transduction of

these neurons using AAV2retro. Further, mouse lines expressing Cre recombinase within defined neuronal populations have been widely and successfully used to functionally study a variety of spinal neuron subtypes [24; 29; 36; 51]. We therefore assessed the reliability of the TPH2::Cre mouse line as a marker of serotonergic neurons and a potential tool to gain genetic access to these cells. To determine whether the Cre expression in this mouse line was specific to serotonergic neurons that project to the spinal cord, we injected the lumbar dorsal horn of TPH2::Cre mice with an AAV2retro serotype vector containing a Cre-dependent eGFP construct (Figure 2A). This resulted in TPH2-expressing neurons of the hindbrain being labelled with eGFP, particularly in the LPGi (Figure 2B). In addition, many neurons were labelled in the ipsilateral DRG (Figure 2C). We observed a similar pattern in cell location as in the AAV2retro.eGFP labelling experiments (Figure 1C), with 72% of eGFP neurons being present bilaterally in the LPGi (Figure 2D and table 3). Furthermore, 77.5% of all neurons labelled with AAV2retro.flex.eGFP contained detectable levels of TPH2 (Figure 2E and table 3). Therefore, among the hindbrain neurons that project to the spinal dorsal horn TPH2::Cre is expressed specifically in serotonergic neurons, but may also be expressed in other neurons of the nervous system, including a subset of DRG neurons.

To specifically manipulate spinally-projecting serotonergic neurons without influencing other neuronal populations, such as those in the DRG, we used an intersectional strategy to induce reporter expression in spinally-projecting TPH2::Cre neurons of the hindbrain (Figure 2F). This strategy utilized AAV2retro serotype vectors containing a Cre-dependent optimized flippase (FLPo), which, when injected intraspinally, results in FLPo expression in Cre-expressing neurons that project to the dorsal horn. The AAV2retro serotype would also enable expression restricted to laterally located serotonin-expressing neurons. This construct also contained the coding sequence for mCherry, allowing these transduced cells to be visualized (AAV2retro.flex.FLPo.mCherry). The hindbrain was injected one week later with an AAV containing a flippase-dependent eGFP viral vector (AAV9.FRT.eGFP), to capture the neurons that were transduced from the spinal cord with AAV2retro (Figure 2F and G). This

approach enables the quantification of intersectionally labelled serotonergic neurons (mCherry- and eGFP-expressing), relative to the population of neurons directly labelled from the spinal dorsal horn (mCherry-expressing). Therefore, the accuracy of the hindbrain injections and the proportion of the descending projection neurons captured within the injection site could be determined.

Different brain injection coordinates were tested to optimize the proportion of spinally-projecting TPH2::Cre neurons that were captured with the stereotaxic brain injection. Bilateral injections of AAV9.FRT.eGFP (-6, ±0.5, -5.9 relative to Bregma) was the most efficient in terms of mCherry+eGFP+ neurons labelled in the hindbrain (Figure 2F, G, and H). With this approach, we labelled 80.6% of all retrogradely transduced neurons with eGFP with the remaining 19.4% containing mCherry only, indicating that most retrogradely transduced neurons are labelled using this strategy (Figure 2H and table 4). Additionally, this proportion was similar between the different areas containing retrogradely-labelled serotonergic neurons (LPGi = 80.6%, Medial = 78.3%, other = 82.4%, labelled neurons containing eGFP), suggesting these injection sites encompass most regions that contain AAV2retro-labelled serotonergic projection neurons. Some cells only expressed eGFP without detectable mCherry staining, either due to non-specific recombination of the FRT sites or low to undetectable levels of mCherry present in these neurons. We found that a smaller proportion of eGFP-only neurons were present in the LPGi compared to the medial serotonergic nuclei (4.6% versus 16.9% respectively, Figure 2H and table 4). We concluded that this intersectional strategy is suitable for the specific manipulation of serotonergic hindbrain neurons that project to the spinal dorsal horn, and that the majority of these traced cells are located in the LPGi.

Lateral and midline serotonergic hindbrain neurons display anatomical differences in their spinal cord innervation

Previous anterograde tracing studies have suggested that within the RVM, the innervation of the superficial dorsal horn originates from the lateral hindbrain, whereas the deeper dorsal horn is innervated by the medial hindbrain [30]. The preferential labelling of lateral serotonergic neurons using our intersectional strategy provided us an opportunity to compare the projection patterns of different serotonergic neurons of the hindbrain.

To assess the serotonergic innervation of the spinal cord that originates from the medial RVM we injected AAVs containing Cre-dependent reporters into the NRM of TPH2::Cre mice (Coordinates from Bregma = -6, 0, 5.9) Figure 3A) and were able to label midline serotonergic neurons with eGFP without transduction of the LPGi neurons (Figure 3B). In addition to the NRM, neighboring midline regions were also labelled, such as the raphe obscuris (ROb) and raphe pallidus (RPa) [19]. Most eGFP-labelled neurons in the injection site contained a detectable level of TPH2 (80.5%), indicating this strategy can be used to label the serotonergic midline neurons of the hindbrain (Figure 3C).

We used tdTomato-expressing vectors to visualize the axon termination pattern of descending serotonergic projections in the spinal cord (Figure 3D and F). For preferential labelling of the LPGi descending neurons, the lumbar spinal cord was injected with AAV2retro vectors containing a Cre-dependent optimized Flippase (AAV2retro.flex.FLPo.BFP), and one week later the hindbrain was injected with AAVs containing flippase-dependent tdTomato (AAV9.FRT.tdTOM). The stronger fluorescence of tdTomato enabled more sensitive labelling of the distal axons of descending projection neurons. The intersectional approach revealed that the densest labelling of axon terminals was present in the superficial laminae of the dorsal horn (Figure 3D, E, and H). When the spinal cords were immunostained for CGRP and PKCγ (to delineate laminae I-IIo and IIi boundaries respectively), most of the labeled axons were seen within the dorsal CGRP plexus, indicating that these neurons innervate laminae I-IIo of the spinal cord (figure 3 - figure supplement 1A). Surprisingly, projections retrogradely labelled from the left lumbar

spinal cord contained axon collaterals that projected to both ipsilateral and contralateral

dorsal horns, as well as to spinal segments caudal to the spinal injection site in the lumbar

segment (Figure 3E and H). This indicates that these descending neurons have a wide-

ranging axon termination pattern that extends over most of the spinal cord.

In contrast, axons from midline serotonergic neurons (including NRM, ROb, and RPa)

directly labelled with tdTomato (from injection of AAVs containing a Cre-dependent tdTOM

(AAV8.flex.tdTOM) into the NRM (-6, 0, 5.9)) were rarely present in the superficial laminae of

the dorsal horn, but they densely innervated the ventral horn and were present in the deep

dorsal horn (Figure 3G and I). This was confirmed by revealing the laminar boundaries with

CGRP and PKCγ staining, which indicated that the axons of these neurons was mostly

present ventral to the lamina IIi-III border (figure 3 - figure supplement 1B) In both labeling

experiments, most of the axon terminals in the spinal cord contained 5-HT, although much of

the spinal 5-HT was not colocalized with tdTomato by either labeling approach (Figure 3H

and I). Together this indicates that the lateral and midline serotonergic neurons in the RVM

project to distinct regions of the spinal cord, in agreement with Gautier *et al* [30].

Midline and lateral serotonergic hindbrain neurons have similar active and passive

membrane properties

Although much is known about the electrophysiological properties of serotonergic hindbrain

neurons, to date there has not been a direct comparison of the membrane properties of

those located in the NRM or LPGi. To test whether these distinct populations have similar

biophysical membrane properties, TPH2::Cre neurons were labelled with tdTomato by

hindbrain injection of AAV9.flex.tdTom and targeted for whole-cell recording.

In contrast to their anatomical differences (such as cell soma location and spinal axon

termination pattern), TPH2::Cre neurons in the LPGi and NRM had similar active membrane

properties and neuronal excitability (Figure 4A and B). In response to depolarizing current

steps, these two groups of neurons discharged action potentials that were similar in terms of

their thresholds, kinetics, and firing rates (Figure 4A, B, D-G). Cells were labelled with

Biocytin during recording, and these groups were distinguished based on the location of

biocytin-labelled cell bodies in the hindbrain slice after recording and tissue processing

(Figure 4C). Similar to previous reports on the electrophysiological properties of serotonergic

RVM neurons, both groups generally had broad action potentials (measured as width at half

maximal) and frequently exhibited a slow afterhyperpolarization (9/10 NRM neurons, 5/8

LPGi neurons), which was similar in amplitude between groups (Figure 4A, E, and H) [73]. A

summary of these electrophysiological measurements can be found in table 5.

Acute activation of descending serotonergic neurons in the LPGi decreases thermal sensitivity

Descending serotonergic projection neurons can both decrease or increase pain sensitivity,

and optogenetic activation of serotonergic neurons of the nucleus raphe magnus produces

long-term hypersensitivity to both mechanical and thermal sensitivity [14; 66]. However,

intrathecal injection of serotonin can produce acute antinociception [64], and currently little is

known of the effect of preferentially activating descending serotonergic neurons in the LPGi.

To clarify whether the acute activation of these serotonergic neurons is pro- or

antinociceptive, we used the intersectional strategy to express the excitatory DREADD

hM3D(q) in descending serotonergic neurons of the LPGi (Figure 5A). Although many

neurons of the hindbrain were labelled with hM3Dq-mCherry and virtually all mCherry-

expressing axon terminals in the lumbar spinal cord contained detectable 5-HT, this only

corresponded to a minority of the serotonergic innervation in this area (Figure 5B and C).

After injecting CNO hM3Dq-mCherry-expressing neurons increased their activity, as

evidenced by an increased expression of c-Fos in their nuclei relative to vehicle injected

controls (Figure 5 - figure supplement 1)

Despite sparse terminal labelling in the spinal cord, withdrawal latencies to thermal stimuli

were increased following CNO injection (Figure 5D). Animals that had received CNO

injections exhibited significantly longer withdrawal thresholds to infrared heat stimulation than

vehicle injected controls (Repeated measures one-way ANOVA, $F_{(3, 9)}$ = 10.55, $P <$

0.0001). Similarly, response latencies to the cold plantar assay were also prolonged following

CNO injection (Repeated measures one-way ANOVA, $F_{(3, 9)}$ = 7.309, P = 0.0012). In

contrast, tactile sensitivity to punctate mechanical stimulation with von Frey filaments was

unaltered (Repeated measures one-way ANOVA, $F_{(3, 9)}$ = 1.402, P = 0.2666) (Figure 6E).

To exclude the possibility that any observed alteration in sensory stimulus-induced behavior

was due to impaired motor control or sedation, we measured the latency to fall from an

accelerating rotarod before and after injection of CNO or vehicle. Latency to fall was

comparable between animals before and after injection indicating no deficits in sensorimotor

function or the absence of sedative effects (repeated measures one-way ANOVA, $F_{(3, 5)}$ =

0.354, P = 0.787)(Figure 5D) Together, these data indicate that acute activation of

descending serotonergic neurons reduces sensitivity to thermal stimuli, but does not strongly

influence mechanical sensitivity.

The presence of axon collaterals on the contralateral side of the spinal cord raises the

possibility that these could also mediate a similar effect to those on the ipsilateral side.

Therefore, we also tested the contralateral paw using the same sensory tests (Figure 5 –

figure supplement 2). We saw a significant increase in withdrawal latencies to heat

stimulation in the Hargreaves assay (Repeated measures one-way ANOVA, $F_{(3, 14)}$ =

7.757, P = 0.0003)(Figure 5 - figure supplement 3), suggesting that the activation of these

neurons has some influence on the sensitivity of the contralateral paw, presumably via axon

collaterals present in the contralateral dorsal horn.

Sex differences are often reported in the context of sensory neurobiology, and serotonin

specifically has been implicated in mediating such differences [13; 39; 48]. To see if similar

differences are observed in our experiment, we divided the group into male and female

animals to determine whether both sexes responded in a similar manner. We find that both

male and female mice show reduced sensitivity to thermal stimuli, and both show unaltered

mechanical sensitivity (Figure 5 - figure supplement 3).

In addition, CNO has been reported to mediate effects independently of hM3Dq expression

[31]. To exclude that the effects seen in our assays were caused by such off-target effects,

we performed the same sensory and motor tests on animals that did not express hM3Dq. In

our hands, the injection of CNO did not produce any noticeable change in any of the

behavioral tests in the absence of hM3Dq expression (Figure 5 - figure supplement 4).

When we examined the hindbrain tissues from animals used in the LPGi activation

experiments, we quantified the proportion of retrogradely labelled and intersectionally

labelled neurons. On average, 67% of BFP-labelled neurons were also labeled with hM3Dq-

mCherry, and very few (1.85%) only expressed mCherry (figure 5 - figure supplement 5).

This shows that the intersectional strategy employed for chemogenetic activation of

descending serotonergic neurons is similarly specific as our previous labelling experiments

(figure 2F, G, and H)

Acute activation of serotonergic neurons in the medial RVM increases mechanical sensitivity

Activation of serotonergic neurons has also been associated with increased sensitivity to

tactile and thermal stimuli and is thought to underly certain forms of neuropathic pain [14;

59]. The differences between our findings and those of Cai *et al* [14] could be due to either of

the mode/strength of neuronal activation (chemogenetic vs optogenetic) or the anatomical

location of the activated neurons (medial hindbrain vs lateral hindbrain). To test this directly,

we used the same chemogenetic receptor as our previous experiments to activate midline

serotonergic neurons (Figure 6A). Using the same strategy as the anatomical tracing

experiment (Figure 3A), we injected AAVs containing Cre-dependent hM3Dq.mCherry into

the NRM. Similarly, we were able to limit transgene expression to the medial serotonergic

neurons including the NRM, the ROb and RPa, but without spread to the LPGi (Figure 6B

and figure 6 – figure supplement 1). This resulted in the presence of mCherry-labelled

terminals in the deep dorsal horn of the spinal cord, with limited expression in the dense

plexus of 5-HT terminals in the superficial spinal laminae (Figure 6C).

In contrast to the activation of serotonergic neurons within the LPGi, the activation of medial

RVM neurons had no effect on thermal withdrawal latencies (Repeated measures one-way

ANOVA, $F_{(3, 6)} = 0.531$, $P = 0.667$) or responses to cooling (Repeated measures one-way

ANOVA, $F_{(3, 6)} = 3.071$, $P = 0.054$) (Figure 6D). However, we observed a significant

decrease in the withdrawal thresholds to punctate mechanical stimuli in the hindpaw

(Repeated measures one-way ANOVA, $F_{(3, 6)} = 13.84$, $P = 0.787$)(Figure 6D). Again,

neither CNO nor vehicle injection impaired sensorimotor coordination or induced sedation,

using latency to fall from an accelerating rotarod as a readout measure (Repeated measures

one-way ANOVA, $F_{(3, 6)} = 1.791$, $P = 0.185$) (Figure 6D).These data indicate that, in

contrast to serotonergic neurons of the LPGi, activating medial serotonergic hindbrain

neurons increases mechanical sensitivity without influencing responses to thermal stimuli.

To test for potential sex differences in mechanical hypersensitivity, we again divided our

group into males and females and compared the withdrawal thresholds/latencies post-vehicle

injection to post-CNO injection. We see that both male and female animals exhibit similar

increases in mechanical sensitivity after CNO injection (figure 6 - figure supplement 2),

suggesting that this effect is not sex specific. The withdrawal latencies to heat and cold were

unchanged and the latency to fall from the accelerating rotarod were similarly unchanged in

both male and female animals.

Additionally, when we inspected the injection sites in the brain stem from animals used in

these behavioral experiments, we found that a similar proportion of neurons expressed TPH2

(85.1%, figure 6 - figure supplement 3) as compared to our previous labelling experiments

(figure 3C). Therefore, we concluded that the chemogenetic activation is largely restricted to

serotonergic neurons of the medial hindbrain in these experiments.

Discussion

Anatomical organization of descending serotonergic innervation of the lumbar spinal cord

The anatomical locations of serotonergic nuclei in the brainstem have been known since the

early studies of Dahlstrom and Fuxe [19]. These areas have later been shown to contain

spinally-projecting neurons and together provide the spinal cord with serotonin [41]. Similarly,

we were also able to label the same regions from the spinal cord with CTb tracing (Figure 1 -

figure supplement 2). However, we found that most of the descending serotonergic neurons

transducible with AAV2retro serotype vectors from the mouse spinal cord are located in the

LPGi, similar to what has been previously demonstrated [65]. We used this selectivity to

devise a strategy to preferentially label midline or lateral serotonergic neurons and

demonstrate that, in agreement with anterograde tracing studies, the innervation of the

superficial laminae of the dorsal horn mainly originates from the LPGi and not the NRM [30].

Unlike Gautier et al [30], we found many axon terminals present within the ventral horn as

well as the deep dorsal horn. Potentially this reflects a species difference (rat versus mouse),

or the labelling of other midline serotonergic regions beyond the NRM, such as the ROb or

the RPa in the present study [34; 49]. However, inclusion of these midline serotonergic

neurons within the injection site demonstrates that they do not project to the superficial

dorsal laminae. Therefore, it is likely that these pathways influence anatomically distinct

neuron populations of the spinal cord, and likely play different functional roles.

The spinal dorsal horn is organized in a laminar structure that is closely linked to its function,

such that thermal and nociceptive information is generally received and processed in the

superficial laminae, whereas non-nociceptive mechanical and proprioceptive information is

processed in deeper spinal laminae [61]. This laminar organization can be seen in terms of

gene expression [33], the termination zones of different primary sensory neurons [17; 42],

and the restricted laminar location of neuron populations [20]. Therefore, the laminar

location of serotonin release will mostly influence different populations of neurons within

these regions, which play diverse functional roles [3; 23; 24; 29].

Antinociceptive function of descending LPGi serotonergic neurons in spinal circuits

Acute activation of spinally-projecting serotonergic neurons using excitatory DREADDs was

able to reduce sensitivity to thermal tests (Figure 5D). In support of this idea, the axons of

these projections are densest in the superficial laminae of the spinal cord (Figure 3E and H),

where temperature-related information is conveyed by TRPV1- and TRPM8-expressing

sensory neurons [16; 17; 22]. The presence of axons from lateral serotonergic neurons within

the region that receives, and processes thermal information is consistent with the

observation that activating these neurons affects thermal sensitivity. Furthermore, 5-HT can

inhibit C fiber-mediated input to the rat superficial dorsal horn and hyperpolarize many types

of excitatory neurons in this area, in agreement with a reduction in neuronal activity that

would inhibit the relay of temperature-related information [44].

Pronociceptive function of medial serotonergic neurons in spinal circuits

Preclinical models of chronic pain can increase the activity of serotonergic hindbrain

neurons, and TPH2 knockdown experiments can transiently reduce spinal 5-HT and

hypersensitivity in neuropathic animals [59; 66]. In these neuropathic models, several

changes in the physiology of the dorsal horn are widely reported [18; 62; 67]. Together, these

pathological changes may alter the way that 5-HT influences the processing of sensory

information within the dorsal horn. In agreement, the spinal serotonin receptors engaged by

endogenous pain control systems such as diffuse noxious inhibitory controls are altered

during the development of chronic pain [4-6]. Additionally, alterations could also occur within

the serotonergic projection neurons, such as their activity and neurotransmitter

content/concentration [63].

However, there are also data that point to a pro-nociceptive role for 5-HT at the spinal level in naïve animals. For example, it was shown that optogenetic stimulation of the NRM of TPH2-ChR2 animals could induce hypersensitivity that lasted several weeks [14]. In our chemogenetic experiments, the activation of NRM neurons with CNO did produce an increase in mechanical sensitivity, but unlike Cai *et al* [14] we did not see an alteration in thermal sensitivity or an effect that outlasted the neuron stimulation. These differences could be explained in part by the intensity and location of the stimulation used in experiments. In support of this idea, activation of the same neuron population with optogenetic or chemogenetic tools can produce distinct behaviors, likely due to differences in the strength or pattern of the neuronal activation [58].

The activation of the midline serotonergic neurons influences multiple functionally different groups of neurons within the injection site, including the NRM, ROb, and RPa [49]. These more caudal nuclei are known to affect the motor system and respiratory functions [12; 34; 37]. Within the sensory assays tested, there was a rather selective change in mechanical responses and unaltered thermal responses, indicating that the effect was modality selective and unlikely due to motor effects alone. However, it cannot be excluded that serotonergic neurons in these distinct midline regions contribute to the reduced mechanical thresholds, either indirectly or by direct altering motor neuron activity in the ventral horn [21; 34; 40]. Further studies of sensory function utilizing intersectional strategies to precisely capture and manipulate the midline populations will help to validate the present findings [49].

Together the results of the chemogenetic experiments indicate that the same transmitter released into different spinal laminae can produce opposing effects on spinal nociception. Potential explanations for these findings include the 5-HTR subtypes that are engaged, the spinal neuron populations that are modulated, or a combination of both. Some 5-HTRs are known to have an excitatory influence when activated, whereas others have an inhibitory effect upon ligand binding [1; 7; 44]. The activation of excitatory neurons and the inhibition of

inhibitory neurons in the deep dorsal horn, corresponding to the termination zone of the

medial serotonergic neurons, are known increase mechanical sensitivity [23; 24; 51; 52].

Conversely, the activation of inhibitory neurons in the superficial dorsal horn, the region

innervated by the lateral serotonergic neurons, and the inhibition of nociceptors through

presynaptic inhibition of their central terminals could promote thermal hyposensitivity [38; 44;

71]. Further studies are required to elucidate precisely which spinal 5-HT receptors and

neuronal elements are mediating these opposing sensory phenomena.

Given the previously reported heterogeneity in biophysical properties of serotonergic neurons

[49], we found surprisingly little difference in the passive and active membrane properties

between NRM and LPGi serotonergic neurons (table 5 and figure 4). Additionally, none of the

recorded neurons displayed spontaneous activity, which has been reported previously for

serotonergic projection neurons of the hindbrain [73]. This may be attributable to differences

in recording conditions, such as recording temperature, animal species, and recording

solution composition. Additionally, these groups of neurons may exhibit differences in

electophysiological properties, such as chemosensitivity, spontaneous activity, and

responses to nociceptive stimuli, that are only detected in *in vivo* preparations or under

recording conditions that differ from the present study [27; 28; 34; 45; 55; 56].

Limitations of transsynaptic rabies tracing of serotonergic RVM neurons

Transsynaptic rabies tracing has been widely used in recent years to identify presynaptic

inputs to genetically defined neuronal populations and has become a frequently used tool for

the study of neuronal circuits. Functional studies are limited due to the toxicity of first

generation ∆G rabies viruses. However, other limitations of this technique are known, such

as the resistance of some neuron types to infection by modified rabies viruses [2]. It is

possible that some types of hindbrain neurons are completely resistant to infection with

modified rabies viruses, although this is not the case for serotonergic neurons, as they can

be infected via their axon terminals (figure 1- figure supplement 4A-C). We show that

although descending TPH2-expressing neurons are not resistant to infection from G-protein

deficient rabies viruses, they are rarely traced transsynaptically from dorsal horn neurons

(figure 1 - figure supplement 4). The lack of transsynaptic labelling from the dorsal horn is

possibly due to the synaptic organization of serotonergic axons at this site, since serotonin is

known to mainly act via volume transmission [57]. Since similar numbers of hindbrain

neurons were labelled using direct rabies infection and transsynaptic rabies labelling, this

could indicate an increased labelling efficiency of other neurons with the latter approach.

Potentially this could be related to the synaptic density from these neurons to the starter

population. However, both direct and transsynaptic rabies labelling were far less sensitive

than the CTb or AAV2retro tracing, making interpretation of these findings complicated due

to the comparatively small sample sizes.

Most serotonergic axons in the rat dorsal horn do not form synaptic contacts [46; 57].

However, some 5-HT-containing axonal boutons do form synapses within the dorsal horn,

and these are observed as symmetrical synapses [53]. Since projection neurons are

approximately 5% of all lamina I neurons [15], the transsynaptic tracing from these neurons

may not be detected in our experiments. Additionally, transsynaptic tracing from a primary

afferent starter population was used to label serotonergic hindbrain neurons, suggesting axo-

axonic synapses between descending serotonergic neurons and their central terminals [75].

In the present study, the number of primary afferent neurons transduced from intraspinal

injection of helper virus is likely small, since the AAV8 serotype and the inclusion of the hSyn

promoter were used to reduce the transduction of sensory neurons [32]. This is consistent

with the low number of mCherry-expressing cells found in the DRG taken from these

animals. Taken together, these may explain why so few serotonergic hindbrain neurons were

traced transsynaptically from the spinal dorsal horn [25; 43].

Preferential transduction of LPGi serotonergic neurons with AAV2retro serotype vectors

The AAV2retro serotype has been developed for the retrograde delivery of genetic material to projection neurons via their axon terminals in a target region [60]. Many projection neurons are found to display some resistance to transduction with these tools, for example dopaminergic neurons that project from the substantia nigra to the striatum and descending noradrenergic neurons of the locus coeruleus to the spinal cord [26; 60]. Like others, we also find that midline neurons including the NRM serotonergic neurons are less sensitive to AAV2retro-mediated transduction [65]. Surprisingly, we observed that the serotonergic neurons of the lateral hindbrain do not display such resistance (Figure 1). This would suggest that these neurons are different in some property that allows the AAV2retro to enter their terminals or enhances the retrograde transport of the viral payload. The increased retrograde labelling efficiency of AAV2retro is not fully understood, although enhanced entry into axon terminals by increased spread of the vector in the injection site, entry into neurons via novel cell surface receptor interactions, and vesicular trafficking have been suggested [60]. AAV2retro-resistant neurons commonly use monoamines as neurotransmitters, suggesting that they likely share common features protecting them from transduction by these vectors.

Primarily, our data demonstrate that serotonergic neurons of the lateral hindbrain are antinociceptive when activated. These neurons project their axons to the superficial laminae of the dorsal horn and can be targeted using AAV2retro vectors. In contrast, the medial serotonergic neurons are largely resistant to AAV2retro-mediated transduction, produce mechanical hypersensitivity upon acute activation, and project their axons to the deeper dorsal horn. Secondly, we highlight some of the limitations and challenges associated with AAV2retro-mediated and rabies virus-based transsynaptic tracing from the spinal dorsal horn. These limitations are likely to have broader implications for transsynaptic rabies-based circuit tracing and the targeting of other projections that use biogenic amines, and possibly other transmitter systems.

Materials and Methods

Animals

Mice of either sex aged between 6-12 weeks were used for experiments. Permission to

perform these experiments was obtained from the Veterinäramt des Kantons Zürich

(154/2018 and 063/2016). The various transgenic mouse lines used in this study are listed in

table 6. TPH2::cre and HoxB8::cre mice are both BAC transgenic lines, and the ROSA^{TVA} is a

knockin reporter line.

Surgeries

The dorsal horn of the lumbar spinal cord was injected in a similar manner to previous

studies [26; 32]. Briefly, anesthesia of mice was induced with 5% Isoflurane in an induction

chamber. Mice were then transferred to a stereotaxic injection setup and anesthesia was

maintained with 1-3% Isoflurane delivered through a face mask. Body temperature was

maintained using a heated mat placed beneath the animal. Vitamin A cream was applied to

the eyes to prevent corneal drying during the operation and buprenorphine (0.1-0.2 mg/kg)

was injected subcutaneously prior to the operation. The back of the animal was shaved, and

the skin was scrubbed with Betadine solution. Once dried, a midline incision was made

above the vertebral column to expose the T13 vertebra, which was clamped using a pair of

spinal adaptors to isolate movements from the animal's breathing. This vertebra was

selected since it is directly above the lumbar L3 spinal cord, which corresponds to the spinal

cord segment receiving innervation from the hindlimbs [32]. A borehole was made in the

center of the left-hand side of the clamped vertebra and viruses, and/or 1% cholera toxin b

(CTb) were injected into the dorsal horn at a depth of 300 μm below the spinal surface

approximately 500 μm left of the central artery. For most injections, 3 x 300 nl virus solution

was injected along the rostrocaudal extent of the spinal cord at an infusion rate of 50 nl/min.

For transsynaptic tracing experiments, the rabies virus was injected 2 x 500 nl at either side

of the T13 vertebra in the same region as the injection of helper virus, which was injected two

weeks previously. For a list of the retrograde tracers and viruses used in this study, please

refer to table 7

For injections into the hindbrain, animals were prepared for surgery in a similar manner to the

spinal cord injections. The head was fixed in position using ear bars mounted on the

stereotaxic frame (Kopf instruments). Coordinates were chosen based on the hindbrain

location of the TPH2::Cre neurons retrogradely labelled from the spinal cord with reference to

a mouse brain atlas (-6, +/-0.5, 5.9, 1 μl per injection site, infusion rate 50 nl/min). All

coordinates are given as rostrocaudal, mediolateral, and dorsoventral (x, y, z) relative to

Bregma. Injections into the midline nucleus raphe magnus were made at coordinates -6, 0,

5.9 and a volume of 300nl was chosen to limit the transgene expression to the medial

serotonergic neurons. The movement of the frame was achieved using motorized axes

controlled by a computer interface, which was also used to select the injection target

(Neurostar). This same software was used to adjust the injection target for tilt and scaling, by

adjusting the mouse brain atlas relative to four points on the surface of the skull (Bregma,

Lambda, 2 mm to the right of the midline, and 2 mm to the left of the midline).

General features of tissue preparation and immunohistochemistry

Animals were perfusion fixed with freshly depolymerized 4% paraformaldehyde (room

temperature, dissolved in 0.1 M PB, adjusted to pH 7.4) following a brief rinse of the mouse

circulatory system with 0.1 M PB. Nervous tissues were quickly dissected and post-fixed in

the same fixative for two hours at 4˚C. After post-fixation, tissues were rinsed 3 times with

0.1 M PB and placed in 30% sucrose solution (w/v dissolved in 0.1 M PB) for 24-72 hours for

cryoprotection. Tissues were rinsed with 0.1 M PB before being embedded in NEG-50

mounting medium and were either cut at 60 μm on a sliding blade microtome (Hyrax KS 34,

histocam AG) and stored as free-floating sections, or were cut at 30 μm using a cryostat

(Hyrax 60, histocam AG) and were mounted directly onto microscope slides (Superfrost Plus, Thermoscientific).

Free floating sections were processed immediately for tissue staining or were stored in antifreeze medium (50 mM sodium phosphate buffer, 30% ethylene glycol, 15% glucose, and sodium azide (200 mg/L) at -20°C until required. Antifreeze medium was removed by rinsing sections three times in 0.1 M PB before further processing. Sections were rinsed in 50% ethanol for 30 minutes at room temperature, followed by three rinses in PBS + 8g/L NaCl Alternatively, slides were prepared by mounting frozen sections directly onto superfrost slides. These were either used immediately for immunostaining or stored at -80°C until further use. Before immunostaining, excess NEG50 freezing medium was removed by rinsing in 0.1 M PB for one hour at room temperature. A hydrophobic barrier was drawn around the sections using a fat pen, and immunoreactions were performed on the slides.

All primary antibodies and dilutions used in this study are included in table 8. Primary antibodies were diluted in a PBS + 8g/L NaCl, 0.3% v/v Triton-X, and 10% v/v normal donkey serum. Secondary antibodies were diluted in a similar solution, but without the 10% normal donkey serum (Table 8). Sections or slides were incubated in primary antibodies for 24 – 72 hours at 4°C, and these were revealed by incubation in secondary antibodies overnight at 4°C. Following immunostaining, sections or slides were rinsed three times for 10 minutes each in PBS + 8g/L NaCl before being mounted in Dako anti-fade medium.

Rabies virus tracing experiments

Rabies virus tracing experiments used either the SAD-G pseudotyped or EnvA pseudotyped RabΔG-GFP viruses, which are deficient in the SAD-glycoprotein required for transsynaptic spread and contain eGFP for identification of infected cells [68; 69]. Rabies viruses pseudotyped with the SAD-G glycoprotein can directly infect most, but not all, types of neurons [2; 68]. The rabies viruses that are pseudotyped with the EnvA glycoprotein are only

able to infect cells through binding to the TVA receptor. Therefore, the restricted expression

of the TVA receptor to certain cell types allows the selective infection of those neurons with

EnvA pseudotyped rabies viruses [69]. For transsynaptic tracing experiments, the helper

virus containing the rabies glycoprotein was injected two weeks prior to rabies virus injection

to enable transsynaptic spread.

Animals were perfused 5 or 7 days after dorsal horn injection of rabies viruses. Hindbrain

sections from rabies virus injected tissue were cut at 30 μm and mounted directly onto

microscope slides. Tissues were immunostained for eGFP and TPH2 to determine the

proportion of labelled hindbrain neurons that contained detectable TPH2.

Image acquisition and analysis

For quantification of retrogradely labelled cells in the hindbrain, image stacks were acquired

at 5 μm z-spacing using a Zeiss lsm 800 confocal microscope with Zen blue software.

Confocal scans were made using 488, 561 and 640 nm lasers and the pinhole was set to 1

Airy Unit for reliable optical sectioning. Care was taken to acquire image stacks up to a depth

where there was clear immunoreactivity of all antigens to avoid false negatives during

quantification due to antibody penetration. Within each experiment, all acquisition settings

were kept constant, and images were analyzed with FIJI using the cell counter plugin.

Alternatively, an automated cell quantification pipeline was designed in CellProfiler to count

retrogradely labelled neurons in the RVM. We noticed that many large neurons within the

RVM contained NeuN immunoreactivity that was present in more than one image in the

image stacks (5 μm z-spacing). Automated counting of all images within each z-stack would

lead to double counting a large proportion of cells present on multiple images within the

image stack. However, if only one image was analyzed per image stack, numerous cells

would be excluded from the analysis. Therefore, the acquired z-stacks were processed into

either one to two orthogonal projections from 2 images to reduce the double counting of cells

whilst ensuring the counting of most visible cells within each stack. One sample projection was then counted manually using the cell counter plugin of ImageJ, to serve as a guide in the development of the CellProfiler pipeline. All orthogonal projections were processed using the same CellProfiler pipeline with set parameters to ensure consistent data collection.

The images were loaded into this CellProfiler pipeline and the three channels (CTb, eGFP and NeuN) were separated from each other. NeuN- and CTb-stained neurons were identified as primary objects using the two-class Otsu method as a global thresholding strategy, whilst eGFP-containing objects were identified using the adaptive thresholding strategy (with an adaptive window of 50 pixels). NeuN stained cells were distinguished from the background based on their intensity, whereas the method used for detecting CTb and eGFP positive cells was based on their shape. The NeuN objects were converted into a binary (black & white) image and the coexpression of eGFP or CTb was assessed. The object counts of eGFP/NeuN, CTb/NeuN, eGFP/CTb/NeuN and NeuN positive objects were then extracted and exported to a spreadsheet.

To determine whether serotonergic neurons were labelled using modified rabies virus tracing, eGFP labelled neurons in the hindbrain were identified and immunostained against TPH2. These were scanned on a Zeiss 710 LSM confocal microscope, either as a short stack through the cell body, or as a single optical section through the center of the neuron. Scan settings were determined by the fluorescent intensity of the surrounding TPH2 immunoreactive regions in the nucleus raphe magnus (NRM) and lateral paragigantocellularis (LPGi) and were kept consistent for all experiments. Cell identities were catalogued and TPH2 immunoreactivity was determined for all sampled cells.

Slice preparation and electrophysiology

Hindbrain slices were prepared from TPH2::Cre animals that had received a bilateral injection of AAV9.flex.tdTomato into the ventral hindbrain. Animals were aged 3-6 weeks at the time of injection and were prepared for electrophysiological recordings 1-2 weeks later.

Slices were prepared in a similar manner to previous studies [50; 73]. Animals were decapitated and the brain was rapidly dissected and placed in ice cold oxygenated dissection solution (containing in mM (65 NaCl, 105 sucrose, 2.5 KCl, 1.25 NaH_2PO_4, 25 $NaHCO_3$, 25 glucose, 0.5 $CaCl_2$, 7 $MgCl_2$). The hindbrain was isolated, glued to a block of 2% agarose and installed in a slicing chamber. Transverse slices of hindbrain were cut at 250 μm on a vibrating blade microtome (D.S.K microslicer DTK1000), which were allowed to recover for at least 30min in oxygenated aCSF at 34°C prior to recording, containing (in mM) 120 NaCl, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 2.5 KCl, 5 HEPES, 14.6 glucose, 2 $CaCl_2$, 1 $MgCl_2$), pH 7.35 - 7.40, osmolarity 305 - 315 mOsm.

During recording, slices were perfused with aCSF at a flow rate of 2-3 ml/min. Targeted recordings were taken from tdTomato-expressing neurons using glass microelectrodes filled with a K-gluconate internal solution (containing 130 K-Gluconate, 5 NaCl, 1 EGTA, 10 HEPES, 5 Mg-ATP, 0.5 Na-GTP, 2 biocytin). Whole-cell recordings were acquired using a HEKA EPC10 amplifier with Patchmaster software at a sampling frequency of 20 kHz (HEKA Elektronik). A biophysical characterization of passive and active membrane properties was performed in current and voltage clamp modes, and the access resistance was monitored between recordings using a 10mV voltage step protocol. Data were excluded if the access resistance changed >30% during recording.

The relative position of the recorded/labelled neurons in each slice was noted, and slices were fixed overnight in 4% PFA at 4°C at the end of each experiment. Slices were immunoreacted with primary antibodies against tdTOM and TPH2, which were revealed the next day with secondary antibodies conjugated to Cy3 or Alexa 647. Biocytin was revealed with Avidin-Alexa A488 and the position of the filled neurons was assigned to the NRM or the LPGi, which could be determined based on the pattern of TPH2 immunoreactivity in the slice. Cells located outside of these two regions were not analyzed further.

Behavioral assays

For specific DREADD-mediated activation of descending serotonergic pathways, an

intersectional approach was used for labelling the lateral hindbrain neurons whereas direct

labelling was used to label the medial neurons. TPH2::Cre mice received an intraspinal

injection of AAV2retro.flex.FLPo.BFP, and one week later received bilateral injections into

the ventral hindbrain with AAV1.FRT.hM3Dq.mCherry. Behavioral tests were performed after

10 days incubation time to allow the expression of the viral transgene. Before experiments

mice were acclimatized to the behavioral setup for one hour in a room maintained at 20-

22°C. For the Hargreaves plantar, cold plantar, electronic von Frey, and Rotarod assays, six

measurements were taken for each time point and an average of these was reported. All

measurements were taken from both hindlimbs of all animals. Alternatively, serotonergic

neurons in the medial hindbrain were labelled by injection of AAV8.hsyn.flex.hM3Dq into the

NRM (injection coordinates -6, 0, 5.9) and the same behavioral tests were performed with the

same experimental design.

Hargreaves

Sensitivity to heat stimuli was assessed with the Hargreaves plantar assay (IITC). Mice were

placed on a pre-heated transparent platform set to a temperature of 30°C, and the

withdrawal latencies were recorded using a timed infrared heat source. A resting intensity of

5% and an active intensity of 20% was used for stimulations, and a maximum cutoff time of

32 s was set to avoid tissue damage. On average, naïve animals will withdraw to this

stimulation at temperatures between 34 - 36°C [11].

Cold plantar assay

Mice were placed on a 5 mm borosilicate glass platform and were stimulated from beneath

with dry ice pellets. The time taken to withdraw the paw was measured using a stopclock and

a maximum stimulation time of 20 s was used to avoid tissue damage. The cold plantar

assay generally produces a withdrawal when the temperature of the glass decreases 2°C for

naïve animals [11].

von Frey

Von Frey thresholds were measured using an electronic von Frey algesiometer (IITC).

Animals were adapted on a mesh surface, and the plantar surface of each paw was

stimulated with a bendable plastic filament attached to a pressure sensitive probe. Pressure

was applied to the plantar surface in a linear manner until the animal withdrew its paw, and

the maximum pressure (the pressure at which the animal withdrew) was displayed on the

device.

Rotarod

Sensorimotor coordination was evaluated using an accelerating rotarod, and the time taken

for animals to fall from the rotating barrel was recorded. The barrel rotated from 4 - 40rpm

over a period of 300s, and increased speed constantly throughout each experiment. Values

were discarded if the animal jumped from the barrel, and if the animal jumped in >50% of

trials for a given time point these data were discarded from the experiment. Two training

sessions were given for all animals prior to the experiment being started to ensure a stable

performance in the absence of treatment.

Drug application

For DREADD activation experiments, clozapine-N-oxide (CNO, 2 mg/kg, Enzo life sciences,

product number BML-NS105-0025) or vehicle controls were injected intraperitoneally, with

the experimenter being blinded to the injected substance. Animals were tested directly before

and 1-3 hours following i.p. injection. Stock CNO was dissolved in DMSO at 100 mg/ml and

kept at room temperature, which was diluted 1:500 in sterile filtered saline immediately prior

to injection with the volume of injected substance being 10 μl/g.

Experimental design and statistical tests

All behavioral assays were performed twice on each animal so that they received an

intraperitoneal injection of CNO on one day and Vehicle injection on the other, with the

experimenter being blinded to the treatment. Response latencies and thresholds following

CNO and Vehicle were compared using a repeated measures one-way ANOVA, a normal

distribution was assumed for response latencies and thresholds and a Bonferroni post-hoc

test was used to compare the mean withdrawal latencies and thresholds between CNO and

vehicle treated groups before and 1-3 hours after injection. Statistical significance was taken

as $P < 0.05$.

To compare membrane properties between medial and lateral serotonergic neurons, a

normal distribution in values was assumed and an unpaired two-tailed t-test was used to test

for differences.

Data collection, storage, and presentation

Data obtained from the quantification of images were collected and processed in Microsoft

Excel and were presented using GraphPad Prism 5. All datapoints on graphs represent

either a number or percentage of cells counted per animal, or as a response for each animal

in behavioral assays. Representative images were produced in Affinity Photo and were

annotated and arranged into figures in Affinity Designer. Raw data acquired in these

experiments are uploaded to www.datadryad.org and are available for download.

Figure 1: Retrograde labelling of spinally-projecting serotonergic neurons with AAV2retro vectors and Cholera toxin b subunit. A. Injection scheme for retrograde labelling of spinally-projecting neurons with AAV2retro.eGFP. B Image of the ventral hindbrain containing eGFP-labelled neurons (scale bar = 200 μm). Inset shows enlargement of the LPGi to reveal eGFP neurons that also express TPH2. C. Quantification of cell location for eGFP-labelled neurons that express TPH2, each datapoint is a count per animal (n = 3) D. Injection scheme for retrograde tracing from the spinal dorsal horn with AAV2retro.eGFP and CTb. E. Representative image of the ventral hindbrain containing CTb-labelled and AAV2retro-transduced projection neurons (scale bar = 200 μm). F. Anatomical locations of retrogradely labelled TPH2+ hindbrain neurons labelled with CTb or AAV2retro (n = 3 animals).

FIgure 1 supplement 1: Defined areas in the RVM used for quantifying the location of retrogradely-labelled neurons. The NRM was identified as a dense cluster of TPH2-expressing neurons around the midline in a triangular shape, whereas the LPGi were distinguished by their lateral location and separation from the midline structures. All other regions were defined as "other". Examples of retrogradely labelled TPH2+ neurons in the NRM are indicated with arrows, the LPGi with filled arrowheads, and other with empty arrowheads, NRM = nucleus raphe magnus, LPGi = lateral paragigantocellularis, Pyr = Pyramids,

Figure 1 supplement 2: Retrograde labelling of serotonergic hindbrain neurons with CTb. A. injection scheme for CTb tracing from the lumbar dorsal horn and an example of the injection site in the lumbar spinal cord (scale bar = 200 μm). B. Example of a hindbrain section from the injection site shown in A (scale bars = 200 μm). The inset shows that many CTb-labelled neurons that express TPH2 are found in the nucleus raphe magnus (indicated by arrows). C. Location of CTb-labelled cells in the RVM that express TPH2, each datapoint is a count per animal (n = 2).

Figure 1 supplement 3. Proportion of neurons retrogradely labelled with CTb and

AAV2retro in different RVM areas. A. Injection scheme for labelling descending projection

neurons with AAV2retro and CTb. B. Quantification of coexpression of CTb and eGFP in all

retrogradely-labelled RVM neurons, and C. Quantification of coexpression of CTb and eGFP

in retrogradely-traced RVM neurons that contain TPH2. The proportion of retrogradely cells

in each area expressing CTb, eGFP, or both eGFP and CTb are illustrated. The cell counts

used to generate these charts are given in Table 1 (n = 3 animals).

Figure 1 supplement 4: labelling of spinally-projecting serotonergic neurons with

direct rabies infection and transsynaptic rabies tracing. A. Injection strategy for the

direct labelling of spinally-projecting hindbrain neurons with a SAD pseudotyped and G-

protein deficient rabies virus. B. Example of a cell body of a labelled hindbrain neuron that is

immunoreactive for TPH2 (scale bar = 50 μm), the ventral border of the hindbrain is indicated

by the dotted line. C. Quantification of the GFP-labelled hindbrain neurons that express

detectable levels of TPH2. Note that 7 days after rabies virus injection there are fewer

labelled neurons that express TPH2. N = 3 animals. D. Injection strategy for the

transsynaptic tracing of neurons from spinal dorsal horn neurons. Hoxb8-Cre is transiently

expressed in all lumbar dorsal horn neurons and is crossed with ROSA26[TVA] to induce TVA

expression in all spinal neurons. E. Example of an RVM neuron traced transsynaptically from

the spinal dorsal horn (scale bar = 50 μm) F. Quantification of GFP-labelled neurons in the

hindbrain that express detectable TPH2. Data from C. "5 days post injection" is included to

allow comparison between directly labelled and transsynaptically traced neurons. N = 4

animals for transsynaptic tracing experiment.

Figure 2: Labelling of spinally-projecting neurons in the TPH2::Cre mouse with

AAV2retro vectors. A. Injection scheme for labelling spinally-projecting TPH2::Cre neurons.

B. An example of a hindbrain section containing neurons labelled with eGFP (scale bar = 200

μm). Inset is an enlargement of the area indicated in the dashed box, with many eGFP

labelled cells found to express detectable TPH2 that are indicated with arrows. Cells that

were labelled with eGFP not containing detectable TPH2 are indicated with arrowheads. C.

Image of an ipsilateral DRG from a TPH2::Cre mouse that received a spinal injection of

AAV2retro.flex.eGFP, showing many eGFP-expressing neurons (scale bar = 100 μm). D.

Quantification of the location of eGFP-labelled cells in the hindbrain. E. Quantification of the

hindbrain neurons labelled with eGFP that also contain TPH2. For D. and E. each datapoint

is a count per animal (n = 3 animals). F. Injection scheme for the intersectional labelling of

spinally-projecting TPH2::Cre neurons. Brain injection coordinates (-6, +/-0.5, 5.9) from

bregma, for the labelling of spinally-projecting TPH2::Cre neurons in the LPGi, according to

the mouse brain atlas. Target injection sites are indicated in red crosshairs. G. Example of a

hindbrain section from a TPH2::Cre mouse that received the injections illustrated in F (scale

bar = 500 μm). Inset is an enlargement of the boxed area and highlights neurons that were

captured with the brain injection and express eGFP (arrowheads) as well as neurons that

were directly labelled from the spinal cord injection (mCherry+) that were not transduced

from the hindbrain injection (arrows). Scale bar = 200 μm H. Quantification of the mCherry-

expressing cells that are labelled with eGFP, which indicated that they were captured with

the hindbrain injection. The percentages of mCherry-only and eGFP-only cells are also

quantified for each hindbrain area that contains serotonergic neurons, and areas that could

not be assigned as either medial or LPGi were classified as "other" (n = 4 animals)

Figure 3: Spinal cord regions innervated by serotonergic hindbrain neurons. A.

Injection scheme for labelling medially located serotonergic neurons of the hindbrain with

AAVs. Stereotaxic injection coordinates for the hindbrain injection of AAVs to label midline

neurons without transducing LPGi serotonergic neurons (-6, 0, 5.9). B. Representative brain

injection site from the hindbrain of a TPH2::Cre mouse that received a single 300nl injection

of AAV8.hSyn.flex.eGFP (Scale bar = 500 μm). A higher magnification image of the injection

site is also shown (scale bar = 100 μm). C. Quantification of the proportion of eGFP-labelled

neurons in the injection site that were immunoreactive for TPH2. D. Intersectional strategy to

preferentially label the spinal axon terminals of descending serotonergic neurons that

originate in the LPGi. E. Representative images of the hindbrain injection site and the axon

termination pattern in the spinal cord of a TPH2::Cre animal that received the injections

depicted in D (scale bars = 500 μm and 100 μm for hindbrain and spinal cord sections

respectively). F. Injection scheme for labelling medially located serotonergic neurons with

tdTOM. G. Representative images of the hindbrain injection site and spinal cord axon

termination pattern of a TPH2::Cre animal that received a single 300nl injection of

AAV8.hSyn.flex.tdTOM depicted in F (scale bars = 500 μm and 100 μm for hindbrain and

spinal cord sections respectively). H. Higher magnification images of the cervical and lumbar

dorsal horns of a TPH2::Cre animal that received the injections depicted in D (scale bar =

100 μm). I. Representative images from cervical and lumbar spinal cord segments of animals

that received an injection with AAV8.hSyn.flex.tdTOM into the NRM (scale bar = 100 μm).

Figure 3 - figure supplement 1. Laminar location of axons originating from descending LPGi

and midline serotonergic neurons of the hindbrain. A. Intersectional labelling of TPH2::Cre

hindbrain neurons in the LPGi, axons and terminals are found within laminae I-IIo,

highlighted by CGRP immunostaining. B. Direct labelling of medial serotonergic hindbrain

neurons with AAV vectors, most labelled axons are found ventral to the lamina II-III border

delineated by PKCγ immunoreactivity. Scale bars = 100 μm, laminae I-IIo and IIi are

indicated by the dotted lines.

Figure 4: Electrophysiological characterization of serotonergic neurons of the NRM

and the LPGi. A. Current step protocol (red) and resulting AP firing (black) from

representative serotonergic NRM and LPGi neurons recorded in current clamp. B.

Current/frequency plot for the AP firing frequency of NRM and LPGi neurons resulting from

increasing 1s depolarizing current injections. C. Representative images of biocytin filled

neurons within the NRM and LPGi of hindbrain slices revealed following recording (scale bar

= 200 μm). D – H comparison of active and passive membrane properties between NRM and

LPGi serotonergic neurons. D – G. n = 10 NRM (from 4 animals) n = 8 LPGi (from 3

animals). H. n = 9 NRM n = 5 LPGi

Figure 5: Chemogenetic activation of descending serotonergic LPGi neurons. A.

Injection scheme for expressing the excitatory DREADD hM3Dq in spinally-projecting

TPH2::Cre neurons. Brain injection coordinates according to the mouse brain atlas (-6, +/-

0.5, 5.9 from bregma). B. Example of the injection site from the hindbrain of a mouse that

received a spinal dorsal horn injection of AAV2retro.flex.FLPo.BFP followed by a bilateral

hindbrain injection of AAV1.FRT.hM3Dq.mCherry one week later (scale bar = 200 μm). C.

Example of the ipsilateral spinal dorsal horn form an animal that had received the injections

indicated in A . Note that most 5-HT-containing terminals are not labelled, but the majority of

labelled terminals contain a detectable level of 5-HT (scale bar = 200 μm). D. Sensory tests

of the ipsilateral hindpaw: Hargreaves plantar assay; repeated measures one-way ANOVA,

$(F_{(3, 42)} = 16.93, P < 0.0001)$, post hoc tests with Bonferroni's correction detected

differences between post-Vehicle and post CNO, as well as pre-CNO and post-CNO

(adjusted P values are p = 0.007 and p < 0.0001 respectively) . Cold plantar assay; repeated

measures one-way ANOVA, $(F_{(3, 39)} = 12.41, P < 0.0001)$ post hoc tests with Bonferroni's

correction detected differences between post-Vehicle and post CNO, as well as pre-CNO

and post-CNO (adjusted P values are p = 0.0122 and p = 0.0103 respectively). von Frey test;

repeated measures one-way ANOVA, $(F_{(3, 39)} = 0.5013 P = 0.6836)$. Rotarod test for

sensorimotor coordination/sedation; repeated measures one-way ANOVA, $(F_{(3, 30)} =$

$0.8684, P = 0.4683)$. Significance: *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 5 figure supplement 1. Chemogenetic activation of hM3Dq-labelled neurons. A.

Hindbrain containing hM3Dq-mCherry labelled neurons that upregulated c-Fos following

intraperitoneal injection of CNO (scale bar = 200 μm). B. Quantification of the proportion of

hM3Dq-mCherry neurons that upregulated c-Fos after CNO or Vehicle injection (CNO n = 2,

Vehicle n = 3).

Figure 5 figure supplement 2: Altered sensitivity of the contralateral paw following LPGi activation. Sensory tests of the contralateral hindpaw: Hargreaves plantar assay; repeated measures one-way ANOVA, ($F_{(3, 9)}$ = 4.833, P = 0.0081). Post-hoc tests with Bonferroni correction detect differences between pre and post CNO injection, and between post-Vehicle and post-CNO injection in the Hargreaves plantar assay. Significance: **$p <$ 0.01, ***$p < 0.001$. (n = 15 animals).

Figure 5 figure supplement 3: Both male and female mice show alterations in thermal thresholds following chemogenetic activation of LPGi serotonergic neurons. Sensory and motor tests with groups divided into male and female animals. Increases in withdrawal latencies are seen in the Hargreaves and cold plantar assay in both male and female animals (paired t-tests post Vehicle vs post CNO with Holm-Sidak correction). N = 6 male and 9 female animals Hargreaves, 6 male and 8 female for Cold plantar assay and von Frey, and 6 male and 5 female for Rotarod. Significance: *$p < 0.05$, **$p < 0.01$, **$p < 0.001$.

Figure 5 figure supplement 4: CNO does not alter response latencies, thresholds or sensorimotor coordination in the absence of hM3Dq. Sensory tests before and after the injection of either CNO or Vehicle (n = 5 animals). post-Vehicle vs post-CNO paired t-test, Hargreaves $p = 0.239$, Cold plantar $p = 0.461$, von Frey $p = 0.424$, Rotarod $p = 0.732$.

Figure 5 figure supplement 5: Proportion of AAV2retro-traced neurons labelled with hM3Dq-mCherry for behavioral experiments. Quantification of the hindbrain neurons labelled from the injection scheme depicted in Figure 5A. n = 4 animals.

Figure 6: Chemogenetic activation of medial serotonergic hindbrain neurons. A. Injection scheme for chemogenetic activation of medial serotonergic hindbrain neurons. Stereotaxic coordinates used for activating midline serotonergic neurons of TPH2::Cre animals with hM3D(q)-containing AAVs (-6, 0, 5.9). B. Example of an injection site from an experiment to activate midline serotonergic neurons with hM3D(q). C. Lumbar spinal cord from the injection site shown in B. showing mCherry-expressing terminals located ventral to

the dense 5-HT innervation of the superficial dorsal horn. D. Sensory and sensorimotor coordination assays of animals following CNO or Vehicle injections. Repeated one-way ANOVA, followed by Bonferroni's post-hoc tests for von Frey ($F_{(3, 6)}$ = 13.84, $P < 0.0001$) show significant decreases between post-Vehicle and post-CNO injections ($p = 0.0005$) as well as pre-CNO and post-CNO ($p < 0.0001$). Significance: *$p < 0.05$, **$p < 0.01$, **$p < 0.001$.

Figure 6 figure supplement 1: Injection sites for chemogenetic activation of medial serotonergic neurons of the hindbrain. In all cases, spread is restricted to the medial neurons with limited spread to the LPGi serotonergic neurons. The targeted region of the hindbrain was the NRM at (-6, 0, 5.9) relative to Bregma according to the mouse brain atlas. In many cases, hM3Dq-mCherry-expressing neurons are also seen in the Raphe Obscuris (ROb), and the Raphe Pallidus (RPa).

Figure 6 figure supplement 2: Both male and female animals show increase in mechanical hypersensivity during chemogenetic activation of medial serotonergic hindbrain neurons. Sensory and motor tests of male and female animals 1-3 hours after CNO or Vehicle injection. A decrease in withdrawal threshold is seen in both male and female animals for the von Frey test (paired t-tests post Vehicle vs post CNO with Holm-Sidak correction). N = 5 male and 6 female animals for all tests. Significance: *$p < 0.05$, **$p < 0.01$, **$p < 0.001$.

Figure 6 figure supplement 3: Proportion of neurons labelled in chemogenetic experiments that express TPH2. Summary of the proportion of hM3Dq-mCherry neurons labelled in the medial hindbrain that also express detectable TPH2 (n = 3 animals)

References

[1] Abe K, Kato G, Katafuchi T, Tamae A, Furue H, Yoshimura M. Responses to 5-HT in morphologically identified neurons in the rat substantia gelatinosa in vitro. Neuroscience 2009;159(1):316-324.

[2] Albisetti GW, Ghanem A, Foster E, Conzelmann KK, Zeilhofer HU, Wildner H. Identification of Two Classes of Somatosensory Neurons That Display Resistance to Retrograde Infection by Rabies Virus. J Neurosci 2017;37(43):10358-10371.

[3] Albisetti GW, Pagani M, Platonova E, Hösli L, Johannssen HC, Fritschy JM, Wildner H, Zeilhofer HU. Dorsal Horn Gastrin-Releasing Peptide Expressing Neurons Transmit Spinal Itch But Not Pain Signals. J Neurosci 2019;39(12):2238-2250.

[4] Bannister K, Dickenson AH. What do monoamines do in pain modulation? Curr Opin Support Palliat Care 2016;10(2):143-148.

[5] Bannister K, Lockwood S, Goncalves L, Patel R, Dickenson AH. An investigation into the inhibitory function of serotonin in diffuse noxious inhibitory controls in the neuropathic rat. Eur J Pain 2017;21(4):750-760.

[6] Bannister K, Patel R, Goncalves L, Townson L, Dickenson AH. Diffuse noxious inhibitory controls and nerve injury: restoring an imbalance between descending monoamine inhibitions and facilitations. Pain 2015;156(9):1803-1811.

[7] Bardoni R. Serotonergic Modulation of Nociceptive Circuits in Spinal Cord Dorsal Horn. Curr Neuropharmacol 2019;17(12):1133-1145.

[8] Basbaum AI, Clanton CH, Fields HL. Opiate and stimulus-produced analgesia: functional anatomy of a medullospinal pathway. Proc Natl Acad Sci U S A 1976;73(12):4685-4688.

[9] Basbaum AI, Fields HL. Endogenous pain control systems: brainstem spinal pathways and endorphin circuitry. Annu Rev Neurosci 1984;7:309-338.

[10] Basbaum AI, Marley NJ, O'Keefe J, Clanton CH. Reversal of morphine and stimulus-produced analgesia by subtotal spinal cord lesions. Pain 1977;3(1):43-56.

[11] Brenner DS, Golden JP, Gereau RWt. A novel behavioral assay for measuring cold sensation in mice. PLoS One 2012;7(6):e39765.

[12] Brust RD, Corcoran AE, Richerson GB, Nattie E, Dymecki SM. Functional and developmental identification of a molecular subtype of brain serotonergic neuron specialized to regulate breathing dynamics. Cell Rep 2014;9(6):2152-2165.

[13] Butkevich IP, Barr GA, Vershinina EA. Sex differences in formalin-induced pain in prenatally stressed infant rats. Eur J Pain 2007;11(8):888-894.

[14] Cai YQ, Wang W, Hou YY, Pan ZZ. Optogenetic activation of brainstem serotonergic neurons induces persistent pain sensitization. Mol Pain 2014;10:70.

[15] Cameron D, Polgár E, Gutierrez-Mecinas M, Gomez-Lima M, Watanabe M, Todd AJ. The organisation of spinoparabrachial neurons in the mouse. Pain 2015;156(10):2061-2071.

[16] Cavanaugh DJ, Chesler AT, Jackson AC, Sigal YM, Yamanaka H, Grant R, O'Donnell D, Nicoll RA, Shah NM, Julius D, Basbaum AI. Trpv1 reporter mice reveal highly restricted brain distribution and functional expression in arteriolar smooth muscle cells. J Neurosci 2011;31(13):5067-5077.

[17] Cavanaugh DJ, Lee H, Lo L, Shields SD, Zylka MJ, Basbaum AI, Anderson DJ. Distinct subsets of unmyelinated primary sensory fibers mediate behavioral responses to noxious thermal and mechanical stimuli. Proc Natl Acad Sci U S A 2009;106(22):9075-9080.

[18] Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sik A, De Koninck P, De Koninck Y. Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. Nature 2003;424(6951):938-942.

[19] Dahlstrom A, Fuxe K. Localization of monoamines in the lower brain stem. Experientia 1964;20(7):398-399.

[20] Das Gupta RR, Scheurer L, Pelczar P, Wildner H, Zeilhofer HU. Neuron-specific spinal cord translatomes reveal a neuropeptide code for mouse dorsal horn excitatory neurons. Sci Rep 2021;11(1):5232.

[21] Depuy SD, Kanbar R, Coates MB, Stornetta RL, Guyenet PG. Control of breathing by raphe obscurus serotonergic neurons in mice. J Neurosci 2011;31(6):1981-1990.

[22] Dhaka A, Earley TJ, Watson J, Patapoutian A. Visualizing cold spots: TRPM8-expressing sensory neurons and their projections. J Neurosci 2008;28(3):566-575.

[23] Duan B, Cheng L, Bourane S, Britz O, Padilla C, Garcia-Campmany L, Krashes M, Knowlton W, Velasquez T, Ren X, Ross S, Lowell BB, Wang Y, Goulding M, Ma Q. Identification of spinal circuits transmitting and gating mechanical pain. Cell 2014;159(6):1417-1432.

[24] Foster E, Wildner H, Tudeau L, Haueter S, Ralvenius WT, Jegen M, Johannssen H, Hösli L, Haenraets K, Ghanem A, Conzelmann KK, Bösl M, Zeilhofer HU. Targeted ablation, silencing, and activation establish glycinergic dorsal horn neurons as key components of a spinal gate for pain and itch. Neuron 2015;85(6):1289-1304.

[25] Francois A, Low SA, Sypek EI, Christensen AJ, Sotoudeh C, Beier KT, Ramakrishnan C, Ritola KD, Sharif-Naeini R, Deisseroth K, Delp SL, Malenka RC, Luo L, Hantman AW, Scherrer G. A Brainstem-Spinal Cord Inhibitory Circuit for Mechanical Pain Modulation by GABA and Enkephalins. Neuron 2017;93(4):822-839.e826.

[26] Ganley RP, Werder K, Wildner H, Zeilhofer HU. Spinally projecting noradrenergic neurons of the locus coeruleus display resistance to AAV2retro-mediated transduction. Mol Pain 2021;17:17448069211037887.

[27] Gao K, Mason P. Serotonergic Raphe magnus cells that respond to noxious tail heat are not ON or OFF cells. J Neurophysiol 2000;84(4):1719-1725.

[28] Gao K, Mason P. The discharge of a subset of serotonergic raphe magnus cells is influenced by baroreceptor input. Brain Res 2001;900(2):306-313.

[29] Gatto G, Bourane S, Ren X, Di Costanzo S, Fenton PK, Halder P, Seal RP, Goulding MD. A Functional Topographic Map for Spinal Sensorimotor Reflexes. Neuron 2020.

[30] Gautier A, Geny D, Bourgoin S, Bernard JF, Hamon M. Differential innervation of superficial versus deep laminae of the dorsal horn by bulbo-spinal serotonergic pathways in the rat. IBRO Rep 2017;2:72-80.

[31] Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, Ellis RJ, Richie CT, Harvey BK, Dannals RF, Pomper MG, Bonci A, Michaelides M. Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. Science 2017;357(6350):503-507.

[32] Haenraets K, Albisetti GW, Foster E, Wildner H. Adeno-associated Virus-mediated Transgene Expression in Genetically Defined Neurons of the Spinal Cord. J Vis Exp 2018(135).

[33] Haring M, Zeisel A, Hochgerner H, Rinwa P, Jakobsson JET, Lonnerberg P, La Manno G, Sharma N, Borgius L, Kiehn O, Lagerstrom MC, Linnarsson S, Ernfors P. Neuronal atlas of the dorsal horn defines its architecture and links sensory input to transcriptional cell types. Nat Neurosci 2018;21(6):869-880.

[34] Hennessy ML, Corcoran AE, Brust RD, Chang Y, Nattie EE, Dymecki SM. Activity of Tachykinin1-Expressing Pet1 Raphe Neurons Modulates the Respiratory Chemoreflex. J Neurosci 2017;37(7):1807-1819.

[35] Hirschberg S, Li Y, Randall A, Kremer EJ, Pickering AE. Functional dichotomy in spinal- vs prefrontal-projecting locus coeruleus modules splits descending noradrenergic analgesia from ascending aversion and anxiety in rats. Elife 2017;6.

[36] Huang J, Polgar E, Solinski HJ, Mishra SK, Tseng PY, Iwagaki N, Boyle KA, Dickie AC, Kriegbaum MC, Wildner H, Zeilhofer HU, Watanabe M, Riddell JS, Todd AJ, Hoon MA. Circuit dissection of the role of somatostatin in itch and pain. Nat Neurosci 2018;21(5):707-716.

[37] Iceman KE, Richerson GB, Harris MB. Medullary serotonin neurons are CO_2 sensitive in situ. J Neurophysiol 2013;110(11):2536-2544.

[38] Jeong HJ, Mitchell VA, Vaughan CW. Role of 5-HT(1) receptor subtypes in the modulation of pain and synaptic transmission in rat spinal superficial dorsal horn. Br J Pharmacol 2012;165(6):1956-1965.

[39] Kaur S, Benton WL, Tongkhuya SA, Lopez CMC, Uphouse L, Averitt DL. Sex Differences and Estrous Cycle Effects of Peripheral Serotonin-Evoked Rodent Pain Behaviors. Neuroscience 2018;384:87-100.

[40] Kawashima T. The role of the serotonergic system in motor control. Neurosci Res 2018;129:32-39.

[41] Kwiat GC, Basbaum AI. The origin of brainstem noradrenergic and serotonergic projections to the spinal cord dorsal horn in the rat. Somatosens Mot Res 1992;9(2):157-173.

[42] Li L, Rutlin M, Abraira VE, Cassidy C, Kus L, Gong S, Jankowski MP, Luo W, Heintz N, Koerber HR, Woodbury CJ, Ginty DD. The functional organization of cutaneous low-threshold mechanosensory neurons. Cell 2011;147(7):1615-1627.

[43] Liu MZ, Chen XJ, Liang TY, Li Q, Wang M, Zhang XY, Li YZ, Sun Q, Sun YG. Synaptic control of spinal GRPR(+) neurons by local and long-range inhibitory inputs. Proc Natl Acad Sci U S A 2019.

[44] Lu Y, Perl ER. Selective action of noradrenaline and serotonin on neurones of the spinal superficial dorsal horn in the rat. J Physiol 2007;582(Pt 1):127-136.

[45] Mason P, Gao K, Genzen JR. Serotonergic raphe magnus cell discharge reflects ongoing autonomic and respiratory activities. J Neurophysiol 2007;98(4):1919-1927.

[46] Maxwell DJ, Leranth C, Verhofstad AA. Fine structure of serotonin-containing axons in the marginal zone of the rat spinal cord. Brain Res 1983;266(2):253-259.

[47] Millan MJ. Descending control of pain. Prog Neurobiol 2002;66(6):355-474.

[48] Mogil JS. Sex differences in pain and pain inhibition: multiple explanations of a controversial phenomenon. Nat Rev Neurosci, Vol. 13. England, 2012. pp. 859-866.

[49] Okaty BW, Commons KG, Dymecki SM. Embracing diversity in the 5-HT neuronal system. Nat Rev Neurosci 2019;20(7):397-424.

[50] Pedersen NP, Vaughan CW, Christie MJ. Opioid receptor modulation of GABAergic and serotonergic spinally projecting neurons of the rostral ventromedial medulla in mice. J Neurophysiol 2011;106(2):731-740.

[51] Peirs C, Williams SG, Zhao X, Arokiaraj CM, Ferreira DW, Noh MC, Smith KM, Halder P, Corrigan KA, Gedeon JY, Lee SJ, Gatto G, Chi D, Ross SE, Goulding M, Seal RP. Mechanical Allodynia Circuitry in the Dorsal Horn Is Defined by the Nature of the Injury. Neuron 2020.

[52] Peirs C, Williams SP, Zhao X, Walsh CE, Gedeon JY, Cagle NE, Goldring AC, Hioki H, Liu Z, Marell PS, Seal RP. Dorsal Horn Circuits for Persistent Mechanical Pain. Neuron 2015;87(4):797-812.

[53] Polgar E, Puskar Z, Watt C, Matesz C, Todd AJ. Selective innervation of lamina I projection neurones that possess the neurokinin 1 receptor by serotonin-containing axons in the rat spinal cord. Neuroscience 2002;109(4):799-809.

[54] Porreca F, Burgess SE, Gardell LR, Vanderah TW, Malan TP, Jr., Ossipov MH, Lappi DA, Lai J. Inhibition of neuropathic pain by selective ablation of brainstem medullary cells expressing the mu-opioid receptor. J Neurosci 2001;21(14):5281-5288.

[55] Potrebic SB, Fields HL, Mason P. Serotonin immunoreactivity is contained in one physiological cell class in the rat rostral ventromedial medulla. J Neurosci 1994;14(3 Pt 2):1655-1665.

[56] Richerson GB, Wang W, Tiwari J, Bradley SR. Chemosensitivity of serotonergic neurons in the rostral ventral medulla. Respir Physiol 2001;129(1-2):175-189.

[57] Ridet JL, Rajaofetra N, Teilhac JR, Geffard M, Privat A. Evidence for nonsynaptic serotonergic and noradrenergic innervation of the rat dorsal horn and possible involvement of neuron-glia interactions. Neuroscience 1993;52(1):143-157.

[58] Sharif B, Ase AR, Ribeiro-da-Silva A, Séguéla P. Differential Coding of Itch and Pain by a Subpopulation of Primary Afferent Neurons. Neuron 2020;106(6):940-951.e944.

[59] Suzuki R, Morcuende S, Webber M, Hunt SP, Dickenson AH. Superficial NK1-expressing neurons control spinal excitability through activation of descending pathways. Nat Neurosci 2002;5(12):1319-1326.

[60] Tervo DG, Hwang BY, Viswanathan S, Gaj T, Lavzin M, Ritola KD, Lindo S, Michael S, Kuleshova E, Ojala D, Huang CC, Gerfen CR, Schiller J, Dudman JT, Hantman AW, Looger LL, Schaffer DV,

Karpova AY. A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. Neuron 2016;92(2):372-382.

[61] Todd AJ. Neuronal circuitry for pain processing in the dorsal horn. Nat Rev Neurosci 2010;11(12):823-836.

[62] Todd AJ. Plasticity of inhibition in the spinal cord. Handb Exp Pharmacol 2015;227:171-190.

[63] Tudeau L, Acuña MA, Albisetti GW, Neumann E, Ralvenius WT, Scheurer L, Poe M, Cook JM, Johannssen HC, Zeilhofer HU. Mice lacking spinal α2GABA(A) receptors: Altered GABAergic neurotransmission, diminished GABAergic antihyperalgesia, and potential compensatory mechanisms preventing a hyperalgesic phenotype. Brain Res 2020;1741:146889.

[64] Wang JK. Antinociceptive effect of intrathecally administered serotonin. Anesthesiology 1977;47(3):269-271.

[65] Wang Z, Maunze B, Wang Y, Tsoulfas P, Blackmore MG. Global Connectivity and Function of Descending Spinal Input Revealed by 3D Microscopy and Retrograde Transduction. J Neurosci 2018;38(49):10566-10581.

[66] Wei F, Dubner R, Zou S, Ren K, Bai G, Wei D, Guo W. Molecular depletion of descending serotonin unmasks its novel facilitatory role in the development of persistent pain. J Neurosci 2010;30(25):8624-8636.

[67] West SJ, Bannister K, Dickenson AH, Bennett DL. Circuitry and plasticity of the dorsal horn--toward a better understanding of neuropathic pain. Neuroscience 2015;300:254-275.

[68] Wickersham IR, Finke S, Conzelmann KK, Callaway EM. Retrograde neuronal tracing with a deletion-mutant rabies virus. Nat Methods 2007;4(1):47-49.

[69] Wickersham IR, Lyon DC, Barnard RJ, Mori T, Finke S, Conzelmann KK, Young JA, Callaway EM. Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. Neuron 2007;53(5):639-647.

[70] Witschi R, Johansson T, Morscher G, Scheurer L, Deschamps J, Zeilhofer HU. Hoxb8-Cre mice: A tool for brain-sparing conditional gene deletion. Genesis 2010;48(10):596-602.

[71] Xie DJ, Uta D, Feng PY, Wakita M, Shin MC, Furue H, Yoshimura M. Identification of 5-HT receptor subtypes enhancing inhibitory transmission in the rat spinal dorsal horn in vitro. Mol Pain 2012;8:58.

[72] Yesilyurt O, Seyrek M, Tasdemir S, Kahraman S, Deveci MS, Karakus E, Halici Z, Dogrul A. The critical role of spinal 5-HT7 receptors in opioid and non-opioid type stress-induced analgesia. Eur J Pharmacol 2015;762:402-410.

[73] Zhang L, Sykes KT, Buhler AV, Hammond DL. Electrophysiological heterogeneity of spinally projecting serotonergic and nonserotonergic neurons in the rostral ventromedial medulla. J Neurophysiol 2006;95(3):1853-1863.

[74] Zhang W, Gardell S, Zhang D, Xie JY, Agnes RS, Badghisi H, Hruby VJ, Rance N, Ossipov MH, Vanderah TW, Porreca F, Lai J. Neuropathic pain is maintained by brainstem neurons co-expressing opioid and cholecystokinin receptors. Brain 2009;132(Pt 3):778-787.

[75] Zhang Y, Zhao S, Rodriguez E, Takatoh J, Han BX, Zhou X, Wang F. Identifying local and descending inputs for primary sensory neurons. J Clin Invest 2015;125(10):3782-3794.

A CTb

B TPH2 CTb NeuN

NRM

LPGi

C

Ctb+TPH2+ cell (%)

LPGi NRM other

A SAD.RabiesΔG.GFP (SAD-G)

● GFP-expressing neuron

D Helper virus + SAD.RabiesΔG.GFP(EnvA) (2 weeks later)

Hoxb8-Cre; ROSA26^TVA

○ starter cell
● rabies-infected starter cell
● monosynaptically traced cell

B TPH2 GFP

LPGi

C % GFP+ cells expressing TPH2

days post injection

E NRM

F % GFP+ cells expressing TPH2

SAD.RabdG-eGFP   EnvA.RabdG-eGFP

A AAV8.hsyn.flex.eGFP

TPH2::cre

B TPH2 eGFP NeuN

eGFP TPH2

C
%eGFP cells expressing TPH2

D AAV2retro.flex. AAV9.FRT.tdTOM
FLPo.BFP

TPH2::cre

E
hindbrain
cervical
thoracic
lumbar
sacral

F AAV8.hsyn.flex.tdTOM

TPH2::cre

G tdTOM

H tdTOM 5-HT NeuN   tdTOM
cervical
lumbar

I tdTOM 5-HT NeuN   tdTOM
cervical
lumbar

A

20pA
0.5s

20mV
0.5s

20mV
50ms

10ms

NRM   LPGi

B

APs
20
10

LPGi
NRM

-50   50   100
Current injected (pA)

C   tdTOM Biocytin

NRM   LPGi

D
Resting membrane potential (mV)
0
-20
-40
-60
-80
-100
LPGi   NRM

E
Action potential threshold (mV)
0
-10
-20
-30
-40
-50
LPGi   NRM

F
Action potential amplitude (mV)
LPGi   NRM

G
Action potential half width (ms)
3
2
1
0
LPGi   NRM

H
slow afterhyperpolarisation amplitude (mV)
35
30
25
20
LPGi   NRM

A

B

CONTRA

A

AAV8.flex.hM3Dq.mCherry

TPH2::cre

● hM3Dq.mCherry-expressing TPH2::cre neuron

B mCherry TPH2 NeuN

C mCherry 5-HT

D

Hargreaves 🌡️

Cold plantar ❄️

von Frey

Rotarod

Brain region	Subset	GFP only	Ctb only	GFP+Ctb+	%GFP only	%Ctb only	%GFP+Ctb+
RVM (all neurons)	-	99 (11 - 57)	679 (149 – 319)	279 (65 – 134)	9.1 (4.0 – 12.1)	65.6 (59.7 – 73.3)	25.3 (22.7 – 28.1)
RVM (TPH2+ neurons)	All	5 (0 – 4)	269 (63 – 123)	62 (12 – 25)	1.5 (0 – 5.1)	80.1 (76.1 – 83.1)	18.5 (15.2 – 22.9)
	LPGi	5 (0 – 4)	79 (16 – 44)	35 (7 – 15)	4.2 (0 – 14.8)	66.4 (54.3 – 77.2)	29.4 (22.8 – 42.9)
	NRM	0 (0 – 0)	143 (37 – 63)	15 (4 -6)	0 (0 – 0)	90.5 (87.8 – 94.0)	9.5 (6.0 – 12.2)
	Other	0 (0 – 0)	47 (10 – 21)	12 (0 – 8)	0 (0 – 0)	79.7 (66.7 – 100)	20.3 (0 – 33.3)

2 Table 1: Quantification of hindbrain neurons traced from the spinal dorsal horn with AAV2retro.eGFP and CTb. The number in each column is

3 the total number of cells counted for each group, with the range of cells counted for each animal in parentheses. For percentages, the average

4 value is presented with the range for each animal given in parentheses.

1
2
3

labelling strategy	rabies virus	animals	total GFP+ cells counted in RVM	GFP neurons expressing TPH2	GFP neurons not expressing TPH2	% GFP+ neurons that are TPH2+
direct infection (5d)	SAD.RabiesΔG-eGFP (SAD-G)	3	25 (5-11)	11 (2-5)	14 (3-6)	44 (40-45)
direct infection (7d)	SAD.RabiesΔG-eGFP (SAD-G)	3	36 (7-17)	4 (1-2)	32 (6-16)	11 (6-16)
monosynaptic tracing	SAD.RabiesΔG-eGFP (EnvA)	4	33 (3-20)	1 (0-1)	32 (3-19)	3 (0-5)

4   Table 2: Neurons in the ventral hindbrain traced from the spinal cord with modified rabies viruses. Number represents the total number of cells

5   counted with the range of cells counted for each animal in parentheses. Percentages of labelled neurons expressing TPH2 are the average for

6   all cells counted per group, with the range for each animal given in parentheses.

GFP (all)	GFP (NRM)	GFP (LPGi)	GFP+ TPH2+ (all)	GFP+TPH2+ (NRM)	GFP+TPH2+ (LPGi)
127 (24-63)	25 (7 – 12)	102 (17 – 57)	96 (22 – 43)	25 (7 – 12)	71 (15 – 37)

2   Table 3: Cell counts for eGFP-labelled cells in the hindbrain from injection of AAV2retro.flex.eGFP into the spinal cord of TPH2::cre animals (n

3   = 3) the total number of cells counted is indicated and the range of cells counted per animal is indicated in parentheses.

Fluorophore	Total	LPGi	medial	Other
eGFP only	94 (19 – 29)	18 (1 – 8)	38 (6 – 16)	38 (5 – 20)
eGFP+mCherry+	558 (72 – 224)	298 (50 – 96)	156 (13 – 69)	104 (6 – 59)
mCherry only	153 (22 – 65)	71 (11 – 23)	55 (4 – 29)	27 (3 – 13)

1   Table 4: Cell counts for neurons labelled in the hindbrain from intersectional labelling experiments, from spinal cord injection of

2   AAV2retro.flex.FLPo.mCherry and hindbrain injection of AAV9.FRT.eGFP (n = 4 animals). Total number of cells counted is given with the range

3   of cells counted in parentheses.

Parameter	NRM	LPGi
Membrane resistance (M□)	1643 ± 240.1	1857 ± 316.9
Resting membrane potential (mV)	-64.69 ± 4.145	-70.32 ± 3.592
Rheobase current (pA)	18.75 ± 2.236	15.00 ± 2.266
Action potential threshold (mV)	-32.45 ± 2.597	-34.18 ± 0.8890
Action potential amplitude (mV)	85.26 ± 3.230	79.14 ± 4.194
Action potential half width (ms)	1.730 ± 0.1960	1.875 ± 0.2198
Slow Afterhyperpolarisation (with AHP / total)	9/10	5/8
Slow Afterhyperpolarisation Amplitude (mV)	27.09 ± 0.7983	27.58 ± 1.304

2 Table 5: Comparison of active and passive membrane properties between serotonergic

3 neurons in the NRM and the LPGi. Data are shown as mean ± SEM (n = 10 cells from 4

4 animals NRM , n= 8 cells from 3 animals LPGi)

Mouse line	Supplier/source	Reference
TPH2::cre	The Jackson Laboratory	MGI: 5435520
Hoxb8-cre	Pawel Pelczar	(Witschi et al. 2010)
ROSA26^{TVA}	Dieter Sauer	(Seidler et al. 2008)

Table 6: Transgenic mouse lines used in this study

Seidler, B., A. Schmidt, U. Mayr, H. Nakhai, R. M. Schmid, G. Schneider & D. Saur (2008) A

Cre-loxP-based mouse model for conditional somatic gene expression and

knockdown in vivo by using avian retroviral vectors. *Proc Natl Acad Sci U S A,* 105**,**

10137-42.

Witschi, R., T. Johansson, G. Morscher, L. Scheurer, J. Deschamps & H. U. Zeilhofer (2010)

Hoxb8-Cre mice: A tool for brain-sparing conditional gene deletion. *Genesis,* 48**,** 596-

602.

Virus/tracer name	Full name	Supplier/Source	Cat#
AAV2retro.eGFP	ssAAV-retro/2-CAG-EGFP-WPRE-SV40p(A)	Viral Vector facility UZH/ETHZ	V24-retro
AAV2retro.flex.eGFP	ssAAV-retro/2-shortCAG-dlox-EGFP(rev)-dlox-WPRE-SV40p(A)	Viral Vector facility UZH/ETHZ	V158-retro
AAV2retro.flex.tdTomato	ssAAV-retro/2-shortCAG-dlox-tdTomato(rev)-dlox-WPRE-hGHp(A)	Viral Vector facility UZH/ETHZ	V167-retro
AAV2retro.flex.FLPo.BFP	ssAAV-retro/2-hSyn1-chI-dlox-EBFP2_2A_FLPo(rev)-dlox-WPRE-SV40p(A)	Viral Vector facility UZH/ETHZ	V175-retro
AAV2retro.flex.FLPo.mCherry	ssAAV-retro/2-hSyn1-chI-mCherry_2A_FLPo-WPRE-SV40p(A)	Viral Vector facility UZH/ETHZ	V173-retro
AAV9.flex.ChR2-YFP	ssAAV-9/2-hEF1a-dlox-hChR2(H134R)_EYFP(rev)-dlox-WPRE-hGHp(A)	Viral Vector facility UZH/ETHZ	v214-9
AAV8.FRT.tdTomato	ssAAV-8/2-hSyn1-dlox-tdTomato(rev)-dlox-WPRE-bGHp(A)	Viral Vector facility UZH/ETHZ	v284-8
AAV9.FRT.eGFP	ssAAV-9/2-hSyn1-chI-dFRT-EGFP(rev)-dFRT-WPRE-	Viral Vector facility UZH/ETHZ	V335-9

	hGHp(A)		
AAV9.FRT.hM3D(q).mCherry	ssAAV-9-hSyn1-dFRT-hM3D(Gq)-mCherry	Viral Vector facility UZH/ETHZ	V189-9
AAV9.FRT.hM4D(i).mCherry	ssAAV-9/2-hSyn1-dFRT-hM4D(Gi)_mCherry(rev)-dFRT-WPRE-hGHp(A)	Viral Vector facility UZH/ETHZ	V190-9
AAV9.FRT.ChR2-YFP	ssAAV-9/2-hSyn1-dFRT-hM4D(Gi)_mCherry(rev)-dFRT-WPRE-hGHp(A)	Viral Vector facility UZH/ETHZ	V190-9
AAV9/2.FRT.eGFP.TeTxLC	ssAAV-9/2-hSyn1.chl-dFRT-EGFP-2A-FLAG:TeTxLC(rev)-dFRT-WPRE-hGHp(A)	Viral Vector facility UZH/ETHZ	v450-9
SAD pseudotyped rabies	SAD.RabiesDG-eGFP (SAD-G)	Karl-Klaus Conzelmann	N/A
EnvA pseudotyped rabies	SAD.RabiesDG-GFP (EnvA)	Karen Haenraets	N/A
CTb	Cholera Toxin b subunit	Sigma Aldrich	C9903-.5MG/

2   Table 7: AAVs, rabies viruses, and retrograde tracers used in the study

Antibody	Host	Supplier/Source	Cat#/RRID	Dilution
GFP	Chicken	LifeTech	A10262/ AB_2619988	1:1000
TPH2	Rabbit	Novus Biologicals	NB100-74555/AB_572263	1:1000
Ctb	Goat	LIST biological laboratories inc.	#703/AB_2314252	1:1000
mCherry	Goat	Sicgen	AB0081-200/AB_2333094	1:500
tdTomato	Goat	Sicgen	AB8181-200/ AB_2722750	1:500
5-HT	Rabbit	ImmunoStar	20080/AB_572263	1:1000
NeuN	Guinea pig	Synaptic systems	266004/AB_2619988	1:1000
Chicken-Alexa 488	Donkey	Jackson ImmunoResearch	703-546-155/AB_2340376	1:500
Goat-Cy3	Donkey	Jackson ImmunoResearch	705-166-147/AB_2340413	1:500

Goat-Alexa 488	Donkey	Jackson ImmunoResearch	795-546-147/ -	1:500
Guinea pig-Alexa 647	Donkey	Jackson ImmunoResearch	706-496-148/ -	1:500
Rabbit-Alexa 647	Donkey	Jackson ImmunoResearch	711-607-003/AB_2340626	1:500
Rabbit-Cy3	Donkey	Jackson ImmunoResearch	711-165-152/AB_2307443	1:500

2   Table 8: Antibodies used in the study