Interface resistance of biomolecular condensates

Yaojun Zhang[1,2,3]*, **Andrew G.T. Pyo**[4], **Yoyo Jiang**[2], **Clifford P. Brangwynne**[5,6], **Howard A. Stone**[7], **Ned S. Wingreen**[8,9]*

*For correspondence:
yaojunz@jhu.edu (YZ);
wingreen@princeton.edu (NSW)

[1]Center for the Physics of Biological Function, Princeton University, Princeton, United States; [2]Department of Physics and Astronomy, Johns Hopkins University, Baltimore, United States; [3]Department of Biophysics, Johns Hopkins University, Baltimore, United States; [4]Department of Physics, Princeton University, Princeton, United States; [5]Department of Chemical and Biological Engineering, Princeton University, Princeton, United States; [6]Howard Hughes Medical Institute, Chevy Chase, United States; [7]Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, United States; [8]Department of Molecular Biology, Princeton University, Princeton, United States; [9]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, United States

Abstract   A hallmark of biomolecular condensates formed via liquid-liquid phase separation is that they dynamically exchange materials with their surroundings, and this can be crucial to condensate function. How is this rate of exchange controlled? Intuitively, the rate can be limited by the flux from the dilute phase or by the mixing speed in the dense phase. Surprisingly, recent experiments suggest that the exchange rate can instead be limited by the dynamics of molecules at the droplet interface, implying the existence of an "interface resistance". We combine theory and simulation to show that interface resistance can arise when incident molecules transiently touch the interface without bonding to the dense phase, i.e., the molecules "bounce" from the interface. This occurs when the molecules can adopt conformations that limit the accessibility of their sticky regions. Our work highlights the underappreciated role of interface resistance, with implications for both natural and synthetic condensates.

Introduction

The interior of cells is organized in both space and time by biomolecular condensates, which form and dissolve as needed *Shin and Brangwynne* (*2017*); *Banani et al.* (*2017*). These condensates play key roles in processes ranging from transcription to translation, metabolism, signaling, and more *An et al.* (*2008*); *Su et al.* (*2016*); *Sabari et al.* (*2018*); *Formicola et al.* (*2019*). The complex interactions among their components endow condensates with distinct physical properties, including low surface tension, viscoelasticity, aging, etc. These distinct properties are crucial to the ability of condensates to carry out their unique biological functions. Here, we focus on one important physical property of condensates – the rate of exchange of material between condensed and dilute phases. This rate of exchange can impact biochemical processes taking place in condensates by limiting the escape of completed products, such as ribosomes produced in nucleoli *Yao et al.* (*2019*), or limiting the availability of components or regulatory molecules (e.g., snoRNAs and ribosomal proteins entering nucleoli, or mRNAs entering P bodies or stress granules). The rate of exchange can also control the dynamical response of condensates to a changing environment, and, as exchange

between dense and dilute phase is central to coarsening via Ostwald ripening, it can regulate the
number, size, and location of condensates within the cell.
The material exchange between a condensate and the surrounding dilute phase can be probed
via FRAP experiments, a commonly used approach for measuring condensate fluidity and molecu-
lar diffusion coefficients. Exchange dynamics are thus readily measurable and have been reported
for a variety of systems *Li et al.* (*2012*); *Patel et al.* (*2015*); *Burke et al.* (*2015*); *Banani et al.* (*2016*);
Jain et al. (*2016*); *Aumiller Jr et al.* (*2016*). However, only a very limited number of studies *Taylor*
et al. (*2019*); *Hubatsch et al.* (*2021*); *Bo et al.* (*2021*); *Folkmann et al.* (*2021*); *Lee* (*2021*) aimed to
understand what controls the timescales of condensate component exchange. Briefly, Taylor *et*
al. *Taylor et al.* (*2019*) combined FRAP experiments on condensates *in vitro* and *in vivo* with differ-
ent theoretical models to examine the impact of model choice on the physical parameters derived
from data fitting. Folkmann *et al.* *Folkmann et al.* (*2021*) and Lee *Lee* (*2021*) proposed that the rate
of molecular absorption to the condensate can be "conversion-limited" instead of diffusion-limited
and established a mathematical framework for the temporal evolution of droplet sizes in this limit.
In all these cases, the modeling of interface resistance *Taylor et al.* (*2019*) or conversion-limited
material transfer *Folkmann et al.* (*2021*); *Lee* (*2021*) was conducted at the phenomenological level,
without aiming to understand the underlying physical mechanism that gives rise to interface resis-
tance. Hubatsch *et al.* *Hubatsch et al.* (*2021*) and Bo *et al.* *Bo et al.* (*2021*) tackled the exchange
dynamics problem by developing, respectively, a continuum theory of macroscopic phase separa-
tion and a stochastic Langevin equation of single-molecule trajectories. However, the mean-field
approaches in *Hubatsch et al.* (*2021*) and *Bo et al.* (*2021*) neglect the potentially complex dynamics
of molecules at the condensate interface, which can slow down material exchange significantly as
shown by *Taylor et al.* (*2019*); *Folkmann et al.* (*2021*).
In the following, we first derive an analytical expression for the timescale of condensate material
exchange, which conveys a clear physical picture of what controls this timescale. We then utilize a
"sticker-spacer" polymer model to investigate the mechanism of interface resistance. We find that
a large interface resistance can occur when molecules bounce off the interface rather than being
directly absorbed.

Results

Mathematical formulation of exchange dynamics

The exchange of molecules between a condensate and the dilute phase can be investigated through
FRAP-type experiments in which, e.g., fluorescence is locally bleached and recovery as a function
of time recorded (Figure 1A and B). Theoretically, the time evolution of the concentration profile
$c(r, t)$ of the molecules initially located in a spherical condensate of radius R (bleached population)
can be described by the following continuum diffusion equations *Taylor et al.* (*2019*):

$$\frac{\partial}{\partial t} c(r, t) = D_{\text{den}} \left(\frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} \right) c(r, t), \quad r < R;$$

$$\frac{\partial}{\partial t} c(r, t) = D_{\text{dil}} \left(\frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} \right) c(r, t), \quad r > R, \tag{1}$$

72 with the initial condition:

$$c(r, 0) = \begin{cases} c_{\text{den}}, & r < R; \\ 0, & r > R, \end{cases} \tag{2}$$

and boundary conditions:

$$\frac{\partial}{\partial r} c(r, t) \bigg|_{r=0} = c(+\infty, t) = 0;$$

$$-D_{\text{den}} \frac{\partial}{\partial r} c(r, t) \bigg|_{r=R_-} = -D_{\text{dil}} \frac{\partial}{\partial r} c(r, t) \bigg|_{r=R_+} = \kappa \left[c(R_-, t) - \frac{c_{\text{den}}}{c_{\text{dil}}} c(R_+, t) \right], \tag{3}$$

73 where D_{den} and D_{dil} are the diffusion coefficients of molecules in the dense and dilute phases,
74 and c_{den} and c_{dil} are the equilibrium concentrations in the dense and dilute phases. The second

Figure 1. FRAP experiments on droplets. (A) Schematic of a FRAP experiment in which an entire droplet is photobleached and the recovery of fluorescence is recorded. Material exchange between the condensate and the surrounding dilute phase can be limited by the flux of unbleached molecules coming from the dilute phase, the speed of internal mixing in the dense phase, or the flux passing through the interface. (B) The recovery time τ is defined as the time required for fluorescence to return 63% (i.e., $1 - 1/e$) of the way back to its original level. (C) Experimental data from Ref. *Taylor et al.* (*2019*) in which a LAF-1 droplet of radius $R = 1\,\mu$m recovers from photobleaching. (Left) Images before bleaching, immediately after bleaching of the entire droplet region, and at two subsequent times. (Right) Expected recovery time $\tau_{\mathrm{dil}} = c_{\mathrm{den}} R^2 / (3 c_{\mathrm{dil}} D_{\mathrm{dil}}) = 4\,$s and $\tau_{\mathrm{den}} = R^2 / (\pi^2 D_{\mathrm{den}}) = 60\,$s if the slowest recovery process was the flux from the dilute phase or diffusion within the droplet, respectively, with $D_{\mathrm{den}} = 0.0017\,\mu$m^2/s, $D_{\mathrm{dil}} = 94\,\mu$m^2/s, and $c_{\mathrm{den}}/c_{\mathrm{dil}} = 1190$ taken from *Taylor et al.* (*2019*). While the timescale associated with interface resistance τ_{int} is unknown, the measured recovery time $\tau \sim 1$ hour is much longer than τ_{dil} and τ_{den}, suggesting the recovery is limited by flux through the interface, with an interface conductance of $\kappa = R/(3\tau) \simeq 10^{-4}\,\mu$m/s.

boundary condition corresponds to flux balance at the interface of the condensate. Specifically, the flux exiting the dense phase (left) equals the flux entering the dilute phase (middle) and also equals the flux passing through the interface (right).

To understand the physical origin of the last term in the second boundary condition in Equation (3), we note that the net outward flux across the interface can be written as $k_- c(R_-, t) - k_+ c(R_+, t)$, where $k_{+/-}$ denotes the entering/exiting rate of molecules at the interface and $c(R_{+/-}, t)$ the concentration of bleached molecules immediately outside/inside of the boundary. At thermal equilibrium, this net flux goes to zero, i.e., $k_- c_{\mathrm{den}} = k_+ c_{\mathrm{dil}}$ so $k_+ = k_-(c_{\mathrm{den}}/c_{\mathrm{dil}})$. The net outward flux is therefore $k_- \left[c(R_-, t) - (c_{\mathrm{den}}/c_{\mathrm{dil}}) c(R_+, t) \right]$. The parameter $\kappa \equiv k_-$ is a transfer coefficient that governs the magnitude of this net flux. When the ratio of the concentrations on the two sides of the interface deviates from the equilibrium ratio, a small κ can kinetically limit the flux going through the interface. We therefore term κ the interface conductance, the inverse of interface resistance.

For the model described by Equations (1)–(3), the fraction of molecules in the condensate which are unbleached at time t is

$$f(t) = 1 - \frac{\int_0^R 4\pi r^2 c(r, t)\, dr}{\int_0^R 4\pi r^2 c_{\mathrm{den}}\, dr}. \tag{4}$$

Clearly, how quickly $f(t)$ recovers from 0 to 1 quantifies the timescale of material exchange between the condensate and the surrounding dilute phase.

Timescale of condensate component exchange

The authors of *Taylor et al.* (*2019*) derived an exact solution for $f(t)$ in an integral form using Laplace transforms. However, it is not directly apparent from the integral expression what physics governs the timescale of fluorescence recovery. In addition, the lengthy integral form of the expression also presents an impediment to its practical experimental applications. To obtain a more intuitive and concise result, we note that diffusion of biomolecules in the dilute phase is typically much faster

97 than diffusion in the dense phase, with measured D_{dil}/D_{den} in the range of 10^2-10^5 *Rosenzweig et al.*
98 (*2017*); *Taylor et al.* (*2019*). We therefore employed the exact solution to derive an approximate
99 solution in the parameter regime $D_{dil} \gg D_{den}$:

$$f(t) = 1 - \exp\left(-\frac{t}{\tau}\right), \tag{5}$$

100 where the timescale of fluorescence recovery is given by:

$$\tau = \frac{R^2}{\pi^2 D_{den}} + \frac{c_{den}R^2}{3c_{dil}D_{dil}} + \frac{R}{3\kappa}. \tag{6}$$

Please refer to Appendix 1 for a detailed derivation. We note that, in practice, $D_{dil} > 20D_{den}$ is
sufficient for the validity of the approximation with the approximate τ in Equation (6) always within
10% of the exact value.

Equation (6) conveys a clear physical picture of what controls the timescale of condensate ma-
terial exchange. First, for large condensates and slow internal diffusion, exchange is limited by
the rate of mixing within the condensate, so that $\tau \simeq R^2/(\pi^2 D_{den})$. Second, if instead diffusion
in the dilute phase is sufficiently slow, or the concentration in the dilute phase is very low, then
$\tau \simeq c_{den}R^2/(3c_{dil}D_{dil})$, which is the time required to replace all molecules in the condensate if
molecules incident from the dilute phase are immediately absorbed (see Appendix 1). Finally, if
the interface conductance κ is very small, the interfacial flux can be rate limiting for exchange,
yielding $\tau \simeq R/(3\kappa)$.

Interface conductance κ from mean-field theory is contradicted by experimental results

What determines the magnitude of the interface conductance κ? From a theoretical perspective,
transitions between dense and dilute phases have been modeled both from the perspective of
continuum theory *Hubatsch et al.* (*2021*) and by considering single-molecule trajectories *Bo et al.*
(*2021*). However, for any particular systems, the magnitude of the interface conductance depends
on microscopic features of the biomolecules, such as internal states, which may not be captured
by Flory-Huggins and Cahn-Hilliard-type mean-field theories. Indeed, if we start with the contin-
uum approach in *Hubatsch et al.* (*2021*), where the concentration of bleached components $c(\mathbf{r},t)$
is governed by

$$\frac{\partial c(\mathbf{r},t)}{\partial t} = \nabla \cdot \left\{ D\left[c_{eq}(\mathbf{r})\right] \left[\nabla c(\mathbf{r},t) - c(\mathbf{r},t)\frac{\nabla c_{eq}(\mathbf{r})}{c_{eq}(\mathbf{r})}\right] \right\} \tag{7}$$

122 with $c_{eq}(\mathbf{r})$ the equilibrium concentration profile and $D[c_{eq}(\mathbf{r})]$ the diffusion coefficient which de-
123 pends on the local equilibrium concentration, one can obtain an expression for κ (see Appendix
124 1):

$$\kappa^{-1} = \int \frac{c_{den}}{c_{eq}(r)D(r)}dr, \tag{8}$$

where the integral is over the interface region. We would then conclude $\kappa^{-1} < \delta/D_{den} + \delta c_{den}/(c_{dil}D_{dil})$,
where δ is the width of the interface. As the interface is typically narrow, this inequality would imply
that in practice the interfacial term in Equation (6) would always be smaller than the sum of the
other two terms, and thus could be neglected.

However, a recent FRAP experiment on LAF-1 protein droplets *Taylor et al.* (*2019*) contradicts
the above mean-field result. In the experiment, a micron-sized LAF-1 droplet ($R = 1\,\mu$m) was
bleached and fluorescence recovery measured as a function of time (Figure 1C). It was observed
that recovery of that droplet occurs on a timescale of ~ 1 hour. Given the measured parame-
ters of the system, one can estimate the recovery time in the mean-field approach to be $\tau =$
$R^2/(\pi^2 D_{den}) + c_{den}R^2/(3c_{dil}D_{dil}) = 64\,$s, much shorter than the measured recovery time. A large in-
terface resistance was proposed as a possible explanation for this discrepancy *Taylor et al.* (*2019*).
However, the underlying mechanism that leads to such an unexpectedly large interface resistance
remains unclear. More generally, how does interface resistance depend on the microscopic fea-
tures of phase-separating molecules?

Coarse-grained simulation of "sticker-spacer" polymer phase separation

Motivated by the question of what could give rise to interface resistance, we employed a "sticker-spacer" polymer model *Choi et al.* (*2020*); *Semenov and Rubinstein* (*1998*) to explore the exchange of molecules between a condensate and the dilute phase. The "sticker-spacer" model provides a conceptual framework for understanding biomolecular phase separation, wherein the "stickers" represent residues or larger domains that are capable of forming saturable bonds, while the "spacers" connect the stickers to form polymers. Specifically, we simulated polymers consisting of type A and type B stickers connected by implicit spacers in the form of stretchable bonds *Kremer and Grest* (*1990*) (Figure 2A):

$$U_b(r) = -\frac{1}{2}KR_0^2 \ln\left[1 - \left(\frac{r}{R_0}\right)^2\right], \quad r < R_0, \tag{9}$$

where r is the distance between two stickers. One-to-one heterotypic bonds between A and B are implemented via an attractive potential:

$$U_a(r) = -\frac{1}{2}U_0\left(1 + \cos\frac{\pi r}{r_0}\right), \quad r < r_0, \tag{10}$$

while stickers of the same type interact through a repulsive potential to prevent many-to-one binding:

$$U_r(r) = 4\epsilon\left[\left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6\right] + \epsilon, \quad r \leq r_c. \tag{11}$$

We take $K = 0.15\,k_B T/\text{nm}^2$, $R_0 = 10\,\text{nm}$, $U_0 = 14\,k_B T$, $r_0 = 1\,\text{nm}$, $\epsilon = 1\,k_B T$, $\sigma = 2\,\text{nm}$, and $r_c = 1.12\sigma$ in all simulations, except in the simulations of Figure 3F where we vary U_0 systematically from 13.5 to $15\,k_B T$. For all simulation results we reported below, the standard error of the mean is smaller than the symbol size and therefore not shown.

For each of the five sequences shown in Figure 2A, we simulated 1000 polymers in a $500\,\text{nm} \times 50\,\text{nm} \times 50\,\text{nm}$ box with periodic boundary conditions using Langevin dynamics (see Appendix 2 for details). Simulations were performed using LAMMPS molecular-dynamics simulator *Plimpton* (*1995*). Figure 2B shows a snapshot of coexisting dense and dilute phases after equilibration of the A6B6 polymers (6 A stickers followed by 6 B stickers), while Figure 2C shows the time-averaged profile of the total polymer concentration. The five different polymer sequences we simulated were chosen to yield a range of dilute- and dense-phase sticker concentrations (Figure 2D) as well as a range of dilute- and dense-phase diffusion coefficients (Figure 2E). As found previously *Weiner et al.* (*2021*), polymers like A6B6 with long blocks of stickers of the same type have low dilute-phase concentrations. This follows because it is entropically unfavorable for these polymers to form multiple self-bonds, which favors the dense phase where these polymers can readily form multiple trans-bonds. These long-block polymers also have low dense-phase diffusion coefficients because of their large number of trans-bonds, which need to be repeatedly broken for the polymers to diffuse.

Interface conductance κ from simulations

Having determined the concentrations and diffusion coefficients in the dense and dilute phases, we are now in a position to extract the values of interface conductance from simulations. Figure 3 depicts a simple protocol that allows us to infer κ by applying the 1D, slab-geometry version of Equations (1)–(6) to simulation results (see Appendices 1 and 2 for details): (i) All polymers in the dilute phase are initially considered "labeled", (ii) any labeled polymer that forms a lasting A-B bond with a polymer in the dense phase becomes permanently unlabeled (Figure 3A), (iii) the remaining fraction of labeled dilute phase polymers is fit to an exponential decay (Figure 3B), and (iv) the resulting decay time constant τ is used together with the known dense and dilute phase parameters to infer κ from:

$$\kappa = \frac{c_{dil}}{c_{den}}\sqrt{\frac{D_{dil}}{\tau}}\tan\frac{d}{\sqrt{\tau D_{dil}}}, \tag{12}$$

Figure 2. Coarse-grained molecular-dynamics simulations of multivalent phase-separating polymers. (A) Each polymer is composed of monomers ("stickers") of type A (blue) or B (red), and modeled as a linear chain of spherical particles each with a diameter of 2 nm, connected by stretchable bonds with an equilibrium length of 3.9 nm. Stickers of different types have an attractive interaction, while stickers of the same type interact repulsively, ensuring one-to-one binding between the A and B stickers. (B) Snapshot of a simulation of 1000 A6B6 polymers in a 500 nm × 50 nm × 50 nm box with periodic boundary conditions. The system undergoes phase separation into a dense phase (middle region) and a dilute phase (two sides), driven by the one-to-one A-B bonds. (C) Polymer concentration profile for the simulation in (B) with the center of the dense phase aligned at $x = 0$ and averaged over time and over ten simulation repeats. (D) Average total polymer concentrations in the dense (top) and dilute (bottom) phases from simulations of the five types of polymers shown in (A). (E) Polymer diffusion coefficients in the dense (top) and dilute (bottom) phases. All simulations were performed and snapshots were obtained using LAMMPS *Plimpton* (*1995*). Please refer to Appendix 2 for simulation details.

where d is the half-width of the dilute phase. As shown in Figure 3C, the resulting values of κ span
more than an order of magnitude for our selected polymer sequences, despite the fact that all five
polymers can in principle form the same number (6) of self-bonds.

We note that one can alternatively obtain κ by directly measuring the flux of molecules that
enter the dense phase. Mathematically, this flux equals $k_+ c_{\text{dil}} = \kappa c_{\text{den}}$. We show in Appendix 2 that
the values of κ found via this method are consistent with results reported in Figure 3C.

"Bouncing" of molecules can lead to large interface resistance

187   What gives rise to the very different values of κ? To address this question, we first consider the
188   predicted interface conductance κ_0 if polymers incident from the dilute phase simply move through
189   the interface region with a local diffusion coefficient that crosses over from D_{dil} to D_{den}. Then
190   according to Equation (8) (see Appendix 1)

$$\kappa_0 = \frac{c_{\text{dil}} D_{\text{dil}}}{\delta c_{\text{den}}}. \tag{13}$$

191   However, as shown in Figure 3D, the actual values of κ in our simulations can be a factor of ~ 50

Figure 3. Determination of interface conductance from simulations. (A) Illustration of simulation protocol: At $t = 0$ only polymers in the dilute phase are "labeled" (solid balls), any polymer that enters the dense phase (forms an A-B bond lasting >10 times the average bond lifetime of an isolated A-B pair) becomes permanently "unlabeled" (hollow balls). (B) Fraction of labeled polymers in the dilute phase as a function of time for simulations of the five types of polymers shown in Figure 2A. (C) Decay time of labeled polymers from exponential fits to curves in (B) (top), and corresponding calculated values of interface conductance κ (bottom). (D) For all simulated polymers, interface conductance scaled by κ_0 [Equation (13)] is approximately a linear function of a parameter u which reflects the fraction of unbound stickers in the dense and dilute phases. Inset illustration: Polymers in the dilute phase with few or no unbound stickers may "bounce" off the dense phase, which contributes to the interface resistance. (E) Example of simulated trajectory in which a dilute-phase A6B6 polymer "bounces" multiple times before finally joining the dense phase. (F) Interface conductance κ of A6B6 system as a function of binding strength U_0 between A-B stickers.

smaller than κ_0. This reduction can be traced to a "bouncing" effect. As shown schematically in the
inset to Figure 3D and for an exemplary simulated trajectory in Figure 3E (more trajectories can
be found in Appendix 2), molecules incident from the dilute phase may fail to form bonds with the
dense phase, effectively "bouncing" off of the condensate. The differing extent of this bouncing
effect for the five sequences we studied reflects differences in their numbers of free stickers in
both their dilute- and dense-phase conformations. The fewer such available stickers, the fewer
ways for a polymer incident from the dilute phase to bond with polymers at the surface of the
dense phase, and thus the more likely the incident polymer is to bounce. In support of this picture,
we find that all our simulation results for κ/κ_0 collapse as a linear function of a lumped parameter

$$u = \frac{4\pi r_0 \delta^2 n^2 c_{\text{den}}}{2 + s + s^{-1}}(f_{\text{dilA}} f_{\text{denB}} + f_{\text{dilB}} f_{\text{denA}}), \tag{14}$$

which expresses the availability of free stickers (see Appendix 1), where n is the number of monomers

in a polymer, s is the global stoichiometry (i.e., c_A/c_B), $f_{dilA/dilB}$ and $f_{denA/denB}$ are the fractions of un-
bound A/B monomers in the dilute and dense phases, all parameters are determined from simu-
lations.

Comparing sequences with unequal sticker stoichiometry A8B6 and A10B6 to their most closely
related equal-stoichiometry sequence A6B6, we find that the extra A stickers substantially increase
the interface conductance κ. Intuitively, the excess As in both dense and dilute phases of A8B6
and A10B6 provide a pool of available stickers for any unbound B to bind to. By contrast, at equal
stoichiometry, both free As and free Bs are rare which maximizes the bouncing effect. This reduc-
tion in potential binding partners at equal stoichiometry has also been observed experimentally
Brassinne et al. (*2017*) and theoretically *Ronceray et al.* (*2022*) to cause an anomalous slowing of
diffusion within condensates at equal stoichiometry in the regime of strong binding.

Finally, we expect the interface resistance to increase approximately exponentially with the
increase of binding strength U_0 between A-B stickers, as the tighter the binding, the fewer available
stickers, and hence the more bouncing of molecules at the interface. We demonstrate in Figure 3F
that the interface conductance κ of the A6B6 system indeed drops by a factor of 5 as the value of
U_0 increases from 13.5 to $15\,k_BT$.

Signatures of interface resistance

Under what circumstances is interface resistance experimentally measurable? If there were no
bouncing effect, i.e., if all molecules incident from the dilute phase that touch the interface get
immediately absorbed into the condensate, then interface resistance would never dominate the
recovery time in FRAP-type experiments, making it very difficult to measure κ. However, as shown
in Figure 3, the bouncing effect can reduce κ substantially. For such systems, the interface con-
ductance can be inferred quantitatively from Equation (6) or by fitting FRAP recovery curves as
in Taylor *et al.* *Taylor et al.* (*2019*), using the experimentally measured dense- and dilute-phase
concentrations and diffusion coefficients.

Even without knowing these parameters, one may still be able to infer the presence of a large
interface resistance by observing the pattern of fluorescence recovery in droplets of different sizes.
According to Equation (6) the recovery time associated with interface resistance increases linearly
with radius R while the other terms increase as R^2 (Figure 4A). Therefore, one expects a cross-over
for the recovery from being interface-resistance dominated (small R) to being either dilute-phase-
diffusion or dense-phase-mixing dominated (large R). In the latter case, the fluorescence profile
during recovery will be notably different in large versus small droplets as shown in Figure 4B –
for large droplets progressive diffusion of fluorescence into the droplet will be apparent, whereas
small droplets will recover uniformly as internal mixing will be fast compared to exchange with
the surroundings. Thus observation of such a cross-over of the recovery pattern as a function
of droplet size provides evidence for the presence of a large interface resistance, which can be
followed up by more quantitative studies. For example, the uniform recovery of the LAF-1 droplet
in Figure 1C is indicative of a large interface resistance, especially because the diffusion in the dilute
phase is too fast to be rate limiting.

Discussion

The dynamic exchange of condensate components with the surroundings is a key feature of mem-
braneless organelles, and can significantly impact condensate biological function. In this work, we
combined analytical theory and coarse-grained simulations to uncover physical mechanisms that
can control this exchange dynamics. Specifically, we first derived an analytical expression for the
exchange rate, which conveys the clear physical picture that this rate can be limited by the flux of
molecules from the dilute phase, by the speed of mixing inside the dense phase, or by the dynam-
ics of molecules at the droplet interface. Motivated by recent FRAP measurements *Taylor et al.*
(*2019*) that the exchange rate of LAF-1 droplets is limited by interface resistance, which contradicts

Figure 4. FRAP recovery patterns for large versus small droplets can be notably different for condensates with a sufficiently large interface resistance. (A) Expected relaxation time as a function of droplet radius for *in silico* "FRAP experiments" on the A6B6 system. The interface resistance dominates recovery times for smaller droplets, whereas dense-phase diffusion dominates recovery times for larger droplets. Parameters matched to the simulated A6B6 system: $c_{den} = 7.7\,mM$, $c_{dil} = 0.05\,mM$, $D_{den} = 0.013\,\mu m^2/s$, $D_{dil} = 17\,\mu m^2/s$, and $\kappa = 0.14\,\mu m/s$. (B) Time courses of fluorescence profiles for A6B6 droplets of radius $1\,\mu m$ (top) and $0.1\,\mu m$ (bottom); red is fully bleached, green is fully recovered.

predictions of conventional mean-field theory, we investigated possible physical mechanisms un-
derlying interface resistance using a "sticker-spacer" model. Specifically, we hypothesized that a
large interface resistance could be caused by incident molecules and/or molecules at the inter-
face adopting conformations that limit the accessibility of their sticky regions, and thus markedly
slowing down bonding of incident molecules to the dense phase. We subsequently demonstrated
via simulations a notable example in which incident molecules have formed all possible internal
bonds, and thus bounce from the interface, giving rise to a large interface resistance. Finally, we
discussed the signatures in FRAP recovery patterns of the presence of a large interface resistance.
What are the microscopic features of phase-separating molecules that can lead to a significant
interface resistance? The essential requirement is that molecules in the dilute phase and molecules
at the interface should not present "sticky" surfaces to each other. Since these same molecules
must be capable of sticking to each other in order to phase separate, a natural scenario is that these
molecules assume non-sticky conformations due to the shielding of interacting regions, e.g., burial
of hydrophobic residues in the core of a protein, or, in the scenario explored in the simulations,
the saturation of sticker-like bonds. Examples of systems with strong enough bonds to allow bond
saturation include SIM-SUMO *Banani et al.* (*2016*) and nucleic acids with strong intramolecular
base-pairing. Interestingly, a recent coarse-grained simulation of RNA droplets of $(CAG)_{47}$ *Nguyen*
et al. (*2022*) illustrated that a $(CAG)_{47}$ molecule in a closed hairpin conformation fails to integrate
into a droplet but rather bounces off the droplet interface. In the case of LAF-1, we note that the
values of interface conductance κ obtained in our simulations are a factor of 10^3 to 10^4 higher than
the experimentally measured κ for the LAF-1 droplet. While we do not aim to specifically simu-
late the LAF-1 system in this work and the value of κ in simulations can in principle be tuned by
adjusting the bond strength U_0, the large disparity between simulation and experiment renders
the mechanism responsible for the inferred large interface resistance in LAF-1 droplets unclear.
Nevertheless, the current study suggests at least one possibility: LAF-1 proteins may adopt con-
formations that significantly slow down the attachment of incident proteins, leading to frequent
bouncing of incident LAF-1s.
Biologically, the bouncing effect may influence the coarsening of condensates. The same in-
terface resistance that governs exchange between phases at equilibrium will control the flux of
material from the dilute phase to the dense phase during coarsening, so that bouncing will slow
down the coarsening process. Indeed, a recent theoretical study *Ranganathan and Shakhnovich*
(*2020*) of coarsening via mergers of small polymer clusters found anomalously slow coarsening dy-
namics due to exhaustion of binding sites, paralleling the single-polymer bouncing effect explored
here. Other mechanisms that may slow coarsening include the Pickering effect in which adsorbed
particles partially cover the interface, reducing the area accessible for exchange and also reduc-

ing overall condensate surface tension *Folkmann et al.* (*2021*). In this same study, additional slow
coarsening of PGL-3 condensates was attributed to a conversion-limited rather than a diffusion-
limited flux of particles from the dilute phase into the dense phase, which we hypothesize could
be a consequence of the bouncing effect. Interestingly, a conversion-limited flux has been shown
to lead to qualitatively distinct scaling of condensate size with time *Lee* (*2021*). As many conden-
sates dissolve and reform every cell cycle (or as needed) we anticipate that the bouncing effect
may constitute an additional means of regulating condensate dynamics.

Acknowledgments

This work was supported in part by the National Science Foundation, through the Center for the
Physics of Biological Function (PHY-1734030), NIH Grants R01 GM140032, U01 DA040601, the Howard
Hughes Medical Institute, and the Air Force Office of Scientific Research (FA9550-20-1-0241 to CPB).
YZ was partially supported by a startup fund at Johns Hopkins University. CPB and HAS were par-
tially supported by Princeton University's Materials Research Science and Engineering Center DMR-
1420541.

References

An S, Kumar R, Sheets ED, Benkovic SJ. Reversible compartmentalization of de novo purine biosynthetic com-
plexes in living cells. Science. 2008; 320(5872):103–106.

Aumiller Jr WM, Pir Cakmak F, Davis BW, Keating CD. RNA-based coacervates as a model for membraneless
organelles: formation, properties, and interfacial liposome assembly. Langmuir. 2016; 32(39):10042–10053.

Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of cellular biochemistry. Nat
Rev Mol Cell Biol. 2017; 18(5):285–298.

Banani SF, Rice AM, Peeples WB, Lin Y, Jain S, Parker R, Rosen MK. Compositional control of phase-separated
cellular bodies. Cell. 2016; 166(3):651–663.

Bo S, Hubatsch L, Bauermann J, Weber CA, Jülicher F. Stochastic dynamics of single molecules across phase
boundaries. Phys Rev Research. 2021; 3(4):043150.

Brassinne J, Cadix A, Wilson J, Van Ruymbeke E. Dissociating sticker dynamics from chain relaxation in
supramolecular polymer networks – The importance of free partner! J Rheol. 2017; 61(6):1123–1134.

Burke KA, Janke AM, Rhine CL, Fawzi NL. Residue-by-residue view of in vitro FUS granules that bind the C-
terminal domain of RNA polymerase II. Mol Cell. 2015; 60(2):231–241.

Choi JM, Holehouse AS, Pappu RV. Physical principles underlying the complex biology of intracellular phase
transitions. Annu Rev Biophys. 2020; 49:107–133.

Folkmann AW, Putnam A, Lee CF, Seydoux G. Regulation of biomolecular condensates by interfacial protein
clusters. Science. 2021; 373(6560):1218–1224.

Formicola N, Vijayakumar J, Besse F. Neuronal ribonucleoprotein granules: dynamic sensors of localized sig-
nals. Traffic. 2019; 20(9):639–649.

Hubatsch L, Jawerth LM, Love C, Bauermann J, Tang TD, Bo S, Hyman AA, Weber CA. Quantitative theory for
the diffusive dynamics of liquid condensates. eLife. 2021; 10:e68620.

Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A, Parker R. ATPase-modulated stress granules contain a
diverse proteome and substructure. Cell. 2016; 164(3):487–498.

Kremer K, Grest GS. Dynamics of entangled linear polymer melts: A molecular-dynamics simulation. J Chem
Phys. 1990; 92(8):5057–5086.

Langevin P. Sur la théorie du mouvement brownien [On the theory of Brownian motion]. C R Acad Sci Paris.
1908; 146:530–533.

Lee CF. Scaling law and universal drop size distribution of coarsening in conversion-limited phase separation.
Phys Rev Research. 2021; 3(4):043081.

330 **Li P**, Banjade S, Cheng HC, Kim S, Chen B, Guo L, Llaguno M, Hollingsworth JV, King DS, Banani SF, et al. Phase
331 transitions in the assembly of multivalent signalling proteins. Nature. 2012; 483(7389):336–340.

332 **Nguyen HT**, Hori N, Thirumalai D. Condensates in RNA repeat sequences are heterogeneously organized and
333 exhibit reptation dynamics. Nat Chem. 2022; 14(7):775–785.

334 **Patel A**, Lee HO, Jawerth L, Maharana S, Jahnel M, Hein MY, Stoynov S, Mahamid J, Saha S, Franzmann TM,
335 et al. A liquid-to-solid phase transition of the ALS protein FUS accelerated by disease mutation. Cell. 2015;
336 162(5):1066–1077.

337 **Plimpton S**. Fast parallel algorithms for short-range molecular dynamics. J Comput Phys. 1995; 117(1):1–19.

338 **Ranganathan S**, Shakhnovich EI. Dynamic metastable long-living droplets formed by sticker-spacer proteins.
339 eLife. 2020; 9:e56159.

340 **Ronceray P**, Zhang Y, Liu X, Wingreen NS. Stoichiometry controls the dynamics of liquid condensates of asso-
341 ciative proteins. Phys Rev Lett. 2022; 128(3):038102.

342 **Rosenzweig ESF**, Xu B, Cuellar LK, Martinez-Sanchez A, Schaffer M, Strauss M, Cartwright HN, Ronceray P,
343 Plitzko JM, Förster F, et al. The eukaryotic CO_2-concentrating organelle is liquid-like and exhibits dynamic
344 reorganization. Cell. 2017; 171(1):148–162.

345 **Sabari BR**, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM, Zamudio AV, Man-
346 teiga JC, et al. Coactivator condensation at super-enhancers links phase separation and gene control. Science.
347 2018; 361(6400):3958.

348 **Semenov AN**, Rubinstein M. Thermoreversible gelation in solutions of associative polymers. 1. Statics. Macro-
349 molecules. 1998; 31(4):1373–1385.

350 **Shin Y**, Brangwynne CP. Liquid phase condensation in cell physiology and disease. Science. 2017;
351 357(6357):eaaf4382.

352 **Su X**, Ditlev JA, Hui E, Xing W, Banjade S, Okrut J, King DS, Taunton J, Rosen MK, Vale RD. Phase separation of
353 signaling molecules promotes T cell receptor signal transduction. Science. 2016; 352(6285):595–599.

354 **Taylor NO**, Wei MT, Stone HA, Brangwynne CP. Quantifying dynamics in phase-separated condensates using
355 fluorescence recovery after photobleaching. Biophys J. 2019; 117(7):1285–1300.

356 **Weiner BG**, Pyo AG, Meir Y, Wingreen NS. Motif-pattern dependence of biomolecular phase separation driven
357 by specific interactions. PLoS Comput Biol. 2021; 17(12):e1009748.

358 **Yao RW**, Xu G, Wang Y, Shan L, Luan PF, Wang Y, Wu M, Yang LZ, Xing YH, Yang L, et al. Nascent pre-rRNA sorting
359 via phase separation drives the assembly of dense fibrillar components in the human nucleolus. Mol Cell.
360 2019; 76(5):767–783.

361 **Zhang Y**, Xu B, Weiner BG, Meir Y, Wingreen NS. Decoding the physical principles of two-component biomolec-
362 ular phase separation. eLife. 2021; 10:e62403.

363 **Appendix 1**

364 ## Derivation of the FRAP recovery curve in Equations (5) and (6)

365 The exact solution for $f(t)$ in Equation (4), the fraction of molecules in a spherical condensate of
366 radius R which are unbleached at time t, is derived by Taylor *et al.* in *Taylor et al.* (*2019*) in an
367 integral form using Laplace transforms,

$$f(t) = 1 - \frac{6\alpha k^2}{\pi \sqrt{\lambda}} \int_0^\infty g(u, t^*) \, du, \tag{15}$$

368 where

$$g(u, t^*) = \frac{(u \cos u - \sin u)^2 \exp(-u^2 t^*)}{u^2 \left\{ u^2 \left[u \cos u + (k-1) \sin u \right]^2 + \lambda \left[(1 + k\frac{\alpha}{\lambda}) u \cos u + (k - 1 - k\frac{\alpha}{\lambda}) \sin u \right]^2 \right\}} \tag{16}$$

369 and $\lambda = D_{\text{dil}}/D_{\text{den}}$, $\alpha = c_{\text{den}}/c_{\text{dil}}$, $t^* = tD_{\text{den}}/R^2$, and $k = R\kappa/D_{\text{den}}$, where κ is the interface conduc-
370 tance.

371 To obtain a more intuitive result, we first rearrange $g(u, t^*)$ as

$$g = \frac{\exp(-u^2 t^*)}{u^4 \left(1 + \frac{k}{u \cot u - 1}\right)^2 + \lambda u^2 \left(1 + k\frac{\alpha}{\lambda} + \frac{k}{u \cot u - 1}\right)^2}. \tag{17}$$

372 We note that the diffusion of biomolecules in the dilute phase is typically much faster than diffusion
373 in the dense phase, i.e., $\lambda = D_{\text{dil}}/D_{\text{den}} \gg 1$. In this parameter regime, $g(u, t^*)$ is sharply peaked at
374 the values of u where

$$1 + k\frac{\alpha}{\lambda} + \frac{k}{u \cot u - 1} = 0, \tag{18}$$

375 i.e., when the second term in the denominator of $g(u, t^*)$ becomes 0. Representative $g(u, t^*)$ curves
376 are shown in Appendix 1—figure 1. We can therefore approximate $g(u, t^*)$ as

$$g(u, t^*) = \sum_n a_n \frac{\lambda^2}{u_n^4 k^2 \alpha^2} \exp(-u_n^2 t^*) \delta(u - u_n), \tag{19}$$

377 where u_n is the n-th solution of Equation (18) and $a_n \sim 1/\sqrt{\lambda}$ is the inverse of effective width of n-th
378 peak of $g(u, t^*)$. Clearly, the prefactor of the delta function $\delta(u - u_n)$ in Equation (19) drops rapidly
379 with increasing values of u_n. Consequently, the integral in Equation (15) is always dominated by the
380 contribution from the first mode, and therefore

$$f(t) \approx 1 - \exp(-t/\tau), \tag{20}$$

381 where

$$\tau = \frac{R^2}{D_{\text{den}} u_1^2}, \tag{21}$$

382 with u_1 the first root of Equation (18).

Appendix 1 Figure 1. Representative curves of $g(u, t^*)$. Parameters matched to the simulated A6B6 system: $c_{den} = 7.7\,\text{mM}$, $c_{dil} = 0.05\,\text{mM}$, $D_{den} = 0.013\,\mu\text{m}^2/\text{s}$, $D_{dil} = 17\,\mu\text{m}^2/\text{s}$, and $\kappa = 0.14\,\mu\text{m/s}$. Droplet radius is $R = 1\,\mu\text{m}$ (top) and $0.1\,\mu\text{m}$ (bottom). Inset: same plot with the y-axis in log scale.

At any given values of k, α, and λ, the solutions of Equation (18) can be obtained numerically. Alternatively, we can obtain an approximate analytical solution by first rewriting Equation (18) as

$$1 - u \cot u = K = \frac{k}{1 + k\alpha/\lambda}. \tag{22}$$

We plot the combined parameter $K = k/(1 + k\alpha/\lambda)$ versus the first root u_1 in Appendix 1—figure 2. For small K, Taylor series expansion around $u = 0$ for the left side of Equation (22) yields

$$1 - u\left(\frac{1}{u} - \frac{u}{3}\right) = \frac{u^2}{3} = K, \tag{23}$$

which yields $u_1 = \sqrt{3K}$. For large K, u_1 plateaus at π. We therefore approximate u_1 as

$$u_1 = \frac{\pi\sqrt{3K}}{\sqrt{\pi^2 + 3K}}. \tag{24}$$

This approximate solution is compared with the exact numerical solution of u_1 in Appendix 1—figure 2. The maximum error of about 5% occurs at an intermediate value of K (Appendix 1—figure 2, inset). The relaxation time τ corresponding to the approximate solution in Equation (24) is

$$\tau = \frac{R^2}{\pi^2 D_{den}} + \frac{c_{den} R^2}{3 c_{dil} D_{dil}} + \frac{R}{3\kappa}. \tag{25}$$

Appendix 1 Figure 2. Comparison of exact and approximate solutions of Equation (22). Inset: percentage error of the approximate solution.

Time required to replace all molecules in a spherical droplet in the absorbing boundary limit

The time evolution of the spherically-symmetric concentration profile $c(r, t)$ of molecules in the dilute phase around a droplet of radius R is given by

$$\frac{\partial}{\partial t} c(r, t) = D_{\text{dil}} \left(\frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} \right) c(r, t), \quad r > R. \tag{26}$$

If molecules incident from the dilute phase are immediately and irreversibly absorbed into the dense phase, the boundary condition is then $c(R, t) = 0$. The steady-state solution of Equation (26) in this absorbing-boundary limit is

$$c(r) = c_{\text{dil}} \left(1 - \frac{R}{r} \right), \quad r > R, \tag{27}$$

where c_{dil} is the concentration at $r \to \infty$, which yields a total steady-state flux into the droplet of

$$J = -4\pi R^2 D_{\text{dil}} \left. \frac{d}{dr} c(r) \right|_{r=R} = 4\pi R D_{\text{dil}} c_{\text{dil}}. \tag{28}$$

It then takes a time

$$\tau = \frac{R^2 c_{\text{den}}}{3 D_{\text{dil}} c_{\text{dil}}} \tag{29}$$

to replace all $4\pi R^3 c_{\text{den}}/3$ molecules in the droplet.

Derivation of the interface conductance in the continuum limit, yielding Equations (8) and (13)

To derive the interface conductance in the continuum limit (which neglects the bouncing effect), we start with the mean-field formulation developed in Hubatsch *et al.* *Hubatsch et al.* (*2021*), where the concentration of bleached components $c(\boldsymbol{r}, t)$ is governed by

$$\frac{\partial}{\partial t} c(\boldsymbol{r}, t) = -\nabla \cdot \boldsymbol{j}(\boldsymbol{r}, t), \tag{30}$$

with the flux

$$\boldsymbol{j}(\boldsymbol{r}, t) = -D \left[c_{\text{eq}}(\boldsymbol{r}) \right] \left[\nabla c(\boldsymbol{r}, t) - c(\boldsymbol{r}, t) \frac{\nabla c_{\text{eq}}(\boldsymbol{r})}{c_{\text{eq}}(\boldsymbol{r})} \right], \tag{31}$$

where $c_{\text{eq}}(\boldsymbol{r})$ is the equilibrium concentration profile, and $D[c_{\text{eq}}(\boldsymbol{r})]$ the diffusion coefficient which depends on the local equilibrium concentration.

For a spherical condensate of radius R, if the interface width is narrow, we can assume that the flux going through the interface is uniform in space along the radial direction, i.e.,

$$\boldsymbol{j}(\boldsymbol{r}, t) = -D(r) \left[\frac{\partial}{\partial r} c(r, t) - \frac{c(r, t)}{c_{\text{eq}}(r)} \frac{d}{dr} c_{\text{eq}}(r) \right] \hat{r} = j(t)\hat{r}, \tag{32}$$

where $\hat{r}$ denotes the unit vector in the radial direction. Therefore

$$\frac{\partial}{\partial r} c(r, t) - \frac{c(r, t)}{c_{\text{eq}}(r)} \frac{d}{dr} c_{\text{eq}}(r) = -\frac{j(t)}{D(r)}. \tag{33}$$

The solution to the above equation is

$$c(r, t) = \frac{c_{\text{eq}}(r) c(R, t)}{c_{\text{eq}}(R)} - \int_R^r \frac{c_{\text{eq}}(r) j(t)}{c_{\text{eq}}(r') D(r')} dr'. \tag{34}$$

We assume that the interface spans a width of δ from R_- to R_+ with $R_{+/-} = R \pm \delta/2$, then $c_{\text{eq}}(R - \delta/2) = c_{\text{den}}$ and $c_{\text{eq}}(R + \delta/2) = c_{\text{dil}}$. Substituting Equation (34) into the second boundary condition in Equation (3), we have

$$j(t) = \kappa \left[c(R - \delta/2, t) - \frac{c_{\text{den}}}{c_{\text{dil}}} c(R + \delta/2, t) \right] = \kappa \int_{R-\delta/2}^{R+\delta/2} \frac{c_{\text{den}} j(t)}{c_{\text{eq}}(r) D(r)} dr. \tag{35}$$

⁴¹⁹ We then obtain an expression for the interface conductance κ:

$$\kappa = \left[\int_{R-\delta/2}^{R+\delta/2} \frac{c_{\text{den}}}{c_{\text{eq}}(r)D(r)} dr \right]^{-1}. \tag{36}$$

⁴²⁰ The equilibrium concentration $c_{\text{eq}}(r)$ transitions from c_{den} to c_{dil} between $R-\delta/2$ and $R+\delta/2$, and
⁴²¹ the corresponding diffusion coefficient $D(r)$ transitions from D_{den} to D_{dil}. Assuming a monotonic
⁴²² sigmoidal transition, along with $D_{\text{den}} \ll D_{\text{dil}}$ and $c_{\text{den}} \gg c_{\text{dil}}$, we obtain

$$\frac{c_{\text{den}}}{c_{\text{eq}}(r)D(r)} \leq \frac{1}{D_{\text{den}}} + \frac{c_{\text{den}}}{c_{\text{dil}}D_{\text{dil}}}, \tag{37}$$

⁴²³ which leads to an interface conductance in the continuum limit

$$\kappa \geq \frac{1}{\delta} \left[\frac{1}{D_{\text{den}}} + \frac{c_{\text{den}}}{c_{\text{dil}}D_{\text{dil}}} \right]^{-1}. \tag{38}$$

⁴²⁴ In the simulations in Figure 3, we are only interested in molecules that remain in the dilute phase
⁴²⁵ without entering the dense phase, and the corresponding interface conductance in the continuum
⁴²⁶ limit is then

$$\kappa_0 = \frac{c_{\text{dil}}D_{\text{dil}}}{\delta c_{\text{den}}}. \tag{39}$$

⁴²⁷ **Derivation of the 1-dimensional, slab-geometry versions of Equations (1)–(6)**

For the case of a quasi-1D slab geometry, we consider the condensate to sit in the middle in the region $-l < x < l$ with the simulation box extending along the x-axis from $-L$ to L. The time evolution of the concentration profile $c(x,t)$ of molecules initially located in the dense phase (bleached population) is then given by the 1D diffusion equations:

$$\frac{\partial}{\partial t}c(x,t) = D_{\text{den}}\frac{\partial^2}{\partial x^2}c(x,t), \quad |x| < l;$$
$$\frac{\partial}{\partial t}c(x,t) = D_{\text{dil}}\frac{\partial^2}{\partial x^2}c(x,t), \quad l < |x| < L, \tag{40}$$

⁴²⁸ with the initial condition:

$$c(x,0) = \begin{cases} c_{\text{den}}, & |x| < l; \\ 0, & l < |x| < L, \end{cases} \tag{41}$$

and boundary conditions:

$$\frac{\partial}{\partial x}c(x,t)\bigg|_{x=0} = \frac{\partial}{\partial x}c(x,t)\bigg|_{x=L} = 0; \tag{42}$$

$$-D_{\text{den}}\frac{\partial}{\partial x}c(x,t)\bigg|_{x=l_-} = -D_{\text{dil}}\frac{\partial}{\partial x}c(x,t)\bigg|_{x=l_+} = \kappa\left[c(l_-,t) - \frac{c_{\text{den}}}{c_{\text{dil}}}c(l_+,t)\right]. \tag{43}$$

⁴²⁹ The general solution for the diffusion Equation (40) with the boundary condition in Equation (42)
⁴³⁰ is:

$$c(x,t) = \begin{cases} \sum_n a_n \cos(p_n x)e^{-D_{\text{den}}p_n^2 t}, & |x| < l; \\ \sum_n b_n \cos[q_n(L-x)]e^{-D_{\text{dil}}q_n^2 t}, & l < |x| < L. \end{cases} \tag{44}$$

Applying the boundary condition in Equation (43) to this general solution yields:

$$D_{\text{den}}p_n^2 = D_{\text{dil}}q_n^2, \tag{45}$$

$$D_{\text{den}}a_n p_n \sin(p_n l) = -D_{\text{dil}}b_n q_n \sin(q_n(L-l)) = \kappa\left[a_n\cos(p_n l) - \frac{c_{\text{den}}}{c_{\text{dil}}}b_n\cos\left(q_n(L-l)\right)\right], \tag{46}$$

⁴³¹ which leads to

$$\frac{1}{\kappa} = \sqrt{\frac{\tau}{D_{\text{den}}}}\cot\frac{l}{\sqrt{\tau D_{\text{den}}}} + \frac{c_{\text{den}}}{c_{\text{dil}}}\sqrt{\frac{\tau}{D_{\text{dil}}}}\cot\frac{L-l}{\sqrt{\tau D_{\text{dil}}}}, \tag{47}$$

where $\tau = (D_{\text{den}} p_n^2)^{-1} = (D_{\text{dil}} q_n^2)^{-1}$ is the relaxation time of the n-th mode of the system. For given parameters c_{dil}, c_{den}, D_{dil}, D_{den}, l, L, and κ, τ can be obtained numerically using the above equation. In the regime where the interface conductance is small, we derive an analytical expression for the relaxation time

$$\tau = \frac{l(L - l)}{\kappa(L - l + c_{\text{den}} l / c_{\text{dil}})}, \tag{48}$$

which resembles the corresponding relaxation time for a spherical droplet when interface conductance is small $\tau = R/(3\kappa)$.

We note that, in principle, Equation (47) can be used to infer the value of κ for a system using the relaxation time τ from simulation. However, due to the relatively small simulation sizes, the interface regime can constitute a significant fraction of the dense-phase condensate. This can result in uncertainties in the determination of the dense phase width l and diffusion coefficient D_{den}, etc., leading to errors in the determination of the interface conductance using Equation (47). Such errors can be significant when slow diffusion in the dense phase becomes rate-limiting for overall system relaxation. Therefore, instead of sticking to the "FRAP protocol", we find it more convenient to track the molecules that remain in the dilute phase without ever entering the dense-phase (Figure 3A), as this minimizes the errors caused by any inaccuracies in dense-phase parameters. To relate κ to the decay time of the dilute-phase molecules, we note that the time evolution of the concentration profile $c(x, t)$ of the dilute-phase molecules which have never entered the dense phase is given by:

$$\frac{\partial}{\partial t} c(x, t) = D_{\text{dil}} \frac{\partial^2}{\partial x^2} c(x, t), \qquad l < |x| < L, \tag{49}$$

with the initial condition $c(x, 0) = c_{\text{dil}}$ for $l < |x| < L$ and boundary conditions:

$$\frac{\partial}{\partial x} c(x, t) \bigg|_{x=L} = c(l_-, t) = 0; \tag{50}$$

$$-D_{\text{dil}} \frac{\partial}{\partial x} c(x, t) \bigg|_{x=l_+} = -\kappa \frac{c_{\text{den}}}{c_{\text{dil}}} c(l_+, t). \tag{51}$$

Going through a similar procedure as for Equations (44)–(47), we obtain

$$\kappa = \frac{c_{\text{dil}}}{c_{\text{den}}} \sqrt{\frac{D_{\text{dil}}}{\tau}} \tan \frac{L - l}{\sqrt{\tau D_{\text{dil}}}}. \tag{52}$$

The interface conductance κ for simulated systems in Figure 3C (bottom) is obtained from the relationship in Equation (52) using the measured relaxation time τ, Figure 3C (top), of molecules that remain in the dilute phase without ever entering the dense phase.

Derivation of the unbound-sticker parameter u in Equation (14)

In the interface resistance dominated regime, the decay time τ of the number of dilute-phase molecules that have not entered the dense phase can be obtained from Equation (52):

$$\tau = \frac{c_{\text{dil}}(L - l)}{c_{\text{den}} \kappa}. \tag{53}$$

In the slab geometry, this decay time is controlled by the flux per unit area j entering the dense phase

$$\tau = \frac{c_{\text{dil}} V}{2jA}, \tag{54}$$

where V is the volume of the dilute phase, and A the cross-sectional area of the interface between the dilute and dense phases (the factor of 2 accounts for the two interfaces). Combining these two equations, we have

$$\kappa = \frac{j}{c_{\text{den}}}. \tag{55}$$

For our simulations of polymers with A and B type stickers, we can approximate j by assuming that a polymer incident from the dilute phase will join the dense phase if and only if an unbound

monomer on the polymer binds to an unbound monomer in the dense phase somewhere in the interface region. To find an approximate formula for j we therefore need to estimate the rate of such binding events per unit area of the interface. To this end, we can use the formula for diffusion-limited monomer-monomer binding, but with some modifications: First, we can write the concentration of unbound monomers of type A in the dilute phase as

$$c_{\text{dilA}} = n_{\text{A}} c_{\text{dil}} f_{\text{dilA}}, \tag{56}$$

where n_{A} is the number of type A monomers per polymer and f_{dilA} is the fraction of these monomers that are unbound. This concentration implies a diffusion-limited binding flux onto each unbound dense-phase type B monomer in the interface region

$$j_{\text{A}\to\text{B}} = 4\pi r_0 D_{\text{dil}} c_{\text{dilA}}, \tag{57}$$

where r_0 is the sticker radius, and we have assumed that the diffusion rate is set by the whole polymer. Now we need an estimate for the areal density $\rho_{\text{denB, unbound}}$ of available unbound B-type monomers in the dense-phase interface region, since each one will contribute the above flux [Equation (57)]. We can write

$$\rho_{\text{denB, unbound}} = \delta n_{\text{B}} c_{\text{den}} f_{\text{denB}}, \tag{58}$$

where δ is the width of the interface region, n_{B} is the number of type B monomers per polymer, and f_{denB} is the fraction of unbound B monomers on dense-phase polymers in this region. Finally, we can combine the above equations, and include the binding of dilute-phase B-type monomers to dense-phase A type monomers, to obtain

$$j = j_{\text{A}\to\text{B}} \rho_{\text{denB, unbound}} + j_{\text{B}\to\text{A}} \rho_{\text{denA, unbound}} = 4\pi r_0 D_{\text{dil}} \delta n_{\text{A}} n_{\text{B}} c_{\text{dil}} c_{\text{den}} \left(f_{\text{dilA}} f_{\text{denB}} + f_{\text{dilB}} f_{\text{denA}} \right). \tag{59}$$

Using this expression for j, we can then estimate the ratio between the true κ and its continuum limit κ_0 to be

$$\frac{\kappa}{\kappa_0} = \frac{4\pi r_0 \delta^2 n^2 c_{\text{den}}}{2 + s + s^{-1}} \left(f_{\text{dilA}} f_{\text{denB}} + f_{\text{dilB}} f_{\text{denA}} \right), \tag{60}$$

where $n = n_{\text{A}} + n_{\text{B}}$ is the length of a polymer and $s = c_{\text{A}}/c_{\text{B}}$ is the global stoichiometry. In practice, we find this approximation to be quite accurate up to a constant prefactor (Figure 3D).

₄₈₄ **Appendix 2**

₄₈₅ **Simulation procedures and data recording**

₄₈₆ We perform coarse-grained molecular-dynamics simulations using LAMMPS *Plimpton* (*1995*) to
₄₈₇ simulate phase separation of "sticker and spacer" polymers. Individual polymers are modeled as
₄₈₈ linear chains of spherical stickers of types A and B connected by implicit spacers (Figure 2A) with
₄₈₉ the interaction potentials in Equations (9)–(11), which ensure one-to-one binding between A and B
₄₉₀ stickers.

₄₉₁ For each of the five selected sequences (Figure 2A), we simulate 1000 polymers in a $500\,\mathrm{nm} \times$
₄₉₂ $50\,\mathrm{nm} \times 50\,\mathrm{nm}$ box with periodic boundary conditions. Following the simulation procedures in *Zhang*
₄₉₃ *et al.* (*2021*), we first initialize the simulation by confining polymers in the region $-90\,\mathrm{nm} < x <$
₄₉₄ $90\,\mathrm{nm}$ to promote phase separation and ensure that only a single dense condensate is formed.
₄₉₅ The attractive interaction between A and B stickers [Equation (10)] is gradually switched on from
₄₉₆ $U_0 = 0$ to 14 over 2.5×10^7 time steps. This annealing procedure leads to the formation of a dense
₄₉₇ phase close to its equilibrated concentration. The dense condensate is equilibrated at fixed $U_0 = 14$
₄₉₈ for another 2.5×10^7 steps and then the confinement is removed. The system is equilibrated for
₄₉₉ 2×10^8 more time steps to allow for the formation of a dilute phase and further relaxation of the
₅₀₀ dense phase. We then record the positions of all particles every 2.5×10^5 steps for 800 recordings.
₅₀₁ Through the entire simulation, we equilibrate the system using a Langevin thermostat imple-
₅₀₂ mented with LAMMPS commands fix nve and fix langevin, i.e., the system evolves according to
₅₀₃ *Langevin* (*1908*)

$$m\frac{d^2\vec{r}_i}{dt^2} = -\gamma\frac{d\vec{r}_i}{dt} - \nabla_{\vec{r}_i}U(\vec{r}_1, ..., \vec{r}_N) + \vec{f}, \qquad (61)$$

₅₀₄ where $\vec{r}_i$ is the coordinate of particle i, m is its mass, γ is the friction coefficient, $\vec{f}$ is random ther-
₅₀₅ mal noise, and the potential energy $U(\vec{r}_1, ..., \vec{r}_N)$ contains all interactions between particles, includ-
₅₀₆ ing bonds and sticker-sticker interactions [Equations (9)–(11)]. We take temperature $T = 300\,\mathrm{K}$,
₅₀₇ damping factor $\tau = m/\gamma = 10\,\mathrm{ns}$, step size $dt = 0.1\,\mathrm{ns}$, and mass of particle $m = 188.5\,\mathrm{ag}$. These
₅₀₈ parameters give each sticker the correct diffusion coefficient $D = k_\mathrm{B}T/(3\pi\eta d)$ where η is the water
₅₀₉ viscosity $0.001\,\mathrm{kg/m/s}$ and $d = 2\,\mathrm{nm}$ is the sticker diameter.

₅₁₀ We perform 10 simulation replicates with different random seeds for each of the five selected
₅₁₁ polymer sequences. Consistency of results is checked across replicates and between the first and
₅₁₂ second halves of the recorded data.

₅₁₃ **Determining the dilute- and dense-phase concentrations**

₅₁₄ To measure the dilute- and dense-phase concentrations, we first group polymers into connected
₅₁₅ clusters in each recording. Two stickers are considered connected if they are part of the same poly-
₅₁₆ mer, or if they are within the attraction distance $r_0 = 1\,\mathrm{nm}$. Connected stickers are then grouped
₅₁₇ into clusters. In all simulations, we observe one large cluster which contains most of the polymers,
₅₁₈ and tens to hundreds of very small clusters (Figure 2B). We consider the large cluster to constitute
₅₁₉ the dense phase, and the smaller clusters to be constituents of the dilute phase.

₅₂₀ To find the concentrations of each phase, we identify the center of mass of the dense cluster
₅₂₁ in each recording, and recenter the simulation box to this center of mass. We then compute the
₅₂₂ polymer concentration histogram along the x axis with a bin size 1/50 of box length. The histogram
₅₂₃ of numbers of stickers per bin is averaged over all recordings and simulation replicates. The poly-
₅₂₄ mer concentration profile is derived as the sticker concentration profile divided by the number of
₅₂₅ stickers per polymer. The resulting polymer concentration profile has high values in the middle
₅₂₆ corresponding to the dense-phase concentration, and low values on the two sides correspond-
₅₂₇ ing to the dilute-phase concentration (Figure 2C). The dilute- and dense-phase concentrations in
₅₂₈ Figure 2D are calculated by averaging the concentration profile over the regions ($x \leq -150\,\mathrm{nm}$ or
₅₂₉ $x \geq 150\,\mathrm{nm}$) and ($-10\,\mathrm{nm} \leq x \leq 10\,\mathrm{nm}$), respectively.

Determining the dilute- and dense-phase diffusion coefficients

To measure the dilute- and dense-phase diffusion coefficients, we perform simulations with a pure dilute phase or dense phase, i.e., with polymers at the measured dilute- and dense-phase concentrations. Specifically, for the dilute-phase case, we simulate 750 $(A2B2)_3$, 285 $(A3B3)_2$, 101 A6B6, 104 A8B6, and 180 A10B6 polymers, each in a $150\,\text{nm} \times 150\,\text{nm} \times 150\,\text{nm}$ box with periodic boundary conditions. For the dense-phase case, we simulate 1000 polymers for all selected sequences in an $W\,\text{nm} \times 50\,\text{nm} \times 50\,\text{nm}$ box with periodic boundary conditions, where $W = 116.1$ for $(A2B2)_3$, 96.5 for $(A3B3)_2$, 85.8 for A6B6, 136.5 for A8B6, and 234.2 for A10B6. To equilibrate the system, the attractive interaction between A and B stickers [Equation (10)] is gradually switched on from $U_0 = 0$ to $14\,k_\text{B}T$ over 2.5×10^7 time steps and equilibrated at fixed $U_0 = 14\,k_\text{B}T$ for 2×10^8 more time steps. We then record the displacement of all particles every 2.5×10^5 steps for 400 recordings. 5 simulation replicates with different random seeds are performed for each selected sequence.

To find the diffusion coefficients, we compute the time-averaged mean squared displacement (MSD) for each polymer as a function of the lag time t_lag, and average over all polymers in a simulation box and over 5 replicates. The time- and ensemble-averaged MSD is then linearly fit to $\text{MSD} = 6Dt_\text{lag}$ to extract the diffusion coefficient.

Appendix 2 Figure 1. Interface conductance inferred from average survival time of particles initially in the dilute phase for different criteria for having "joined" the dense phase (cf. Figure 3). The interface conductance is shown for three criteria: a polymer is considered to have joined the dense phase if it is in the dense-phase cluster for a continuous duration of τ, 10τ, or 20τ, where τ is the average bond lifetime of an isolated A-B sticker pair.

Determining the interface conductance

To measure the interface conductance κ, we follow the simple protocol depicted in Figure 3A. This scheme, based on the rate that particles in the dilute phase join the dense phase, is both computationally efficient and allows us to infer the interface conductance even when slow diffusion in the dense phase is rate-limiting for overall system relaxation. Specifically, in this protocol we first define a "survival" variable S for each polymer as a function of time: $S = 1$ if the polymer belongs to any dilute-phase cluster (including a solo cluster), and $S = 0$ if the polymer is in the dense-phase cluster. Next, for all polymers starting with $S = 1$ (i.e., in the dilute phase), we check if there is a period of time (chosen here to be 10 times the average bond lifetime of an isolated A-B pair) for which its S value is always 0 (i.e., the polymer has joined the dense-phase cluster). If yes, we set $S = 1$ at the time points before the joining event and $S = 0$ at all times afterward. If not, we set $S = 1$ for this polymer for all time points. We then average $S(t)$ over all polymers starting with $S = 1$ and over the 10 simulation replicates. The obtained $S(t)$ is the average survival probability of polymer in the dilute-phase polymer that have never entered the dense phase. We fit $S(t)$ to a decaying exponential to extract the decay time τ. The interface conductance κ is then calculated using Equation (52) with the measured decay time and dilute- and dense-phase parameters.

In Figure 3, we set the criterion for a polymer to have entered the dense phase as being contin-
uously connected to the dense-phase cluster for a duration longer than 10τ where τ is the average
bond lifetime of an isolated A-B sticker pair. In Appendix 2—figure 1, we compare the results for
the interface conductance κ using alternative durations, $> \tau$ and $> 20\tau$, as criteria for joining the
dense phase. The value of κ changes very little between the 10τ and 20τ criteria, suggesting that
the results in Figure 3 are robust to the definition of "joining" the dense phase, provided very short-
lived bonds are neglected.

As an alternative approach, we calculated κ by directly measuring the flux j of molecules that
enter the dense phase and then using $\kappa = j/c_{\text{den}}$ [Equation (55)]. To find this flux, we first define an
entering event as occurring when a molecule starting from the dilute phase joins the dense-phase
cluster and stays for a duration longer than 10 times the average bond lifetime for an isolated
A-B sticker pair. We count the number of total entering events N in a simulation (note that some
molecule can enter the dense phase multiple times), and the flux is then $N/(2AT)$, where A is
the cross-sectional area of the interface and T is the duration of the simulation. We show in Ap-
pendix 2—figure 2 that the values of κ obtained via this method are consistent with the results
reported in Figure 3C.

Appendix 2 Figure 2. Comparison of interface conductance κ obtained with the flux method and with
Equation (52) as reported in Figure 3C.

580   We show in Appendix 2—figure 3 a few representative trajectories of A6B6 (top) and A10B6
581   (bottom) polymers "bouncing" off the interface between dilute and dense phases. More bouncing
582   events per unit time are observed in the A6B6 system compared to A10B6 system, consistent with
583   the presence of a larger interface resistance in the A6B6 system.

Appendix 2 Figure 3. Representative trajectories of molecules "bouncing" multiple times at the interface of the A6B6 system (top) and A10B6 system (bottom).