Chlamydia trachomatis induces the transcriptional activity of host YAP in a Hippo-independent fashion

Liam T. Caven[1,2], Amanda J. Brinkworth[1], Rey T. Carabeo[1*]

[1] Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE,
USA

[2] School of Molecular Biosciences, College of Veterinary Medicine, Washington State University,
Pullman, WA, USA

* Correspondence:
Rey Carabeo
rey.carabeo@unmc.edu

Keywords: *Chlamydia trachomatis*, **host response, YAP, pathogen-directed host transcription,**
Hippo kinase

Abstract

The obligate intracellular pathogen *Chlamydia trachomatis* is the causative agent of the most
common bacterial sexually transmitted disease worldwide. While the host response to infection by
this pathogen has been well characterized, it remains unclear to what extent host gene expression
during infection is the product of *Chlamydia*-directed modulation of host transcription factors. In this
report, we show the transcriptome of *Chlamydia*-infected epithelial cells exhibits gene expression
consistent with activity of YAP, a transcriptional coactivator implicated in cell proliferation, organ
morphogenesis, wound healing, and fibrosis. After confirming induction of YAP target genes during
infection, we observed increased YAP nuclear translocation in *Chlamydia*-infected epithelial cells.
We show that increased YAP activation is a *Chlamydia*-directed process occurring during midcycle
infection; critically, this phenotype is sensitive to inhibition of protein synthesis by the pathogen.
Infection-mediated YAP activation bypasses YAP inhibition by the Hippo kinase cascade, instead
involving YAP tyrosine phospho-activation and the activity of host Src family kinases. Taken
together, our results define a mechanism for *Chlamydia*-directed modulation of host gene expression
independent of the host inflammatory response to infection, as well as introduce novel therapeutic
targets for treatment of this pervasive disease.

1   Introduction

Chlamydia trachomatis is the most common bacterial cause of both sexually-transmitted
disease and infectious blindness worldwide. An obligate intracellular pathogen, *C. trachomatis* binds
to and induces its own internalization by epithelial cells, establishing a replicative niche within a
pathogen-remodeled vacuole termed the chlamydial inclusion. Chronic or repeated *C. trachomatis*
infection is associated with severe inflammatory and fibrotic sequelae. *C. trachomatis* serovars A-C,
which preferentially infect the conjunctiva, are associated with excess collagen production and
contraction of infected tissues, leading to the inward turning of the upper eyelid and subsequent
corneal abrasion by the eyelashes (Taylor et al., 2014; Lansingh, 2016). *Chlamydia*-mediated
trachoma is endemic to Africa, Latin America, Asia, and the Middle East (Hu et al., 2010), and is
responsible for the blindness or visual impairment of up to 1.9 million people according to current
WHO estimates. Infection by *C. trachomatis* serovars D-K, which preferentially colonize the
reproductive tract, are associated with pelvic inflammatory disease as well as progressive scarring of
the fallopian tubes (Ness et al., 2008; Haggerty et al., 2010a). Blockage of the upper female genital
tract by scar tissue can consequently lead to ectopic pregnancy or tubal factor infertility (Chow et al.,
1990). In the United Kingdom, *C. trachomatis* is estimated to account for 29% of all cases of tubal
factor infertility, which in turn represents 25% of all cases of female infertility (Price et al., 2016).
Collectively, the urgent threat to public health presented *Chlamydia*-associated sequelae has made
understanding the underlying mechanisms of the host-*Chlamydia* interaction a subject of ongoing
investigation.

Initial study of the host response to infection has made a persuasive argument for infection-
associated pathology being the product of cytokine signaling cascade originating in infected
epithelial cells, termed the cellular paradigm of pathogenesis (Stephens, 2003). Clinical infections
have revealed substantial infiltration of infected tissues by immune cells *in vivo*, suggesting that
infected epithelial cells recruit components of the innate and adaptive immune systems to the site of
infection (Kiviat et al., 1990; Stephens, 2003). Indeed, upregulation of an expansive portfolio of
inflammatory cytokines has been consistently observed in chlamydial infections *in vitro*, including
GROα, GM-CSF, IL-1α, IL-6, IL-11, and IL-8 (Eckmann et al., 1993; Rasmussen et al., 1997; Cheng
et al., 2008). However, the pro-inflammatory host response fails to account for the fibrotic outcomes
of asymptomatic infection; it is estimated that up to 75% of *C. trachomatis* infections go unreported
due to absent or subclinical symptoms, and that up to 18% of such cases are believed to cause
infertility (Haggerty et al., 2010b).

One explanation for this discrepancy in the data is that *C. trachomatis* induces pathology via
direct manipulation of host gene expression; indeed, mounting evidence suggests that regulation of
host transcription is targeted by the pathogen in a variety of ways. Past work has shown that *C.
trachomatis* antagonizes pro-inflammatory signaling mediated by the transcription factor NF-κB by
stabilizing its inhibitory subunit IκBα, through the activity of the chlamydial deubiquitnase
ChlaDub1 (Le Negrate et al., 2008). The chlamydial proteinase CPAF has also been shown to inhibit
the NF-κB complex, in that ectopic expression of this chlamydial effector leads to degradation of the
complex subunit p65/RelA (Christian et al., 2010). However, the specific role of CPAF in inhibiting
pro-inflammatory gene expression remains unclear, given that infection with CPAF-deficient mutant
strains does not exhibit increased NF-κB activation (Snavely et al., 2014), likely due to compensatory
activity of ChlaDub1 and other potential mechanisms of chlamydial NF-κB antagonism. In similar
fashion, infection with the related species *C. pneumonia*e has been shown to induce expression and
phosphorylation of the AP-1 transcription factor c-Jun (Krämer et al., 2015). Intriguingly, this effect
did not occur during infection with heat-killed *C. pneumoniae*, indicating that c-Jun activation is a

pathogen-directed phenotype with implications for virulence, given that c-Jun knockdown negatively
impacted bacterial load as well. Collectively, these data illustrate the potential for *Chlamydiae* to
induce changes in host gene expression to facilitate pathogenesis, demonstrating a need to better
characterize how the pathogen acts on host transcription factors.

To that end, we have examined the host transcriptome of *C. trachomatis*-infected immortalized
endocervical epithelial cells, to perform unbiased *in silico* discovery of pathogen-modulated host
transcription factors. In this report, we show that *C. trachomatis* infection induces gene expression
consistent with the function of YAP (<u>Y</u>es-<u>a</u>ssociated <u>p</u>rotein), a transcriptional coactivator. After
empirically confirming infection- and YAP-dependent induction of target genes identified by this
analysis, we subsequently demonstrate that infection promotes YAP activity by enhancing its nuclear
translocation. Increased YAP nuclear translocation was first detectable at 18 hours post-infection and
exhibited sensitivity to inhibition of bacterial protein synthesis, indicating that YAP induction is a
pathogen-directed process. Intriguingly, chlamydial activation of YAP occurs irrespective of the
Hippo kinase cascade (a well characterized inhibitor of YAP activity), instead relying upon YAP
phosphorylation at Y357, a post-translational modification shown to enhance YAP activation (Smoot
et al., 2018; Sugihara et al., 2018). We show that phosphorylation at this residue is essential for
infection-mediated YAP nuclear translocation. Additionally, we find that enhanced YAP nuclear
translocation and Y357 phosphorylation in infected cells is sensitive to inhibition of Src-family
kinases, consistent with prior reports implicating this kinase family in Hippo-independent YAP
activation. With these data, we present an initial framework by which *Chlamydia trachomatis*
manipulates the activity of a transcription factor to drive changes in host gene expression.

2   Materials and Methods

2.1   Eukaryotic Cell Culture

Human endocervical epithelial HPV-16 E6/E7 transformed End1s (End1 E6/E7, ATCC CRL-
2615) were cultured at 37° C with 5% atmospheric CO_2 in Keratinocyte Serum-Free Medium
(KSFM, Thermo Fisher Scientific) supplemented with human recombinant epidermal growth factor,
bovine pituitary extract, 5 micrograms/mL gentamicin, and 0.4 mM $CaCl_2$ (unless otherwise
indicated). For all experiments, End1s were cultured between passages 3 and 15. Primary human
cervical epithelial cells (HCECs, ATCC PCS-0480-011, Lot 80306190) were cultured at 37° C with
5% atmospheric CO_2 in Cervical Epithelial Cell Basal Medium (CECBM, ATCC PCS-480-032)
supplemented with all contents of a Cervical Epithelial Growth Kit (ATCC PCS-080-042). Raft
cultures were prepared as previously described by our laboratory (Nogueira et al., 2017), exchanging
HaCaT cells for End1s where indicated, and cultured over 19 days before being fixed for 30 minutes
in 4% paraformaldehyde, OCT-embedded, cryo-sectioned, and stained with hematoxylin and eosin.

2.2   Cloning and DNA/siRNA Transfection

Both FLAG-YAP1-WT and FLAG-YAP1-Y357F were generated via the simplified
FastCloning method of Gibson assembly (Li et al., 2011); briefly, PCR-amplified recombinant DNA
was treated with DpnI (NEB Biolabs) for 3 h, then transformed into Stellar chemically competent *E.*
coli (Clontech 636763) via heat shock, followed by Kanamycin selection, screening via colony PCR
(where applicable), and sequence confirmation. FLAG-YAP1-WT was generated via amplification of
pEGFP-C3-HYAP1, a gift from Marius Sudol (Addgene plasmid # 17843 ;
http://n2t.net/addgene:17843 ; RRID:Addgene_17843), using the following primers: FWD: 5'-
GACTACAAAGACGATGACGACAAGGATCCCGGGCAGCAGCCG-3'; REV: 5'-
CTTGTCGTCATCGTCTTTGTAGTCCATGGTGGCGACCGGTAGCG-3'. FLAG-YAP1-Y357F

was generated from FLAG-YAP1-WT via site-directed mutagenesis of the Y357 codon (TAC to
TTC), using the following primers: FWD: 5'-GCAGCTTCAGTGTCCCTCGA-3', REV: 5'-
ACACTGAAGCTGCTCATGCTTAGTC-3'.
End1s were transfected with plasmid DNA via electroporation: after harvest via
trypsinization, $2x10^6$ End1s per construct were resuspended in 400 μL Opti-MEM at room
temperature (Thermo Fisher Scientific 31985062) containing 20 ug of plasmid DNA and 3 μL
sheared salmon sperm DNA (Thermo Fisher Scientific AM9680). Suspensions were then transferred
to individual GenePulser 0.4 cm cuvettes (Bio-Rad 1652088), then electroporated using a GenePulser
XCell (Bio-Rad) using the following parameters: exponential decay program template, 225 V, 850
μF capacitance, infinite resistance. 100 μL aliquots of the resulting electroporants were transferred to
individual wells of a 24-well plate containing cover slips and pre-warmed KSFM, then incubated
until confluent monolayers had formed (24-48h) for subsequent infection as previously described.
For siRNA-mediated knockdown experiments, End1s were transfected with either an ON-
TARGETplus non-targeting siRNA pool (Horizon Discovery D-001810-10-05) or an ON-
TARGETplus YAP1-targeting siRNA SMARTpool (Horizon Discovery L-012200-00-0005) using
Lipofectamine 3000 (Thermo Fisher Scientific L3000008), per manufacturer's instructions, at an
empirically determined optimal concentration of 10 nM. At 16 h post-seeding of End1s at 125% of
confluence on 6-well plates as described above, siRNA was combined in Opti-MEM (Thermo Fisher
Scientific 31985062) with the Lipofectamine 3000 reagent, incubated for 5 m at room temperature to
allow for liposome formation, then added to wells dropwise with mixing. Transfected End1s were
then incubated for 24 h prior to infection with *Chlamydia* (see below).

2.3   Chlamydial Infections

Chlamydia trachomatis serovar L2 (434/Bu) was originally obtained from Dr. Ted Hackstadt
(Rocky Mountain National Laboratory, NIAID). Chlamydial EBs were isolated from infected,
cycloheximide-treated McCoy cells at 36-40 hours post-infection (hpi) and purified by density
gradient centrifugation as previously described (Caldwell et al., 1981). For infection of 6- and 24-
well tissue culture plates (Greiner Bio-One 657160 and 662160), End1s seeded at 125% of
confluence (16 h post-seeding) or HCECs grown to confluence (6-7 days) were washed with pre-
warmed Hanks Buffered Saline Solution (HBSS) prior to inoculation with *Chlamydia*-containing
KSFM or CECBM at a multiplicity of infection (MOI) of 5, unless otherwise indicated. Tissue
culture plates were centrifuged at 4° C and 500 rcf (Eppendorf 5810 R tabletop centrifuge, A-4-81
rotor) for 15 minutes to synchronize infection. Inoculum was then aspirated, and cells were washed
with chilled HBSS prior to the addition of pre-warmed KSFM or CECBM.

2.4   Bulk RNA-Sequencing and Analysis

End1s seeded on fibronectin-coated 6-well plates (Corning 354402) and infected at an MOI of
2 as described above were harvested for RNA using TRIzol (Thermo Fisher Scientific 15596026) and
the DNA-free DNA removal kit (Thermo Fisher Scientific AM1906), according to manufacturers'
protocols. Extracted RNA was subsequently enriched for polyadenylated mRNA transcripts using the
NucleoTrap mRNA enrichment kit (Macherey-Nagel) according to the manufacturer's protocol. For
the comparison of mock- and *Ct* serovar L2-infected transcriptomes, RNA samples were
subsequently assayed for fragmentation using an Agilent 5200 Fragment Analyzer; intact samples
were enriched for polyadenylated transcripts via the NucleoTrap kit a second time due to the high
incidence of ribosomal RNA in the fragment analysis results. cDNA library preparation was
performed using the Ion Xpress Plus Fragment Library kit (Thermo Fisher Scientific 4471269), and
sequencing of cDNA libraries was performed using the Ion Torrent system (Life Technologies). For

the bulk RNA-sequencing of primary cells, HCECs infected at an MOI of 2 were harvested for RNA
samples via TRIzol and the DNA-free DNA removal kit as described above, and were subsequently
assayed for fragmentation using an Agilent Bioanalyzer 2100. cDNA library preparation of intact
samples was performed using the NuGEN Universal mRNA-Seq Library Preparation kit, and
sequencing of cDNA libraries was performed using the NextSeq 550 system (Illumina).
Read alignment and downstream analysis of both experiments was performed using CLC
Genomics Workbench (Qiagen); each treatment group was comprised of libraries from three
biological replicates, each with a minimum of 30 million reads (unstranded single read, mean length
150 bp), genes with an FDRP ≤ 0.05 were considered differentially expressed. ChEA
crossreferencing was performed in Python, identifying the number of genes induced (FC ≥ 1.5, FDRP
≤ 0.05) and repressed (FC ≤ 1.5, FDRP ≤ 0.05) by infection for each transcription factor's gene
targets (Lachmann et al., 2010). Pearson's correlation coefficients between End1 and HCEC
expression of YAP-responsive genes were subsequently calculated in R, using $\log_2$-transformed fold
changes of all genes differentially expressed in at least one data set.

2.5　Reverse Transcription Quantitative Real-Time PCR

End1s seeded on fibronectin-coated 6-well plates (Corning 354402) and infected or cocultured
as described above were harvested for RNA using TRIzol (Thermo Fisher Scientific 15596026) and
the DNA-free DNA removal kit (Thermo Fisher Scientific AM1906), according to manufacturers'
protocols. cDNA libraries were subsequently prepared using SuperScript IV Reverse Transcriptase
(Thermo Fisher Scientific 11766050) according to the manufacturer's protocol. Quantitative real-
time PCR was performed on a QuantStudio 3 (Thermo Fisher Scientific) using TaqMan assay kits
(Thermo Fisher Scientific) of the following genes: CTGF (Hs01026927_g1), INHBA
(Hs01081598_m1), BMP2 (Hs00154192_m1), MMP9 (Hs00957562_m1), and the housekeeping
gene HPRT (Hs02800695_m1). Statistical analysis was performed in R, using pairwise Student's t-
tests and Bonferroni's correction for multiple comparisons; p-values less than 0.05 were considered
statistically significant.

2.6　Immunofluorescence Microscopy

End1s or HCECs seeded on glass cover slips (VWR) coated with fibronectin (Corning 354008)
or fibronectin-deposited micropatterned chips (CYTOO) and infected as described above were fixed
in 4% paraformaldehyde in phosphate-buffered saline (PBS) for 10 minutes at 37° C, washed in PBS,
then blocked in 5% bovine serum albumin (BSA) in PBS for 1 hour at room temperature. Fixed and
blocked cover slips were subsequently incubated overnight at 4° C with primary antibodies in 1%
BSA-PBS: rabbit anti-YAP (CST 4912, 1:100 dilution), rabbit anti-E-Cadherin (CST 3195, 1:200
dilution). Cover slips were again washed in PBS, then incubated for 1 hour at room temperature with
the following fluorophore-conjugated antibodies/dyes in 1% BSA-PBS: goat anti-rabbit Alexa-488
conjugate (Thermo Fisher Scientific A-11034, 1:1000 dilution), phalloidin Alexa-594 conjugate
(Thermo Fisher Scientific A-12381, 1:120 dilution), DAPI (Sigma-Aldrich 10236276001, 1:1000
dilution). Afterward, cover slips were washed in PBS and ultrapure water, then mounted on
microscope slides using Shandon Immu-Mount (ThermoFisher Scientific 9990402).
A minimum of 5 fields per cover slip were imaged using a SP-8 Lightning Confocal
Microscope (Leica) or CSU-W1 Spinning-Disk Confocal Microscope (Nikon). To account for
variations in nuclear size and avoid biased selection of cells for measurement, blinded image
quantification was performed by assigning image filenames randomized number codes, selecting 10
nuclei at random per field using only the DAPI channel, manually masking the nuclear area, and
recording the mean YAP fluorescence intensity per nucleus. To account for variation in total YAP

between cells, staining efficiency between cover slips, and compression of the nuclear/cytosolic
compartments by the chlamydial inclusion, 5 cytosolic regions not occluded by an inclusion body
and adjacent to measured nuclei were selected per field, with the mean YAP fluorescence intensity of
these regions averaged to produce a per-field measurement of mean cytosolic YAP fluorescence
intensity; nuclear translocation of YAP was thereby expressed as a ratio of mean nuclear
fluorescence to mean cytosolic fluorescence. Statistical analysis was performed in R, using a
Kruskal-Wallis test to first verify a statistically significant (p-value < 0.05) difference between
treatment groups. Subsequent pairwise comparisons were performed using a Wilcoxon rank sum test
and Bonferroni's correction for multiple comparisons, with p-values less than 0.05 being considered
statistically significant.

2.7   SDS-PAGE and Western Blotting

To minimize activity of the chlamydial protease CPAF, End1s seeded on 6-well plates and
infected as described above were subsequently lysed in 1% SDS buffer heated to 95° C, as previously
described (Johnson et al., 2015). After treatment with Pierce Universal Nuclease (Thermo Fisher
Scientific, 1:1000 dilution) for 5 minutes at room temperature, lysates were combined with 4X
Laemmli Sample Buffer (Bio-Rad 1610747) for loading on a 7.5% acrylamide/bis-acrylamide gel for
SDS-PAGE (1.5 hours, 120V). Gels were then transferred to PVDF membranes (Bio-Rad 1620177)
using a semi-dry transfer method. After blocking in 5% BSA in PBST (PBS containing 0.1% Tween-
20) for 1 hour at room temperature, membranes were incubated overnight at 4° C with primary
antibodies in 5% BSA-PBST: rabbit anti-YAP (CST 4912, 1:1000 dilution), rabbit anti-S127-pYAP
(CST 13008, 1:1000 dilution), rabbit anti-Y357-pYAP (Abcam ab62571, 1:1000 dilution).
Membranes were subsequently washed in PBST, then incubated with a goat anti-rabbit HRP-
conjugated secondary antibody (Dako P0448, 1:2000 dilution in 5% BSA-PBST) for 2 hours at room
temperature. After additional washing in PBST, membranes were imaged using Immobilon HRP
Substrate (Millipore Sigma WBKLS0500) or an Azure Biosystems c600. Images were analyzed
using the ImageJ gel analysis tool to quantify the fluorescence density of phospho-analyte bands
relative to the YAP total protein loading control. Statistical analysis was performed in R, using
pairwise Student's t-tests with Bonferroni's correction for multiple comparisons; p-values less than
0.05 were considered statistically significant.

3   Results

3.1   *Chlamydia* infection induces the expression of a subset of YAP target genes.

To identify host transcription factors modulated by *Chlamydia* infection, we opted to model
infection of the upper female genital tract using End1/E6E7 immortalized, non-transformed
endocervical epithelial cells (Fichorova et al., 1997). We first confirmed the non-tumorigenic growth
properties of End1 cells via organotypic raft culture: first seeding End1 cells on collagen gels in a
transwell insert, then inducing differentiation via exposure to the air-liquid interface (Nogueira et al.,
2017). Transformed cervical epithelial cells form irregular layers when induced to differentiate via
air-liquid interface (Aasen et al., 2003); by contrast, differentiating End1s grown via raft culture
formed epithelial tissues of uniform thickness (Supplementary Figure S1), consistent with previous
reports of organotypic culture with this cell line (Gali et al., 2010). Having thus validated our
experimental model, we then infected End1s with the lymphogranuloma venereum *Chlamydia*
trachomatis serovar L2, then harvested RNA for bulk RNA-sequencing at 24 hours post-infection
(hpi). This timepoint occurs well after chlamydial differentiation into proliferative and metabolically
active reticulate bodies (Lee et al., 2018), thus allowing for production of any chlamydial factors
modulating host gene expression. Critically, infection induced the differential expression (FDRP ≤

0.05) of 3611 genes relative to a mock-infected control (Figure 1A, Supplementary Data S1, GEO
Accession: GSE180784).

To then determine potential transcription factors driving the host response to infection, we then
cross-referenced this differentially expressed gene set with the ChEA database of transcription factor
gene targets, derived from gene set enrichment analysis of published ChIP-chip, ChIP-seq, ChIP-
PET, and DamID data (Lachmann et al., 2010). Of the 199 transcription factors included in ChEA,
149 exhibited twice as many induced target genes (FC ≥ 1.5) as repressed target genes (FC ≤ 1.5),
suggesting chlamydial induction of transcription factor activity (Supplementary Data S2). Critically,
targets of transcription factors known to regulate inflammation (IRF1, STAT1, STAT3) and fibrosis
(SMAD2/3, EGR1, TEAD4, YAP1) appeared to be highly induced in the host (Table 1), consistent
with prior reports of infection-dependent induction of pro-inflammatory and pro-fibrotic gene
expression (Carlson et al., 2005; Lad et al., 2005; Humphrys et al., 2013; Sun et al., 2017).

Intriguingly, the fibrosis-associated transcription factors identified by this analysis (SMAD2/3,
EGR1, and TEAD4) are known binding partners of the transcriptional coactivator YAP1
(Zagurovskaya et al., 2009; Zanconato et al., 2015; Szeto et al., 2016). While YAP1 target gene
induction was modest compared to other transcription factors identified by this method, the more
robust target gene induction of transcription factors upregulated by YAP suggests the potential for
chlamydial modulation of this transcriptional coactivator. To assess the physiological relevance of
apparent YAP-associated gene expression, we then sequenced polyadenylated RNA from primary
human endocervical epithelial cells (HCECs) infected with *Ct* serovar L2 (Supplementary Data S3,
GEO Accession: GSE180784). Infected HCECs exhibited differential expression (FDRP ≤ 0.05) of
8241 genes (Figure 1B), of which 839 mapped to the ChEA database of YAP-responsive genes
(Figure 1C). Expression of YAP-responsive genes by infected End1s and HCECs exhibited
significant correlation (Figure 1D), suggesting that infection of these cell types activates a conserved,
putatively YAP-mediated transcriptional program. Intriguingly, only a subset of YAP target genes is
upregulated in either End1s or HCECs, suggesting that *Chlamydia* may alter the YAP regulon in host
cells, or potentially modulate YAP transcriptional activity (Figure 1E). Taken together, these data
imply that the transcriptome of *Chlamydia*-infected epithelial cells may be in part be driven by YAP
activity.

3.2 *Chlamydia* **infection promotes YAP nuclear translocation.**

To confirm our RNA-sequencing analysis suggesting infection may promote YAP-dependent
gene expression, we next assayed expression of connective tissue growth factor (CTGF) – a
consistent marker of YAP target gene induction (Futakuchi et al., 2018). We observed host CTGF
expression was significantly enhanced by infection, as was expression of additional genes known to
be YAP-responsive (Figure 2A), including inhibin beta A (INHBA), bone morphogenetic protein 2
(BMP2), and matrix metallopeptidase 9 (MMP9) (Mo et al., 2012; Huang et al., 2016; Pan et al.,
2017). Critically, CTGF expression was sensitive to siRNA-mediated YAP knockdown (Figure 2B,
Supplementary Figure S1), suggesting that infection modulates gene expression in a YAP-dependent
fashion.

YAP's ability to access the nuclear compartment and its DNA-binding partners correlates with
YAP-mediated gene expression (Pocaterra et al., 2020). Thus, we opted to confirm that chlamydial
infection promotes YAP activity by assaying the nuclear incidence of YAP in serovar L2-infected
cells via immunofluorescence. Consistent with our initial observation of YAP target gene induction,
nuclear YAP was increased in *Chlamydia*-infected End1/E6E7 cells at 24 hpi, relative to mock-

infected control cells (Figure 2C). To account for variations in nuclear size and staining efficiency,
we subsequently performed blind quantification of randomly selected nuclei in each sample,
measuring ratios of nuclear to cytosolic YAP, analogous to previous study of YAP activation (Zhao
et al., 2007; Dupont et al., 2011; Das et al., 2016). Consistently, infection was concomitant with a
1.5-fold increase in nuclear-to-cytosolic YAP ratio relative to mock-infected cells (Figure 2D). Given
that expression of HPV protein E6 has been shown to stabilize YAP and thereby modulate its activity
(Strickland et al., 2018), we assayed nuclear YAP translocation in primary HCECs as well, observing
an equivalent increase to that in E6-expressing End1s (Figure 2E, 2F). Collectively, these data
indicate chlamydial induction of YAP is irrespective of HPV E6/E7 expression.

To confirm that the observed increase in YAP activity was a *Chlamydia*-driven process
requiring the action of one or more chlamydial effectors, we utilized chloramphenicol to inhibit
chlamydial protein synthesis. Over the course of a 24-hour infection, we observed a significant
increase in YAP nuclear localization by immunofluorescence between mock- and serovar L2-infected
End1/E6E7 cells, starting as early as 8 hpi. A statistically significant increase between infected cells
was first detected between 12 and 18 hpi, indicating that synthesis of YAP-modulating chlamydial
effectors may occur at this stage of infection (Figure 2G). We next attempted to attenuate chlamydial
YAP induction at 18 hpi via treatment with chloramphenicol. In agreement with our initial
hypothesis, chloramphenicol treatment for one hour prior to fixation at 18 hpi was sufficient to
significantly inhibit YAP nuclear translocation relative to a DMSO-treated control (Figures 2H, 2I).
Given these data, we conclude that YAP activation in infected cells requires *de novo* synthesis of
chlamydial effectors.

3.3 ***Chlamydia* infection bypasses Hippo-mediated YAP inhibition by enhancing YAP**
phosphorylation at Y357.

In confluent epithelial cells, YAP's access to the nuclear compartment is inhibited by the Hippo
kinase complex, which in turn is stabilized by cell-cell contact (Gumbiner and Kim, 2014). The
phosphorylation cascade of Hippo complex members terminates in YAP phosphorylation at S127 and
other serine residues, which drives cytoplasmic sequestration of YAP via binding to 14-3-3 (Piccolo
et al., 2014). To determine if *Chlamydia* induced YAP activity by relieving Hippo-mediated
inhibition, we examined YAP activation in experimentally enforced culture conditions of inhibited
cell-cell contact. By seeding End1/E6E7 cells on glass treated with fibronectin in 30- and 45-μm
micropatterns (Degot et al., 2010), the resulting deposition of small cell clusters allowed us to assay
YAP nuclear translocation in an environment where cell-cell contact (and, by extension, Hippo-
mediated YAP inhibition) was absent or severely limited. In both 30-μm patterns (1-3 cells/cluster)
and 45-μm patterns (2-4 cells/cluster), we observed a significant increase in YAP nuclear
translocation in serovar L2-infected cells at 24 hpi, relative to mock-infected controls (Figure 3A).
Quantification of nuclear translocation by immunofluorescence indicated that infection promotes a
1.5-fold increase in YAP activation relative to uninfected cells, equivalent to our observations in
confluent monolayers (Figure 3B). This phenotype was additionally observed when adherens
junctions were disassembled by calcium chloride withdrawal in confluent *Chlamydia*-infected
End1/E6E7 monolayers (Supplementary Figure S2) (O'Keefe et al., 1987). Taken together, these data
suggest that chlamydial activation of YAP occurs independent of cell-cell contact, indicating this
phenotype is not dependent on destabilization of the YAP-inhibitory Hippo kinase complex.

The recruitment of Hippo cascade members to adherens junctions promotes the formation of a
YAP-inhibitory complex, resulting in YAP phosphorylation at serine residues by Hippo complex
members LATS1/2 (Zhao et al., 2007). In contrast, recent studies have implicated phosphorylation of

YAP at Y357 and other tyrosine residues in Hippo-independent YAP activation (Elbediwy et al.,
2018; Sugihara et al., 2018, 2018). To confirm that chlamydial YAP activation occurs via Hippo-
independent signaling, we thus surveyed YAP S127 and Y357 phosphorylation in confluent,
Chlamydia-infected End1s via Western blotting. Serovar L2-infected End1/E6E7 cells exhibited
increased YAP Y357 phosphorylation at 24 hpi relative to a mock-infected control. This result was
consistent with mock-infected cells treated with sodium orthovanadate (Na_3VO_4), a tyrosine
phosphatase inhibitor that serves as a positive control for increased YAP tyrosine phosphorylation
(Figure 3C) (Sugihara et al., 2018). Surprisingly, L2-infected cells did not exhibit altered YAP S127,
suggesting infection does not relieve Hippo-mediated YAP inhibition (Figure 3D). By contrast,
pY357-YAP levels were increased in an infection-dependent manner (Figure 3E). Collectively, these
data demonstrate that infection-dependent YAP activation bypasses S127 phospho-inhibition by the
Hippo kinase cascade, instead promoting alternative phospho-activation of YAP at Y357 to facilitate
its nuclear translocation.

3.4   SFK-mediated YAP phosphorylation at Y357 is required for chlamydial YAP activation.

In contrast with the YAP-inhibitory effect of Hippo kinase at cell-cell junctions, YAP
activation has been associated with cell-substrate contact, as well as the activity of Src-family
kinases. Cell adhesion to fibronectin has been shown to promote YAP nuclear translocation in a Src-
dependent, Hippo-inhibitory fashion (Kim and Gumbiner, 2015). Intriguingly, Src and the related
kinase Lck are also implicated in Hippo-independent YAP activation via Y357 phosphorylation (Li
et al., 2016; Elbediwy et al., 2018; Smoot et al., 2018; Sugihara et al., 2018). Further, it has been
shown that *Chlamydia* induces Src activity (Mital and Hackstadt, 2011), and that Src-family kinase
(SFK) modulation is critical to multiple stages of chlamydial pathogenesis (Elwell et al., 2011). Thus,
to determine if infection-mediated YAP activity was the product of SFK-mediated YAP activation,
we assayed YAP nuclear translocation of infected cells treated with the pan-SFK inhibitor PP2.
Consistent with our hypothesis, serovar L2-infected cells treated with PP2 over the course of a 24-
hour infection exhibited statistically significant attenuation of YAP nuclear translocation relative to a
DMSO-treated control (Figures 4A, 4B). YAP phosphorylation at Y357 in infected cells was also
severely attenuated by SFK inhibition (Figure 4C), consistent with studies directly implicating SFKs
in YAP tyrosine phosphorylation (Li et al., 2016; Sugihara et al., 2018). To confirm that YAP
phosphorylation at this residue is indeed required for infection-dependent YAP nuclear translocation,
we ectopically expressed FLAG-tagged, Y357F-mutant YAP in infected End1s, comparing nuclear
translocation to an equivalent mock-infected control. Importantly, nuclear translocation of the
phospho-dead YAP-Y357F mutant in infected cells was indistinguishable from mock-infected cells
expressing the same construct, whereas infection significantly increased the nuclear translocation of
FLAG-tagged, wild-type YAP relative to a mock-infected control (Figures 4D, 4E). Taken together,
these data indicate that chlamydial induction of YAP activity operates via SFK- and Y357-dependent
means of enhancing YAP nuclear translocation.

4   Discussion

In this report, we present a novel mechanism for *Chlamydia*-directed modulation of host gene
expression. Infection of primary and immortalized epithelial cells with *C. trachomatis* serovar L2
resulted in a host transcriptome consistent with induction of the transcriptional coactivator YAP and
its DNA-binding partners. Consistent with this observation, infected cells also exhibit increased YAP
nuclear translocation. This phenotype was concomitant with increased expression of YAP target
genes INHBA, BMP2, MMP9, and CTGF (which in the latter case was sensitive to YAP siRNA-
mediated knockdown). Enhanced YAP activation is first detectable at 18 hours post-infection;

critically, this phenotype was attenuated by inhibition of bacterial protein synthesis, indicating YAP
activation is a pathogen-directed effect of infection. Intriguingly, *Chlamydia* infection induced YAP
nuclear translocation in epithelial cells that were sparsely seeded on micropatterned chips – a
scenario in which canonical YAP inhibition by the Hippo kinase cascade is minimal – suggesting that
infection enhances YAP activation in a Hippo-independent fashion. Accordingly, Hippo-dependent
YAP phospho-inhibition at S127 was unaffected by infection, whereas YAP phosphorylation at Y357
was enhanced. Inhibition of host Src-family kinases significantly attenuated both YAP Y357
phosphorylation and nuclear translocation, consistent with previous reports of SFK-mediated
phosphorylation at this residue enhancing YAP activity. Importantly, we observe that infection does
not promote nuclear translocation of a YAP Y357F mutant, indicating that phosphorylation at this
residue is required for chlamydial YAP activation. Taken together, our data show *Chlamydia* induces
YAP-dependent gene expression by acting upon an alternative, Hippo-independent regulatory
pathway in the host, involving phosphorylation of YAP at Y357 and the activity of host Src-family
kinases.

Our observation of *Chlamydia*-directed induction of YAP activity is consistent with prior
reports of YAP modulation by other bacterial pathogens. The Gram-negative pathogen *Helicobacter*
pylori was recently shown to enhance YAP expression through the action of the effector CagA (Li et
al., 2018). It is unclear how CagA interacts with the steady-state inhibition of YAP by the Hippo
kinase cascade; given the prior finding that CagA destabilizes host cell-cell junctions – a common
site of Hippo kinase complex recruitment – one possible explanation is that CagA antagonizes
Hippo-mediated YAP inhibition by indirectly preventing stabilization of the Hippo complex (Song et
al., 2013). Alternatively, the observation that YAP expression is increased by CagA in a dose-
dependent fashion suggests that *H. pylori* infection may instead circumvent Hippo inhibition by
overwhelming either Hippo's capacity to phospho-inhibit YAP or 14-3-3's capacity to sequester
serine-phosphorylated YAP in the cytoplasm (Li et al., 2018). The observation that *H. pylori*
enhances EMT in gastric epithelia in a CagA-dependent fashion suggests a potential for pathogen-
directed YAP activation to have downstream effects on pathology (Song et al., 2013; Li et al., 2018).
Regardless, these results illustrate that a component of the altered host transcriptome observed during
infection is the product of pathogen manipulation of host transcription factors, rather than a general
host response to the pathogen.

Our finding that chlamydial YAP activation is sensitive to inhibition of bacterial protein
synthesis indicates that one or more chlamydial effectors facilitate YAP activation. One possible
candidate for this effect is the chlamydial inclusion protein IncG, which has been previously shown
to interact with host 14-3-3β (Scidmore and Hackstadt, 2001). Given that 14-3-3-family proteins are
known to act downstream of Hippo-mediated YAP inhibition by binding and sequestering serine-
phosphorylated YAP in the cytosol (Basu et al., 2003; Zhao et al., 2007), potential inhibition of 14-3-
3 by IncG may complement infection-dependent Y357 phosphorylation in decoupling YAP from
Hippo-mediated phospho-inhibition at S127. Additional YAP-activating factors may operate via
enhancement of YAP tyrosine phosphorylation, an event observed to be essential for infection-
mediated YAP translocation. Given that chlamydial YAP phospho-activation is demonstrably
sensitive to inhibition of Src-family kinases, screening for chlamydial modulators of SFKs or
signaling pathways that lead to increased SFK activity may identify additional YAP-activating
effectors.

Although we do not observe chlamydial inhibition of the Hippo kinase cascade at the level of
YAP S127 phosphorylation, we cannot entirely dismiss a role for this negative regulator in
determining levels of nuclear YAP during infection. The recent report of chlamydial disruption of

cell-cell junctions in an organoid model of infection (Dolat and Valdivia, 2021) suggests that YAP
inhibition by Hippo may be relevant in three-dimensional models of infection. Our laboratory has
previously reported on chlamydial modulation of focal adhesion kinase (FAK) (Thwaites et al., 2014;
Pedrosa et al., 2020); given that FAK has been shown to antagonize Hippo-mediated YAP
phosphorylation at S397 (Hu et al., 2017), infection may potentially impact YAP inhibition by Hippo
in this way as well. Though beyond the scope of this initial study, further investigation of chlamydial
YAP activation will likely require use of three-dimensional culture models to determine which of the
extensive portfolio of subcellular structures and mechanical inputs co-opted by the pathogen
ultimately drive enhancement of YAP activity during infection.

The effect of *Chlamydia*-induced YAP nuclear translocation on host gene expression is still
somewhat unclear. Critically, a substantial component of the YAP-responsive gene set was
downregulated in response to infection – an outcome inconsistent with YAP's reported function as a
transcriptional coactivator. One explanation for this result is the activity of transcriptional repressors
induced by infection: while infection-downregulated genes designated by ChEA as YAP targets may
be regulated YAP-cofactor complexes, this event may be superseded by infection-induced repressors
also binding to the same promoters. Ultimately, future work to define the true extent of the
Chlamydia-induced YAP regulon will require comparing the infected host transcriptome reported
here to an equivalent infection in a YAP-negative background. Given that the relatively modest
siRNA-mediated knockdown we report here is consistent with past use of siRNA in End1s
(Govender et al., 2014), generating a CRISPR/Cas9-mediated YAP-knockout from an alternative cell
line may be necessary.

Our results illustrate a mechanism by which chlamydial infection promotes YAP-dependent
gene expression in the host; however, the benefit of this interaction to the pathogen is unclear. YAP
activation in infected epithelial cells can be viewed as a downstream process relevant to
pathogenesis, but not necessarily one beneficial to chlamydial growth in host cells. Given past work
demonstrating *Chlamydia*-infected cells exhibit increased expression of genes encoding pro-fibrotic
cytokines and ECM components (Humphrys et al., 2013; Porcella et al., 2015), and our own
observation that infection induces expression of fibroblast-activating signaling factors (CTGF,
INHBA, BMP2), the potential effect of infection-mediated YAP activation on *Chlamydia*-associated
scarring warrants further study. Fibroblasts in particular are understood to play a critical part in other
types of scar-forming disease (Kendall and Feghali-Bostwick, 2014); however, their contribution to
chlamydial fibrosis has yet to be fully assessed. Understanding the yet uncharacterized effect of
Chlamydia-infected cells on fibroblasts and other uninfected cell types will likely be critical to
elucidating mechanisms of chlamydial fibrosis *in vivo*, as well as contextualizing the broader role of
YAP induction in chlamydial pathogenesis.

5   Acknowledgments

The authors acknowledge the members of the Carabeo laboratory for their critical feedback on
the development of this project. We also acknowledge the generosity of Dr. Ted Hackstadt for
providing the chlamydial strain used in this study, Dr. Marius Sudol for providing pEGFP-C3-
HYAP1, as well as the WSU SMB Genomics Core and UNMC Genomics Core for performing
mRNA library preparation and bulk RNA-sequencing, and the UNMC Tissue Science Facility for
RAFT culture sectioning and slide preparation.

6   Contribution

Our study presents a means by which *C. trachomatis* induces gene expression in host epithelial
cells, providing an explanation for how infection mediates a host transcriptomic response above and
beyond pathogen recognition. We identify chlamydial induction of YAP activity as a principal
mediator of this effect, and thus a potential therapeutic target for *Chlamydia*-associated pathology.
Our finding that infection bypasses Hippo-mediated YAP inhibition by enhancing YAP Y357
phosphorylation in a pathogen-directed, Src-family kinase-dependent fashion corroborates recent
discoveries indicating this relatively novel axis of YAP activation can circumvent its canonical
inhibition by the Hippo kinase cascade. Collectively, our data illustrates an underappreciated aspect
of the host-pathogen interaction: pathogen-directed regulation of transcription factor activity driving
changes in the gene expression of the host.

7   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

8   Author Contributions

Conceptualization, L.C. and R.C.; Methodology, L.C., A.B., and R.C.; Validation, L.C. and
A.B.; Formal Analysis, L.C.; Investigation, L.C. and A.B.; Resources, R.C.; Data Curation, L.C.;
Writing – Original Draft, L.C.; Writing – Review & Editing, R.C. and A.B.; Visualization, L.C.;
Supervision, R.C.; Project Administration, R.C.; Funding Acquisition, R.C.

9   Funding

This work was supported by NIAID grant R01 AI065545 to RAC. The University of Nebraska
DNA Sequencing Core receives partial support from the National Institute for General Medical
Science (NIGMS) INBRE - P20GM103427-19 grant as well as The Fred & Pamela Buffett Cancer
Center Support Grant - P30 CA036727. This publication's contents are the sole responsibility of the
authors, and do not necessarily represent the official views of the NIH.

10   References

Aasen, T., Hodgins, M. B., Edward, M., and Graham, S. V. (2003). The relationship between connexins, gap junctions, tissue architecture and tumour invasion, as studied in a novel in vitro model of HPV-16-associated cervical cancer progression. *Oncogene* 22, 7969–7980. doi: 10.1038/sj.onc.1206709.

Basu, S., Totty, N. F., Irwin, M. S., Sudol, M., and Downward, J. (2003). Akt phosphorylates the Yes-associated protein, YAP, to induce interaction with 14-3-3 and attenuation of p73-mediated apoptosis. *Mol Cell* 11, 11–23. doi: 10.1016/s1097-2765(02)00776-1.

Caldwell, H. D., Kromhout, J., and Schachter, J. (1981). Purification and partial characterization of the major outer membrane protein of Chlamydia trachomatis. *Infect Immun* 31, 1161–1176.

Carlson, J. H., Porcella, S. F., McClarty, G., and Caldwell, H. D. (2005). Comparative genomic analysis of Chlamydia trachomatis oculotropic and genitotropic strains. *Infect. Immun.* 73, 6407–6418. doi: 10.1128/IAI.73.10.6407-6418.2005.

Cheng, W., Shivshankar, P., Zhong, Y., Chen, D., Li, Z., and Zhong, G. (2008). Intracellular Interleukin-1α Mediates Interleukin-8 Production Induced by Chlamydia trachomatis Infection via a Mechanism Independent of Type I Interleukin-1 Receptor. *Infect. Immun.* 76, 942–951. doi: 10.1128/IAI.01313-07.

Chow, J. M., Yonekura, M. L., Richwald, G. A., Greenland, S., Sweet, R. L., and Schachter, J. (1990). The association between Chlamydia trachomatis and ectopic pregnancy. A matched-pair, case-control study. *JAMA* 263, 3164–3167.

Christian, J., Vier, J., Paschen, S. A., and Häcker, G. (2010). Cleavage of the NF-κB Family Protein p65/RelA by the Chlamydial Protease-like Activity Factor (CPAF) Impairs Proinflammatory Signaling in Cells Infected with Chlamydiae *. *Journal of Biological Chemistry* 285, 41320–41327. doi: 10.1074/jbc.M110.152280.

Das, A., Fischer, R. S., Pan, D., and Waterman, C. M. (2016). YAP Nuclear Localization in the Absence of Cell-Cell Contact Is Mediated by a Filamentous Actin-dependent, Myosin II- and Phospho-YAP-independent Pathway during Extracellular Matrix Mechanosensing. *J Biol Chem* 291, 6096–6110. doi: 10.1074/jbc.M115.708313.

Degot, S., Auzan, M., Chapuis, V., Béghin, A., Chadeyras, A., Nelep, C., et al. (2010). Improved Visualization and Quantitative Analysis of Drug Effects Using Micropatterned Cells. *J Vis Exp*, 2514. doi: 10.3791/2514.

Dolat, L., and Valdivia, R. H. (2021). An endometrial organoid model of interactions between Chlamydia and epithelial and immune cells. *Journal of Cell Science* 134, jcs252403. doi: 10.1242/jcs.252403.

Dupont, S., Morsut, L., Aragona, M., Enzo, E., Giulitti, S., Cordenonsi, M., et al. (2011). Role of YAP/TAZ in mechanotransduction. *Nature* 474, 179–183. doi: 10.1038/nature10137.

Eckmann, L., Kagnoff, M. F., and Fierer, J. (1993). Epithelial cells secrete the chemokine
interleukin-8 in response to bacterial entry. *Infect Immun* 61, 4569–4574. doi:
10.1128/iai.61.11.4569-4574.1993.

Edgar, R., Domrachev, M., and Lash, A. E. (2002). Gene Expression Omnibus: NCBI gene
expression and hybridization array data repository. *Nucleic Acids Res* 30, 207–210.

Elbediwy, A., Vanyai, H., Diaz-de-la-Loza, M.-C., Frith, D., Snijders, A. P., and Thompson, B. J.
(2018). Enigma proteins regulate YAP mechanotransduction. *J Cell Sci* 131, jcs221788. doi:
10.1242/jcs.221788.

Elwell, C. A., Kierbel, A., and Engel, J. N. (2011). Species-Specific Interactions of Src Family
Tyrosine Kinases Regulate Chlamydia Intracellular Growth and Trafficking. *mBio* 2, e00082-
11. doi: 10.1128/mBio.00082-11.

Fichorova, R. N., Rheinwald, J. G., and Anderson, D. J. (1997). Generation of Papillomavirus-
Immortalized Cell Lines from Normal Human Ectocervical, Endocervical, and Vaginal
Epithelium that Maintain Expression of Tissue-Specific Differentiation Proteins. *Biol Reprod*
57, 847–855. doi: 10.1095/biolreprod57.4.847.

Futakuchi, A., Inoue, T., Wei, F.-Y., Inoue-Mochita, M., Fujimoto, T., Tomizawa, K., et al. (2018).
YAP/TAZ Are Essential for TGF-β2-Mediated Conjunctival Fibrosis. *Invest Ophthalmol Vis
Sci* 59, 3069–3078. doi: 10.1167/iovs.18-24258.

Gali, Y., Ariën, K. K., Praet, M., Van den Bergh, R., Temmerman, M., Delezay, O., et al. (2010).
Development of an in vitro dual-chamber model of the female genital tract as a screening tool
for epithelial toxicity. *Journal of Virological Methods* 165, 186–197. doi:
10.1016/j.jviromet.2010.01.018.

Govender, Y., Avenant, C., Verhoog, N. J. D., Ray, R. M., Grantham, N. J., Africander, D., et al.
(2014). The injectable-only contraceptive medroxyprogesterone acetate, unlike norethisterone
acetate and progesterone, regulates inflammatory genes in endocervical cells via the
glucocorticoid receptor. *PLoS One* 9, e96497. doi: 10.1371/journal.pone.0096497.

Gumbiner, B. M., and Kim, N.-G. (2014). The Hippo-YAP signaling pathway and contact inhibition
of growth. *J Cell Sci* 127, 709–717. doi: 10.1242/jcs.140103.

Haggerty, C. L., Gottlieb, S. L., Taylor, B. D., Low, N., Xu, F., and Ness, R. B. (2010a). Risk of
Sequelae after *Chlamydia trachomatis* Genital Infection in Women. *The Journal of Infectious
Diseases* 201, 134–155. doi: 10.1086/652395.

Haggerty, C. L., Gottlieb, S. L., Taylor, B. D., Low, N., Xu, F., and Ness, R. B. (2010b). Risk of
Sequelae after Chlamydia trachomatis Genital Infection in Women. *The Journal of Infectious
Diseases* 201, S134–S155. doi: 10.1086/652395.

Hu, J. K.-H., Du, W., Shelton, S. J., Oldham, M. C., DiPersio, C. M., and Klein, O. D. (2017). A
FAK-YAP-mTOR signaling axis regulates stem cell-based tissue renewal in mice. *Cell Stem
Cell* 21, 91-106.e6. doi: 10.1016/j.stem.2017.03.023.

Hu, V. H., Harding-Esch, E. M., Burton, M. J., Bailey, R. L., Kadimpeul, J., and Mabey, D. C. W.
(2010). Epidemiology and control of trachoma: systematic review. *Trop Med Int Health* 15,
673–691. doi: 10.1111/j.1365-3156.2010.02521.x.

Huang, Z., Hu, J., Pan, J., Wang, Y., Hu, G., Zhou, J., et al. (2016). YAP stabilizes SMAD1 and
promotes BMP2-induced neocortical astrocytic differentiation. *Development* 143, 2398–2409.
doi: 10.1242/dev.130658.

Humphrys, M. S., Creasy, T., Sun, Y., Shetty, A. C., Chibucos, M. C., Drabek, E. F., et al. (2013).
Simultaneous Transcriptional Profiling of Bacteria and Their Host Cells. *PLOS ONE* 8,
e80597. doi: 10.1371/journal.pone.0080597.

Johnson, K. A., Lee, J. K., Chen, A. L., Tan, M., and Sütterlin, C. (2015). Induction and inhibition of
CPAF activity during analysis of Chlamydia-infected cells. *Pathog Dis* 73, 1–8. doi:
10.1093/femspd/ftv007.

Kendall, R. T., and Feghali-Bostwick, C. A. (2014). Fibroblasts in fibrosis: novel roles and
mediators. *Frontiers in Pharmacology* 5. Available at:
https://www.frontiersin.org/article/10.3389/fphar.2014.00123 [Accessed March 15, 2022].

Kim, N.-G., and Gumbiner, B. M. (2015). Adhesion to fibronectin regulates Hippo signaling via the
FAK–Src–PI3K pathway. *J Cell Biol* 210, 503–515. doi: 10.1083/jcb.201501025.

Kiviat, N. B., Wølner-Hanssen, P., Eschenbach, D. A., Wasserheit, J. N., Paavonen, J. A., Bell, T. A.,
et al. (1990). Endometrial histopathology in patients with culture-proved upper genital tract
infection and laparoscopically diagnosed acute salpingitis. *Am. J. Surg. Pathol.* 14, 167–175.

Krämer, S., Crauwels, P., Bohn, R., Radzimski, C., Szaszák, M., Klinger, M., et al. (2015). AP-1
Transcription Factor Serves as a Molecular Switch between Chlamydia pneumoniae
Replication and Persistence. *Infect Immun* 83, 2651–2660. doi: 10.1128/IAI.03083-14.

Lachmann, A., Xu, H., Krishnan, J., Berger, S. I., Mazloom, A. R., and Ma'ayan, A. (2010). ChEA:
transcription factor regulation inferred from integrating genome-wide ChIP-X experiments.
Bioinformatics 26, 2438–2444. doi: 10.1093/bioinformatics/btq466.

Lad, S. P., Fukuda, E. Y., Li, J., de la Maza, L. M., and Li, E. (2005). Up-regulation of the
JAK/STAT1 signal pathway during Chlamydia trachomatis infection. *J Immunol* 174, 7186–
7193. doi: 10.4049/jimmunol.174.11.7186.

Lansingh, V. C. (2016). Trachoma. *BMJ Clin Evid* 2016, 0706.

Le Negrate, G., Krieg, A., Faustin, B., Loeffler, M., Godzik, A., Krajewski, S., et al. (2008).
ChlaDub1 of Chlamydia trachomatis suppresses NF-κB activation and inhibits IκBα
ubiquitination and degradation. *Cellular Microbiology* 10, 1879–1892. doi: 10.1111/j.1462-
5822.2008.01178.x.

Lee, J. K., Enciso, G. A., Boassa, D., Chander, C. N., Lou, T. H., Pairawan, S. S., et al. (2018).
Replication-dependent size reduction precedes differentiation in Chlamydia trachomatis. *Nat
Commun* 9, 45. doi: 10.1038/s41467-017-02432-0.

Li, C., Wen, A., Shen, B., Lu, J., Huang, Y., and Chang, Y. (2011). FastCloning: a highly simplified,
purification-free, sequence- and ligation-independent PCR cloning method. *BMC*
Biotechnology 11, 92. doi: 10.1186/1472-6750-11-92.

Li, N., Feng, Y., Hu, Y., He, C., Xie, C., Ouyang, Y., et al. (2018). Helicobacter pylori CagA
promotes epithelial mesenchymal transition in gastric carcinogenesis via triggering oncogenic
YAP pathway. *J Exp Clin Cancer Res* 37, 280. doi: 10.1186/s13046-018-0962-5.

Li, P., Silvis, M. R., Honaker, Y., Lien, W.-H., Arron, S. T., and Vasioukhin, V. (2016). αE-catenin
inhibits a Src-YAP1 oncogenic module that couples tyrosine kinases and the effector of
Hippo signaling pathway. *Genes Dev* 30, 798–811. doi: 10.1101/gad.274951.115.

Mital, J., and Hackstadt, T. (2011). Diverse Requirements for Src-Family Tyrosine Kinases
Distinguish Chlamydial Species. *mBio*. doi: 10.1128/mBio.00031-11.

Mo, J.-S., Yu, F.-X., Gong, R., Brown, J. H., and Guan, K.-L. (2012). Regulation of the Hippo-YAP
pathway by protease-activated receptors (PARs). *Genes Dev* 26, 2138–2143. doi:
10.1101/gad.197582.112.

Ness, R. B., Soper, D. E., Richter, H. E., Randall, H., Peipert, J. F., Nelson, D. B., et al. (2008).
Chlamydia Antibodies, Chlamydia Heat Shock Protein, and Adverse Sequelae After Pelvic
Inflammatory Disease: The PID Evaluation and Clinical Health (PEACH) Study. *Sexually*
Transmitted Diseases 35, 129–135. doi: 10.1097/OLQ.0b013e3181557c25.

Nogueira, A. T., Braun, K. M., and Carabeo, R. A. (2017). Characterization of the Growth of
Chlamydia trachomatis in In Vitro-Generated Stratified Epithelium. *Front Cell Infect*
Microbiol 7, 438. doi: 10.3389/fcimb.2017.00438.

O'Keefe, E. J., Briggaman, R. A., and Herman, B. (1987). Calcium-induced assembly of adherens
junctions in keratinocytes. *J Cell Biol* 105, 807–817. doi: 10.1083/jcb.105.2.807.

Pan, Z., Tian, Y., Zhang, B., Zhang, X., Shi, H., Liang, Z., et al. (2017). YAP signaling in gastric
cancer-derived mesenchymal stem cells is critical for its promoting role in cancer progression.
Int J Oncol 51, 1055–1066. doi: 10.3892/ijo.2017.4101.

Pedrosa, A. T., Murphy, K. N., Nogueira, A. T., Brinkworth, A. J., Thwaites, T. R., Aaron, J., et al.
(2020). A post-invasion role for Chlamydia type III effector TarP in modulating the dynamics
and organization of host cell focal adhesions. *J Biol Chem* 295, 14763–14779. doi:
10.1074/jbc.RA120.015219.

Piccolo, S., Dupont, S., and Cordenonsi, M. (2014). The biology of YAP/TAZ: hippo signaling and
beyond. *Physiol Rev* 94, 1287–1312. doi: 10.1152/physrev.00005.2014.

Pocaterra, A., Romani, P., and Dupont, S. (2020). YAP/TAZ functions and their regulation at a
glance. *Journal of Cell Science* 133. doi: 10.1242/jcs.230425.

Porcella, S. F., Carlson, J. H., Sturdevant, D. E., Sturdevant, G. L., Kanakabandi, K., Virtaneva, K.,
et al. (2015). Transcriptional Profiling of Human Epithelial Cells Infected with Plasmid-
Bearing and Plasmid-Deficient Chlamydia trachomatis. *Infect. Immun.* 83, 534–543. doi:
10.1128/IAI.02764-14.

Price, M. J., Ades, A. E., Soldan, K., Welton, N. J., Macleod, J., Simms, I., et al. (2016). *Pelvic*
inflammatory disease and tubal factor infertility. NIHR Journals Library Available at:
https://www.ncbi.nlm.nih.gov/books/NBK350656/ [Accessed April 7, 2018].

Rasmussen, S. J., Eckmann, L., Quayle, A. J., Shen, L., Zhang, Y. X., Anderson, D. J., et al. (1997).
Secretion of proinflammatory cytokines by epithelial cells in response to Chlamydia infection
suggests a central role for epithelial cells in chlamydial pathogenesis. *J Clin Invest* 99, 77–87.

Scidmore, M. A., and Hackstadt, T. (2001). Mammalian 14-3-3beta associates with the Chlamydia
trachomatis inclusion membrane via its interaction with IncG. *Mol Microbiol* 39, 1638–1650.
doi: 10.1046/j.1365-2958.2001.02355.x.

Smoot, R. L., Werneburg, N. W., Sugihara, T., Hernandez, M. C., Yang, L., Mehner, C., et al.
(2018). Platelet-derived growth factor regulates YAP transcriptional activity via Src family
kinase dependent tyrosine phosphorylation. *Journal of Cellular Biochemistry* 119, 824–836.
doi: https://doi.org/10.1002/jcb.26246.

Snavely, E. A., Kokes, M., Dunn, J. D., Saka, H. A., Nguyen, B. D., Bastidas, R. J., et al. (2014).
Reassessing the role of the secreted protease CPAF in Chlamydia trachomatis infection
through genetic approaches. *Pathog Dis* 71, 336–351. doi: 10.1111/2049-632X.12179.

Song, X., Chen, H.-X., Wang, X.-Y., Deng, X.-Y., Xi, Y.-X., He, Q., et al. (2013). H. pylori-encoded
CagA disrupts tight junctions and induces invasiveness of AGS gastric carcinoma cells via
Cdx2-dependent targeting of Claudin-2. *Cell Immunol* 286, 22–30. doi:
10.1016/j.cellimm.2013.10.008.

Stephens, R. S. (2003). The cellular paradigm of chlamydial pathogenesis. *Trends in Microbiology*
11, 44–51. doi: 10.1016/S0966-842X(02)00011-2.

Strickland, S., Brimer, N., Lyons, C., and Vande Pol, S. B. (2018). Human Papillomavirus E6
interaction with cellular PDZ domain proteins modulates YAP nuclear localization. *Virology*
516, 127–138. doi: 10.1016/j.virol.2018.01.003.

Sugihara, T., Werneburg, N. W., Hernandez, M. C., Yang, L., Kabashima, A., Hirsova, P., et al.
(2018). YAP Tyrosine Phosphorylation and Nuclear Localization in Cholangiocarcinoma
Cells Are Regulated by LCK and Independent of LATS Activity. *Mol Cancer Res* 16, 1556–
1567. doi: 10.1158/1541-7786.MCR-18-0158.

Sun, Y., Zhou, P., Chen, S., Hu, C., Bai, Q., Wu, H., et al. (2017). The JAK/STAT3 signaling
pathway mediates inhibition of host cell apoptosis by Chlamydia psittaci infection. *Pathog*
Dis 75. doi: 10.1093/femspd/ftx088.

Szeto, S. G., Narimatsu, M., Lu, M., He, X., Sidiqi, A. M., Tolosa, M. F., et al. (2016). YAP/TAZ
Are Mechanoregulators of TGF-β-Smad Signaling and Renal Fibrogenesis. *JASN*,
ASN.2015050499. doi: 10.1681/ASN.2015050499.

Taylor, H. R., Burton, M. J., Haddad, D., West, S., and Wright, H. (2014). Trachoma. *The Lancet*
384, 2142–2152. doi: 10.1016/S0140-6736(13)62182-0.

Thwaites, T., Nogueira, A. T., Campeotto, I., Silva, A. P., Grieshaber, S. S., and Carabeo, R. A.
(2014). The Chlamydia Effector TarP Mimics the Mammalian Leucine-Aspartic Acid Motif
of Paxillin to Subvert the Focal Adhesion Kinase during Invasion. *J. Biol. Chem.* 289, 30426–
30442. doi: 10.1074/jbc.M114.604876.

Zagurovskaya, M., Shareef, M. M., Das, A., Reeves, A., Gupta, S., Sudol, M., et al. (2009). EGR-1
forms a complex with YAP-1 and upregulates Bax expression in irradiated prostate carcinoma
cells. *Oncogene* 28, 1121–1131. doi: 10.1038/onc.2008.461.

Zanconato, F., Forcato, M., Battilana, G., Azzolin, L., Quaranta, E., Bodega, B., et al. (2015).
Genome-wide association between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic
growth. *Nat Cell Biol* 17, 1218–1227. doi: 10.1038/ncb3216.

Zhao, B., Wei, X., Li, W., Udan, R. S., Yang, Q., Kim, J., et al. (2007). Inactivation of YAP
oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth
control. *Genes Dev.* 21, 2747–2761. doi: 10.1101/gad.1602907.

11   Data Availability Statement

The bulk RNA-sequencing datasets generated and analyzed for this study can be found in the Gene
Expression Omnibus (Edgar et al., 2002), GEO Series accession number GSE180784
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE180784).

12   Figure Legends

12.1 **Figure 1:** *Chlamydia* infection induces expression of a subset of YAP target genes.

(A, B) Volcano plot of gene expression changes identified via bulk RNA-sequencing of End1/E6E7
immortalized epithelial cells (A) and primary human endocervical epithelial cells (B) during
infection with *Chlamydia trachomatis* (*Ct*) serovar L2 compared to mock infection, at 24 hours post-
infection (hpi). n = 3, with a minimum of $3\text{x}10^7$ unstranded single reads per replicate with a mean
length of 150 bp. All fold changes are relative to the mock-infected control; red dots: false discovery
rate p-value (FDRP) ≤ 0.05, labels: top 20 genes whose expression differed most significantly
(lowest FDRP) from to the mock infection.
(C) Venn diagram of differentially expressed genes identified in (A) and (B) cross-referenced with
the ChIP Enrichment Analysis (ChEA) database of YAP target genes.
(D) Scatter plot of gene expression of YAP-responsive (ChEA), differentially expressed genes in
either *Ct* serovar L2-infected End1/E6E7 cells (End1s, x-axis) or primary human cervical epithelial
cells (HCECs, y-axis). All fold changes are relative to each cell type's respective mock-infected
control; blue line: linear regression model of correlation; grey shading: 95% confidence interval. R^2
and p-values calculated using Pearson's correlation.
(E) Heatmap of YAP target gene expression in *Ct* serovar L2-infected End1s (left columns) and
HCECs (right columns). All fold changes are relative to each cell type's respective mock-infected
control; only genes differentially expressed (FDRP ≤ 0.05) in both cell types are shown.

12.2 Figure 2: *Chlamydia* infection promotes YAP nuclear translocation.

722
(A) Expression of CTGF, INHBA, BMP2, and and MMP9 at 24 hpi in mock- and *Ct* L2-infected
End1 cells, as measured by RT-qPCR. n = 6 biological replicates; fold changes are relative to mean
expression of the mock-infected and untreated control. Whiskers: minimum to maximum; asterisks:
p-values ≤ 0.05, using pairwise Student's t-tests and Bonferroni's correction for multiple
comparisons.
(B) Expression of YAP1 and CTGF at 24 hpi in mock- and *Ct* L2-infected End1 cells transfected
with non-targeting (NT) or YAP1-targeting siRNA (10 nM for 24 h prior to infection), as measured
by RT-qPCR. n = 3 biological replicates; fold changes are relative to mean expression of the mock-
infected and untreated control. Whiskers: minimum to maximum; asterisks: p-values ≤ 0.05, using
pairwise Student's t-tests and Bonferroni's correction for multiple comparisons.
(C) Representative micrographs of YAP (green) translocation into the nuclei (blue) of confluent
mock- and *Ct* L2-infected End1 cells at 24 hpi. Scale bar: 20 μm.
(D) Quantification of YAP nuclear translocation in (A) as a ratio of nuclear to cytosolic YAP
fluorescence. n = 5 biological replicates, 50 cells measured per sample. Black bars: group means;
asterisks: p-values ≤ 0.05, using pairwise Wilcoxon rank sum tests and Bonferroni's correction for
multiple comparisons.
(E) Representative micrographs of YAP (green) translocation into the nuclei (blue) of confluent
mock- and *Ct* L2-infected primary human cervical epithelial cells at 24 hpi. Scale bar: 20 μm.
(F) Quantification of YAP nuclear translocation in (C) as a ratio of nuclear to cytosolic YAP
fluorescence. n = 5 biological replicates, 50 cells measured per sample. Black bars: group means;
asterisks: p-values ≤ 0.05, using pairwise Wilcoxon rank sum tests and Bonferroni's correction for

multiple comparisons.

(G) Quantification of YAP nuclear translocation at 2, 4, 8, 12, 18, and 24 hpi in confluent mock- and

Ct L2-infected End1 cells as a ratio of nuclear to cytosolic YAP fluorescence. n = 5 biological

replicates, 50 cells measured per sample. Blue dots: mock-infected cells, red dots: *Ct* L2-infected

cells, black bars: group means, asterisks: p-value ≤ 0.05, using pairwise Wilcoxon rank-sum tests and

Bonferroni's correction for multiple comparisons.

(H) Representative micrographs of YAP (green) translocation at 18 hpi in confluent mock- and *Ct*

L2-infected End1 cells treated with chloramphenicol (Cm, 50 µg/mL for 1 h at 17 hpi) or DMSO.

Scale bar: 20 µm.

(I) Quantification of YAP nuclear translocation in (B). n = 3 biological replicates, 50 cells measured

per sample. Black bars: group means; asterisks: p-values ≤ 0.05, using pairwise Wilcoxon rank-sum

tests and Bonferroni's correction for multiple comparisons.

12.3   **Figure 3:** *Chlamydia* infection bypasses Hippo-mediated YAP inhibition by enhancing YAP
phosphorylation at Y357.

(A) Representative micrographs of YAP (green) translocation into the nuclei (blue) at 24 hpi of
mock- and *Ct* L2-infected End1 cells cultured in small cell clusters using 30 μm (1-2 cells) and 45
μm (2-4 cells) micropatterns. Asterisks: chlamydial inclusions; scale bar: 20 μm.
(B) Quantification of YAP nuclear translocation in (A) as a ratio of nuclear to cytosolic YAP
fluorescence. n = 3 biological replicates, with a minimum of 5 fields measured per sample. Black
bars: group means; asterisks: p-values $\leq$ 0.05, using pairwise Wilcoxon rank sum tests and
Bonferroni's correction for multiple comparisons.
(C) Representative Western blot of YAP phosphorylation of mock- and *Ct* L2-infected or sodium
orthovanadate-treated (Na$_3$VO$_4$, 10 mM for 1 h at 23 hpi) End1 cells at 24 hpi.
(D, E) Densitometric quantification of YAP phosphorylation at S127 (D) and Y357 (E), normalized
to total YAP. n = 3 biological replicates; whiskers: minimum to maximum; asterisks: p-values $\leq$ 0.05,
using pairwise Student's t-tests and Bonferroni's correction for multiple comparisons.

12.4 **Figure 4:** Infection-induced phosphorylation at Y357 is required for YAP nuclear
translocation.

(A) Representative Western blot of YAP tyrosine phosphorylation of mock- and *Ct* L2-infected End1
cells at 24 hpi, treated with the Src-family kinase inhibitor PP2 (10 μM for 24 h), or DMSO.
(B) Representative micrographs of YAP translocation at 24 hpi in confluent mock- and *Ct* L2-
infected End1 cells treated with the Src-family kinase inhibitor PP2 (10 μM for 24 h), or DMSO.
Scale bar: 20 μm.
(C) Quantification of YAP nuclear translocation in (B). n = 4 biological replicates, 50 cells measured
per sample. Black bars: group means; asterisks: p-values ≤ 0.05, using pairwise Wilcoxon rank sum
tests and Bonferroni's correction for multiple comparisons.
(D) Representative micrographs of the translocation of FLAG-tagged YAP wild-type or the phospho-
dead Y357F mutant (green) into the nuclei (blue) of confluent mock- and *Ct* L2-infected End1 cells
at 24 hpi. Scale bar: 20 μm, dotted lines: nuclear area of transfected cells.
(E) Quantification of YAP nuclear translocation in (D). n = 4 biological replicates, 50 cells measured
per sample. Black bars: group means; asterisks: p-values ≤ 0.05, using pairwise Wilcoxon rank sum
tests and Bonferroni's correction for multiple comparisons.

13   Tables

13.1   Table 1: Cross-referencing of RNA-sequencing results with ChEA transcription factor target
790   database reveals induction of transcription factor targets.

Transcription Factor	Induced Targets	Repressed Targets	Mean Fold Change
IRF1	55	0	2.6
STAT1	34	9	1.44
STAT3	255	80	2.46
SMAD2	126	31	9.11
SMAD3	162	57	4.89
TEAD4	67	29	1.21
EGR1	292	101	2.11
YAP1	81	36	1.24
MYCN	61	56	-62.65
MYBL1	2	4	-0.49
CRX	21	18	-2.02