Chimpanzees organize their social relationships like humans

Diego Escribano,[1] Victoria Doldán-Martelli,[1] Katherine A. Cronin,[2]
Daniel B.M. Haun,[3,4] Edwin J.C. van Leeuwen,[5,6] José A. Cuesta,[1,7]
Angel Sánchez[1,7,*]

[1]Grupo Interdisciplinar de Sistemas Complejos (GISC), Departamento de Matemáticas,
Universidad Carlos III de Madrid, 28911 Leganés, Spain

[2]Animal Welfare Science Program, Lincoln Park Zoo,Chicago, IL 60614, USA

[3]Department for Comparative Cultural Psychology,
Max Planck Institute for Evolutionary Anthropology,
Deutscher Platz 6, 04103 Leipzig, Germany

[4]Leipzig Research Centre for Early Child Development, Leipzig University,
Jahnallee 59, 04109 Leipzig, Germany

[5]Behavioural Ecology and Ecophysiology Group, Department of Biology,
University of Antwerp, 2610 Wilrijk, Belgium

[6]Animal Behaviour and Cognition, Department of Biology, University Utrecht,
3584 CH Utrecht, The Netherlands

[7]Instituto de Biocomputación y Física de Sistemas Complejos (BIFI),
Universidad de Zaragoza, 50018 Zaragoza, Spain

[*]To whom correspondence should be addressed; E-mail: anxo@math.uc3m.es

**Human relationships are structured in a set of layers, ordered from higher
(intimate relationships) to lower (acquaintances) emotional and cognitive in-
tensity. This structure arises from the limits of our cognitive capacity and the
different amounts of resources required by different relationships. However, it**

is unknown whether nonhuman primate species organize their affiliative relationships following the same pattern. We here show that the time chimpanzees devote to grooming other individuals is well described by the same model used for human relationships, supporting the existence of similar social signatures for both humans and chimpanzees. Furthermore, the relationship structure depends on group size as predicted by the model, the proportion of high intensity connections being larger for smaller groups.

One-sentence summary: Chimpanzees distribute their time to groom other individuals in the same manner as humans structure their relationships.

Social network analysis has been a very active field for about a century, revealing the complex set of relationships that connect individuals (*1, 2*). Among the main objects of interest of social network analysis are personal or egonetworks, which consist of the social networks surrounding selected actors (*3*). A very general observation is that human egonetworks show a layered structure where each layer corresponds to relationships of different emotional closeness (*4–6*). These layers have a definite emotional closeness: there is a layer of very close friends, a subsequent one of good friends, and so on. It is convenient to introduce the concept of nested circles, i.e., the sets of all the relationships up to a certain closeness. Typical circles established in the literature contain 5, 15, 50 and 150 individuals—with a scaling ~ 3 between a circle and the next one. There is also evidence for a subsequent circle, formed by acquaintances, of about 500-600 people (*7*).

Social networks have also been studied in a diverse array of species, including mammals, birds, fish, amphibians, reptiles and invertebrates (*8–13*). In this context, the study of nonhuman

primate social networks is of particular interest in light of the complexity of their societies, the variability between species, and their evolutionary proximity to us (*14*). Layered structures have been reported in both the distribution of primate social group sizes (*15*) and in groups of mammals living in multilevel social systems (mainly baboons, chimpanzees, elephants and dolphins) (*16,17*). However, while this suggests a possible connection to the layers of emotional closeness observed in humans, those results are not about individual egonetworks but about group-level social structures. To date, nobody has considered whether a continuous analysis of nonhuman primate social interactions reveals an underlying structure consistent with the structure that emerges in humans due to inherently limited resources of cognition and time.

To fill this gap, in this paper we present strong evidence that chimpanzees organize their relationships very much like humans do by means of a continuous version (*18*) of the theory introduced in (*19*), consisting of a resource allocation model based on two widely accepted assumptions: the capacity that an individual can invest in social relationships is finite, and relationships of a different intensity carry different costs. This mathematical approach allows us to advance our thinking beyond circles and assign a continuum value to a relationship, which is more reflective of real life and can include, for example, frequency of contact (*20*), number of messages exchanged (*21*), or duration of time spent together (*21*). The formalism developed in (*18*) was applied to face-to-face contact time (*22*), number of messages between Facebook users (*23*), and number of phone calls (*21*), showing a structure similar to that arising when intensities are regarded as discrete categories. This implies that it is not necessary to arbitrarily categorize the data to unveil its structure. As a consequence, egonetworks turn out to be characterized by a new universal scale parameter η, which plays the role of (and is consistent with) the scale factor ~ 3 typically found in the discrete setup. We here apply the same formalism to grooming data extracted from over four years of observations of four groups of chimpanzees living in the Chimfunshi Wildlife Orphanage in Zambia, taking grooming as a proxy of the

effort devoted to pairwise relationships. Grooming behavior is characterized by one individual manually or orally manipulating the hair or skin of another individual. While this behavior does serve a hygienic function, grooming is well-known to facilitate and reflect social bonding between individual chimpanzees (*24*). As we will see below, our results confirm that the time chimpanzees devote to grooming other individuals is well described by the continuous probability distribution predicted by the model, supporting the existence of similar social signatures for both humans and chimpanzees.

Generally speaking, the amount of grooming between primates is considered to be an indicator of their relationship quality (*25*). Therefore, we apply the formalism summarized above to grooming data of chimpanzees living in the Chimfunshi Wildlife Orphanage in Zambia between 2015 and 2019. At this sanctuary, chimpanzees live in four different populations without any interaction between individuals from different groups. The number of chimpanzees living in each group differs considerably (groups 1–4, $n = 26, 60, 11, 14$, respectively), and after preparing the data the number of individuals considered in each group for the analysis is reduced (groups 1–4, $n = 21, 32, 10, 10$, respectively). Full information about the chimpanzee population studied is provided in the Supporting Material (SM), *Materials and Methods*.

To analyze the data, we resort to the theoretical approach in (*18*), where the results for the discrete case (*19*) were extended to relationships characterized by continuous values. Briefly, the discrete approach assumes that L is the total number of relationships in an ego-network and σ is the average cognitive cost of a relationship. Relationships belong to r different categories, each of them bearing a different cost $s_{\max} = s_1 > s_2 > \cdots > s_r = s_{\min}$. As described in detail in (*19*), using a maximum entropy approach it is possible to show that the number of relationships in one circle divided by the previous, smaller one, behaves approximately as

$$\frac{\chi_{k+1}}{\chi_k} \sim \begin{cases} e^\mu, & \mu \to \infty, \\ 1, & \mu \to -\infty. \end{cases} \tag{1}$$

where $\mu \equiv \hat{\mu}(s_{max} - s_{min})/(r-1)$, with $\hat{\mu}$ a Lagrange multiplier connected to the cognitive capacity constraint σ. Therefore, the circles satisfy an approximate scaling relation; in particular, for $\mu \approx 1$ the usual value of 3 found on empirical data is recovered. On the other hand, the theory also predicts a so-called "inverse" regime, when $\mu < 0$, in which most of the relationships are in the closest circle. This second behavior had not been described prior to the publication of (*19*), when it was checked against empirical data of small migrant communities, confirming its existence.

In the continuum case (*18*), circles are defined as the fraction of links $\chi(t)$ whose "distance" to the individual is not larger than a specified value t ($0 \leq t \leq 1$). Their scaling ratio turns out to be controlled by a new parameter, η, the equivalent of μ (in fact, it can be shown that $\eta \approx (r-1)(e^{\mu} - 1)$). In this continuous approach, the separation between the two regimes, the normal and the inverted ones, also takes place at $\eta = 0$. On the other hand, setting the number of circles to the usual value $r = 4$ and assuming, as empirically observed, that $e^{\mu} \approx 3$ (*5*), we then find $\eta \approx 6$. More details can be found in the SM, *Theoretical background*, and in the original references (*18, 19*).

With the above approach in mind, given a dataset of relationships with continuous weights, the scaling parameter η can be estimated using the maximum-likelihood method (see SM, *Parameter estimation*). The type of fits of the function $\chi(t)$, giving the size of the continuous circles, to the data on individual chimpanzees is exemplified in Fig. 1 (plots of the fits for all the individuals considered in the study are provided in the *Supplementary Material*). The plots clearly show that the fits are not perfect, but on the other hand most points lie within the 95% confidence interval for the fitted distribution, and those that do not are not far from it. As the *Supplementary Material* shows, there are examples of better and worse fits than those shown here—which have been selected because they had more data for the fitting. One should bear in mind that the chimpanzee data are quite noisy because they have been obtained from chim-

panzees living in large, naturalistic enclosures that lead to varying levels of animal visibility, and data have been collected by many different observers over the four-year period (see SM, *Materials and Methods*). Under those circumstances, the fits can be actually considered to be very good, and in fact if we compare them to those reported in (*18*), they are of a similar quality. Therefore, we can conclude that the continuum theory is a good description of how a chimpanzee distributes the time it devotes to grooming other individuals.

Our analysis of the results for the parameters characterizing all the individuals studied is summarized in Fig. 2. The values for the parameter η obtained from the fits have a mode of approximately $\eta = 4$, while the mean value is close to the mode except for Group 2 where it is closer to 6. The range of obtained values for η falls within the range of expected values, and the mode indicates that typically the scaling ratio of the circles for chimpanzees would be somewhat smaller than for humans, except in Group 2. We believe that the reason for the difference of this group with respect to the others arises from the fact that it is distinctly larger than the rest, both pre- and post-filtering, and that this allows group members to develop a richer social life as far as grooming is concerned—i.e., it allows individuals to devote small intervals of time to grooming many others, leading to higher values of η and more low intensity relationships.

The histograms presented show that there are no fits yielding negative values of η, which would indicate the existence of chimpanzees with an inverted structure of relationships, like those observed in humans (*19*). However, the values of η close to zero are on the border between regimes, so they should give rise to a larger fraction of individuals in the inner part of the distribution $\chi(t)$. Figure 3 confirms that this is indeed the case by showing the relationship structure and the $\chi(t)$ fitted function for two very different examples. On one side of the η scale we have the distribution of grooming times by Kit, whose $\eta = 0.14$ is rather typical of a structure that is intermediate between regimes. Kit devotes quite some time to grooming Kambo, Commander, Bobby and Val, and also a noticeable amount of time to other chimpanzees. In-

terestingly, this agrees with the fact that Kit is in group 4, the smallest one, in agreement with the situation in which the inverted regime is expected to arise: not so many possible individuals available to groom. On the opposite extreme of the η scale, Fig. 3 shows the results for Genny, in Group 1, with $\eta = 10.3$. In this case, we have a situation which is typical of the normal regime, with a lot of grooming devoted to her baby, some amount to her daughter Gonzaga, and very small amounts to many other individuals. Once again, this is to be expected in so far as Genny lives in Group 1, where there are many chimpanzees she can relate with. Therefore, the results corresponding to different values of η are also quite similar to those found for humans (*18*).

The results reported in this study show that chimpanzees appear to organize their grooming time in very good agreement with the prediction of the continuum resource allocation theory applied to humans in (*18*). In other words, chimpanzees distribute the time they devote to grooming other individuals in their group very much like humans organize their relationships when the relationship intensity is treated as a continuous value. These findings are in line with previous accounts that grooming can be considered a resource allocation problem (*26–30*). As we see in humans, some chimpanzees solve this allocation problem by investing large amounts of grooming in a few other individuals (more so when the group is small), while others invest small amounts of grooming in many other individuals. In other words, they show the same two allocation strategies observed in human relationships and in the same situations, with inverted structures being more likely in small groups. We note that these findings corroborate grooming as an expression of friendship in non-human primates (*31*), yet more importantly indicate that different social strategies might be at play among chimpanzees, dependent on their immediate group structure. Our results show that chimpanzees living in larger groups employ their social capital differently than chimpanzees in smaller groups, like in humans (*19, 32*). This suggests that chimpanzees navigate their social environment flexibly, distributing their social resources across many group members when needed, yet investing more intensely in a

few others when possible. In this light, any social forces that affect levels of group cohesiveness (e.g., to what extent a group forms a whole versus a modular, sub-grouping structure) indirectly shape a primate's resource allocation strategy, a link we only know from the science of human sociology (*1*).

The novel finding here is that a nonhuman primate species, chimpanzees, appears to organize their affiliative relationships, or friendships, following the same pattern that we previously characterized for humans. Indeed, it appears that the mean group sizes of all primate species follow the same pattern as we see in the circles of human egonetworks (*15*). Furthermore, the global network structures of some primate species have a similar internal structure (*16, 17, 33*). By means of our continuum analysis of grooming data, we show that egonetworks in chimpanzees also exhibit a specific organization in terms of (the continuous equivalent of) circles. Our results thus establish that social networks in humans and chimpanzees show similar relationship structure. Further research could leverage the behavioral data of other primate species to determine whether the continuum analysis of egonetworks reveals a consistent pattern in other species living in large social groups.

References and Notes

1. Wellman, B., Berkowitz, S.D. (1988) *Structural Analysis in the Social Sciences 2: Social Structures: a Network Approach* (Cambridge University Press, Cambridge, UK).

2. Freeman, L.C. (2004) *The Development of Social Network Analysis.* (Empirical Press, Vancouver, Canada).

3. McCarty, C., Lubbers, M.J., Vacca, R., Molina, J.L. (2019) *Conducting Personal Network Research: A Practical Guide* (The Guilford Press, New York, USA).

4. Zhou, W. X., Sornette, D., Hill, R. A., Dunbar, R. I. M. (2005) Discrete hierarchical organization of social group sizes. *Proc. R. Soc. B* **272**, 439–444.

5. Dunbar, R. I. M., Arnaboldi, V., Conti, M., Passarella, A. (2015) The structure of online social networks mirrors those in the offline world. *Social Networks* **43**, 39–47.

6. Dunbar, R. I. M., Sosis, R. (2017). Optimising human community sizes. *Evol. Human Beh.* **39**, 106–111.

7. Lubbers, M. J., Molina, J. L., Valenzuela, H. (2019) When networks speak volumes: Variation in the size of broader acquaintanceship networks. *Social Networks* **59**, 55–69.

8. Krause, J., James, R., Franks, D. W., Croft, D. P. (2015) *Animal Social Networks* (Oxford University Press, Oxford, UK).

9. Dunbar, R.I.M. (2020) Structure and function in human and primate social networks: implications for diffusion, network stability and health. *Proc. R. Soc. A* **476**, 20200446.

10. Brask, J.B, Ellis, S., Croft, D.P. (2021) Animal social networks: An introduction for complex system scientists. *J. Complex Networks* **9**, cnab001.

11. Godfrey, S.S., Moore, J.A., Nelson, N.J., Bulla, C.M. (2010) Social network structure and parasite infection patterns in a territorial reptile, the tuatara (*Sphenodon punctatus*). *Int. J. Parasitology* **40, 1575–1585**.

12. Sah, P., Méndez, J.D., Bansal, S. (2019) A multi-species repository of social networks. *Sci. Data* **6**, 44.

13. Lamb, S.D.,Altobelli, J.T., Easton, L.J., Godfrey, S.S., Bishop, P.J. (2021) Captive Hamilton's frog (*Leiopelma hamiltoni*) associates non-randomly under retreat sites: preliminary insights into their social networks. *New Zealand J. Zool.*, 1–16.

14. Kappeler, P.M., van Schaik, C. P. (2002) Evolution of primate social systems. *Int. J. Primatol.* **23**, 707–740.

15. Dunbar R.I.M., MacCarron P., Shultz S. (2018) Primate social group sizes exhibit a regular scaling pattern with natural attractors. *Biol. Lett.* **14**, 20170490.

16. Kudo H., Dunbar R.I.M. (2001) Neocortex size and social network size in primates. *Anim. Behav.* **62**, 711–722.

17. Hill R., Bentley A., Dunbar R.I.M. (2008) Network scaling reveals consistent fractal pattern in hierarchical mammalian societies. *Biol. Lett.* **4**, 74–751.

18. Tamarit, I., Sánchez, A., Cuesta, J.A. (2022). Beyond Dunbar circles: a continuous description of social relationships and resource allocation. *Sci. Rep.* **12**, 2287 (2022).

19. Tamarit, I., Cuesta, J.A., Dunbar, R.I.M., Sánchez, A. (2018) Cognitive resource allocation determines the organization of personal networks. *Proc. Nat'l. Acad. Sci. USA* **115**, 8316–8321 (2018).

20. Roberts, S. G., Dunbar, R. I., Pollet, T. V., Kuppens, T. (2009) Exploring variation in active network size: Constraints and ego characteristics. *Soc. Netw.* **31**, 138–146.

21. Saramäki J., Leicht, E.A., López, E., Roberts, S.G.B., Reed-Tsochas, F., Dunbar, R.I.M. (2014) Persistence of social signatures in human communication. *Proc. Natl. Acad. Sci. USA* **111**, 942–947.

22. Isella, L., Stehlé, J., Barrat, A., Cattuto, C., Pinton, J.-F., Van den Broeck, W. (2011) What's in a crowd? Analysis of face-to-face behavioral networks. *J. Theor. Biol.* **271**, 166–180.

23. Arnaboldi, V., Conti, M., Passarella, A., Pezzoni, F. (2012) Analysis of ego network structure in online social networks. In: *2012 International Conference on Social Computing (SocialCom) Privacy, Security, Risk and Trust (PASSAT)*, 31–40 (IEEE).

24. Watts, D.P. (2002) Reciprocity and interchange in the social relationships of wild male chimpanzees. *Behavior* **139**, 343–370.

25. Massen J.J.M., Sterck E.H.M., de Vos, H. (2010) Close social associations in animals and humans: functions and mechanisms of friendship. *Behaviour* **147**, 1379–1412.

26. Dunbar, R. I. M. (1992) Time: a hidden constraint on the behavioural ecology of baboons. *Behavioral Ecol. Sociobiol.* **31**, 35–49.

27. Waal, F. B. M. (1997) The chimpanzee's service economy: food for grooming. *Evol. Human Behav.* **18**, 375–386.

28. Barrett, L., Henzi, S. P., Weingrill, T., Lycett, J. E., Hill, R. A. (1999) Market forces predict grooming reciprocity in female baboons. *Proc. R. Soc. London B* **266**, 665–670.

29. Fruteau, C., Lemoine, S., Hellard, E., van Damme, E., Noë, R. (2011) When females trade grooming for grooming: testing partner control and partner choice models of cooperation in two primate species. *Animal Behav.* **81**, 1223–1230.

30. Kaburu, S. S. K., Newton-Fisher, N. E. (2015) Egalitarian despots: hierarchy steepness, reciprocity and the grooming-trade model in wild chimpanzees, Pan troglodytes. *Animal Behav.* **99**, 61–71.

31. Silk, J. B. (2002). Using the F-word in primatology. Behaviour, 139(2-3), 421–446.

32. Bernard HR, Killworth PD (1973) On the social structure of an ocean-going researchvessel and other important things. *Social Sci. Res.* **2**, 145–184.

33. Shultz, S., Dunbar, R.I.M. (2008) Social complexity and the fractal structure of group size in primate social evolution. *Biol. Rev.* **96**, 1889–1906.

34. Ron, T., McGrew, W. C. (1988). Ecological assessment for a chimpanzee rehabilitation project in Northern Zambia. *Primate Conservation* **9**, 37–41.

35. van Leeuwen, E. J. C., Cronin, K. A., Haun, D. B. M. (2019). Reply to Farine and Aplin: Chimpanzees choose their association and interaction partners. *Proceedings of the National Academy of Sciences of the United States of America* 116(34), 16676-–16677.

36. van Leeuwen, E. J. C., Cronin, K. A., Haun, D. B. M. (2018). Population-specific social dynamics in chimpanzees. *Proceedings of the National Academy of Sciences of the United States of America* **115(45)**, 11393–11400.

37. Lincoln Park Zoo. (2020). ZooMonitor (version 3.2.) [Mobile app]. https://zoomonitor.org.

Acknowledgments: We are thankful to Robin Dunbar and José L. Molina for discussions on our results.

Funding:

MCIN/AEI/ 10.13039/501100011033 and "ERDF A way of making Europe" grant BASIC (PGC2018-098186-B-I00) (DE, VD-M, JAC, AS).

Flanders Research Foundation (FWO) postdoctoral fellowship (EJCvL).

Max Planck Society for the Advancement of Science (DBMH).

Author contributions:

Conceptualization: DE, VD-M, JAC, AS

Resources: KAC, DBMH, EJCvL

Data curation: DE, KAC

Formal analysis: DE, VD-M

Software: DE, KAC

Visualization: DE

Investigation: DE, VD-M, KAC, DBMH, EJCvL, JAC, AS

Project administration: AS

Writing — original draft: DE, AS

Writing — review and editing: DE, KAC, DBMH, EJCvL, JAC, AS

Competing interests: The authors declare no competing interests

Data availability: Data are available upon reasonable request from KAC, DBMH, EJCvL

Supplementary Materials

Materials and Methods

Figures S1 to S12

Tables S1 to S4

References (34–37)

Figure 1: **Examples of fittings for an individual of each group.** Selected individuals are those for which there were more available data points. From left to right, individuals belong to groups 1, 2, 3, and 4. Shown in each case is $\chi(t)$, the fraction of links whose "distance" to the individual is not larger than t. Red dots are actual data, representing the number of individuals who receive no more grooming than a fraction t of the maximum. The blue dashed line is the fitted function $\chi(t)$, and the blue-shadowed region is the interval of confidence.

Figure 2: **Histograms for η parameter distribution in each group.** The red dashed line represents the change of regime $\eta = 0$ and the blue dashed line the mean value for each group.

Figure 3: **Examples of the relationship between the η parameter and the proportion of relationships of different intensity.** $\chi(t)$ functions and egonetworks for Genny (Group 1 and $\eta = 10.3$) and Kit (Group 4 and $\eta = 0.14$). Arrows connect the focal individual with those it grooms, while the width of the arrows represents the total amount of time devoted to grooming a specific chimpanzee. Orange ovals represent females, blue ones represent males.