Humans trade-off energetic cost with fatigue avoidance while walking

Kirsty A. McDonald[1,2,3,*], Joseph P. Cusumano[4], Andrew Hieronymi[5], Jonas Rubenson[2,3,6]

[1]Department of Exercise Physiology, School of Health Sciences, University of New South Wales, Sydney, New South Wales, Australia, 2052

[2]School of Human Sciences, The University of Western Australia, Crawley, Perth, Western Australia, Australia, 6009

[3]Biomechanics Laboratory, Department of Kinesiology, The Pennsylvania State University, University Park, Pennsylvania, United States of America, 16802

[4]Department of Engineering Science and Mechanics, The Pennsylvania State University, University Park, Pennsylvania, United States of America, 16802

[5]School of Visual Arts, The Pennsylvania State University, University Park, Pennsylvania, United States of America, 16802

[6]Integrative and Biomedical Physiology, The Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, Pennsylvania, United States of America, 16802

[*]Kirsty A. McDonald

Email: kirsty.mcdonald@unsw.edu.au

Abstract

Metabolic cost minimization is widely regarded to underscore the neural control of locomotion and
the evolution of human bipedalism. Yet, minimizing muscle fatigue often outperforms energy
minimization in computational predictions of human gait. Discerning the relative importance of
these key criteria in human walking has proved elusive, in part, because they have not been
empirically decoupled. Here we explicitly decouple metabolic cost and muscle fatigue (estimated
from electromyography-derived muscle activations) by pitting them against one another using two
distinct gait tasks. When experiencing these competing costs, participants (*N*=10) chose the task
that systematically protected against muscle fatigue at the expense of higher metabolic power (19%
penalty, $p<0.05$). This held true when fatigue was expressed as the sum of muscle activations
squared (66% saving, $p<0.05$), and as the maximal activation across muscles (44% saving,
$p<0.05$), both of which penalize overburdening any individual muscle and thus indicate fatigue
avoidance. Activation cost expressed as the sum of muscle volume-normalized activation more
closely models energy use and was also protected by the participants' decision (23% saving,
$p<0.05$) demonstrating that muscle activation was, at best, an inaccurate proxy signal for metabolic
energy. Energy minimization was only observed when there was no adverse effect on muscle
fatigue. By decoupling metabolic and fatigue costs, we provide among the first empirical evidence
of humans embracing non-energetic optimality in favor of a clearly defined neuromuscular fatigue
objective. This finding indicates that fatigue may well be a key factor dictating human walking
behavior and its evolution.

Introduction

Humans often move in ways that save metabolic energy. For example, preferred walking speed (1–5), step frequency (6–12), step width (13) and arm swing (14,15) are classic examples known to minimize the metabolic cost of locomotion. More recently, it has been demonstrated that humans continue to exhibit energy-favoring behaviors, even after normal gait mechanics are disrupted (8,16,17). Given the consistency of these observations in human locomotion, and the selective advantage that they purportedly bestow, it is argued that whole-body energy minimization is a central factor dictating human gait behavior and its evolution (18,19).

However, whether metabolic cost minimization is itself the principal, overriding, optimality criterion during gait remains a matter of debate. For example, it has been proposed that other factors, such as fatigue avoidance, may be prioritized in the control scheme of locomotion (20–22), with low metabolic cost arising as a byproduct. Similarly, others suggest that economical human gait arises not from minimization of metabolic cost alone, but rather via the control of metabolic energy in conjunction with several additional optimality criteria (23,24).

Of the possible non-energetic optimality criteria, a compelling argument can be made for a muscle activation-based control signal in human locomotion. Several groups have developed successful simulations of human locomotion based on minimizing total muscle activation. This approach has proven particularly powerful in predicting realistic locomotor mechanics and energetics, often outperforming simulations based on minimizing total energy cost alone (21,22,25,26). For example, Miller et al. (21) discovered that more realistic gait biomechanics and whole-body metabolic cost could be predicted by minimizing muscle activation compared to metabolic cost itself, despite the longstanding view that humans consistently prioritize energy optimizing gaits (5,11). Likewise, Falisse et al. (23), and more recently, Veerkamp et al. (24), demonstrated that the inclusion of muscle activation in a multi-objective performance criterion is essential for generating the most physiologically realistic simulations of human gait. There also exists experimental data that point to muscle activation as a control signal in human locomotion. For example, the metabolic cost of walking on inclines of up to 10% can be predicted using muscle activation data (27). Preferred stride frequencies (28,29) and the walk-run transition (30–32)—a fundamental locomotor behavior that is not always tracked by whole-body energy cost (20)—can also be predicted by a muscle activation minimizing criterion. Similarly, self-selected cadence in cycling has been found to coincide with low muscle activity (33), more so than with minimization of metabolic energy expenditure (34,35). Together, these simulation and experimental data indicate that muscle activation may indeed be among the primary optimality criteria governing human gait.

On closer examination of the aforementioned studies, the activation cost functions that have been most successful in predicting gait mechanics and energetics (often better than energy optimization) penalize high activations in any given muscle. These cost functions minimize the sum of individual muscle activations (A) raised to an integer power (e.g., A^2, A^3 *etc.*), where the activation of a muscle is defined as the fraction (between 0 and 1) of the total muscle fibers activated (21,22). Other successful simulations of muscle control have used a similar, but more extreme formulation, whereby the maximum activation of any single muscle is minimized (the min-max model; (36)). As above, the resulting recruitment pattern is one that favors a uniform distribution of activations across muscles as opposed to high activations among a select few muscles—even in cases where the latter results in lower total energy consumption (22,26). This avoidance of high muscle activation that arises from non-uniform muscle recruitment has been regarded as a fatigue-minimizing (endurance maximizing) strategy (22,37,38). Thus, the nervous system may prioritize a 'division of labor' across muscles that minimizes muscle fatigue, as opposed to a strict 'economy of labor'. In this scenario, economical gait may arise not as the principal optimization criterion, but rather secondary to a control paradigm favoring even muscle recruitment (22).

Establishing the mechanism(s) driving locomotor behavior is key to understanding the control of human gait and the resulting energetics. Previous investigations have largely been limited to simulation studies that assign objective functions (optimality criteria) *a priori* and subsequently

assess their efficacy based on their ability to reproduce a high number of human gait features. To
the best of our knowledge, no study has empirically addressed whether whole-body energy use or
fatigue-like muscle activation costs (e.g., sum of A^2; (22)) are prioritized during locomotion.
Therefore, we designed an experiment to pit fatigue-like muscle activation costs against the total
metabolic cost during walking by disrupting the distribution of activation across the lower limb
muscles. Importantly, in this design, moving with low energy expenditure while simultaneously
favoring a low muscle activation cost (i.e., low muscle fatigue) was unfeasible, therefore providing
an opportunity to empirically test whether one criterion was favored over the other.

Materials and methods

Participants
Ten healthy adults (5F, 5M; age 31.1±7.2 years; mass 69.6±11.1 kg; height 1.70±0.07 m;
mean±SD) participated in this study after providing informed consent. The study was approved by
the Institutional Review Board at The Pennsylvania State University and received reciprocal
approval by the Human Research Ethics Committee at the University of Western Australia.
Volunteers were required to be free from medical conditions that could make moderate-intensity
exercise unsafe (e.g., heart, respiratory and/or musculoskeletal conditions).

Experimental design overview
The primary objective of our experimental design was to create two competing conditions
(Competing-Cost-Pairs) that required the participants to select between either a low whole-body
metabolic cost or a low activation cost (see below for cost functions). Importantly, in selecting
between the Competing-Cost-Pairs, the metabolic cost and activation cost could not be
simultaneously reduced. Through pilot testing, two locomotor conditions were established to meet
this experimental design goal. The first condition, crouch walking on a level treadmill, was
determined to elicit a moderate metabolic cost and a high activation cost due to the uneven
distribution of muscle recruitment across lower limb muscles. The second condition, incline walking,
included five levels of treadmill incline that ranged from 0-24% and that predominantly taxed
metabolic cost. As the incline level was increased, a Competing-Cost-Pair was determined by
identifying the incline level that elicited a metabolic cost exceeding that of crouch walking, while at
the same time having a lower muscle activation cost (see Fig. 1).

To establish the Competing-Cost-Pairs, five independent comparison trials were performed in a
random order between flat (i.e., level treadmill) crouch walking and upright walking on one of the
five incline levels (0%, 6%, 12%, 18% or 24%; Fig. 1). For each comparison, participants were
asked to explore both crouch walking and upright walking on the assigned incline for three minutes.
After this time, they selected their preferred state (crouch or upright incline walking), in which they
then walked for an additional five minutes. The steepest incline level that was selected over crouch
walking was referred to as the pre-transition incline. The first incline level for which the crouch
walking was preferred was referred to as the post-transition incline (data were collected at the end
of the five comparison trials for all incline levels that were rejected in favor of crouch walking).

Crouch walking
Crouch walking was achieved by having participants walk on a level treadmill (0% incline) while
avoiding contact between their body and a virtual ceiling, with the specific instruction to maintain
their trunk and head in a normal, upright walking posture (i.e., crouched, but not hunched or with
lowered head). A motion sensing device (Xbox Kinect, Microsoft, WA, USA), situated behind the
participant, was programmed to detect the participant's body shape and produce a two-dimensional
body silhouette. This silhouette was displayed on a large screen in front of the treadmill together
with the virtual ceiling projection (Fig. 2). The ceiling height was programed to equal 93% of the
participant's standing height. This height was established in pilot testing to result in a crouch posture
that elicited the desired increases in our metric of activation cost (see below). During periods of
silhouette-ceiling collision, the outline of the silhouette was filled blue to alert the participant of
contact. The number and duration of collisions was recorded for each trial. The real-time motion-

tracking and visual feedback was implemented in a custom program developed with the
openFrameworks open-source C++ toolkit.

At the beginning of each experiment, participants were provided a crouch walking familiarization
trial. Participants walked at a set speed of 1.0 m s^{-1}, crouching below the virtual ceiling. After three
minutes of crouch walking with unlimited silhouette-ceiling collisions, the participant was informed
that, to end the familiarization session, they must perform one minute of walking with no recorded
collisions.

All participants achieved the one-minute benchmark and progressed to the next phase of data
collection, in which their crouch preferred walking speed (PWS$_{C/W}$; m s^{-1}) was assessed in
accordance with the method outlined by Dingwell and Marin (39). All crouch and incline walking
trials described below were administered at the PWS$_{C/W}$ so as not to disadvantage metabolic
energy use in the crouch condition: i.e., it was assumed that preferred walking speed selection
would minimize metabolic cost of transport for the crouch condition (2,40).

Comparison trials
Following PWS$_{C/W}$ determination, a five-minute trial of crouch walking was recorded (initial crouch).
In a randomized order, five levels of upright incline walking (on treadmill inclines of 0%, 6%, 12%,
18% and 24%) were then pitted against crouch walking, one by one. Each comparison trial (crouch
versus a single incline level) commenced with 30 seconds of crouch walking and 30 seconds of
incline walking (the order of which was reversed for consecutive comparison trials), followed by a
further two minutes where the participant was free to explore both the crouch and incline. During
the two-minute exploratory period, transition between the crouch and incline walking was verbally
requested by the participant and manually initiated by the investigator, taking less than 10 seconds
to implement.

At the conclusion of the exploratory period, the participant verbally expressed the condition (crouch
or incline) in which they would prefer to walk for the following five minutes. Their preference was
then imposed for a five-minute period, during which metabolic, surface electromyography (EMG),
and ground reaction force (GRF) data were collected (details below). On discussions with
participants at the conclusion of the experiment, it was confirmed that, together, the one minute
(total) of imposed walking in the two conditions and the two minutes of free exploration was
adequate for all participants to conclusively decide on a preferred condition.

Once crouch walking had been compared to the five incline levels, an additional five-minute crouch
walking trial was recollected to account for fatigue/familiarization effects (final crouch). Any
outstanding incline levels for which data was not collected as part of the comparison trials (because
crouch walking was preferred) were then completed at PWS$_{C/W}$, each for five minutes (randomized
order), to provide insight into the participant's motivation for transitioning (rejecting these inclines).
To minimize any persistent muscle fatigue effects, participants commenced each consecutive trial
when they had returned to a baseline (resting) level of exertion. When describing the protocol,
participants were made aware that they may be asked to complete at most 12 walking trials.
However, the nature of the trials following the comparison trials were not revealed to the participants
to ensure it did not influence their prior decisions.

Whole-body energetics and electromyography measurements
Breath-by-breath oxygen uptake and carbon dioxide production were sampled using a portable
metabolic system (Cosmed K4b^2, Rome, Italy). A portable metabolic device was necessary to
facilitate adherence to the required crouching posture and free regulation of body position on the
treadmill. The final minute of data was used to compute a steady-state mass-specific metabolic
power ($C_{met,P}$; W kg^{-1}) for each trial (details below). Surface EMG recordings were made
synchronously from gluteus maximus (Gmax), biceps femoris (BF), rectus femoris (RF), vastus
medialis (VM), medial gastrocnemius (MG), soleus (SOL) and tibialis anterior (TA) of the right leg,
using silver-tipped surface electrodes placed according to SENIAM guidelines (41). Signals were
recorded using a Bortec Octopus AMT-8 system (Calgary, Canada). Twenty strides of EMG data
were collected at 2000 Hz. Ground reaction forces from the custom-built instrumented treadmill

were also obtained (again sampled at 2000 Hz) to determine foot contacts and were later used to
crop the EMG data into strides during data processing. Metabolic, EMG and GRF data were
collected during the initial and final crouch trials, all comparison trials where the incline was
selected, and any outstanding incline trials (thus, there were seven total recording trials).

Data processing
Data processing was performed in MATLAB 2018a (MathWorks, Natick, USA) using in-built
functions and custom scripts.

Metabolic power $\left(C_{met,P}\right)$
The rate of oxygen uptake ($\dot{V}O_2$; L s^{-1}) and carbon dioxide production ($\dot{V}CO_2$; L s^{-1}) were averaged
from the final minute of each trial and the following stoichiometric equation was applied to determine
metabolic power (42,43):

201
$$C_{met,P} = \frac{\left(16.89(\dot{V}O_2) + 4.82(\dot{V}CO_2)\right)1000}{M}$$

Where M represents participant body mass. Because speed was constant across trial, we did not
convert to a cost per distance traveled.

Muscle activation
Two participants' activation data were eliminated from further analyses due to notable and
persistent signal artefact in one of their processed electromyograms. For the remaining participants
(*N*=8), the raw EMG data were DC offset, band-pass filtered (20-350 Hz) to remove any movement
artefact and high frequency noise, and full-wave rectified. A linear envelope was applied to the
rectified data using a low-pass filter with 6 Hz cut-off frequency (EMG_{env}). All data filtering utilized
fourth-order Butterworth underdamped filters (44). For each of the seven trials for which data was
collected, five strides ($j = 1,2,\dots5$) of EMG_{env} data per muscle ($i = 1,2,\dots7$) were individually
integrated to produce scalar muscle activation values for each muscle, for each stride $\left(a_{ij}\right)$, as
below:

214
$$a_{ij} \triangleq \int_0^{tf} (EMG_{env})_{ij}\, dt$$

where $(EMG_{env})_{ij}$ represents the processed linear envelop of the i^{th} muscle for the j^{th} stride, 0 is
the time at the initial contact of the right foot and tf is the time at the consecutive contact of the
same (ipsilateral) foot determined from force plate thresholds (2% of peak signal).

To take into consideration fluctuations in the activation integral $\left(a_{ij}\right)$ that can occur due to changes
in stride time $\left(T_j\right)$, each stride's muscle activation values were expressed as a rate $\left(\frac{a_{ij}}{T_j}\right)$. A
normalized (unitless) activation $\left(A_{ij}\right)$ was computed for each muscle, for each stride by normalizing
by the average value obtained from the five strides of the 0% incline trial:

222
$$A_{ij} \triangleq \frac{a_{ij}}{T_j} \bigg/ \left(\frac{1}{5}\sum_{j=1}^{5}\frac{a^0{}_{ij}}{T^0{}_j}\right)$$

where a^0 and T^0 are the activation integral and stride time of the 0% incline trial, respectively.

Activation cost functions
We used what have previously been classified as 'fatigue-like' activation cost functions that
penalized large, fatigue-inducing, muscle activations irrespective of the size of the muscle (21,22).
Our cost functions are based on those previously adopted, minimizing a) the sum of individual

muscle activations raised to an integer power $(p > 1)$ (22) or b) the individual maximal muscle
activation across all muscles $(p \to \infty)$ (Rasmussen et al., 2001).

a) Sum of activations squared $(\overline{C}_{a^2})$**;** For each trial, a total muscle activation cost was computed
as the sum of the seven muscles' normalized activations (A_{ij}) squared. This value was then
averaged across five strides:

233
$$\overline{C}_{a^2} \triangleq \frac{1}{5} \sum_{j=1}^{5} \sum_{i=1}^{7} \left(A_{ij}^2\right)$$

234 **b) Maximal muscle activation** $(\overline{C}_{a,max})$**;** At the extreme end of fatigue-like activation cost
235 functions, i.e., the integer power limit $(p \to \infty)$, the cost function depends only on the maximally
236 activated muscle (22). Thus, for each trial, the maximum activation cost was computed as the
237 maximum value (max_i) from the seven muscles' normalized activation (A_{ij}). This value was then
238 averaged across five strides:

239
$$\overline{C}_{a,max} \triangleq \frac{1}{5} \sum_{j=1}^{5} \left[max_i\left(A_{ij}\right)\right]$$

By adopting these two cost functions we capture a range of fatigue-like activation cost formulations,
with the latter most strongly penalizing high activation in individual muscles. Both activation costs
represent fatigue through a combined effect of the amount of fibers active within the muscles (a_{ij}),
as well as the rate at which they are used during continuous walking $\left(\frac{1}{T_j}\right)$, similar to the fatigue
activation cost function in Miller et al. (21).

c) Sum of the volume-weighted activations $(\overline{C}_{a,vol})$**;** In addition to the two aforementioned
fatigue-like activation costs, we also included an 'effort-like' activation cost (22). This cost function
does not penalize an uneven distribution of muscle activation. Rather, it takes into consideration
active muscle volume, a key determinant of muscle energy expenditure (45,46), by weighting the
activations by muscle size. Therefore, unlike the fatigue-like functions, this cost is more sensitive
to activation of large muscles (22). This cost was included to assess the possibility that muscle
activation serves as a proxy for metabolic energy expenditure (47).

The normalized activation rate (A_{ij}) for each muscle was weighted (w_i) according to the muscle
volumes documented in the supplementary material of Handsfield et al. (48). These weightings
were 0.33, 0.08, 0.10, 0.17, 0.10, 0.17 and 0.05 for Gmax, BF, RF, VM, MG, SOL, and TA,
respectively, and represent each muscle's volume as a proportion of the sum of all seven muscles.
The seven volume-weighted normalized activation rates were then summed, and this value was
averaged across five strides:

258
$$\overline{C}_{a,vol} \triangleq \frac{1}{5} \sum_{j=1}^{5} \sum_{i=1}^{7} w_i \left(A_{ij}\right)$$

Data analysis
For each metric described above, five values were initially analyzed; mean crouch (the mean value
of the initial crouch and final crouch trials), initial crouch, final crouch, pre-transition incline (the
steepest incline level which was selected over crouch walking) and post-transition incline (the
lowest incline level that was rejected in favor of crouch walking). Whether the mean, initial or final
crouch data were compared to the pre- and post- transition inclines had no bearing on the
interpretation of the results discussed herein. Thus, the mean crouch data is represented by all
future references to the crouch walking data.

Competing-Cost-Pairs
Each participant's mean crouch walking data was compared to the pre-transition incline to
determine whether Competing-Cost-Pairs had been established (i.e., the two trials were designed
to provide mutually exclusive $C_{met,P}$ and $\overline{C}_{a^2}$ advantages). We accepted a Competing-Cost-Pair
when $C_{met,P}$ decreased between the pre-transition incline and crouch walking trials, and $\overline{C}_{a^2}$
increased. The $C_{met,P}$ and $\overline{C}_{a^2}$ advantages were assessed because our aim was to compare
metabolic and fatigue-like muscle activation costs, and $\overline{C}_{a^2}$ represents the less extreme (least
biased) of our fatigue-like cost functions. A secondary analysis was then performed to determine
whether Competing-Cost-Pairs were also present between $C_{met,P}$ and $\overline{C}_{a,max}$, and $C_{met,P}$ and $\overline{C}_{a,vol}$.

Statistical analyses
Summary statistics (group mean±SD) were computed for crouch walking performance variables
(number and duration of silhouette-ceiling collisions), and for key cost metrics ($C_{met,P}$, $\overline{C}_{a^2}$, $\overline{C}_{a,max}$
and $\overline{C}_{a,vol}$) during the crouch and incline walking trials. Normality was assessed using the Shapiro-
Wilk test. One-way repeated measures analysis of variance (ANOVA) tests were used to confirm
main effects of the normally distributed incline data (0, 6, 12, 18, 24% inclines) for $C_{met,P}$, $\overline{C}_{a^2}$, $\overline{C}_{a,max}$
and $\overline{C}_{a,vol}$.

For the crouch, pre-transition incline and post-transition incline comparisons, key cost metrics were
each analyzed via a one-way repeated measures ANOVA (normally distributed) or Friedman test
(non-normally distributed). *Post hoc* testing was then administered as paired-samples t-tests
(normally distributed) or Wilcoxon tests (non-normally distributed). Results were compared against
an adjusted alpha level (p=0.017; Bonferroni correction). All other analyses employed a standard
0.05 alpha level. In group-level analyses, all available data (*N*=10 for $C_{met,P}$ and *N*=8 for $\overline{C}_{a^2}$, $\overline{C}_{a,max}$,
$\overline{C}_{a,vol}$) was used. Paired samples t-tests/Wilcoxon tests analyzed $C_{met,P}$, $\overline{C}_{a^2}$, $\overline{C}_{a,max}$ and $\overline{C}_{a,vol}$ in
the sub-sample of participants for whom a Competing-Cost-Pair was successfully established
(*N*=7).

To discern whether the use of the initial, final or mean crouch data to represent crouch walking
$C_{met,P}$, $\overline{C}_{a^2}$, $\overline{C}_{a,max}$ and $\overline{C}_{a,vol}$ would make a difference to our interpretation of results, all relevant
statistical tests (above) were performed three times: i) using the initial crouch data, ii) using the final
crouch data, and iii) using the mean of the initial and final crouch data. No difference in
interpretation was present and so mean crouch data is presented in the Results section below.
Initial and final crouch data comparisons can be found in the Supplementary Material.

Any non-normally distributed data is denoted by a caret (^), in the Results section below. SPSS
26.0 (IBM, Chicago, USA) was used to perform all statistical analyses.

Results

Gait selections
The current experiment used the selection of one locomotor mode over another—in this case,
walking in a crouch posture versus walking upright at one of five incline levels (0, 6, 12, 18 and
24%)—to reveal underlying prioritization objectives. Thus, in total, five decisions were made by our
participants, i.e., one for each incline level. At low levels of incline (0-12%), all participants (*N*=10)
chose the incline over the crouch. At 18% incline, the decision divided our sample; seven
participants continued to select the incline, while three preferred the crouch (Fig. 3A). Those three
participants also chose crouch over the 24% incline. The remaining seven participants all chose
crouch walking over walking on the 24% incline.

The task performance of crouch walking was assessed by the total number and duration of
silhouette-ceiling collisions during a single five-minute trial. To determine potential learning effects,
these metrics were recorded at the beginning (initial crouch trial) and end (final crouch trial) of the
data collection session. Total number (start: 1.2±1.5; end: 2.3±2.4) and duration (start: 0.11±0.15

seconds; end: 0.22±0.25 seconds) of collisions did not change significantly (p=0.131^ and
p=0.075^, respectively).

Incline and crouch effects on activation and energy costs

All key cost metrics ($C_{met,P}$, $\overline{C}_{a^2}$, $\overline{C}_{a,max}$ and $\overline{C}_{a,vol}$) increased with increasing incline level (Fig. 3A-
D). This was confirmed by one-way repeated measures ANOVAs, which demonstrated a main
effect of incline for $C_{met,P}$ (p<0.001), $\overline{C}_{a^2}$ (p=0.001), $\overline{C}_{a,max}$ (p<0.001), and $\overline{C}_{a,vol}$ (p<0.001).

Crouch walking disrupted the distribution of muscle activation compared to upright walking, such
that select muscles had proportionately higher activation values. In particular, Figure 4 highlights
the increased activation requirements of knee extensor musculature (vastus medialis and rectus
femoris) to facilitate the crouch gait, while the steepest incline gaits are achieved with a more even
distribution of activation.

When statistical comparisons were made between the crouch, pre-transition incline and post-
transition incline, the one-way repeated measures ANOVAs/Friedman tests demonstrated a main
effect for $C_{met,P}$ (p<0.001, N=10), $\overline{C}_{a^2}$ (p=0.008^; N=8), $\overline{C}_{a,max}$ (p=0.001, N=8) and $\overline{C}_{a,vol}$ (p=0.003;
N=8; Fig. 3E-H).

Competing-Cost-Pairs

Competing-Cost-Pairs (i.e., two trials providing mutually exclusive $C_{met,P}$ and $\overline{C}_{a^2}$ advantages) were
effectively established for seven of the eight participants for whom we acquired a complete set of
EMG data (Table 1; Fig. 5). Competing-Cost-Pairs could also be established for these participants
between $C_{met,P}$ and $\overline{C}_{a,max}$, and $C_{met,P}$ and $\overline{C}_{a,vol}$ (Table 1).

All seven participants (1-4 and 8-10) chose incline walking over crouch walking and thus selected
low muscle activation at the expense of a high $C_{met,P}$. The average $\overline{C}_{a^2}$ advantage in the Competing-
Cost-Pairs was 66% (N=7; p=0.018^; range: 31 to 95%) and for $\overline{C}_{a,max}$ was 44% (N=7; p=0.003;
range: 15 to 89%; Table 1). The average $C_{met,P}$ penalty was 19% (N=7; p=0.002; range: 4% to
31%; Table 1). Participant #6 had a $C_{met,P}$ that was nearly identical in the crouch and incline
condition (1% difference) and therefore did not present a clear energetic advantage/disadvantage
for gait selection (Table 1). However, Participant #6 achieved the group's largest activation
advantage by selecting the incline condition (210% reduction in $\overline{C}_{a^2}$; 103% reduction in $\overline{C}_{a,max}$). We
also found an advantage in the effort-like activation cost ($\overline{C}_{a,vol}$) when the participants selected
incline walking over crouch walking (Table 1). The average $\overline{C}_{a,vol}$ advantage in the Competing-
Cost-Pairs was 23% (N=7; p=0.007; range: 8 to 38%).

In addition to the Competing-Cost-Pairs, prioritization objectives can be further assessed from
comparing the group mean $C_{met,P}$ (N=10) and activation costs (N=8) of the crouch walking and pre-
and post-transition incline conditions (Fig. 3E-H). Crouch walking incurred a $C_{met,P}$ of 7.84±1.38 W
kg^{-1}, which was significantly lower than both the pre- (9.46±0.91 W kg^{-1}; p=0.007) and post-
(11.92±1.31 W kg^{-1}; p=0.005) transition incline levels, with the pre- and post-transition inclines
being significantly different from one another (p=0.005; Fig. 3E). In contrast, the $\overline{C}_{a^2}$ and $\overline{C}_{a,max}$ for
the crouch walking were significantly greater (84%; p=0.012^ and 52% p=0.006, respectively) than
the pre-transition incline (Fig. 3F-G). These data show, at a group level, a selection (pre-transition
incline) for lower fatigue-like activation costs at the expense of higher $C_{met,P}$. We also found that
the $\overline{C}_{a^2}$ and $\overline{C}_{a,max}$ in crouch walking were statistically equivalent to the post-transition incline
(p=0.263^ and p=0.214, respectively), indicating that the disadvantage in the fatigue-like activation
costs was no longer present when crouch walking was finally selected (crouch walking versus post-
transition incline walking). The effort-like activation metric ($\overline{C}_{a,vol}$ followed a similar pattern (Fig.
3H). $\overline{C}_{a,vol}$ for crouch walking was significantly greater (27%) than for the pre-transition incline
(p=0.003) but not for the post-transition incline (p=0.779).

Discussion

Metabolic energy expenditure is widely regarded as a principal determinant of animal locomotor
behavior (5,8,11,49). However, simulation studies of human locomotion suggest muscle activation
may also be a key control parameter (21,22). It is difficult to discern how these criteria are weighted
in the control scheme of locomotion due to their highly coupled nature. The current study utilized
an experimental design that systematically pitted whole-body metabolic cost against muscle
activation costs. In these competing conditions, global energy expenditure was clearly not
prioritized, and our empirical data provide promising support for muscle activity prioritization.

Energy expenditure is not always the principal objective in human locomotion
There are a handful of previous studies in which humans have been observed to expend more
energy than minimally required for both upper and lower limb movement tasks (40,50–56). These
studies suggest that energy minimization may be task-dependent. Perhaps the most well
documented example of non-energetically optimal movement is self-selected cycling cadence (for
a review see Vercruyssen and Brisswalter, 2010). In the context of gait, Yandell and Zelik (56) show
inconsistent prioritization of metabolic cost when participants used their preferred step frequency
during barefoot walking. The authors propose this may be related to the discomfort/pain associated
with an unshod gait (56). Some evidence also suggests humans do not prioritize metabolic cost
minimization when moving in less stable environments, e.g., walking downhill (50) (but this remains
inconclusive (57)) or walking with experimentally imposed asymmetrical leg lengths (40). Humans
have also been observed to perform walking tasks in more effortful ways to eliminate a cognitive
burden (51). Furthermore, there exist several studies of human movement in which immediate,
independent detection and/or acute prioritization of the energetic minima is not evident
(8,40,58,59). The costs driving gait selection in the aforementioned studies remain speculative. The
current study is, in contrast, among the first to explicitly and experimentally demonstrate an
alternative criterion, namely muscle activation, that is prioritized over metabolic cost in the control
scheme of locomotion (at least under the conditions studied here).

While it is clear that a $C_{met,P}$ penalty occurs with the selection of incline walking in the Competing-
Cost-Pairs (and in the pre-transition incline selection more generally; Fig. 3E), we do not suggest
that energy minimization is unimportant to the control of walking, even in less common situations
such as the crouch walking investigated in this study. Indeed, in the final incline condition (post
transition incline), when the $C_{met,P}$ was the highest and the activation penalty of crouch walking was
reduced or eliminated, participants opted to save energy by selecting crouch walking. Rather, our
study supports the idea that locomotion tasks are regulated in a multi-objective manner that
includes metabolic cost as one important, but not the sole, factor. Taken together, our data across
all conditions suggest that a priority for energy minimization is not an inevitability, but also that as
$C_{met,P}$ increases so might the relative weighting of $C_{met,P}$ in the control scheme of walking.

Does a 'division of muscle labor' dictate gait control?
The non-volume-weighted exponential activation cost function, $\overline{C}_{a^2}$, penalizes high (fatiguing)
activations in any single muscle and thus favors an even muscle recruitment strategy (22). The
hypothesis that gait is under a control scheme that minimizes muscle fatigue is not new, but has
been mostly limited to simulation studies (21,22). To the best of our knowledge, the current study
is the first to observe this phenomenon empirically. Our finding that subjects protected their $\overline{C}_{a^2}$ at
the direct expense of $C_{met,P}$ was observed both in the individual participants' Competing-Cost-Pair
trials (Table 1), and also at a group (average) level (Fig. 3E-F).

The clear preference for protecting $\overline{C}_{a^2}$ over energy cost suggests a strong control objective for
maintaining an even muscle recruitment and thus avoiding high, fatiguing, activations that can
cause local muscle exhaustion (22). This is also evident from the more extreme cost function,
$\overline{C}_{a,max}$, which showed that all participants' gait selection in the Competing-Cost-Pairs protected
their would-be maximally activated muscle, in line with a *min/max* cost function (36). This further
suggests that the decision to avoid crouch walking may, in fact, have been specifically predicated

on not exhausting the primary contributing muscle. Even if highly activated individual muscles represent a small fraction of the total muscle mass, they can have an important limit on performance because maximally fatigued muscles set the bottle neck for the endurance of the whole muscle system (22,60).

Further support for activation-driven gait selection comes from our observations of the post-transition incline gait (i.e., when crouch walking was finally chosen). When crouch walking is compared to the post-transition incline, we observe a diminished activation advantage. This can be seen in the group average data, where similar and not statistically different $\overline{C}_{a^2}$, $\overline{C}_{a,max}$ and $\overline{C}_{a,vol}$ values were observed between conditions (Fig. 3F-H). This suggests that not only avoiding, but also selecting crouch walking, involved information from muscle activation. It may be that only when there is minimal/no activation disadvantage, do participants accept a $C_{met,P}$ advantage (whether this was selecting incline over crouch walking, or vice versa, selecting crouch walking over very steep incline walking). It is also worth pointing out that the high $\overline{C}_{a^2}$ cost associated with crouch walking and the steepest incline walking resulted from different muscle recruitment patterns (Fig. 4); high activations in crouch walking are primarily in the vastus medialis (for 6/8 participants), while in the post-transition incline the muscles that exhibit the high activations are more variable between subjects.

Why should muscle activation be prioritized over energy cost? One answer may be that minimizing high activations that fatigue a given muscle will likely permit longer movement duration, a performance criterion that is regarded to be fundamental to the evolution of human bipedalism and with known ecological relevance (61–63). For instance, to effectively travel long distances in persistence hunting (regarded as a key feature of human evolution; 62) there is a clear selective advantage to resist fatigue. Muscle activation is a compelling control signal for sensing a division of labor among muscles and for avoiding overburdening individual muscles. It is thought that muscle effort can be readily sensed using central (e.g., via an efference copy) and peripheral (e.g., Group III and IV muscle afferents) mechanisms (64,65).

Does muscle activation represent a proxy for metabolic energy use?
Notwithstanding the clear acceptance of a $C_{met,P}$ penalty, could it be possible that the nervous system was attempting to prioritize total metabolic energy expenditure using a proxy sensor, but failed? In their recent work, Wong and colleagues (66) concluded that blood-gas receptors are an unlikely sensor for whole-body locomotor energy expenditure and raised the possibility of a muscle-level sensor for metabolic cost. Muscle activation is a promising candidate for sensing energy cost because it is related to the fraction of the muscle that is metabolically active and because activation itself exacts a substantial metabolic cost in the form of calcium pumping (67–69). If activation serves as a proxy sensor for energy expenditure, the high volume-weighted activation signal associated with crouch walking may have 'tricked' the nervous system into registering a higher metabolic cost than that which was actually present. This could occur, in part, due to the inflated activation contribution of the large (high volume) knee-extensor muscles (Fig. 4). It is worth noting that in tasks where energy minimization is ubiquitously demonstrated (e.g., steady level walking), muscle activation and whole-body metabolic costs respond similarly to changing task parameters [e.g., step frequency; (11,28)], thus indicating that the muscle activations can, indeed, serve as a proxy for the energy consumption. This was, however, not the case in the present study. Both $\overline{C}_{a^2}$, $\overline{C}_{a,max}$, and $\overline{C}_{a,vol}$ were disassociated from $C_{met,P}$, with $\overline{C}_{a,vol}$ considered a proxy for energy use because it takes into account active muscle volume (22)—a key determinant of locomotor energy use (46,69,70).

Thus, if activation was used as a proxy for energy, it resulted is a misrepresentation of the relative energy cost of the crouch versus incline walking conditions. This scenario affects our interpretation of the energy optimization hypothesis in important ways. First, it would indicate that locomotor energy sensing is imperfect and may be cruder than previously thought. This implies that high-precision, rapid, energy sensing and subsequent gait control may be confined to common and predictable steady-state movements (e.g., steady level ground walking), and, in-turn, that energy

optimization may not be particularly robust outside of a narrow range of conditions. Other
experiments that have altered the mechanics of walking in novel ways have found that subjects
often need guidance to discover energy minimizing movements (8,16). This suggests that while
energy minimization may be achieved *eventually*, after sufficient motor learning has taken place, it
may not be prioritized *acutely*. Whether trial-and-error learning can bring about versatile energy
sensing across a myriad of natural conditions for human locomotion remains unclear. This question
will prove key, both for practical reasons (e.g., development of assistive devices), as well as
informing theories into the evolution of human locomotor energetics.

Among the most interesting questions that remains is *how* the central nervous system uses muscle-
level signaling to determine the physiological state of the body. For example, does the central
nervous system achieve this by summing the activity of all muscles involved in a given task,
weighted according to their size and tracked over a given distance or time period, and finally
referencing this against a threshold representing global (whole-body) effort or energy use (i.e.,
$\overline{C}_{a,vol}$)? Or, at the other end of the spectrum, is the control signal based on assessing individual
muscle activation levels against local thresholds unique to each actuator (i.e., $\overline{C}_{a,max}$)? Luu et al.
(71) provide support for the latter strategy. The authors observed that when elbow flexors and knee
extensors were isometrically contracted in unison (at 25, 50 and 100% of their individual maximal
voluntary contraction values), participants' ventilation did not increase to a level that reflected the
cumulative effect of contracting the individual muscle groups. In fact, the ventilatory response did
not change significantly between the single and combined muscle group contractions at any
contraction intensity level (71).

We posit a strategy assessing individual muscle activation levels against local thresholds is also a
simpler control strategy for the following reasons. First, $\overline{C}_{a,max}$ (and also, to a lesser extent, $\overline{C}_{a^2}$)
has a computational advantage, limiting the synthesis of information required by the central nervous
system. Individual muscle activation thus presents a promising sensory modality when considering
the credit-assignment problem (66,72), which refers to the direct or indirect nature of the signal(s)
used by the central nervous system to steer gait selection behavior. The internal monitoring of
individual muscle activations to minimize muscle fatigue is an example of a highly direct signal-to-
optimality criterion coupling. Conversely, weighting and summing the individual activations requires
the additional collation of information and is therefore a less direct signaling method for ascribing
locomotor 'cost'. The latter would therefore likely require greater computational effort and time.
Whilst we are not aware of any literature explicitly detailing preferences for computationally
inexpensive control strategies in locomotion, a reduction in cognitive load at the expense of physical
effort was previously observed by Rosenbaum and colleagues (51).

We also do not, at present, have an effective way to rank the likelihood that one activation
parameter is a more likely target of control than the other, although we do note that the pre-
transition incline selection saw participants avoid greater $\overline{C}_{a^2}$ and $\overline{C}_{a,max}$ penalties than that
associated with $\overline{C}_{a,vol}$ (84% and 52% versus 27%, respectively; *N*=8). Simulation studies that have
established both mass- or volume-weighted activation and distribution cost functions also support
this perspective, with more realistic gait patterns predicted by a non-volume-weighted or low
volume-weighted muscle activation criterion (21,22). Irrespective of what optimality criterion
activation minimization serves (e.g., a proxy for metabolic cost minimization, improved distribution
of muscle effort, fatigue minimization, etc.), activation itself seems like a reasonable control target
that can utilize relatively simple computational cost strategies.

Alternate explanations for the observed gait selection
Despite the systematic prioritization of a gait with low muscle activation in the Competing-Cost-
Pairs, we cannot rule out alternate hypotheses for the observed gait selection. For example,
discomfort, pain, or other psychological motivators of gait selection behaviors (56,73) could, in
theory, be at the source of the selection of incline walking over crouch walking in the Competing-
Cost-Pairs. Clearly delineating the role of discomfort versus activation *per se* in our observed gait
selections is challenging because the two are related. In crouch walking, muscles with

disproportionately high activations are prone to fatigue, which is known to cause discomfort (74).
High muscle activations may also result in high local mechanical stresses that could trigger a pain
response. It is conceivable, therefore, that high muscle activations are the proxy signals used in a
neural computation of a 'comfort' cost. In this regard, the minimization of muscle activation in the
Competing-Cost-Pairs could underlie a higher-level comfort objective.

Limitations

The current study captured the activity of seven lower limb muscles that play a key role in human
locomotion. Expanding upon previous empirical studies (47), we also include a weighting factor
when determining the total activation cost. However, incorporating a greater selection of muscles
when computing $\overline{C}_{a^2}$ and $\overline{C}_{a,vol}$ would likely provide further insight into prioritization behaviors.

While $C_{met,P}$ is susceptible to changes in both walking speed and step frequency (5,11), the latter
was not imposed for any condition, and the former may have provided a metabolic advantage to
crouch walking because all trials occurred at the preferred speed established for crouch walking.
Yet, irrespective of this, participants still accepted a metabolic penalty through their pre-transition
incline selection. It must be noted that whether one variable was closer to its physiological
maximum than the other (e.g., $\dot{V}O_2max$ versus maximal activation) was not measured in the current
study. How close a variable is to its maximum may impact the priority for minimizing that variable.
In this sense it is possible that selectively impairing other (more) important optimality criteria through
energetic minimization will only occur in situations where a very high metabolic cost needs to be
avoided. We acknowledge that the relative importance of activation and $C_{met,P}$ may also be different
in steady-state level gait, and that it is likely task- and condition-dependent.

It is of course possible that other physiological signals exist whose distribution across muscles may
parallel that of activation, for example afferent signals from muscle metabolism receptors (class III
and IV afferents), or proprioceptive signals from muscle spindles and/or Golgi tendon organs.
These may also be involved in the control scheme of walking and deserve further exploration
together with muscle activation.

Conclusion

Here, we provide among the first empirical evidence of locomoting humans accepting a metabolic
cost penalty in favor of a predefined, alternative neuromuscular criterion. When choosing between
crouch walking and walking on a series of increasing inclines, our healthy participants protected
both fatigue-like and effort-like activation cost metrics. This observation is concomitant with
activation cost prioritization in the control of steady-state walking (22) and running (21). Further
research is required to determine whether muscle activation is used to estimate global energy or
effort, or if local muscle costs are of greater importance in the control scheme of gait, independent
of their relationship to energy costs *per se*.

Acknowledgements

548 The authors would like to thank Dr. Suzanne Cox and Mr. Brekke Green for data collection
549 assistance and Drs. Brian Umberger, Ton van den Bogert, Lex Gidley, Jessica Selinger, Max
550 Donelan, Greg Sawicki, Monica Daley, Young-Hui Chang, and Joseph Hamill for thoughtful
551 discussions and/or feedback on earlier drafts of this manuscript. The authors also thank Mr. Randall
552 McCullough for his technical support.

References

1.   Browning RC, Kram R. Energetic cost and preferred speed of walking in obese vs. normal
weight women. Obesity. 2005;13(5):891–9.

2.   Browning RC, Baker EA, Herron JA, Kram R. Effects of Obesity and Sex on the Energetic
Cost and Preferred Speed of Walking. J Appl Physiol. 2006;100(2):390–8.

3.   Holt KG, Jeng SF, Ratcliffe R, Hamill J. Energetic cost and stability during human walking
at the preferred stride frequency. J Mot Behav. 1995;27(2):164–78.

4.   Willis WT, Ganley KJ, Herman RM. Fuel oxidation during human walking. Metabolism.
2005;54(6):793–9.

5.   Ralston HJ. Energy-speed relation and optimal speed during level walking. Eur J Appl
Physiol Occup Physiol. 1958;17(4):277–83.

6.   Minetti AE, Capelli C, Zamparo P, Di Prampero PE, Saibene F. Effects of stride frequency
on mechanical power and energy expenditure of walking. Med Sci Sport Exerc.
1995;27(8):1194–202.

7.   Molen NH, Rozendal RH, Boon W. Graphic representation of the relationship between
oxygen-consumption and characteristics of normal gait of the human male. Proc K Ned
Akad van Wet. 1971;75(4):305–14.

8.   Selinger JC, O'Connor SM, Wong JD, Donelan JM. Humans can continuously optimize
energetic cost during walking. Curr Biol. 2015;25(18):2452–6.

9.   Snaterse M, Ton R, Kuo AD, Donelan JM. Distinct fast and slow processes contribute to the
selection of preferred step frequency during human walking. J Appl Physiol.
2011;110(6):1682–90.

10.   Umberger BR, Martin PE. Mechanical power and efficiency of level walking with different
stride rates. J Exp Biol. 2007;210(18):3255–65.

11.   Zarrugh MY, Todd FN, Ralston HJ. Optimization of energy expenditure during level walking.
Eur J Appl Physiol Occup Physiol. 1974;33(4):293–306.

12.   Zarrugh MY, Radcliffe CW. Predicting Metabolic Cost of Level Walking. Eur J Appl Physiol
Occup Physiol. 1978;38(3):215–23.

13.   Donelan JM, Kram R, Kuo AD. Mechanical and metabolic determinants of the preferred step
width in human walking. Proc R Soc B. 2001;268(1480):1985–92.

14.   De Graaf ML, Hubert J, Houdijk H, Bruijn SM. Influence of arm swing on cost of transport
during walking. Biol Open. 2019;8(6).

15.   Ortega JD, Fehlman LA, Farley CT. Effects of aging and arm swing on the metabolic cost
of stability in human walking. Journal of biomechanics, 41(16), 3303-3308. J Biomech.
2008;41(16):3303–8.

16.   Abram SJ, Selinger JC, Donelan JM. Energy optimization is a major objective in the real-
time control of step width in human walking. J Biomech. 2019;91:85–91.

17.   Roemmich RT, Leech KA, Gonzalez AJ, Bastian AJ. Trading symmetry for energy cost
during walking in healthy adults and persons poststroke. Neurorehabil Neural Repair.
2019;33(8):602–13.

18.   Wang WJ, Crompton RH, Li Y, Gunther MM. Energy transformation during erect and 'bent-
hip, bent-knee'walking by humans with implications for the evolution of bipedalism. J Hum
Evol. 2003;44(5):563–79.

19.   Rodman PS, McHenry HM. Bioenergetics and the origin of hominid bipedalism. Am J Phys
Anthropol. 1980;52(1):103–6.

20.   Hreljac A. Preferred and energetically optimal gait transition speeds in human locomotion.

Med Sci Sport Exerc. 1993;25(10):1158–62.

21.   Miller RH, Umberger BR, Hamill J, Caldwell GE. Evaluation of the minimum energy
hypothesis and other potential optimality criteria for human running. Proc R Soc B.
2012;279(1733):1498–505.

22.   Ackermann M, van den Bogert AJ. Optimality principles for model-based prediction of
human gait. J Biomech. 2010;43(6):1055–60.

23.   Falisse A, Serrancolí G, Dembia CL, Gillis J, Jonkers I, De Groote F. Rapid predictive
simulations with complex musculoskeletal models suggest that diverse healthy and
pathological human gaits can emerge from similar control strategies. J R Soc Interface.
2019;16(157):20190402.

24.   Veerkamp K, Waterval NFJ, Geijtenbeek T, Carty CP, Lloyd DG, Harlaar J, et al. Evaluating
cost function criteria in predicting healthy gait. J Biomech. 2021;110530.

25.   Gidley AD, Marsh AP, Umberger BR. Performance criteria for generating predictive optimal
control simulations of bicycle pedaling. Comput Methods Biomech Biomed Engin.
2019;22(1):11–20.

26.   Song S, Geyer H. Predictive neuromechanical simulations indicate why walking
performance declines with ageing. J Physiol. 2018;596(7):1199–210.

27.   Silder A, Besier T, Delp SL. Predicting the metabolic cost of incline walking from muscle
activity and walking mechanics. J Biomech. 2012;45(10):1842–9.

28.   Russell DM, Apatoczky DT. Walking at the preferred stride frequency minimizes muscle
activity. Gait Posture. 2016;45:181–6.

29.   Jaramillo J. Are muscles tuned to the preferred stride frequency of walking? University of
Massachusetts Amherst; 2020.

30.   Prilutsky BI, Gregor RJ. Swing-and support-related muscle actions differentially trigger
human walk–run and run–walk transitions. J Exp Biol. 2001;204(13):2277–87.

31.   Stenum J, Choi JT. Neuromuscular effort predicts walk–run transition speed in normal and
adapted human gaits. J Exp Biol. 2016;219(18):2809–13.

32.   Segers V, Lenoir M, Aerts P, De Clercq D. Influence of M. tibialis anterior fatigue on the
walk-to-run and run-to-walk transition in non-steady state locomotion. Gait Posture.
2007;25(4):639–47.

33.   Neptune RR, Hull ML. A theoretical analysis of preferred pedaling rate selection in
endurance cycling. J Biomech. 1999;32(4):409–15.

34.   Marsh AP, Martin PE. The association between cycling experience and preferred and most
economical cadences. Med Sci Sport Exerc. 1993;25(11):1269–74.

35.   Vercruyssen F, Brisswalter J. Which factors determine the freely chosen cadence during
submaximal cycling? J Sci Med Sport. 2010;13(2):225–31.

36.   Rasmussen J, Damsgaard M, Voigt M. Muscle recruitment by the min/max criterion—a
comparative numerical study. J Biomech. 2001;34(3):409–15.

37.   Crowninshield RD, Brand RA. A physiologically based criterion of muscle force prediction
in locomotion. J Biomech. 1981;14(11):793–801.

38. Redfield R, Hull ML. Prediction of pedal forces in bicycling using optimization methods. J
Biomech. 1986;19(7):523–40.

39. Dingwell JB, Marin LC. Kinematic variability and local dynamic stability of upper body
motions when walking at different speeds. J Biomech. 2006;39(3):444–52.

40. McDonald KA, Cusumano JP, Peeling P, Rubenson J. Multi-objective control in human
walking: insight gained through simultaneous degradation of energetic and motor regulation
systems. J R Soc Interface. 2019;16(158):20190227.

41. Stegeman D, Hermens H. Standards for surface electromyography: The European project
Surface EMG for non-invasive assessment of muscles (SENIAM). 2007;

42. Peronnet F, Massicotte D. Table of nonprotein respiratory quotient: an update. Can J Sport
Sci. 1991;16(1):23–9.

43. Kipp S, Byrnes WC, Kram R. Calculating metabolic energy expenditure across a wide range
of exercise intensities: the equation matters. Appl Physiol Nutr Metab. 2018;43(6):639–42.

44. Devaprakash D, Weir GJ, Dunne JJ, Alderson JA, Donnelly CJ. The influence of digital filter
type, amplitude normalisation method, and co-contraction algorithm on clinically relevant
surface electromyography data during clinical movement assessments. J Electromyogr
Kinesiol. 2016;31:126–35.

45. Kram R, Taylor CR. Energetics of running: a new perspective. Nature. 1990;346(6281):265–
7.

46. Roberts TJ, Kram R, Weyand PG, Taylor CR. Energetics of bipedal running. I. Metabolic
cost of generating force. J Exp Biol. 1998;201(19):2745–51.

47. Carrier D, Anders C, Schilling N. The musculoskeletal system of humans is not tuned to
maximize the economy of locomotion. Proc Natl Acad Sci. 2011;108(46):18631–6.

48. Handsfield GG, Meyer CH, Hart JM, Abel MF, Blemker SS. Relationships of 35 lower limb
muscles to height and body mass quantified using MRI. J Biomech. 2014;47(3):631–8.

49. Alexander RM. Optimization and gaits in the locomotion of vertebrates. Physiol Rev.
1989;69(4):1199–227.

50. Hunter LC, Hendrix EC, Dean JC. The cost of walking downhill: is the preferred gait
energetically optimal? J Biomech. 2010;43(10):1910–5.

51. Rosenbaum DA, Gong L, Potts CA. Pre-crastination: hastening subgoal completion at the
expense of extra physical effort. Psychol Sci. 2014;25(7):1487–96.

52. Kistemaker DA, Wong JD, Gribble PL. The central nervous system does not minimize
energy cost in arm movements. J Neurophysiol. 2010;104(6):2985–94.

53. Westling G, Johansson RS. Factors influencing the force control during precision grip. Exp
Brain Res. 1984;53(2):277–84.

54. Balasubramanian R, Howe RD, Matsuoka Y. Task performance is prioritized over energy
reduction. IEEE Trans Biomed Eng. 2009;56(5):1310–7.

55. De Rugy A, Loeb GE, Carroll TJ. Muscle coordination is habitual rather than optimal. J
Neurosci. 2012;32(21):7384–91.

56. Yandell MB, Zelik KE. Preferred Barefoot Step Frequency is Influenced by Factors Beyond

Minimizing Metabolic Rate. Sci Rep. 2016;6:23243.

57.   Monsch ED, Franz CO, Dean JC. The effects of gait strategy on metabolic rate and
681   indicators of stability during downhill walking. J Biomech. 2012;45(11):1928–33.

682   58.   Finley JM, Bastian AJ, Gottschall JS. Learning to be economical: the energy cost of walking
683   tracks motor adaptation. J Physiol. 2013;591(4):1081–95.

684   59.   Huang HJ, Ahmed AA. Older adults learn less, but still reduce metabolic cost, during motor
685   adaptation. J Neurophysiol. 2014;111(1):135–44.

686   60.   Whiting CS, Allen SP, Brill JW, Kram R. Steep (30°) uphill walking vs. running: COM
687   movements, stride kinematics, and leg muscle excitations. Eur J Appl Physiol.
688   2020;120(10):2147–57.

689   61.   Pontzer H. Economy and endurance in human evolution. Curr Biol. 2017;27(12):R613-21.

690   62.   Bramble DM, Lieberman DE. Endurance running and the evolution of Homo. Nature.
691   2004;432(7015):345–52.

692   63.   Noakes TDO. Fatigue is a brain-derived emotion that regulates the exercise behavior to
693   ensure the protection of whole body homeostasis. Front Physiol. 2012;3:82.

694   64.   Proske U, Allen T. The neural basis of the senses of effort, force and heaviness. Exp Brain
695   Res. 2019;237(3):589–99.

696   65.   Monjo F, Terrier R, Forestier N. Muscle fatigue as an investigative tool in motor control: A
697   review with new insights on internal models and posture–movement coordination. Hum Mov
698   Sci. 2015;44:225–33.

699   66.   Wong JD, O'Connor SM, Selinger JC, Donelan JM. Contribution of blood oxygen and carbon
700   dioxide sensing to the energetic optimization of human walking. J Neurophysiol.
701   2017;118(2):1425–33.

702   67.   Woledge RC, Curtin NA, Homsher E. Energetic aspects of muscle contraction. Monogr
703   Physiol Soc. 1985;41:1–357.

704   68.   Umberger BR, Gerritsen KG, Martin PE. A model of human muscle energy expenditure.
705   Comput Methods Biomech Biomed Engin. 2003;6(2):99–111.

706   69.   Beck ON, Punith LK, Nuckols RW, Sawicki GS. Exoskeletons improve locomotion economy
707   by reducing active muscle volume. Exerc Sport Sci Rev. 2019;47(4):237–45.

708   70.   Roberts TJ, Marsh RL, Weyand PG, Taylor CR. Muscular force in running turkeys: the
709   economy of minimizing work. Science (80-). 1997;275(5303):1113–5.

710   71.   Luu BL, Smith JL, Martin PG, McBain RA, Taylor JL, Butler JE. Feedforward consequences
711   of isometric contractions: effort and ventilation. Physiol Rep. 2016;4(15):e12882.

712   72.   Berniker M, Kording K. Estimating the sources of motor errors for adaptation and
713   generalization. Nat Neurosci. 2008;11(12):1454–61.

714   73.   Xiang Y, Arora JS, Abdel-Malek K. Physics-based modeling and simulation of human
715   walking: a review of optimization-based and other approaches. Struct Multidiscip Optim.
716   2010;42(1):1–23.

717   74.   Taylor JL, Amann M, Duchateau J, Meeusen R, Rice CL. Neural contributions to muscle
718   fatigue: from the brain to the muscle and back again. Med Sci Sport Exerc.

719   2016;48(11):2294.

721 **Figures and tables**

722
Figure 1. Experimental design. Participants were asked to select between crouch walking, which
moderately affects metabolic cost and penalizes activation cost, and a series of incline levels that
incrementally increase both metabolic and activation costs. Competing-Cost-Pairs were
established when participants walked on an incline level that incurred a higher metabolic cost but
provided an activation cost advantage relative to the crouch condition. A virtual ceiling and the
participant's silhouette, displayed on a screen anterior to the treadmill, provided feedback regarding
crouch walking task performance.

730
Figure 2. Experimental setup for crouch walking. Participants walked on a split-belt treadmill with
their silhouette projected onto an anteriorly placed screen. Their objective was to avoid collisions
between their silhouette and a virtual ceiling, which was set to 93% of their standing height. Real-
time motion-tracking and visual feedback was implemented in a custom program developed with
the openFrameworks open-source C++ toolkit, using data from a motion sensing device (Xbox
Kinect, Microsoft, WA, USA), situated behind the participant.

737
Figure 3. Walking data for the (A-D) mean (±SD) crouch trial versus five levels of incline (from 0-
24%) and (E-H) mean (±SD) crouch trial versus pre- and post-transition inclines. (A, E) Metabolic
power ($C_{met,P}$; W kg^{-1}; N=10), (B, F) sum of seven lower limb muscles' normalized activations
squared ($\bar{C}_{a^2}$; unitless; N=8), (C, G) maximum activation value from seven lower limb muscles'
normalized activation ($\bar{C}_{a,max}$; unitless; N=8), and (D, H) sum of the seven lower limb muscles'
volume-weighted activations ($\bar{C}_{a,vol}$; unitless; N=8). The number of participants that selected incline
(i) or crouch (c) in each condition are indicated in (A) (grey text).

*Significantly different (p<0.017) from crouch (E-H only); ^Significantly different (p<0.017) from pre-
transition (E-H only).

Figure 4 Average (*N*=8) normalized activation costs (A_{ij}) during crouch walking (level treadmill) and incline walking on treadmill slopes of 0-24%. Seven lower limb muscles were analyzed; gluteus maximus (Gmax), biceps femoris (BF), rectus femoris (RF), vastus medialis (VM), medial gastrocneuimus (MG), soleus (SOL) and tibialis anterior (TA). The unitless quantity A_{ij} was established by integrating the linear envelope (activation integral (a_{ij})), expressing this as a rate $\left(\frac{a_{ij}}{T_j}\right)$ and normalizing data for each muscle to the average value obtained from the 0% incline trial.

Figure 5. Competing-Cost-Pairs were established for seven of the eight participants for which this
was feasible. All seven participants selected the incline condition that favored the sum of activations
squared ($\bar{C}_{a^2}$) and penalized metabolic power ($C_{met,P}$). In doing this, they rejected the crouch
walking condition that favored metabolic power and penalized the sum of the activations squared.
Data is presented as the mean±SD.

*Significantly different (p<0.05) from crouch $C_{met,P}$; ^Significantly different (p<0.05) from crouch $\bar{C}_{a^2}$.

Table 1. Competing-Cost-Pairs (CCP) were established for seven of the eight participants for which this was feasible. The presence of a Competing-Cost-Pair was determined by comparing the mean crouch and pre-transition incline's percentage change from upright (normal) walking on a 0% incline for: i) metabolic power ($C_{met,P}$) and the sum of activations squared ($\bar{C}_{a^2}$), ii) $C_{met,P}$ and the maximal muscle activation ($\bar{C}_{a,max}$), and iii) $C_{met,P}$ and the sum of the volume-weighted activations ($\bar{C}_{a,vol}$). The pre-transition incline is defined as the steepest incline that was selected over crouch.

Participant ID	$C_{met,P}$* % Crouch	$C_{met,P}$* % Pre-transition incline	$\bar{C}_{a^2}$* % Crouch	$\bar{C}_{a^2}$* % Pre-transition incline	$\bar{C}_{a^2}$* % CCP established?	$\bar{C}_{a,max}$* % Crouch	$\bar{C}_{a,max}$* % Pre-transition incline	$\bar{C}_{a,max}$* % CCP established?	$\bar{C}_{a,vol}$* % Crouch	$\bar{C}_{a,vol}$* % Pre-transition incline	$\bar{C}_{a,vol}$* % CCP established?
1	172.5	189.5	566.0	290.8	Y	292.2	196.8	Y	167.1	125.7	Y
2	226.4	236.3	1382.3	794.3	Y	575.9	399.7	Y	308.1	236.9	Y
3	229.9	266.9	1410.1	799.4	Y	533.8	463.1	Y	222.5	161.4	Y
4	137.6	182.5	605.2	373.1	Y	429.4	227.2	Y	169.5	155.9	Y
5	170.9	200.6	NA	NA	-	NA	NA	-	NA	NA	-
6	204.9	202.5	1409.4	454.2	N	747.6	368.4	N	239.7	154.4	N
7	174.6	245.6	NA	NA	-	NA	NA	-	NA	NA	-
8	137.6	199.3	284.4	211.4	Y	219.5	165.6	Y	166.9	143.4	Y
9	169.6	215.9	584.3	308.0	Y	380.7	249.3	Y	198.8	156.8	Y
10	160.1	217.0	469.2	359.1	Y	297.3	233.9	Y	167.2	154.2	Y

*Significant difference ($p<0.05$) between the crouch and pre-transition incline trials for participants with a CCP ($N=7$).

Supplementary material

768 **Table S1.** Competing-Cost-Pairs (CCP; $N=7$) and whole-group ($C_{met,P}$: $N=10$; $\bar{C}_{a^2}$: $N=8$; $\bar{C}_{a,max}$: $N=8$; $\bar{C}_{a,vol}$: $N=8$) statistical analyses using the
769 mean crouch, initial crouch, and final crouch data. The pre-transition incline is defined as the steepest incline that was selected over crouch, whereas
770 the post-transition incline is the first rejected incline in favor of crouch.
771

			$C_{met,P}$		$\bar{C}_{a^2}$		$\bar{C}_{a,max}$		$\bar{C}_{a,vol}$	
			CCP	Group	CCP	Group	CCP	Group	CCP	Group
Mean crouch	Distribution		Normal	Normal	Non-normal	Non-normal	Normal	Normal	Normal	Normal
	Main effect	p-value	-	<0.001*	-	0.008*	-	0.001*	-	0.003*
	Crouch vs. pre-transition incline		0.002*	0.007*	0.018*	0.012*	0.003*	0.006*	0.007*	0.003*
	Crouch vs. post-transition incline		-	0.005*	-	0.263	-	0.214	-	0.779
	Pre-transition incline vs. post-transition incline		-	0.005*	-	0.017*	-	0.003*	-	0.002*
Initial crouch	Distribution		Normal	Normal	Non-normal	Non-normal	Normal	Normal	Normal	Normal
	Main effect	p-value	-	<0.001*	-	0.003*	-	0.002*	-	0.002*
	Crouch vs. pre-transition		0.002*	0.007*	0.018*	0.008*	0.003*	0.008*	0.008*	0.003*
	Crouch vs. post-transition		-	0.005*	-	0.071	-	0.117	-	0.329
	Pre-transition incline vs. post-transition incline		-	0.005*	-	0.011*	-	0.003*	-	0.002*
Final crouch	Distribution		Normal	Normal	Non-normal	Non-normal	Normal	Normal	Normal	Normal
	Main effect	p-value	-	<0.001*	-	0.013*	-	0.001*	-	0.002*
	Crouch vs. pre-transition		0.002*	0.005*	0.018*	0.013*	0.004*	0.005*	0.009*	0.005*
	Crouch vs. post-transition		-	0.005*	-	0.698	-	0.513	-	0.453
	Pre-transition incline vs. post-transition incline		-	0.005*	-	0.011*	-	0.003*	-	0.002*

772 *Significant difference ($p<0.05$ for CCP; $p<0.05$ for whole-group main effect; $p<0.017$ for whole-group *post hoc* test comparisons).